1 **Single capsid mutations modulating phage adsorption, persistence, and plaque**
2 **morphology shape evolutionary trajectories in ΦX174**
3
4
5 Key words:
6 **Adsorption, persistence, ΦX174, lytic bacteriophage, phage therapy, phage evolution**
7

8 Manuela Reuter[1], Michael Sieber[1], Octavio Reyes-Matte[1], Christina Vasileiou[1], Christopher
9 Böhmker[1], Jordan Romeyer Dherbey[2], Frederic Bertels[1,*] and Javier Lopez-Garrido[1,*]
10
11 [1]Max Planck Institute for Evolutionary Biology, Plön, Germany
12 [2]University of Cambridge, Cambridge, United Kingdom
13
14 *Correspondence: frederic.bertels.publications@gmail.com; lopezgarrido@evolbio.mpg.de

15
16

1 Abstract

The evolutionary success of lytic bacteriophages depends on key life-history traits, including
adsorption, lysis time, burst size, and persistence in the environment. However, how these traits
evolve to allow adaptation to different environments remains poorly understood. Here, we
explored this question in ΦX174, combining experimental evolution and mathematical
modelling. By investigating how serial transfer conditions shape evolutionary outcomes in
liquid culture, we found that the time between transfers imposes divergent selection on
adsorption and context-dependent directional selection on persistence. Longer transfer
intervals, which allow multiple infection cycles until host depletion, favoured fast-adsorbing,
highly persistent mutants that could rapidly initiate infections and remained viable in the
absence of the host. In contrast, shorter transfer intervals selected for slower adsorption without
substantially altering persistence. Mathematical modelling of phage population dynamics
predicted that adsorption evolution during short transfers reflects a trade-off between two
opposing selective forces within each transfer: an early phase in which susceptible hosts are
abundant and adsorption is productive, favouring fast adsorption, and a later phase in which
most hosts are already infected and adsorption primarily removes phage particles via
attachment to already infected cells, favouring slower adsorption. A single-point mutation in
the major capsid protein was sufficient to drive these changes in adsorption. In the case of fast-
adsorbing mutants, this mutation was positively pleiotropic and also enhanced environmental
persistence. Our findings show how simple changes in propagation conditions can steer phage
phenotypes, providing insights relevant to evolutionary biology and phage therapy.

Introduction

The success of lytic bacteriophages relies on four central life-history traits, each of which may evolve under different selective pressures to maximise fitness [1–6]. These traits are: (I) adsorption constant, which determines the ease with which a phage irreversibly attaches to a host cell, (II) latent period, or time required to complete the infection cycle inside the host, (III) burst size, defined as the number of new phage particles produced per infected host cell, and (IV) decay rate, representing the loss of infectious phage particles in the absence of susceptible host cells [7–12].

Interest in these traits has grown due to the need to predict phage virulence for medical and biocontrol applications [13–16]. However, such predictions remain challenging because the fitness effects of individual traits are often highly context-dependent [17–19]. For example, high adsorption rates are typically considered beneficial in well-mixed liquid environments, where they facilitate rapid infection initiation [20, 21]. However, in spatially-structured environments such as biofilms, these same traits can hinder phage spread to uninfected neighbouring cells by promoting phage attachment to already infected cells at the centre of infection [17, 22]. Understanding how life-history traits influence phage fitness across environments is therefore essential for predicting and potentially optimising phage performance.

The model coliphage ΦX174 provides a powerful platform for dissecting how the relationship between trait and environment impacts fitness [23–27]. ΦX174 is a strictly lytic, tailless phage of the *Microviridae* family, best known as the first DNA-based biological entity whose entire genome was sequenced [28]. Its 5386 bp circular single-stranded DNA genome encodes only 11 genes, most of which are necessary for phage DNA replication, morphogenesis, and host lysis [28–31]. Only four genes (F, G, H, and J) encode for proteins that form the mature virion [32, 33]. The icosahedral capsid is made of 60 F, 60 J and 12 H protein monomers, decorated at each of its 12 vertices by 5 G spike monomers [32, 34]. G likely mediates host recognition and subsequent attachment to host lipopolysaccharide (LPS), after which it dissociates, allowing F to bind irreversibly to the host cell surface [35–37]. Irreversible binding triggers a conformational change that results in the injection of phage DNA into the host cell through the pilot protein H [37, 38].

Here, we have explored how life-history traits influence phage fitness across different

environments by evolving ΦX174 under different serial transfer regimes. Through a

combination of life-history trait quantification and mathematical modelling, we have identified

individual mutations that alter adsorption constant and persistence, leading to fitness effects

that are strongly dependent on the environment. By disentangling these context-dependent

fitness effects, our results improve our understanding of phage ecology and evolution and also

provide insights for rational trait optimisation in medical and biocontrol applications.

Materials and Methods

Strains and culture conditions

The ancestral ΦX174 and its host *E. coli* C were provided by H.A. Wichman (Department of Biological Sciences, University of Idaho). All evolved phage strains were derived from the same ΦX174 genotype (GenBank: AF176034). Cultures were grown in Lysogeny Broth (LB, Miller) supplemented with 10 mM $MgCl_2$ and 5 mM $CaCl_2$ and incubated at 37°C, shaking at 250 rpm (orbital shaking diameter 2.5 cm). Mid-exponential *E. coli* C cultures (~$1\text{-}2*10^8$ colony-forming units [CFU] per ml) were prepared from overnight cultures and used for all transfer experiments and life history quantification assays.

Phage lysate preparation and titration

Phage lysates were generated as described in Romeyer Dherbey *et al.* [39] by co-culturing exponentially growing *E. coli* C with ΦX174 for 3 hours in a final volume of 5 ml. After 3 hours, the remaining bacteria were removed with 10-12 drops of chloroform and subsequent centrifugation at 5000 rpm for 10 minutes at 4°C. Phage-containing supernatants were collected and stored as lysate at 4°C or at -80°C as glycerol stocks for long-term storage.

Phage titers were quantified by plaque assays in which plaque-forming units (PFUs) were assessed by plating 100 µl of serially diluted phage lysate with 100 µl of *E. coli* C stationary-phase culture pre-mixed in 4 ml semi-solid agar (SSA, 0.5% agar, supplemented with 10 mM $MgCl_2$ and 5 mM $CaCl_2$). The mixture was overlayed on a plate of LB-agar (1.5%) and incubated at 37°C for 18 hours. Phage lysates were diluted in phage buffer juice (PBJ: 2.03% Tris-HCl, 0.61% $MgCl_2$).

Serial transfer regimes

We implemented two transfer regimes, the 3-h regime and the 30-min regime, which are described in detail below. For both transfer regimes, five independent selection lines were initiated by infecting mid-exponential phage-sensitive *E. coli* C cultures (~$1\text{-}2*10^8$ CFU/ml) with ancestral ΦX174 extracted from independent plaques. Each transfer was performed in a final volume of 5 ml of supplemented LB medium (see "Strains and culture conditions"). All transfer cultures were incubated at 37°C, shaking at 250 rpm (orbital shaking diameter 2.5 cm).

The 3-h and 30-min regimes are described in detail below. Additional details on the transfer
regimes can be found in **Fig. S1**.

3-h transfer regime

The 3-h regime was chosen based on a previous finding in our lab, which showed that
small plaques begin to emerge after approximately seven 3-hour transfers of ancestral ΦX174
[39]. This regime was initiated by infecting a 5-ml culture of exponentially growing *E. coli* C
(~1-2*10^8 CFU/ml) at an MOI of ~0.1, followed by a 3-h incubation at 37°C with shaking at
250 rpm. After the incubation period, phages were extracted with chloroform. The resulting
phage lysate was titrated and used to inoculate the next transfer on the following day at MOI_{input}
of ~0.1 on a fresh, mid-exponential host culture (~1-2*10^8 CFU/ml) (**Fig. 1**; **Fig. S1A**). We
repeated this cycle every day and performed titration assays after every transfer to quantify
titers and screen for changes in plaque morphology. Serial transfers were continued until
heritable small-plaque mutants emerged. When detected, these small plaques were isolated and
replated to confirm the phenotype.

30-min transfer regime

The 30-min regime was a modification of the 3-h regime in which 3-h transfers were
interspersed by blocks of four 30-min transfers (**Fig. 1**; **Fig. S1B**). The first 30-min transfer of
each block was initiated by infecting a 5-ml culture of exponentially growing *E. coli* C (~1-
2*10^8 CFU/ml) at an MOI of ~0.1. The culture was incubated for 30 min at 37°C with shaking
at 250 rpm, after which 2% of the phage-bacteria co-culture was transferred to 5 ml of a fresh,
mid-exponential *E. coli* C culture. This transfer was repeated after each 30-min incubation for
a total of four consecutive 30-min transfers. At the end of the fourth 30-min transfer, 2% of the
phage-bacteria co-culture was transferred to 5 ml of a fresh, mid-exponential *E. coli* C culture
and incubated for 3h at 37°C with shaking at 250 rpm. After this incubation, phages were
extracted with chloroform. The lysate was titrated and used to seed the next 30-min transfer
block on the following day. Plaques larger than the ancestral ones emerged in all five lineages
(see results). Phage lysate of each transfer day was stored for Sanger sequencing.

Sanger sequencing of evolved phages

Whole-genome Sanger sequencing was conducted for individual phage isolates from both transfer regimes according to Biggs *et al.* [40]. DNA was extracted from high-titer lysates (QIAprep Spin Miniprep Kit® Qiagen, Hilden, Germany) and amplified via PCR in two genome halves. PCR products were purified (QIAprep PCR Purification Kit® Qiagen, Hilden, Germany) and sequenced at the Max Planck Institute for Evolutionary Biology. Primers and settings used for amplification and sequencing are in **Tables S1, S2, and S3**. Sequences were aligned against the reference ΦX174 genome using Geneious Prime 2023.0.1 (https://www.genious.com). Identified amino acid positions in which single-point mutations were located were numbered according to the residue number, which differs from the codon number by minus one because the first amino acid (methionine) of the F protein is cleaved [32, 41]. Raw Sanger Sequencing files are deposited in Zenodo (see Zenodo doi: 10.5281/zenodo.17076220).

Adsorption assays

Adsorption constants were determined following a modified version of the Hyman and Abedon assay [42]. Mid-exponential, phage-sensitive *E. coli* C cultures were co-cultured with phages at MOI_{input} ~0.05 at 37°C in a shaking water bath at 200 rpm. Phages were sampled 2, 4, and 6 minutes after adsorption initiation and diluted 10-fold into non-supplemented LB. Samples were cooled on ice to avoid temperature-dependent decay effects, treated with chloroform to stop infection, and quantified. Adsorption constants were calculated by fitting linear models to log-transformed phage titers and dividing slope estimates by bacterial density. Differences between genotypes were tested using a linear mixed model (replicate as a random effect) and post hoc Tukey tests for multiple comparisons (**Supplementary Methods**).

Phage persistence assays

Persistence was measured over a 5-hour co-culturing period of ΦX174 and an exponentially-growing *E. coli* C culture infected with phages at MOI_{input} ~0.1. Phage samples were taken every 30 minutes, extracted with chloroform, and phage titers were determined by PFU counting. Decay rates were estimated by fitting exponential decay models to the titer decline period after peak productivity. Model fitting procedures are detailed in the **Supplementary Methods**.

1   Simulations of phage evolution

The following system of delay differential equations (DDEs) describes the dynamics of

the bacterial host B (cells per ml), the infected bacterial population B_I (cells per ml), and the

three phage types P_A (ancestral, particles per ml), P_L (large-plaque type evolved in vitro in the

30-min regime, particles per ml), and P_S (small-plaque type evolved in vitro in the 3-h regime,

particles per ml). In addition, $B_\tau = B(t - \tau)$ denotes the bacterial density τ minutes in the past,

reflecting the delay introduced by the lysis time. We use a similar notation for the phage

densities, e.g. $P_{A,\tau}$ denotes the density of the ancestral phage τ minutes in the past. For $t < \tau$,

we set the corresponding densities to zero. Annotations of small and large plaque types were

based on the in-vitro phenotypes specified in the "Serial Transfer Section" to simplify labelling.

However, plaque morphology was not included in the in-silico modelling.

Bacterial growth throughout the transfer period was found not to play a significant role

in the simulations, so it is neglected in the model for simplicity. All parameter values for phage

life traits and their biological meaning are given in **Table 1**. Beyond the values given in Table

1, we tested the robustness of the in-silico models for the 30-min and 3-h regimes to variation

in central phage life history traits (i.e. lysis time and burst size) (**Fig. S2**).

$$\frac{dB}{dt} = -(\phi_A P_A + \phi_L P_L + \phi_S P_S)B$$

$$\frac{dB_I}{dt} = \phi_A(BP_A - B_\tau P_{A,\tau}) + \phi_L(BP_L - B_\tau P_{L,\tau}) + \phi_S(BP_S - B_\tau P_{S,\tau})$$

$$\frac{dP_A}{dt} = \phi_A(\beta B_\tau P_{A,\tau} - (B + B_I)P_A) - \delta_A P_A$$

$$\frac{dP_L}{dt} = \phi_L(\beta B_\tau P_{L,\tau} - (B + B_I)P_L) - \delta_L P_L$$

$$\frac{dP_S}{dt} = \phi_S(\beta B_\tau P_{S,\tau} - (B + B_I)P_S) - \delta_S P_S$$

The DDE system was solved numerically in Python using the *scipy*-based *ddeInt* solver.

After 30 minutes or 3 hours of simulated time, depending on the transfer regime, the system

was reset to the initial density of fresh bacteria B (10^8 CFU). In the 3-h regime, each transfer

started with 10^7 PFU from the previous transfer. In the 30-min regime, the first transfer was

also initiated with 10^7 PFU, and subsequent transfers with 2% of the final densities of infected

bacteria B_I. For the competition simulations, phages with different life history traits were

introduced at different initial densities, as indicated in the corresponding figure legends. The

model makes the following simplifying assumptions. First, it assumes a fixed lysis time of 17 min, according to our experimental calculations (**Fig. S3**). Second, it assumes that infected host cells disappear upon lysis and do not contribute to phage attachment as debris. Third, the model assumes a constant phage decay rate. The code to reproduce the figures is available at https://github.com/OReyesMatte/Reuter_2025/tree/main/Simulations.

Image analysis of lysis plaques

Lysis plaque morphology was quantified using a custom image analysis pipeline. Plaques were imaged under standardised conditions (ChemiDoc: Bio-Rad, California, USA; 0.1s exposure time) and processed using Ilastik [43] for pixel classification based on intensity, texture, and edge features. Post-processing steps included erosion, whole-filling, and watershed transformation using SciPy [44] and scikit-image [45]. Each plaque was assigned an ID and analysed for area (cm²), distance to periphery (via distance transform), and grayscale intensity (overlaying mask on original image) (details see **Supplementary Methods**).

Data analysis and visualisation

Statistical analysis of adsorption constants, model fitting for adsorption and decay, as well as all data visualisations, were performed in R (version 4.2.1) [46]. Details of the packages and scripts are in https://github.com/OReyesMatte/Reuter_2025/tree/main/Data_analysis and **Table S4**. The protein structure, resolved by McKenna *et al.* [32], shown in **Fig. 4A**, was taken from RCSB PDB (reference 2BPA: https://www.wwpdb.org/pdb?id=pdb_00002bpa), and the mutation visualisation was done in UCSF Chimera (version 1.18). All cartoons were created using BioRender® as indicated in the corresponding figure legends.

Results

Differences in serial transfer regimes drive the evolution of contrasting lysis plaque

phenotypes

In the context of a previous experimental evolution study [39], we observed that small,

turbid plaques frequently emerged when ΦX174 was serially transferred after 3-hour co-

culturing periods with host cells in liquid, shaking conditions. To confirm this observation, we

evolved multiple independent selection lines of the ancestral ΦX174 under the same selection

regime (hereafter, 3-h regime) and plated the evolving phage populations onto lawns of phage-

sensitive host cells at every transfer (**Fig. 1; Fig. S1A**). Small, turbid plaques appeared in all

lines after four to seven transfers (**Fig. S4A**). In parallel, we tested a modified regime in which

the 3-h incubation periods were interspersed by four consecutive 30-min transfers, each

initiated by transferring 2% of the phage-bacteria mixture into fresh, sensitive host cells

(hereafter, referred to as the 30-min regime). In contrast to the 3-h transfer regime, no small

plaques were detected under the 30-min regime, even after 50 transfers (**Fig. 1, Fig. S1B**).

However, we began to observe plaques larger than those formed by the ancestral phage after

approximately 20 transfers (**Fig. S4B**).

Large- and small-plaque phenotypes are caused by different mutations in the major

capsid (F) protein

Since differences in plaque morphology often reflect underlying changes in life history

traits [2, 17, 22, 47–49], we hypothesised that our experimental regimes were selecting for

different life-history trait changes in the 3-h and 30-min regimes. To identify the genetic

modifications leading to the small- and large-plaque phenotypes, we isolated five phages of

each type, one per selection line, and sequenced their genomes. Each evolved isolate differed

from the ancestor by only one or two single-point mutations (**Fig. 2**).

All five large-plaque mutants isolated from the 30-min regime lines carried an A-to-G

transition at position 1301 in the phage genome, resulting in a threonine-to-alanine substitution

at residue 100 of the F gene [A1301G/F(T100A); we use this notation throughout the

manuscript: original nucleotide, nucleotide position, new nucleotide/protein (original amino

acid, residue position, new amino acid)] (**Fig. 2A**). The mutant isolated from line 3 carried only

this substitution. The other four isolates carried one additional mutation in either the F or the

H protein (**Fig. 2A**) acquired after the F(T100A) mutation. Phenotypically, however, large-plaque-formers resembled each other, suggesting that the T100A mutation in the F protein is sufficient for the manifestation of the large-plaque phenotype.

None of the mutations observed in large-plaque mutants were found in the small-plaque-forming phages isolated from the 3-h regime lines. However, all five small-plaque phages also differed from the ancestor by a single-point mutation in the F gene (**Fig. 2B**), each producing a distinct small-plaque phenotype (**Fig. 2B; Fig. S5**). The most drastic phenotype was caused by the G1965A/F(G321D) mutation, resulting in the formation of extremely small and turbid plaques. Phages carrying the C2280A/F(S426*) mutation, which introduces an early stop codon and truncates the major capsid protein by the last amino acid, produced slightly larger and clearer plaques (**Fig. S5**). The G1639A/F(M212I) mutation led to a heterogeneous phenotype, with both small and large plaques, and was excluded from further analysis.

Altogether, the sequencing results indicate that different modifications of the major capsid protein are responsible for the contrasting plaque phenotypes emerging in our selection regimes.

Plaque size negatively correlates with the adsorption constant

The major capsid protein is critical for the irreversible adsorption of ΦX174 to the host cells [37], suggesting the large- and small-plaque-forming phages may have evolved different adsorption constants compared to the ancestral ΦX174. To test this hypothesis, we measured the adsorption constant of the ancestral phage, a large-plaque-forming mutant carrying the single F(T100A) mutation evolved in the 30-min regime, and the two small-plaque mutants, F(G321D) and F(S426*), isolated from the 3-h regime. We focused on mutants with single substitutions to ensure that each mutation could be directly linked to specific changes in life history traits.

The adsorption constants of the evolved phages differed significantly from that of the ancestral ΦX174 and correlated inversely with plaque size (**Fig. 3**). The ancestral mean adsorption constant was 4.15×10^{-9} ml^{-1} min^{-1} (**Table S5**), in close agreement with previous estimations [50, 51]. The adsorption constants of the small-plaque-forming mutants evolved in the 3-h regime were slightly but significantly higher than those of the ancestral. Conversely,

the large-plaque mutant evolved in the 30-min regime had an absorption rate constant approximately five-fold lower than the ancestral (**Table S5**).

The relationship between adsorption constant and plaque size was not linear. The large-plaque mutant with a five-fold reduction in adsorption constant produced plaques that were 1.5-fold larger than those of the ancestor, whereas the small-plaque mutants with less than two-fold increase in adsorption constant formed plaques about ten times smaller (**Table S5**). These results are in line with previous predictions of a non-linear relationship between adsorption constant and plaque size [49].

These results demonstrate that the adsorption constants evolved in opposite directions under the 3-h and 30-min regimes, and suggest that the differences in plaque size may be caused by variation in phage-host cell binding affinity.

3-h transfers select for high persistence outside the host

We mapped the mutations underpinning the small- and large-plaque phenotypes onto the ΦX174 F-protein structure. The F(T100A) substitution in the large-plaque-forming mutant evolved in the 30-min regime was located in the centre of the F monomer (**Fig. 4A**). However, the two small-plaque-associated mutations F(G321D) and F(S426*) found in isolates of the 3-h regime were at the peripheral interfaces between F monomers in the assembled capsid (**Fig. 4A**). These peripheral mutations might influence capsid symmetry or stability by altering inter-subunit interactions [52], potentially contributing to the persistence of small-plaque-forming mutant phages during extended incubation periods, such as our 3-h regime.

To test this, we performed in vitro time-course phage-bacteria co-culturing using the ancestral ΦX174, the large-plaque mutant F(T100A) from the 30-min regime, and the two small-plaque mutants (i.e. F(G321D and F(S426*)) evolved in the 3-h regime. Co-cultures were initiated at an MOI_{input} of ~0.1, and infectious phage particles were quantified at regular intervals over five hours following chloroform extraction. All phage populations exhibited an initial productive phase, during which infectious particles increased, followed by a decay phase after the depletion of susceptible host cells (**Fig. S6**). The productive phase differed slightly between the different phages. The ancestor and the large-plaque-forming mutant evolved in the 30-min regime reached their maximum titer of infective phages ~60 minutes post-infection, with a ~5,000-fold increase in population size. Small-plaque mutants evolved in the 3-h regime

peaked slightly later but achieved a 10,000-fold increase in population size. The higher population size peak of small-plaque mutants may be attributed to their higher adsorption constant, allowing for earlier infection initiation. The most striking differences, however, were observed during the decay phase (**Fig. 4B**). Both the ancestor and the large-plaque-forming mutant evolved in the 30-min regime showed a drastic reduction in infective phage particles, with titers declining more than 15-fold by the end of the experiment. In contrast, the small-plaque mutants evolved in the 3-h regime maintained a relatively constant number of infective particles throughout the remainder of the assay, showing only a minor decline over time.

In silico, adsorption and persistence contribute to the selection of small-plaque mutants in the 3-h regime

To explore the roles of absorption rate and persistence in phage fitness under different transfer regimes, we developed a mathematical model of phage infection dynamics. The model captures the interaction between the bacterial host and phage with different adsorption constants or decay rates during the different transfer experiments. Upon adsorption of a phage to an uninfected host cell, the cell moves into the infected class. Phages of all types continue to adsorb to already infected cells, but only phages corresponding to the type that infected first are released upon lysis. We explicitly consider the lysis time, reflecting the time delay between adsorption of a phage and lysis of the infected cell, triggering the release of new viral particles. The full model is given in the **Methods**.

We started by exploring the role of the absorption constant in the 3-h regime. We simulated 3-h transfers with phages with an intermediate adsorption constant representing the ancestral phage, and introduced phages with higher and lower adsorption constants at low frequency to monitor changes in their frequencies throughout the simulations. All other life-history traits, including persistence, were held constant (**Table 1**). The simulations predicted an increase in the frequency of fast adsorbing phages at the expense of phages with intermediate and low adsorption constants (**Fig. 4C, top graph**), indicating that fast adsorption could explain the evolution of the small plaque-mutants under the 3-h regime.

Next, we tested if differences in decay rates could also explain the evolution of the small-plaque mutants in the 3-h regime. We incorporated the in-vitro decay rates into the in-silico 3-hour serial transfer model (**Table 1**), this time keeping adsorption constant at intermediate levels. The simulation predicted the rapid increase in the frequency of phages with

higher persistence (**Fig. 4C, bottom graph**), indicating that increased persistence provides a larger fitness gain than increased adsorption in the 3-h regime. Thus, the F(G321D) and F(S426*) mutations evolved in the 3-h regime are positively pleiotropic and enhance fitness by increasing both adsorption and persistence.

The adsorption optimum reflects a within-transfer trade-off between productive and unproductive adsorption in the 30-min regime

While the evolution of fast adsorption and increased persistence in the 3-h regime is intuitively understandable, the evolution of a lower adsorption constant under the 30-min regime is less intuitive. We expected fast adsorption to be beneficial in well-mixed liquid cultures as it would allow rapid infection of host cells [20, 21]. Hence, we were surprised that phages with dramatically reduced adsorption constants were selected under the 30-min regime.

We used our model to explore the role of the adsorption constant on fitness under the 30-min regime. First, we simulated competition between the ancestral phage and hypothetical mutants with adsorption constants ranging from the ancestor value to 1000-fold lower over a single 30-min transfer (**Fig. 5A**). Each simulation began with a 1:1 ratio of ancestral and mutant phages at an input multiplicity of infection (MOI_{input}) of 0.1, and predicted the relative mutant titer after 30 minutes, corresponding the first 30-min transfer of our regime. The simulation predicted an optimal adsorption constant approximately 6-fold lower than that of the ancestor, very close to the value observed in the F(T100A) mutant. This optimum was robust across a range of burst sizes and lysis times that allow fewer than two infection cycles (**Fig. S2**). In silico, mutants with very low adsorption constants performed poorly due to limited infection initiation, whereas mutants with very high adsorption also performed poorly because rapid adsorption increases loss to binding of already infected cells, which acted as a superinfection sink. As a result, slower-adsorbing phages were predicted to remain free at a higher frequency at the end of the transfer (**Fig. 5B**) and therefore be available to initiate infections after dilution into fresh hosts. Consistent with this, in vitro measurements showed a higher number of free phages for slow-adsorbing mutants evolved in the 30-min regime than for ancestral phages (**Fig. 5C**).

To explore this further, we simulated phage dynamics over multiple consecutive 30-min transfers (**Fig. 6A**). Within-transfer dynamics converged after two transfers to a repeating pattern that recurred from transfer to transfer, characterized by a major burst of phage release

and two phases within each transfer: an early phase with high host availability, in which many phages attach to uninfected cells and can generate progeny (hereafter, the productive phase), followed by a longer phase with low host availability, in which additional adsorption increasingly target cells that are already infected and, in our model, does not contribute to progeny production (hereafter, the unproductive phase) (**Fig. 6B-D**). The sharp separation between these phases is likely accentuated by the deterministic assumptions of our model, including a fixed 17-min lysis time. In vitro, lysis times are probably more broadly distributed, which would smooth the transition between phases. Regardless, this within-transfer structure imposes opposing selection pressures on adsorption. Faster adsorption increases productive infections during the productive phase, whereas lower adsorption reduces losses to an unproductive adsorption sink during the late unproductive phase. Hence, the optimal adsorption constant depends on the balance between these opposing selective pressures and should vary with the strength and duration of each phase under a given transfer protocol.

We reasoned that changing the dilution factor between transfers should modify the relative weight of the early productive phase versus the late unproductive phase, thereby shifting the adsorption optimum (**Fig. 6D; Fig. S8**). To test this, we simulated competitions between phages with different adsorption constants (low, intermediate, and high) under regimes with different dilution factors and transfer times, and mapped the fitness landscape (**Fig. 6E**). As expected, different dilution factors favoured different adsorption strategies. At high dilution, fast adsorption was generally favoured because each transfer begins with abundant susceptible hosts and relatively low phage densities, increasing the payoff of quickly initiating productive infections. At intermediate dilution, where the early productive phase and the late unproductive phase are more comparable in duration and intensity, phages with intermediate and even low adsorption constants were favoured. At very low dilution, fast adsorption again became advantageous in our simulations because transfers begin with high phage densities, susceptible host cells are rapidly depleted, and there is a strong selection to capture the short initial window in which productive infections are still possible. These outcomes should be interpreted in the context of model assumptions. In particular, we do not include potential additional adsorption sinks such as cell debris, which could further penalise fast-adsorbing phages, especially at low dilution factors.

1   Overall, the simulations indicate that selection on adsorption can be tuned by transfer-
2   regime parameters that alter the relative contribution of productive adsorption early in the
3   transfer versus adsorption losses later in the transfer.
4

Discussion

Bacteriophage life-history traits are commonly studied using experimental evolution under a variety of selection pressures, both in vitro and in silico [26, 27, 53–61]. Variation in host supply is a frequently used selection pressure providing limited or excess hosts at the onset of phage-bacteria co-culturing in serial-transfer experiments [2, 4, 62, 63]. In these experiments, evolution often targets the latency period and burst size, with excess hosts favouring shorter latency periods to maximise the rate of host exploitation, whereas host limitation selects for higher burst sizes to maximise yield per infection. Selection for an increased burst size often comes at the cost of longer latency periods, and vice versa, due to a lysis time-burst size trade-off [1, 64–67]. Our transfer regimes begin with hosts in excess, but host availability still changes markedly within transfers. In the 3-h regime, phages deplete susceptible hosts and must persist outside the host for much of each transfer, whereas in the 30-min regime, hosts are initially abundant but most become infected during the transfer. Our experimental results and simulations indicate that adaptation to our transfer regimes is not mediated by changes in latency period and burst size, but instead by the modulation of adsorption and persistence (**Fig. 3; Fig. 4**). We will dissect the reasons for this in the next paragraphs.

Longer transfer times select for phages with higher adsorption constants and greater ability to persist outside the host. This pattern can be explained by the ecological dynamics of the 3-h regime: given lysis time and host availability, multiple infection cycles can occur per transfer, leading to rapid host exhaustion. Fast-adsorbing phages initiate these cycles more efficiently. Moreover, since phage populations increase by more than two orders of magnitude per infection cycle and transfers are performed at an MOI_{input} of ~0.1, the vast majority of sensitive host cells are lysed after two cycles, forcing phages to persist in the absence of host cells for extended periods before being supplied with new susceptible hosts (**Fig. S5**). Thus, both enhanced adsorption and environmental persistence are selected in this regime.

Two different F mutations evolved in the 3-h regime increased adsorption and dramatically increased persistence. Both mutations were located at the monomer-monomer interface in the mature capsid (**Fig. 4A**). Changes in this region can affect capsid symmetry and stability in icosahedral phages [52, 68]. The positively pleiotropic nature of these mutations makes it difficult to disentangle the relative fitness contributions of adsorption and persistence. However, the drastic increase in persistence compared to the modest elevation in the adsorption

constant suggests that persistence is the primary trait under selection. This is further supported by our simulations, which show that although both increased adsorption and persistence are favoured under the 3-h regime, selection on persistence alone leads to a more rapid evolution of small-plaque mutants than selection on adsorption alone (**Fig. 4C**).

The rapid decay of the ancestral ΦX174 under our experimental conditions, with more than 90% of infectious particles lost in less than two hours (**Fig. 4B**), contrasts with the lower decay estimates previously reported [69]. Particle loss due to attachment to debris could contribute to our observed in vitro decay. However, the rapid decay was also observed when we filtered or chloroform-extracted the phages before the decay quantification (data not shown), suggesting that decay may be an intrinsic property of our ancestral phage and not exclusively related to debris attachment. Although differences between our ancestral strain and the strains used in other studies could partially account for the discrepancy in decay rates, there may also be physiologically relevant explanations. Populations of phage λ have been reported to show a biphasic decay pattern, with a rapid decay phase immediately after bursting and a slower phase thereafter [70]. If the same applies to ΦX174, it could explain the discrepancy between our results and previous decay measurements. These earlier studies may have missed the unstable fraction and primarily captured the more stable phage fraction. Further work will be needed to clarify this point.

In contrast to the 3-h regime, the shorter transfers of the 30-min regime do not impose strong selection on persistence but instead select for drastically reduced adsorption (**Fig. 3A**). The low-adsorption mutants evolved in the 30-min regime carried an F(T100A) substitution in the major capsid protein, located in an α-helix previously implicated in host binding and specificity or temperature adaptation [23, 25, 27, 39, 71, 72]. A decrease in the adsorption constant may seem counterintuitive because it reduces the rate at which infections can be initiated. However, the selection for slow adsorption can be understood as an adaptation to the specific timing of the bottlenecks in the 30-min transfer regimes. These bottlenecks favour strategies for maintaining a high density of free phages at the time of the bottleneck (transfer in our case). One such strategy is to reduce losses to adsorption during periods when most host cells are already infected. More broadly, our simulations indicate that selection on adsorption is shaped by the relative strength of two within-transfer phases imposing opposing selection pressures: a productive phase, in which uninfected cells are abundant and fast adsorption is favoured, and an unproductive sink phase, in which most cells are already infected and adsorption primarily

removes phage particles via attachment to infected cells (**Fig. 6D**). Importantly, this unproductive sink may be stronger in vitro than in our model, because additional processes not explicitly included in the model—such as adsorption to cell debris or other non-productive binding interactions [50, 73, 74]—could further increase phage losses during the unproductive phase. Modulating the relative strengths of these phases (for example, by changing the dilution factor between transfers) could, in principle, allow the selection of phages with different adsorption constants (**Fig. 6E**).

Even though the complex interplay between bottlenecks and the success of adsorption rate mutants has been shown theoretically before [75], it is not usually observed in phenotype-fitness maps of bacteria-phage interactions [76]. This is because the unexpected benefit of a reduced adsorption rate in the 30-min regime requires taking into account (i) exhaustion of host cells, (ii) unproductive adsorptions on already infected cells, and (iii) frequent population bottlenecks on the order of the lysis time of the phage. A recent theoretical study showed a similar effect resulting from the need to balance productive and unproductive phases of phage growth with respect to lysis time evolution [77].

The contrasting effects of different adsorption constants across transfer regimes suggest ecological trade-offs that may drive divergent selection in natural populations. For instance, sewage water, a major source of phages, contains both a liquid phase where bacteria grow pelagically and a semi-solid phase where bacteria form biofilms [78, 79]. High adsorption may be favoured in the liquid phase to exploit transient host availability. In contrast, slow adsorption may be advantageous in biofilms or host-rich environments to avoid attachment to already infected cells [17]. Slow adsorption could also be beneficial in particle-rich environments, where phages risk irreversible binding to inert material [20, 73, 80]. Recurrent transitions between these environments may have shaped the evolutionary plasticity of adsorption in ΦX174, where single-point mutations in the F protein can lead to changes in adsorption with profound fitness consequences in different environments.

Plaque phenotypes mirrored the contrasting selective pressures in our two regimes, with small- and large-plaque-forming phages evolving under the 3-h and 30-min regimes, respectively. In theory, plaque size reduction can result from changes in the three different life history traits: (1) extended lysis time, (2) reduction in burst size, or (3) increased (or extremely low) adsorption constant [1, 22, 48, 49, 81, 82]. In our study, the differences in plaque size

may reflect adsorption differences, as we found an inverse correlation between adsorption constant and plaque size (**Fig. 3**). This pattern is consistent with previous findings in phage λ, where mutations in the host recognition protein J that increased the adsorption constant led to reduced plaque size [17, 22]. Similar observations were also made in phage U136B, where fast adsorbers often formed smaller plaques [83]. A likely explanation for this is that rapid adsorption promotes local reinfection, trapping phages at the focus of infection and preventing them from reaching uninfected regions of the bacterial lawn [22, 81].

The findings presented in this manuscript have potential practical relevance. Phages with different life-history traits may be more or less suited for treating different types of infections, such as biofilm or bloodstream infections. Our results illustrate how experimental evolution can uncover key environmental trade-offs, which may inform the selection or engineering of phages for therapy. In addition, understanding how phage life-history traits evolve under different selective pressures is essential for anticipating phage evolutionary dynamics once they are administered to patients. Thus, experimental evolution serves not only as a valuable research approach but also as a practical tool to guide the development of safer, more effective, and more predictable phage-based therapies.

Acknowledgements

We acknowledge the Sequencing Service Team from the Max Planck Institute for Evolutionary Biology in Plön for performing the Sanger Sequencing. We thank Daniel Martens and Anja Baade for their support regarding lab logistics as well as the entire Microbial Population Biology Department for their support. We also want to thank David Rogers for helping with PCR troubleshooting and Eva Lievens and Stephen T. Abedon for the insightful discussions on viral life history evolution. MR, CB, and ORM were supported by the International Max-Planck Research School for Evolutionary Biology (IMPRS EvolBio). Finally, we would like to thank our reviews and editor for their helpful comments.

Funding

The funding for this study was received from the Max Planck Society. Work in JLG lab was also supported by ERC starting grant 853323.

Data availability

Raw sequencing reads used to generate figures **Fig. 2** and **Fig. S4**, plaque assay images used for plaque morphology image analysis and to generate figures **Fig. 1**, **Fig. 2**, **Fig. 3B**, **Fig. S4**, **Fig. S5,** and **Fig. S11** and raw bacteria and phage counting data and measurements of plaque area and turbidity used to generate figures **Fig. 3, Fig. 4B, Fig. 5C, Fig. S3, Fig. S5, Fig. S6, Fig. S7, Fig. S9, and Fig. S10** are found in Zenodo (doi: 10.5281/zenodo.17076220).

The analysis script used for analysing and visualising the raw counting and image analysis data is in https://github.com/OReyesMatte/Reuter_2025/tree/main/Data_analysis.

Analysis information for the simulations used to generate figures **Fig. 4C, Fig. 5A, Fig. 5B, Fig. 6, Fig. S2, Fig. S8** is deposited in the repository found in https://github.com/OReyesMatte/Reuter_2025/tree/main/Simulations.

A list of processed plaque assay images for which plaque area and turbidity are found in https://github.com/OReyesMatte/Reuter_2025/tree/main/Plaque_analysis.

Supplementary material

All supplementary material is found in the file "**Supplementary.pdf**". The file contains supplementary figures **Fig. S1-S11**, supplementary tables **Table S1-S13**, and **Supplementary Methods**.

Conflicts of interest

The authors declare no conflict of interest.

Authors Contributions

Conceptualisation: MR, MS, JLG, FB

Methodology: MR, MS, ORM

Investigation: MR, CV, CB, JRD

Data Curation: MR, ORM

Formal analysis: MR, ORM, MS

Validation: MR, MS

Visualisation: MR, MS

Writing- Original Draft/Review & Editing: MR, JLG, MS

Supervision: JLG, FB

Author ORCIDs

Manuela Reuter: https://orcid.org/0009-0000-5191-3414

Michael Sieber: https://orcid.org/0000-0003-2703-3665

Octavio Reyes-Matte: https://orcid.org/0009-0006-7031-6213

Christopher Böhmker: https://orcid.org/0009-0003-0098-9816

Jordan Romeyer Dherbey: https://orcid.org/0000-0002-6125-1722

Frederic Bertels: https://orcid.org/0000-0001-6222-4139

Javier Lopez-Garrido: https://orcid.org/0000-0002-8907-502X

1 **Tables**

2 **Table 1: Parameters and corresponding parameter values used in the mathematical model.**

Parameter	Value
Adsorption constant, ancestor Φ_A	$4*10^{-9}$ (ml^{-1} min^{-1})
Adsorption constant, large-plaque type evolved in vitro in the 30-min regime Φ_L	$2*10^{-9}$ (ml^{-1} min^{-1})
Adsorption constant, small-plaque type evolved in vitro in the 3-h regime Φ_S	$6*10^{-9}$ (ml^{-1} min^{-1})
Burst size, all types β	200 (per infected host cell)
Lysis time, all types τ	17 min
Decay rate, ancestor δ	$1.7*10^{-2}$ (min^{-1})
Decay rate, ancestor and large-plaque type δ_L	$1.4*10^{-2}$ (min^{-1})
Decay rate, small-plaque type δ_S	$6*10^{-3}$ (min^{-1})

Figures

Fig. 1: 30-min and 3-h transfer regimes select for large- and small-plaque-forming phages, respectively. Phage ΦX174 was propagated under 30-min (top) or 3-h (bottom) regimes, leading to the evolution of large- and small-plaque-forming phages, respectively. Ancestral plaques are shown on the left, and evolved large (top) and small (bottom) plaques on the right. Scale bars, 1 cm. All plaque images have the same magnification. Diagrams of the 30-min and 3-h regimes are in the middle. The approximate input multiplicity of infection (MOI$_{input}$) and incubation times are the same for each transfer. For illustration purposes, only four transfers are shown, but the experiment involved more transfers. For the 30-min regime, 20 transfers (fifteen 30-min transfers, interrupted by a 3-h transfer after every fourth transfer) were done before large-plaque mutants were observed in plaque assays. In the 3-h regime, transfers were done until heritable small plaques emerged, which occurred after four to seven transfers. See methods and **Fig. S1** for details about the individual transfer protocols. Figure created in BioRender. Reuter, M. (2025) https://BioRender.com/q39e4in.

Fig. 2: Mutations linked to large- and small-plaque phenotypes occur in the major capsid protein. (A)
Large-plaque phenotypes (top) and corresponding genotypes (bottom) of mutants isolated from independent lines
of the 30-min regime. All the mutants carried an A-to-G transition at position 1301 (A1301G) in the phage
genome, resulting in a threonine-to-alanine substitution at position 101 of the F protein [F(T100A)]. Four isolates
carried an additional mutation, either in the F (green) or the H (blue) gene. The ancestor is shown in grey (WT).
(B) Small-plaque phenotypes (top) and genotypes (bottom) of mutants isolated from independent lines of the 3-h
regime. All mutants differed from the ancestor by different mutation in the F gene (green), resulting in different
amino acid changes: G321D in the clones from lines 1, 2, and 5; M212I in the clone from line 3; and the formation
of a premature stop codon at the very last codon of the F gene (S426*) in the clone from line 4. The nucleotide
change in the phage genome is shown at the top and the corresponding amino acid change at the bottom. The
ancestor is shown in grey (WT). Scale bars, 1 cm. Genome illustrations were generated using Geneious Prime
2023.0.1. Figure created in BioRender. Reuter, M. (2025) https://BioRender.com/dck2411.

Fig. 3: The adsorption constant is inversely correlated with plaque size. (A) Adsorption constant "k" of the

ancestor (grey), the large-plaque mutant F(T100A) (purple) evolved from the 30-min regime and the two small-

plaque mutants F(G321D) (orange) and F(S426*) isolated from the 3-h regime (green). At least three independent

experiments were performed for each genotype. Each black dot represents the adsorption constant of an individual

experiment, the horizontal lines indicate the median, and the boxes span the middle 50% of the data. The whiskers

span the lower and upper 25% of the data. Different letters on top of the boxes indicate significant adsorption

constant differences between genotypes (significance level $p<0.05$). (B) Lysis plaque area in cm^2 (y-axis)

measured in vitro for the ancestor (grey, n=76), and the evolved phages F(T100A) (purple, n=60, 30-min regime),

F(G321D) (green, n=303, 3-h regime) and F(S426*) (orange, n=431, 3-h regime). Violin plots show distribution

shape; boxplots display medians (black lines), interquartile range (boxes), and upper/lower 25% (whiskers). See

Methods and **Supplementary Methods** for details of adsorption and plaque phenotype analysis.

Fig. 4: Single-point mutations in the major capsid protein modulate adsorption and persistence. (A)
Location of point mutations linked to plaque size differences in the F protein monomer (left) or multimer (right)
structure. Mutations F(G321D) and F(S426*) from 3-h regime isolates leading to small plaque phenotypes are
shown in green and orange, respectively. The 30-min regime mutation F(T100A) leading to a large plaque
phenotype is shown in purple. Mutations associated with the mall plaque phenotype are localised at monomer-
monomer interfaces, whereas mutation F(T100A) is located at the F protein centre. The protein structure, resolved
by McKenna *et al.* [32], was taken from RCSB PDB (reference 2BPA:
https://www.wwpdb.org/pdb?id=pdb_00002bpa). The figure was prepared in UCSF Chimera (version 1.18). (B)
In vitro measured decay of infectious phage particles of the ancestor (grey), large-plaque mutant T100A (purple),
and small-plaque mutants G321D (green) and S426* (orange). The decay is visualised as the remaining fraction
of infectious particles (y-axis) relative to the maximum phage titer, reached 60 minutes (ancestor and F(T100A))
or 120 min (F(G321D) and F(S426*)) post-infection initiation. The loss of infectious particles is shown for a five-
hour infection period (x-axis). The full phage population dynamics, including the productive infection period,
used to calculate the infection phage particle fraction, are shown in **Fig. S6**. See **Supplementary Methods** for
model selection and parameter extraction. (C) Simulations of free phage population dynamics (y-axis) across
transfers (x-axis) for the 3-h regime, incorporating differences in adsorption constants (top) or decay rates
(bottom) according to values shown in Table 1 for the ancestor (grey), 30-min mutant T100A (purple), and 3-h
mutant G321D (green). The simulations started with a genotype frequency of 99% of the ancestral phages and
0.5% of each mutant phage. The graphs show the predicted frequency of each type at the end of each transfer.

Fig. 5: Simulations and in-vitro quantification of phage populations over single 30-min transfers. (A)
Predicted optimal adsorption constants based on simulated competitions between the ancestor and hypothetical
adsorption constant mutants (x-axis) over a single 30-min transfer, starting from an initial 1:1 ratio. The y-axis
shows the predicted mutant-to-ancestor titer ratios after one transfer. The adsorption constant of the F(T100A)
mutant evolved in the 30-min regime (purple line) is close to the predicted optimum. (B) Simulation of free phage
densities (y-axis) within a single 30-min transfer (x-axis) for the ancestor (grey) and the slow-adsorbing mutant
F(T100A) (purple). The free phage dynamics within one transfer predict a higher number of free phages of the
slow-adsorbing mutant at the end of a transfer. The 30-min transfer is separated into a first and second adsorption
period, interrupted by a burst phase, based on the in vitro determined lysis time of 17 minutes (**Fig. S3**). (C) In-
vitro measured free phages of the slow-adsorbing F(T100A) mutant (purple bars) relative to ancestral free phages
(y-axis) after consecutive 30-min transfers (x-axis). The number of free phages of mutant F(T100A) was
normalised to the free phages of the ancestor (dashed black line). Data represent the average and standard error
of three independent biological replicates. Individual titer measurements for mutant F(T100A) and the ancestral
phage after each 30-min transfer are shown in **Fig. S7**.

Fig. 6: The relative strength of productive and unproductive adsorptions modulates the optimal adsorption constant in serial transfers. (A) Simulation of phage (top) and bacterial (bottom) densities (y-axis) over six 30-min transfers (x-axis) in which 2% of the bacteria-phage co-culture of the previous transfer is propagated to fresh bacteria every 30 minutes. The densities of not infected (blue) and infected (brown) bacterial cells are shown. The simulated phage and bacteria dynamics reach a stable within-transfer pattern that then repeats transfer after transfer. (B) and (C) Density (y-axis) of free phage (B) and non-infected (blue) and infected (brown) host cells (C) over the stable 30-minute transfer pattern that emerges after a few transient transfers. The pattern is characterized by overlapping infection cycles that produce three phage burst events: a major burst after ~17 min originating from phages that infected host cells at the beginning of the current transfer (first infection cycle), another after ~4 min originating from the lysis of virocells infected in the previous transfer (2nd infection cycle), and another after ~21 min originating from infections from phages that infected after the first burst (3rd infection cycle). (D) Productive (blue) and unproductive (brown) adsorptions per minute (y-axis) during a stable 30-minute co-culturing period (x-axis) for regimes with different dilution factors between transfers: 50-fold (left), 500-fold (middle), and 2000-fold (right). See also **Fig. S8**. (E) Adsorption fitness landscape across a range of transfer times (x-axis) and transfer dilution factors (y-axis). Phages with high (green), intermediate (blue) or low (purple) adsorption constants were competed in silico under transfer regimes with different co-culturing times and dilution factors between transfers. Competitions were initiated with equal frequencies of the three phages. The colour corresponding to the type fixed after 100 transfers is indicated in the plot.

References

1. Abedon ST, Hyman P, Thomas C. Experimental examination of bacteriophage latent-period evolution as a response to bacterial availability. *Appl Environ Microbiol* 2003;**69**:7499–7506. https://doi.org/10.1128/AEM.69.12.7499-7506.2003

2. Eshelman CM et al. Unrestricted migration favours virulent pathogens in experimental metapopulations: Evolutionary genetics of a rapacious life history. *Philosophical Transactions of the Royal Society B: Biological Sciences* 2010;**365**:2503–2513. https://doi.org/10.1098/rstb.2010.0066

3. Dessau M et al. Selective pressure causes an RNA virus to trade reproductive fitness for increased structural and thermal stability of a viral enzyme. *PLoS Genet* 2012;**8**:e1003102. https://doi.org/10.1371/journal.pgen.1003102

4. Heineman RH, Bull JJ. Testing optimality with experimental evolution: Lysis time in a bacteriophage. *Evolution (N Y)* 2007;**61**:1695–1709. https://doi.org/10.1111/j.1558-5646.2007.00132.x

5. DeLong JP et al. Towards an integrative view of virus phenotypes. *Nat Rev Microbiol* 2022;**20**:83–94. https://doi.org/10.1038/s41579-021-00612-w

6. Abedon ST. Impact of phage properties on bacterial survival. In: Adams HT (ed.), Contemporary Trends in Bacteriophage Research. Nova Science Publishers, Inc., 2009, 217–235.

7. Ellis EL, Delbrück M. The growth of bacteriophage. *J Gen Physiol* 1939;**22**:365–384. https://doi.org/10.1085/jgp.22.3.365

8. Delbrück M. Adsorption of bacteriophage under various physiological conditions of the host. *J Gen Physiol* 1940;**23**:631–642. https://doi.org/10.1085/jgp.23.5.631

9. Delbrück M. The growth of bacteriophage and lysis of the host. *J Gen Physiol* 1940;**23**:643–660. https://doi.org/10.1085/jgp.23.5.643

10. Suttle CA, Chen F. Mechanisms and rates of decay of marine viruses in seawater. *Appl Environ Microbiol* 1992;**58**:3721–3729. https://doi.org/10.1128/aem.58.11.3721-3729.1992

11. Abedon ST. Phage Ecology. In: Calendar R, Abedon ST (eds.), The bacteriophages. Oxford: Oxford University Press, 2006, 37–46.

12. Adams MH, Wassermann FE. Frequency distribution of phage release in the one-step growth experiment. *Virology* 1956;**2**:96–108. https://doi.org/10.1016/0042-6822(56)90079-4

13. Abedon ST. Kinetics of phage-mediated biocontrol of bacteria. *Foodborne Pathog Dis* 2009;**6**:807–815. https://doi.org/10.1089/fpd.2008.0242

14. Lindberg HM, McKean KA, Wang I-N. Phage fitness may help predict phage therapy efficacy. *Bacteriophage* 2014;**4**:e964081. https://doi.org/10.4161/21597073.2014.964081

15. Abedon S. Phage therapy pharmacology: Calculating phage dosing. In: Laskin A, Gadd G, Sariaslani S (eds.), Advances in Applied Microbiology. Academic Press Inc., 2011, 1–40.

16. Kowalska JD et al. Growing trend of fighting infections in aquaculture environment—opportunities and challenges of phage therapy. *Antibiotics* 2020;**9**:301. https://doi.org/10.3390/antibiotics9060301

17. Gallet R, Shao Y, Wang IN. High adsorption rate is detrimental to bacteriophage fitness in a biofilm-like environment. *BMC Evol Biol* 2009;**9**:241. https://doi.org/10.1186/1471-2148-9-241

18. Heineman RH, Brown SP. Experimental evolution of a bacteriophage virus reveals the trajectory of adaptation across a fecundity/longevity trade-off. *PLoS One* 2012;**7**:e46322. https://doi.org/10.1371/journal.pone.0046322

19. Mäkelä K, Laanto E, Sundberg LR. Determinants in the phage life cycle: The dynamic nature of ssDNA phage FLiP and host interactions under varying environmental conditions and growth phases. *Environ Microbiol* 2024;**26**:e16670. https://doi.org/10.1111/1462-2920.16670

20. Daniels LL, Wais AC. Virulence in phage populations infecting *Halobacterium cutirubrum*. *FEMS Microbiol Ecol* 1998;**25**:129–134. https://doi.org/10.1016/S0168-6496(97)00089-5

21. Shao Y, Wang IN. Bacteriophage adsorption rate and optimal lysis time. *Genetics* 2008;**180**:471–482. https://doi.org/10.1534/genetics.108.090100

22. Gallet R, Kannoly S, Wang IN. Effects of bacteriophage traits on plaque formation. *BMC Microbiol* 2011;**11**:181. https://doi.org/10.1186/1471-2180-11-181

23. Bull JJ et al. Exceptional convergent evolution in a virus. *Genetics* 1997;**147**:1497–1507. https://doi.org/https://doi.org/10.1093/genetics/147.4.1497

24. Wichman HA et al. Different trajectories of parallel evolution during viral adaptation. *Science (1979)* 1999;**285**:422–424. https://doi.org/10.1126/science.285.5426.422

25. Crill WD, Wichman HA, Bull JJ. Evolutionary reversals during viral adaptation to alternating hosts. *Genetics* 2000;**154**:27–37. https://doi.org/10.1093/genetics/154.1.27

26. Wichman HA, Brown CJ. Experimental evolution of viruses: *Microviridae* as a model system. *Philosophical Transactions of the Royal Society B: Biological Sciences* 2010;**365**:2495–2501. https://doi.org/10.1098/rstb.2010.0053

27. Wichman HA, Millstein J, Bull JJ. Adaptive molecular evolution for 13,000 phage generations: A possible arms race. *Genetics* 2005;**170**:19–31. https://doi.org/10.1534/genetics.104.034488

28. Sanger F et al. The nucleotide sequence of bacteriophage φX174. *J Mol Biol*
1978;**125**:225–246. https://doi.org/10.1016/0022-2836(78)90346-7

29. Sinsheimer RL. A single-stranded deoxyribonucleic acid from bacteriophage φX174. *J*
Mol Biol 1959;**1**:43–53. https://doi.org/10.1016/S0022-2836(59)80006-1

30. Fane B. ΦX174 et al. the *Microviridae*. In: Calendar R (ed.), The Bacteriophages.
Oxford University Press, Inc., 2006, 129–145.

31. Baas PD. DNA replication of single-stranded *Escherichia coli* DNA phages. *Biochim*
Biophys Acta 1985;**825**:111–139. https://doi.org/10.1016/0167-4781(85)90096-X

32. McKenna R et al. Atomic structure of single-stranded DNA bacteriophage ΦX174 and
its functional implications. *Nature* 1992;**355**:137–143.
https://doi.org/10.1038/355137a0

33. Sun L et al. Icosahedral bacteriophage Φx174 forms a tail for DNA transport during
infection. *Nature* 2014;**505**:432–435. https://doi.org/10.1038/nature12816

34. Burgess AB. Studies on the proteins of ΦX174 II. The protein composition of the ΦX
coat. *Proceedings of the National Academy of Sciences* 1969;**64**:613–617.
https://doi.org/10.1073/pnas.64.2.613

35. Incardona NL, Selvidge L. Mechanism of adsorption and eclipse of bacteriophage
ΦX174 II. Attachment and eclipse with isolated *Escherichia coli* cell wall
lipopolysaccharide. *J Virol* 1973;**11**:775–782. https://doi.org/10.1128/jvi.11.5.775-
782.1973

36. Incardona NL, Tuech JK, Murti G. Irreversible binding of phage ΦX174 to cell-bound
lipopolysaccharide receptors and release of virus-receptor complexes. *Biochemistry*
1985;**24**:6439–6446.

37. Sun Y et al. Structural changes of tailless bacteriophage ΦX174 during penetration of
bacterial cell walls. *Proceedings of the National Academy of Sciences*
2017;**114**:13708–13713. https://doi.org/10.1073/pnas.1716614114

38. Ilag LL et al. Calcium ion-induced structural changes in bacteriophage ΦX174.
Journal of Molecular Biologie 1994;**244**:291–300.
https://doi.org/10.1006/jmbi.1994.1730

39. Romeyer Dherbey J et al. Stepwise evolution of *E. coli* C and φX174 reveals
unexpected lipopolysaccharide (LPS) diversity. *Mol Biol Evol* 2023;**40**:msad154.
https://doi.org/10.1093/molbev/msad154

40. Biggs KRH et al. Ecological approach to understanding superinfection inhibition in
bacteriophage. *Viruses* 2021;**13**:1389. https://doi.org/10.3390/v13071389

41. Hirel Ph-H et al. Extent of N-terminal methionine excision from *Escherichia coli*
proteins is governed by the side-chain length of the penultimate amino acid.
Proceedings of the National Academy of Sciences 1989;**86**:8247–8251.

42. Hyman P, Abedon ST. Practical methods for determining phage growth parameters. In: Clokie MRJ, Kropinski AM (eds.), Methods in molecular biology, 1st ed. Totowa, NJ: Humana Press, 2009, 175–202.

43. Berg S et al. Ilastik: interactive machine learning for (bio)image analysis. *Nat Methods* 2019;**16**:1226–1232. https://doi.org/10.1038/s41592-019-0582-9

44. Virtanen P et al. SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat Methods* 2020;**17**:261–272. https://doi.org/10.1038/s41592-019-0686-2

45. Van Der Walt S et al. scikit-image: Image processing in Python. *PeerJ* 2014;**2**:e453. https://doi.org/10.7717/peerj.453

46. R Core Team. A language and environment for statistical computing. R Foundation for Statistical Computing. 2022. Vienna, Austria, 2022.

47. Abedon ST. Detection of Bacteriophages: Phage Plaques. In: Harper DR et al. (eds.), Bacteriophages: biology, technology, therapy. Springer Cham, 2021, 507–538.

48. Lee Y, Yin J. Detection of evolving viruses. *Nat Biotechnol* 1996;**14**:491–493. https://doi.org/10.1038/nbt0496-491

49. Abedon ST, Yin J. Bacteriophage plaques: theory and analysis. In: Clokie MRJ, Kropinski AM (eds.), Bacteriophages: Methods and Protocols, Volume 1: Isolation, Characterization, and Interactions. Totowa, NJ: Humana Press, 2009, 161–174.

50. Fujimura R, Kaesberg P. The adsorption of bacteriophage φX174 to its host. *Biophys J* 1962;**2**:433–449. https://doi.org/10.1016/S0006-3495(62)86866-0

51. Bayer ME, Starkey TW. The adsorption of bacteriophage ΦX174 and its interaction with *Escherichia coli*; a kinetic and morphological study. *Virology* 1972;**49**:236–256. https://doi.org/10.1016/S0042-6822(72)80026-6

52. Lee KH et al. First-step mutations for adaptation at elevated temperature increase capsid stability in a virus. *PLoS One* 2011;**6**:e25640. https://doi.org/10.1371/journal.pone.0025640

53. Tom EF et al. Experimental evolution of UV resistance in a phage. *PeerJ* 2018;**6**:e5190. https://doi.org/10.7717/peerj.5190

54. Heineman RH, Bull JJ, Molineux IJ. Layers of evolvability in a bacteriophage life history trait. *Mol Biol Evol* 2009;**26**:1289–1298. https://doi.org/10.1093/molbev/msp037

55. Pepin KM, Wichman HA. Experimental evolution and genome sequencing reveal variation in levels of clonal interference in large populations of bacteriophage φX174. *BMC Evol Biol* 2008;**8**. https://doi.org/10.1186/1471-2148-8-85

56. Rokyta DR, Abdo Z, Wichman HA. The genetics of adaptation for eight microvirid bacteriophages. *J Mol Evol* 2009;**69**:229–239. https://doi.org/10.1007/s00239-009-9267-9

57. Coberly LC et al. Space, time, and host evolution facilitate coexistence of competing bacteriophages: Theory and experiment. *American Naturalist* 2009;**173**:E121–E138. https://doi.org/10.1086/597226

58. Hall JPJ, Harrison E, Brockhurst MA. Viral host-adaptation: Insights from evolution experiments with phages. *Curr Opin Virol* 2013;**3**:572–577. https://doi.org/10.1016/j.coviro.2013.07.001

59. Domingo-Calap P, Sanjuán R. Experimental evolution of RNA versus DNA viruses. *Evolution (N Y)* 2011;**65**:2987–2994. https://doi.org/10.1111/j.1558-5646.2011.01339.x

60. Rokyta D et al. Experimental Genomic Evolution: Extensive Compensation for Loss of DNA Ligase Activity in a Virus. *Mol Biol Evol* 2002;**19**:230–238. https://doi.org/10.1093/oxfordjournals.molbev.a004076

61. Heineman RH, Molineux IJ, Bull JJ. Evolutionary Robustness of an Optimal Phenotype: Re-evolution of Lysis in a Bacteriophage Deleted for Its Lysin Gene. *J Mol Evol* 2005;**61**:181–191. https://doi.org/10.1007/s00239-004-0304-4

62. Abedon ST, Herschler TD, Stopar D. Bacteriophage latent-period evolution as a response to resource availability. *Appl Environ Microbiol* 2001;**67**:4233–4241. https://doi.org/10.1128/AEM.67.9.4233-4241.2001

63. Kerr B et al. Local migration promotes competitive restraint in a host-pathogen 'tragedy of the commons'. *Nature* 2006;**442**:75–78. https://doi.org/10.1038/nature04864

64. Nabergoj D, Modic P, Podgornik A. Effect of bacterial growth rate on bacteriophage population growth rate. *Microbiologyopen* 2018;**7**:e558. https://doi.org/10.1002/mbo3.558

65. Wang IN. Lysis timing and bacteriophage fitness. *Genetics* 2006;**172**:17–26. https://doi.org/10.1534/genetics.105.045922

66. Josslin R. The Lysis Mechanism of Phage T4: Mutants Affecting Lysis'. *Virology* 1970;**40**:719–726. https://doi.org/10.1016/0042-6822(70)90216-3

67. Abedon ST. Selection for bacteriophage latent period length by bacterial density: a theoretical examination. *Microb Ecol* 1989;**18**:79–88. https://doi.org/10.1007/BF02030117

68. Bull JJ, Badgett MR, Wichman HA. Big-benefit mutations in a bacteriophage inhibited with heat. *Mol Biol Evol* 2000;**17**:942–950. https://doi.org/10.1093/oxfordjournals.molbev.a026375

69. De Paepe M, Taddei F. Viruses' life history: Towards a mechanistic basis of a trade-off between survival and reproduction among phages. *PLoS Biol* 2006;**4**:e193. https://doi.org/10.1371/journal.pbio.0040193

70. Seurat J et al. Design, optimization, and inference of biphasic decay of infectious virus particles. *J Theor Biol* 2025;**600**:112042. https://doi.org/10.1016/j.jtbi.2025.112042

71. Wichman HA et al. Experimental evolution recapitulates natural evolution.
Philosophical Transactions of the Royal Society B: Biological Sciences
2000;**355**:1677–1684. https://doi.org/10.1098/rstb.2000.0731

72. Pepin KM, Domsic J, McKenna R. Genomic evolution in a virus under specific
selection for host recognition. *Infection, Genetics and Evolution* 2008;**8**:825–834.
https://doi.org/10.1016/j.meegid.2008.08.008

73. Harvey RW, Ryan JN. Use of PRD1 bacteriophage in groundwater viral transport,
inactivation, and attachment studies. *FEMS Microbiol Ecol* 2004;**49**:3–16.
https://doi.org/10.1016/j.femsec.2003.09.015

74. Rabinovitch A, Aviram I, Zaritsky A. Bacterial debris - An ecological mechanism for
coexistence of bacteria and their viruses. *J Theor Biol* 2003;**224**:377–383.
https://doi.org/10.1016/S0022-5193(03)00174-7

75. Patwa Z, Wahl LM. Fixation probability for lytic viruses: The attachment-lysis model.
Genetics 2008;**180**:459–470. https://doi.org/10.1534/genetics.108.090555

76. Bull JJ, Heineman RH, Wilke CO. The phenotype-fitness map in experimental
evolution of phages. *PLoS One* 2011;**6**:e27796.
https://doi.org/10.1371/journal.pone.0027796

77. Smith A et al. Live and let die: lysis time variability and resource limitation shape lytic
bacteriophage fitness. *bioRxiv* 2025. https://doi.org/10.1101/2025.07.21.665848

78. Jensen H, Biggs CA, Karunakaran E. The importance of sewer biofilms. *Wiley
Interdisciplinary Reviews: Water* 2016;**3**:487–494. https://doi.org/10.1002/wat2.1144

79. Balleste E et al. Bacteriophages in sewage: abundance, roles, and applications. *FEMS
microbes* 2022;**3**:xtac009. https://doi.org/10.1093/femsmc/xtac009

80. Richter Ł et al. Adsorption of bacteriophages on polypropylene labware affects the
reproducibility of phage research. *Sci Rep* 2021;**11**:7387.
https://doi.org/10.1038/s41598-021-86571-x

81. Yin J, McCaskill JS. Replication of viruses in a growing plaque: a reaction-diffusion
model. *Biophys J* 1992;**61**:1540–1549. https://doi.org/10.1016/S0006-3495(92)81958-
6

82. Yin J. Evolution of Bacteriophage T7 in a Growing Plaque. *J Bacteriol*
1993;**175**:1272–1277. https://doi.org/10.1128/jb.175.5.1272-1277.1993

83. Burmeister AR et al. Experimental evolution of the TolC-receptor phage U136B
functionally identifies a tail fiber protein involved in adsorption through strong parallel
adaptation. *Appl Environ Microbiol* 2023;**89**:e0007923.
https://doi.org/10.1128/aem.00079-23