Cryo-EM structure of a methanogen nitrogenase-PII protein supercomplex

Rajnandani Kashyap[1], Thomas M. Deere[2], Ahmed Dhamad[2#], Melissa Chanderban[2], Monika Tokmina-Lukaszewska[3], Brian Bothner[3], Daniel J. Lessner[2,*], and Edwin Antony[1,*]

Affiliations:

[1]Department of Biochemistry and Molecular Biology, St. Louis University School of Medicine, St. Louis, MO, USA

[2]Department of Biological Sciences, University of Arkansas, Fayetteville, AR, USA

[3]Department of Chemistry and Biochemistry, Montana State University, Bozeman, MT, USA

#Current address: Department of Biological Sciences, Wasit University, Wasit, Iraq

*Corresponding authors. Email: edwin.antony@health.slu.edu; dlessner@uark.edu

Abstract: Nitrogenases are metalloenzymes that catalyze the reduction of atmospheric dinitrogen to ammonia, sustaining the global nitrogen cycle. While bacterial nitrogenase has been extensively characterized, the architecture and regulation of archaeal nitrogenases remain unknown despite longstanding evidence of nitrogen fixation in methanogens. Here we report a 3.1 Å cryo-electron microscopy structure of a native nitrogenase–PII protein supercomplex from *Methanosarcina acetivorans*. The structure reveals an unprecedented assembly of three NifDK heterotetramers bridged by six NifI$_{1,2}$ heterotrimeric PII complexes, which sterically block NifH association and lock the enzyme in an inactive state. The NifI complexes display asymmetric binding of ADP and 2-oxoglutarate, coupling nitrogenase inhibition directly to cellular energy and nitrogen status. Addition of 2-oxoglutarate and ATP releases the NifI complexes, stimulating a threefold increase in NifDK activity in vitro. This higher-order architecture uncovers a previously unrecognized regulatory strategy in methanogens, in which PII proteins drive nitrogenase oligomerization to control activity. The discovery that nitrogenase activity may be modulated through direct assembly into higher-order structures opens new avenues for exploring nitrogenase evolution, regulation, and biotechnological applications.

One sentence summary: Discovery of a nitrogenase-PII protein supercomplex in methanogens, uncovering a metabolite-gated assembly mechanism for nitrogenase inhibition.

Introduction.

Biological nitrogen fixation (diazotrophy) is a critical process in the global nitrogen cycle, yet it is confined to a limited group of bacteria and archaea. All known and predicted diazotrophs rely on molybdenum (Mo)-dependent nitrogenase to catalyze the energetically demanding reduction of atmospheric dinitrogen (N_2) to ammonia (NH_3), following the stoichiometry[1,2]:

$$\textbf{N}_2 + \textbf{8H}^+ + \textbf{8e}^- + \textbf{16ATP} \rightarrow \textbf{2NH}_3 + \textbf{H}_2 + \textbf{16ADP} + \textbf{16P}_i.$$

Mo-nitrogenase is a highly oxygen-sensitive metalloenzyme composed of two components: dinitrogenase (a NifD-NifK heterotetramer, also known as the MoFe protein) and dinitrogenase reductase (a NifH homodimer, or Fe protein). NifH contains a [4Fe-4S] cluster that transfers electrons to the P-cluster (8Fe-7S) of NifDK, which in turn delivers electrons to the FeMo-cofactor (FeMo-co or M-cluster), the active site for N_2 reduction[3-5]. Transfer of one electron

by NifH requires the hydrolysis of two ATP molecules[2,6,7]. Given the high energy cost, nitrogenase expression and activity are tightly regulated at multiple levels—transcriptional, translational, and post-translational—and are closely integrated with cellular carbon metabolism and energy status[8-10].

PII signal transduction proteins are central regulators of nitrogen fixation and assimilation in diazotrophs[11]. These ancient and highly conserved proteins are found across all domains of life[12]. Canonical PII proteins (e.g., GlnK, GlnB, GlnZ) form homotrimers, each with three ligand-binding sites located at the interface between subunits. Each monomer adopts a ferredoxin-like α/β fold with three characteristic loops (T, B, and C). Ligand-induced conformational changes in the T-loop are key to modulating interactions with target proteins. PII proteins sense the cellular carbon and nitrogen status by binding 2-oxoglutarate (2OG), a TCA cycle intermediate and universal indicator of nitrogen limitation[13]. They also bind ADP and ATP to monitor the cell's energy state. These ligand interactions influence PII binding to various targets, including enzymes, transporters, and regulatory proteins[14].

PII proteins are known to indirectly control the activity of nitrogenase in diverse bacteria, including anaerobes, aerobes, and oxygenic phototrophs. The best example is the DraT-DraG system, where under nitrogen-replete conditions the PII protein GlnB activates DraT, which reversibly inactivates NifH via ADP-ribosylation, thereby preventing association with NifDK. This modification is triggered by low intracellular 2OG, signaling nitrogen sufficiency[11].

In contrast, nitrogenase in archaea is restricted to anaerobic methanogens and methanotrophs. Methanogens, among the most ancient lineages of life, are thought to be the progenitors of nitrogenase [15,16]. They typically possess a minimal *nif* gene cluster (*nifHI₁I₂DKEN*), encoding the catalytic components (NifHDK), the FeMo-co assembly scaffold (NifEN), and two PII proteins (NifI$_1$ and NifI$_2$)[1,16]. Methanogens operate near the thermodynamic limits of life, conserving significantly less energy during methane production than aerobic diazotrophs like *Azotobacter vinelandii* or cyanobacteria[17]. Thus, efficient regulation of nitrogenase is vital for survival under fluctuating nitrogen conditions.

In studied methanogens—including *Methanococcus maripaludis*, *Methanosarcina barkeri*, *Methanosarcina mazei*, and more recently *Methanosarcina acetivorans*—nitrogenase production is regulated by fixed nitrogen (e.g., ammonia) availability[18-24]. Unlike proteobacteria, methanogens control nitrogenase activity through direct interaction of the NifI$_{1,2}$ complex with NifDK. Studies with *M. maripaludis* have shown that under nitrogen-replete conditions, the heteromeric NifI$_{1,2}$ complex binds NifDK, likely blocking electron transfer from NifH and thereby inactivating nitrogenase. The presence of 2OG and ATP promotes dissociation of this complex, restoring activity[22,25,26]. Notably, NifI$_{1,2}$ is the only known heteromeric PII complex and the only PII variant that directly interacts with nitrogenase.

Here, we demonstrate that in *M. acetivorans*, the NifI$_{1,2}$ complex forms a heterotrimer composed of two NifI$_1$ subunits and one NifI$_2$ subunit. Six NifI$_{1,2}$ complexes assemble into a novel 30 subunit ~888 kDa supercomplex by binding six NifDK heterodimers. Importantly, this interaction not only prevents NifH from accessing NifDK but also likely induces oligomerization of NifDK itself, forming a unique inhibitory structure. These findings reveal a previously unrecognized mechanism of nitrogenase regulation in methanogens, mediated by PII protein-dependent oligomerization.

Materials and Methods.

Cell Growth. *M. acetivorans* strains were grown in anoxic high-salt (HS) medium at 35º C as previously described[19]. HS medium was prepared inside an anaerobic chamber (Coy Laboratories)

containing 75% N_2, 20% CO_2, and 5% H_2. The medium was supplemented with 2 µg/mL puromycin when required. Growth comparison experiments were performed in Balch tubes containing 10 ml of HS medium containing 125 mM methanol, 1 mM sulfide and 3 mM cysteine. Fixed nitrogen (18 mM NH_4Cl) was added from anaerobic sterile stock solutions prior to inoculation where indicated. The optical density was measured at 600 nm using a spectrophotometer. For purification of Strep-NifD, *M. acetivorans* cells were grown in 1.5 L HS medium lacking NH_4Cl with 125 mM methanol and 0.025% Na_2S. Cells were grown to late log phase (OD_{600} = ~0.5), harvested anaerobically by centrifugation, and the pellet was stored at -80°C in an anoxic Hungate tube. *E. coli* strains were grown maintained with solid or liquid LB medium with the appropriate antibiotic. For recombinant Strep-NifH purification, *E. coli* SufFeScient cells[27] containing pDL906 were grown in four liters of LB medium with 100µg/mL kanamycin, 1 mM L-cysteine, and 1 mM ferric citrate, shaking at 240 rpm at 35° C. At an OD_{600} of 0.5-0.6, expression was induced with 0.2 mM IPTG and supplemented with 0.5 M D-sorbitol and set to shake at 150 rpm at 16° C. After 16 hours, cultures were transferred to sterile GL45-closure reagent bottles with 1.7 mM sodium dithionite, sealed tightly, and incubated at 16° C for 3 hours. These cells were then harvested anaerobically by centrifugation, and the pellets stored at -80°C in anoxic containers.

Strain Construction. Plasmids, primers, and strains used in this study are listed in **Supplementary Tables 1-2.** *M. acetivorans* strain DJL70 that expresses Strep-NifD was generated using the CRISPR-Cas9 system[28] with a few modifications. All primers and gBlocks were designed using Geneious Prime software and purchased from IDT. The gRNA-*nifI₂* was designed to target Cas9 to cut within *nifI₂* (MA3897) upstream of *nifD* (MA3898). Homology regions upstream and downstream of *nifD* were obtained as synthetic gBlocks and contained sequence to add a twin-Strep-tag with linker, identical to as described for Mcr[29], to the N-terminus of NifD. A single nucleotide change (C to A) in the PAM sequence was added to prevent further Cas9 targeting of DNA by gRNA-*nifI₂*. The C to A change does not alter the amino acid sequence of NifI₂. The homology repair templates and gRNA-*nifI₂* were assembled into *Asc*I-digested pDL238 using Gibson assembly Ultra Kits (Telesis Bio) followed by transformation into *E. coli* WM4489 competent cells. Transformants were screened by PCR using primers P7.3 and P6.53 to confirm insert assembly into pDL238. The resultant plasmid pDL705 was retrofitted with pAMG40 using Gateway BP Clonase II enzyme mix (Invitrogen) followed by transformation into WM4489 cells. Transformants were screened by PCR using the primer set P6.64 /P6.65 and the resultant plasmid was designated pDL707.

M. *acetivorans* WWM73 cells were transformed with pDL707. Transformants were selected on anaerobic HS agar plates containing 125 mM methanol and 2 µg/mL puromycin as previously described[30]. Colonies were screened by PCR using primers P7.20 and P7.18 that anneal to the Strep-tag and the 3'-end of *nifD*. Cells from a positive colony were passed through HS medium lacking puromycin then plated on HS agar plates containing 125 mM methanol and 50 µg/mL 8-aza-2,6-diaminopurine sulfate (8-ADP) as described[28] to select for the loss of pDL707. Colonies were assessed for the loss of pDL707 by PCR using primers P7.45 and P7.46. The resultant plasmid-free and puromycin-sensitive strain was designated strain DJL70. PCR amplification of *nifI₂-strep-nifD* from the chromosome of strain DJL70 using primers P7.14 and P7.18, followed by sequencing of the PCR product confirmed the addition of the twin-Strep-tag to NifD.

An *E. coli* strain capable of expressing Strep-tagged NifH (Strep-NifH) was generated by PCR amplifying *nifH* from *M. acetivorans* C2A genomic DNA using Q5 polymerase and primers P8.67 and P8.43 to add restriction sites for *Nco*I and *Xho*I as well as a N-terminal strep tag. The

PCR product was digested with *Nco*I and *Xho*I according to manufacturers' instructions (NEB) and ligated into *Nco*I/*Xho*I-digested pET28a using T4 DNA ligase and incubated for 16 hours at 16°C. The ligation reaction was used to transform *E. coli* DH5α cells and selection done using LB agar plates containing 100 μg/mL kanamycin. Colony PCR and sequencing confirmed *strep-NifH* insertion into pET28a, resulting in plasmid pDL906. Plasmid pDL906 was used to transform *E. coli* "SufFeScient" cells and transformants selected by plating on LB agar plates containing 100 μg/mL kanamycin. *E. coli* "SufFeScient" (pDL906) was used for recombinant production and purification of Strep-NifH.

Protein Purification. All purification steps were performed under anoxic conditions unless otherwise stated. For purification of Strep-NifD, a *M. acetivorans* strain DJL70 cell pellet was resuspended in resuspension buffer (50 mM Tris, 150 mM NaCl pH = 7.2 with 1 mM each PMSF and benzamidine HCl, plus 1.7 mM sodium dithionite). Cells were lysed by sonication on ice with a Qsonica Q55 using a CL-188 probe. Five pulses of sonication (10 seconds each) at an amplitude of 30% with at least one minute on ice between pulses) was followed by anaerobic centrifugation for 10 minutes at 16,000 x *g*, and the supernatant was passed through a 0.45 μm filter. Cleared lysate was loaded onto IBA Strep-Tactin Superflow resin 2 times. The column (0.5 mL) was washed 5 times with Buffer W (50 mM Tris, 150 mM NaCl pH = 7.2 with 1.7 mM sodium dithionite) and protein eluted with Buffer E (Buffer W with 2.5 mM desthiobiotin). Elutions were concentrated using Nanosep 10K Omega columns according to manufacturers' instructions (Pall Corp.).

For purification of Strep-NifH, a SufFeScient (pDL906) cell pellet was resuspended in Resuspension Buffer plus 10 μg/mL DNase I (Sigma) before lysis by three passages through a French pressure cell at > 1100 psi. Lysate was centrifuged at 39,000 x *g* and 4 °C for 10 min, then the supernatant was passed through a 0.45 μm filter. Strep-Tactin column loading, washing with Buffer W, and elution with Buffer E were performed as described above for Strep-NifD.

Strep-NifD and Strep-NifH proteins were quantitated using a Qubit 2.0 fluorometer (Invitrogen). Protein purity was analyzed by SDS-PAGE using a 15% gel (**Supplementary Fig. 1**). A Broad Range (10–250 kDa) color prestained protein standard (New England Biolabs) was used to approximate the molecular weight of each protein. SDS-PAGE gels were stained with Coomassie Brilliant Blue solution, then destained prior to imaging. The protein bands were excised and sent for trypsin digestion followed by mass spectrometry analysis by the IDeA National Resource for Quantitative Proteomics facility at University of Arkansas for Medical Sciences (Source data file).

Nitrogenase activity assay. Nitrogenase activity was assayed by measuring reduction of acetylene to ethylene using methods similar to those described previously[25]. The assay mixtures contained 25 mM HEPES, pH = 7.5, with 5 mM ATP, 12.5 mM MgCl$_2$, 25 mM creatine phosphate, 100 U/mL creatine phosphokinase (Sigma), 20 mM sodium dithionite in reaction volumes of 1 mL. Where indicated, these mixtures were supplemented with 10 mM 2-oxoglutarate. All reactions contained 50 μg purified Strep-NifD and were initiated by the addition of 50 μg purified Strep-NifH. Reaction vials were 10 mL serum vials (Kimble) with a total internal volume measured at 13 mL. These were closed with butyl rubber stoppers and sealed with 20 mm crimps before purging with argon for 5 minutes. 2 mL of acetylene was added by syringe to each vial, then 800 μL assay mix was dispensed by needle to each, and reactions initiated by adding 200 μL Strep-NifH solution by syringe. Ethylene production was monitored by periodic sampling of 200 μL headspace which was injected onto a GC-2030 gas chromatograph (Shimadzu) with a CP-PoraPLOT Q capillary column (Varian, #CP7551). Ethylene standards were used to quantify results.

Western blot. The production of Strep-NifD in *M. acetivorans* strain DJL70 was assessed by the detection of Strep-NifD using a custom polyclonal NifD-specific antibody and a monoclonal Strep-tag-specific antibody (Qiagen, #34850) as previously described[19].

Cryo-EM Sample Preparation and Data-Collection: Cryo-EM samples were prepared following a protocol previously developed for the nitrogenase-like enzyme DPOR[31]. The eluted NifD protein sample (0.7 mg/ml), which co-purifies with NifK, $NifI_1$, and $NifI_2$, was diluted to 0.4 mg/ml using Buffer W in a sealed glass vial inside an anaerobic chamber. Quantifoil R2/2 holey carbon grids (Q350CR2, Electron Microscopy Sciences) were negatively glow discharged for 60 seconds using a Gloqube (Quorum). Immediately following grid cleaning, 3 µL of the protein sample was applied to each grid using a Hamilton gas-tight syringe to minimize air exposure. Grid preparation was performed using a Vitrobot Mark IV (Thermo Fisher Scientific) with the chamber maintained at 4°C and 100% humidity. Without additional incubation, grids were blotted for 2.5 seconds (blot force 2) and plunge-frozen into liquid ethane cooled by liquid nitrogen. The total exposure time of each sample to ambient air during transfer and freezing was less than 30 seconds. Grids were stored in liquid nitrogen until data collection.

For imaging, grids were clipped and loaded into a 200 kV Thermo Fisher Scientific Glacios transmission electron microscope equipped with image correction. Movies were collected using a Falcon IV direct electron detector (4k × 4k) operated with EPU software v2.12.1 (Thermo Fisher Scientific). Data were acquired at a calibrated pixel size of 0.928 Å, with a nominal defocus range of −1.0 to −2.4 µm. Each movie was recorded over a 8.73 second exposure at a dose rate of 5.25 e⁻/Å²/s, resulting in a total dose of 52 e⁻/Å² across 50 frames. This preparation strategy consistently yielded thin, uniform vitreous ice with no evidence of air–water interface artifacts. All samples exhibited excellent particle distribution and orientation diversity in 2D classification analyses.

Data processing: NifDK bound to $NifI_{1a,b}$ and $NifI_2$ structure: Single-particle cryo-EM data were processed using cryoSPARC v4.7.0. For the NifDK dataset, 3,826 raw movies were motion-corrected using patch motion correction, followed by contrast transfer function (CTF) estimation via patch CTF estimation (**Supplementary Fig. 2**). Initial particle picking was performed using the blob picker, and particles were extracted with a box size of 400 pixels (binning factor 4). This yielded an initial stack of 1,798,035 particles. Multiple rounds of 2D classification were conducted to remove false positives and damaged particles, resulting in a cleaned subset of 899,239 particles. These particles were used to generate ab-initio 3D reconstructions.

The 2D classification revealed three distinct particle populations, none of which corresponded to the canonical NifDK heterotetramer. Instead, we observed: (1) a NifDK dimer, (2) a NifDK dimer bound to the inhibitors $NifI_1$ and $NifI_2$, and (3) a higher-order complex consisting of a NifDK heterotrimer associated with $NifI_{1a,b}$ and $NifI_2$. Unfortunately, the volumes corresponding to the NifDK dimer and NifDK dimer bound to the inhibitors $NifI_1$ and $NifI_2$ could not be refined to high-resolution models. These classes exhibited significant conformational flexibility and yielded low-resolution, diffuse densities, consistent with dynamic and heterogeneous behavior.

We were able to successfully process and refine the higher-order complex comprising a NifDK heterotrimer in association with $NifI_{1a,b}$ and $NifI_2$, which displayed well-defined features suitable for detailed structural analysis. The particle stack corresponding to the heterotrimeric complex (124,698 particles) was subjected to multiple rounds of 2D classification to remove poorly aligned particles and improve homogeneity. An initial 3D reconstruction was generated

using ab-initio modeling in cryoSPARC. The resulting volume was further refined using non-uniform refinement to enhance resolution and account for structural variability. Subsequently, 3D classification was performed to resolve conformational heterogeneity, particularly focusing on differences in particle populations based on the presence or absence of associated subunits. 3D classification resulted in the separation of the dataset into three distinct particle classes. Among these, one class, comprising the largest number of particles, yielded the highest-quality 3D reconstruction and clearly resolved all three components of the heterotrimeric complex. This class was further refined using homogeneous refinement followed by non-uniform refinement, during which the particles were re-extracted at full resolution (binning factor of 1). The final map, generated from a subset of 74,083 particles, was further processed using DeepEMhancer and reached a resolution of 3.16 Å at a Fourier shell correlation (FSC) threshold of 0.143 (**Supplementary Table 3**).

Results

M. acetivorans strain DJL70 expresses functional Strep-NifD that co-purifies with NifK, NifI$_1$ and NifI$_2$.

To facilitate the purification of NifDK (MoFe protein) from *M. acetivorans*, strain DJL70 was generated using the CRISPR-Cas9 system to replace native *nifD* in parent strain WWM73 with *nifD* encoding a tandem Strep tag (*strep-nifD*) (**Fig. 1A**). Importantly, strain DJL70 grows similar to strain WWM73 when using N$_2$ as the sole N source, indicating that Strep-NifD does not alter Mo-nitrogenase expression and activity (**Fig. 1B**). Also, like strain WWM73, and consistent with previous results[18,19], Strep-NifD is only detected by Western blot in DJL70 cells grown without NH$_4$Cl (**Fig. 1A**). Affinity purification of Strep-NifD results in the co-purification of NifK and NifI$_1$ and NifI$_2$, as seen by SDS-PAGE (**Fig. 1C**) and confirmed by mass spectrometry. In *M. maripaludis*, NifI$_1$ and NifI$_2$ form an inhibitory complex that binds NifDK to block N$_2$ reduction in response to depletion of 2OG, an indicator of fixed N sufficiency[26]. Despite strain DJL70 grown only with N$_2$ as the sole N source, a significant portion of NifDK appears in the NifI$_1$-NifI$_2$ (NifI$_1$I$_2$) bound inhibited state, similar to results obtained with *M. maripaludis*[25]. Importantly, the addition of 2OG was shown to the release of NifI$_1$I$_2$ from NifDK, allowing electron transfer from NifH thereby facilitating the reduction of N$_2$ by NifDK[25,26]. To ascertain if co-purified NifI$_1$I$_2$ performs the same function in *M. acetivorans*, the acetylene reduction activity of purified NifDK was assayed with purified recombinant NifH (**Supplementary Fig. 1**) in reaction mixtures with or without 2OG (**Fig. 1D**). An approximate 2-fold increase in the specific activity (nmol ethylene reduced min^{-1} mg^{-1}) of NifDK with 2OG was observed [537 ± 97 (-2OG) vs 1239 ± 67 (+2OG)], indicating that a significant portion of *in vivo* NifDK is in the inactive NifI$_1$I$_2$ bound inhibited state. To understand how NifI$_1$I$_2$ binds to NifDK, the affinity-purified complex was directly analyzed by single-particle cryo-EM without the addition of 2OG.

M. acetivorans NifDK forms a supercomplex with NifI$_1$-NifI$_2$

2D-classification of the cryo-EM images revealed the presence of three distinct particle populations in the dataset (**Supplementary Fig. 2**). Two of these were smaller in size, but these volumes resulted in poorly defined density due to conformational heterogeneity and thus did not yield high-resolution reconstructions. Manual docking of NifDK subunits to these volumes suggest the likely presence of NifDK dimers and NifDK dimers bound to the NifI$_1$I$_2$ inhibitor complex (**Supplementary Fig. 3**). However, the third volume showcased a more structurally homogeneous larger complex which resulted in a high-resolution structure of the *M. acetivorans* NifDK-NifI$_1$-NifI$_2$ supercomplex at 3.16 Å (**Fig. 1E, F; Supplementary Table 3**). The supercomplex reveals a non-canonical assembly of nitrogenase comprising three NifDK heterotetramers, each associated

Figure 1. Genetic, biochemical, and structural analysis of nitrogenase function in *M. acetivorans*. A. Schematic comparison of the nifHI$_1$I$_2$DKEN operon in strains WWM73 and DJL70. A tandem N-terminal Strep-tag was introduced at the *nifD* locus in WWM73 using CRISPR-Cas9 to generate strain DJL70. The sub-panel below shows Western blot analysis of lysates from cells grown in **(B)**, probed with antibodies against NifD and the Strep tag. **B.** Growth of each strain in HS medium with methanol (125 mM) with either NH$_4$Cl or N$_2$ as the nitrogen source. **C.** SDS-PAGE analysis of a representative purification of Strep-tagged NifD, showing co-elution of NifK, NifI$_1$, and NifI$_2$ proteins, from strain DJL70, as described in the Materials and Methods. Two additional bands below the full-length NifD and NifK, marked with asterisks (*), were later identified as truncated forms of NifD and NifK. **D.** Effect of 2-oxoglutarate (2OG) on acetylene reduction activity of purified NifDK. Ethylene production was measured per mg of NifDK (75 µg) in the presence of recombinant NifH (150 µg), with or without 10 mM 2OG. Reactions without added NifH were included as a negative control. Data represent the mean±SD from three biological replicates. **E.** Representative Cryo-EM map of the nitrogenase super-complex resolved at 3.13 Å, revealing the overall architecture and spatial organization of its core components. The supercomplex consists of three NifDK units, each adopting an α$_2$β$_2$ architecture and coordinated by a NifI heterotrimer positioned on either side of the tetrameric assembly. In each NifDK unit, NifD (α) and NifD′ (α′) are shown in eggplant and teal, while NifK (β) and NifK′ (β′) are colored coral and gold, respectively. The associated inhibitor proteins-NifI$_{1a}$, NifI$_{1b}$, and NifI$_2$ are depicted in pale violet-red, pink, and maroon, respectively. **F.** Cryo-EM structure of the supercomplex highlighting a single NifDK tetrameric unit bound by two heterotrimeric NifI complexes on either end (colors are the same as in panel E).

with two heterotrimeric NifI-inhibitor complexes consisting of two NifI$_1$ (NifI$_{1a}$ and NifI$_{1b}$)

subunits and one NifI$_2$ subunit. For clarity in presentation, we here define each NifDK tetramer along with the associated NifI$_{1a,b}$-NifI$_2$ complexes bound on either half as one unit. Three such units form the supercomplex. In addition, we also define the two halves within each unit: For

Figure 2. Structural features of individual supercomplex units and overlay with *A. vinelandii* NifDK bound to NifH and FeSII. A. Cryo-EM structure of an individual *M. acetivorans* (Ma) NifDK tetramer highlighting the core components of a single unit. Each unit adopts an α$_2$β$_2$ architecture to form the core NifDK complex and is bound by two NifI heterotrimers positioned on either side of the tetrameric assembly. In the NifDK unit, NifD (α) and NifD′ (α′) are shown in eggplant and teal, while NifK (β) and NifK′ (β′) are shown in coral and gold, respectively. The associated inhibitor proteins NifI1a, NifI1b, and NifI2 are depicted in pale violet-red, pink, and maroon, respectively. The callouts highlight the P-cluster, the M-cluster containing homocitrate (HCA), and the His-485 ligand. Each cluster is displayed with its corresponding carved density map. **B.** Overlay of a single Ma NifI (teal)–NifDK (pink) complex (Unit 1) from our cryo-EM structure with the asymmetric *A. vinelandii* nitrogenase NifDK (MoFe protein; grey; PDB: 7UT8) bound to NifH (Fe protein; blue). **C.** Overlay of a single Ma NifI (teal)–NifDK (pink) complex (Unit 1) from the supercomplex onto the *A. vinelandii* NifDK (grey) tetramer (MoFe protein; PDB: 8RHP) bound to FeSII (yellow) and NifH (Fe protein; blue).

example, NifD (α) and NifK (β) subunits in unit 1 are labeled as α1, β1 and α1′, β1′. The NifI proteins bound in unit 1 are correspondingly labeled as NifI$_{1a}$, NifI$_{1b}$, NifI$_2$, and NifI$_{1a′}$, NifI$_{1b′}$, NifI$_{2′}$, respectively (**Fig. 2A**).

Despite the unusual higher-order assembly, the core architecture of each NifDK unit remains conserved and resembles that of the well-characterized *Azotobacter vinelandii* (Av) nitrogenase (**Fig. 2A & B**). The NifD and NifK subunits form an $\alpha_2\beta_2$ heterotetramer (MoFe-protein) and each half houses a M-cluster ([Mo-Fe$_7$S$_9$C•homocitrate]) and P-cluster ([Fe$_8$S$_7$]) (**Fig. 2A and Supplementary Figs. 4 & 5**). The M-cluster in *M. acetivorans* NifDK is coordinated by His-485, and Cys-267 from NifD while Lys-469 interacts with homocitrate (HCA) (**Supplementary Fig. 4A**). The P-cluster is coordinated by Cys-48, Cys-106, Cys-23, and Ser-141 from NifK along with Cys-58, Cys-84, and Cys-149 from NifD (**Supplementary Fig. 4B**). These contacts are identical in each NifDK half in all three units observed in the supercomplex. While the overall structure of NifDK in all three units are similar, minor structural changes are observed, especially around the M-cluster. His-485, positioned near the M-cluster, adopts different conformations in the NifDK halves in units 1 & 2 of the supercomplex (**Supplementary Fig. 5**).

The NifI heterotrimeric inhibitor complex coordinates assembly of multiple MoFe-proteins in the supercomplex

PII proteins are a conserved group of nitrogen-regulatory proteins that form homotrimeric complexes in structural studies[12,32]. In contrast, in the supercomplex, the NifI proteins form a heterotrimer (NifI complex) with one subunit of NifI$_2$ wedged between two copies of NifI$_1$ (NifI$_{1a}$ & NifI$_{1b}$; **Fig. 2A**). Each NifDK half in the tetramer is bound to a NifI complex. Comparison of the *A. vinelandii* (*Av*) nitrogenase and the NifI-bound NifDK complex from *M. acetivorans* shows that the NifH and NifI binding interfaces on the NifDK complex overlap (**Fig. 2B**) and reveal how nitrogenase activity is switched-off by NifI complex binding in methanogens. Recently, *Av* nitrogenase was shown to form inactive and oxygen-resistant assemblies promoted by Shethna protein II (FeSII)[33]. Like NifI in methanogens, FeSII is an inhibitor of nitrogenase in *Av* and drives the assembly of NifDK, along with NifH, into higher order complexes [33]. However, the binding of FeSII and the NifI complex are distinct (**Fig. 2C**). These structures suggest that the mechanism of regulation of nitrogenase by the PII proteins in methanogens is different than inhibition by FeSII in *Av*. Moreover, assembly of the supercomplex in *M. acetivorans* involves direct MoFe protein interaction, whereas Av MoFe proteins do not interact in higher order structures. These differences highlight the evolutionary divergence in regulatory strategies to inactivate nitrogenase between anaerobic methanogens and aerobic bacteria.

The nitrogenase supercomplex from *M. acetivorans* reveals extensive inter-subunit interactions among neighboring NifDK heterotetramers that contribute to the stability of the higher-order assembly (**Fig. 3**). The three NifDK complexes are arranged such that each tetramer is tilted ~48º with respect to each other. The buried surface area is ~1000 Å^2 between adjacent NifDK tetramers in the supercomplex. An array of charged and polar residues forms a dense network of acid-base salt bridges and hydrogen bonds between the NifD and NifK subunits from adjacent units, with the majority of the interactions mediated by NifK residues (**Fig. 3**). The spatial distribution of these interactions is symmetrical and therefore forms a cooperative electrostatic interface that stabilizes the trimeric arrangement. This supercomplex of *Ma* nitrogenase is distinct from other structures of bacterial nitrogenases that are typically observed as single NifDK heterotetramers[34]. There is no other reported evidence for such direct interactions between NifDK tetramers either under *in vivo* or *in vitro* conditions. Nitrogenase from aerobic bacteria have recently been shown to form a filamentous higher order structure, but this complex is formed through protein-protein interactions between nitrogenase and regulatory factors (e.g., FeSII)[33]. The presence of extensive electrostatic interfaces in *Ma* NifDK suggests a distinct regulatory mechanism unique to NifI-containing anaerobes. The stabilization of the trimeric assembly

through inter-subunit salt bridges could lock the enzyme in a catalytically inactive state, representing a previously unobserved strategy for nitrogenase inhibition and assembly regulation. However, a NifH homodimer can be manually docked on NifDK within the supercomplex architecture with no steric hinderance if the NifI complex is removed (**Supplementary Fig. 6A**). Therefore, it is also possible that this supercomplex could be the functional state of methanogenic nitrogenase capable of binding NifH and reducing N_2 within this higher order architecture.

Figure 3. Interactions between NifDK tetrameric units within the supercomplex. A. Surface representation of NifDK subunits in the supercomplex. The interaction interface between Unit 1 and Unit 2 is highlighted. **B.** Close-up view of residues mediating this interaction. **C.** The interface between the NifDK subunits in Unit 1 and Unit 3 is indicated. **D**. Detailed view of residues involved in this interaction. **E.** Top view of the NifDK tetrameric units within the supercomplex, showing extensive interactions between the NifK (β) subunits. **F.** Enlarged view highlighting the individual residues involved in these interactions. In this figure, NifD (α), NifD′ (α′), NifK (β), and NifK′ (β′) are colored eggplant, teal, coral, and gold, respectively.

Each NifI inhibitor in our structure is arranged as a NifI$_{1a}$-NifI$_2$-NifI$_{1b}$ complex and is positioned to mediate interactions with two adjacent NifDK units in the supercomplex. An

example of the NifI complex connecting two distinct NifDK units is shown in **Fig. 4**. This spatial

Figure 4. Interactions between the NifDK tetrameric unit and the NifI complex within the nitrogenase supercomplex. A. Cartoon representation of a single NifI complex interacting with NifDK tetramers of Unit 1 and Unit 3 and overlaid on the surface-rendered supercomplex. Interaction interfaces with Unit 1 and Unit 3 are highlighted by the blue and green demarcations, respectively. **B.** Close-up view of the NifDK units and associated NifI complex, illustrating their spatial arrangement and interface architecture. **C.** Detailed view of residues mediating the interaction between $NifI_{1a}$ and $NifI_{1b}$ of the NifI complex with NifDK of Unit 3. **D. & E.** Detailed views of residues involved in the interaction between $NifI_2$ and NifDK. NifD (α), NifD′ (α′), NifK (β), and NifK′ (β′) are colored eggplant, teal, coral, and gold, respectively. The NifI subunits $NifI_{1a}$, $NifI_{1b}$, and $NifI_2$ are colored pale violet-red, pink, and maroon, respectively.

configuration facilitates extensive inter-subunit contacts within the NifI complex, predominantly mediated by salt bridges, with additional contributions from hydrogen bonding and hydrophobic interactions (**Fig. 4C & D**). $NifI_2$ interacts in *cis* with NifDK in the primary unit (**Fig. 4D**). In

contrast, NifI₁ₐ-NifI₁ᵦ make extensive contacts in *trans* with NifDK of the adjacent unit (**Fig. 4C**). The NifI complex and NifDK interactions are largely conserved across all three units, albeit with some differences in the total buried surface area within the interaction interface (**Supplementary Fig. 7**). Thus, in addition to extensive interactions between NifDK tetramers within the three units, each NifI complex also serves to connect and stabilize the three units within supercomplex.

The NifI heterotrimeric complex is structurally similar to canonical homotrimeric PII proteins.

Canonical PII proteins are homotrimers, consisting of monomeric units each typically 11-18 kDa. Each monomer exhibits a ferredoxin-like fold with three protruding flexible loops (T, B, and C; **Supplementary Fig. 8**). In the trimeric state, ligand binding sites are formed at the dimeric interface of two subunits, whereby the B-loop and the base of the T-loop of one subunit and the C-loop of the adjacent subunit form a pocket capable of binding ADP, ATP, and/or 2OG[12,14] (**Supplementary Fig. 9A**). As such, the trimer has three intercommunicating ligand binding sites. For example, GlnK in bacteria and archaea controls ammonium transport by the Amt transporter. At low intracellular levels of 2OG, which indicates ammonium sufficiency, GlnK binds ADP which results in the T-loop having a rigid extended confirmation that specifically binds within the channel of Amt effectively blocking ammonium transport. When ammonium is depleted, the intracellular levels of 2OG rise and GlnK subsequently binds 2OG along with ATP, resulting in the T-loop changing conformation and disassociation of GlnK from Amt[35,36]. Thus, the ADP-bound form of PII proteins typically binds target proteins and the ATP/2OG bound form typically does not.

NifI₁ (11 kDa) and NifI₂ (12 kDa) are similar in size to canonical PII proteins and contain the conserved residues involved in binding ADP/ATP and 2OG (**Supplemental Fig. 10**). A distinguishing difference is that NifI₂ has a longer T-loop and an extended C-loop compared to NifI₁. Importantly, both the T-loops of NifI₁ and NifI₂ were shown to be required for NifI₁-I₂ association with NifDK in *M. maripaludis*, consistent with the T-loops modulating target protein interaction[25]. Interestingly, the T-loops of NifI₁ from both methanogens are quite similar in sequence, whereas the longer NifI₂ T-loop sequences a markedly different, indicating there are likely differences in target protein specificity (**Supplemental Fig. 10**). Consistent with other canonical homotrimeric PII proteins, the NifI heterotrimer complex exhibits a ferredoxin-like fold, with the trimerization interface stabilized by a central β-sheet core as shown for the NifI complex bound to unit 1 in the supercomplex (**Fig. 5**). The T-loops for both NifI₁ₐ and NifI₂ are structured and extended, whereas the T-loop for NifI₁ᵦ is unresolved/flexible (**Fig. 5B**). Several residue-specific interactions reinforce the stability of the complex, including salt bridges and hydrophobic interactions (**Fig. 5C**). Collectively, these interactions promote formation of a stable heterotrimeric NifI complex. The NifI₂ monomeric unit is identical in structure to that of GlnK, (**Supplementary Fig. 8A**), except for the T-loop and extended C-loop. All six NifI complexes are largely identical when superimposed (**Supplementary Fig. 12)**, with only differences in the T-loop conformation observed. Thus, although the NifI complex is composed of two separate proteins, the assembled heterotrimeric complex is indistinguishable to canonical PII homotrimers, indicting the NifI T-loop confirmation is also responsive to 2OG and ATP/ADP binding.

The NifI complexes exhibit differential 2OG and ADP occupancy that reveals functional asymmetry.

ATP/ADP and 2OG function as metabolic signals that regulate PII protein interaction with targets[12]. In canonical PII proteins, 2OG binding typically requires co-binding of Mg^{2+} and ATP that results in disassociation from the target protein. Depletion of 2OG has been shown to increase

Figure 5. Structural organization and intra-complex interactions within the NifI heterotrimer. A. Cartoon representation of a single NifI heterotrimeric complex, composed of NifI1a (pale violet-red), NifI1b (pink), and NifI2 (maroon), shown against the grey surface representation of the remaining supercomplex. **B.** Close-up view of the NifI complex illustrating the architecture of NifI$_{1a}$, NifI$_{1b}$, and NifI$_2$, with their respective T-, B-, and C-loops colored blue, green, and purple, respectively. **C.** Detailed view of residues mediating intra-complex interactions among the NifI$_{1a}$, NifI$_{1b}$, and NifI$_2$ subunits.

the ATPase activity of PII proteins, converting ATP to ADP that promotes T-loop alteration and association with the target protein[37]. Previous studies with *M. maripaludis* showed that 2OG/ATP promotes the *in vitro* dissociation of the NifI complex from NifDK, establishing the nucleotide/2OG dependent regulation of nitrogen fixation and nitrogenase activity in methanogens[25,26]. As such, ADP-bound NifI complex should be associated with NifDK. However, the binding of ADP and how these ligands engage with the NifI complex when in complex with nitrogenase has not been established. The six NifI complexes in our CryoEM structure display

asymmetric and heterogeneous binding to ADP and/or 2OG (**Fig. 7 & Supplemental Fig. 11**). ADP and/or 2OG interactions are mediated by the B-, C-, and T-loops of the NifI subunits (**Supplemental Fig. 12**). Notably, ADP is found in at least one ligand-binding site in five of the six NifI complexes, consistent with the ADP-bound form driving association with NifDK.

Consistent with the conservation of the ligand binding residues in the primary sequence (**Supplemental Fig. 10**), the architecture of the three ligand binding pockets of the NifI complex is similar to that of GlnK (**Supplemental Fig. 9**). Pocket 1 is formed by the base of the T-loop and B-loop from $NifI_{1a}$ and the C-loop of $NifI_2$. Pocket 1 of each NifI complex contains a ligand, with five containing ADP and one with 2OG. Pocket 2 is formed by the base of the T-loop and B-loop from $NifI_2$ and the C-loop of $NifI_{1b}$. Pocket 2 displays the most heterogeneity in ligand occupancy, with two NifI complexes containing ADP, two containing 2OG, and two free of any ligands (**Fig. 6**). Pocket 3 is formed by the base of the T-loop and B-loop from $NifI_{1b}$ and the C-loop of $NifI_{1a}$. Pocket 3 is free of ligand in all six NifI complexes. These results indicate that the ligand affinity differs between the three binding pockets, with pocket three likely having the lowest affinity for ligand (**Fig. 6 and Supplemental Fig. 11**). Moreover, the asymmetric binding of ligand by the NifI complexes is a feature of the stable supercomplex. The asymmetry of ligand association may poise the complex for rapid release in response to small changes in 2OG and ATP/ADP levels.

Interestingly, binding of 2OG is tolerated in ligand binding pockets 1 and 2 that still allows stable association of the NifI complex with NifDK. Moreover, 2OG binding by the NifI complexes is independent of ATP. However, previous results with *M. maripaludis* showed that the addition of both ATP and 2OG were required for the *in vitro* disassociation of the NifI complex from NifDK[25]. Similarly, washing resin-bound strep-NifD-NifK with 2OG and ATP removed associated $NifI_1$ and $NifI_2$ from the complex, resulting in the elution of NifDK that is free of $NifI_1$ and $NifI_2$ (**Supplementary Fig. 13**). Taken together, these results support that model that ADP binding by the NifI complexes triggers association with NifDK and likely the assembly of the supercomplex. NifI complex binding is reversible with elevated levels of 2OG/ATP releasing the NifI complex from NifDK. The disassociation of the NifI complex from NifDK could lead to disassembly of the supercomplex. However, we cannot rule out the possibility that the trimeric NifDK complex could also be the active form of the enzyme.

The α-loop insertion of *M. acetivorans* NifD directly interacts with the T-loop of $NifI_2$.

Although all known and predicted structures of the NifDK proteins are very similar, there is some sequence and structural heterogeneity (**Fig. 7A and Supplementary Fig. 14**). For example, most aerobic diazotrophic bacteria have an N-terminal extension in their NifK (β) subunit that is hypothesized to stabilize the heterotetramer[38]. In contrast, not only do diazotrophic bacteria and archaea that are anaerobes lack the NifK extension, but many instead have a ~ 50 amino acid insertion within NifD (α-subunit) called the α-loop insertion[38] (**Fig. 7A**). For example, the crystal structure of NifDK from *Clostridium pasteurianum* shows the α-loop insertion adjacent to the predicted binding site of NifH, indicating that the α-loop is a specificity determinant for interaction with NifH[39]. All methanogens known to or predicted to fix nitrogen possess a *nif* operon that contains both $NifI_1$ and $NifI_2$, indicating that NifI-dependent control of nitrogenase activity is a universal feature of nitrogen fixation in methanogens[16]. However, deeply rooted methanogens, such as *M. maripaludis*, lack the α-loop insertion within NifD while later evolving Methanosarcinales, including *M. acetivorans*, have the NifD α-loop insertion (**Supplemental Fig. 14**)[38]. However, *C. pasteurianum* lacks NifI proteins.

Figure 6. Ligand Binding occupancy of the six NifI complexes bound to the three NifDK tetrameric units within the supercomplex. (A–C) Structural views of Units 1, 2, and 3 of the supercomplex, respectively, showing the core NifDK subunits in grey and the corresponding NifI complexes in pink, cyan, and sandy brown. For each unit, the T-loop (blue), B-loop (green), and C-loop (magenta) of the $NifI_2$ subunit are highlighted. Rectangles indicate the occupancy of ADP or 2-oxoglutarate (2OG) within the NifI complex. The electron density for each of the bound nucleotide/2OG molecule is also depicted. **D**. Detailed view of residues mediating the interaction with ADP in one binding site (left ligand-binding pocket at the top of Unit 1 in panel A) is shown. **E**. Detailed view of residues mediating the interaction between 2OG and the right ligand-binding pocket (right ligand-binding pocket at the top of Unit 1 in panel A) is shown.

In our supercomplex structure, not only does the NifI complex binding site overlap with the putative NifH binding site (**Fig. 2**), but the T-loop of $NifI_2$ also extends out and associates directly with the NifD α-loop insertion in five of the six the NifI complexes (**Figs. 6 and 7**). Thus, the α-loop is a key specificity determinant in the association of the NifI complex with NifDK in *M. acetivorans*. This result is consistent with substantial differences in the $NifI_2$ T-loop sequences

Figure 7. α-loop features in NifDK. A. Schematic comparison of NifDK subunit architecture in *Azotobacter vinelandii* (Av), *Clostridium pasteurianum* (Cp), and *Methanosarcina acetivorans* (Ma), highlighting the N-terminal extension in *Av* (red) and α-loop insertion in *Cp* and *Ma* (green). **B.** Overall structure of the three NifDK tetrameric units (Unit 1, pink; Unit 2, cyan; Unit 3, orange) with α- and α′-loops (blue) mapped. The NifI complexes are surface-rendered (grey). **C–E.** Detailed structures of NifDK and NifI complexes in Unit 1, Unit 2, and Unit 3, respectively, with α- and α′-loops highlighted. Inset in panel C showcases residues in the α-loop of Unit 1 and $NifI_2$ that promote the interaction. **F.** Superposition of the three NifDK tetrameric units illustrating variability in α- and α′-loop conformations. **G**. Structural overlay of the α- and α′-loops in this study (blue) with those from *C. pasteurianum* (PDB: 8DBY) and *A. vinelandii* (PDB: 4WES), showing loop conformational diversity.

between *M. acetivorans* and *M. maripaludis* (**Supplemental Fig. 14**) Thus, the binding of the NifI complex to NifDK, and possibly supercomplex formation, is likely different between methanogens that contain or lack the α-loop insertion. The T-loop of *M. acetivorans* $NifI_2$ is uniquely suited to interact with the α-loop insertion, and this interaction is likely depending on ligand binding by the NifI complex. Specifically, binding of ADP is likely critical for facilitating the $NifI_2$ T-loop in an extended confirmation to allow the association with the α-loop of NifDK. Notably, the only NifI complex with an unstructured $NifI_2$ T-loop and a NifDK α-loop is the NifI complex that lacks any ADP and contains only 2OG in ligand binding pocket 2 (**Fig. 7E**). Moreover, the loss of the T-loop/ α-loop interaction decreases the buried surface area between the NifI complex and NifDK in this unit by one-half (**Supplemental Fig. 7C**), indicating that this interaction is substantial but not essential for the stable association of the NifI complex to NifDK, at least in the supercomplex. The shorter $NifI_{1a}$ T-loop also has a structured confirmation. However, the T-loop of $NifI_{1b}$ in all NifI complexes is unstructured, indicating that this T-loop is not involved in association of the NifI complex with NifDK in the supercomplex. These results reveal that the T-loop of $NifI_2$ and the α-loop of NifD are critical, and unique, factors involved in the association of the NifI complex with NifDK in *M. acetivorans*.

Discussion

Our cryo-EM structure reveals a previously uncharacterized mechanism of nitrogenase regulation in methanogens, involving the formation of a higher-order complex through direct interactions between NifDK catalytic subunits. The non-canonical PII proteins $NifI_1$ and $NifI_2$ assemble into a heterotrimeric complex that binds each half of the NifDK heterotetramer, resulting in (1) inhibition of nitrogenase activity by blocking electron transfer from NifH, and (2) formation of an inactive supercomplex comprising three NifDK units.

Given that all known and predicted diazotrophic methanogens encode $NifI_1$ and $NifI_2$, this supercomplex likely represents a conserved regulatory feature in these organisms. Indeed, similar-sized complexes were observed by size-exclusion chromatography after purification of $NifI_{1,2}$-NifDK from *M. maripaludis*[25].   While no analogous structures have been observed in bacteria, the presence of NifI homologs in several diazotrophic bacterial species suggests that similar regulatory mechanisms may exist outside of methanogens[10,40]. Notably, NifI proteins are restricted to anaerobes, raising the question of why anaerobic organisms—including all methanogens—utilize NifI-mediated regulation, whereas aerobes rely on alternative systems to control nitrogenase activity.

Methanogens operate near the thermodynamic limits of life, deriving minimal energy from methanogenesis compared to aerobic respiration[17]. Consequently, energy-efficient regulation of nitrogen fixation is essential. Direct inhibition of NifDK by NifI complex binding in response to fixed nitrogen availability is likely more rapid and energetically favorable than indirect mechanisms. Supporting this, ammonia addition leads to a swift loss of nitrogenase activity in methanogens[20,22]. In contrast, many aerobic bacteria employ the DraTG system, which inactivates nitrogenase via ADP-ribosylation of NifH—a process that requires additional enzymes and substrates and is likely slower and less efficient[11,13].

The formation of the supercomplex appears to stabilize the inactive state of NifDK beyond simply preventing NifH binding. Two lines of evidence support this: (1) the NifI complex bridges two NifDK heterotetramers, increasing the interaction surface and enhancing complex stability, as evidenced by successful purification of the supercomplex, whereas single NifDK tetramer-NifI complex assemblies were not observed by cryo-EM; and (2) recent cryo-EM studies of bacterial nitrogenase reveal conformational changes in NifDK during catalysis[41], which are likely restricted in the NifDK-NifI supercomplex. This structural constraint may further prevent catalysis and limit

substrate access to the active site. Importantly, NifI complex binding is reversible, as shown by dissociation from resin-bound NifDK and increased activity in the presence of 2OG and ATP.

Unexpectedly, in the native state, we observed asymmetrical binding of ADP and 2OG within the NifI complexes. In canonical PII proteins, ADP binding typically promotes target association, while MgATP-2OG binding typically promotes dissociation[12]. Here, 2OG binds in the absence of ATP, indicating that partially 2OG occupancy is insufficient to trigger dissociation of the NifI complex. Ligand occupancy was detected in binding pockets 1 and 2—each partially formed by $NifI_2$—while pocket 3, formed between $NifI_{1a}$ and $NifI_{1b}$, remained unoccupied. Despite this asymmetry, the T-loops across the six NifI complexes are largely symmetric, with all six the $NifI_{1b}$ T-loops disordered/unstructured, all six Nif_{1a} T-loops structured, and five of the six longer $NifI_2$ T-loops structured (**Fig. 6, Supplemental Figs. 9–10**). The lone exception is the disordered/unstructured $NifI_2$ T-loop in the top half of Unit 3, which is the only NifI complex to have only 2OG bound (**Fig. 7**). This asymmetry suggests a stable feature of the supercomplex, potentially allowing it to respond more sensitively to fluctuations in ATP/ADP and 2OG levels than a uniformly ADP-bound complex.

The T-loop of canonical PII proteins plays a central role in both ligand binding and target interaction. Consistent with this, deletion of the T-loop region in either $NifI_1$ or $NifI_2$ from *M. maripaludis* disrupts co-purification with NifDK—completely abolishing it with $NifI_1$-ΔT and significantly reducing it with $NifI_2$-ΔT[25]. The *M. acetivorans* NifI-NifDK supercomplex supports these findings. While the $NifI_{1a}$ and $NifI_{1b}$ T-loops do not directly contact NifDK, the $NifI_{1a}$ T-loop is essential for ligand binding at pocket 1, which is consistently occupied by ADP or 2OG across all six NifI complexes (**Supplementary Fig. 11**). This suggests that ligand occupancy at pocket 1 is required for NifI complex association with NifDK, and without the NifI -T-loop ligand cannot be bound. *M. maripaludis* $NifI_2$-ΔT retains some binding to NifDK, consistent with the *M. acetivorans* supercomplex structure that reveals $NifI_2$ contributes most of the interaction surface with NifDK, with approximately half of it located outside the T-loop (**Supplemental Fig. 7**).

Importantly, the $NifI_2$ T-loop directly interacts with the α-loop insertion in NifD (**Fig. 7**), identifying the α-loop as a key specificity determinant for NifI complex interaction with NifDK. This insertion is absent in *M. maripaludis* and other Methanococci but present in all Methanosarcinales and in *Clostridium pasteurianum*, although the latter lacks $NifI_1$ and $NifI_2$ and shows significant sequence divergence in the α-loop (**Supplemental Fig. 14**). Despite sequence differences, the α-loops of *M. acetivorans* NifD and *C. pasteurianum* NifD are structurally conserved (**Fig. 7**F), suggesting a conserved structural role. The $NifI_2$ T-loop to NifD α-loop interaction appears to be a defining feature of nitrogenase regulation in Methanosarcinales. In contrast, Methanococci must rely on alternative NifD regions for $NifI_2$ T-loop interaction. The acquisition of the α-loop insertion in Methanosarcinales may have evolved to enhance specificity of NifDK for NifI and NifH, particularly given their unique capacity to express all three nitrogenase isoforms—Mo-, V-, and Fe-nitrogenases[18]. We previously demonstrated that *M. acetivorans* expresses all three simultaneously[18], and the α-loop likely ensures selective interaction of NifH and NifI with NifDK, excluding VnfDGK and AnfDGK. Supporting this, the Vnf and Anf operons in *M. acetivorans* encode distinct PII proteins with divergent T-loops (**Supplemental Fig. 10**), despite each using the same Fe-protein[15]. Similarly, the α-loop in *Cp*NifD may serve to restrict NifH interaction to NifDK, despite the absence of NifI proteins. Thus, the supercomplex structure highlights the coevolution of nitrogenase (e.g. α-loop insertion) with factors (NifI complex) that control activity.

Finally, this study uncovers a previously unrecognized mechanism of nitrogenase regulation in methanogens, characterized by the formation of a stable, inactive supercomplex mediated by non-canonical PII proteins $NifI_1$ and $NifI_2$. The direct inhibition of NifDK, coupled

with structural stabilization through higher-order assembly, represents an energy-efficient strategy uniquely suited to the metabolic constraints of anaerobic life. The discovery of the α-loop insertion in NifD as a specificity determinant for NifI$_2$ interaction further reveals a molecular basis for selective nitrogenase regulation in Methanosarcinales, particularly in organisms expressing multiple nitrogenase isoforms. These findings expand our understanding of nitrogenase control, highlight evolutionary adaptations in anaerobic archaea, and suggest broader implications for nitrogen fixation strategies across diverse microbial lineages.

Acknowledgments: We thank Drs. Summer Brock, Katherine Basore, and Bradley Readnour at the Washington University Center for Cellular Imaging and Drs. Liguo Wang, Guobin Hu, and Jake Kaminsky at the Laboratory for BioMolecular Structure (LBMS) at Brookhaven National Labs for Cryo-EM data collection. LBMS is supported by the Department of Energy, Office of Biological and Environmental Research (KP1607011). This work was supported by grants from the Department of Energy, Office of Basic Energy Sciences, DE-SC0020965 (EA) for structure determination, DE-SC0019226 (DJL) for strain construction, protein purification, and activity analyses, DE-SC0018143 (BB) for mass spectrometry. Funding for the Montana State Mass Spectrometry Facility used in this publication was made possible in part by the MJ Murdock Charitable Trust, MSU office of the VPRED, and the National Institute of General Medical Sciences of the National Institutes of Health under Award Numbers P20GM103474 and S10OD28650. We also acknowledge generous financial support from the Doisy Research Fund of the Edward A. Doisy Department of Biochemistry and Molecular Biology at Saint Louis University School of Medicine.

Author contributions:

Conceptualization: EA and DJL

Methodology: RK, TMB, AD, MC, MTK, and BB

Investigation: RK, TMB, AD, MC, and MTK

Visualization: RK, EA, and DJL

Funding acquisition: EA, DJL, BB

Project administration: EA and DJL

Supervision: EA, DJL, and BB

Writing – original draft: DJL, EA, and RK

Writing – review & editing: all authors

Competing interests: All authors declare that they have no competing interests.

Data and materials availability: The single particle cryo-EM maps and models have been deposited into the PDB and EMDB with the following codes **PDB: 9P1X & EMDB-EMD-71144**

Supplementary Materials

Materials and Methods

Figs. S1 to S14

Tables S1 to S3

References

Dos Santos, P. C., Fang, Z., Mason, S. W., Setubal, J. C. & Dixon, R. Distribution of nitrogen fixation and nitrogenase-like sequences amongst microbial genomes. *BMC genomics* **13**, 162 (2012). https://doi.org:10.1186/1471-2164-13-162

Peters, J. W., Fisher, K. & Dean, D. R. Nitrogenase structure and function: a biochemical-genetic perspective. *Annual review of microbiology* **49**, 335-366 (1995). https://doi.org:10.1146/annurev.mi.49.100195.002003

Seefeldt, L. C. *et al.* Reduction of Substrates by Nitrogenases. *Chemical reviews* **120**, 5082-5106 (2020). https://doi.org:10.1021/acs.chemrev.9b00556

Seefeldt, L. C. *et al.* Energy Transduction in Nitrogenase. *Acc Chem Res* **51**, 2179-2186 (2018). https://doi.org:10.1021/acs.accounts.8b00112

Rutledge, H. L. & Tezcan, F. A. Electron Transfer in Nitrogenase. *Chemical reviews* **120**, 5158-5193 (2020). https://doi.org:10.1021/acs.chemrev.9b00663

Einsle, O. & Rees, D. C. Structural Enzymology of Nitrogenase Enzymes. *Chemical reviews* **120**, 4969-5004 (2020). https://doi.org:10.1021/acs.chemrev.0c00067

Hu, Y. & Ribbe, M. W. Nitrogenase and homologs. *Journal of biological inorganic chemistry : JBIC : a publication of the Society of Biological Inorganic Chemistry* **20**, 435-445 (2015). https://doi.org:10.1007/s00775-014-1225-3

Halbleib, C. M. & Ludden, P. W. Regulation of biological nitrogen fixation. *The Journal of nutrition* **130**, 1081-1084 (2000). https://doi.org:10.1093/jn/130.5.1081

Martin Del Campo, J. S. *et al.* Overview of physiological, biochemical, and regulatory aspects of nitrogen fixation in Azotobacter vinelandii. *Critical reviews in biochemistry and molecular biology* **57**, 492-538 (2022). https://doi.org:10.1080/10409238.2023.2181309

Leigh, J. A. & Dodsworth, J. A. Nitrogen regulation in bacteria and archaea. *Annual review of microbiology* **61**, 349-377 (2007). https://doi.org:10.1146/annurev.micro.61.080706.093409

Huergo, L. F. *et al.* PII signal transduction proteins: pivotal players in post-translational control of nitrogenase activity. *Microbiology* **158**, 176-190 (2012). https://doi.org:10.1099/mic.0.049783-0

Forchhammer, K., Selim, K. A. & Huergo, L. F. New views on PII signaling: from nitrogen sensing to global metabolic control. *Trends in microbiology* **30**, 722-735 (2022). https://doi.org:10.1016/j.tim.2021.12.014

Huergo, L. F., Chandra, G. & Merrick, M. P(II) signal transduction proteins: nitrogen regulation and beyond. *FEMS microbiology reviews* **37**, 251-283 (2013). https://doi.org:10.1111/j.1574-6976.2012.00351.x

Selim, K. A. & Alva, V. PII-like signaling proteins: a new paradigm in orchestrating cellular homeostasis. *Current opinion in microbiology* **79**, 102453 (2024). https://doi.org:10.1016/j.mib.2024.102453

Boyd, E. S. *et al.* A late methanogen origin for molybdenum-dependent nitrogenase. *Geobiology* **9**, 221-232 (2011). https://doi.org:10.1111/j.1472-4669.2011.00278.x

Boyd, E. S., Hamilton, T. L. & Peters, J. W. An alternative path for the evolution of biological nitrogen fixation. *Frontiers in microbiology* **2**, 205 (2011). https://doi.org:10.3389/fmicb.2011.00205

Deppenmeier, U. & Muller, V. Life Close to the Thermodynamic Limit: How Methanogenic Archaea Conserve Energy. *Results Probl Cell Differ* (2007).

Chanderban, M., Hill, C. A., Dhamad, A. E. & Lessner, D. J. Expression of V-nitrogenase and Fe-nitrogenase in Methanosarcina acetivorans is controlled by molybdenum, fixed nitrogen, and the expression of Mo-nitrogenase. *Applied and environmental microbiology* **89**, e0103323 (2023). https://doi.org:10.1128/aem.01033-23

Dhamad, A. E. & Lessner, D. J. A CRISPRi-dCas9 System for Archaea and Its Use To Examine Gene Function during Nitrogen Fixation by *Methanosarcina acetivorans*. *Applied and environmental microbiology* **86** (2020). https://doi.org:10.1128/AEM.01402-20

Lobo, A. L. & Zinder, S. H. Diazotrophy and Nitrogenase Activity in the Archaebacterium Methanosarcina barkeri 227. *Applied and environmental microbiology* **54**, 1656-1661 (1988).

Lobo, A. L. & Zinder, S. H. Nitrogenase in the archaebacterium Methanosarcina barkeri 227. *Journal of bacteriology* **172**, 6789-6796 (1990). https://doi.org:10.1128/jb.172.12.6789-6796.1990

Kessler, P. S., Daniel, C. & Leigh, J. A. Ammonia switch-off of nitrogen fixation in the methanogenic archaeon Methanococcus maripaludis: mechanistic features and requirement for the novel GlnB homologues, NifI(1) and NifI(2). *Journal of bacteriology* **183**, 882-889 (2001). https://doi.org:10.1128/JB.183.3.882-889.2001

Kessler, P. S. & Leigh, J. A. Genetics of nitrogen regulation in Methanococcus maripaludis. *Genetics* **152**, 1343-1351 (1999).

Ehlers, C., Veit, K., Gottschalk, G. & Schmitz, R. A. Functional organization of a single nif cluster in the mesophilic archaeon Methanosarcina mazei strain Go1. *Archaea* **1**, 143-150 (2002).

Dodsworth, J. A. & Leigh, J. A. Regulation of nitrogenase by 2-oxoglutarate-reversible, direct binding of a PII-like nitrogen sensor protein to dinitrogenase. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 9779-9784 (2006). https://doi.org:10.1073/pnas.0602278103

Dodsworth, J. A. & Leigh, J. A. NifI inhibits nitrogenase by competing with Fe protein for binding to the MoFe protein. *Biochemical and biophysical research communications* **364**, 378-382 (2007). https://doi.org:10.1016/j.bbrc.2007.10.020

Corless, E. I., Mettert, E. L., Kiley, P. J. & Antony, E. Elevated Expression of a Functional Suf Pathway in Escherichia coli BL21(DE3) Enhances Recombinant Production of an Iron-Sulfur Cluster-Containing Protein. *Journal of bacteriology* **202** (2020). https://doi.org:10.1128/JB.00496-19

Nayak, D. D. & Metcalf, W. W. Cas9-mediated genome editing in the methanogenic archaeon *Methanosarcina acetivorans*. *Proceedings of the National Academy of Sciences of the United States of America* **114**, 2976-2981 (2017). https://doi.org:10.1073/pnas.1618596114

Nayak, D. D. & Metcalf, W. W. Genetic techniques for studies of methyl-coenzyme M reductase from Methanosarcina acetivorans C2A. *Methods in enzymology* **613**, 325-347 (2018). https://doi.org:10.1016/bs.mie.2018.10.012

Metcalf, W. W., Zhang, J. K., Apolinario, E., Sowers, K. R. & Wolfe, R. S. A genetic system for Archaea of the genus *Methanosarcina*: liposome-mediated transformation and construction of shuttle vectors. *Proc. Natl. Acad. Sc.i USA* **94**, 2626-2631 (1997).

Kashyap, R. *et al.* Cryo-EM captures the coordination of asymmetric electron transfer through a di-copper site in DPOR. *Nat Commun* **16**, 3866 (2025). https://doi.org:10.1038/s41467-025-59158-7

Yildiz, O., Kalthoff, C., Raunser, S. & Kuhlbrandt, W. Structure of GlnK1 with bound effectors indicates regulatory mechanism for ammonia uptake. *The EMBO journal* **26**, 589-599 (2007). https://doi.org:10.1038/sj.emboj.7601492

Franke, P., Freiberger, S., Zhang, L. & Einsle, O. Conformational protection of molybdenum nitrogenase by Shethna protein II. *Nature* **637**, 998-1004 (2025). https://doi.org:10.1038/s41586-024-08355-3

Warmack, R. A. & Rees, D. C. Nitrogenase beyond the Resting State: A Structural Perspective. *Molecules* **28** (2023). https://doi.org:10.3390/molecules28247952

Muller, M. C. & Wagner, T. The Oxoglutarate Binding Site and Regulatory Mechanism Are Conserved in Ammonium Transporter Inhibitors GlnKs from Methanococcales. *International journal of molecular sciences* **22** (2021). https://doi.org:10.3390/ijms22168631

Maier, S. *et al.* Mechanism of disruption of the Amt-GlnK complex by P(II)-mediated sensing of 2-oxoglutarate. *PloS one* **6**, e26327 (2011). https://doi.org:10.1371/journal.pone.0026327

Radchenko, M. V., Thornton, J. & Merrick, M. P(II) signal transduction proteins are ATPases whose activity is regulated by 2-oxoglutarate. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 12948-12953 (2013). https://doi.org:10.1073/pnas.1304386110

Cuevas-Zuviría Bruno *et al.* Structural evolution of nitrogenase enzymes over geologic time *eLife* **14:RP105613** (2025). https://doi.org:https://doi.org/10.7554/eLife.105613.3

Zhang, L. M., Morrison, C. N., Kaiser, J. T. & Rees, D. C. Nitrogenase MoFe protein from Clostridium pasteurianum at 1.08 A resolution: comparison with the Azotobacter vinelandii MoFe protein. *Acta crystallographica. Section D, Biological crystallography* **71**, 274-282 (2015). https://doi.org:10.1107/S1399004714025243

Leigh, J. A. Nitrogen fixation in methanogens: the archaeal perspective. *Current issues in molecular biology* **2**, 125-131 (2000).

Rutledge, H. L., Cook, B. D., Nguyen, H. P. M., Herzik, M. A., Jr. & Tezcan, F. A. Structures of the nitrogenase complex prepared under catalytic turnover conditions. *Science* **377**, 865-869 (2022). https://doi.org:10.1126/science.abq7641