Co-isolation of genetically distinct *Burkholderia pseudomallei* strains from a single patient in North Queensland

Short title: Co-isolated *Bp* strains with distinct genomic, proteomic, and phenotypic profiles

Pauline M.L. Coulon[1], Piklu Roy Chowdhury[1], Ian Gassiep[3], Kay Ramsay[3], Aven Lee[2], Edita Ritmejeryte[2], Miranda E. Pitt[1], Joyce To[1], Sarah Reed[2], Patrick N. A. Harris[3] and Garry S.A. Myers[1]

Affiliations:

[1]Australian Institute for Microbiology and Infection, Faculty of Science, University of Technology Sydney, NSW, Australia

[2]Mass Spectrometry Facility, University of Queensland, Centre for Clinical Research, Queensland, Australia

[3]The University of Queensland Centre for Clinical Research (UQCCR), Faculty of Health, Medicine and Behavioural Sciences, The University of Queensland, Queensland, Australia

Corresponding author:

Pauline M.L. Coulon, Australian Institute for Microbiology and Infection, Faculty of Science, University of Technology Sydney, NSW, Australia, +61390353555,

pauline.coulon@uts.edu.au

Abstract

While investigating colony morphology variation (CMV) within clinical isolates of *Burkholderia pseudomallei* (*Bp*) isolated from a single infection, we identified two distinct strains (based on their multi-locus sequence type) from *Bp* TSV292. Simultaneous co-infection is rare, previously reported in only 2 of 133 cases. Here, we present the first comprehensive multi-omics characterization of co-infecting *Bp* strains isolated from a single infection at a hospital in Townsville, Australia in 2018. This finding impacts the design of future diagnosis and treatment of *Bp*.

Importance

Melioidosis, a severe disease caused by the bacterium *Burkholderia pseudomallei*, is rarely caused by more than one strain at the same time. In this study, we examined two bacterial strains isolated at once from a patient treated for melioidosis at a hospital in northern Queensland (Australia). By comparing their genomes, proteins, and physical traits, we found important strain differences that may affect how they adapt and survive during infection. Considering strain diversity when studying how this pathogen causes disease is essential, as it directly impacts future diagnosis and treatment.

Keywords: Melioidosis, diagnosis, colony morphotype variation, multi-omics

Introduction

Burkholderia pseudomallei (*Bp*), is an opportunistic pathogen and the causative agent of melioidosis, a severe and often fatal disease in humans and animals (1, 2). Previously endemic to tropical and subtropical regions of Asia, Africa, Central and South America, and

northern Australia (1), it is now recognized as established in the United States (3). In most

cases, *Bp* acquisition and infection will not lead to the development of melioidosis in healthy

populations. In people with underlying conditions, such as diabetes, chronic kidney or lung

disease, alcoholism, or compromised immune systems, melioidosis often manifests as skin

abscesses or severe pneumonia, which can progress to fatal sepsis with mortality rates

reaching up to 52% (4). With climate change intensifying extreme weather events such as

heavy rainfall and flooding, the incidence of *Bp* infections is projected to rise (5). In fact,

Queensland (Australia), has recently reported its worst year on record for melioidosis, with

over 239 confirmed infections and 31 deaths linked to dispersal of the pathogen during

flooding events (Queensland Health). While direct transmission between animals and humans

has not yet been confirmed, growing concerns have emerged regarding potential transmission

from pets, livestock or tropical animals to human (2), either through contact with infected

wounds or via shared exposure to contaminated environments (6). *Bp* is naturally highly

resistant to several antibiotic classes. Resistance to antibiotics used as primary treatment

(meropenem, ceftazidime and trimethoprim/sulfamethoxazole) is uncommon (7), however

emerging resistance against these drugs is increasingly described (8-11).

A key yet often overlooked feature of *Burkholderia* species is their ability to undergo colony

morphotype variation (CMV), a reversible process by which bacteria switch between distinct

colony phenotypes (12). This phenomenon reflects rapid adaptation to environmental

stressors and is often associated with changes in outer membrane proteins, antibiotic

production, and virulence factors (13-17). Mechanisms underlying CMV often include

mutations in global regulators, two-component systems, genome reduction and duplication,

bacteriophage integration, and DNA methylation (recently reviewed by Coulon et al. (12)).

Bp is known to exhibit up to seven different macrocolony morphologies which can be

modulated under laboratory conditions (13, 17, 18). Importantly, CMV is not limited to

adaptation *in vitro* as it also occurs during infection (19). Approximatively 10% of 450 *Bp*

clinical isolates produce mixed mucoid and non-mucoid CMV populations on blood agar

(19). In a long-term murine infection model, small colony variants (scv) were observed (20,

21). During human infection, differences in O-antigen lipopolysaccharides (LPS; [OPS])

production explains the mucoid phenotype (as observed on blood agar) even in the absence of

mutations or altered expression of the *wbiA* O-antigen acetylase (19). Additionally, genes

within the LPS biosynthesis cluster are upregulated during persistent infection in mice (20,

21). The upregulation of the σ-54 dependent regulator *yelR*, is involved in the emergence of

two reversible yellow *Bp* variants, which display enhanced resistance to hypoxic stress and

increased colonization and persistence in the murine stomach (22).

While investigating CMV within a cohort of *Bp* isolates (23) collected from clinical

melioidosis cases presented at Townsville University Hospital (Australia) (24), we identified

a polyclonal infection initially mistaken for colony morphotype variants. The apparent rough

and smooth 'variants' were in fact two genetically distinct strains isolated from the same

infection site. Polyclonal *Bp* infection is rarely reported, with an apparent occurrence of 1.5%

in a cohort of 133 patients (25); to date no integrated multi-omics analyses have been

performed at the bacterial level. Here, we present the first study combining genomic,

proteomic, and phenotypic analyses of two co-infecting *Bp* strains from a melioidosis patient

in Australia.

Materials and Methods

Ethical Approval

This study received ethical approval from the Royal Brisbane and Women's Hospital Ethics

Committee (LNR/2020/QRBW/65573), with site-specific authority obtained from the

Townsville Hospital and Health Service and approval under the Queensland Public Health

Act.

Colony morphology screening

Bp TSV292 was streaked on LB agar plates containing 0.01% Congo red (CRLA) from -

80°C stock and incubated at 37°C for two overnight periods.

Genomic analysis

gDNA extraction. Using either gDNA spin or high molecular weight gDNA extraction kits

(NEB), gDNA was extracted from multiple macrocolonies grown for three overnights at 37°C

on CRLA. Short-read sequencing (Illumina) was conducted on all CMVs. Briefly, libraries

were prepared using 1 ng of gDNA via the Nextera XT kit and sequenced using the NovaSeq

XPlus platform (35). On select isolates, long-read sequencing (Oxford Nanopore

Technologies) was implemented. Libraries were prepared using 400 ng gDNA with the SQK-

NBD114.24 kit and sequenced using R10.4.1 flow cells on the GridION platform. Reads

were base called using Dorado 7.6.7.

Genome assembly. Nanopore and Illumina reads were filtered and trimmed using respectively

nanofilt (v.2.8.0; (26)) and fastp (v0.24.1; (27)). Each set of Nanopore long reads were then

assemble using autocycler tool (v0.4.0; (28)). Next, the resulting long-read consensus

assemblies were polished with Illumina short reads using polypolish (v0.6.0; (29)) and then

pypolca (v3.0.1; (30)). Finally, the obtained assembly was annotated using batka (v1.11;

(31)). For the second CMV isolated from each isolate, Illumina short reads were assembled

using unicycler short read assembler (v.0.5.1; (32)) generating contigs.

Genomic modification analyses. Each set of Illumina reads, belonging to the second CMV,

were mapped on their respective first CMV assembly using snippy tools with haploid settings

(v4.6.0; (33) available on Galaxy) to determine SNPs and indels, to confirm the results, the

first CMV Illumina reads were also mapped onto their own assembly, resulting in showing no

genomic variation.

Genotyping and Phylogeny analysis. 6,259 genomic allelic profiles of 1,225 available *Bp*

isolates on pubMLST were retrieved and used to generate a distance phylogenetic tree with

GrapeTree using allelic profile differences (34). The resulting tree was generated using ggtree

R package. PhyloSift v1.0.1 was used to resolve subclade phylogeny of the two clusters

(identified from the wgMLST clustering analysis) by computing a maximum-likelihood tree

based on the genomes (available on pubMLST) using FastTree2.2 (35). FigTree v1.4.4 was

used to draw the unrooted phylogenetic trees and FastANI was used to identify the pairwise

average nucleotide identity with the respective ST identified in this analysis. BRIG-0.95 (36)

was used to map pairwise BLASTn results of 5 to 6 six genomes (37)- selected from the

PhyloSift phylogeny.

Prediction of antibiotic resistance genes. AMR gene orthologs present in genomes was

identified using the CARD database (38) and outputs were filtered to tabulate data of

candidates which had 100% coverage of subjects. Antibiotic Resistance Detection and

Prediction (ARDaP) tool was used to determined antibiotic resistance profile of both isolates

(39).

Whole genome alignment. Both MAUVE (40) and LASTZ (41, 42) tools were used to aligned

TSV292_1 (rough) and TSV292_2 (smooth) together, using the default settings. To correct

aligned genomes with LASTZ tool, the first residue numbers for each chromosome were

changed as followed, to set up the same origin. For TSV292_1 rough contig 2 residue

1,249,185 became 1 while TSV292_2 (smooth) contig 1 was reversed and residue 285,486

became 1; TSV292_2 (smooth) contig 2 residue 490,391 became 1.

Methylome analysis

Identification of genes encoding for DNA methyltransferase and their linked motifs. Pod5

files obtained from Nanopore sequencing were merged, using pod5 merge function and

analyzed   for   DNA   methylation   motifs   using   the   Dorado   with   both

dna_r10.4.1_e8.2_400bps_hac@v4.3.0_6mA@v2,

dna_r10.4.1_e8.2_400bps_hac@v4.3.0_5mC_5hmC@v1   models

(https://github.com/nanoporetech/dorado). To identify the location and number of detected

motifs, as well as the genes encoding DNA methyltransferases, the call_methylation and

annotate_rm functions from MicrobeMod (43) were used. These tools were provided with the

corresponding genome assembly ".fasta" and annotation ".gff3" files for each isolate.

Proteomic analysis

Protein extraction and peptide preparation. Bacterial cells in 1M Tris-HCL containing 1%

SDS were sonicated in an ultrasonic bath YJ5120-1 (Labtex) for 3 minutes, followed shaking

at 1,000 rpm at 95°C for 15 minutes on a Thermomixer Compact (Eppendorf). The mixture

was allowed to cool down to room temperature and 30 µL was mixed with 270 µL of Buffer I

(6 M guanidine chloride, 50 mM Tris pH 8.0, 10 mM DTT) and incubated at 30°C for 40

min. Following addition of 9 µL of 0.5 M iodoacetamide, the sample was incubated at 30°C

for 50 min. The reduced and alkylated protein sample was transferred to a 10 kDa cut-off

Amicon Ultra device (Merck) which was placed into a collection tube and centrifuged at

14,000 × g for 30 min. The flow-through from the column was removed and 300 µL of 100

mM ammonium bicarbonate was added to the column followed by centrifugation at 14,000 ×

g for 30 min. To digest proteins in the column, 130 µl of 100 mM ammonium bicarbonate

containing 2 µg Trypsin was added. The column was incubated at 37°C for 24 h and then

centrifuged in a new collection tube at 14,000 × g for 30 min. Fifty µL of water with 0.1%

formic acid was then added to the spin column followed by centrifugation at 14,000 × g for

30 min to elute the remaining digested peptides. Enzymes and salts were removed using C_{18}

ZipTips (Merck) following the manufacturer's instructions and desalted peptides were eluted

with 80% acetonitrile/0.1% formic acid into microcentrifuge tubes. Digested peptide samples

were dried down using a vacuum concentrator (Concentrator plus, Eppendorf) and

reconstituted with 50 µL water with 0.1% formic acid prior to LC-MS/MS analysis.

LC-MS/MS analysis. The samples were chromatographically separated and resulting

proteomics data was acquired using an Acquity M-class micro-LC system (Waters, USA)

coupled with ZenoTOF 7600 LC-MS/MS system using the OptiFlow Turbo V ion source.

(AB Sciex). Two µL of sample were loaded onto Micro TRAP C18 column (0.3 × 10 mm;

Phenomenex) and washed for 10 min at 10 µL/min with 97% solvent A (water with 0.1%

[v/v] formic acid) and 3% solvent B (99.9% [v/v] acetonitrile, 0.1% [v/v] formic acid).

Liquid chromatography was performed at 5 µL/min using a 2.7 µm Peptide C18 160Å; (0.3 ×

150 mm; Bioshell) column with the column oven temperature maintained at 40°C. The

gradient started at 3% solvent B which was increased to 5% by 0.5 min, followed by an

increase to 35% solvent B by 38.9 min and then another increase to 80% solvent B by 39

min. The column was washed with 80% solvent B for 3.9 min before equilibrating with 3%

solvent B from 43 to 45 min. Data was acquired using Zeno SWATH DIA, using a 65

variable width SWATH windows. The TOF MS parameters were as follows: precursor mass

range was 400 – 1500 m/z; declustering potential (DP) was set to 80 V and accumulation

time was set at 0.25 s. The TOF MS/MS parameters were as follows: fragmentation mass

range was set to 100 – 1500 m/z; with an accumulation time of 0.025s; fragmentation mode

was set to CID with Zeno pulsing selected. The collision energy for the MS/MS acquisition

was automatically adjusted according to the m/z and charge of the peptide.

Data analyses. Raw label free quantification (LFQ) LC-MS/MS data were converted to

.mzML files, using MS convert, to allow the data search using Fragpipe v22 (44). Protein

identification was conducted against the respective annotated proteomes using the

DIA_SpecLib_Quant workflow, which builds a spectral library with MSFragger-DIA and

performs quantification with DIA-NN. The output "diann-output.pg_matrix.tsv" was filtered

to remove contaminants and retain proteins with at least two non-missing values for at least

one group. Data was normalized using the variance stabilization normalization (45). Proteins

with two or more missing values in one group were categorized as "detected/not detected."

For proteins with a single missing value in one group, missing data were imputed using the

mean of the other replicates in that group. Differential protein expression between groups was

assessed using Welch's t-test, with Benjamini–Hochberg correction applied to control the

false discovery rate (adjusted $p < 0.05$) (46, 47). Analyses were performed on three biological

replicates (Table S5-9).

Proteomic representation. To assign homologous proteins between TSV292 rough, smooth

and K96243 (*Bp* model), a megablastp analysis was set up with only one output match for

each gene in query using the following parameters: -evalue 10 -outfmt 6 -num_threads 3 -

max_target_seqs 1 -max_hsps 1 (37). Clustering of orthologous groups (COGs) for annotated

proteins were determined using EggNOG tool v5 ((48)). To represent the data in a volcano

plot, detected/not detected data were imputed using the lower normalized Log2 value for

intensity with an imputed adjusted pvalue randomly assigned between 0.01 and 0.00001.

Enrichment analysis was done using a Fisher exact test based on the COGs, using Claude ai

for the script (REF).

Phenotypic assays

Production of virulence factors. Individual colonies (four biological replicates) were picked

and stabbed into LB agar plate containing 1.5% skimmed milk (assessing proteolytic activity;

(49)), Cas assay plates made with 10X chelated LB (23); 0.5 g/L yeast extract supplemented

with 4g/L D-mannitol (YEM) and agar (assessing mucoidy; (50)) and Columbian horse blood

agar plate (assessing hemolysis activity; (51)). Plates were incubated at 37°C for 3 days.

Inhibition circles were measured and confirmed by statistical analyses.

Antimicrobial sensitivity assay. Individual colonies were resuspended in 0.9% NaCl solution

to obtain a turbidity of 0.5-0.63 McFarland. Bacterial suspensions were then spread onto

Muller Hinton II agar plates and incubated overnight at 37°C with 0.002-32 µg/mL

Meropenem (MEM) ETEST (Biomérieux), 10µg Ceftazidime (CAZ) and 25µg

Sulphamethoxazole-Trimethoprim (TMP-SMX) Antimicrobial Susceptibility discs (Oxoid).

Antimicrobial susceptibility was determined by following the recommendation from

EUCAST   (v15;

https://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/v_15.0_

Breakpoint_Tables.pdf). For CMVs from the same isolate showing considerable visual

changes in antibiotic susceptibility, three biological replicates were conducted on independent

days, and the average of inhibition data were used to determine their susceptibility

characteristic.

Data availability

Genomes from this study are available in GenBank under the Bioproject: PRJNA1295108

(52, 53). Proteomic raw data deposited to the ProteomeXchange Consortium via the PRIDE

partner repository with the dataset identifier PXD066041 (54).

Results

Isolation of TSV292_1 rough and TSV292_2 smooth colony morphotypes

In *Burkholderia,* CMV typically occurs upon removal of environmental pressure or during

aging (13, 17). Because selective media for *Burkholderia pseudomallei* often contain

antibiotics that can induce colony morphology changes (e.g., the wrinkled phenotype in *Bp*),

the clinical isolate TSV292 was streaked directly onto LB agar supplemented with 0.1%

congo red (CRLA) from a −80 °C glycerol stock to facilitate CMV distinction. On this

medium, a mixed population was observed, consisting of flat colonies with a red center (type

RI) and smoother, raised colonies (type SI) (**Fig. 1A**). The rough and smooth phenotypes

were confirmed by isolating single colonies and subculturing them on CRLA or Ashdown

agar (**Fig. 1B**). Both media yielded CMV types consistent with those previously reported for

isolate TSV82 (23).

Polyclonal strains masquerading as CMV in melioidosis infection

Genomic variation, master regulator expression or epigenetics are known drivers of CMV in

Burkholderia; hence, we first investigated whether SNPs or insertion and deletions (indels)

accounted for the distinct morphologies of TSV292. Illumina reads from TSV292 rough were

mapped against the assembled genome of TSV292 smooth, using Snippy (33), which

revealed 23,229 genomic modifications (**Table S1**). This unexpectedly high level of variation

suggested that the two morphotypes were polyclonal rather than colony morphotype variants

of the same isolate. Multi-locus sequence typing (MLST) further supported this conclusion:

the rough isolate was identified as ST-2319 (TSV292_1; PubMLST sample ID:7631;

previously reported by Gassiep and colleagues (24)), whereas the smooth isolate represented

a novel sequence type (TSV292_2; PubMLST sample ID:7519; ST-2323) (**Table S2**). As

simultaneous infection with multiple distinct *Bp* clones from the same patient is considered

rare (25), we further investigated the genomic and proteomic profiles of these isolates to

better understand their divergence.

Comparative analysis of *Bp* ST-2319 and ST-2323:

The average nucleotide identity (ANI) between the two genomes was 99.4%. TSV292_1

(rough) had a genome size of 7,262,464 bp with a GC content of 68.1%, while TSV292_2

(smooth) had a genome size of 7,352,845 bp with a GC content of 68.0%. Clustering analysis

of whole-genome MLST (wgMLST) based on 6,259 allelic profiles of 1,225 available

isolates in the PubMLST database (**Table S3**; as of August 2025) revealed two distinctly

separate groups (**Fig. 2A**). Maximum-likelihood phylogenetic analysis showed that

TSV292_1 rough clustered with *Bp* strains previously isolated from melioidosis patients

admitted at Townsville University Hospital between 1997 and 2020 (24), whereas TSV292_2

smooth was more closely related to *Bp* strains isolated in Cambodia (**Fig. 2B**). TSV292_1

rough and TSV292_2 smooth shared >99.2% ANI with their closely related *Bp* isolates but

also contain variable regions representing strain-specific regions (**Fig. S1**).

Whole-genome alignment of the TSV292 co-isolates, using MAUVE algorithm (40),

revealed synteny across the locally co-linear blocks and highlight genomic rearrangements

likely as a result of recombination events in both chromosomes (**Fig. S2**). Alignment using

the LASTZ tool (41, 42), further revealed divergence between TSV292_1 rough and

TSV292_2 smooth in several regions (**Fig. S3–4**).

In TSV292_1 rough, regions absent in TSV292_2 smooth included: (i) a 7,465 bp segment

containing a type II restriction enzyme and DNA Methyltransferase (DNA MTase) predicted

to be responsible for m5C methylation on GTCGAC pattern (TSV292_R_RS_00500-00501)

surrounded by IS3 family transposase; (ii) a 36,612 bp region including predicted phage

elements and a singleton DNA MTase (TSV292_R_RS_02319); and (iii) a 61,103 bp region

with 26 putative transposases consistent with lateral gene transfer, that also encoded the *Bp*

adhesion (*bpa*) gene cluster—implicated in bacterial attachment and immune evasion (55) —

suggesting functional differences in host interaction and pathogenic potential.

On chromosome 2 TSV292_1 rough carried a 43,430 bp region with an integrase

(TSV292_R_RS_04679) and phages elements; a 36,084 bp region rich in transposases and

viral genes, including a type III restriction DNA methyltransferase (TSV292_R_RS_03252;

Table S7); a 21,283 bp region containing a putative origin of replication with a PAAR gene

and a type III restriction DNA MTase (TSV292_R_RS_03525) suggestive of plasmid

integration; and a 43,340 bp region containing phage elements (**Tables S4 and S5**).

By contrast, TSV292_2 smooth lacked multiple phage-associated and transposon-rich regions

present in TSV292_1 rough, including loci encoding type II, III, and IV restriction MTases

predicted to modify m4C, m5C, and m6A motifs (e.g., TSV292_RS_03205,

TSV292_RS_03345, TSV292_RS_15530, TSV292_RS_15535, TSV292_RS_18580,

TSV292_RS_26185; Tables S4, S6–S7). Differences also extended to the flagellar

biosynthesis cluster, where the two genomes displayed altered gene order (Fig. S5).

Taken together, these results indicate that divergence between both strains involved

acquisition or loss of phage elements, transposons, plasmid-associated regions, and DNA

methylation systems. Such changes not only underscore the genome plasticity of *Bp* but also

suggest functional consequences for immune evasion, horizontal gene transfer, and regulation

of surface structures critical for infection.

Distinct phenotypes in antibiotic sensitivity, mucoidy, hemolysis, siderophore production

and proteolytic activity

To confirm the proteomic analyses, phenotypic assays assessing virulence-related traits,

including antibiotic susceptibility were performed (**Table 1**), revealing notable differences

between the two isolates. TSV292_2 smooth exhibited a higher resistance to trimethoprim-

sulfamethoxazole (TMP-SMX), with no apparent difference in predicted antibiotic resistance

by ARDaP (**Table S8-9**; (39)), and displayed increased mucoidy, consistent with higher

exopolysaccharide (EPS) production when grown on YEM agar. In contrast to TSV292_1

rough, the TSV292_2 smooth strain lost its ability to lyse red blood cells and lacked

proteolytic activity. Interestingly, both isolates produced a smooth-edged sector around the

main colony when grown on Ashdown and blood agar, suggesting an active role in CMV

emergence (**Fig. 3**).

Table 11 Phenotypic profile of both *Bp* TSV292_1 rough and TSV292_2 smooth

Isolate	Colony morphology	Antibiotic Susceptibility test			Mucoidy	Hemolysis	Proteolytic Activity (cm^2)	Siderophore production (cm^2)
		MEM	CAZ	TMP-SMX				
TSV 292	Strain 1 (rough)	S	I	I	+	beta	3.26 ± 0.75 **	0.399±0.093**
	Strain 2 (smooth)	S	I	R^a	++	gamma	0.88 ± 0.09	1.811±0.33

EUCAST: S = Susceptible; I= Susceptible increased exposure; R= Resistant; Meropenem (MEM): S≤2; R>2; Ceftazidime (CAZ; disk

10µg): S≥50mm; R<18mm; Trimethoprim-Sulphamethadoxone (TMP-SMX; disk 1.25-23.75µg): S≥50mm; R<17mm;

"a" represent a TMP-SMX susceptibility statistically different between TSV292_1 and TSV292_2 using a ttest;

Mucoidy: + = mucoid; ++ = more mucoid;

Hemolysis at 8 days: beta = complete lysis of red blood cell; gamma = no lysis;

Proteolytic activity and Siderophore production: average ± stand deviation;

 "**" represent statistical significance (ttest) in both proteolytic activity or siderophore production with a pvalue comprised between 0.05 and

 0.01 when comparing both TSV292_1 and TSV292_2.

Proteomics confirms profile respectively belonging to rough and smooth *Bp* CMV

morphotypes despite being two isolates

Proteomic profiling confirmed that the two isolates correspond to the expected profiles of

rough and smooth *Bp* morphotypes, with differences in virulence- and resistance-associated

pathways. In total, 1,015 proteins were differentially produced between the two isolates (**Fig.**

3A; Table S10–11).

In the TSV292_2 smooth strain, proteins associated with antimicrobial gene cluster

biosynthesis were more abundant or exclusively detected. These included those involved in

the production of pyochelin, malleobactin, malleilactone, bactobolin, and syrbactins (56-58).

Notably SyrC/GlbC, part of the syrbactin cluster, is also known to influence *Bp* survival and

replication within immune cells (56, 59). Additionally, proteins related to flagellar

biosynthesis, the quorum sensing system 2 (BpsR2; belonging to the bactobolin biosynthesis

cluster), and the Hmq system, which produces HMAQs with antimicrobial activity and

known interaction with QS systems, were more abundant or exclusively detected (58, 60, 61).

In contrast, the TSV292 rough morphotype showed higher abundance or exclusive detection

of proteins involved in exopolysaccharide (EPS) production, type III O-polysaccharide (OPS)

synthesis, DNA methyltransferase (MTase) responsible for the CACAG methylation motif,

the quorum sensing system 3 (BpsI3/BpsR3), and the master regulator ScmR (**Fig. 3B**).

Proteins from antibiotic resistance genes (11, 39, 62, 63), predicted using the CARD database

(**Table S12;** (38)), were generally not detected or detected at lower abundance in TSV292_2

smooth compared with the rough isolate (**Table 2**). This includes both AmrR, the TetR family

transcriptional regulator that controls the expression of the AmrAB-OprA efflux pump, and

BpeT, the transcriptional regulator of the BpeEF-OprC efflux pump, potentially explaining

the observed TMP-SMX resistance in the smooth strain in Table 1 (39).

Together these results indicate a difference in adaptations between the two strains, with

TSV292_2 smooth adopting an environmental-like smooth form and TSV292_1 rough

displaying   a   pathogenic   rough   form   (15,   58).

Table 2: Significance differences in the abundance of protein involved in antibiotic resistance

Protein in *Bp* K96243	Protein in TSV292 R	Protein in TSV292 S	Gene Name	Product	Mean intensity in TSV292_1 rough (Log2)	Mean intensity in TSV292_2 smooth (Log2)	Differential abundance TSV292_1 rough vs TSV292_2 smooth (Log2)	pvalue (ttest)	Adjusted pvalue (fdr=0.01)	Significance	Resistance	Reference
BPSL1440	RS_05656	RS_24985		hypothetical protein	14.4	12.2	-2.178	0.007	0.019	Lower abundance in rough	SXT	(62)
BPSL1357	RS_04972	RS_21485	*folP*	dihydropteroate synthase	11.3	13.0	1.729	0.001	0.006	Lower abundance in rough	SXT	(63)
BPSS0946	RS_02455	RS_12455	*penA*	beta-lactamase precursor	12.5	14.4	1.903	0.001	0.005	Lower abundance in rough	CAZ	(62)
BPSL0814	RS_04343	RS_18740	*bpeA*	RND family acriflavine resistance protein A	13.9	15.7	1.778	0.000	0.002	Lower abundance in rough	MEM	(62)
BPSS1898	RS_01406	RS_02545	*fadH*	2,4-dienoyl-CoA reductase	NA	9.1	NA	NA	NA	Not detected in rough	MEM	(62)
BPSL1805	RS_05301	RS_23130	*amrR*	TetR family regulatory protein	NA	10.0	NA	NA	NA	Not detected in rough	MEM	(11, 62)
BPSS0290	RS_00441	RS_08815	*bpeT*	LysR family transcriptional regulator	NA	10.9	NA	NA	NA	Not detected in rough	MEM/CAZ/SXT	(11, 39, 62)
BPSS1269	RS_02119	RS_00900	*syrC/glbC*	peptide synthase/polyketide synthase	NA	12.5	NA	NA	NA	Not detected in rough	AMC/CAZ	(62)
BPSS1095	RS_02298	RS_13015		heat-shock chaperone protein	NA	13.0	NA	NA	NA	Not detected in rough	SXT	(62)
BPSS1654	RS_01708	RS_01070		cytochrome P450	9.4	NA	NA	NA	NA	Not detected in smooth	AMC	(62)
BPSL0812	RS_04342	RS_18735	*bpeR*	TetR family regulatory protein	10.5	NA	NA	NA	NA	Not detected in smooth	MEM	(11)
BPSS2246	RS_01043	RS_06515		sensor kinase/response regulator fusion protein	11.2	NA	NA	NA	NA	Not detected in smooth	SXT	(62)
BPSS1197	RS_02196	RS_00505		non-ribosomal peptide synthase	12.3	NA	NA	NA	NA	Not detected in smooth	CAZ/MEM/SXT	(62)
BPSL0816	RS_04346	RS_18755	*oprB*	outer membrane efflux protein	13.6	NA	NA	NA	NA	Not detected in smooth	CAZ	(62)
BPSL2553	RS_02674	RS_28050		siderophore receptor protein	10.5	13.4	2.850	0.007	0.018	Lower abundance in smooth	CAZ/MEM	(62)
	RS_01304	RS_05215	*blaOXA*	OXA-42 family class D beta-lactamase	NA						TET	

376 *NA: not   applicable

377 **Methylome showed no overall difference in methylated sites between TSV292 rough and**

378 **smooth**

379 Predicted restriction enzyme methylated motifs were consistent with the one described by

380 Nandi et al. (64) (**Table S7&13**). Given the observed difference in abundance of the DNA

381 MTase responsible for the CACAG methylation motif (TSV292_R_RS_03660/03661 in

382 rough colony morphotype TSV292_RS_15080/15085 in the smooth, for which proteins were

383 not detected in TSV292_2 smooth while detected in TSV292_1 rough isolates), we further

384 investigated the methylome profiles of both isolates (**Table S13**). Despite the presence of

385 distinct abundance of DNA MTases, no significant differences were observed in methylated

386 sites across the genomes of TSV292 rough and smooth for motifs CACAG, GTWWAC and

387 CATCAG.

388

389

390 # Discussion

391 Melioidosis causes an estimated 179,000 cases and 90,000 deaths worldwide each year (4).

392 As *Bp* is intrinsically resistant to antibiotics, treatment relies on last-resort agents. Emerging

393 resistance remains a major concern (8-10). While simultaneous polyclonal *Bp* infections are

394 considered rare (25), here we report the first such case in Queensland, isolated from a patient

395 swab in 2018.

396 A key diagnostic challenge revealed by this study is that polyclonality can be mistaken for

397 CMV. Current workflows, typically single-colony purification with selection of a typical

398 wrinkled morphotype followed by MALDI-TOF or sequencing, cannot reliably distinguish

399 between true CMV and mixed-strain infections. This diagnostic blind spot risks

400 underestimating strain diversity and may lead to inappropriate clinical decisions and

401 treatment failures (12). Because *Burkholderia* species exhibit extensive colony morphotype

plasticity with a particular mucoid/wrinkled colony morphotype (13, 17, 19), relying on

phenotypic assessment alone can mask genetically distinct pathogens. In this case, two

morphotypes that appeared as rough and smooth variants on Congo red agar and

mucoid/wrinkled on Ashdown selective media were in fact distinct strains.

We also investigated DNA methylation, given its known role in CMV and intraspecies

recombination (65). While global methylation patterns appeared similar between the isolates,

the absence of detectable DNA MTases by LC-MS/MS in the smooth morphotype mirrored

our prior findings in smooths CMVs (23). This could reflect biological differences in MTase

stability or abundance but may also stem from limitations of single-replicate methylome

analysis in detecting subtle variation.

By integrating genomic, proteomic, and phenotypic analyses, we demonstrate how distinct

co-infecting strains can mimic CMV, with smooth and rough colony morphotypes differing in

the production of proteins involved in virulence, stress tolerance, drug responses and

adaptation—features with direct implications for patient management. Importantly, outbreak

tracing and phylogenomic inference are highly vulnerable to such hidden co-infection; clonal

verification should therefore precede any evolutionary or epidemiological analyses of *Bp*.

Our findings suggest that covert polyclonality, rather than ultrafast evolution, is a major

driver of observed genomic diversity in this pathogen. High-quality, closed genomes from

long-read assemblies, combined with read-level deconvolution using short-read sequencing,

should be standard practice before calling variants.

Ultimately, this work highlights the urgent need for strain-level diagnostic approaches that

integrate multi-omics and phenotypic data. Building comprehensive reference databases of

such profiles will enable predictive tools to guide diagnosis, treatment, and surveillance.

These advances will not only improve patient outcomes but also yield more accurate

epidemiological insights, deepening our understanding of *Bp* biology.

Acknowledgment

This work was supported by AIMI and UTS Faculty of Science seed funds to Dr Pauline

M.L. Coulon, UTS Strategic Research Accelerator funding (SRA 2726229) to Prof Garry

S.A. Myers and EL2 Investigator grant from the NHMRC (APP2033851) to Dr Patrick N.A.

Harris. Experimental work was done at UQCCR which would not have been possible without

the ISME scholar funding awarded to Dr Pauline M.L. Coulon.

P.M.L.C: Concept and visualization, Funding, Experimental work; K.M: Experimental work

and reviewing, P.R.C: Genomic comparison; M.P and J.T: DNA libraries and sequencing;

A.L, E.R and S.R: sample preparation and LCMS runs for the proteomic experiment; I.G &

P.N.A.H: Reviewing and Funding; G.S.A.M: Reviewing and Funding.

References

1. Meumann EM, Limmathurotsakul D, Dunachie SJ, Wiersinga WJ, Currie BJ. 2024. *Burkholderia pseudomallei* and melioidosis. Nat Rev Microbiol 22:155-169.

2. Sprague LD, Neubauer H. 2004. Melioidosis in animals: a review on epizootiology, diagnosis and clinical presentation. J Vet Med B Infect Dis Vet Public Health 51:305-20.

3. Petras JK, Elrod MG, Ty MC, Dawson P, O'Laughlin K, Gee JE, Hanson J, Boutwell C, Ainsworth G, Beesley CA, Saile E, Tiller R, Gulvik CA, Ware D, Sokol T, Balsamo G, Taylor K, Salzer JS, Bower WA, Weiner ZP, Negrón ME, Hoffmaster AR, Byers P. 2023. Locally Acquired Melioidosis Linked to Environment - Mississippi, 2020-2023. N Engl J Med 389:2355-2362.

4. Limmathurotsakul D, Golding N, Dance DA, Messina JP, Pigott DM, Moyes CL, Rolim DB, Bertherat E, Day NP, Peacock SJ, Hay SI. 2016. Predicted global distribution of *Burkholderia pseudomallei* and burden of melioidosis. Nat Microbiol 1.

5. Currie BJ, Jacups SP. 2003. Intensity of rainfall and severity of melioidosis, Australia. Emerg Infect Dis 9:1538-42.

6. Inglis TJ, Garrow SC, Adams C, Henderson M, Mayo M, Currie BJ. 1999. Acute melioidosis outbreak in Western Australia. Epidemiol Infect 123:437-43.

7. Dance DA, Davong V, Soeng S, Phetsouvanh R, Newton PN, Turner P. 2014. Trimethoprim/sulfamethoxazole resistance in *Burkholderia pseudomallei*. Int J Antimicrob Agents 44:368-9.

8. Sarovich DS, Price EP, Limmathurotsakul D, Cook JM, Von Schulze AT, Wolken SR, Keim P, Peacock SJ, Pearson T. 2012. Development of ceftazidime resistance in an acute *Burkholderia pseudomallei* infection. Infect Drug Resist 5:129-32.

9. Sivabalan P, Satyaputra F, Gassiep I, Forde B, Frazer J, Glover M, Adams B, Norton R. 2024. Meropenem-resistant *Burkholderia pseudomallei*: a concerning single case in Australia with no prior meropenem exposure. Access Microbiol 6.

10. Wu PH, Chen CH, Hsih WH, Chou CH, Chi CY, Ho MW, Lin YT, Lin HH, Tseng KH, Hsueh PR. 2023. Emergence of meropenem and levofloxacin resistance in *Burkholderia pseudomallei* in Taiwan. J Infect 86:e161-e163.

11. Sarovich DS, Webb JR, Pitman MC, Viberg LT, Mayo M, Baird RW, Robson JM, Currie BJ, Price EP. 2018. Raising the Stakes: Loss of efflux pump regulation decreases meropenem susceptibility in *Burkholderia pseudomallei*. Clin Infect Dis 67:243-250.

12. Coulon PML, Agnoli K, Myers GSA. 2025. Colony morphotype variation in *Burkholderia*: implications for success of applications and therapeutics. Journal of Bacteriology 207:e00521-24.

13. Chantratita N, Wuthiekanun V, Boonbumrung K, Tiyawisutsri R, Vesaratchavest M, Limmathurotsakul D, Chierakul W, Wongratanacheewin S, Pukritiyakamee S, White NJ, Day NPJ, Peacock SJ. 2007. Biological Relevance of Colony Morphology and Phenotypic Switching by *Burkholderia pseudomallei*. Journal of Bacteriology 189:807--817.

14.   Coulon PML, Groleau M-C, Hachani A, Padula MP, Stinear TP, Déziel E.
2024. Quorum sensing and DNA methylation play active roles in clinical
Burkholderia phase variation. Journal of Bacteriology bioRxiv 207:e00531-24.
15.   Vial L, Groleau M-C, Lamarche MG, Filion G, Castonguay-Vanier J, Dekimpe
V, Daigle F, Charette SJ, Déziel E. 2009. Phase variation has a role in
Burkholderia ambifaria niche adaptation. The ISME Journal 4:49.
16.   Mariappan V, Barathan M, Zulpa AK, Vadivelu J, Vellasamy KM. 2023. Small
colony variants of *Burkholderia pseudomallei*: alteration of the virulence
factors. Journal of Taibah University for Science 17:2244657.
17.   Häußler S, Rohde M, Steinmetz I. 1999. Highly resistant *Burkholderia*
pseudomallei small colony variants isolated in vitro and in experimental
melioidosis. Medical Microbiology and Immunology 188:91--97.
18.   Chantratita N, Tandhavanant S, Wikraiphat C, Trunck LA, Rholl DA, Thanwisai
A, Saiprom N, Limmathurotsakul D, Korbsrisate S, Day NPJ, Schweizer HP,
Peacock SJ. 2012. Proteomic analysis of colony morphology variants of
Burkholderia pseudomallei defines a role for the arginine deiminase system in
bacterial survival. Journal of Proteomics 75:1031--1042.
19.   Wikraiphat C, Saiprom N, Tandhavanant S, Heiss C, Azadi P, Wongsuvan G,
Tuanyok A, Holden MTG, Burtnick MN, Brett PJ, Peacock SJ, Chantratita N.
2015. Colony morphology variation of *Burkholderia pseudomallei* is
associated with antigenic variation and o-polysaccharide modification.
Infection and Immunity 83:2127--2138.
20.   Bernhards RC, Cote CK, Amemiya K, Waag DM, Klimko CP, Worsham PL,
Welkos SL. 2017. Characterization of in vitro phenotypes of *Burkholderia*
pseudomallei and *Burkholderia mallei* strains potentially associated with
persistent infection in mice. Archives of Microbiology 199:277--301.
21.   Al-Maleki AR, Vellasamy KM, Mariappan V, Venkatraman G, Tee TS, Vadivelu
J. 2019. Transcriptome analysis of *Burkholderia pseudomallei* SCV reveals an
association with virulence, stress resistance and intracellular persistence.
Genomics 112:501--512.
22.   Austin CR, Goodyear AW, Bartek IL, Stewart A, Sutherland MD, Silva EB,
Zweifel A, Vitko NP, Tuanyok A, Highnam G, Mittelman D, Keim P, Schweizer
HP, Vázquez-Torres A, Dow SWC, Voskuil MI. 2015. A *Burkholderia*
pseudomallei colony variant necessary for gastric colonization. mBio
6:e02462--14.
23.   Coulon PML, Ramsay K, Lee A, Ritmejeryte E, Pitt ME, To J, Mediati DG,
Reed S, Harris PNA, Myers GSA. 2025. Profiling of *Burkholderia*
pseudomallei variants derived from Queensland's clinical isolates. bioRxiv
doi:10.1101/2025.08.10.669548:2025.08.10.669548.
24.   Gassiep I, Chatfield MD, Permana B, Burnard D, Bauer MJ, Cuddihy T, Forde
BM, Mayer-Coverdale J, Norton RE, Harris PNA. 2024. The Genomic
Epidemiology of Clinical *Burkholderia pseudomallei* Isolates in North
Queensland, Australia. Pathogens 13:584.
25.   Limmathurotsakul D, Wuthiekanun V, Chantratita N, Wongsuvan G, Thanwisai
A, Biaklang M, Tumapa S, Lee S, Day NP, Peacock SJ. 2007. Simultaneous
infection with more than one strain of *Burkholderia pseudomallei* is
uncommon in human melioidosis. J Clin Microbiol 45:3830-2.
26.   De Coster W, D'Hert S, Schultz DT, Cruts M, Van Broeckhoven C. 2018.
NanoPack: visualizing and processing long-read sequencing data.
Bioinformatics 34:2666-2669.

27. Chen S. 2023. Ultrafast one-pass FASTQ data preprocessing, quality control,
and deduplication using fastp. iMeta 2:e107.
28. Wick RR, Howden BP, Stinear TP. 2025. Autocycler: long-read consensus
assembly for bacterial genomes. bioRxiv
doi:10.1101/2025.05.12.653612:2025.05.12.653612.
29. Bouras G, Judd LM, Edwards RA, Vreugde S, Stinear TP, Wick RR. 2024.
How low can you go? Short-read polishing of Oxford Nanopore bacterial
genome assemblies. Microbial Genomics 10.
30. Zimin AV, Salzberg SL. 2020. The genome polishing tool POLCA makes fast
and accurate corrections in genome assemblies. PLOS Computational
Biology 16:e1007981.
31. Schwengers O, Jelonek L, Dieckmann MA, Beyvers S, Blom J, Goesmann A.
2021. Bakta: rapid and standardized annotation of bacterial genomes via
alignment-free sequence identification. Microb Genom 7.
32. Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: Resolving bacterial
genome assemblies from short and long sequencing reads. PLoS Comput
Biol 13:e1005595.
33. Seemann T. 2015. Snippy: rapid haploid variant calling and core SNP
phylogeny. GitHub Available at: github com/tseemann/snippy.
34. Zhou Z, Alikhan N-F, Sergeant MJ, Luhmann N, Vaz C, Francisco AP, Carriço
JA, Achtman M. 2018. GrapeTree: visualization of core genomic relationships
among 100,000 bacterial pathogens. Genome Research 28:1395 1404.
35. Price MN, Dehal PS, Arkin AP. 2010. FastTree 2--approximately maximum-
likelihood trees for large alignments. PLoS One 5:e9490.
36. Alikhan N-F, Petty NK, Ben Zakour NL, Beatson SA. 2011. BLAST Ring Image
Generator (BRIG): simple prokaryote genome comparisons. BMC Genomics
12:402.
37. Johnson M, Zaretskaya I, Raytselis Y, Merezhuk Y, McGinnis S, Madden TL.
2008. NCBI BLAST: a better web interface. Nucleic Acids Research 36:W5
W9.
38. Alcock BP, Huynh W, Chalil R, Smith KW, Raphenya AR, Wlodarski MA,
Edalatmand A, Petkau A, Syed SA, Tsang KK, Baker SJC, Dave M, McCarthy
MC, Mukiri KM, Nasir JA, Golbon B, Imtiaz H, Jiang X, Kaur K, Kwong M,
Liang ZC, Niu KC, Shan P, Yang JYJ, Gray KL, Hoad GR, Jia B, Bhando T,
Carfrae LA, Farha MA, French S, Gordzevich R, Rachwalski K, Tu MM,
Bordeleau E, Dooley D, Griffiths E, Zubyk HL, Brown ED, Maguire F, Beiko
RG, Hsiao WWL, Brinkman FSL, Van Domselaar G, McArthur AG. 2023.
CARD 2023: expanded curation, support for machine learning, and resistome
prediction at the Comprehensive Antibiotic Resistance Database. Nucleic
Acids Res 51:D690-d699.
39. Madden DE, Webb JR, Steinig EJ, Currie BJ, Price EP, Sarovich DS. 2021.
Taking the next-gen step: Comprehensive antimicrobial resistance detection
from Burkholderia pseudomallei. EBioMedicine 63:103152.
40. Darling AC, Mau B, Blattner FR, Perna NT. 2004. Mauve: multiple alignment
of conserved genomic sequence with rearrangements. Genome Res 14:1394-
403.
41. Harris RS. 2007. Improved pairwise alignment of genomic DNA. The
Pennsylvania State University.

42.   Schwartz S, Kent WJ, Smit A, Zhang Z, Baertsch R, Hardison RC, Haussler
583   D, Miller W. 2003. Human–mouse alignments with BLASTZ. Genome
584   research 13:103-107.
585   43.   Crits-Christoph A, Kang SC, Lee HH, Ostrov N. 2023. MicrobeMod: A
586   computational toolkit for identifying prokaryotic methylation and restriction-
587   modification   with   nanopore   sequencing.   bioRxiv
588   doi:10.1101/2023.11.13.566931:2023.11.13.566931.
589   44.   Kong AT, Leprevost FV, Avtonomov DM, Mellacheruvu D, Nesvizhskii AI.
590   2017. MSFragger: ultrafast and comprehensive peptide identification in mass
591   spectrometry-based proteomics. Nat Methods 14:513-520.
592   45.   Huber W, von Heydebreck A, Sultmann H, Poustka A, Vingron M. 2002.
593   Variance stabilization applied to microarray data calibration and to the
594   quantification of differential expression. Bioinformatics 18 Suppl 1:S96-104.
595   46.   Coombs KM, Berard A, Xu W, Krokhin O, Meng X, Cortens JP, Kobasa D,
596   Wilkins J, Brown EG. 2010. Quantitative Proteomic Analyses of Influenza
597   Virus-Infected Cultured Human Lung Cells ▯ †. Journal of Virology 84:10888--
598   10906.
599   47.   Kammers K, Cole RN, Tiengwe C, Ruczinski I. 2015. Detecting significant
600   changes in protein abundance. EuPA Open Proteomics 7:11--19.
601   48.   Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK,
602   Cook H, Mende DR, Letunic I, Rattei T, Jensen Lars J, von Mering C, Bork P.
603   2019. eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated
604   orthology resource based on 5090 organisms and 2502 viruses. Nucleic Acids
605   Research 47:D309-D314.
606   49.   Chapalain A, Vial L, Laprade N, Dekimpe V, Perreault J, Déziel E. 2013.
607   Identification of quorum sensing‐controlled genes in *Burkholderia ambifaria*.
608   MicrobiologyOpen 2:226 242.
609   50.   Zlosnik JE, Hird TJ, Fraenkel MC, Moreira LM, Henry DA, Speert DP. 2008.
610   Differential mucoid exopolysaccharide production by members of the
611   *Burkholderia cepacia* complex. J Clin Microbiol 46:1470-3.
612   51.   Buxton R. 2005. Blood agar plates and hemolysis protocols. American Society
613   for Microbiology 15:1-9.
614   52.   Sayers EW, Bolton EE, Brister JR, Canese K, Chan J, Comeau DC, Connor
615   R, Funk K, Kelly C, Kim S, Madej T, Marchler-Bauer A, Lanczycki C, Lathrop
616   S, Lu Z, Thibaud-Nissen F, Murphy T, Phan L, Skripchenko Y, Tse T, Wang J,
617   Williams R, Trawick BW, Pruitt KD, Sherry ST. 2022. Database resources of
618   the national center for biotechnology information. Nucleic Acids Res 50:D20-
619   D26.
620   53.   Sayers EW, Cavanaugh M, Clark K, Ostell J, Pruitt KD, Karsch-Mizrachi I.
621   2019. GenBank. Nucleic Acids Res 47:D94-D99.
622   54.   Perez-Riverol Y, Bai J, Bandla C, Garcia-Seisdedos D, Hewapathirana S,
623   Kamatchinathan S, Kundu DJ, Prakash A, Frericks-Zipper A, Eisenacher M,
624   Walzer M, Wang S, Brazma A, Vizcaino JA. 2022. The PRIDE database
625   resources in 2022: a hub for mass spectrometry-based proteomics evidences.
626   Nucleic Acids Res 50:D543-D552.
627   55.   Lazar Adler NR, Stevens MP, Dean RE, Saint RJ, Pankhania D, Prior JL,
628   Atkins TP, Kessler B, Nithichanon A, Lertmemongkolchai G, Galyov EE. 2015.
629   Systematic mutagenesis of genes encoding predicted autotransported
630   proteins of *Burkholderia pseudomallei* identifies factors mediating virulence in

mice, net intracellular replication and a novel protein conferring serum
resistance. PLoS One 10:e0121271.

56. Biggins JB, Kang HS, Ternei MA, DeShazer D, Brady SF. 2014. The chemical
arsenal of *Burkholderia pseudomallei* is essential for pathogenicity. J Am
Chem Soc 136:9484-90.

57. Biggins JB, Ternei MA, Brady SF. 2012. Malleilactone, a polyketide synthase-
derived virulence factor encoded by the cryptic secondary metabolome of
Burkholderia pseudomallei group pathogens. Journal of the American
Chemical Society 134:13192--13195.

58. Klaus JR, Coulon PML, Koirala P, Seyedsayamdost MR, Déziel E, Chandler
JR. 2020. Secondary metabolites from the *Burkholderia pseudomallei*
complex: structure, ecology, and evolution. Journal of Industrial Microbiology
\& Biotechnology doi:10.1007/s10295-020-02317-0:1--11.

59. Wagley S, Vanaporn M, Rinchai D, Conejero L, Lertmemongkolchai G,
Bancroft GJ, Titball RW. 2017. A proteasome inhibitor produced by
Burkholderia pseudomallei modulates intracellular growth. Microb Pathog
107:175-180.

60. Coulon PML, Groleau M-C, Déziel E. 2019. Potential of the *Burkholderia
cepacia* complex to produce 4-hydroxy-3-methyl-2-alkyquinolines. Frontiers in
Cellular and Infection Microbiology 9:33.

61. Piochon M, Coulon PML, Caulet A, Groleau M-C, Déziel E, Gauthier C. 2020.
Synthesis and antimicrobial activity of *Burkholderia* -related 4-hydroxy-3-
methyl-2-alkenylquinolines (HMAQs) and Their N -oxide counterparts. Journal
of Natural Products 83:2145–2154.

62. Hinwan Y, Nithimongkolchai N, Trisakul K, Kaewseekhao B, Wonglakorn L,
Chetchotisakd P, Chareonsudjai S, Sirichoat A, Nithichanon A, Phelan J, Clark
TG, Faksri K. 2025. Whole-genome sequencing analysis *of Burkholderia
pseudomallei* comparing drug-resistant and pan-susceptible isolates reveals
novel biomarkers for drug resistance. Infect Genet Evol 133:105779.

63. Podnecky NL, Rhodes KA, Mima T, Drew HR, Chirakul S, Wuthiekanun V,
Schupp JM, Sarovich DS, Currie BJ, Keim P, Schweizer HP. 2017.
Mechanisms of resistance to folate pathway inhibitors in *Burkholderia
pseudomallei*: Deviation from the Norm. mBio 8.

64. Nandi T, Holden MT, Didelot X, Mehershahi K, Boddey JA, Beacham I, Peak I,
Harting J, Baybayan P, Guo Y, Wang S, How LC, Sim B, Essex-Lopresti A,
Sarkar-Tyson M, Nelson M, Smither S, Ong C, Aw LT, Hoon CH, Michell S,
Studholme DJ, Titball R, Chen SL, Parkhill J, Tan P. 2015. *Burkholderia
pseudomallei* sequencing identifies genomic clades with distinct
recombination, accessory, and epigenetic profiles. Genome Res 25:129-41.

65. Nandi T, Ong C, Singh AP, Boddey J, Atkins T, Sarkar-Tyson M, Essex-
Lopresti AE, Chua HH, Pearson T, Kreisberg JF, Nilsson C, Ariyaratne P,
Ronning C, Losada L, Ruan Y, Sung W-K, Woods D, Titball RW, Beacham I,
Peak I, Keim P, Nierman WC, Tan P. 2010. A Genomic Survey of Positive
Selection in *Burkholderia pseudomallei* provides insights into the evolution of
accidental virulence. PLoS Pathogens 6:e1000845.

Figure 1: Colony morphologies of the two *Bp* strains isolated from TSV292. A) -80°C
glycerol stock streak on CRLA, white arrow indicates the presence of second colony
morphotype corresponding to TSV292_2 smooth strain. B) Colony morphology of both
isolates after 3-days growth on both CRLA and Ashdown agar.

Figure 2: Two distinct *Bp* strains co-infecting the same patient suffering from
melioidosis in 2017.
A) Circular phylogenetic tree constructed using GrapeTree software showing relationships
between isolates based on wgMLST allelic profiles. The tree represents a minimum spanning
tree where branch lengths correspond to the number of allelic differences between isolates
across genome loci. Each tip represents an individual isolate, with closer proximity indicating
greater genetic similarity. Two reference strains are highlighted: TSV292_1 rough (blue) and
TSV292_2 smooth (green). The scale bar represents 1 allelic difference. Isolates are colored
by source as indicated in the legend. The analysis included 1,225 isolates characterized using
6,259 loci from the wgMLST scheme. B) Zoomed on closest isolated for both TSV292_1
rough (blue, right top panel) and TSV292_2 smooth (green, right bottom panel), using
maximum likelihood phylogenetic analysis, using Fasttree, based on closest relative
genomes.

Figure 3: CMV emergence from both isolates occur after 8 days of growth on various
media. White arrows indicate variants emergence from the main colony when culture on
different media to study bacterial phenotypes.

Figure 4: Proteomic profiling of both TSV292_1 rough and TSV292_2 smooth isolates.
A) Volcano plot representing the significant abundance changes in proteins between both
isolates. B) Enrichment analysis of virulence factors and COGs reflecting the proteins
changes.

A

B

TSV292

CRLB Agar

Strain 1
rough

Strain 2
smooth

Ashdown Agar

Strain 1
rough

Strain 2
smooth

A

B

Country ST

TSV292_2 smooth

0.974 — TSV23 — 2157
— TSV308 — 2074
0.478 — TSV33
0.39 — TSV160 — 2186
0.903 — 0.759 — TSV292_2 smooth — 2323
0.537 — TSV282 — 2139
— TSV213 — 2107
0.962 — TSV67
0.787 — TSV58 — 2166
— TSV201 — 2100
0.882 — TSV7
— TSV57 — 2165
0.274 — TSV261 — 2117

3.0E-4

60–103 — 34
DMG2200737
40–043 — 34
DMG2200739
1042 — 1542
0.94 — 70–062 — 1866
0.85 — 0.87 — DMG2200758
50–027 — 535
0.85 — DMG2200802
0.92 — TSV292_1 rough — 2319
0.91 — DMG2200741
0.53 — DMG2200761
0.95 — DMG2200759
0.62 — 0.76 — SR–039 — 694
0.93 — DMG2200745
0.9 — 90–008 — 197
DMG2200801

TSV292_1rough

2.0E-4

1

Source (Tips) ● Animal ● Environmental ● Human ○ Unknown

Countries ● Australia ● Cambodia ● Thailand ● Vietnam
Source (Tips) ● Human ○ Unknown

	0.01% Congo red LB Agar		**Ashdown Agar**		**Columbian Horse Blood agar**		**1.5% skimmed milk LB agar**		**Yeast Extract Mannitol Agar**	
	Strain 1 rough	**Strain 2 smooth**	**Strain 1 rough**	**Strain 2 smooth**	**Strain 1 rough**	**Strain 2 smooth**	**Strain 1 rough**	**Strain 2 smooth**	**Strain 1 rough**	**Strain 2 smooth**
TSV292

 |

 |

 |

 |

 |

 | | |

 |

 |

A

B