Differential Immunomodulatory Properties of Langya and Nipah Virus

Proteins

Stéphanie DURAND§, Dimitrios MAROUSIS§, Branka HORVAT*, and Louis-

Marie BLOYET*

Centre International de Recherche en Infectiologie, Inserm U1111, CNRS UMR5308,

Université de Lyon, Université Claude Bernard Lyon 1, École Normale Supérieure de Lyon,

Lyon, France

§ Contributed equally

* Corresponding authors: branka.horvat@inserm.fr; louis-marie.bloyet@cnrs.fr

Differential Immunomodulatory Properties of Langya and Nipah Virus Proteins

ABSTRACT

Langya virus (LayV) is a shrew-borne emerging parahenipavirus first identified in 2018 in 35 febrile patients in China. The closely related Nipah virus (NiV) is a highly pathogenic emerging bat-borne henipavirus that has caused numerous outbreaks with public health concerns in Asia. Among other symptoms, NiV causes severe acute respiratory syndrome and encephalitis, leading to high lethality. Thus, although closely related, infections with these two emerging and zoonotic viruses have distinct pathogenicity. Since the interplay with the host's immune system is a key determinant of species barrier crossing and pathogenicity, we aimed at deciphering the ability of LayV to counteract the human intrinsic immunity using the better-characterised NiV as a prototype. NiV expresses the P, V, and W proteins, known to hinder the host's innate immune response during infection. We thus compared the immunomodulatory properties of LayV and NiV proteins in human cells and showed that, similarly to NiV, the C-terminal domain of LayV V proteins inhibits the response to the activation of the pattern recognition receptor MDA5. However, although the N-terminal region of LayV P can inhibit the interferon signalling, it is not as efficient as its NiV counterpart. Moreover, only NiV W inhibits MDA5 and RIG-I signalling pathways. Similarly, unlike NiV, LayV W cannot efficiently block the activation of the NF-κB promoter after stimulation with IL-1β. These results suggest that LayV is less efficient than NiV in counteracting the human intrinsic immunity, which may contribute to the difference in severity observed between NiV and LayV-infected patients.

IMPORTANCE

Langya virus (LayV) is a shrew-borne emerging parahenipavirus recently identified in patients in China with symptoms such as fever, fatigue, cough, anorexia, headache, and vomiting. Since the interaction between a virus and its host's immune system is an essential parameter influencing host adaptation and disease severity, we investigated the interplay between LayV and the human intrinsic immunity using as a prototype the better-characterized and closely related Nipah virus (NiV), a highly pathogenic henipavirus. We compared the immunomodulatory properties of LayV and NiV proteins in human cells and showed that, similarly to NiV, some of LayV proteins can inhibit human signalling pathways, while, unlike NiV, LayV W protein is unable to block essential immune pathways. This suggests that LayV is less efficient than NiV in counteracting the human intrinsic immunity, which may contribute to the difference in severity between NiV and LayV-infected patients.

KEYWORDS: Langya virus, Nipah virus, zoonosis, innate immunity, interferon, NF-κB

INTRODUCTION

Langya and Nipah viruses are closely related zoonotic paramyxoviruses responsible for outbreaks in Asia. Langya virus (LayV) was first identified in 2018 in 35 patients with acute infections in China's Henan and Shandong provinces [1]. Symptoms included fever, fatigue, cough, anorexia, myalgia, nausea, headache, and vomiting. Shrews are likely the natural reservoir of LayV, although seropositivity has also been detected in dogs and goats. LayV has also been recently identified in shrews in the Republic of Korea [2], and closely related viruses have been discovered in shrews and rodents in Asia, Africa, Europe, and North America [3–10]. Due to its phylogenetic proximity with Nipah and Hendra viruses, LayV was initially described as a henipavirus, but the discovery of several related shrew-borne viruses led to the creation of the new *Parahenipavirus* genus [11] (Figure 1A). The ability of LayV to infect humans, the severity of the symptoms, the seropositivity of domestic animals, and its widespread geographic area raise concerns about its zoonotic potential. To date, no reverse genetic system for LayV is available, only the structures of LayV glycoproteins have been studied [12–17], and the interplay between LayV and its hosts remains unexplored.

Nipah virus (NiV) is a closely related emerging and highly pathogenic virus responsible for regular zoonotic outbreaks in Asia [18]. Discovered during an outbreak in Malaysia in 1998, NiV causes severe symptoms, including severe acute respiratory syndrome and encephalitis, with a case-fatality rate ranging from 40% to 75%. Since 2001, almost annual outbreaks in Bangladesh and India have been caused by spillovers from NiV natural reservoir: *Pteropus* fruit bats. In the absence of approved vaccines and treatments, and reports of human-to-human transmission, the World Health Organisation placed NiV on its Blueprint list of priority pathogens requiring urgent research and development efforts [19].

NiV and LayV share a similar genome organisation with the conserved six adjacent genes (Figure 1B) [2]. Most henipavirus P genes encode four proteins: P, V, W, and C proteins

[20]. V and W proteins are expressed from mRNAs edited by the viral polymerase during

transcription with the addition of one (V) or two (W) guanosine residues at a specific editing

site. Thus, the P, V, and W proteins share a common N-terminal domain (P_{NTD}) but differ in

their unique C-terminal domains (CTD) (Figure 1B). Although there is no direct evidence for

the existence of LayV V and W proteins, the conservation of the editing site and the sequences

of the V and W CTDs strongly suggests these proteins are produced during the infection (Figure

S1B-D).

NiV is well known to interfere with the host's intrinsic immunity, notably by inhibiting

the interferon type I (IFN-I) induction and signalling pathways, contributing to its high

pathogenicity [21–26]. Indeed, P_{NTD}, common to P, V, and W, binds and inhibits STAT1, thus

preventing the expression of interferon-stimulated genes (ISGs) [27–31,26,32,33].

Additionally, NiV V protein binds to MDA5, disrupts its folding, and thus prevents its

activation and the expression of interferon beta (IFN-β) [34–36]. Via its CTD, NiV W also

inhibits the NF-κB pathway by preventing the accumulation of the p65 subunit in the nucleus

[37].

Although closely related, LayV and NiV have different pathogenicity and infection

outcomes in humans. Since the interplay between viruses and the innate immune response plays

a critical role in pathogenicity, to examine these differences we compared the ability of the P,

V, and W proteins of LayV and NiV to counteract several pathways of the intrinsic immunity

in human cells.

MATERIALS AND METHODS

Cells

Human carcinoma HeLa cell line (Cat# ATCC® CCL-2™), Human Embryonic Kidney HEK293T (Cat# ATCC® CRL-3216™) and Human hepatocellular carcinoma cell line Huh-7.5 (kind gift from Charles Rice (Rockefeller University, New York, USA)) were grown in Dulbecco's modified Eagle's medium (DMEM), high glucose, GlutaMAX™ (LifeTechnologies™, Cat# 61965-026) supplemented with 10% of heat-inactivated (30 min at 56°C) foetal calf serum (Gibco, Cat #A5256701), and 1% of Penicillin-Streptomycin solution (Life Technologies™, Cat# 15140-130). All cell types were incubated at 37°C with 5% CO_2 and tested negative for Mycoplasma *spp*. (Lonza, Cat#LT07-318). Depending on their features, the different cell lines were used in specific assays: since they have a very high transfection rate, HEK293T cells were used in the IFN-I response luciferase assay to maximize the signal-to-noise ratio; due to their large cytoplasm, HeLa cells were used for microscopy and, to be consistent with the p65 localization assay, for the NF-κB pathway luciferase assay; and Huh7.5 cells were used to study RIG-I and MDA5 since they do not express RIG-I and only a low level of MDA5, thus allowing us to study one sensor or the other without interference from endogenous proteins [38].

Plasmids

The plasmids expressing the Firefly luciferase under the control of the IFN-β promoter (pβ-IFN-Luc [39]), the NF-κB promoter (pSGNluc, Addgene, Cat# 186705) [40], and the interferon response element (pISRE-Luc, Stratagene, Cat # 219092) were used to monitor the activation of the promoters. The plasmid phRL-TK expressing the Renilla luciferase under the HSV-thymidine kinase promoter (Promega, Cat# E6241) was used for normalisation. The plasmids

pEF_c-myc-MDA5 and pEF-BOS-HA-RIG-I encode myc-tagged human MDA5 and HA-tagged human RIG-I, respectively [41].

Plasmids coding for LayV P, V, and W (GenBank #OM101129) were cloned in pcDNA3.1 and ordered from GeneScript. These proteins contain an N-terminal Flag tag, separated by a GGGSG linker. Plasmids coding for NiV proteins (GenBank #AY988601) were cloned with the same tag in the same backbone using InFusion-mediated recombination (Takara, Cat# 638933). To express P/V/W$_{NTD}$, P$_{CTD}$ was removed from the plasmids expressing P. V$_{CTD}$ and W$_{CTD}$ were expressed as fusion proteins with a Flag-tagged mCherry protein fused at their N-terminus (separated from the CTDs by a single G residue). V and W mutants were generated from the plasmids encoding the non-mutated proteins using InFusion-mediated recombination.

The open reading frames of all plasmids were sequenced by Sanger sequencing (Eurofins Genomics).

Luciferase reporter assays

For the investigation of the MDA5 and RIG-I pathways, 2×10^4 Huh-7.5 cells were seeded per well in ninety-six-well plates. Each well was transfected the day after with 50 ng of pβ-IFN-Luc, 20 ng of phRL-TK, 50 ng of MDA5 or RIG-I expressing plasmid, together with 80 ng of plasmids expressing the viral proteins or the mCherry control, using JetOptimus reagent (Ozyme, Cat# POL101000006) according to the manufacturer's recommendations. Cells were transfected 24 h later with 100 ng of Poly(I:C) (Invivogen, Cat# PIC-39-05) using Lipofectamine 2000 (ThermoFisher Scientific, Cat #11668027). Medium was replaced four hours after stimulation with DMEM 10% FCS. Activities of the luciferases were measured 24 h after stimulation.

To investigate the IFN-I response pathway, ninety-six-well plates were coated with 50

μg/mL of poly-D-lysine (ThermoFisher Scientific, Cat# A3890401) for 30 min at 37°C, then

washed with water and dried before being seeded with 4×10^4 HEK293T cells per well. Each

well was transfected the day after with 12.5 ng of pISRE-Luc, 5 ng of phRL-TK, together with

35.4 ng of plasmids expressing the viral proteins or the mCherry control, using JetOptimus

reagent (Ozyme, Cat# POL101000006) according to the manufacturer's recommendations.

Cells were treated the day after with 1000 U/mL of IFNα-2a (ImmunoTools, Cat#11343506)

for 18 h before measuring the activity of the luciferases.

To monitor the NF-κB pathway, 2×10^4 HeLa cells per well were seeded in ninety-six-

well plates. Each well was transfected the day after with 12.5 ng of pSGNluc, 6.25 ng of phRL-

TK, together with 31.25 ng of plasmids expressing the viral proteins or the mCherry control,

using JetOptimus reagent (Ozyme, Cat# POL101000006) according to the manufacturer's

recommendations. Cells were treated 24 h later with 10 ng/μL of human IL-1β (Peprotech, Cat#

200-01B) for 2 h before measuring the luciferase activities.

For all assays, the activities of the luciferases were measured using the Dual-Glo

Luciferase Assay system according to the manufacturer's protocol (Promega, Cat# E2920) and

a microplate reader (Tristar 5, Berthold) with acquisition for 1 s. For each well, the relative

luciferase activity was calculated as the ratio between the Firefly and Renilla luciferase

activities. For each condition, experiments were performed in triplicate and repeated in four

separate experiments.

Immunofluorescence

For the detection of viral proteins, 2×10^5 HeLa cells were seeded on 12 mm glass coverslips

placed in a 12-well plate. Twenty-four hours later, cells were transfected with 0.5 μg of plasmid

and fixed 48 h after transfection with 4% paraformaldehyde. Cells were permeabilised with

0.1% Triton X-100 in PBS for 20 min and blocked for 30 min in PBS - 2% BSA. Staining was done overnight at 4°C using mouse anti-Flag antibody (Merck, Cat# F1804) diluted in PBS – 0.5% BSA, followed by incubation for 1 h at room temperature with DAPI (Thermo FisherTM, Cat# 62248) and Alexa Fluor 488 goat anti-mouse antibody (ThermoFisher Scientific) diluted at 1.3 μg/mL in the same buffer.

For the detection of p65, $2x10^5$ HeLa cells were seeded on 12 mm glass coverslips placed in a 12-well plate and transfected 24 h later with 2 μg of plasmid expressing NiV or LayV W or W_{CTD} proteins. Twenty-four hours after transfection, cells were stimulated or not with 10 ng/mL of human IL-1β (Peprotech, Cat# 200-01B) for 20 min at 37°C before being washed with ice-cold PBS and fixed in 4% paraformaldehyde for 30 min on ice. Cells were then permeabilised for 10 min in TBS – 0.1% Triton X-100 and blocked for 10 additional minutes in TBS – 2.5% BSA. Then, cells were stained overnight at 4°C with rabbit anti-NF-κB p65 (Cell Signalling, Cat#8242) and mouse anti-Flag antibody (Merck, Cat# F1804) diluted in TBS– 2% FCS - 0.1% Tween. The day after, cells were incubated for 1 h at room temperature with DAPI (ThermoFisher Scientific, Cat# 62248) together with Alexa Fluor 488 donkey anti-rabbit (ThermoFisher Scientific, Cat# A21206) and Alexa Fluor 555 goat anti-mouse antibodies (ThermoFisher Scientific, Cat# A21422) diluted at 1.3 μg/mL in the same buffer.

Coverslips were mounted with Fluoromount-G (SouthernBiotech, Cat# 0100-01) before observation on a confocal microscope (Zeiss, LSM800Airy Scan) hosted in the PLATIM microscopy facility (SFR Biosciences, Lyon, France). Images were treated using ImageJ software.

Quantification and statistical analysis

Luciferase assay analysis

All assays with statistical analysis were performed in four independent experiments, with each condition tested in triplicate (three technical replicates per plate). For each well, the Firefly luciferase signal was normalised by the Renilla luciferase signal. Statistical analyses were done using the non-parametric Mann-Whitney test.

Image quantification

Fluorescence microscopy images were analysed and quantified using QuPath 0.5.1 software (https://qupath.github.io/). Cells were identified using a cell detection program based on the staining of the nuclei (DAPI staining). For the quantification of the NF-κB p65 signal, the nuclear signal is defined as the signal present in the mask created by the DAPI staining, and the cytoplasmic signal as the signal present in a 2 μm radius outside of the nucleus mask. Each condition was tested in four independent experiments with three images analysed per experiment, leading to a total of 150 cells analysed per condition. Statistical analysis was done using the non-parametric Mann-Whitney test.

Phylogenetic tree

The L protein sequences from Nipah virus (NC_002728), Hendra virus (NC_001906), Cedar virus (NC_025351), Ghana virus (NC_025256), Yunnan bat henipavirus 1 (PQ621839), Yunnan bat henipavirus 2 (PQ621840), Angavokely henipavirus (ON613535), Gamak virus (MZ574407), Crocidura tanakae henipavirus 2 (PP272531), Jingmen Crocidura shantungensis henipavirus 1 (OM030314), Wufeng Crocidura attenuata henipavirus 1 (OM030317), Crocidura tanakae henipavirus 1 (OQ970176), Resua virus (OR713876), Wufeng Chodsigoa smithii henipavirus 1 (OM030316), Chodsigoa hypsibia henipavirus (OQ236120), Camp Hill virus (PQ140948), Ninorex virus (OQ438286), Denwin virus (OR713883), Crocidura olivieri henipavirus 2 (PQ541139), Melian virus (OK623353), Jingmen Crocidura shantungensis henipavirus 2 (PP272750), Daeryong virus (MZ574409), Crocidura olivieri henipavirus 1

223 (PQ541138), Lechcodon virus (OR713879), Wenzhou shrew henipavirus 1 (OQ715593),

224 Hasua virus (OR713881), Mojiang virus (NC_025352), and Langya virus (OM101125), were

225 used to build a multiple sequence alignment using MAFFT [42]. The phylogenetic tree was

226 calculated with the Neighbor Joining algorithm in MAFFT.

RESULTS

Similar intracellular localisation of NiV and LayV P and V proteins, but different localisation of their W proteins

While NiV P and V proteins are found in the cytoplasm, W is concentrated in the nucleus [29,43,44,37]. To investigate if LayV proteins harbour the same cellular localisation, we transfected NiV and LayV proteins fused to a Flag tag in HeLa cells and observed their localisations by immunofluorescence. Similar to NiV, LayV P and V proteins are both localised in the cytoplasm (Figure 2A, B). On the contrary, NiV and LayV W proteins are found in different cell compartments, with NiV W being restricted to the nucleus while LayV W is diffused in the cytoplasm (Figure 2C). These results correlate with the presence of a nuclear localisation signal (NLS) located on NiV W_{CTD} and the absence of conservation of these residues in LayV W (Figure S1D) [45].

NiV and LayV P proteins inhibit the IFN-I signalling pathway

Via their common N-terminal domain (P_{NTD}), NiV P, V, and W proteins inhibit the transcription factor STAT1, leading to a decrease in IFN-I signalling [27–31,26,32,33]. To monitor this effect, the highly transfectable HEK293T cells were transfected with plasmids encoding the viral proteins and a reporter gene under the control of the Interferon-Stimulated Response Element (ISRE). After stimulation with IFN-2α, the reporter signal was significantly decreased for cells expressing NiV and LayV P proteins compared to the control cells expressing the fluorescent protein mCherry (Figure 3A). Similarly to NiV, LayV P_{NTD} is sufficient to mediate this inhibition (Figure 3B). However, although LayV P and P_{NTD} expression levels were higher or similar to NiV proteins, LayV proteins were less efficient in blocking the signalling (Figure

3, S2A). This difference may be linked to the low conservation of the STAT1-binding region

identified on NiV P and thus to a lower affinity for human STAT1 (Figure S1A).

NiV and LayV V proteins inhibit MDA5 but not RIG-I signalling pathways

The C-terminal domain of V proteins is capable of binding MDA5, disrupting its folding, and

inhibiting the subsequent activation of the IFN-β promoter [34–36,46]. To compare the effect

of NiV and LayV V proteins on the IFN-β induction pathway after MDA5 or RIG-I transfection

and activation, we used Huh-7.5 cells since they do not express RIG-I and only a low level of

MDA5 [38]. NiV and LayV V proteins are both capable of inhibiting the induction of the IFN-

β promoter upon MDA5 activation (Figure 4A, S2B). This effect is even stronger when only

the V_{CTD} is expressed as a fusion protein with mCherry (mCh-V_{CTD}) (Figure 4B, S2B). As

expected, a mutant of NiV V_{CTD} carrying the mutations E411A and I414A, known to impair the

binding to MDA5 [47], is no longer able to inhibit the activation of the IFN-β promoter (Figure

4B). On the contrary, neither NiV nor LayV V proteins nor the mCh-V_{CTD} constructs inhibited

the activation of the IFN-β promoter after RIG-I stimulation (Figure 4C, D).

Overall, these results indicate that both NiV and LayV V can inhibit the IFN-β induction

pathway after the activation of MDA5 but not RIG-I.

The W protein of NiV, but not LayV, inhibits MDA5 and RIG-I signalling pathways

When V proteins were replaced with W in the MDA5 and RIG-I activation assays, NiV W was

able to inhibit the activation of the IFN-β promoter upon both MDA5 (Figure 5A) and RIG-I

(Figure 5C) activation. This effect was stronger when only the CTD of NiV W protein was

expressed (mCh-W_{CTD}) (Figure 5B, D). Conversely, LayV W protein did not have any

inhibitory effect on the activation of the IFN-β promoter upon MDA5 and RIG-I activation

(Figure 5). To compensate for the lower expression of LayV W compared to NiV W (Figure

S2C), we transfected a range of plasmid quantities. While we detected an inverse correlation

between the amount of NiV W and the reporter signal intensity, the activation of the reporter's

promoter was insensitive to the increasing amount of LayV W, even when expressed at higher

levels than NiV W, thus indicating that the difference in inhibitory effect is protein-specific and

not due to the protein expression levels (Figure S3). This is further confirmed with the mCh-

W_{CTD} constructs, which are similarly expressed, with only NiV W_{CTD} being able to inhibit the

induction (Figure 5B, D, S2C).

While LayV W is found in the cytoplasm, NiV W is concentrated in the nucleus thanks

to an NLS present on W_{CTD} (Figure 1B, S1D). Since this signal is required for NiV W inhibitory

activity of immune pathways [37,43], we examined whether the absence of inhibitory activity

of LayV W would be due to its cytoplasmic localisation. Mutating NiV W NLS (NLSko)

relocated the mCh-W_{CTD} fusion protein in the cytoplasm, and adding back the exogenous NLS

of the simian vacuolating virus 40 ($SV40_{NLS}$) partially restored the nuclear localisation (Figure

6A). As expected, the inhibitory effect of these NiV mCh-W_{CTD} constructs on the IFN-β

promoter correlated with their intracellular localisation (Figure 6A, B). On the contrary, while

the addition of the $SV40_{NLS}$ to LayV mCh-W_{CTD} also partially relocated the protein to the

nucleus, this relocalisation did not provide any significant inhibitory effect to LayV W_{CTD}

(Figure 6A, C). Thus, the absence of an inhibitory effect of LayV W is not solely due to its

cytoplasmic localisation.

The NF-κB pathway and the translocation of p65 into the nucleus are impaired by NiV but

not LayV W protein

We have previously demonstrated that NiV W can inhibit the NF-κB pathway via its CTD [37].

Here, we analysed the effect of LayV and NiV proteins on the NF-κB pathway after stimulation

of cells with IL-1β (Figure 7A, B). Although weakly for the full-length proteins, the expression

of NiV W and W_{CTD} decreased the activation of the NF-κB promoter, while LayV proteins did

not and even increased the activity (Figure 7A, B).

Since NiV W decreases the nuclear concentration of NF-κB p65 subunit in infected

cells, we monitor the localisation of p65 after NiV or LayV W expression and IL-1β stimulation

(Figure 7C, D). While NiV W decreased the concentration of p65 in the nucleus via its CTD,

LayV W did not prevent the relocalisation of p65 in the nucleus upon IL-1β stimulation. These

results suggest that, unlike NiV W, LayV W is not able to inhibit the accumulation of p65 in

the nucleus and thus the subsequent activation of the NF-κB promoter.

DISCUSSION

The presence of the entry receptor(s) and the ability to counteract the host defence are among

the most important barriers for a virus to successfully infect a new host. For paramyxoviruses,

the proteins translated from the P gene (P, V, W, and C) are the main anti-host defence factors.

In the absence of a reverse genetic system for Langya virus, we used cell transfection assays to

assess the ability of LayV P, V, and W proteins to block several pathways of the intrinsic

immunity in human cells and compared them to NiV proteins, a closely related, zoonotic, and

highly pathogenic virus.

We report that both LayV and NiV V proteins can efficiently inhibit transcription

activation after MDA5 stimulation. Since the amino acids essential for the folding of the V_{CTD}

and the binding to MDA5 are highly conserved between NiV and LayV, and among henipa-

and parahenipaviruses (Figure S1C), LayV V likely directly binds to and blocks the activation

of MDA5. Although NiV V was described to decrease RIG-I activation via the inhibition of the

TRIM25-mediated ubiquitination of RIG-I [48], we did not observe any effect in our assays.

This discrepancy likely reflects differences in the assays, mainly the absence of overexpression

of TRIM25 and the use of different cell lines (HEK293T vs Huh-7.5). Thus, we cannot conclude

on the effect of LayV V on the TRIM25-mediated ubiquitination of RIG-I.

328   Similarly, LayV P can inhibit the signalling under the IFN-I receptor, but less efficiently

329   than NiV P. This difference does not seem correlated with P expression levels and may reflect

330   differences in their ability to bind human STAT proteins, as suggested by the rather low

331   conservation of the STAT1-binding region of NiV P.

332   Unlike NiV W, LayV W was unable to inhibit transcription activation after activation

of MDA5, RIG-I, and the IL-1β receptor. Since the transcription factor NF-κB is activated by

each of these signalling pathways, and is known to be inhibited by NiV W, NiV W may inhibit

these three pathways using the same molecular mechanism: recruiting the cellular protein 14-

3-3 to the nucleus to favour the export of p65 [37,49]. The relocation of LayV W to the nucleus,

where NiV W is concentrated, did not provide any functional effect, and LayV W lacks the

residues of NiV W known to bind 14-3-3. Thus, unlike NiV W, LayV W is most likely not able

to recruit 14-3-3 to the nucleus to inhibit p65 accumulation.

340   Since the C protein also participates in the immune evasion, and LayV is thought to

341   express an additional protein upstream of the F open reading frame (Figure 1B), these two

342   proteins may contribute to the immune evasion and compensate for the absence of activity of

343   LayV W in inhibiting the human immunity pathways.

344   Overall, this study suggests that the P and W proteins of LayV are not as efficient as

345   NiV proteins in counteracting human intrinsic immunity. Thus, since the inhibition of the innate

346   immune system is a major factor of pathogenicity, this distinction between NiV and LayV may

347   be one of the reasons for the difference in pathogenicity observed in humans, with the high

348   lethality of NiV-infected patients, in contrast to LayV infection.

ACKNOWLEDGEMENTS

We acknowledge the contribution of the SFR Biosciences (Université Claude Bernard Lyon 1, CNRS UAR3444, Inserm US8, ENS de Lyon) LYMIC-PLATIM microscopy platform for training and the use of the LSM800 AS confocal microscope. We thank Mathieu Iampietro and Lucia Amurri (CIRI, Lyon) for their advice on the NF-κB experiments and all current and former members of the team Immunobiology of Viral Infections at CIRI for their help in the realisation of the study. We thank Omran Allatif (CIRI, bioinformatics and biostatistics (BIBS) platform) for his help with statistical analysis.

FUNDING STATEMENT

The work was supported by the Institut National de la Santé et de la Recherche Médicale (Inserm), the Centre National de la Recherche Scientifique (CNRS), the Aviesan Sino-French agreement on Nipah virus study, and is part of the "Investissements d'Avenir ExcellencES" program from the French National Research Agency (Shape-Med@Lyon; ANR-22-EXES-0012).

DISCLOSURE STATEMENT

The authors report no conflict of interest.

DATA AVAILABILITY STATEMENT

Data is available upon request.

REFERENCES

[1]   Zhang X-A, Li H, Jiang F-C, et al. A Zoonotic Henipavirus in Febrile Patients in China. N Engl J Med. 2022;387(5):470–472.

[2]   Natasha A, Pye SE, Park K, et al. Detection and characterization of Langya virus in *Crocidura lasiura* , Republic of Korea [Internet]. 2024 [cited 2024 Dec 17]. Available from: http://biorxiv.org/lookup/doi/10.1101/2024.12.02.626520.

[3]   Parry RH, Yamada KYH, Hood WR, et al. Henipavirus in Northern Short-Tailed Shrew, Alabama, USA. Emerg Infect Dis. 2025;31(2):392–394.

[4]   Zhang J-T, Hu Z-Y, Tang F, et al. Decoding the RNA viromes in shrew lungs along the eastern coast of China. NPJ Biofilms Microbiomes. 2024;10(1):68.

[5]   Wu Z, Yang L, Yang F, et al. Novel Henipa-like virus, Mojiang Paramyxovirus, in rats, China, 2012. Emerg Infect Dis. 2014;20(6):1064–1066.

[6]   Lee S-H, Kim K, Kim J, et al. Discovery and Genetic Characterization of Novel Paramyxoviruses Related to the Genus Henipavirus in Crocidura Species in the Republic of Korea. Viruses. 2021;13(10):2020.

[7]   Vanmechelen B, Meurs S, Horemans M, et al. The characterization of multiple novel paramyxoviruses highlights the diverse nature of the subfamily Orthoparamyxovirinae. Virus Evol. 2022;8(2):veac061.

[8]   Horemans M, Van Bets J, Joly Maes T, et al. Discovery and genome characterization of six new orthoparamyxoviruses in small Belgian mammals. Virus Evol. 2023;9(2):vead065.

[9]   Chen Y-M, Hu S-J, Lin X-D, et al. Host traits shape virome composition and virus transmission in wild small mammals. Cell. 2023;186(21):4662-4675.e12.

[10] Haring VC, Litz B, Jacob J, et al. Detection of novel orthoparamyxoviruses, orthonairoviruses and an orthohepevirus in European white-toothed shrews. Microb Genomics. 2024;10(8):001275.

[11] Vanmechelen B, Balkema-Buschmann A, Drexler F, et al. Taxonomic reorganization of the family Paramyxoviridae [Internet]. 2023. Available from: https://www.google.com/url?sa=t&source=web&rct=j&opi=89978449&url=https://ictv.global/system/files/proposals/approved/Animal_dsRNA_and_ssRNA-_viruses/2023.018M.Paramyxoviridae_reorg.docx&ved=2ahUKEwjVq8SBiZ6OAxVIcKQEHTWZE9cQFnoECD4QAQ&usg=AOvVaw0lkIr3W06nbxmNzrDMife-.

[12] Byrne PO, Blade EG, Fisher BE, et al. Prefusion stabilization of the Hendra and Langya virus F proteins. J Virol. 2024;98(2):e0137223.

[13] Guo Y, Wu S, Li W, et al. The cryo-EM structure of homotetrameric attachment glycoprotein from langya henipavirus. Nat Commun. 2024;15(1):812.

[14] Isaacs A, Low YS, Macauslane KL, et al. Structure and antigenicity of divergent Henipavirus fusion glycoproteins. Nat Commun. 2023;14(1):3577.

[15]   May AJ, Pothula KR, Janowska K, et al. Structures of Langya Virus Fusion Protein
Ectodomain in Pre- and Postfusion Conformation. J Virol. 2023;97(6):e0043323.

[16]   Wang C, Li M, Wang Y, et al. Structural insights into the Langya virus attachment
glycoprotein. Struct Lond Engl 1993. 2024;32(8):1090-1098.e3.

[17]   Wang Z, McCallum M, Yan L, et al. Structure and design of Langya virus glycoprotein
antigens. Proc Natl Acad Sci U S A. 2024;121(16):e2314990121.

[18]   Pelissier R, Iampietro M, Horvat B. Recent advances in the understanding of Nipah
virus immunopathogenesis and anti-viral approaches. F1000Research. 2019;8.

[19]   Mehand MS, Al-Shorbaji F, Millett P, et al. The WHO R&D Blueprint: 2018 review of
emerging infectious diseases requiring urgent research and development efforts.
Antiviral Res. 2018;159:63–67.

[20]   Lo MK, Harcourt BH, Mungall BA, et al. Determination of the henipavirus
phosphoprotein gene mRNA editing frequencies and detection of the C, V and W
proteins of Nipah virus in virus-infected cells. J Gen Virol. 2009;90(Pt 2):398–404.

[21]   Fontana JM, Bankamp B, Rota PA. Inhibition of interferon induction and signaling by
paramyxoviruses. Immunol Rev. 2008;225(1):46–67.

[22]   Pelissier R, Iampietro M, Horvat B. Recent advances in the understanding of Nipah
virus immunopathogenesis and anti-viral approaches. F1000Research. 2019;8.

[23]   Mathieu C, Guillaume V, Volchkova VA, et al. Nonstructural Nipah virus C protein
regulates both the early host proinflammatory response and viral virulence. J Virol.
2012;86(19):10766–10775.

[24]   Satterfield BA, Cross RW, Fenton KA, et al. The immunomodulating V and W proteins
of Nipah virus determine disease course. Nat Commun. 2015;6:7483.

[25]   Satterfield BA, Cross RW, Fenton KA, et al. Nipah Virus C and W Proteins Contribute
to Respiratory Disease in Ferrets. J Virol. 2016;90(14):6326–6343.

[26]   Satterfield BA, Borisevich V, Foster SL, et al. Antagonism of STAT1 by Nipah virus P
gene products modulates disease course but not lethal outcome in the ferret model. Sci
Rep. 2019;9(1):16710.

[27]   Rodriguez JJ, Parisien J-P, Horvath CM. Nipah Virus V Protein Evades Alpha and
Gamma Interferons by Preventing STAT1 and STAT2 Activation and Nuclear
Accumulation. J Virol. 2002;76(22):11476–11483.

[28]   Park M-S, Shaw ML, Muñoz-Jordan J, et al. Newcastle disease virus (NDV)-based
assay demonstrates interferon-antagonist activity for the NDV V protein and the Nipah
virus V, W, and C proteins. J Virol. 2003;77(2):1501–1511.

[29]   Shaw ML, García-Sastre A, Palese P, et al. Nipah Virus V and W Proteins Have a
Common STAT1-Binding Domain yet Inhibit STAT1 Activation from the Cytoplasmic
and Nuclear Compartments, Respectively. J Virol. 2004;78(11):5633–5641.

[30] Rodriguez JJ, Cruz CD, Horvath CM. Identification of the nuclear export signal and
STAT-binding domains of the Nipah virus V protein reveals mechanisms underlying
interferon evasion. J Virol. 2004;78(10):5358–5367.

[31] Ciancanelli MJ, Volchkova VA, Shaw ML, et al. Nipah virus sequesters inactive
STAT1 in the nucleus via a P gene-encoded mechanism. J Virol. 2009;83(16):7828–
7841.

[32] Jensen MR, Yabukarski F, Communie G, et al. Structural Description of the Nipah
Virus Phosphoprotein and Its Interaction with STAT1. Biophys J. 2020;118(10):2470–
2488.

[33] Keiffer TR, Ciancanelli MJ, Edwards MR, et al. Interactions of the Nipah Virus P, V,
and W Proteins across the STAT Family of Transcription Factors. mSphere. 2020;5(6).

[34] Andrejeva J, Childs KS, Young DF, et al. The V proteins of paramyxoviruses bind the
IFN-inducible RNA helicase, mda-5, and inhibit its activation of the IFN-β promoter.
Proc Natl Acad Sci U S A. 2004;101(49):17264–17269.

[35] Childs K, Stock N, Ross C, et al. mda-5, but not RIG-I, is a common target for
paramyxovirus V proteins. Virology. 2007;359(1):190–200.

[36] Wagner ND, Liu H, Rohrs HW, et al. Nipah Virus V Protein Binding Alters MDA5
Helicase Folding Dynamics. ACS Infect Dis. 2022;8(1):118–128.

[37] Enchéry F, Dumont C, Iampietro M, et al. Nipah virus W protein harnesses nuclear 14-
3-3 to inhibit NF-κB-induced proinflammatory response. Commun Biol.
2021;4(1):1292.

[38] Louber J, Brunel J, Uchikawa E, et al. Kinetic discrimination of self/non-self RNA by
the ATPase activity of RIG-I and MDA5. BMC Biol. 2015;13:54.

[39] King P, Goodbourn S. The beta-interferon promoter responds to priming through
multiple independent regulatory elements. J Biol Chem. 1994;269(48):30609–30615.

[40] Kuri P, Ellwanger K, Kufer TA, et al. A high-sensitivity bi-directional reporter to
monitor NF-κB activity in cell culture and zebrafish in real time. J Cell Sci.
2017;130(3):648–657.

[41] Louber J, Kowalinski E, Bloyet L-M, et al. RIG-I Self-Oligomerization Is Either
Dispensable or Very Transient for Signal Transduction. Boudinot P, editor. PLoS ONE.
2014;9(9):e108770.

[42] Katoh K, Rozewicki J, Yamada KD. MAFFT online service: multiple sequence
alignment, interactive sequence choice and visualization. Brief Bioinform.
2019;20(4):1160–1166.

[43] Shaw ML, Cardenas WB, Zamarin D, et al. Nuclear localization of the Nipah virus W
protein allows for inhibition of both virus- and toll-like receptor 3-triggered signaling
pathways. J Virol. 2005;79(10):6078–6088.

[44]   Lo MK, Harcourt BH, Mungall BA, et al. Determination of the henipavirus
484   phosphoprotein gene mRNA editing frequencies and detection of the C, V and W
485   proteins of Nipah virus in virus-infected cells. J Gen Virol. 2009;90(2):398–404.

486   [45]   Smith KM, Tsimbalyuk S, Edwards MR, et al. Structural basis for importin alpha 3
487   specificity of W proteins in Hendra and Nipah viruses. Nat Commun. 2018;9(1):3703.

488   [46]   Motz C, Schuhmann KM, Kirchhofer A, et al. Paramyxovirus V Proteins Disrupt the
489   Fold of the RNA Sensor MDA5 to Inhibit Antiviral Signaling. Science.
490   2013;339(6120):690–693.

491   [47]   Ramachandran A, Horvath CM. Dissociation of Paramyxovirus Interferon Evasion
492   Activities: Universal and Virus-Specific Requirements for Conserved V Protein Amino
493   Acids in MDA5 Interference. J Virol. 2010;84(21):11152–11163.

494   [48]   Sánchez-Aparicio MT, Feinman LJ, García-Sastre A, et al. Paramyxovirus V Proteins
495   Interact with the RIG-I/TRIM25 Regulatory Complex and Inhibit RIG-I Signaling. J
496   Virol. 2018;92(6):e01960-17.

497   [49]   Edwards MR, Hoad M, Tsimbalyuk S, et al. Henipavirus W Proteins Interact with 14-
498   3-3 To Modulate Host Gene Expression. J Virol. 2020;94(14):e00373-20.

FIGURES

Figure 1. Phylogenetic tree and genome organisation of henipaviruses and parahenipaviruses.

(A) Phylogenetic tree of henipavirus and parahenipaviruses based on the protein sequence of

the L protein. Red dots indicate viruses known to be pathogenic in humans, and the red circle

506   indicates Mojiang virus, suspected to be pathogenic for humans [5]. (B) Genome organisation

507   of NiV and LayV. The proteins produced from the P gene are indicated below the genomes. A

508   STAT1-binding site (STAT), a nucleus export signal (NES), and a nucleus localisation signal

509   (NLS), described for NiV, are indicated.

Figure 2. Localisation of NiV and LayV P, V, and W proteins.

HeLa cells were transfected to express NiV or LayV Flag-tagged P (A), V (B), or W (C). At 24

h.p.t., proteins were stained using an anti-Flag antibody and imaged by confocal microscopy.

Viral proteins are in green, and nuclei are in blue. Scale bars correspond to 20 μm.

Figure 3. NiV and LayV P/V/W$_{NTD}$ inhibit the IFN-I signalling pathway.

HEK293T cells were co-transfected with three plasmids coding for (1) a viral protein or the

mCherry (mCh) control, (2) the Firefly luciferase under an ISRE-dependent promoter, and (3)

the Renilla luciferase under a constitutive promoter. Either the full-length P proteins are

expressed (A) or only the N-terminal domain common to P, V, and W (P/V/W$_{NTD}$) (B). At 18

h.p.t., cells were stimulated with 1000 U/mL IFN-2α for 24 h before measuring the luciferase

activities. Mean and SD were calculated from four independent experiments.

Figure 4. NiV and LayV V proteins inhibit MDA5 but not RIG-I signalling pathway.

Huh-7.5 cells were co-transfected with four plasmids coding for (1) human MDA5 (A, B) or RIG-I (C, D), (2) a viral protein or the mCherry control, (3) the Firefly luciferase under the IFN-β promoter, and (4) the Renilla luciferase under a constitutive promoter. Viral proteins are expressed either as full-length proteins (A, C) or as fusion proteins between mCherry and the V C-terminal domain (mCh-V$_{CTD}$) (B, D). Cells were stimulated 24 h.p.t. by transfecting 100 ng Poly(I:C), and the luciferase activities were measured 48 h.p.t. Mean and SD were calculated from four independent experiments. mCh-V$_{CTD}$* carries the mutations E411A and I414A.

Figure 5. Only NiV W protein inhibits MDA5 and RIG-I signalling pathways.

Huh-7.5 cells were co-transfected with four plasmids coding for (1) human MDA5 (A, B) or

RIG-I (C, D), (2) a viral protein, (3) the Firefly luciferase under the IFN-β promoter, and (4)

the Renilla luciferase under a constitutive promoter. Viral proteins are expressed either as full-

length proteins (A, C) or as fusion proteins between mCherry and the W C-terminal domain

(mCh-W_{CTD}) (B, D). Cells were stimulated 24 h.p.t. by transfecting 100 ng of Poly(I:C), and

luciferase activities were measured 48 h.p.t. Mean and SD were calculated from four

independent experiments.

Figure 6. The cytoplasmic localisation of LayV W is not responsible for the absence of inhibition of the MDA5 and RIG-I pathways.

(A) HeLa cells were transfected to express fusion proteins between mCherry and NiV or LayV W C-terminal domain (mCh-W$_{CTD}$). In some constructs, the nuclear localisation signal (NLS) of NiV W was mutated (NLSko), and the NLS from SV40 was added to NiV and LayV W$_{CTD}$ (SV40$_{NLS}$). Twenty-four hours post-transfection, stained cells were imaged by confocal

microscopy. Viral proteins are in red and nuclei in blue. Scale bars correspond to 20 μm. (B,

C) Huh-7.5 cells were co-transfected with four plasmids coding for (1) human MDA5 or RIG-

I, (2) a viral protein or the mCherry control, (3) the Firefly luciferase under the IFN-β promoter,

and (4) the Renilla luciferase under a constitutive promoter. Cells were stimulated 24 h.p.t. by

transfecting 100 ng of Poly(I:C), and relative activities of the luciferases were measured 48

h.p.t. Mean and SD were calculated from four independent experiments.

Figure 7

Figure 7. NiV but not LayV W protein can block the NF-κB pathway.

(A and B) HeLa cells were co-transfected with three plasmids coding for (1) a viral protein or

the mCherry control, (2) the Firefly luciferase under an NF-κB-dependent promoter, and (3) the

Renilla luciferase under a constitutive promoter. Viral proteins are expressed either as full-

length proteins (A) or as fusion proteins between mCherry and the W C-terminal domain (mCh-W$_{CTD}$) (B). At 18 h.p.t., cells were incubated with 10 ng/mL IL-1β for 2 h before measuring the luciferase activities. Mean and SD were calculated from four independent experiments. (C) HeLa cells were transfected to express NiV or LayV W or mCh-W$_{CTD}$ viral proteins. At 24 h.p.t., cells were stimulated with 10 ng/mL IL-1β for 20 min before being fixed. W was stained with an anti-Flag antibody and endogenous p65 protein with an anti-NF-κB p65 antibody. Stained cells were imaged by confocal microscopy. Nuclei are in blue, p65 in green, and viral proteins in red. Scale bars correspond to 20 μm. (D) The mean intensity of nuclear and cytoplasmic NF-κB p65 was evaluated using QuPath software on an average of 150 events per condition. Mean and SD were calculated from four independent experiments.