1 **TipA, a *Bdellovibrio bacteriovorus* BPI-like double-TULIP is opened by its adapter**
2 **protein, TipB.**
3

4 **Simon G. Caulton[1*], Ian T. Cadby[2], Gareth W. Hughes[3], Hannah Johnston[1], Timothy J.**
5 **Knowles[1], Andrew L. Lovering[1]**
6

7 **Author Affiliations**

8
9 [1]School of Biosciences, University of Birmingham, Birmingham B15 2TT
10 [2]Bristol Veterinary School, University of Bristol, Langford, Bristol BS40 5DU, UK
11 [3]Cancer and Genomic Sciences, University of Birmingham, Birmingham B15 2TT
12 ***corresponding author**
13

14 **Abstract**

15
Bdellovibrio bacteriovorus is a predatory bacterium that invades the periplasm of other Gram-
negative bacteria and liberates prey biomolecules for replication. *Bdellovibrio* has a wealth of
genes that encode unique proteins to enable this lifestyle. Using a series of x-ray structures, we
show that the operonal pairing of *bd2538* and *bd2539,* encode a double <u>T</u>ULIP (tubular lipid
binding protein) <u>i</u>nvasion <u>p</u>rotein A (TipA) and a small beta sandwich (TipB), respectively.
TipA has a specialised N-terminal TULIP domain with a beta hairpin and helix that mediate
homodimerisation through hairpin interdigitation. This dimerisation creates a large,
continuous, enclosed lumen that we demonstrate to contain multiple lipid molecules. In
addition, we show that TipB functions as a small adapter protein that binds to TipA using
specialised loops that bury into the TipA hydrophobic core. This binding forms a clamp on the
edge of the N-terminal beta sheet and induces a large 20 Å conformational change, opening the
TULIP fold to create an accessible interior. This study presents the first structural
characterisation of lipid binding proteins in *Bdellovibrio*, and the first example of
conformational change in TULIPs mediated by an adaptor protein.

Synopsis

Bdellovibrio bacteriovorus is a predatory bacterium that invades the periplasm of other Gram-
negative bacteria in order to consume them from within. To enable this lifestyle, it has an
arsenal of predation-associated genes, including the operonal pair *bd2538* and *bd2539*. We use
structural and biophysical techniques to characterise two proteins produced by these genes,
TipA and TipB.

•   TipA is a double tubular lipid binding protein (TULIP) that forms a homodimer that
sequesters lipids in its lumen and binds to lipid bilayers
•   TipB is a small beta sandwich that interacts with TipA via two specialised loops
•   Interaction of TipA with TipB opens the TULIP fold of TipA

Acronyms

ASU – Asymmetric unit

TULIP – Tubular lipid binding protein

SMP – Synaptotagmin-like mitochondrial-lipid-binding protein

LPT – Lipopolysaccharide transport

MLA - Maintenance of OM lipid asymmetry

BPI – Bacterial permeability-increasing protein

CETP – Cholesteryl ester transfer protein

LBP – LPS-binding protein

E-Syt – Extended Syntaptotagmin

ERMES – ER mitochondrial encounter structure

RMSD – Root mean square deviation

TLC – Thin layer chromatography

ITC – Isothermal titration calorimetry

QCM-D – Quartz crystal microbalance with dissipation

PDB – protein data bank

CL – cardiolipin

PE – Phosphatidylethanolamine

PG – phosphatidylglycerol

AFDB50 – AlphaFold Database 50

Introduction

Bdellovibrio bacteriovorus is a predatory bacterium that preys on other Gram-negative bacteria. Upon prey encounter, *B. bacteriovorus* attaches to its prey and invades its periplasm, where it remodels the host peptidoglycan to form a rounded bdelloplast. Here, it liberates prey biomolecules for its filamentous growth, followed by septation into progeny. Finally, when the prey cell is exhausted of nutrients, it becomes lysed to release free swimming *B. bacteriovorus* to repeat the cycle anew [1]. This remarkable prey-killing lifestyle, and its indifference to prey antibiotic-resistance status gives *Bdellovibrio* huge potential as a novel biological control agent [2]. However, this lifestyle also requires a wealth of specialised tools [3]. Indeed, *Bdellovibrio* utilizes a number of liberated biomolecules from its prey [4-7],

including both lipid A [8] and fatty acids [9]. These lipids can either remain in their original

form or undergo metabolic alterations by *Bdellovibrio*.

Bdellovibrio's intricate association with membranes is central to its predatory lifestyle.

Membrane and cell wall remodelling [10-12], as well as lipid uptake [8, 9] are critical facets

of host cell penetration and subsequent intraperiplasmic survival. The mechanisms through

which *Bdellovibrio* obtains lipids from its prey remain elusive, highlighting the complexity of

lipid transport in bacteria, which typically necessitates dedicated machinery, such as the Lpt or

Mla systems for lipid movement across the aqueous periplasmic environment [13]. However,

these systems are responsible for intracellular transport of lipids across the periplasm of a cell,

shuttling between inner and outer membranes. *Bdellovibrio* has the additional need for

intercellular lipid transport, to extract and transfer lipids from prey cell membranes to self.

Tubular lipid-binding proteins (TULIPs) exhibit unique structural features comprising a

cylindrical fold that surrounds a hydrophobic cavity, enabling them to sequester lipids from the

surrounding environment. These include the extracellular bacterial permeability increasing

protein-like (BPI-like) double-TULIP family, Takeout-like proteins and intracellular

Synaptotagmin-like mitochondrial-lipid-binding protein (SMP) family [14, 15]. TULIPs have

gained attention for their diverse roles in membrane tethering, lipid transfer, hormone binding

and innate immunity in eukaryotes [15, 16]. Recently, widespread identification in prokaryotes

has made it clear that TULIPs have important roles in prokaryote physiology [17, 18].

Structurally characterised prokaryotic TULIPS are few, and include *E. coli* YceB, and

Clostridium botulinum neurotoxins/neurotoxin-associated proteins, P47 and OrfX proteins [19,

20]. Understanding the molecular mechanisms governing TULIP-lipid interactions is crucial

for deciphering their role in prokaryotic organisms, inclusive of alternative lifestyles like the

predator *Bdellovibrio*.

In this study, we provide X-ray crystal structures of two proteins encoded from the operonal

gene cluster *bd2538* and *bd2539,* TULIP invasion protein A (TipA) and TULIP-associated

invasion protein B (TipB), respectively. These proteins are upregulated during the invasion

process [21] and secreted into prey [22]. We show that TipA forms a novel dimeric BPI-like

double-TULIP, longer than classical characterized family representatives, and that TipB binds

to TipA to induce a conformational opening of the TULIP fold. These results provide the first

example of a lipid-binding protein in *Bdellovibrio* and provide insight into the molecular

mechanisms of lipid trafficking during its predatory lifestyle.

Results

Bdellovibrio TipA is a BPI-like double TULIP divergent from other family

members

The operonal genes *bd2538* and *bd2539* (Fig. 1A) encode two proteins we term TULIP

invasion protein A (TipA) and TULIP-associated invasion protein B (TipB), respectively. To

identify the role of TipA in predation we sought to determine its structure (mature enzyme,

minus signal peptide) and elucidate its function. The crystal structure of selenomethionine-

incorporated TipA was solved in space group P 3_1 2 at 4 Å resolution with three molecules in

the asymmetric unit (ASU). The low resolution of diffraction was likely a consequence of the

87% solvent content caused by the end-to-end packing of the proteins in the unit cell.

Nevertheless, maps of sufficient quality were obtained (Fig. 1B) and in all three chains, at least

residues N35-A566 were traced with no unmodelled gaps in the model (Fig. 1C). TipA was

annotated to contain a DUF2785 domain, however the crystal structure confirms mis-

annotation, as the protein fold shows it to be of the lipid binding protein BPI/LBP superfamily

(IPR030675) [23].

TipA shows similar gross features to those of other BPI/LBP family structures: a double-

TULIP fold, comprising a central beta sheet sandwiched by two tubular domains each formed

by a beta sheet that wraps around an extended helix (Fig. 1C). Each TULIP domain has the

typical secondary structure characteristic of that domain: an N-terminal helix (helix A), an

elongated beta sheet, and a C-terminal helix (helix B). The single protomer forms a cylindrical

protein with dimensions of approximately 142 x 35 x 40 Å enveloping a central hydrophobic

cavity throughout the entire protein.

TipA was submitted to the DALI server [24] to identify proteins with similar structures. The

top hits were all double TULIP proteins: Bacterial permeability-increasing protein (BPI; Z-

score 16.5; 21% seq ID), Cholesteryl ester transfer protein (CETP; Z-scores 13.5-15) and LPS-

binding protein (LBP; Z-scores 10.4-11). Superposition of the individual N- and C-terminal

domains of TipA with those of BPI shows significant divergence of the protein fold, with

RMSD values of 5.7 Å over 176 residues for the N-terminal domain and 5.8 Å over 112

residues for the C-terminal domain (Fig. 1D-E). Hence, we conclude that TipA is a divergent

TULIP representing a grouping of the superfamily previously uncharacterised.

The N-terminal domain shows three inserts relative to that of BPI: a beta hairpin between the

last beta strand and helix B, a loop containing an additional helix between the first and second

strands, and a small helix between the second and third strands (Fig. 1D). The first two of these

inserts comprise the dimerisation interface of TipA, whereas the function of the latter is

unclear.

Compared to BPI, the C-terminal domain has a truncated loop between strands 1 and 2, a larger

triangular-shaped loop between strands 2 and 3, and two small helices inserted between the last

strand and helix B. Superposition reveals that the central beta sheet and helix A align well with

the BPI C-terminal domain, but upon exit from helix A, TipA diverges greatly (Fig. 1E). In

particular, both the sheet and helix B are much shorter than BPI, and this causes a relative

displacement.

TipA forms a cylindrical homodimer

A surprising finding of the crystal structure of TipA was that unlike any other characterised

TULIP, TipA forms a C2 symmetrical dimer (Fig. 2A), with one complete dimer in the ASU

and one protein forming a dimer with a symmetry copy. TipA has a novel beta hairpin and

helix on the N-terminal domain that form a dimer interface, producing an extended

hydrophobic tube through an end-on-end interaction (Fig. 2A-B). The channel throughout

dimeric TipA is almost continuous, containing a pinch point in each of the central sheet

domains, and has a total volume of approximately 25000 $Å^3$ as calculated by 3vee [25] (Fig.

2B). The dimerisation hairpins form a single antiparallel beta sheet and produce hydrophobic

and polar faces with two intermolecular electrostatic interactions formed by E251 and K256

(Fig. 2C). A sequence alignment of TipA shows a number of highly conserved residues that

are important structurally (Fig. EV1). The tip of the dimerisation hairpin contains a highly

conserved asparagine (N253) that hydrogen bonds with the backbone of F184. N260 is also

highly conserved and serves to propagate helix B following the hairpin. The highly conserved

P246 kinks the backbone to bend the hairpin away from the beta sheet and angle it parallel to

the protein tube, allowing interdigitation with the hairpin of the second molecule. The N-

terminus of the dimerisation helix packs against its own helix B and extends outwards to pack

its C-terminal tryptophan into the hydrophobic face of the other protomer's helix B (Fig. 2D).

The dimerisation helix makes hydrogen bonds between Q104 and Q104, and E263 and Q111.

The associated loop (L113-S117) forms backbone hydrophobic interactions through L113 and

F116 and backbone hydrogen bonds between V114 and Y258. A highly conserved glycine

(G115) allows the chain to form a sharp turn to thread back to the beta sheet of the N-terminal

domain.

Due to the spacious packing of the TipA dimer crystals, and consequent relatively low

resolution, we subsequently sought to obtain a higher resolution structure of TipA by altering

the packing of the crystal lattice. To do this we substituted the hairpin region (A249-F258) with

a triple glycine linker, termed TipA$_{\Delta Hairpin}$. We expected the loss of dimerisation to allow better

packing of the protein, reducing solvent content to produce higher resolution diffraction. This

substitution shifted the elution volume of the protein from 164 mL to 200 mL on a Superdex

200 pg 26/60 column, showing a drastic decrease in hydrodynamic radius of the protein, and

indicating that the protein is primarily monomeric in solution (Fig. 2E). As predicted, the

TipA$_{\Delta Hairpin}$ protein crystallised with different packing in space group P 2_1 2_1 2_1, with a solvent

content of 60% and higher resolution diffraction of 2.6 Å. The crystal structure of TipA$_{\Delta Hairpin}$

contains a single monomeric copy in the ASU (Fig. 2F). The lack of dimer interface introduced

disorder into the N-terminal domain and the regions of G107-M118 and the glycine linker

region P246-N253 (N260 in wild-type (WT)) so these residues could not be modelled. The C-

terminal domain and central sheet are very similar to that of the WT protein, but the N-terminal

TULIP sheet is pulled inward, mediated by a bend in Helix B at G277 (G284 in WT). This

suggests that dimerisation is integral to maintaining correct folding of the N-terminal domain.

TULIP partner TipB is a β-sandwich with exposed hydrophobic loops

Since the TipA-encoding *bd2538* gene is in a predicted operon with the downstream *bd2539*

gene, which it overlaps by 4 nucleotides and which we name *tipB*, we hypothesised that TipB

could be functionally linked to TipA. Following our structural studies of TipA, we purified and

crystalized the cryptic operonal TipB protein. TipB crystallised in space group C 2 2 2_1 with

three copies in the ASU and diffracted to 2.17 Å resolution. The structure reveals that TipB

forms a compact β-sandwich, in which one four-strand anti-parallel sheet forms the central

globular core of the protein and is packed on one side by a pair of short beta sheets and the

other side by an elongated capping-loop extending from β7 and β8 (Fig. 3A). This loop folds

back on the sheet burying Y120, W122 and M133 into the core. The protein has another pair

of loops that extend between β3 and β4, and β6 and β7, termed interaction loops 1 and 2,

respectively (Fig. 3A). Importantly, both loops have hydrophobic residues at the ends: F65 and

I66 in loop 1 and F103 in loop 2, and demonstrate a high degree of flexibility (as inferred by

electron density and B-factor values; Fig 3B). Finally, the protein contains a short helical turn

between strands 2 and 3 and an 11-residue helix between strands 8 and 9.

We used the DALI server [24] to identify structural homologues of TipB and the highest

scoring structure was *P. aeruginosa* ExsB [26] (PDB code 2YJL), with a Z-score of 7.3. Both

TipB and ExsB show a topology similar to that of type II C2 domains (Fig. 3C-E) [27], a

calcium-dependent membrane-localisation domain found in multiple eukaryotic signalling

proteins including kinases [28], lipases [29] and membrane trafficking proteins [30]. In

comparison, TipB lacks the 8^{th} strand that runs antiparallel between the 1^{st} and 7^{th} strands,

instead packing its last strand parallel to the 2^{nd} strand (the 1^{st} strand is broken into two smaller

strands due to a broken beta sheet in TipB), forming a mixed sheet. The other top-scoring hits

all had the standard type II C2-domain topology, with varying additional loops and secondary

structure components. Most of the C2-domain proteins have an exposed face on the beta sheet,

however some homologues show extensions in the β7 and β8 loop that cap off this sheet in a

similar manner to TipB (Fig. 3E). None of the homologues had extensions between β3 and β4

or β6 and β7 strands where the interaction loops are observed in TipB.

ExsB has an exposed beta sheet with a basic face predicted to mediate binding to acidic lipids

(Fig. 3E). In TipB, this sheet face is not exposed and forms part of the hydrophobic core that

is enclosed by an extended loop between β7 and β8. To investigate whether TipB would adhere

to a lipid bilayer, we used quartz crystal microbalance with dissipation monitoring (QCM-D)

measurements (Fig. 3F). This surface-based technique monitors changes to the resonant

frequency of a surface (Δf) under an applied voltage. Δf is inversely proportional to the mass

deposited on the surface, and as such allows us to monitor interactions between biomolecules

[31]. In the case of TipB-lipid interactions, a stable POPC bilayer was deposited onto a silicon

QCM-D sensor, following well-established methods [32, 33]. Despite not being a constituent

of bacterial membranes, a POPC bilayer was used here due to its well-reported ability to form

highly stable bilayers when deposited onto a surface [34]. Once lipid bilayers were deposited,
TipB was introduced and its interaction with the lipid bilayer was assessed in a semi-
quantitative manner. Upon flow of TipB over the bilayer, negligible frequency (0.3 Hz) and
dissipation (0.48 ppm) changes occurred, thus confirming that TipB alone does not bind
directly to lipid membranes.

C2 domains commonly bind calcium, such as the high-scoring DALI hit *Homo sapiens*
phospholipase 2 [35]. We wanted to confirm that TipB doesn't bind calcium as it does not
contain obvious calcium-binding residues nor was calcium observed in the crystal structure.
After treatment with 5 mM EGTA and 5 mM EDTA, and subsequent gel filtration, no binding
was seen using isothermal titration calorimetry (ITC; Fig. 3G), confirming that TipB is a
calcium-independent C2-like β-sandwich.

TipA and TipB form a TULIP:partner complex

Since TipB did not display any phospholipid binding activity we next hypothesised that TipA
and TipB might interact with each other in a functional manner (whilst also being stable in their
isolated states, as inferred from our structures). Isothermal titration calorimetry (ITC) of TipB
into TipA resulted in a clear binding curve (Fig. 4A), and the -TΔS value suggests the
interaction is largely driven through entropy effects, possibly through hydrophobic
contributions. However, due to the low stoichiometry observed (0.25), the curve wasn't
suitable to calculate an accurate Kd. The relatively poor stability of the TipA protein would not
allow for higher concentrations to be used. The low stoichiometry could result from TipA
existing as multiple species in solution. Although TipA elutes as a single peak in gel filtration
experiments, this peak is broad (when compared with ΔHairpin TipA), potentially the result of
multiple TipA conformers (Fig. 2E). It is therefore likely that only a subpopulation of TipA is
competent for binding and may further demonstrate the flexible nature of the protein.

We subsequently desired to obtain a crystal structure of the complex of TipA and TipB. The
two proteins were purified and mixed with a molar excess of TipB in the crystallisation
conditions in order to ensure a complex of the two proteins. This complex crystallised in space
group P 2 2 2$_1$ and diffracted to 2.6 Å with one copy of each protein in the ASU. The TipA
component formed a dimer on a crystallographic 2-fold axis exhibiting the same hairpin-helix

dimer interface as the selenomethionine-labelled TipA crystals (Fig. 4B). TipB can be observed

binding centrally on TipA, straddling the unusually kinked helix A, which protrudes away from

the beta sheet, forming a triangular seat (Fig. 4C-E). The structure of complexed TipB is very

similar to that of TipB alone, with RMSDs ranging from 1.43 to 1.81 Å over 139 amino acids

for the three chains of the TipB crystal structure. Differences are mostly observed in interaction

loops 1 and 2, which fold down over the seat and bury residues F65, I66 and F103 into the

hydrophobic lumen of TipA (Fig. 4F-G). In addition to the hydrophobic components, there are

a number of hydrogen and electrostatic bonds from side chains and backbone atoms throughout

the 1346 Å^2 interface (Fig. 4H-I), as calculated by PISA [36].

Partner TipB binds to open up the TULIP fold and central cavity

The most striking observation of the complex structure is the conformational change seen in

TipA's N-terminal domain (Fig. 5A-B). While the superposition of the C-terminal domains of

the apo and complexed TipA structures show an RMSD of 1.26 Å over 261 residues, the N-

terminal domain shows an RMSD of 6.13 Å over 271 residues. The N-terminal domain of the

TipA apo structure comprises an almost completely closed tube, with very little access to

solvent (Fig. 5C). However, the N-terminal domain of the complex is in an open conformation,

with a gap of approximately 20 Å (from Y80-S282; Fig. 5D). This opening is mediated by an

approximate 45° rotation of the protruding TipA helix A (Fig. 5A), which acts like a lever,

pulling the N-terminal sheet open. This opening is also stabilised by the anchoring of

interaction loop 1 into the C-terminal domain, and the packing of interaction loop 2 F103 under

the lifted N-terminal sheet. TipB F103 acts as a clamp underneath the TipA beta sheet in

concert with K100 and R34, which form electrostatic interactions with E72 and E129,

respectively, on the top of the sheet (Fig. 5E). Analysis by DynDom [37] characterises this as

a hinge mechanism, comprising three moving domains (Fig 5F): the C-terminal domain and

most of the N-terminal helix B (Domain 1); most of the N-terminal sheet and dimerisation

hairpin (Domain 2); and the end of the N-terminal sheet and dimerisation helix (Domain 3).

The analysis identifies two hinge points between these domains which include the rotation

point of the N-terminal helix A seat. Domain 2 rotates approximately 29° relative to Domain 1

and Domain 3 rotates 36° relative to Domain 2, keeping it a similar relative position to Domain

1 in both structures (Fig. 5G). Hence, the function of TipB is to convert TipA to a more open

form and, to the best of our knowledge, represents a new type of TULIP binding-partner.

Using Foldseek [38] to search for homologous proteins, we observe double-TULIPs with
additional fused domains (Fig. EV2). These include beta-rich domains, helical bundles and
mixed domains including enzymes and entire double-TULIP repeats. Of particular interest is
the presence of a double-TULIP in the *Planctomyces sp*. SH-PL14 protein, Uniprot accession
A0A142WTQ2. This protein contains a fused N-terminal esterase-like enzyme in addition to a
C-terminal beta-sandwich. This beta-sandwich has similar topology to TipB, and is also
predicted to localise on the N-terminal Helix A and sheet, suggesting that it could have a similar
role to TipB in this system.

TipA binds rapidly to membrane bilayer surfaces

During the purification process of TipA, we observed that under conventional buffer conditions
(<500 mM NaCl) the protein remained largely in the pellet after lysis and centrifugation.
However, upon addition of high salt concentrations or detergents, the protein was released into
the supernatant. This led us to believe that the protein was adhering to the membrane fraction
(Fig. 6A).

To confirm this, we again used QCM-D (Fig. 6B-E). To investigate TipA-lipid interactions,
stable bilayers consisting of either *E. coli* polar lipids, POPC or mixed POPC:LPS lipids were
deposited onto a silicon QCM-D sensor. Upon introduction of TipA to all three bilayer
compositions monitored here, it exerted a dramatic negative Δf, suggesting it was rapidly
deposited into the bilayer systems. This interaction was most pronounced for TipA deposition
into POPC:LPS bilayers (Fig. 6D), with a $\Delta f3$ of -1106 Hz, compared to -1064 Hz and -686
Hz for POPC and polar lipid bilayers, respectively. In all bilayers tested, a critical point was
reached upon which further addition of TipA resulted in removal of mass from the surface.
This suggests an ability to first insert into lipid bilayers followed by membrane disruption.
Furthermore, in bilayers containing POPC a double-dip insertion of TULIP was observed (Fig.
6C-D), which may result from the ability of POPC to form more stable bilayers on silicon
surfaces than *E. coli* polar lipids. Interestingly, this distinctive mode of interaction was not
observed when TipA was introduced to an empty silicon surface generated through the inability
to deposit polar lipid:LPS bilayers, with no major mass deposition or removal observed (Fig.
6E).

Lipids are observed in the TipA TULIP lumen in multiple states

In order to confirm that TipA is a classic TULIP that binds lipids within its cavity, thin layer chromatography (TLC) was performed (Fig. 7A). TipA clearly copurifies with all three common *E. coli* polar lipids, cardiolipin (CL), phosphatidylethanolamine (PE) and phosphatidylglycerol (PG). These have likely been extracted from *E. coli* during the expression or extraction process. Density for lipids was observed in all three crystal structures, however the low resolution of the selenomethionine form prevented their modelling (Fig. EV2A). In both the 2.6 Å TipAB complex structure (Fig. 7B) and the 2.6 Å monomeric TipA$_{\Delta\text{Hairpin}}$ structure (Fig. 7C), clear densities are observed for two lipid head groups in the N-terminal TULIP domain. It was also possible to trace the lipid tails in the TipA$_{\Delta\text{Hairpin}}$ structure but not in the complex structure. The lipids were modelled as PE as this headgroup fit the density well (Fig. 7D), and the density would not accommodate the glycerol moiety of PG, nor did the density indicate the presence of the double headgroup of CL.

In both structures the lipid headgroups are seen in similar regions in the protein (as are the Fo-Fc peaks in the selenomethionine TipA structure; Fig. EV2A), one underneath Helix A, and the other between the sheet and Helix B (Fig. 7E-F). However, contributing amino acids that bind the ligand are different between the structures. The placement of the inner lipid in the TipAB structure means that the headgroup makes a H-bond and hydrophobic interactions with G102 and F103. In the TipA$_{\Delta\text{Hairpin}}$ structure (where TipB is not present), it makes H-bonds with Y132 and K69, but additional interactions are formed by V493 of the symmetry copy, providing a backbone H-bond and sidechain hydrophobic interactions. These modelling differences suggest that TipB affects lipid interactions with TipA.

Interestingly, in both structures one of the lipids makes contact with the C-terminal domain of a symmetry copy (Fig. EV2B-D), with polar contributions from the symmetry copy K457 and K503 to the phosphate moiety of the outer lipid in the TipAB structure. When visualising higher order packing in all three TipA structures, this is a common occurrence – in three different space groups the C-terminal hydrophobic residues M454, F455, L504 and F505 pack into the N-terminal domain 'seam' between Helix B and the 1$^{\text{st}}$ beta strand (Fig. EV2E-G).

Further structures of close TipA homologues are needed to determine if this mode of interaction

is a conserved feature.

Discussion

Tubular lipid binding proteins (TULIP) proteins are a universal lipid-binding fold,

characterised by their ability to sequester lipids within their internal hydrophobic cavities.

Whilst these folds are ubiquitous to all domains of life, prokaryotic TULIPS present an under-

represented group within the literature [17]. We have structurally characterised the first

example of a lipid-binding protein from *Bdellovibrio bacteriovorus* HD100 - the predation-

associated TULIP, TipA, and identified an associated adapter protein TipB.

TipA is in the same superfamily as the eukaryotic bacterial permeability increasing protein

(BPI), which also forms a double-TULIP fold. The protein shows similar structural features to

that of other BPI-like proteins, such as the classical Helix-ββββ-Helix TULIP core secondary

structure, two TULIP domains back-to-back with an intervening central sheet, and longer N-

terminal than C-terminal TULIP domain. However, modifications to the protein have made the

overall fold divergent, with the addition of molecular features that enable the formation of a

dimeric double-TULIP, and for interaction with the TipB adapter protein. The dimerisation

motif includes an additional hairpin and helix that lock the homodimer together to form a

hugely elongated ~260 Å tube.

While no other double-TULIP has been observed in a dimeric form, the extended

synaptotagmins (E-Syts) and the eukaryotic ER-mitochondria encounter structure (ERMES)

proteins do form single-TULIP homodimers, but they pack using a different interface where

helix B packs against the beta sheet edge [39]. It is interesting that a heterotetrameric form of

TULIP is also seen between ERMES proteins Mdm12 and Mmm1, where heterodimers stack

head-to-head through an Mmm1-Mmm1 interface [40]. This creates an equivalent chain of four

TULIP domains, highlighting that higher-order arrangements of TULIPs are functionally

significant. It is possible that this increase in TULIP length is needed in order to efficiently

transport lipids effectively at some juncture between prey and *Bdellovibrio*. The length of the

dimer is between 250 and 260 Å which is equivalent to that of the periplasmic spanning

transporter complexes, Yeb and Pqi [41] as well as the predicted structure of the putative

transporter YhdP [42]. ERMES proteins and E-Syts tether membranes and transfer lipids

between them, which is dependent on the presence of their C2 domains for membrane tethering
[43], and therefore proximity of membranes to their single-TULIP is required. The increased
length of the TipA dimer could allow for lipid transfer between membranes without the need
for direct membrane tethering by additional domains.

It is also noteworthy that we consistently observe TipA packing C-terminal hydrophobic
residues into the N-terminal domain – which creates a network of TULIPs in the crystal lattice.
This could indicate that during predation TipA could form higher order structures that facilitate
transport of lipids from TULIP to TULIP. Higher order structures also occur in the ERMES
complex as the Mmm1-Mdm12 complex also interacts with another TULIP Mdm34 [40, 44],
but structural information and exact stoichiometry is unknown for these larger complexes.
Further work could confirm whether there is a physiological function for this consistent C-
terminal-N-terminal TULIP packing of TipA.

The *tipA* gene is in an operon with *tipB*, which encodes for a small C2-like sandwich domain.
The functional link between TULIPs and C2 domains is observed in ERMES and E-Syt
systems, where the C2 domains mediate membrane tethering. TipB, while C2-like, does not
bind calcium or membranes, and functions in an entirely different manner. The TipB sandwich
has been adorned with two extended loops that are used for interaction with TipA by insertion
into its lumen. The crystal structure of the complex of TipA and TipB shows that upon binding
of the two proteins there is a large conformational change in TipA, mediated by a clamping
mechanism that lifts its N-terminal beta sheet. This creates an opening of the TULIP seam,
creating an accessible core across both N-terminal domains of the TipA dimer. Previously, a
gate-and-latch mechanism has been proposed to account for lipid binding into the lumen of the
single-TULIP juvenile-hormone binding protein from the domestic silk moth [45]. Here, the
N-terminal helix A opens in the apo form and closes on the buried sesquiterpene molecule
when bound. As the N-terminal helix of double-TULIP proteins is locked into the central beta
sheet by an additional N-terminal beta strand, this latch mechanism is not possible. Instead, it
is possible that the opening of TipA by TipB is what facilitates loading or unloading of lipids;
observation of lipid-loaded recombinant TipA by TLC indicates that it would not be dependent
on TipB. The intriguing finding of a planctomycete double-TULIP with a fused beta-sandwich
that binds to the N-terminal domain Helix A and sheet proposes the idea that this functional
relationship is repeated across diverse bacteria. In addition, it is possible that TipA
homodimerisation is also required to ensure the stability of the overall fold during this opening

and closing process as we have observed a high degree of flexibility in the N-terminal sheet (very pronounced in TipA$_{\Delta\text{hairpin}}$), and the lumen of the N-terminal domain is larger than typically observed in other double-TULIPs.

We were unsuccessful in attempts to monitor TipA localisation – trials with a C-terminal mCherry tag resulted in frequent mutational escape, so TipA could not be investigated by fluorescence microscopy; future studies could use antibodies or computational-design binders to address this issue. TipA's ability to adhere to membranes and destabilise them as shown via QCM-D suggests that it could be used to extract lipids directly from prey bacterial membranes. Binding and membrane destabilisation by TipA could aid *Bdellovibrio*, and could work in a similar fashion to BPI [46] by increasing membrane permeability. TipA is not the first example of a TULIP protein adhering to membranes without C2 domain partners - tricalbin 2 (a yeast E-syt homologue) and nucleus-vacuole junction protein 1 TULIP domains were shown to bind to membranes *in vitro* using liposomes [47], and the tricalbins are known to transfer lipids between membrane contact sites. Therefore, membrane adherence (as observed by both our purification procedure and QCM-D experiments) is likely an effective way for TipA to uptake lipids into its core ready for transfer to another protein or membrane.

In conclusion, our findings underscore the evolutionary significance of TULIPs as universal lipid-binding folds across both eukaryotes and prokaryotes. The crystal structures of TipA, TipB and their binary complex show that *B. bacteriovorus* has novel lipid-binding machinery for its predatory lifestyle in the form of a modified dimeric double-TULIP protein. Through the mechanistic interdependence of TipA and TipB, we give new insight into the conformational changes possible in double-TULIP proteins and suggest an adaptor-mediated mechanism for promoting lipid loading and unloading.

Methods

Cloning, expression and purification

TipA-His was cloned into a modified version of the pET41c overexpression vector using restriction-free cloning methods. The final construct contained residues of 26-572 of TipA (Uniprot accession Q6MK73) fused to a C-terminal octa-His-tag. TipB was cloned into pET26b

using NdeI and NotI. A tagless form of TipA was made by mutagenesis to remove the octa-
Histag. A monomeric form of tagless TipA (TipA$_{\Delta Hairpin}$) was created using Q5 mutagenesis,
replacing the dimer-hairpin residues A249-Y-258 (AVEVNGKKFY) with a triple glycine
linker. TipB was cloned into pET26b using NdeI and XhoI restriction sites. The final construct
contained an N-terminal MAHHHHHH tag and residues 29-171 of TipB (Uniprot accession
Q6MK72).

To express selenomethionine-labelled TipA, a 60 mL overnight culture of BL21 (DE3) cells in
LB was centrifuged and resuspended in minimal media supplemented with kanamycin and
chloramphenicol to a final volume of 1 L. At OD$_{600}$ 0.4, 0.1 g of lysine, 0.1 g of threonine, 0.1
g of phenylalanine, 0.05 g leucine, 0.05 g of isoleucine, 0.05 g of valine and 0.06 g of
selenomethionine were added to the culture. After 15 minutes, IPTG was added to 1 mM. Cells
were harvested by centrifugation after 16 h at 18 ˚C and stored at -20 ˚C.

TipA$_{\Delta Hairpin}$, TipA-His and TipB constructs were expressed in *Escherichia coli* BL21 RIPL
(DE3). Cells were grown in LB supplemented with kanamycin to an OD$_{600}$ of 0.5-0.7 and
induced with 0.5 mM IPTG at 18 °C for 16 h. Cells were harvested by centrifugation at 4000g
for 15 minutes. During trial purifications we noted that the majority of TipA protein sedimented
with insoluble cell components following lysis and centrifugation, but could be liberated from
the insoluble pellet by the addition of high (1M) NaCl. We exploited this characteristic of TipA
in our subsequent purification protocols. Cells expressing untagged selenomethionine-labelled
TipA and TipA$_{\Delta Hairpin}$ were resuspended in wash buffer (50 mM HEPES and 0.15 M NaCl, pH
7.5), sonicated and centrifuged at 48384 g for 1 h. The pellet was again resuspended, sonicated
and centrifuged to wash away soluble proteins. A third pellet wash was performed and finally
the pellet was resuspended in 50 mM HEPES and 1 M NaCl, pH 7.5. Cells expressing his-
tagged proteins were resuspended in buffer A (TipA-His - 50 mM HEPES and 1M NaCl, pH
7.5; TipB - 50 mM HEPES and 300 mM NaCl, pH 7.5). The cells were lysed by sonication on
ice and centrifuged at 48384 g for 1 hour. The pellet was resuspended. The lysates were loaded
onto a 5 mL Ni-NTA column (GE Healthcare) and washed with 10 CV of lysis buffer. The
protein was eluted with a gradient of 20–500 mM imidazole. All proteins were finally passed
through a Superdex 200 pg or 75 pg 26/60 column (GE Healthcare), which was pre-equilibrated
with 20 mM HEPES, 150 mM NaCl, pH 7.5. Fractions containing the protein of interest were
pooled, concentrated and stored frozen at -70 °C if not required for immediate use.

Crystallisation

Selenomethionine-lebelled TipA crystallisation screens were prepared by mixing a 1.5 μL of 8 mg/mL TipA and 1.5 μL precipitant solution. TipA crystallised in space group P 31 2 1 (a =252.4, b=252.4, c=181.1, α=90, β=90, γ=120) in a precipitant solution from the Jena Bioscience JBS Classic Screen consisting of 15% w/v PEG 4000; 100 mM trisosodium citrate, pH 5.6; 200 mM ammonium sulphate. Diffraction data were collected at Diamond Light Source. Data reduction and processing were performed using XDS.

TipA$_{\Delta Hairpin}$ crystallisation screens were prepared by mixing a 0.4 μL of 10 mg/mL TipA and 0.4 μL precipitant solution. TipA crystallised in space group P 21 21 21 (a =58.0, b=97.8, c=132.0, α=90, β=90, γ=90) in a precipitant solution from the BCS screen consisting of 0.1 M MES pH 6.5, 2% v/v PEG 400, 2% v/v PEG 500 MME, 2% v/v PEG 600, 2% w/v PEG 1000, 2% w/v PEG 2000, 2% w/v PEG 3350, 2% w/v PEG 4000, 2% w/v PEG 5000 MME, 2% w/v PEG 6000, 2% w/v PEG 8000, 2% w/v PEG 10000. Diffraction data were collected at Diamond Light Source. Data reduction and processing were performed through the Xia-2 Dials pipline, and Aimless.

TipB crystallisation screens were prepared by mixing a 0.4 μL of 10 mg/mL TipB and 0.4 μL precipitant solution. TipB crystallised in space group C 2 2 21 (a =106.6, b=126.2, c=78.0, α=90, β=90, γ=90) in a precipitant solution from the Morpheus screen consisting of 0.03 M Diethylene glycol; 0.03 M Triethylene glycol; 0.03 M Tetraethylene glycol; 0.03 M Pentaethylene glycol, 0.1 M Bicine, 0.1 M Tris, pH 8.5, 20% v/v PEG 500 MME, 10 % PEG 20000. Diffraction data were collected at Diamond Light Source. Data reduction and processing were performed through the Autoproc suite, and Aimless.

TipAB crystallisation screens were prepared by mixing a 0.5 μL of 8 mg/mL TipA and 9.5 mg/mL TipB (approx. 5X molar ratio) and 0.5 μL precipitant solution. The TipAB complex crystallised in space group C 2 2 21 (a =89.3, b=230.7, c=131.6, α=90, β=90, γ=90) in a precipitant solution from the Proplex screen consisting of 0.1 M Sodium acetate pH 5, 0 2 % w/v PEG 4000 15 % v/v MPD. Diffraction data were collected at European Synchrotron Radiation Facility. Data reduction and processing were performed through the Xia2-Dials pipeline.

Structure determination and model building

Diffraction data processing and refinement statistics are given in Table EV1.

The crystal structure of Selenomethionine-labelled TipA was solved using Phenix, with manual
interpretation and improvement of non-crystallographic symmetry operators.

The crystal structure of TipB was solved by molecular replacement with a Colabfold model
[48] in Phenix MR [49] Likewise, TipA$_{\Delta hairpin}$ was solved by molecular replacement with the
C-terminal domain of the selenomethionine-labelled TipA and manual boot-strap building and
refinement of the N-terminal domain. The complex TipAB structure was solved using
molecular replacement of the TipB structure with the interaction loops removed, and the C-
terminal domain of the selenomethionine-labelled TipA structure. The N-terminal domain was
also manually built through boot-strap building and refinement. All structures were refined
through iterative cycles of real-space building in Coot [50] and refinement in Phenix Refine
[51].

Pellet binding assays

Untagged TipA was expressed in *Escherichia coli* BL21 RIPL (DE3). Cells were grown in 10
mL LB supplemented with kanamycin to an OD$_{600}$ of 0.5-0.7 and induced with 0.5 mM IPTG
at 18 °C for 16 h. 5 X 1 mL of cells were centrifuged at 13000 RPM for 15 seconds and the
supernatant was removed. The cells were resuspended in 500 μL of either 0.15 M NaCl, 0.05
M HEPES pH 7.5; 0.5 M NaCl, 0.05 M HEPES pH 7.5; 1 M NaCl, 0.05 M HEPES pH 7.5;
0.15 M NaCl, 0.05 M HEPES pH 7.5, 0.05% Triton X100; 0.15 M NaCl, 0.05 M HEPES pH
7.5, 0.05% Tween 20. sonicated for 15 seconds and centrifuged at 13000 RPM for 15 minutes.
The pellet obtained from the 0.15 M NaCl buffer was resuspended in 500 μL SDS-PAGE
loading buffer for the pellet control. The pellet control and supernatant of each buffer was run
on SDS-PAGE.

QCM-D

QCM-D was performed using a Q-sense Analyzer QCM-D System (Biolin Scientific, Gothenburg, Sweden). Silicon QCM-D sensors were fixed into the QCM-D flow cell at a constant temperature of 20 °C and attached to a peristaltic pump. ddH$_2$O was pumped through the flow cell at a constant rate of 0.1 ml/min, a rate which was subsequently maintained throughout the rest of the experiment. Frequency and dissipation changes (Δf and ΔD) were monitored using multiple harmonics (n = 3, 5, 7, 9, 11, 13) of the resonant frequency. For simplicity we report here only the changes to the 3rd harmonic. Prior to data acquisition flow cells were equilibrated in H$_2$O for 5 mins to allow for a stable baseline to be acquired. H$_2$O was then exchanged for buffer, which was pumped through the flow cell until the change in frequency of the 3rd harmonic was stabilised. Bilayers consisting of either POPC, *E. coli* polar lipids or POPC:LPS (LPS from *E. coli* O111:B4) were then deposited onto the silicon sensor. This initially involves the formation of small unilamellar vesicles (SUVs), which were prepared by resuspension of a desiccated POPC/E. coli Polar Lipids/POPC:LPS film in buffer in 150 mM NaCl, 20 mM HEPES pH 7.5 to a concentration of 0.2 mg/mL and subsequent sonication for 30 min until an optically transparent solution was obtained. POPC:LPS SUVs were prepared at a 10:1 (w/w) ratio. These SUVs were subsequently flowed over the silicon sensor and then ruptured through osmotic shock with H2O, resulting in the characteristic formation of a deposited lipid bilayer. Exchange back into buffer 150 mM NaCl, 20 mM HEPES pH 7.5 aids the formation of a stable bilayer, with Δf3 ranging between -35 to -45 Hz. Once bilayers were deposited and baselines stabilized, TipA was introduced into the flow cell at a concentration of 1 mg/ml and flowed over the deposited bilayers for ~45 mins. Excess TipA was then removed by washing sensors in 150 mM NaCl, 20 mM HEPES pH 7.5. QCM-D with TipB was performed as above for TipA with POPC bilayers.

An *E. coli* Polar Lipid:LPS (10:1 w/w) mixed bilayer was also attempted to be formed and deposited onto silicon surfaces as outlined above. These bilayers were unable to be successfully generated, with SUVs unable to rupture and subsequently washed from the sensor. These surfaces were treated as 'unformed bilayers' and used as control surfaces to monitor TipA interaction with empty silicon sensors.

ITC

To test whether TipB binds to calcium, 5 mM EDTA and 5 mM EGTA was added to the protein

following size exclusion. After 16 hours the protein was passed through a Superdex 75 pg

26/60 column (GE Healthcare), which was pre-equilibrated with 20 mM HEPES, 150 mM

dialysed against 1 L of 20 mM HEPES, 150 mM NaCl, pH 7.5. $CaCl_2$ was dissolved in the

dialysis buffer to a final concentration of 500 μM. ITC was performed by titrating 500 μM

$CaCl_2$ into 50 μM TipB in 13 injections (1 x 0.4, 12 x 3 μL injections).

To test whether TipB binds to TipA-His, purified protein was dialysed against 1 L of 20 mM

HEPES, 150 mM NaCl, pH 7.5. ITC was performed by titrating 200 μM TipB into 20 μM TipA

in 13 injections (1 x 0.4, 12 x 3 μL injections).

TLC

TLC was employed to assess the lipid binding ability of TipA. Lipids were extracted from

protein samples through chloroform/methanol extraction. Briefly, 2 ml of Bd2538 at 0.5 mg/ml

was added to 2 ml methanol and 1 ml chloroform (2:2:1, v/v/v). Samples were vortexed for 5

min, followed by a 30 min incubation at 50 °C and then vortexed again for 5 min.

Centrifugation at 2,000 x g for 10 min resulted in the formation of two phases, the lower

chloroform layer containing lipidic material. This lower phase was extracted and then

evaporated. The dried lipids were resuspended in 100 μl chloroform with 5 μl then loaded onto

a Silica TLC plate (Sigma) and run in a 6.5:2.5:1 (chloroform:methanol:acetic acid) solvent

system. The TLC plate was dried, stained with 10% (w/v) phosphomolybdic acid in ethanol,

and finally heated until stained spots could be visualised.

Figure 1. The structure of *Bdellovibrio* TULIP protein TipA

(A) Schematic of the forward strand ORFs for the gene operon of *bd2538* (green) and *bd2539* (blue). (B) 2Fo-Fc map at 2.5 σ surrounding chain C of the crystal structure of TipA after the final refinement. (C) Top; Crystal structure of TipA, showing the double-TULIP fold of the Bactericidal permeability-increasing protein, alpha/beta domain superfamily, coloured by secondary structure (beta – red, helix – light blue, coil – white). The lengths of the protein are shown for all three dimensions. Middle; a surface view of TipA with hydrophobic residues shown

in stick form, highlighting the hydrophobic nature of the central cavity. Bottom; A clipped surface view showing the inner cavity. **(D)** Top;

structural features of the N-terminal TipA TULIP domain, with the dimerisation motif coloured blue. The expanded view shows residues in

stick form that cause the N-terminal Helix A to protrude. Bottom; The structure of the N-terminal TULIP domain of bacterial permeability

increasing protein (BPI, code 1BP1) for comparison. **(E)** Top; structural features of the C-terminal TipA TULIP domain, with the helix

positions highlighted with arrows. The expanded view shows residues in stick form that cause the N-terminal Helix A to protrude. Bottom;

The structure of the C-terminal TULIP domain of BPI for comparison.

Figure 2. Dimerisation of TipA into a four TULIP-long assembly

(A) Crystal structure of TipA, showing two chains forming an end-on-end homodimer via interaction of their N-terminal TULIP domains.
One protomer is shown in light blue and the other is in green with the 2-fold symmetry axis shown as a black oval and dotted line. **(B)** Upper;
the dimer forms an elongated cavity within (grey surface, calculated by Mole [52]). Lower; the electrostatic surface view shows that the dimer
forms a continuous tube with very few solvent accessible gaps. **(C)** The view down the two-fold axis shows how the two interdigitated
dimerisation hairpins (light blue and green) create a polar surface on the outside of the TULIP dimer and a hydrophobic inner face for a
continuous hydrophobic channel. **(D)** The dimerisation helix forms polar and hydrophobic interactions with the other protomer's Helix B and
dimerisation helix. **(E)** Mutation of the dimerisation hairpin causes a shift in the elution volume of TipA on gel filtration from 165 mL to 200
mL. **(F)** The crystal structure of TipA$_{\Delta Hairpin}$ (magenta) superimposed onto the TipA wild type (WT) structure (green). The C-terminal domain
overlays well, but the N-terminal domain is contracted due to twisting of the beta-sheet and bending of the N-terminal Helix B at G277.

663
664
665
666

Figure 3. Crystal structure of the TipB adapter protein

(A) Crystal structure of TipB, coloured by secondary structure (beta – red, helix – light blue, coil – white) with the two specialised interaction loops coloured yellow. Key hydrophobic amino acids in the interaction loops are shown in stick form. **(B)** The three chains of TipB in the ASU are superposed to highlight flexibility in the interaction loops (shown blue, green and yellow for each chain). **(C)** The topology of TipB, with strands in red, helices as blue ovals and loops as blue lines. **(D)** The topology of type II C2 domains. **(E)** Crystal structures of proteins identified in structural homology searches with DALI [24], shown in same orientation as TipB (left, light blue). Arginines that form the basic face of the sheet in *Pseudomonas aeruginosa* ExsB [26] (coloured salmon) are shown in stick form; the sheet capping loop of *Porphyromonas gingivalis* [53] (coloured pink) is shown in yellow; and the two calcium ions of human phospholipase A2 [35] (coloured green) are shown as

675   spheres. **(F)** QCM-D trace showing f3 and d3 signals. Upon protein addition, no significant change in f3 or d3 is observed. **(G)** ITC thermogram
676   of a titration of calcium chloride into TipB showing no binding.

678
Figure 4. The interaction of the TipA and TipB TULIP:adapter pairing

(A) ITC thermogram of a titration of TipB into TipA, showing binding of the two proteins in solution. The reported energies for binding are
shown as an embedded bar graph. **(B)** Crystal structure of the TipAB complex. Tip A is shown in green and forms the same 2-fold dimer as
the isolated form through a crystallographic axis. TipB is shown in blue on top of TipA. **(C)** A surface view of TipA is shown, highlighting
burial of the two interaction loops of TipB (coloured yellow). **(D)** A clipped surface view of the TipAB complex, showing TipA sat on the
TipA N-terminal helix A. **(E)** A cartoon schematic of the interaction with parts of the TipA N-terminal TULIP domain secondary structure
visible, and key hydrophobic residues of TipB shown as sticks. **(F)** Hydrophobic interactions of the TipB interaction loop 1. **(G)** Hydrophobic

interactions of the TipB interaction loop 2. **(H)** polar interactions of the TipB interaction loop 1. **(I)** Polar interactions of the TipB interaction
loop 2.

689
Figure 5. Conformational change of TipA when bound to the TipB adapter

(A) A superimposition of apo TipA (magenta) with the TipAB complex. Large motions in the N-terminal domain of TipA are visible. **(B)**
contact plots of TipA for the apo TipA and TipAB structures, generated by DISTEVAL [54]. Both plots have a box surrounding residues that
differ significantly between the two structures, which are only found in the N-terminal domain. **(C)** Three orientations of the apo TipA
homodimer (one protomer green, one protomer red), showing a mostly-closed tube. **(D)** The same three orientations of the TipA homodimer
as **(C)** for the TipAB complex. TipB is shown in blue, and the large opening of the TULIP homodimer stretching between the two TipB
proteins is visible. **(E)** The clamp formed by interaction loop 2 and charged residues grasp the bottom and top of the TipA N-terminal beta
sheet, respectively. **(F)** Analysis of the apo TipA and complexed TipA structure with DynDom to investigate the motions of the conformational
change identifies three domains (coloured green, orange and blue), with motions in the hinge regions (magenta). **(G)** Upper; superpositions
show the 29° rotation between Domain 1 and Domain 2 rotate by 29. Lower; superpositions show the 36° movement between domain 2 and
domain 3.

701
702

703
Figure 6. TipA can insert into a range of lipid bilayers

(A) Pellet-binding assays with TipA. SDS-PAGE of pellet and supernatants from various buffer washes. A prominent band for soluble TipA
is observed in high salt concentrations or with the presence of detergents. **(B)** QCM-D measurements showing the changes in frequency (blue)
and dissipation (red) in response to the mass deposition associated with *E. coli* polar lipid bilayer formation (polar lipid liposomes were
deposited onto a silicon surface and ruptured through osmotic shock), followed by the addition of TipA, and subsequent bilayer washing with
excess buffer. **(C)** QCM-D measurements as specified in **(A)** showing POPC bilayer formation, TipA addition, and buffer wash. **(D)** QCM-D
measurements as specified in **(A)** indicating the formation of a stable POPC:LPS bilayer, TipA addition and subsequent buffer wash. **(E)** A
comparison of the changes to frequency (indicative of mass changes), as measured through QCM-D, in response to the addition of TipA to
various lipid bilayer compositions (1) and subsequent buffer wash (2). Note we were unable to successfully form a mixed polar lipid, LPS
bilayer.

Figure 7. TipA sequesters lipids in the TULIP lumen

(A) TLC of lipids extracted from TipA. All three *E. coli* polar lipids are visible in the TipA lane (cardiolipin (CL), phosphatidylglycerol (PG),

phosphatidylethanolamine (PE)), as are some additional species. **(B)** Crystal structure of the TipAB complex, with two lipid head groups

shown as spheres (magenta carbons). **(C)** Crystal structure of TipA$_{\Delta Hairpin}$, with two modelled PE molecules shown as spheres (magenta

carbons). **(D)** 2Fo-Fc electron density of the two lipids in the TipA$_{\Delta Hairpin}$ structure at 0.8 σ. **(E)** The molecular interactions of TipAB complex

and the lipid head groups for the inner PE molecule (top) and outer PE molecule (bottom). TipA is coloured green, TipB is blue, with a

symmetry TipA copy coloured salmon. Lipids are shown in stick form and coloured as heteroatoms (carbon – magenta, oxygen – red, nitrogen

– blue, phosphorus – orange) **(F)** The molecular interactions of TipA$_{\Delta Hairpin}$ and the lipid head groups for the inner PE molecule (top) and outer

PE molecule (bottom). TipA is coloured green with a symmetry TipA copy coloured salmon. Lipids are shown in stick form and coloured as

heteroatoms (carbon – magenta, oxygen – red, nitrogen – blue, phosphorus – orange)

Figure EV2. Additional domains fused to TipA structural homologues

(A) Using Foldseek [38], structural TipA homologues were identified within the AlphaFold [58] AFDB50 database. A number of those homologues had additional domains to the double-TULIP. Representative examples of fusions are shown and comprise beta-rich domains (upper) helical domains (middle) and other domains including second double-TULIPs and enzymes (lower). The double-TULIP domains are coloured light blue, and the accessory domains are coloured orange. Of particular interest is the *Planctomyces sp*. SH-PL14 protein (bottom right) in which there is a similar beta-sandwich domain to TipB fused C-terminally to the TULIP (B) Superposition of the TipAB with *Planctomyces sp*. SH-PL14 (Uniprot accession: A0A142WTQ2). TipA is coloured green, TipB is blue, the TULIP domains of A0A142WTQ2 are white and the beta sandwich is orange. The enzymatic domain has been removed for clarity. The beta-sandwiches interact with the N-terminal Helix A in both structures and make interactions with the top of the N-terminal TULIP sheet. (C) Superposition of TipB (blue) with the beta sandwich of A0A142WTQ2 (orange), showing similar topology.

 Figure EV3. Lipid binding and higher order packing of TipA(A) Fo-Fc electron density (blue mesh)

observed in the apo selenomethionine TipA structure. Large peaks are observed in similar regions to lipid density in the TipAB complex and

TipA$_{\Delta Hairpin}$. **(B)** Crystal packing in the apo selenomethionine structure. **(C)** Crystal packing in the TipAB complex structure. **(D)** Crystal

packing in the TipA$_{\Delta Hairpin}$ structure. In all structures shown **(B-D)** the C-terminal domain of TipA buries into the N-terminal domain of another

TipA protein. **(E)** Molecular interactions of the C-terminal domain and N-terminal of the apo selenomethionine TipA structure. **(F)** Molecular

interactions of the C-terminal domain and N-terminal of the TipAB structure. **(G)** Molecular interactions of the C-terminal domain and N-

terminal of the TipA$_{\Delta Hairpin}$ structure.

770 Table EV1. Data collection and refinement statistics

	Apo SeMet TipA	TipB	TipAB complex	TipA$_{\Delta Hairpin}$
Wavelength		0.979 Å	0.968 Å	0.968 Å
Resolution range	48.74 - 4.046 (4.191 - 4.046)	29.24 - 2.17 (2.23 - 2.17)	57.66 - 2.6 (2.693 - 2.6)	66.01 - 2.6 (2.72 - 2.6)
Space group	P 31 2 1	C 2 2 21	C 2 2 21	P 21 21 21
Unit cell	252.415 252.415 181.058 90 90 120	106.578 126.162 77.975 90 90 90	89.2741 230.652 131.606 90 90 90	57.9688 97.8339 132.021 90 90 90
Total reflections		376137	596458	307502
Unique reflections	45222 (538)	28055 (2231)	42141 (4178)	23555 (2618)
Multiplicity		13.4 (11.7)	14.1 (13.7)	13.1 (9.0)
Completeness (%)	82.85 (9.99)	99.30 (91.9)	98.36 (85.22)	98.8 (91.5)
Mean I/sigma(I)		16.8 (3.3)	11.6 (0.3)	18.6 (2.7)
Wilson B-factor	188.01	35.30	86.67	60.45
R-merge		0.102 (0.733)	0.117 (3.31)	0.083 (0.681)
R-meas		0.110 (0.798)	0.125 (3.55)	0.089 (0.770)
R-pim		0.041 (0.312)	0.034 (0.946)	0.033 (0.350)
CC1/2		0.999 (0.896)	0.999 (0.333)	0.999 (0.832)
Reflections used in refinement	45184 (538)	28025 (2640)	41504 (3563)	23500 (2098)
Reflections used for R-free	2364 (23)	1360 (115)	2105 (196)	1132 (117)
R-work	0.2766 (0.4538)	0.1774 (0.2220)	0.2463 (0.3946)	0.2283 (0.2910)
R-free	0.3200 (0.5542)	0.2160 (0.2798)	0.2874 (0.4209)	0.2526 (0.3627)
Number of non-hydrogen atoms	12575	3394	5382	4178
macromolecules	12575	3196	5267	3990
ligands	0	0	55	135
solvent	0	198	60	53
Protein residues	1600	415	674	507
RMS(bonds)	0.005	0.009	0.005	0.007
RMS(angles)	1.33	1.01	0.75	1.04
Ramachandran favored (%)	79.99	96.58	94.48	93.01
Ramachandran allowed (%)	18.51	2.93	5.07	6.39
Ramachandran outliers (%)	1.51	0.49	0.45	0.60
Rotamer outliers (%)	0.58	0.55	2.25	3.22
Clashscore	30.35	5.33	12.23	13.31
Average B-factor	239.19	45.22	110.85	83.37
macromolecules	239.19	45.05	111.03	83.25
ligands			115.13	96.48
solvent		47.96	91.11	59.35
Number of TLS groups	1	1	16	1

771   Statistics for the highest-resolution shell are shown in parentheses.

References

1. Negus, D., et al., *Predator Versus Pathogen: How Does Predatory Bdellovibrio bacteriovorus Interface with the Challenges of Killing Gram-Negative Pathogens in a Host Setting?* Annu Rev Microbiol, 2017. **71**: p. 441-457.
2. Bratanis, E., et al., *Biotechnological Potential of Bdellovibrio and Like Organisms and Their Secreted Enzymes.* Front Microbiol, 2020. **11**: p. 662.
3. Rendulic, S., et al., *A predator unmasked: life cycle of Bdellovibrio bacteriovorus from a genomic perspective.* Science, 2004. **303**(5658): p. 689-92.
4. Matin, A. and S.C. Rittenberg, *Kinetics of deoxyribonucleic acid destruction and synthesis during growth of Bdellovibrio bacteriovorus strain 109D on pseudomonas putida and escherichia coli.* J Bacteriol, 1972. **111**(3): p. 664-73.
5. Hespell, R.B., G.F. Miozzari, and S.C. Rittenberg, *Ribonucleic acid destruction and synthesis during intraperiplasmic growth of Bdellovibrio bacteriovorus.* J Bacteriol, 1975. **123**(2): p. 481-91.
6. Thomashow, M.F. and S.C. Rittenberg, *Intraperiplasmic growth of Bdellovibrio bacteriovorus 109J: solubilization of Escherichia coli peptidoglycan.* J Bacteriol, 1978. **135**(3): p. 998-1007.
7. Rosson, R.A. and S.C. Rittenberg, *Regulated breakdown of Escherichia coli deoxyribonucleic acid during intraperiplasmic growth of Bdellovibrio bacteriovorus 109J.* J Bacteriol, 1979. **140**(2): p. 620-33.
8. Nelson, D.R. and S.C. Rittenberg, *Incorporation of substrate cell lipid A components into the lipopolysaccharide of intraperiplasmically grown Bdellovibrio bacteriovorus.* J Bacteriol, 1981. **147**(3): p. 860-8.
9. Kuenen, J.G. and S.C. Rittenberg, *Incorporation of long-chain fatty acids of the substrate organism by Bdellovibrio bacteriovorus during intraperiplasmic growth.* J Bacteriol, 1975. **121**(3): p. 1145-57.
10. Abram, D., J. Castro e Melo, and D. Chou, *Penetration of Bdellovibrio bacteriovorus into host cells.* J Bacteriol, 1974. **118**(2): p. 663-80.
11. Banks, E.J., et al., *Asymmetric peptidoglycan editing generates cell curvature in Bdellovibrio predatory bacteria.* Nat Commun, 2022. **13**(1): p. 1509.
12. Kuru, E., et al., *Fluorescent D-amino-acids reveal bi-cellular cell wall modifications important for Bdellovibrio bacteriovorus predation.* Nat Microbiol, 2017. **2**(12): p. 1648-1657.
13. Giacometti, S.I., et al., *Lipid Transport Across Bacterial Membranes.* Annu Rev Cell Dev Biol, 2022. **38**: p. 125-153.
14. Reinisch, K.M. and P. De Camilli, *SMP-domain proteins at membrane contact sites: Structure and function.* Biochim Biophys Acta, 2016. **1861**(8 Pt B): p. 924-927.
15. Alva, V. and A.N. Lupas, *The TULIP superfamily of eukaryotic lipid-binding proteins as a mediator of lipid sensing and transport.* Biochim Biophys Acta, 2016. **1861**(8 Pt B): p. 913-923.
16. Wong, L.H. and T.P. Levine, *Tubular lipid binding proteins (TULIPs) growing everywhere.* Biochim Biophys Acta Mol Cell Res, 2017. **1864**(9): p. 1439-1449.
17. Levine, T.P., *Remote homology searches identify bacterial homologues of eukaryotic lipid transfer proteins, including Chorein-N domains in TamB and AsmA and Mdm31p.* BMC Mol Cell Biol, 2019. **20**(1): p. 43.
18. Barrio-Hernandez, I., et al., *Clustering predicted structures at the scale of the known protein universe.* Nature, 2023. **622**(7983): p. 637-645.

19. Gustafsson, R., et al., *Crystal structures of OrfX2 and P47 from a Botulinum neurotoxin OrfX-type gene cluster.* FEBS Lett, 2017. **591**(22): p. 3781-3792.

20. Gao, L., et al., *Crystal structures of OrfX1, OrfX2 and the OrfX1-OrfX3 complex from the orfX gene cluster of botulinum neurotoxin E1.* FEBS Lett, 2023. **597**(4): p. 524-537.

21. Lambert, C., et al., *The first bite--profiling the predatosome in the bacterial pathogen Bdellovibrio.* PLoS One, 2010. **5**(1): p. e8599.

22. Tyson, J., et al., *Prey killing without invasion by Bdellovibrio bacteriovorus defective for a MIDAS-family adhesin.* Nat Commun, 2024. **15**(1): p. 3078.

23. Paysan-Lafosse, T., et al., *InterPro in 2022.* Nucleic Acids Res, 2023. **51**(D1): p. D418-D427.

24. Holm, L., *Dali server: structural unification of protein families.* Nucleic Acids Res, 2022. **50**(W1): p. W210-W215.

25. Voss, N.R. and M. Gerstein, *3V: cavity, channel and cleft volume calculator and extractor.* Nucleic Acids Res, 2010. **38**(Web Server issue): p. W555-62.

26. Izore, T., et al., *Structural characterization and membrane localization of ExsB from the type III secretion system (T3SS) of Pseudomonas aeruginosa.* J Mol Biol, 2011. **413**(1): p. 236-46.

27. Nalefski, E.A. and J.J. Falke, *The C2 domain calcium-binding motif: structural and functional diversity.* Protein Sci, 1996. **5**(12): p. 2375-90.

28. Kohout, S.C., et al., *C2 domains of protein kinase C isoforms alpha, beta, and gamma: activation parameters and calcium stoichiometries of the membrane-bound state.* Biochemistry, 2002. **41**(38): p. 11411-24.

29. Nalefski, E.A., et al., *Delineation of two functionally distinct domains of cytosolic phospholipase A2, a regulatory Ca(2+)-dependent lipid-binding domain and a Ca(2+)-independent catalytic domain.* J Biol Chem, 1994. **269**(27): p. 18239-49.

30. Saheki, Y. and P. De Camilli, *The Extended-Synaptotagmins.* Biochim Biophys Acta Mol Cell Res, 2017. **1864**(9): p. 1490-1493.

31. Dixon, M.C., *Quartz crystal microbalance with dissipation monitoring: enabling real-time characterization of biological materials and their interactions.* J Biomol Tech, 2008. **19**(3): p. 151-8.

32. Hughes, G.W., et al., *Evidence for phospholipid export from the bacterial inner membrane by the Mla ABC transport system.* Nat Microbiol, 2019. **4**(10): p. 1692-1705.

33. Richter, R., A. Mukhopadhyay, and A. Brisson, *Pathways of lipid vesicle deposition on solid surfaces: a combined QCM-D and AFM study.* Biophys J, 2003. **85**(5): p. 3035-47.

34. Shen, H.H., et al., *Reconstitution of a nanomachine driving the assembly of proteins into bacterial outer membranes.* Nat Commun, 2014. **5**: p. 5078.

35. Perisic, O., et al., *Crystal structure of a calcium-phospholipid binding domain from cytosolic phospholipase A2.* J Biol Chem, 1998. **273**(3): p. 1596-604.

36. Krissinel, E. and K. Henrick, *Inference of macromolecular assemblies from crystalline state.* J Mol Biol, 2007. **372**(3): p. 774-97.

37. Lee, R.A., M. Razaz, and S. Hayward, *The DynDom database of protein domain motions.* Bioinformatics, 2003. **19**(10): p. 1290-1.

38. van Kempen, M., et al., *Fast and accurate protein structure search with Foldseek.* Nat Biotechnol, 2024. **42**(2): p. 243-246.

39. Schauder, C.M., et al., *Structure of a lipid-bound extended synaptotagmin indicates a role in lipid transfer.* Nature, 2014. **510**(7506): p. 552-5.

40. AhYoung, A.P., et al., *Conserved SMP domains of the ERMES complex bind phospholipids and mediate tether assembly.* Proc Natl Acad Sci U S A, 2015. **112**(25): p. E3179-88.

879   41.   Ekiert, D.C., et al., *Architectures of Lipid Transport Systems for the Bacterial Outer*
880   *Membrane.* Cell, 2017. **169**(2): p. 273-285 e17.
881   42.   Kumar, S. and N. Ruiz, *Bacterial AsmA-Like Proteins: Bridging the Gap in*
882   *Intermembrane Phospholipid Transport.* Contact (Thousand Oaks), 2023. **6**: p.
883   25152564231185931.
884   43.   Yu, H., et al., *Extended synaptotagmins are Ca2+-dependent lipid transfer proteins at*
885   *membrane contact sites.* Proc Natl Acad Sci U S A, 2016. **113**(16): p. 4362-7.
886   44.   Jeong, H., J. Park, and C. Lee, *Crystal structure of Mdm12 reveals the architecture and*
887   *dynamic organization of the ERMES complex.* EMBO Rep, 2016. **17**(12): p. 1857-1871.
888   45.   Suzuki, R., et al., *Structural mechanism of JH delivery in hemolymph by JHBP of*
889   *silkworm, Bombyx mori.* Sci Rep, 2011. **1**: p. 133.
890   46.   Weiss, J., et al., *Purification and characterization of a potent bactericidal and*
891   *membrane active protein from the granules of human polymorphonuclear leukocytes.* J
892   Biol Chem, 1978. **253**(8): p. 2664-72.
893   47.   Toulmay, A. and W.A. Prinz, *A conserved membrane-binding domain targets proteins*
894   *to organelle contact sites.* J Cell Sci, 2012. **125**(Pt 1): p. 49-58.
895   48.   Mirdita, M., et al., *ColabFold: making protein folding accessible to all.* Nat Methods,
896   2022. **19**(6): p. 679-682.
897   49.   McCoy, A.J., *Solving structures of protein complexes by molecular replacement with*
898   *Phaser.* Acta Crystallogr D Biol Crystallogr, 2007. **63**(Pt 1): p. 32-41.
899   50.   Emsley, P. and K. Cowtan, *Coot: model-building tools for molecular graphics.* Acta
900   Crystallogr D Biol Crystallogr, 2004. **60**(Pt 12 Pt 1): p. 2126-32.
901   51.   Afonine, P.V., et al., *Towards automated crystallographic structure refinement with*
902   *phenix.refine.* Acta Crystallogr D Biol Crystallogr, 2012. **68**(Pt 4): p. 352-67.
903   52.   Sehnal, D., et al., *MOLE 2.0: advanced approach for analysis of biomacromolecular*
904   *channels.* J Cheminform, 2013. **5**(1): p. 39.
905   53.   Hall, M., et al., *Structural and functional characterization of shaft, anchor, and tip*
906   *proteins of the Mfa1 fimbria from the periodontal pathogen Porphyromonas gingivalis.*
907   Sci Rep, 2018. **8**(1): p. 1793.
908   54.   Adhikari, B., et al., *DISTEVAL: a web server for evaluating predicted protein*
909   *distances.* BMC Bioinformatics, 2021. **22**(1): p. 8.
910   55.   Altschul, S.F., et al., *Basic local alignment search tool.* J Mol Biol, 1990. **215**(3): p.
911   403-10.
912   56.   Sievers, F. and D.G. Higgins, *Clustal Omega for making accurate alignments of many*
913   *protein sequences.* Protein Sci, 2018. **27**(1): p. 135-145.
914   57.   Robert, X. and P. Gouet, *Deciphering key features in protein structures with the new*
915   *ENDscript server.* Nucleic Acids Res, 2014. **42**(Web Server issue): p. W320-4.
916   58.   Jumper, J., et al., *Highly accurate protein structure prediction with AlphaFold.* Nature,
917   2021. **596**(7873): p. 583-589.
918

Acknowledgements
ALL, SGC, and ITC were funded by the Wellcome Trust Investigator Award in Science
(209437/Z/17/Z); AL and SGC acknowledge additional funding from a BBSRC award
(BB/X006298/1). GH and TJK were funded by BBSRC award BB/S017283/1, and HJ by
BBSRC studentship BB/T00746X/1. We are grateful to Paul Radford and Liz Sockett for
attempts to monitor functional role and localization in vivo. We thank Diamond Light Source
for beamtime and for facility support during data collection.
926