An Aurora Kinase A/TPX2 complex phosphorylates CKAP2 to control mitotic spindle growth

Thomas J Kucharski[1], Tatiana Lyalina[1] and Susanne Bechstedt[1,2*]

1) Department of Anatomy and Cell Biology, McGill University, Québec, Montréal, Canada, H3A 0C7

2) Centre de recherche en biologie structurale, McGill University, Québec, Montréal, Canada, H3G 0B1

*Lead Contact

Correspondence address:

Department of Anatomy and Cell Biology

3640 Rue University W-309

Montréal, Québec

Canada

H3A 0C7

Keywords: microtubule, mitosis, kinase, mitotic spindle, phosphorylation, Aurora, phosphatase

Summary

The formation of the mitotic spindle is a critical step in the equal separation of DNA from the

mother cell to the two daughter cells. Cytoskeleton Associated Protein-2 (CKAP2) is a

microtubule polymerase that associates with microtubules and centrosomes during interphase

and mitosis. Loss of CKAP2 results in chromosomal instability and onset of aneuploidy. To

identify proteins interacting with CKAP2, we immunoprecipitated endogenous CKAP2 together

with interacting proteins from mitotic RPE1 cells and analysed the samples by mass

spectrometry. This experiment identified the mitotic protein kinase Aurora Kinase A, together

with its cofactor TPX2, as interacting proteins of CKAP2. Notably, CKAP2 co-localizes and

interacts with Aurora Kinase A and TPX2 throughout mitosis, but not with the related protein

Aurora Kinase B. Consistently, Aurora Kinase A phosphorylates CKAP2 both in mitotic cells

and *in vitro*. Phosphorylated CKAP2 has reduced affinity for microtubules *in vitro*. Importantly,

expression of Aurora A phospho-mimic mutants of CKAP2 in RPE1 cells results in phenotypes

of a small mitotic spindle and reduced microtubule stability. Thus, Aurora Kinase A

phosphorylates CKAP2 to regulate mitotic spindle growth and stability.

Introduction

The objective of mitosis is the equal partitioning of chromosomes into the two daughter cells. Upon initiation of mitosis, the nuclear envelope breaks down, and the mitotic spindle is formed. The mitotic spindle is comprised of bundles of microtubules (MTs), which are polymers of tubulin subunits[1]. During early mitosis, MTs attach to chromosomes at the kinetochore and pull them toward the metaphase plate. When all chromosomes are correctly attached to the spindle and aligned on the metaphase plate, such that both sister kinetochores are attached to MTs emanating from opposite spindle poles, the MTs then pull the sister chromatids apart into the two daughter cells[2,3]. Importantly, MTs are highly dynamic structures which can grow or shrink to generate pulling forces on chromosomes[1]. The growth and shrinkage rates of MTs must be tightly controlled to correctly align and segregate the chromosomes during mitosis[4–7]. The growth and shrinkage rates are controlled by a group of proteins known as Microtubule-Associated Proteins (MAPs)[1]. Amongst MAPs, the protein CKAP2 (alias TMAP) has generated substantial interest because it potently induces MT growth, its overexpression contributes to the onset of chromosomal instability and aberrant centrosome numbers, and its overexpression is associated with worse survival in cancer patients[8–14].

Studies on CKAP2 have revealed that the protein is phosphorylated on multiple sites during mitosis by both Aurora and Cyclin Dependent Kinases (CDKs). Phosphorylation of CKAP2 on conserved c-terminus sites by these kinases has been proposed to control the subcellular localization of CKAP2 during mitosis[8,15–19]. From early mitosis through anaphase, CKAP2 is localized on centrosomes and pole-proximal MTs. However, in late anaphase, CKAP2 abruptly changes localization and binds to chromosomes before it is finally degraded through the action of the Anaphase Promoting Complex/Cyclosome ubiquitin ligase during telophase/G1[8,15–

[20]. Thus, phosphorylation/dephosphorylation of CKAP2 has been proposed to function as a

switch to control the observed changes in protein localization. Phosphorylation sites in the n-

terminus of CKAP2 have also been detected in large-scale proteomics studies[21–23], but roles for

these phosphorylation sites have not yet been determined.

To gain insights into the molecular mechanisms that govern CKAP2 function, we

performed unbiased proteomic screening to identify CKAP2-interacting proteins. This approach

identified TPX2 and Aurora Kinase A (AurKA) as important interacting proteins of CKAP2. We

found that AurKA/TPX2 binds to and phosphorylates CKAP2 during mitosis to regulate its MT

binding and polymerase activity and thus control mitotic spindle growth and mitotic fidelity.

These new findings effectively integrate CKAP2 into existing spindle regulation pathways and

shed light on the regulation of this potent MT growth factor.

Results

CKAP2 interacts with AurKA/TPX2 during mitosis

To identify proteins that associate with CKAP2 during mitosis, we immunoprecipitated

endogenous CKAP2 together with interacting proteins from normal RPE-1 cells that were

synchronized in mitosis by sequential Palbociclib-STLC block (Figure 1A). 672 proteins were

identified by mass spectrometry in either or both CKAP2 immunoprecipitation or control

immunoprecipitation. To identify proteins that specifically associate with CKAP2, we applied

criteria of at least 4 identified peptides, and a 4:1 signal-to-noise ratio of peptides observed in the

CKAP2 vs control immunoprecipitations. This resulted in a high-confidence list of 90 unique

proteins that associate with CKAP2 in mitosis. To identify pathways in which CKAP2 might

participate, we performed gene ontology analysis on the set of 90 proteins. This analysis revealed

that CKAP2 participates in various cell division processes, as determined previously, as well as

RNA-related processes (Figure 1B). Search Tool for the Retrieval of Interacting Genes/Proteins

interaction network analysis revealed that CKAP2 associates with proteins that participate in

mitosis/chromatin organization (e.g. NCAPG), mRNA processing (e.g. CPSF1), and possibly

translation (e.g. RSL1D1) (Figure 1C). Amongst MAPs, we identified KIF2A, TXLNA,

KIF20A, TPX2, and KIF23 as CKAP2 potential interacting proteins. We also identified AurKA

as a CKAP2-interacting protein, which, although not a MAP as such, exists in cells mainly as a

heterodimer with TPX2, which is a MAP. We also identified numerous chromatin-bound proteins

that might mediate the interaction between CKAP2 and chromatin in late mitosis, including

TOP2A, SMC2, SMC4, NCAPG, NCAPD2, NCAPH, MACROH2A1 and KI67 (Figure 1C, 1D,

supplemental table 1).

Amongst CKAP2 interacting proteins that we detected, AurKA/TPX2 were of particular

interest. Aurora Kinase A phosphorylates numerous substrates at spindle poles and on MTs

during mitosis that broadly contribute to mitotic spindle growth, stability and function[24–27]. TPX2

binds directly to MTs and functions as an AurKA cofactor to localize AurKA to MTs and spindle

poles[28–30]. AurKA has been shown to be overexpressed in numerous cancers and contributes to

oncogenesis[31–33]. Critically, AurKA does not localize to MTs in the absence of TPX2[28].

Additionally, TPX2 has been demonstrated to directly contribute to MT growth, suggesting that

it may function in conjunction with CKAP2 in this capacity[29,30,34]. We first confirmed that

CKAP2 binds to the AurKA/TPX2 complex. We performed another CKAP2

immunoprecipitation from Palbociclib/STLC synchronized RPE-1 cells, but this time performed

Western blot analysis to probe for the presence of TPX2 and AurKA. This experiment confirmed

that both TPX2 and AurKA interact with CKAP2 (Figure 1E). We also performed the reciprocal

experiment, where we immunoprecipitated TPX2 and blotted for CKAP2 and Aurora A. This

experiment revealed a robust interaction with AurKA, as expected, and with CKAP2 (Figure

S1A).

AurKA is closely related to another mitotic kinase, Aurora Kinase B (AurKB). These two

kinases share a consensus motif R/K-R/K-x-S/T. Thus, both kinases are generally capable of

phosphorylating the same substrates. However, while TPX2 localizes AurKA to spindle poles

and MTs, INCENP localizes AurKB to centromeres and kinetochores[24–27,35–37]. Thus, AurKA and

AurKB are spatially separated in cells, which governs their substrate sets. AurKB has been

previously suggested to phosphorylate CKAP2 during mitosis[19]. Therefore, we sought to

determine if CKAP2 co-localizes with AurKA/B during mitosis. We employed RPE-1 cells with

GFP-CKAP2 knock-in[8], and stained for AurKA and AurKA phosphorylated on T288, which is a

requirement for and marker of AurKA activity[38,39]. This experiment revealed clear co-

localization of CKAP2 with AurKA and AurKA PT288 at centrosomes in prophase cells and

throughout mitosis. CKAP2 also substantially colocalized with AurKA and AurKA PT288 on

pole-proximal MTs during prometaphase and metaphase (Figure 2A). Consistently, CKAP2 also

largely colocalized with TPX2 on pole-proximal MTs during prometaphase and metaphase

(Figure 2B) in accordance with our CKAP2 interactome and immunoprecipitation data (Figure 1,

supplemental table 1). In contrast to AurKA/TPX2, which were present at spindle poles and

MTs, AurKB localized to centromeres during prophase-metaphase, and then changed localization

to the midbody during anaphase. CKAP2 did not co-localize with AurKB at any time during

mitosis (Figure S1B, S1C).

AurKA/TPX2 phosphorylate CKAP2 during mitosis

We next sought to determine if CKAP2 contains AurK consensus motifs. We identified

possible AurK phosphorylation sites at amino acids 39, 76, 209 and 347 (equivalent to mouse

CKAP2 sites 52, 86, 206 and 337) plus the sites in the c-terminus that had already been

identified[16–18]. The n-terminal phosphorylation sites are largely conserved in mammals, but not

other species (Figure 2C). We next examined previously published proteomic datasets to see if

phosphorylation of these sites has been previously detected. T209 phosphorylation has been

detected in previous studies, confirming that this site is phosphorylated in cells[21–23]. We next

examined the phosphorylation state of CKAP2 purified from mitotic cells. We

immunoprecipitated CKAP2 from Palbociclib-STLC synchronized RPE1 cell lysates. We boiled

the samples in the presence of SDS to avoid loss of CKAP2 in insoluble chromatin/MT fractions.

To determine if CKAP2 is phosphorylated in mitotic cells, we treated the immunoprecipitated

protein +/- lambda protein phosphatase before separation on SDS-PAGE. We then stained the gel

with ProQ Diamond to detect phosphorylated species and finally with Coomassie blue to

determine total protein levels. This experiment revealed that CKAP2 is phosphorylated in mitotic

cells as the phosphorylation signal disappears following treatment with lambda phosphatase

(Figure 2D). We then performed a similar experiment but treated the cells concurrently with

STLC and either DMSO solvent control or the AurKA inhibitor MLN8237 prior to harvesting the

cells. Interestingly, this experiment revealed a substantial loss of phosphorylation in MLN8237

treated cells, suggesting that AurKA is a key regulator of CKAP2 phosphorylation (Figure 2E)

during mitosis. Lastly, we tested if AurKA can phosphorylate CKAP2 *in vitro*. Therefore, we

incubated mouse CKAP2 together with purified GFP-Aurora A-6XHis and examined the

phosphorylation state of CKAP2 with ProQ Diamond stain. This experiment shows that AurKA

phosphorylates CKAP2 *in vitro* in an ATP and kinase-dependent manner (Figure 2F). Taken

together, these data demonstrate that CKAP2 colocalizes with, binds to and is phosphorylated by

AurKA/TPX2 during mitosis.

We next asked which protein phosphatase might antagonize CKAP2 phosphorylation by

AurKA. Obvious candidates to regulate CKAP2 include the PP1 and PP2A phosphatases.

Okadaic acid inhibits PP2A and PP1 to a lesser, though considerable extent[40,41]. Therefore, we

treated RPE1 cells with okadaic acid to see if it would affect the extent of CKAP2 binding to

MTs in mitotic cells. Although Okadaic acid treatment had a drastic effect on mitotic spindle

morphology owing to hyperstable kinetochore-MT attachments, CKAP2 remained bound to the

mitotic spindle to a similar extent both Okadaic acid and DMSO control treated cells (Figure

S2A, S2B). We also tested if Okadaic acid treatment directly affected the extent of CKAP2

phosphorylation during mitosis. Therefore, we synchronized RPE1 cells by sequential

Palbociclib-STLC synchronization and treated the cells with Okadaic acid or DMSO control for

2 hours prior to harvesting the cells. We then immunoprecipitated CKAP2 and measured the

change in CKAP2 phosphorylation using the phosphospecific ProQ Diamond stain. Okadaic acid

did not strongly affect the extent of CKAP2 phosphorylation (Figure S2C), consistent with the

lack of effect on CKAP2 binding to the mitotic spindle. Therefore, it is not likely that PP1 or

PP2A are the main phosphatase regulators of CKAP2 phosphorylation during mitosis.

CKAP2 phosphorylation affects MT binding

Phosphorylation is known to affect the ability of many MAPs to bind to MTs because it

carries a negative charge that repels similarly charged areas on MTs. For example, the protein

Hec1 is phosphorylated on up to 10 sites to control its affinity for MTs[42–45]. Therefore, we sought

to determine if phosphorylation of CKAP2 by AurKA affects its ability to bind MTs. We treated

RPE1 cells with 200 nM MLN8237 and examined the extent of CKAP2 overlap with the mitotic

spindle. This experiment revealed a substantial increase in signal overlap between CKAP2 and

the spindle in cells treated with the Aurora A inhibitor. Although the intensity ofCKAP2 staining

did not strongly change in treated cells, the intensity of β-tubulin staining was strongly

decreased, resulting in a normalized increase in CKAP2 binding to the spindle (Figure 3A-E).

We next tested if CKAP2 phosphorylation directly affects its ability to bind to MTs using

Total Internal Reflection Microscopy (TIRF) to visualize MTs and bound proteins. We again

incubated mouse CKAP2 together with AurKA +/- ATP to generate phosphorylated or sham

phosphorylated CKAP2. Then, we mixed these proteins with guanylylmethylene diphosphonate

(GMPCPP) MT seeds and taxol-stabilized MTs to determine if phosphorylation of CKAP2

affects its ability to bind to MTs *in vitro*. Interestingly, CKAP2 that had been phosphorylated by

AurKA was profoundly defective in binding microtubules *in vitro*. At 5 nM CKAP2, binding of

unphosphorylated CKAP2 to MTs was visible, but no binding of phosphorylated CKAP2 was

detectable. However, at 50 nM concentration, some binding of phosphorylated CKAP2 to MTs

was detectable, albeit at a much lower level than unphosphorylated CKAP2 (Figure 3F, 3G).

Taken together, our data indicate that phosphorylation of CKAP2 controls its affinity for MT

binding.

TPX2 contributes to CKAP2 phosphorylation

Since we detected CKAP2 in complex with TPX2 (Figure 1C-E, S1A), we next asked if

CKAP2 contributes to TPX2 localization to the mitotic spindle. Therefore, we compared the

intensity of TPX2 on the mitotic spindle in CKAP2-knockout[8] compared to Wild-type RPE1

cells. This experiment revealed a decrease in TPX2 on the mitotic spindle in 2 out of 3 clones of

CKAP2 knockout cells, while the third clone trended downwards in TPX2 intensity (Figure 4A,

4B). All 3 CKAP2 KO clones displayed less intense tubulin staining (Figure 4A, 4C). Therefore,

it is possible that CKAP2 either directly contributes to TPX2 levels on the spindle, or that the

low tubulin levels reduce TPX2 levels on the spindle since there are fewer MTs available for

TPX2 to bind to.

Our immunoprecipitation data suggest that CKAP2 binds directly to TPX2 and indirectly

to AurKA as we detected more TPX2 peptides by mass spectrometry, and we detected relatively

more TPX2 compared to Aurora A by Western blot. Therefore, we asked if the presence of TPX2

would enhance the phosphorylation of CKAP2 by Aurora Kinase A by functioning as a binding

intermediate between AurKA and CKAP2. We repeated our kinase assay in the presence or

absence of TPX2. Interestingly, the presence of TPX2 resulted in a clear increase in the amount

of CKAP2 phosphorylation by AurKA compared to CKAP2 incubated with AurKA alone (Figure

4D). In this assay, TPX2 was also robustly phosphorylated by AurKA.

To Determine how TPX2 and CKAP2 interact, we tested the ability of CKAP2 fragments

to pull down TPX2 and AurKA. CKAP2 was divided into four fragments: the N-terminal alpha-

helical domain (1-60), disordered domain (346-549), alpha-helical bundle (346-549) and C-

terminus (550-664)[46]. Interestingly, the disordered domain interacted robustly with TPX2 and

AurKA, and the N-terminal alpha-helical and C-terminal domains contributed to a lesser extent.

The alpha-helical bundle region of mmCKAP2 did not interact at all with TPX2 and AurKA

(Figure 4E). We then performed an Alphafold2[47] binding simulation between mmCKAP2 and

TPX2. Consistent with our pull-down experiment, Alphafold2 predicted that the basic residues of

the N-terminal alpha-helix, some residues within the disordered domain and a combination of

basic and acidic residues in the C-terminal domain all interact with TPX2 (Figure 4F).

CKAP2 phosphorylation controls mitotic spindle growth

To determine if phosphorylation of CKAP2 affects mitotic spindle functions, we

transfected RPE1 CKAP2 knockout cells with wild-type, phospho-null or phospho-mimic

mutants and measured the intensity of β-tubulin in STLC arrested cells with monopolar spindles.

In constructs containing a phospho-mimic mutation, all other potential phosphorylation sites

were mutated to alanine. Interestingly, cells expressing a phospho-null CKAP2 mutant displayed

slightly higher tubulin intensity. In contrast, cells expressing individual site phosphomimic

mutants displayed slightly lower tubulin staining, suggesting that phosphorylation of CKAP2 is

important for controlling its MT growth functions (Figure 5A, 5B). We also measured the

spindle diameter of STLC-arrested cells expressing the same mutants. Although in general the

spindle diameter was not strongly changed, cells expressing a CKAP2 mutant with all five

potential sites mutated to mimic phosphorylation, as well as individual site phosphomimic

mutants for T86 or T206, showed somewhat smaller mitotic spindle diameter (Figure 5A, 5C).

We next examined TPX2 and AurKA expression in cells expressing CKAP2 phospho-

null or phospho-mimic mutants. Several mutants showed decreased presence of TPX2 on the

mitotic spindle, including the 5A, 5D, 3D and S192 and T206 individual phospho-mimic sites

(Figure S3A, S3C), suggesting that either the reduced MT stability seen in cells expressing these

mutants affects TPX2 binding or perhaps that expression of these mutants directly affects TPX2

spindle binding. We also measured the intensity of AurKA on the mitotic spindle in cells

expressing our set of mutants. Unfortunately, our AurKA measurements showed numerous

outliers in one experimental replicate, resulting in inconclusive results. However, cells

expressing the S192 and T206 individual phospho-mimic proteins trended towards lower AurKA

levels on the mitotic spindle, which would be consistent with the TPX2 measurements (Figure

S3B, S3D). Taken together, these data suggest that CKAP2 phosphorylation affects the stability

of the mitotic spindle and AurKA/TPX2 activity.

Since CKAP2 contributes to TPX2 spindle localization, we considered the possibility that

TPX2 and CKAP2 might function together to regulate MT growth. Therefore, we analysed how

TPX2 and CKAP2 contribute to microtubule dynamics individually and together in an *in vitro*

MT growth assay. In the absence of CKAP2 and TPX2 we observed MT growth and occasional

depolymerization. As shown previously, 50 nM CKAP2 potently increased microtubule growth

rates and suppressed MT catastrophe[9]. Also as shown previously, 50 nM TPX2 suppressed MT

off-rates when applied alone[34]. When applied together, the effect of CKAP2 was dominant since

it prevented catastrophe events, masking the ability of TPX2 to suppress MT off-rates. No

synergistic effects were observed between CKAP2 and TPX2, as MT dynamics were similar

between the CKAP2 only and CKAP2 plus TPX2 conditions (Figure 5D, 5E).

CKAP2 phosphorylation dynamics are required for cell viability

We next sought to determine how CKAP2 phosphorylation is important for mitotic

progression and cell viability. Although we were unable to use puromycin to select for RPE1

cells expressing wild-type or AurKA site mutants of CKAP2, we were able to do so for cancer

cell lines. Therefore, we transfected HeLa cells with either empty vector expressing mNeon

Green, CKAP2 wild-type, or CKAP2 AurKA site mutants fused to mNeon Green. We then added

puromycin selection, incubated for 12 days and finally fixed and stained the cells with crystal

violet to assess colony formation. Transfection of the empty vector supported robust colony

formation, as did transfection of CKAP2 wild-type. Interestingly, a phospho-null mutant (5A)

supported viability to a similar extent as CKAP2 wild-type. A mutant containing T86E, S192D

and T206E (3E) and the single mutants T52E and T86E also supported cell viability to a similar

extent as CKAP2 wild type. The construct containing T52E, T86E, S192D, T206E and T337E

(5E), T52E and T337E (2E) and the single site mutants S192D and T206E poorly supported

colony formation, suggesting that CKAP2 phosphorylation dynamics are important for

supporting cell viability (Figure 5F). Consistently, these two mutants single site mutants S192D

and T206E also had substantially diminished spindle microtubule density in mitotic cells (Figure

5A, 5B).

Discussion

280   Using unbiased proteomics, we have discovered numerous proteins that interact with

CKAP2 during mitosis (Figure1, supplemental table 1). Interestingly, most of these proteins are

not obviously components of the mitotic spindle or MAPs, despite the obvious spindle

localization of CKAP2 during mitosis. Thus, it is likely that CKAP2 has other roles during

mitosis, or alternatively that the interacting proteins have uncharacterized spindle roles in

mitosis. However, many of these proteins, such as the cohesins and condensins, could potentially

serve as receptors for CKAP2 to bind to chromatin in late mitosis. Importantly, AurKA/TPX2

were the only mitotic kinase or phosphatase that we detected in our screen despite the fact that

phosphatase regulation of proteins is essential to dampen kinase activity, and Cdk

phosphorylation has already been shown to regulate CKAP2, consistent with the presence of

multiple potential proline-directed phosphorylation sites in the C-terminus[16–18].

291   We envision a model for CKAP2 function where AurKA/TPX2 phosphorylates CKAP2

in response to cues from the spindle state to constrain mitotic spindle growth as part of a

negative feedback loop. CKAP2 is an extremely potent nucleator and stabilizer of MTs, and so

logically, such a protein would require a negative regulator to dampen its activity to prevent

unrestrained spindle growth. Since phosphorylation adds a negative charge to proteins, this

would cause a reduction in binding affinity of CKAP2 for MTs that would result in a reduction in

MT polymerization activity (Figure 3, Figure 6). In this scenario, AurKA/TPX2 kinase activity

would cause spindle growth, which would in turn allow more binding of AurKA/TPX2 to the

spindle, resulting in more kinase activity and increased CKAP2 phosphorylation, which will then

start to dampen spindle growth.

Although the structure of CKAP2 bound to MTs has not yet been solved, the structure of

TPX2 bound to MTs has been determined. Unlike CKAP2, which binds to MTs both during

interphase and mitosis, TPX2 only binds to MTs during mitosis[28]. TPX2 uses two linked regions

to bind across both longitudinal and lateral MT interfaces[29]. It will be interesting to compare and

contrast the binding mode of CKAP2 to MTs to determine if it competes with TPX2, binds to an

independent site or synergizes with TPX2 for MT binding. However, in terms of activity, CKAP2

appears to directly facilitate MT growth, whereas TPX2 largely reduces MT disassembly rates by

slowing the MT-off -rate. Thus, CKAP2 and TPX2 perform different roles in promoting MT

growth and stability. Importantly, the n-terminus of CKAP2 is largely disordered, and so it might

be able to bind MTs in multiple conformations, which could be impacted by phosphorylation.

The disorder of CKAP2 also likely contributes to the ability of AurKA/TPX2 to phosphorylate

CKAP2. We found that it is largely the disordered domain of CKAP2 which binds to TPX2

(Figure 4E). We speculate that this would position CKAP2 in such a way that the N-terminal

disordered domain of CKAP2 can be optimally phosphorylated by AurKA, as we found

substantially enhanced phosphorylation of CKAP2 in the presence of TPX2 and AurKA

compared to AurKA alone (Figure 4D).

We identified up to 5 AurKA phosphorylation sites in the N-terminus of mouse CKAP2

and 4 in human CKAP2. Of these, the most ideal consensus motif based on the amino acid

sequence is in T39 (mouse T52). However, this site is located within an alpha-helix and might

not be accessible to the kinase. This site has previously been identified in a screen for AurKB

substrates, but not directly in cell lysates[48]. This experiment was performed by mixing lysate

with purified kinase, which might allow the kinase to encounter substrates that it might not

normally encounter due to the lack of spatial organization present in a lysate. Although it is not

inconceivable that soluble CKAP2 might encounter soluble AurKB, considering that the bulk of

these proteins are localized to different structures (Figure S1B, S1C), it is unlikely that AurKB

can substantially phosphorylate CKAP2. In contrast, since AurKA/TPX2 binds to CKAP2 as

detected by MS and Western blot and largely co-localizes throughout mitosis; it is much more

likely that CKAP2 is regulated by AurKA. Human CKAP2 T76 phosphorylation (mouse T86)

has not yet been detected directly, despite forming a good AurK consensus site in an unstructured

region. Human CKAP2 S190 phosphorylation has been detected. However, it is unclear which

kinase might phosphorylate it. Human CKAP2 S209 (mouse T206) has also been detected and

forms a good AurK consensus site. Consistently, our cell biology and *in vitro* experiments point

to the T206 phosphorylation site being the most important one for regulating CKAP2 functions.

Cells expressing this mutant have substantially less stable MTs compared to cells expressing

wild-type CKAP2 and mutation of this site does not efficiently support cell viability (Figure 5A,

5B, 5F). Future studies will aim to determine exactly how phosphorylation of this site impairs

the ability of CKAP2 to bind MTs and stimulate their growth.

Overall, our study provides a valuable resource for further investigation of CKAP2 and

the cellular pathways in which it participates. Future studies will be necessary to validate the

binding of the other proteins we detected and to determine their contributions to the functions of

CKAP2 in mitosis and interphase.

Resource availability

The cell biology datasets generated during and/or analysed during the current study are available

from the corresponding author on reasonable request. The original datasets for CKAP2

interactome has been deposited to the ProteomeXchange Consortium via the PRIDE[49] partner

repository with the dataset identifiers to be determined following acceptance of the article for

publication.

Acknowledgements

We thank Arnold Hayer and Jose Teodoro for providing reagents. We thank Denis Faubert and

Josée Champagne for assistance with the proteomics analysis. We also thank members of the

Bechstedt laboratory and Gary Brouhard for helpful suggestions and feedback. The laboratory of

SB is supported by grants from the Canadian Institutes of Health Research (CIHR PJT-189995),

the Natural Sciences and Engineering Research Council of Canada (NSERC; RGPIN-2024-

05603), and in part by a Fonds de Recherche du Québec (Health Sector) Research Centres Grant

#288558.

Author Contributions

Conceptualization: SB and TJK. Investigation: TJK, TL. Formal analysis: TJK, TL.

Methodology: TJK, SB. Funding acquisition: SB. Supervision: SB. Writing- original draft: TJK,

SB. Writing- review & editing: TJK, SB. Project administration: SB.

Declaration of interests

The authors declare that they have no conflicts of interest

Materials and Methods

<u>Cell Culture</u>

RPE1 were a gift from Arnold Hayer (McGill University). RPE1 CKAP2 GFP-CKAP2 and

CKAP2 knockout cells were described previously[8]. All cells were grown at $37^{\circ}C$ in a humidified

environment with 5% CO_2. HeLa cells and U20S cells were purchased from the ATCC and

grown in Dulbecco's modified Eagle medium (Life Technologies #11995073) containing 10%

Fetal bovine serum (Wisent #098150), 50 U/mL penicillin and 50 μg/mL streptomycin (Life

Technologies #15140122). RPE1 cells were grown in DMEM-F12 (Life Technologies

#11320082) with the same supplements. Cells were verified to be free of mycoplasma by

frequent staining of plated cells with DAPI. No contamination was observed.

<u>Inhibitors and reagents</u>

STLC (Fisher Scientific #T25225G) was used at 25 μM. Okadaic acid (Cell Signaling

Technology #5934) was used at 100 nM. MLN8237 (Selleck #1028486-01-2) was used at 25-500

nM. Palbociclib (MilliporeSigma #PZ0383-5MG) was used at 1 μM. Taxol (Fisher Scientific

#AAJ62734MC) was used at varying concentrations. Puromycin (BioBasic # PJ593) was used at

1 μg/ml.

<u>Plasmids and cloning</u>

Mouse CKAP2 and codon optimized mouse TPX2 were cloned into pHAT His-USER-GFP-Strep

via USER cloning[50]. It was also cloned into pTWIST Puro EF1alpha-USER-mNeonGreen via

USER cloning. Aurora A site mutants of mouse CKAP2 were created by custom DNA synthesis

by Twist Bio and then cloned into plasmids via USER cloning[50]. mmCKAP2 fragments were

described previously[46]. The DNA constructs were then verified by Sanger sequencing.

<u>Transfections</u>

RPE1 or HeLa cells were transfected using Xfect™ transfection reagent (Takara Bio #631318)

according to the manufacturer's directions. Briefly, media on the cells was changed for 500 μl

for a 12-well plate. 4 μg of DNA was mixed with 100 μl of buffer. 1.2 μl of Xfect reagent was

then added, and the solution mixed by vortex. After 5 minutes incubation, 50 µl of transfection

mix was added to cells overnight. The media was then changed 24 hours later.

<u>Immunofluorescence</u>

Following treatments, cells were fixed in 4% paraformaldehyde in PBS at room temperature for

20 minutes. The cells were washed twice with PBS and then permeabilized in PHEM buffer (60

mM PIPES, 25 mM HEPES, pH 6.9, 10 mM EGTA, and 4 mM $MgSO_4$, 1% Triton X-100, 10

mM glycerol 2-phosphate) for 10 minutes. The cells were then washed twice with PBS for before

blocking with ADB (10% serum, 0.1% Triton X-100 in PBS) for 15 minutes. Primary antibodies

were then diluted in ADB and used at the indicated dilution. Secondary antibodies were diluted

in ADB. After incubation with antibodies, the cells were washed three times with PBS 0.1%

Triton X-100. Finally, the cells were counterstained with DAPI and mounted on slides using

FluorSave^{TM} antifade reagent (Millipore #345789-20 ml).

<u>Immunoprecipitation</u>

For protein-protein interactions, 500,000 RPE1 cells were plated on 10 cm dishes (5 per

condition). 24 hours later, palbociclib was added. 18 hours following palbociclib addition, it was

washed out with 2 4 ml washes with PBS. STLC was then added 5 hours later. 20 hours post-

release, the cells were harvested by scraping, washed once with cold PBS, and lysed in the

following buffer: 0.5% NP40, 100 mM NaCl, 50 mM tris pH 7.4, 10 mM glycerol 2-phosphate.

Cells were lysed on ice for 20 minutes before centrifuging at 15,000 x g for 20 minutes at 4°C.

The supernatant was transferred to a fresh tube, and input sample set aside. 10 µl of packed

volume protein G beads were added to each sample together with 5 µg of antibody for 2 hours on

a rotating platform at 4°C. The beads were then washed 5 times in lysis buffer and dried using a

30 g needle attached to a vacuum line. 30-100 µl of 1X sample buffer were added to the beads

before elution by boiling and loading on gel.

For phosphorylation analysis, cells were prepared and synchronized as above. Following

harvesting, the cells were lysed in 50 mM Tris pH 7.5 and 1% SDS. The lysates were then boiled

for 5 minutes. SDS was then inactivated by 10-fold dilution with 1.5% Triton X-100 in TBS.

Debris was then pelleted via centrifugation at 15,000 x g for 20 minutes at 4°C. CKAP2 was then

immunoprecipitated using anti-CKAP2 and protein G agarose as above.

<u>Mass Spectrometry</u>

CKAP2 was purified from palbociclib-STLC synchronized cells by immunoprecipitation as

described above. The entire reaction was then eluted in sample buffer and separated on a 10%

SDS-PAGE gel. The gel was then stained with Coomassie blue, and each lane was cut into 12

slices. Gel bands were excised under a clean bench and cut into 1 mm³ pieces. For all subsequent

steps, reagent volumes were adjusted according to the volume of the gel pieces. Gel pieces were

first washed with water for 5 minutes, then destained twice using a destaining buffer composed

of 50 mM ammonium bicarbonate and 50 µL of acetonitrile, each for 10 minutes. Following this,

the gel pieces were dehydrated with acetonitrile. Proteins were reduced by incubating the gel

pieces in reduction buffer (10 mM DTT, 100 mM ammonium bicarbonate) for 30 minutes at

40 °C, and subsequently alkylated with alkylation buffer (55 mM iodoacetamide, 100 mM

ammonium bicarbonate) for 20 minutes at 40 °C in the dark. The gel pieces were then

dehydrated and washed at 40 °C with acetonitrile for 5 minutes before removing all reagents.

They were dried for 5 minutes at 40 °C and then rehydrated at 4 °C for 40 minutes with trypsin

solution (6 ng/µL sequencing-grade trypsin from Promega in 25 mM ammonium bicarbonate).

The low concentration of trypsin was used to minimize signal suppression and background from

autolysis products during LC-MS/MS analysis. Protein digestion was carried out at 58 °C for 1

hour and quenched with 15 µL of 1% formic acid / 2% acetonitrile. The supernatant was

transferred to a 96-well plate. Peptide extraction was performed in two 30-minute steps at room

temperature using an extraction buffer (1% formic acid / 50% acetonitrile). All peptide extracts

were completely dried using a vacuum centrifuge. The LC column used was a PicoFrit fused

silica capillary column (15 cm × 75 µm i.d.; New Objective, Woburn, MA), self-packed with

C18 reversed-phase material (Jupiter, 5 µm particles, 300 Å pore size; Phenomenex, Torrance,

CA) using a high-pressure packing cell. This column was installed on the Easy-nLC II system

(Proxeon Biosystems, Odense, Denmark) and coupled to a Q Exactive mass spectrometer

(ThermoFisher Scientific, Bremen, Germany) equipped with a Proxeon nanoelectrospray Flex

ion source. The chromatography buffers were 0.2% formic acid in water (buffer A) and 100%

acetonitrile with 0.2% formic acid (buffer B). Peptides were loaded on-column and eluted using

a three-step gradient at a flow rate of 250 nL/min. Solvent B increased from 2% to 35% over 15

minutes, then from 35% to 50% over 7 minutes, and finally from 50% to 90% over 5 minutes.

LC-MS/MS data were acquired using a data-dependent Top12 method combined with a dynamic

exclusion window of 7 seconds. The mass resolution for full MS scans was set to 70,000 (at m/z

400), and lock masses were used to improve mass accuracy. The mass range for MS scanning

was from 360 to 2000 m/z, with a target value of 1×10^6 and a maximum ion fill time of 100 ms.

Data-dependent MS2 scan events were acquired at a resolution of 17,500, with a maximum ion

fill time of 50 ms, a target value of 1×10^5, an intensity threshold of 1.0×10^4, and an underfill

ratio of 0.5%. The normalized collision energy was set to 27%, and the capillary temperature was

maintained at 250 ºC. Nanospray and S-lens voltages were set to 1.3–1.7 kV and 50 V,

respectively. Protein database searches were performed using Mascot 2.6 (Matrix Science)

against the UniProt human database. The mass tolerance was set to 10 ppm for precursor ions

and 0.5 Da for fragment ions. Trypsin was specified as the enzyme, allowing up to two missed

cleavages. Carbamidomethylation of cysteine was set as a fixed modification, and methionine

oxidation was included as a variable modification.

<u>SDS-PAGE and Western blot</u>

Gels and blots were performed as described previously [51]. Prior to loading samples on gel, 4X

Laemmli buffer (200 mM Tris pH 6.8, 4% SDS, 40% glycerol, 4% 2-mercaptoethanol, 0.12

mg/ml bromophenol blue) was added to a final dilution of 1X, and the samples were boiled for 5

minutes. Proteins were separated on SDS-PAGE gel using stacking gel (4% 29:1 acrylamide:

Bis-acrylamide, 125mM Tris pH 6.8, 0.1% SDS, 0.1% ammonium persulfate, 0.1% TEMED)

and (8-15% 29:1 acrylamide: Bis-acrylamide, 400 mM Tris pH 8.8, 0.1% SDS, 0.1% ammonium

persulfate, 0.1% TEMED) at 120V until the bromophenol blue has run off or longer, as needed.

Transfer onto nitrocellulose membrane was performed for at least 24 hours at 30V under wet

conditions in 1X transfer buffer (14.4 g/L glycine, 3.0 g/L Tris, 20% methanol). Conditions for

western blots includes the use of 5% nonfat dry milk in TBS-T 0.5% (50 mM Tris [pH 7.2], 150

mM NaCl, 0.5% Tween 20) for blocking and TBS-T 0.5% for washing. 4 washes for 10 minutes

each were performed after primary and secondary antibody incubation periods. The bands were

visualized by enhanced chemiluminescence using Clarity (Bio-Rad #1705060) using a Bio-Rad

ChemiDoc™ Touch Imaging System.

<u>*In vitro* analysis of CKAP2</u>

Mouse CKAP2 protein variants were produced in the E. Coli strain BL21. The bacteria were

suspended in in 25 ml buffer A (50 mM NaPO$_4$, 10 mM Imidazole, 300 mM NaCl, 10%

Glycerol, pH 8) and processed in an emulsiflex high-pressure homogenizer (Avestin). The lysate

was then ultracentrifuged at 45Ti rotor at 25000 RPM for 45 minutes. 200 μl of packed

glutathione agarose was then added for 2 hours on a rotating platform. 1 ml Ni-NTA agarose was

then added to the lysate for 2 hours on a rotating platform. The agarose was then washed, and

purified proteins eluted with buffer A containing 300mM imidazole and then dialyzed against

0.4X TBS. Glycerol was then added to 10% and the proteins frozen until further use. Kinase

reactions were performed by mixing CKAP2, GFP-Aurora A-6XHis, 100 μM ATP in kinase

buffer 50 mM Tris pH 7.2, 25 mM NaCl, 10mM $MgCl_2$ 1mM DTT, 0.01% Triton X-100 for 1

hour at 37°C. The reactions were stopped by addition of 4X SDS sample buffer, and half the

reaction loaded on SDS-PAGE gel. The gels were then stained with ProQ Diamond

phosphoprotein stain (Thermo Scientific) according to the manufacturer's instructions and

subsequently with Coomassie blue.

In vitro MT assays were performed as described previously[9]. CKAP2 fragments were produced

in BL21(DE3) *E. coli* grown in LB media to OD_{600} and then induced with 0.5 mM Isopropyl-D-

1-thiogalactopyranoside (IPTG) at 20°C overnight. The bacteria were then harvested and lysed in

Buffer A (50 mM NaH2PO4, 300 mM NaCl, 0 mM imidazole, pH 7.8) using an EmulsiFlex-C5

(Avestin Inc, Canada). Debris was then pelleted by ultracentrifugation at 30,000 x g for 1 hour at

4°C. The lysate was then loaded onto a column containing His60 Ni SuperFlow Resin (Takara

Bio, USA). Purified protein was eluted with Buffer B (50 mM NaH2PO4, 300 mM NaCl, 200

mM imidazole, pH 7.8). Eluted proteins were then loaded onto a column containing Strep-Tactin

Sepharose resin (IBA Lifesciences, Germany), washed with 5 column volumes of Buffer W (100

mM Tris-HCl, 150 mM NaCl, 1 mM EDTA, pH 7.8) and finally eluted with Buffer W

supplemented with 2.5 mM Desthiobiotin. Protein concentration was determined by measuring

the absorbance at 280 or 488 nm using a DS- 11 FX spectrophotometer (DeNovix Inc, USA).

SDS-PAGE was used to determine the purity of the proteins. Tubulin was purified from bovine

brains as previously described[52] with the difference of having used Fractogel EMD SO3- (M)

resin (Millipore-Sigma) instead of phosphocellulose. Tubulin was labelled using CF640R-NHS-

Ester (Biotium, USA) and tetramethylrhodamine (TAMRA, Invitrogen Inc, Canada), as

previously described[53]. Before use, an additional cycle of MT polymerization was performed.

<u>Microscope Slide Channel Preparation</u>

Glass coverslips (22x22 and 18x18 mm) were washed with acetone and then sonicated in 50%

methanol with 0.5 M KOH and finally rinsed with water. The coverslips were then dried and

exposed to air plasma (Plasma Etch) for 3 minutes before silanizing by soaking them in 0.1%

dichlorodimethylsilane in *n*-heptane before sonicating them in *n*-heptane and finally in ethanol.

Flow channels with a volume of 7 μl were created by using two pieces of silanized coverslips

held by double-sided tape in custom coverslip holders. The channels were incubated with anti-

TAMRA antibodies for 5 minutes, blocked with 5% Pluronic F-127 for 10 minutes and washed

with BRB80 before incubation with MTs. The channels were used at 35°C using a heated

microscope objective and incubation chamber.

<u>Binding to Paclitaxel-stabilized MTs</u>

GDP-MTs stabilized by paclitaxel were prepared by polymerizing a 1:15 ratio of TAMRA-

labelled to unlabeled tubulin in the presence of 4 mM $MgCl_2$, 1mM GTP, 5% DMSO in BRB80

buffer. The mixture was then incubated for 30 minutes at 37°C. The MTs were then diluted using

BRB80 plus 10 μM paclitaxel, pelleted at 100,000 RPM at 30 psi for 7 minutes using an Airfuge

(Beckman-Coulter) and finally resuspended in BRB80 plus μM paclitaxel.

To assess binding of CKAP2 to MTs, A combination of GMPCPP double-cycled MTs and

paclitaxel-stabilized MTs were mixed and flowed into microscope slide channels. Since the

paclitaxel-stabilized MTs contain less TAMRA label, they possess a dimmer signal compared to

GMPCPP MTs.

Dynamic Microtubule Assay

GMPCPP MT seeds were created by polymerizing a 1:4 ratio of TAMRA-labelled to unlabeled

tubulin in the presence of 1 mM guanosine-5'- [(α, β)-methyleno] triphosphate (GMPCPP, Jena

Biosciences, Germany), a non-hydrolyzable GTP analogue during two cycles. On the day of an

experiment, unlabeled and CF642R-labeled tubulin were mixed at a 1:17 ratio, aliquoted and

flash-frozen in liquid nitrogen. To start an experiment, GMPCPP MT seeds were flowed into a

channel and then washed with BRB80 buffer. MT growth was initiated by introducing tubulin (8

μM) in BRB80 buffer containing 1 mM GTP, 40 mM D-glucose, 64 nM catalase, 250 nM

glucose oxidase, 10 mM DTT, 0.1 mg/ml BSA together with CKAP2. Images were then taken at

suitable time points to track MT growth and catastrophe.

Colony formation assays

300, 000 HeLa cells per well were plated on 6-well plates. 24 hours following plating, the cells

were transfected overnight as described above. 18 hours following transfection, the media was

changed. 42 hours following transfection, the media was changed to include puromycin selection

(1 μg/ml) for 3 days. The media was then changed for media containing 0.2 μg/ml puromycin,

and the cells were allowed to proliferate for another 12 days before fixation and staining with

crystal violet.

Microscopy

Images were acquired with either a Zeiss Axio Observer 7 fitted with a TIRF

equipped with a Prime 95B CMOS camera (Photometrics) with a pixel size of 107 nm controlled

by Zeiss Zen 2.5 Blue Edition software version or a Nikon Eclipse Ti microscope spinning disc

confocal (Cicero; Crest Optics) microscope equipped with AURA III Light Engine (Lumencor®)

and Hamamatsu ORCA-fusion C14440 camera with 110 nm pixel size controlled by Nikon-

Elements AR 6.10.01 software. Images were acquired in 0.15-0.4 μm sections using a plan apo

1.4 numerical aperture 100X or 60X oil-immersion objective using 1X1 binning. All image

analysis, adjustments, measurements and cropping were performed using Fiji software [54].

Widefield Image deconvolution was performed using Zen Blue Edition 2.5 software. All protein

intensity analysis was performed on the raw images. Line scans were performed on the

deconvoluted images in Fiji and the data was then exported to Graphpad Prism version 8.4.3. or

version 10.2.1

<u>Antibodies</u>

The following antibodies were used for immunofluorescence (IF) and/or immunoblotting (IB):

mouse anti-Aurora A (Cell Signaling Technology 1F8; IF at1:1000, blot at 1:500), Rabbit anti-

Aurora A pT288 (Cell Signaling Technology C39D8; IF at 1:500), mouse anti-TPX2 (Cell

Signaling Technology D9Y1V; IF at 1:1000, blot at 1:500), rabbit anti-Aurora B (Cell Signaling

Technology 3094T; IF at 1:50), mouse anti-β−Tubulin (Sigma clone 2.1 T4026; IF at 1:200),

rabbit anti-CKAP2 (Proteintech 25486-1-AP; blot at 1:500), Rabbit anti-CKAP2 (AbClonal

A9706; IF at 1:200), Mouse anti-TAMRA (Invitrogen 5G5; label at 1:100). Secondary antibodies

used were highly cross-adsorbed Alexa Fluor® 488, 594 and 647 raised in donkey or goat

(Invitrogen; IF at 1:1000), horseradish peroxidase (Bio-Rad; IB at 1:10000-10,0000), Anti-IgG

(H+L) Goat Polyclonal Antibody (Horseradish Peroxidase), Peroxidase AffiniPure Goat Anti-

Rabbit IgG (H+L) (IB at 1:5000), Goat Polyclonal Antibody (Horseradish Peroxidase),

Peroxidase AffiniPure Goat Anti-Mouse IgG (H+L) (IB at 1:5000).

<u>Statistical analysis of cell biology and MT data</u>

All statistical tests were performed using Graphpad Prism version 8.4.2. Except where indicated,

all experiments were performed as 2-3 independent biological repeats. In most IF experiments,

25 cells for each condition were measured. Experiments with other numbers are noted in the

figure legends.

Figure legends

Figure 1: Proteomic screening identifies AurKA/TPX2 as CKAP2 interacting proteins

A) Cartoon diagram showing the experimental scheme for determining CKAP2 binding

partners. RPE1 cells were first synchronized in mitosis by Palbociclib-STLC block. The

cells were then harvested and lysed. Endogenous CKAP2 was then immunoprecipitated

from the mitotic lysate together with interacting proteins. The purified proteins were then

separated via SDS-PAGE. The gel lanes were then cut into 12 slices and analysed by

mass spectrometry to identify CKAP2 interaction partners. A single experiment was

performed.

B) Proteins from the phospho-proteomic screen from (A) that that met criteria of at least 4

peptides detected and 4:1 signal to noise ratio were analyzed for gene ontology. The data

were then transferred to Prism software and plotted.

C) STRING analysis of the proteins from (A) meeting detection criteria showing protein

interaction networks.

D) Selected interacting proteins detected in the control immunoprecipitation and CKAP2

immunoprecipitations from (A) showing the number of total detected peptides.

E) RPE1 cells were prepared as in panel (A), except that purified proteins were analyzed by

Western blot. CKAP2 was immunoprecipitated from Palbociclib-STLC mitotic lysates

and the purified proteins were then separated by SDS-PAGE, transferred to nitrocellulose

membrane and blotted as indicated. The panels were adjusted for brightness and contrast.

An experiment representative of two independent biological repeats is shown.

Figure 2: CKAP2 co-localizes with and is phosphorylated by AurKA/TPX2

A) Immunofluorescence images from RPE1 eGFP-CKAP2 cells in various stages of mitosis

stained for AurKA, AurKA pT288 and DAPI. The CKAP2-eGFP, AurKA and AurKA

pT288 channels were adjusted evenly for brightness and contrast. The DAPI channel of

each condition was adjusted independently. Representative images from 2 independent

biological experiments are shown. Scale bars, 10 μm and 1 μm (inset).

B) Immunofluorescence images from RPE1 eGFP-CKAP2 cells in various stages of mitosis

stained for TPX2 and DAPI. The CKAP2-eGFP and TPX2 channels were adjusted evenly

for brightness and contrast. The DAPI channel of each condition was adjusted

independently. Representative images from 2 independent biological experiments are

shown. Scale bars, 10 μm and 1 μm (inset).

C) Amino acid sequence alignment of CKAP2 orthologs showing predicted AurKA

phosphorylation consensus motifs. Residues highlighted in green indicate potential AurK

phospho-acceptor residues.

D) RPE1 cells were synchronized by Palbociclib-STLC block. The cells were then harvested

and lysed by boiling in SDS-containing buffer. The SDS was then inactivated by 10-fold

dilution with 1.5% Triton X-100 buffer. CKAP2 was then purified by

immunoprecipitation. The protein-G agarose beads were either sham treated or treated

with Lambda protein phosphatase for 30 minutes at 30°C. The beads were then boiled in

sample buffer to elute purified proteins. Purified proteins were then separated by SDS-

PAGE, which was then stained with with ProQ Diamond phosphoprotein stain and

imaged. Finally, the gel was stained with Coomassie blue and imaged again. An

experiment representative of two independent biological repeats is shown.

E) Cells were prepared as in (D), except that one set of cells was treated concurrently with

STLC and DMSO control and one set with MLN 8237 AurKA inhibitor. CKAP2 was

then purified and analysed as in (D). An experiment representative of two independent

biological repeats is shown.

F) Purified GFP-Aurora A-6XHis and His-mmCKAP2-GFP-Strep were mixed with reaction

components as indicated and incubated at 37°C for 1 hr. The reaction was then separated

by SDS-PAGE gel. The gel was then stained with ProQ Diamond phosphoprotein stain,

imaged and subsequently stained with Coomassie blue and re-imaged. An experiment

representative of two independent biological repeats is shown. High-brightness/contrast

and low-brightness/contrast images are shown for presentation.

Figure 3: Phosphorylation the mitotic spindle localization of CKAP2

A) Immunofluorescence images from asynchronous RPE1 cells in late prometaphase were

treated with either DMSO or MLN8237 for 3 hours and then fixed and stained for

CKAP2, β-tubulin and DAPI. The CKAP2 and β-tubulin channels were adjusted evenly

for brightness and contrast. The DAPI channel of each condition was adjusted

independently. Representative images from 2 independent biological experiments are

shown. Scale bars, 10 μm and 1 μm (inset).

B) Line scans across the indicated paths from the images in panel (A).

C) Quantification of the CKAP2-β-tubulin ratio from panel (A). The Z-section with the

brightest signal was chosen, and an outline traced around the mitotic spindle. The levels

of CKAP2 and β-tubulin were then measured. Background signal was taken as the

minimum measured value in the mitotic spindle area and subtracted from the measured

signal. At least 50 cells per condition for each of 3 independent biological repeats were

measured. Error bars indicate the mean +/- SEM. Statistical significance was calculated

between the indicated conditions using Student's T- test.

D) Quantification of the CKAP2 levels from panel (A) as for panel (C).

E) Quantification of the β-tubulin levels from panel (A) as for panel (C).

F) TIRF microscopy images of CKAP2 together with guanosine-5'[(α,β)-methleno]

triphosphate GMPCPP-seeds and paclitaxel-stabilized MTs. CKAP2 was incubated *in*

vitro with AurKA in the presence or absence of ATP. Channels in between silanized-glass

coverslips were prepared by flowing-in anti-TAMRA antibodies. After washing, TAMRA

labelled GMPCPP-seeds and paclitaxel-stabilized MTs were flowed-into the chamber.

Then, CKAP2 prepared as indicated was also flowed into the chamber and images were

taken. Representative images from 2 independent biological experiments are shown.

Scale bar, 5 μm.

G) Quantification of the results presented in (F). The fluorescence intensity of CKAP2 on

each of 20 GMPCPP seeds or 20 paclitaxel stabilized microtubules was quantified.

Figure 4: TPX2 and CKAP2 reciprocally regulate their microtubule binding activities

A) Immunofluorescence images of asynchronous RPE1 wild-type and CKAP2 KO clones 8,

9 and 12. The cells were plated on glass coverslips, allowed to proliferate and then fixed

and stained for TPX2, β–tubulin and DAPI. The CKAP2 and β-tubulin channels were

adjusted evenly for brightness and contrast. The DAPI channel of each condition was

adjusted independently. Representative images from 2 independent biological

experiments are shown. Scale bars, 10 μm.

B) Quantification of the TPX2 levels from panel (A). The Z-section with the brightest signal

was chosen, and an outline traced around the mitotic spindle. The levels of TPX2 were

then measured. Background signal was taken as the minimum measured value in the

mitotic spindle area and subtracted from the measured signal. At least 25 cells per

condition for each of 3 independent biological repeats were measured. Error bars indicate

the mean +/- SEM. Statistical significance was calculated between the indicated

conditions using Dunnett's multiple comparison test.

C) As for panel (B) but the β–tubulin levels quantified

D) Purified GFP-Aurora A-6XHis, His-mmTPX2-GFP-Strep and His-mmCKAP2-GFP-

Strep were mixed with reaction components as indicated and incubated at 37°C for 1 hr.

The reaction was then separated by SDS-PAGE gel. The gel was then stained with ProQ

Diamond phosphoprotein stain, imaged and subsequently stained with Coomassie blue

and re-imaged. The images were adjusted for brightness and contrast for presentation. An

experiment representative of two independent biological repeats is shown.

E) His-mmCKAP2-GFP-Strep fragments were purified by one-step Strep affinity gel

purification. Concurrently, HeLa cells were synchronized by thymidine-nocodazole

block. The cells were then harvested and lysed in % Triton X-100 in TBS. The lysates

were then added to the Strep affinity gel bound to mmCKAP2 fragments and incubated.

The purified proteins were then separated by SDS-PAGE and transferred to nitrocellulose

membrane and finally stained with Ponceau and blotted as indicated. The images were

adjusted for brightness and contrast for presentation. An experiment representative of two

independent biological repeats is shown.

F) Alphafold2 model of mmTPX2 (gold) bound to mmCKAP2 (cyan) showing interacting

residues labelled in red.

Figure 5: CKAP2 phosphorylation dynamics are required for spindle stability and cell

proliferation

A) Immunofluorescence images of cells expressing CKAP2 Aurora site mutants fused to

mNeon Green. RPE1 CKAP2 KO Clone 8 cells were plated on glass coverslips. They

were then transfected with the indicated constructs. The cells were treated with STLC,

and after 24 hours of treatment the cells were fixed and stained for β–tubulin. The

CKAP2 and β-tubulin channels were adjusted evenly for brightness and contrast. The

DAPI channel of each condition was adjusted independently. Representative images from

3 independent biological experiments are shown. Scale bars, 10 μm and 1 μm (inset).

B) Quantification of the β-tubulin levels from the cells in panel (A). The Z-section with the

brightest signal was chosen, and an outline traced around the mitotic spindle. The levels

of β-tubulin were then measured. Background signal was taken as the minimum

measured value in the mitotic spindle area and subtracted from the measured signal. At

least 25 cells per condition for each of 3 independent biological repeats were measured.

Error bars indicate the mean +/- SEM. Statistical significance was calculated between the

indicated conditions using Dunnett's multiple comparison test.

C) Quantification of the spindle diameter of the cells from panel (A). The diameter of the β-

tubulin signal was measured for each of the cells quantified in panel (B). Error bars

indicate the mean +/- SEM. Statistical significance was calculated between the indicated

conditions using Dunnett's multiple comparison test.

D) Kymographs of TIRF microscopy images of His-mmCKAP2-GFP-Strep and/or His-

mmTPX2-GFP-Strep together with GMPCPP microtubule seeded microtubules.

GMPCPP seeds were flowed into a slide chamber and washed. Free tubulin subunits were

then flowed into the chamber along with His-mmCKAP2-eGFP-Strep, His-mmTPX2-

eGFP-Strep or both proteins together. Images were then taken at the indicated time

points. Representative images of one of two independent biological repeats are shown.

E) Quantification of the data from (D). The growth rate from at 51 (control) 35 (CKAP2) 32

(TPX2) 34 (CKAP2+TPX2) microtubules per condition from two independent

experiments were measured and plotted in Prism.

F) HeLa cells were plated and then transfected with various mmCKAP2 constructs fused to

mNeon Green as indicated. Two days following transfection, 1 µg/ml puromycin was

added for 3 days and then changed for 0.2 µg/ml puromycin. 12 days later, the cells were

fixed and stained with crystal violet. A single experiment was performed. Scale bar, 4 cm.

Figure 6: Mitotic spindle growth is controlled by an AurKA/TPX2/CKAP2 complex

Cartoon diagram showing how CKAP2 is affected by phosphorylation via an

AurKA/TPX2 complex. By itself, CKAP2 robustly mediates microtubule growth and

stabilization. During mitosis, CKAP2 is bound by TPX2, which results in

phosphorylation via the TPX2 binding partner AurKA. Phosphorylation of CKAP2

results in lower microtubule binding affinity and polymerization activity.

Figure S1: AurKB does not co-localize with CKAP2 during mitosis; TPX2 is a CKAP2

binding partner

A) TPX2 was immunoprecipitated together with interacting proteins from RPE1 cells

synchronized via Palbociclib-STLC block. The purified proteins were then separated by

SDS-PAGE, transferred to nitrocellulose membrane and blotted as indicated. The panels

were adjusted for brightness and contrast. An experiment representative of two

independent biological repeats is shown.

B) Immunofluorescence images from RPE1 eGFP-CKAP2 cells in various stages of mitosis

stained for CKAP2, AurKB, and DAPI. The CKAP2 and AurKB channels were adjusted

evenly for brightness and contrast. The DAPI channel of each condition was adjusted

independently. Representative images from 2 independent biological experiments are

shown. Scale bars, 10 μm and 1 μm (inset).

C) Line scans across the indicated paths from the images in panel (B).

Figure S2: PP2A phosphatase activity does not affect CKAP2 mitotic spindle localization

A) Immunofluorescence images from RPE1 cells treated with Okadaic acid. RPE1 cells were

plated on glass coverslips and allowed to proliferate. The cells were then treated with

DMSO control or 100 nM Okadaic acid for 2 hours before fixation and staining for

CKAP2 and β-tubulin. The CKAP2 and β-tubulin channels were adjusted evenly for

brightness and contrast. The DAPI channel of each condition was adjusted independently.

Representative images from 2 independent biological experiments are shown. Scale bars,

10 μm and 1 μm (inset).

B) Line scans across the indicated paths from the images in panel (A).

C) RPE1 cells were synchronized by Palbociclib-STLC block. The cells were then treated

with 100 nM Okadaic acid for 2 hours before harvesting. The cells were then collected

and lysed by boiling in SDS-containing buffer. The SDS was then inactivated by 10-fold

dilution with 1.5% Triton X-100 buffer. CKAP2 was then purified by

immunoprecipitation. The protein G agarose with bound proteins were then boiled in

sample buffer to elute purified proteins. Purified proteins were then separated by SDS-

PAGE, which was then stained with with ProQ Diamond phosphoprotein stain and

imaged. Finally, the gel was stained with Coomassie blue and imaged again. Two

independent experiments were performed with similar results.

Figure S3: CKAP2 phosphorylation reduces TPX2/AurKA mitotic spindle localization

A) Immunofluorescence images of cells expressing CKAP2 Aurora site mutants fused

to mNeon Green. RPE1 CKAP2 KO Clone 8 cells were plated on glass coverslips.

They were then transfected with the indicated constructs. The cells were then treated

with STLC, and after 24 hours of treatment the cells were fixed and stained for

TPX2. The CKAP2 and TPX2 channels were adjusted evenly for brightness and

contrast. The DAPI channel of each condition was adjusted independently.

Representative images from 3 independent biological experiments are shown. Scale

bars, 10 μm and 1 μm (inset).

B) As for panel (A) but the cells were stained for AurKA

C) Quantification of the TPX2 levels from the cells in panel (A). The Z-section with the

brightest signal was chosen, and an outline traced around the mitotic spindle. The

levels of TPX2 were then measured. Background signal was taken as the minimum

measured value in the mitotic spindle area and subtracted from the measured signal.

At least 25 cells per condition for each of 3 independent biological repeats were

measured. Error bars indicate the mean +/- SEM. Statistical significance was

calculated between the indicated conditions using Dunnett's multiple comparison

test.

D) As for panel (C), but the cells from panel (B) were measured and the levels of

AurKA quantified.

References
1. Kapoor, T. Metaphase Spindle Assembly. *Biology* **6**, 8 (2017).

2. Compton, D. A. SPINDLE ASSEMBLY IN ANIMAL CELLS. (2000).

3. Heald, R. & Khodjakov, A. Thirty years of search and capture: The complex simplicity of

mitotic spindle assembly. *J. Cell Biol.* **211**, 1103–1111 (2015).

4. Bakhoum, S. F. & Compton, D. A. Kinetochores and disease: keeping microtubule dynamics

in check! *Curr. Opin. Cell Biol.* **24**, 64–70 (2012).

5. Bakhoum, S. F. *et al.* The mitotic origin of chromosomal instability. *Curr. Biol.* **24**, R148–

R149 (2014).

6. Ertych, N. *et al.* Increased microtubule assembly rates influence chromosomal instability in

colorectal cancer cells. *Nat. Cell Biol.* **16**, 779–791 (2014).

7. Gomes Paim, L. M. & Bechstedt, S. Regulation of microtubule growth rates and their impact

on chromosomal instability. *Cell Cycle* **23**, 872–891 (2024).

8. Paim, L. M. G., Lopez-Jauregui, A. A., McAlear, T. S. & Bechstedt, S. The spindle protein

CKAP2 regulates microtubule dynamics and ensures faithful chromosome segregation. *Proc.*

Natl. Acad. Sci. **121**, e2318782121 (2024).

9. McAlear, T. S. & Bechstedt, S. The mitotic spindle protein CKAP2 potently increases

formation and stability of microtubules. *eLife* **11**, e72202 (2022).

10.   Yu, G. *et al.* RSK Promotes Prostate Cancer Progression in Bone through ING3, CKAP2,

and PTK6-Mediated Cell Survival. *Mol. Cancer Res.* **13**, 348–357 (2015).

11.   Guo, Q., Song, Y., Hua, K. & Gao, S. Involvement of FAK-ERK2 signaling pathway in

CKAP2-induced proliferation and motility in cervical carcinoma cell lines. *Sci. Rep.* **7**, 2117

(2017).

12.   Dos Santos, A., Ouellete, G., Diorio, C., Elowe, S. & Durocher, F. Knockdown of

CKAP2 Inhibits Proliferation, Migration, and Aggregate Formation in Aggressive Breast

Cancer. *Cancers* **14**, 3759 (2022).

13.   Hayashi, T. *et al.* Cytoskeleton-associated protein 2 is a potential predictive marker for

risk of early and extensive recurrence of hepatocellular carcinoma after operative resection.

Surgery **155**, 114–123 (2014).

14.   Yoo, B. H., Kang, D.-S., Park, C.-H., Kang, K. & Bae, C.-D. CKAP2 phosphorylation by

CDK1/cyclinB1 is crucial for maintaining centrosome integrity. *Exp. Mol. Med.* **49**, e354–

e354 (2017).

15.   Seki, A. & Fang, G. CKAP2 Is a Spindle-associated Protein Degraded by APC/C-Cdh1

during Mitotic Exit. *J. Biol. Chem.* **282**, 15103–15113 (2007).

16.   Hong, K. U. *et al.* Cdk1-Cyclin B1-mediated Phosphorylation of Tumor-associated

Microtubule-associated Protein/Cytoskeleton-associated Protein 2 in Mitosis. *J. Biol. Chem.*

284, 16501–16512 (2009).

17.   Hong, K. U., Kim, H.-J., Bae, C.-D. & Park, J. Characterization of mitosis-specific

phosphorylation of tumor-associated microtubule-associated protein. *Exp. Mol. Med.* **41**, 832

(2009).

18.   Hong, K. U. *et al.* Transient phosphorylation of tumor associated microtubule associated

protein (TMAP)/cytoskeleton associated protein 2 (CKAP2) at Thr-596 during early phases of

mitosis. *Exp. Mol. Med.* **40**, 377 (2008).

19.   Kim, H.-J., Kwon, H.-R., Bae, C.-D., Park, J. & Hong, K. U. Specific primary sequence

requirements for Aurora B kinase-mediated phosphorylation and subcellular localization of

TMAP during mitosis. *Cell Cycle* **9**, 2027–2036 (2010).

20.   Hong, K. U. *et al.* Functional Importance of the Anaphase-Promoting Complex-Cdh1-

Mediated Degradation of TMAP/CKAP2 in Regulation of Spindle Function and Cytokinesis.

Mol. Cell. Biol. **27**, 3667–3681 (2007).

21.   Kettenbach, A. N. *et al.* Quantitative Phosphoproteomics Identifies Substrates and

Functional Modules of Aurora and Polo-Like Kinase Activities in Mitotic Cells. *Sci. Signal.* **4**,

rs5–rs5 (2011).

22.   Olsen, J. V. *et al.* Quantitative Phosphoproteomics Reveals Widespread Full

Phosphorylation Site Occupancy During Mitosis. *Sci. Signal.* **3**, ra3–ra3 (2010).

23.   Kucharski, T. J. *et al.* An Aurora kinase A-BOD1L1-PP2A B56 axis promotes

chromosome segregation fidelity. *Cell Rep.* **44**, 115317 (2025).

24.   Karthigeyan, D., Prasad, S. B. B., Shandilya, J., Agrawal, S. & Kundu, T. K. Biology of

Aurora A kinase: Implications in cancer manifestation and therapy: BIOLOGY OF AURORA

A KINASE. *Med. Res. Rev.* **31**, 757–793 (2011).

25.   Barr, A. R. & Gergely, F. Aurora-A: the maker and breaker of spindle poles. *J. Cell Sci.*

120, 2987–2996 (2007).

26.   Nikonova, A. S., Astsaturov, I., Serebriiskii, I. G., Dunbrack, R. L. & Golemis, E. A.

Aurora A kinase (AURKA) in normal and pathological cell division. *Cell. Mol. Life Sci.* **70**,

661–687 (2013).

27.   Carmena, M., Ruchaud, S. & Earnshaw, W. C. Making the Auroras glow: regulation of

843   Aurora A and B kinase function by interacting proteins. *Curr. Opin. Cell Biol.* **21**, 796–805

844   (2009).

845   28.   Kufer, T. A. *et al.* Human TPX2 is required for targeting Aurora-A kinase to the spindle.

846   *J. Cell Biol.* **158**, 617–623 (2002).

847   29.   Zhang, R., Roostalu, J., Surrey, T. & Nogales, E. Structural insight into TPX2-stimulated

848   microtubule assembly. *eLife* **6**, e30959 (2017).

849   30.   Brunet, S. *et al.* Characterization of the TPX2 Domains Involved in Microtubule

850   Nucleation and Spindle Assembly in *Xenopus* Egg Extracts. *Mol. Biol. Cell* **15**, 5318–5328

851   (2004).

852   31.   Bischoff, J. R. A homologue of Drosophila aurora kinase is oncogenic and amplified in

853   human colorectal cancers. *EMBO J.* **17**, 3052–3065 (1998).

854   32.   Zhou, H. *et al.* Tumour amplified kinase STK15/BTAK induces centrosome

855   amplification, aneuploidy and transformation. *Nat. Genet.* **20**, 189–193 (1998).

856   33.   Sen, S., Zhou, H. & White, R. A. A putative serine/threonine kinase encoding gene

857   BTAK on chromosome 20q13 is ampli®ed and overexpressed in human breast cancer cell

858   lines.

859   34.   Reid, T. A. *et al.* Suppression of microtubule assembly kinetics by the mitotic protein

860   TPX2. *J. Cell Sci.* jcs.178806 (2016) doi:10.1242/jcs.178806.

861   35.   Wheelock, M. S., Wynne, D. J., Tseng, B. S. & Funabiki, H. Dual recognition of

862   chromatin and microtubules by INCENP is important for mitotic progression. *J. Cell Biol.*

863   **216**, 925–941 (2017).

36.   Broad, A. J., DeLuca, K. F. & DeLuca, J. G. Aurora B kinase is recruited to multiple

discrete kinetochore and centromere regions in human cells. *J. Cell Biol.* **219**, e201905144

(2020).

37.   Krenn, V. & Musacchio, A. The Aurora B Kinase in Chromosome Bi-Orientation and

Spindle Checkpoint Signaling. *Front. Oncol.* **5**, (2015).

38.   Eyers, P. A., Erikson, E., Chen, L. G. & Maller, J. L. A Novel Mechanism for Activation

of the Protein Kinase Aurora A. *Curr. Biol.* **13**, 691–697 (2003).

39.   Bayliss, R., Sardon, T., Vernos, I. & Conti, E. Structural Basis of Aurora-A Activation by

TPX2 at the Mitotic Spindle. *Mol. Cell* **12**, 851–862 (2003).

40.   A. B. Dounay & C. J. Forsyth. Okadaic Acid: The Archetypal Serine / Threonine Protein

Phosphatase Inhibitor. *Curr. Med. Chem.* **9**, 1939–1980 (2002).

41.   Takai, A. *et al.* Inhibitory effect of okadaic acid derivatives on protein phosphatases. A

study on structure-affinity relationship. *Biochem. J.* **284**, 539–544 (1992).

42.   Kucharski, T. J. Chromosome segregation fidelity is controlled by small changes in

phospho-occupancy at the kinetochore-microtubule interface. *J. Cell Biol.* (2022).

43.   Zaytsev, A. V. *et al.* Multisite phosphorylation of the NDC80 complex gradually tunes its

microtubule-binding affinity. *Mol. Biol. Cell* **26**, 1829–1844 (2015).

44.   DeLuca, K. F., Lens, S. M. A. & DeLuca, J. G. Temporal changes in Hec1

phosphorylation control kinetochore-microtubule attachment stability during mitosis. *J. Cell*

Sci. **124**, 622–634 (2011).

45.   Guimaraes, G. J., Dong, Y., McEwen, B. F. & DeLuca, J. G. Kinetochore-Microtubule

Attachment Relies on the Disordered N-Terminal Tail Domain of Hec1. *Curr. Biol.* **18**, 1778–

1784 (2008).

46.   Lyalina, T., Paim, L. M. G. & Bechstedt, S. Enzymatic Function of an Intrinsically

Disordered Protein.

47.   Yang, Z., Zeng, X., Zhao, Y. & Chen, R. AlphaFold2 and its applications in the fields of

biology and medicine. *Signal Transduct. Target. Ther.* **8**, 115 (2023).

48.   Hengeveld, R. C. C. *et al.* Development of a Chemical Genetic Approach for Human

Aurora B Kinase Identifies Novel Substrates of the Chromosomal Passenger Complex. *Mol.*

Cell. Proteomics **11**, 47–59 (2012).

49.   Perez-Riverol, Y. *et al.* The PRIDE database resources in 2022: a hub for mass

spectrometry-based proteomics evidence. *Nucleic Acids Res.* **50**, D543–D552 (2022).

50.   Geu-Flores, F., Nour-Eldin, H. H., Nielsen, M. T. & Halkier, B. A. USER fusion: a rapid

and efficient method for simultaneous fusion and cloning of multiple PCR products. *Nucleic*

Acids Res. **35**, e55–e55 (2007).

51.   Kucharski, T. J., Minshall, P. E., Moustafa-Kamal, M., Turnell, A. S. & Teodoro, J. G.

Reciprocal Regulation between 53BP1 and the Anaphase-Promoting Complex/Cyclosome Is

Required for Genomic Stability during Mitotic Stress. *Cell Rep.* **18**, 1982–1995 (2017).

52.   Ashford, A. J., Andersen, S. L. A. & Hyman, A. A. Preparation of Tubulin from Bovine

Brain. in *Cell Biology, a Laboratory Handbook* vol. 2 205–212 (Academic Press, New York,

1998).

53.   Hyman, A. *et al.* Preparation of Modified Tubulins. *METHODS Enzymol.* **196**, 478–485

(1991).

54.   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat.*

Methods **9**, 676–682 (2012).

A

B

C