Adaptation is less accessible through mutations in promoters than in coding sequences when large effect sizes are needed

Simon Aubé[1,2,3,4,5]*, Alexandre K. Dubé[1,2,3,4,5] & Christian R. Landry[1,2,3,4,5]*

1. Département de Biochimie, microbiologie et bio-informatique, Université Laval, 1045 Avenue de la Médecine, Québec, QC, Canada, G1V 0A6
2. Institut de Biologie Intégrative et des Systèmes (IBIS), 1030, Avenue de la Médecine, Université Laval, Québec, QC, Canada, G1V 0A6
3. PROTEO-Regroupement Québécois de Recherche sur la Fonction, l'Ingénierie et les Applications des Protéines, Université Laval, 1030, Avenue de la Médecine Québec, QC, Canada, G1V 0A6
4. Centre de Recherche en Données Massives de l'Université Laval, Université Laval, 1065, Avenue de la Médecine, Québec, QC, Canada, G1V 0A6
5. Centre de recherche en infectiologie, Université Laval, 2705 boul. Laurier, Québec, QC, G1V 4G2

 Correspondance: simon.aube.2@ulaval.ca and christian.landry@bio.ulaval.ca

Abstract

Mutations within promoters and coding sequences are both involved in adaptation, but their relative contributions remain to be compared directly. Using the fungal enzyme cytosine deaminase, we examine if adaptation to 5-fluorocytosine (5-FC), which requires reduced cytosine deamination, can be reached by single promoter mutations. We generated all single-nucleotide substitutions and indels in the yeast *FCY1* promoter and assayed the resulting mutants in presence of 5-FC. This revealed that no promoter mutation is sufficient for adaptation to occur. In contrast, 29% of single-nucleotide substitutions in the *FCY1* gene have previously been reported as conferring adaptation to 5-FC. We hypothesized that this discrepancy could arise if *FCY1* expression was robust to promoter mutations or if fitness was insensitive to expression level changes. Combining large-scale expression measurements with the experimental characterization of the corresponding expression-fitness function, we found that, while 24% of promoter mutations significantly affect expression, the shape of this function prevents them from being adaptive. Because this relationship is step-like, the curve is flat around wild-type level and adaptation can only emerge from a severe reduction of expression. These results highlight how adaptive phenotypes may be more mutationally accessible in coding sequences than promoters, especially when large effects are beneficial.

Introduction

For adaptation to occur, advantageous phenotypes must be accessible through mutations. This largely depends on the underlying fitness landscape or genotype-phenotype map [1,2], defined by the respective fitness of all possible genotypes. Although evolution could occur through any path created by single mutations between neighboring sequences, only immediate fitness increases are favored by natural selection. Adaptation is therefore mostly accessible through trajectories where each successive mutation is beneficial [3,4]. For this reason, any state of higher fitness will be difficult to reach if no single mutant from a starting genotype is adaptive (**Fig. 1a**). Whether beneficial mutations are immediately available thus largely determines the adaptive potential of a population of organisms.

A powerful approach to assess the accessibility of adaptation is hence to perform saturation mutagenesis, through which the effect of all single mutations in a sequence can be mapped. When focused on protein-coding sequences, such work is known as Deep Mutational Scanning or DMS [5]. This type of experiment is especially effective when an adaptive phenotype can readily be measured in the laboratory, as is the case for resistance to drugs such as antimicrobials. Multiple DMS studies have accordingly shown that resistance is accessible through single substitutions in the coding sequences of drug targets [6–11]. This work may however represent an incomplete sampling of the mutational paths that lead to adaptation, since mutations in the regulatory elements of genes can also be impactful (**Fig. 1b**). Multiple reports of adaptive mutations linked to changes in gene expression or gene dosage [12–18], as well as the identification of clear relationships between fitness and the expression level of many genes [16,19–24], indeed demonstrate that regulatory mutations can contribute to adaptation.

Numerous studies have investigated the effect of mutations within promoters and cis-regulatory sequences on expression levels using random or saturation mutagenesis, for instance in bacteria[25], yeast [26–28], *Drosophila* [29] and mammals [30–33]. Yet, much less is known about the fitness impact of mutations within promoters, especially in comparison with the effects of coding mutations in the corresponding genes. The relative accessibility of adaptive phenotypes through such changes thus remains to be assessed thoroughly (**Fig. 1c**). Even in few cases where the fitness consequences of promoter mutations are well known, for instance in *TDH3* [16,34] and *SUL1* [14] in yeast, only a handful of coding sequence mutations have comparatively been assayed [35,36].

To compare the relative contributions of promoter and coding sequence changes to adaptation, we focused on the adaptive loss of Fcy1 activity in yeast exposed to prodrug 5-fluorocytosine (5-FC). While this enzyme's canonical activity is cytosine deamination, it also converts 5-FC into 5-fluorouracil, which leads to growth arrest and cell death [37]. Accordingly, complete or partial *FCY1* loss-of-function (LOF) – which is readily available through single-nucleotide mutations within the coding sequence [6] – confers resistance and is adaptive. To estimate whether promoter mutations are similarly impactful, we performed saturation mutagenesis of the *FCY1* promoter. This experiment revealed no measurable fitness gain in medium with 5-FC, although a quarter of single-nucleotide changes significantly affect gene expression level.

Through the characterization of the underlying expression-fitness function, we show that this
discrepancy emerges from the shape of this relationship. Because this fitness function is largely
flat around the endogenous expression level, only coding sequence changes can sufficiently
decrease the amount of functional *Fcy1* to confer resistance. These findings highlight how
adaptive phenotypes may be less accessible through mutations within promoters, at least when
large changes to the underlying molecular traits are beneficial.

90
Fig. 1: Adaptation can occur through mutations in coding and regulatory sequences. a,
Not all adaptative routes are equally accessible. Two equally advantageous genotypes (1 and
2) are shown on a hypothetical fitness landscape. Both require the acquisition of three mutations
(dots) from the wild-type (WT) genotype (star), but only one of the mutational trajectories is
accessible. Because it first involves the fixation of a mildly deleterious mutation, mutational
path 1 is not favored by natural selection and hence difficult to reach from the chosen starting
point, as illustrated by dashed lines. This illustrates the importance of characterizing the effects
of single mutations from a WT genotype to study its adaptive potential. **b,** Comparison of
protein and promoter evolution, over a landscape defined by expression level and protein
function. Single mutations in the coding sequence (left) affect protein function (stability,
activity, etc), while single mutations in the promoter modify gene expression level and thus
protein abundance. As such, they explore two complementary slices of the landscape.
Saturation mutagenesis experiments on a gene and its promoter respectively study the same two
slices, at WT expression level or WT protein function. Such studies can thus assess whether
high fitness regions can be reached through each of the two types of mutations. **c,** The current
knowledge of fitness landscapes is incomplete. Since systematic studies of the fitness effects
of mutations have mostly explored protein function, only a small region corresponding to WT

expression level is known in most cases. The orthogonal region is unknown, such that the accessibility of adaptive (high-fitness) mutations in promoters remains to be assessed thoroughly. Here, we examine the fitness effects of regulatory mutations for a gene whose fitness landscape has previously been studied in the protein dimension.

Results

Measuring the effects of mutations within the *FCY1* promoter

To identify promoter regions to mutate, we performed deletions of successive 77 bp fragments of the intergenic region upstream of *FCY1*. This revealed that only the most proximal fragment, among four, is essential for gene function (**Extended Data Fig. 1**). For this reason, and because most regulatory elements in yeast promoters are found within the first 150 bp [38], we focused on the 154 bp immediately upstream of *FCY1*. This corresponds to two fragments hereafter named F3 and F4 (**Fig. 2a**). Within this region, we performed all single-nucleotide mutations – substitutions, insertions and deletions. The corresponding variants of F3 and F4 were inserted at the endogenous *FCY1* locus using a CRISPR-Cas9-based pooled approach (**Fig. 2b**, *Methods* [39,40]). This resulted in 1021 unique mutated sequences (462 substitutions, 463 insertions and 96 deletions), of which 1020 were successfully recovered by sequencing of the mutant pools (**Supplementary Fig. 2**). This process was repeated four times for each of the three classes of mutations (**Fig. 2c**). The corresponding pools of substitutions, insertions and deletions were then combined, resulting in four independent transformation replicates, each containing all mutations. Because the set of substitutions was obtained by successively degenerating each of the 154 nucleotide positions, the wild-type (WT) *FCY1* promoter was also (re)constructed along with the desired mutants. Within each combined transformation replicate, its initial frequency was ∼17%.

From each of the four transformation replicates, two culture replicates were made, which were both grown in liquid media with and without 5-FC, resulting in 16 cultures. The prodrug was added at a final concentration of 12 μM, which has been shown to inhibit the growth of *Saccharomyces cerevisiae* by ∼50% and was used in the previous DMS of the *FCY1* coding sequence [6] (**Supplementary Fig. 3**). Additional replicates of all 16 cultures were also done by combining the promoter mutants with 960 previously characterized coding sequence variants[6], to ensure that the fitness effects of the two types of mutations could be directly compared (**Extended Data Fig. 2**).

All 32 cultures were grown for ∼20 generations in liquid media, with dilution every ∼7 generations to maintain exponential growth (*Methods*). Samples were taken at each dilution step, as well as before the start of the experiment, resulting in a series of timepoints from T0 to T3 (**Fig. 2c**). The relative frequencies of all mutations were monitored by sequencing of the F3-F4 region (**Fig. 2a**). Due to the toxicity of Fcy1 activity in the presence of 5-FC, we expected that mutations reducing gene expression would increase in frequency, outcompeting those resulting in a higher expression level. Accordingly, we estimated the selection coefficient (*s*) associated with each promoter sequence from the slope of its log-scaled frequencies relative to the WT [41] (**Fig. 2c**).

Fig. 2: Experimental design of the *FCY1* promoter deep mutational scanning. a, 5-FC is converted into 5-FU, a toxic compound, by the cytosine deaminase (Fcy1p), such that *FCY1* expression can have a strong effect on fitness. Fragments 3 (F3) and 4 (F4) of the *FCY1* promoter were replaced with a library of synthetic sequences corresponding to all possible single-nucleotide substitutions (Sub), insertions (Ins) and deletions (Del). The oligonucleotides used to sequence the region and genotype mutants are marked in red (short fragment) and black (extended fragment, including the recombination breakpoints). **b,** Mutated F3 and F4 sequences were inserted at the endogenous yeast *FCY1* locus by CRISPR-Cas9 genome editing. A NAT resistance cassette was first introduced in F3 or F4 by homologous recombination. Using a specific guide RNA, Cas9 could be targeted to the cassette, such that synthetic DNA fragments would be inserted at F3 or F4. This step was repeated four independent times. **c,** The three types of pools (Sub, Ins, Del) were combined and grown in synthetic complete media (SC) and then transferred to media with 5-FC to allow for competition based on the abundance of Fcy1p. Loss of function of *FCY1* has no detectable effect in SC media without 5-FC. At every timepoint, DNA was extracted and the edited locus was sequenced to follow the frequency of each mutant in the culture, allowing to estimate selection coefficient for each mutation from their enrichment or depletion through time. Schematics created in BioRender. Aubé, S. (2025) https://BioRender.com/frtbee2.

Mutant pools sampled from the competition experiment were first genotyped by sequencing a 174 bp fragment, which covered the full F3-F4 region and the first 20 bp of *FCY1* (red oligonucleotide pair; **Fig. 2a**). There were however major reproducibility issues. While estimated selection coefficients were consistent between culture replicates of the same transformation, they diverged substantially across transformation replicates (**Supplementary Fig. 1a**). In addition, comparisons of the distributions of selection coefficients across the different time intervals revealed a gradual shift of the mean. From ~0 for the T0-T1 interval, it decreased below -0.15 for the full competition experiment (**Supplementary Fig. 1b**). Through further experiments, we found that these effects were caused by secondary mutations introduced during library construction (**Supplementary Fig. 1c-d**). We modified our sequencing protocol to genotype a longer region (black oligonucleotide pair; **Fig. 2a**) and repeated the analyses to eliminate these effects. We explain the observations in detail in the supplementary material (**Supplementary Note**) as we suspect these could be common in genome editing studies and thus be of broad interest.

Fitness is robust to *FCY1* promoter mutations, but sensitive to coding mutations

After correcting for the impact of secondary mutations, the inferred selection coefficients for the F3-F4 mutants revealed only small fitness effects (**Fig. 3a**). Across all assayed promoter mutations, the median *s* estimates obtained from replicate experiments ranged from ~-0.06 to ~0.06 gen^{-1}. Experimental noise was moreover of a similar magnitude as these deviations from 0, as shown by an average difference of ~0.04 gen^{-1} between the highest and lowest *s* among replicate estimates for the same genotype, up to ~0.18 gen^{-1} (**Supplementary Table 8**). A small number of secondary mutations remained undetected (not found within the longer sequencing fragment), resulting in outlier selection coefficients for one pair of culture replicates within some F3-F4 genotypes. Overall, following a stringent removal of these rare outliers (**Supplementary Fig. 9**), none of the 1020 *s* estimates for *FCY1* promoter mutations appeared significantly different from 0 ($p > 0.05$; one-sample t-test with Benjamini-Hochberg correction at FDR=5%).

This lack of fitness effects is not due to a failure of the competition experiment, as confirmed by two layers of controls. When considering all mutations (F3-F4 and secondary), the variance of estimated selection coefficients is indeed much narrower for cultures without 5-FC (**Supplementary Fig. 10**). More importantly, the fitness inferred for the added coding sequence variants replicate previous findings on the effect of amino acid substitutions in Fcy1p [6] (**Fig. 3b**). There is in addition almost perfect agreement between the two experiments regarding the number and identity of resistance-conferring mutations (**Extended Data Fig. 3**). Although samples which contained coding sequence variants also included the complete pool of promoter mutations (**Extended Data Fig. 2**), the resulting cultures were not used to precisely estimate F3-F4 mutational effects. This is because the numerous resistant genotypes within the coding sequence pool outcompeted promoter mutants and rapidly took over the population (**Extended Data Fig. 4a**), thus decreasing the resolution of F3-F4 fitness estimates. Although much

213   noisier, the corresponding selection coefficients were qualitatively similar to those reported in
214   **Fig. 3A** (**Extended Data Fig. 4b**).

215
Fig. 3: Mutations in the *FCY1* promoter do not have a measurable impact on 5-FC
resistance. a, Fitness effects of single-nucleotide mutations in the *FCY1* promoter in 5-FC
conditions. All substitutions (Sub; n=462), insertions (Ins; n=462) and deletions (Del; n=96)
are shown. Points display the median selection coefficient for each mutant genotype, while
error bars show the minimal and maximal values observed at each position along the mutated
region. Nucleotide positions which appear to be missing are due to mutations which result in
the same genotype (**Supplementary Fig. 2**), for instance deletion of any single base in a
homopolymer region. Only data from two unbiased transformation replicates (each duplicated
into two culture replicates; four samples in total) are shown (**Supplementary Note**,
Supplementary Fig. 7). **b,** Comparison of median selection coefficients for *FCY1* coding
sequence mutants added to our experiment with previously published variant effect scores
measured in 5-FC [6]. Values have been aggregated into medians for all genotypes which result
in the same amino acid sequence, such that n=311 points are shown. A LOWESS fit highlights
a mostly monotonically increasing trend, as shown by a high Spearman correlation ($p <$
10^{-64}). **c,** Distribution of variant effect scores obtained in 5-FC [6] for all single-nucleotide
substitutions in *FCY1*. For each genotype (n=1411), the score reported for the corresponding
amino acid sequence was used [6]. Only 20 of the possible nucleotide substitutions are missing,
mostly in the first and last codons. Variants which are susceptible and resistant to 5-FC are
respectively identified with "S" and "R". About 29% are classified as resistant (*Methods*).

235
236   Negative and positive controls thus confirm that 5-FC selected against cytosine deaminase
237   activity, as expected, and that the absence of measurable fitness effects is not artefactual. This

weak impact of promoter mutations contrasts markedly with the strong effects of mutations in the *FCY1* coding sequence. Within this gene, ~29% (408/1411) of single-nucleotide substitutions indeed result in codon changes which confer resistance to 5-FC in the same conditions (**Fig. 3c**)[6]. Moreover only 67 of these 408 mutations introduce a premature stop codon. Notably, this comparison underestimates the accessibility of adaptation through mutations in the *FCY1* gene. Most indels in this coding sequence, which were not assayed, are indeed likely to grant resistance by introducing frameshifts.

Within our *FCY1* model system, single-nucleotide mutations in the promoter thus appear to provide little to no paths to adaptation, contrary to changes in the corresponding coding sequence. Such weak fitness effects of promoter mutations could emerge through two mechanisms. First, the abundance of Fcy1p could be robust to changes in its promoter. Second, fitness in 5-FC conditions could be insensitive to variations in *FCY1* expression level, such that no mutational effects would be measurable through bulk competition assays. Discriminating between these two competing hypotheses first requires measuring the impact of *FCY1* promoter mutations on expression level.

Many mutations in the *FCY1* promoter substantially affect expression level

To obtain expression level measurements for all promoter mutants, we used a sort-seq approach[42,43]. The complete pool of F3-F4 single mutants was reconstructed in a genetic background where the distal part of the *FCY1* gene – starting from its 43rd base – was replaced with a mEGFP sequence (*Methods*, **Extended Data Fig. 5**). As such, the translation initiation context could be maintained while adding a fluorescent reporter to measure promoter activity. Yeast cells of the corresponding mutant library were sorted into five bins (numbered 0 to 4) according to their log-scaled fluorescence intensity (**Fig. 4a**). All the resulting samples were sequenced and read counts were used to compute a weighted average of bins for each genotype (*Methods*). The effect of each mutation was quantified using a Δ_{Ref} score equal to the difference between the corresponding mean bin and the average obtained for the WT. This was performed separately within each of six replicates (three transformation replicates, each subdivided in two culture replicates).

A narrow range of scores was obtained, with the median Δ_{Ref} of most promoter mutations falling between -1 and 0.5 (**Fig. 4b**). Of the 1020 F3-F4 mutants, 1019 were covered in at least two replicates after abundance-based filtering (*Methods*). Many secondary mutations were also recovered (**Fig. 4b**), including indels in the first 42 bp of *FCY1*. Interestingly, while most such mutations were associated with a low fluorescence intensity and thus a negative Δ_{Ref}, not all completely abolished expression. We specifically found out that indels which occurred before the second methionine codon of *FCY1* maintained ~10% of the WT fluorescence level (**Extended Data Fig. 6 a,b**). This is consistent with a previous report of a minor translation initiation site at this codon [44].

Fig. 4: Mutations in the *FYC1* promoter impact expression level. a, Schematics of the sort-seq experiment. Yeast cells from pools of promoter mutants were sorted into five bins according to their fluorescence intensity (representative distribution shown). From the sequencing of each bin, WT-relative fluorescence intensity could be estimated. Created in BioRender. Aubé,S. (2025) https://BioRender.com/uzzv5xy. **b,** Median sort-seq scores obtained for all variants detected across six replicate experiments. In addition to 1019 F3-F4 variants ("Promoter"), 28 single mutations in the first 42 bp of *FCY1* were identified ("CDS"), of which the vast majority are indels. The full distribution of scores ("All") as well as the 30 mutants selected for reconstruction ("Validation") are shown, split between the two categories in each case. **c,** Calibration curve of sort-seq scores. Points display the medians of sort-seq scores (n=6) and WT-relative fluorescence intensities (n=3) for all 30 validation genotypes. Horizontal and vertical error bars show the minimal and maximal observations associated with each median. The high correlation between medians is shown with Spearman's rho ($p < 10^{-23}$). The line

and the corresponding shaded area display a LOWESS fit with its 95% confidence interval
(*Methods*). **d,** Distributions of WT-relative expression levels for all single-nucleotide promoter
mutations, as inferred from the calibration of sort-seq scores. At the top, distributions are shown
separately for substitutions (Sub; n=461), insertions (Ins; n=460) and deletions (Del; n=96),
with a dashed line indicating WT expression level (1.0). At the bottom, the full distribution,
combining all three types of single mutants, is shown (n=1019). **e,** Distribution of promoter
mutations with a significant effect on expression (n=245), along the mutagenized region.
Separately for Sub (n=113), Ins (n=111) and Del (n=21) variants, the estimated WT-relative
expression (point) and its corresponding 95% prediction interval (error bar) are shown for each
mutant affecting FCY1 expression by 5% or more (minimum or maximum of the corresponding
95% prediction interval; *Methods*). Shaded rectangles display the minima and maxima across
all prediction intervals for each position along the sequence, including variants which are not
shown. In the upper plot, vertical lines show the midpoint of hits (n=41) to known TF
recognition motifs, identified using YeTFaSCo [45]. In the lower plot, the black and white
rectangles respectively display A/T and G/C positions. Promoter positions corresponding with
the two main clusters (sites "1" and "2"), the approximate AT-rich region and the TATA-like
element ("TA") [46,47] are also identified.

To validate the sort-seq assays, we selected 30 mutants for independent reconstruction (**Fig.
4b**). For each newly constructed strain as well as for the WT reference, fluorescence intensity
was measured by flow cytometry (*Methods*). The resulting WT-relative intensities are in strong
agreement with the corresponding median Δ_{Ref} scores (**Fig. 4c, Extended Data Fig. 7**). We
additionally used this validation to draw up a calibration curve, enabling the conversion of sort-
seq scores into more easily interpretable relative fluorescence intensities. To this end, we fit a
LOWESS regression (**Fig. 4c**), to avoid making specific assumptions about the relationship
between the two variables. This was performed within a bootstrapping routine which involved
repeated resampling of Δ_{Ref} values as well as validation fluorescence measurements, thus
generating 95% prediction intervals for the curve itself and for the relative fluorescence
intensities of all genotypes (*Methods*, **Supplementary Table 9**). We hereafter refer to the
medians of the latter intervals as WT-relative expression levels for the corresponding
mutations, where a value of 1 denotes the WT level.

Examining the inferred relative expression levels for all F3-F4 mutants reveals a slight bias
towards expression increases, with medians lying between 1.04 and 1.06 (**Fig. 4d**). That
mutations are more likely to increase expression than to decrease it may appear surprising,
since Fcy1p is already highly expressed and is among the 10% most abundant proteins in *S.
cerevisiae* [48]. While this bias could be artefactual, the y-intercept of the calibration curve
suggests otherwise (**Fig. 4c**). It is very close to 1 (~1.00, with 95% prediction interval from
0.98 to 1.02) – as expected for a Δ_{Ref} of 0 –, although no WT measurements were used in the
LOWESS fitting.

The overall distribution of effects highlights that most single mutations in the *FCY1* promoter
weakly affect expression levels, with 75% of variants lying between values of 0.98 and 1.11
(**Fig. 4d**). Yet, we also observe a small number of mutations with larger effects, decreasing
expression to ~50% of its WT level, or even lower. This tail appears more pronounced than

previously reported in yeast [26,27], potentially because we specifically mutated the most proximal
part of the *FCY1* promoter, where mutations are more likely to disrupt processes such as
transcription initiation. Among the 1019 F3-F4 mutations, 245 significantly affect *FCY1*
expression, by 5% or more (95% prediction intervals). In agreement with a previous study of a
yeast promoter [14], these significant mutations are not distributed uniformly along the
mutagenized promoter region (**Fig. 4e**). This distribution reveals important determinants of
FCY1 expression level.
344
The fact that significant mutations form two main clusters, from positions -132 to -127 (site 1)
and from -125 to -120 (site 2), first highlights the importance of transcription factor (TF)
binding. Within these two sites, 92.4 % (73/79) of all variants change expression by 5% or
more, in a consistent direction within each cluster. Both sites are notably within a 80 bp
segment known to be critical for TF binding in yeast promoters [28,42,49], while the whole -132 to
-120 region is additionally a hotspot of predicted TF binding sites [45] (**Fig. 4e top;**
Supplementary Table 10). As such, the clusters likely reveal TF binding sites. Interestingly,
the deletion of either of the two sites is not more impactful than the strongest mutations within
the corresponding cluster (**Extended Data Fig. 8**), suggesting that single mutations are
sufficient to disrupt them.
355
The position of the two most impactful mutations additionally highlights how alterations of
transcription initiation may severely reduce *FCY1* expression. These variants, a substitution
and an insertion which respectively decrease expression to ~16% and ~13% of its WT level
(cytometry; **Fig. 4c**), both occur at position -9 (**Fig. 4e**). This position happens to be
immediately downstream of the previously identified TATA-like element of the *FCY1*
promoter [46,47] – from -17 to -10 –, where the transcription pre-initiation complex assembles.
The corresponding mutational effects might thus reflect the disruption of transcription
initiation. The conflicting report of a most abundant transcript isoform beginning further
upstream at -29 [50], however makes the underlying mechanism unclear.
365
Finally, the complete set of significant mutations underscores that most single-nucleotide
changes within the poly(dA:dT) region of the *FCY1* promoter weakly affect its transcriptional
activity. The corresponding AT-rich sequence, approximately between positions -118 and -73
(17.39% GC; **Fig. 4e**, bottom), is indeed comparatively depleted in strong mutational effects.
Only 17.6 % (52/295) of variants within it are significant, compared to 26.7 % (193/724) in the
remainder of the mutagenized region. This may seem contradictory, considering that such
regions are important for the establishment of nucleosome-depleted regions within promoters
[51,52], which favor transcription and can be sufficient to induce expression from a downstream
core promoter [53]. It is however consistent with previous work on the *URA3* promoter, which
showed that deletion of more than half of the corresponding 50 bp poly(dA:dT) was necessary
to abolish expression [54].
377
378 Overall, we show that 24% (245/1019) of single-nucleotide mutations in the *FCY1* promoter
379 have a significant effect on expression level, many of which appear to disrupt the binding of

380 TFs at two putative sites. As such, *FCY1* expression is sensitive to mutations, meaning that
381 robustness cannot explain the lack of fitness effects in 5-FC conditions.
382
The expression-fitness relationship for *FCY1* is mostly flat

To investigate the mechanism underlying the absence of measurable fitness effects, we set out
to characterize experimentally the expression-fitness relationship for *FCY1* (**Fig. 5a**). Six yeast
promoters covering a wide range of expression levels were used to construct two sets of strains,
in which each promoter was inserted upstream of either the native *FCY1* sequence or the
previously described 42bp-mEGFP construct. The *FCY1* strains were grown in media with 5-
FC (12 μM; like in the competition experiment) – to obtain growth rate measurements –, while
mEGFP strains were used to measure expression levels (*Methods*). For each promoter, the
measurements from the two strains were combined to draw the corresponding expression-
fitness curve (**Fig. 5a**).
393
This experiment revealed a very steep fitness function, where growth rate decreases sharply
until *FCY1* expression reaches ~18% of its WT level and then plateaus (**Fig. 5b**). Consequently,
this fitness curve is flat around the WT, such that small expression changes have no fitness
effect. The shape of this function is thus consistent with the fact that no single-nucleotide
promoter mutation conferred resistance to 5-FC in our bulk competition experiment.
Interestingly, varying the concentration of the prodrug does not impact the sensitivity of fitness
to changes in *FCY1* expression. Across 5-FC concentrations resulting in ~25% to ~75%
inhibition for the WT, growth rate variation is restricted to the same narrow range of relative
expression levels and only the height of the corresponding plateau varies (**Extended Data Fig.
9a**). The same phenomenon is observed in media lacking uracil, where the deamination of
cytosine into uracil by Fcy1p is essential for growth. Across multiple cytosine concentrations,
growth rate increases rapidly until ~18% of WT expression is reached and then remains
constant as expression is increased further (**Extended Data Fig. 9b**). The expression-fitness
function in 5-FC therefore also reflects the canonical deamination function of the Fcy1 enzyme.
408
To increase the resolution of the experimental fitness function around WT expression, we
leveraged previously characterized validation mutants. The set of alleles used in the calibration
of sort-seq measurements, which also includes a small number of indels in the coding sequence,
was reconstructed in the native *FCY1* background and the resulting strains were phenotyped in
5-FC conditions (*Methods*). In combination with the corresponding relative fluorescence
intensities, these growth measurements confirm the flatness of the curve around WT expression
level, with all promoter mutants displaying WT-like growth rates (**Fig. 5c**). Among three
assayed coding sequence mutations, the only one which reached the high fitness region is
associated with a ~98% decrease in fluorescence intensity (**Supplementary Table 11**). This
highlights how resistance – and thus adaptation – is only possible through large reductions in

Fig. 5: The *FCY1* expression-fitness function is mostly flat, making fitness robust to promoter mutations. a, Strains with varying expression levels were constructed to measure the fitness function. A set of yeast promoters covering a wide range of activities were introduced in the genome in place of the endogenous *FCY1* promoter, either upstream of FCY1 or of a fluorescent reporter. Created in BioRender. Aubé, S. (2025) https://BioRender.com/1p60mp7. **b,** Expression-fitness relationship for *FCY1* in 5-FC conditions. All strains with *FCY1* expression equal to or higher than ~18% of its WT level display a growth rate which is indistinguishable from the WT. For each promoter, medians of growth rate and WT-relative fluorescence (n=3) are shown along with the corresponding minima and maxima (error bars). The fitness function is displayed as a fitted exponential decay function, with its corresponding 95% confidence interval (*Methods*). **c,** Growth rates of reconstructed mutants are consistent with the fitness function. For validation mutants (**Fig 4c**) which could also be reconstructed in the *FCY1* background, medians (n=3) are shown for growth rates in 5-FC [12µM] media and WT-relative fluorescence intensities. This includes

single promoter mutations ("Promoter"; n=25) and indels in the first 42 bp of the gene ("CDS"; n=3). The dashed line identifies the WT expression level (1.0). **d,** Reconstructed promoter mutations are mostly neutral. For the same set of single mutations in the promoter (n=25), medians (n=3) are shown for selection coefficients and relative fluorescence intensities, with error bars displaying the corresponding minimal and maximal replicates. Selection coefficients are computed from growth measurements (*Methods*) shown in the previous panel, relative to the WT strain. The previously fitted fitness function (panels **b** and **c**) is displayed with its corresponding 95% confidence interval, after conversion into selection coefficients. Only the region of the expression-fitness curve where promoter mutants lie is shown.

Fcy1 activity, in this case due to an almost complete loss of protein expression. Zooming in on the promoter mutants further illustrates this point. Among them, the two single mutations with the largest effect barely reach the threshold below which fitness rapidly increases, although they respectively decrease *FCY1* expression to ~16% and ~13% of its WT level (**Fig. 5d**). These two mutants nonetheless display a slight growth advantage, corresponding to selection coefficients on the order of ~0.03 (**Fig. 5d**) after conversion (*Methods* [55]), which is near the detection limit of our bulk competition assay (**Fig. 3** and accompanying text). Even the two most fit promoter variants are thus readily outcompeted by the many resistance-conferring mutations which can emerge in the coding sequence, associated with selection coefficients over 0.5 (**Extended Data Fig. 6c**). As such, the shape of the fitness function, combined with the fact that no single-nucleotide change in the promoter completely abolishes expression, severely restricts the accessibility of adaptation through cis-regulatory mutations in this *FCY1* system.

Discussion

Adaptive mutations can in principle occur in coding and regulatory regions but the contribution of each remains difficult to estimate [56]. One powerful approach to do so is to focus on a trait whose relationship with fitness is known and systematically assay the adaptiveness of all mutations in both the coding and regulatory regions. To this end, we focused on the promoter of the *FCY1* gene, a protein-coding region within which single-nucleotide changes have been shown to confer resistance to the antifungal 5-FC [6,57]. We surprisingly show that no promoter mutation has a measurable fitness effect in yeast exposed to this molecule, although large changes in expression level (reaching -87%) are observed. In this system, adaptive phenotypes are thus much more readily available through nucleotide variation in the coding sequence than within the corresponding promoter.

In contradiction with this result, our initial data instead suggested that most mutations were highly deleterious. Our investigations revealed that this was due to secondary resistance-conferring mutations introduced during library construction, outside of the sequenced region (**Supplementary Note; Supplementary Fig. 1,4-8**). These beneficial mutations largely occurred in the WT, since this genotype was generated at high frequency within the mutant pool. This comparatively made all true single mutants appear less fit. Interestingly, bias thus arose from the reconstruction of the WT within the CRISPR library, which is widely regarded as a best practice to ensure the accuracy of mutant fitness estimates [58]. Such artifacts may be common in large-scale genome editing experiments, due to the increased frequency of single-

nucleotide mutations – and especially indels – in synthetic DNA oligonucleotides [59]. This
highlights the importance of designing library construction workflows which minimize the use
of genic regions as homology arms to guide recombination. Alternatively, these segments
should at least be fully included in the genotyping region, as we did to correct our experiment.

In this study, we observed a striking contrast between promoter and coding sequence mutations,
with none of the former conferring resistance, while 30% of the latter do so [6]. This discrepancy
can be explained from the shape of the corresponding fitness function, which is mostly flat, but
displays steep variation below 18% of the WT amount of functional protein. Due to the step-
like nature of this relationship, most promoter mutations have no impact on fitness. While many
similar expression-fitness curves have been reported in yeast [20,22], functions where small
variations around the WT are immediately impactful are also observed [21,22]. If the abundance-
fitness relationship for *FCY1* was instead in the latter category, many promoter mutations
would also be able to reach the state of higher fitness. As such, a different shape of fitness
function could result in more equal proportions of adaptive mutations in the gene and its
promoter. Yet, there is an additional factor underlying the reduced accessibility of adaptation
through changes in the *FCY1* promoter: that mutations in the corresponding coding sequence
have much larger effects at the molecular level. In accordance with the fitness function we
report, the many resistance-conferring mutations in the *FCY1* gene indeed suggest that almost
a third of substitutions in this sequence decrease protein abundance and/or activity below 5%
of its WT level. In contrast, only two promoter mutations – among more than 1000 – approach
the low expression threshold below which fitness begins to increase. Whether coding or
regulatory mutations have the most potential to contribute to adaptive changes therefore likely
depends on both the shape of the corresponding fitness function and the respective distributions
of mutational effects (DME) for promoter and coding mutations. If the latter generally have
larger effects on molecular phenotypes like protein abundance and activity, they would be the
main contributor to adaptation when large changes are needed.

While the lack of strong mutational effects in the *FCY1* promoter could mean that this gene's
expression level is especially robust to mutations, multiple observations suggest that it is not
an exception in that regard. The substantial number of weak but significant expression changes
we report is first largely consistent with previous work. Large-scale mutagenesis of promoters
and regulatory elements in yeast [26,27] and mammals [30–33], have indeed shown that single
mutations at most positions have weak to moderate impact. The *FCY1* promoter is in addition
a typical yeast promoter, being TATA-less like the majority of such regions [46,60] and featuring
the most common poly-T/poly-A configuration [52]. Although bidirectional, it drives much
higher expression of *FCY1* than of the opposite gene and hundreds of such promoters have
been identified in the *S. cerevisiae* genome [61]. As such, the distribution of mutational effects
we report is likely generally representative of yeast promoters. Second, our work mirrors
previous accounts associating weaker functional impacts with changes within promoter
regions. A study of fluorescent reporter proteins for instance reported many more coding
sequence than cis-regulatory changes among fluorescence-decreasing mutations [62]. In addition,
a recent compilation of over 14 000 resistance-conferring mutations in 94 fungi uncovered only
34 promoter mutations, all of which involve insertions of tens of nucleotides [63].

More importantly, the discrepancy between the respective DME of promoter and coding
sequence changes in *FCY1* is consistent with the general picture which emerged from earlier
DMS studies of protein-coding sequences and regulatory regions [64]. While reported DMEs for
proteins are often strikingly bimodal – with the abundance of strongly deleterious mutations
sometimes comparable to that of WT-like mutants –, those of cis-regulatory elements tend to
be unimodal and centered around the WT or to display a minor mode of a small number of
strongly deleterious effects [64]. More recent mutagenesis studies have continued to frequently
report distributions of effects which are consistent with this assertion [25,65–70], sometimes even
showing that most mutations in a protein sequence are either highly deleterious or strongly
beneficial [7].

This multimodality of coding sequence DMEs likely emerges from nonlinear relationships
between the biophysical properties of proteins and fitness-relevant traits [64,71]. This can for
instance be the case of protein stability, which dictates the fraction of correctly folded – and
thus functional – proteins through a sigmoid function [70,72–74]. Due to this relationship, most
mutations affecting stability would either have little effect or drastically decrease activity, for
a WT protein which stably folds and is thus at the peak of the corresponding curve. This
phenomenon likely underlies the high availability of 5-FC resistance through LOF mutations
in the *FCY1* gene. Among resistance-conferring single-nucleotide substitutions in *FCY1*, only
16% (67/408) introduce a premature stop codon, while the predicted impact of mutations on
protein stability is strongly correlated with fitness effects and resistant variants with reduced
stability have been characterized experimentally [6,75]. Although similar nonlinearities are
expected in the binding of regulators to DNA [64,76], almost entirely abolishing gene expression
may be comparatively difficult to achieve. This is illustrated both by spurious transcription of
the yeast genome [77] and by the fact that exogenous or random DNA introduced in yeast is
generally transcribed [78,79]. As such, the unequal contributions of promoter and coding changes
to 5-FC adaptation likely arise from fundamental differences in the effects of the two types of
mutations. Although it is possible that this only applies when adaptation involves LOF,
previous studies have shown that mutations in promoters can also be insufficiently adaptive in
gain of function contexts. For instance, while increasing the expression of antifungal target
ERG11 through gene copy number variation (CNV) can confer resistance in fungi [80,81], no
equally adaptive single promoter mutation has been reported [63]. Similarly, whereas CNVs of
the sulfate transporter *SUL1* consistently arise in yeast evolved in sulfate-limited conditions
[12,13], single mutations in the corresponding promoter have been shown to provide much less
adaptive benefit [14].
Overall, it thus appears plausible that adaptation is generally more accessible through mutations
in protein-coding sequences than in their cis-regulatory regions, at least when large changes to
molecular traits like protein abundance and activity are needed (**Fig. 6a**). This may however
vary across genes and adaptive phenotypes, since the availability of beneficial fitness effects
through mutations in promoter and coding sequences depends on their respective DME and on
the shape of the fitness function. A higher curvature around WT expression / activity levels
could for instance make adaptation accessible through promoter as well as coding mutations,
even if the former had smaller effects as in the *FCY1* model system (**Fig. 6b**). Other shapes of

fitness functions may in contrast restrict adaptation to small variations of molecular traits, potentially favoring cis-regulatory mutations (**Fig. 6c**). Additional studies are hence needed to fully map which contexts may favor one type of change over the other. With the current work, we have shown that combining the DMS of genes and cis-regulatory regions with the characterization of the corresponding fitness functions is a powerful approach to do so.

Fig. 6: Different shapes of fitness function may favor mutational effects of different magnitudes. Assuming that mutations in coding sequences have larger effects on molecular traits than changes within promoters, as is the case for *FCY1*, this can modulate whether adaptation is accessible through one type of mutations or the other. Although we only measured *FCY1* expression, we generalize expression-fitness relationships to total enzymatic activity. For each of the three scenarios, the corresponding fitness curve is shown as a solid line. **a,** A step-like function, which is flat around WT, favors mutations with large effects on total activity. Accordingly, adaptation is accessible through coding sequence changes (arrow), while promoter mutations do not increase fitness and are not selected for (blunt arrow). This is what we report for *FCY1*. **b,** A fitness function with high curvature around WT allows mutations with small and large effects to be adaptive. Here, less impactful promoter mutations are also beneficial, such that adaptation can occur through one large-effect coding sequence mutation or through successive promoter mutations. **c,** A non-monotonic fitness function may favor small changes in total activity. Because they overshoot the curve's optimum, coding mutations with large effects are deleterious and thus not selected for (blunt arrow), while promoter changes with smaller effects can be adaptive (arrow). Created in BioRender. Aubé, S. (2025) https://BioRender.com/enzcymq.

Methods

Strains, oligonucleotides, plasmids and culture media

All *S. cerevisiae* strains constructed in this study were generated from a BY4742 (*his3Δ leu2Δ lys2Δ ura3Δ*) background [82] harboring a hygromycin resistance marker at the *HO* locus (*hoΔ::HPHMX*). This AKD1123 parental background and all other strains are listed in **Supplementary Table 1**, along with the identity of the corresponding genomic inserts and PCR primers as well as the selective media used. The sequences and references for all oligonucleotides used in this study can be found in **Supplementary Table 2**, while references for all plasmids are listed in **Supplementary Table 3**. Recipes for culture media are detailed in **Supplementary Table 4**. When specified, selective media was prepared by adding

one or multiple of the following agents: geneticin [200 μg/mL] (G418; Bioshop Canada), nourseothricin [100 μg/mL] (NAT; Jena Bioscience) and hygromycin B [250 μg/mL] (HYG; Bioshop Canada). All Sanger sequencing was performed at the sequencing platform of Centre de recherche du CHU de Québec-Université Laval (Université Laval, Québec, Canada).

Yeast transformation

Whenever inserts which included a selection marker had to be inserted in the yeast genome, a standard lithium acetate yeast transformation protocol was used [83]. After selection of transformants on the appropriate selective media, four to eight clones were selected for validation PCR, depending on the strain. Unless stated otherwise, colony PCR was performed on each side of the integration site to confirm proper integration, using Taq polymerase (Bioshop Canada). All genomic inserts were amplified using the KAPA HiFi HotStart DNA polymerase (Roche Molecular Systems).

This approach was for instance used to delete each fragment F1 to F4 of the intergenic region between *JID1* (*YPR061C*) and *FCY1* (*YPR062W*), as well as the full intergenic segment and the complete *FCY1* region (including the corresponding gene), by the insertion of a *natNT2* cassette [84]. This resulted in a set of six strains harboring a different deletion of the *FCY1* region (SA0127, SA0128, SA0129, SA0130, SA0131, SA0132). Similar deletions of regions F3 and F4 were also performed in the 42bp-mEGFP construct obtained by CRISPR-Cas9 yeast transformation (see below), generating fluorescent strains SA0246, SA0247 and SA0254

CRISPR-Cas9 yeast transformation

When sequences needed to be inserted in the yeast genome without any flanking selection marker, a CRISPR-Cas9 approach was used. This required the prior generation of a strain harboring a *natNT2* cassette at the chosen insertion site, which was done through standard yeast transformation as described above. Competent cells from this genetic background could then be cotransformed with the pCAS-NAT plasmid (**Supplementary Table 3**) – encoding a guide RNA engineered to target the Cas9 endonuclease to the *natNT2* cassette [39] – and the desired repair DNA, amplified with KAPA HiFi HotStart DNA polymerase (Roche Molecular Systems). A previously published CRISPR-Cas9 transformation protocol was followed [85]. Briefly, competent cells were prepared as described, and transformed with unpurified PCR amplification of the insert, DNA carrier [10 mg/ml], pCAS-NAT and PLATE solution (polyethylene glycol 2000 [40%], lithium acetate [0.1 M], Tris-EDTA [10 mM Tris-Cl, 1 mM EDTA] buffer [pH=8.0]). After mixing, cells were incubated for 30 minutes at 30°C, heat shocked for 15 minutes at 42°C, resuspended in YPD media and incubated for 4-6 hours at 30°C. Transformation reactions were plated on YPD media containing G418, to select for uptake of the pCas plasmid, and incubated at 30°C for 48 hours. Eight colonies from each transformation were then grown overnight (30°C with shaking) in liquid YPD and streaked on solid YPD. After 48 hours of incubation at 30°C, one isolated clone from each streaking was grown overnight in YPD (30°C with shaking) and spotted on both YPD and YPD+G418 solid media, to confirm the loss of the pCAS plasmid. Because the AKD1123 parental background is resistant to hygromycin, HYG was added to the YPD throughout. Correct integration was then validated by colony PCR and Sanger sequencing.

Using this protocol, the reinsertion of deleted regions F2, F4 and F1-F4 (full intergenic) was performed, resulting in strains SA0162, SA0166 and SA0170. It was also used to generate the 42bp-mEGFP construct (SA0228) to measure *FCY1* expression level and all validation clones expressing either *FCY1* or mEGFP from the *FCY1* locus.

Growth assays

Unless stated otherwise, the following protocol was used each time the growth of *FCY1* strains was assessed in media containing 5-FC (Alfa Aesar) or cytosine (Alfa Aesar). Each yeast strain of interest was first inoculated in three replicates in SC [pH=6] liquid media – potentially with HYG, NAT and/or G418, depending on each strain's resistance marker(s) – and grown overnight at 30°C with shaking. After ~16 hours of incubation, a sample (100 µl) of each culture was centrifuged and resuspended in the same volume of sterile water, before being aliquoted and diluted again in sterile water to obtain up to 250 µl of cells at 1 optical density (OD) unit/ml. Growth assays were then assembled in Greiner flat-bottom transparent 96-well plates as follows: 160 µl of media (SC complete for 5-FC or SC-Ura for cytosine); 20 µl of diluted cells [OD=1]; 20 µl of 5-FC or cytosine diluted to obtain the desired final concentration in a volume of 200 µl. Each plate was incubated for 48 hours at 30°C without shaking in a Tecan Infinite M Nano (Tecan Life Sciences) and OD_{595} was measured at 15 minutes intervals.

To obtain exponential growth rates from the resulting growth curves, the complete time-series of OD_{595} was first log-transformed, after adjustment of values to a volume of 1 ml. Then, the slope between each consecutive OD_{595} value was computed, scaled according to the time interval of 0.25 h separating each measurement. For each unique growth curve, the median of the ten largest slopes was used as growth rate for the corresponding culture. When converting the growth rate of any strain i into selection coefficient s_i, the following formula was used [55]:

$$s_i = \frac{\mu_i - \mu_{WT}}{\mu_{WT}} \log 2$$

Variables μ_i and μ_{WT} are the growth rates for the strain of interest and the corresponding WT reference, respectively. This comparison was always made using WT measurements obtained in the same experiment as the growth rate of strain i.

Construction of the mutant libraries

Amplification of mutated F3 and F4 promoter fragments

All single-nucleotide substitutions, insertions and deletions in the F3 and F4 regions were ordered as six separate oPools Oligo Pools (Integrated DNA Technologies Inc.) containing sequences of 117 bp, 118 bp and 116 bp, respectively. Within the different pools, each oligonucleotide is a variant of the full-length F3 or F4 region (77 bp) with flanking 20 bp homology regions at both ends. The latter sequences are respectively the 20 bp immediately upstream or downstream of the corresponding fragment in the reference genome of *S. cerevisiae* S288C, as retrieved from the Saccharomyces Genome Database [86]. For substitutions, each position of each fragment was iteratively degenerated into all four possible nucleotides by specifying a N base, while for insertions, the same mixed base was inserted in 3' of each position in the reference F3 and F4 sequences. When such a N is encountered, an equal ratio of each of the four standard nucleotides (A, T, G, C) is dispensed during synthesis. For the two pools of deletions, single-nucleotide removal was performed iteratively at each position of F3 and F4 and duplicates (occurring whenever two or more consecutive nucleotides are identical in the reference) were removed to obtain non-redundant sets. All sequences ordered as part of the six oPools are listed in **Supplementary Table 5**.

Upon reception, each oPool was spun down and resuspended in fresh PCR-grade water, to obtain a final concentration of 50 µM. Aliquots of 100 µl at 0.05 µM were then obtained by serial dilution from 2 µl of the initial resuspension. From these aliquots, PCR amplification of each oPool was performed in four replicates using KAPA HiFi HotStart DNA polymerase (Roche Molecular Systems) and the following

691   cycle: 98°C for 3 min; {98°C for 30 s; 60°C for 15 s; 72°C for 10 s} x 22; 72°C for 20 s; Hold at 10°C.
692   Each reaction was done in 50 µl:

Volume (µl)	Reagent
33	PCR-grade water
10	KAPA HF buffer [5X]
1.5	dNTPs [10 mM]
1.5	Forward oligo [10 µM]
1.5	Reverse oligo [10 µM]
1.0	KAPA Hi-Fi HotStart DNA polymerase
1.5*	oPool [0.05 µM] (or PCR-grade water)
50	**Total**

In each case, the diluted oPool resuspension was not added to the reaction mix and was instead added
individually in each reaction tube. In addition, one control without the oPool dilution was performed
for each primer combination. Both primer pairs were designed to extend the 20 bp homology regions
included in the oPool, such that mutated F3 and F4 featured 60 bp homology arms upstream and
downstream of the corresponding genomic region. The two pairs were CLOP299-C3 / CLOP299-C4
and CLOP299-D3 / CLOP299-D4, respectively to amplify F3 and F4 oPools.
699
Genomic insertion of mutated fragments in *S. cerevisiae*
To insert promoter mutations upstream of *FCY1*, competent yeast cells harboring a *natNT2* deletion of
the F3 or F4 region (strains SA0129 and SA0130) were transformed with the corresponding PCR-
amplified oPools. For each mutation category and each region, four replicate transformations were
performed from the corresponding four PCR reactions described above – for a total of 24
transformations. The protocol described by Ryan *et al.* (2016) [85] was followed (see *CRISPR-Cas9 yeast*
transformation, above). After incubation, G418-positive colonies from each agar plate were collected
separately. Liquid YPD (5 ml) was added to the plate, cells were resuspended by scraping with a glass
rake and the suspension was transferred to a 15 ml Falcon tube. From this suspension, glycerol stocks
(687 µl cells mixed with 313 µl sterile [80%] glycerol) were made for each transformation and stored
at -80°C. Two 500 µl aliquots from each tube were spun down and stored at -20°C after aspiration of
the supernatant. From these samples, mutant libraries were validated by PE150 sequencing on an
Illumina MiSeq instrument (Genomics Analysis Platform [PAG] of Institut de Biologie Intégrative et
des Systèmes, Université Laval, Québec). This approach was also employed to insert the same set of
promoter mutations upstream of the 42bp-mEGFP construct, using strains SA0246 and SA0254. In this
case, three replicate transformations were instead performed per amplified oPool.
716
Selection of coding sequence mutations
Fifteen codon positions of the *FCY1* gene, for which all possible states had previously been generated
by DMS [6], were selected to obtain a diversity similar to that of the complete mutational scanning of the
F3-F4 region (15 positions x 64 codons = 960 mutants). Among these positions, six were chosen for
their wide range of variant effects and nine others were selected randomly, to fully cover the previously
reported range of mutational effects. The 15 corresponding pools of mutants were each grown for ~24
hours in 4 ml SC complete media (30°C with shaking), from 25 µl of glycerol stock. Optical densities

were measured and all cultures were mixed in equal amounts to make four replicate 1 ml glycerol stocks
(687 μl mixed cultures and 313 μl sterile [80%] glycerol) of this pool of Fcy1p mutants, which were
stored at -80°C.
727
Pooling of mutants
The different sets of promoter mutations (substitutions, insertions and deletions, separately for F3 and
F4) were grown separately in 4 ml SC complete media, from 50-100 μl of the corresponding glycerol
stock. After ~16 hours of incubation at 30°C with shaking, OD_{595} was measured for all cultures.
Subpools were then mixed to obtain equal proportions of all single mutants, according to their OD_{595}
and how many unique genotypes they each contained.
734
For mutations in the *FCY1* background, the four replicate pools of coding sequence mutations were also
grown from 75 μl of glycerol stock. All subpools were then combined as described, according to their
replicate number (i.e. replicate 1 for F3 substitutions was mixed with replicate 1 of all other subpools).
Each subpool of promoter mutations was thus included in only two final pools: one with added coding
sequence mutations and the other without. For mutations in the 42bp-mEGFP background, all subpools
of promoter variants were combined as described without the addition of other mutants, except that
there were only three replicates of each.
742

Bulk competition experiment in 5-FC media
Each mutant pool was grown in two replicates in 4 ml SC complete media (30°C, 200 rpm) for ~20
hours, from 111 μl or 65 μl of the corresponding glycerol stock. These inoculum volumes were chosen
to have at least 2000x cells for each mutant, respectively for pools with and without added coding
sequence mutations. The competition experiment was initiated by diluting each preculture at final
OD_{595} of 0.01 in 25 ml of SC complete media (with and without 5-FC [12 μM]) in 50 ml Falcon tubes.
T0 samples were also prepared by centrifuging two microtubes containing 5 OD_{595} units for each
preculture (850 g for 5 minutes; 4°C). After aspiration of supernatant, cell pellets were stored at -20°C.
Newly inoculated cultures were incubated (30°C with shaking) in the dark until OD_{595} reached ~1.0.
At this first timepoint T1, each culture was diluted in at final OD_{595} of 0.01 in 25 ml of fresh media.
Additional cell pellets were prepared from the remaining cultures. The full volume was first centrifuged
at 4°C (1000 g; 5 minutes). Cell pellets were resuspended in ~5 ml of supernatants and two samples of
~5 OD_{595} units were prepared as previously. This process was repeated at timepoint T2 when the new
cultures reached OD_{595} ~1.0 and final T3 samples were prepared when the newly inoculated T2-T3
cultures reached the same OD_{595}. The timings of each timepoint and the corresponding OD_{595} values
for all mutant pools are listed in **Supplementary Table 6**.
759

Sort-seq experiment
One final pool of F3-F4 mutants in the 42bp-mEGFP background was grown overnight from 100 μl
glycerol stock in 5 ml of SC complete [pH = 6.0] media, at 30°C with shaking. After measurement of
its OD₅₉₅, the culture was diluted to 0.15 OD/ml in 20 ml of fresh SC complete [pH = 6.0] and incubated
for 5 hours at 30°C with shaking, to reach exponential phase (~0.5 OD/mL). From this, 4.5 ml were
transferred to a 5 ml polystyrene round-bottom Falcon tube. The remaining 15.5 ml were centrifuged
to collect cells, so that genotype frequencies before sorting could also be measured by sequencing.
767
A FACSMelody cell-sorter (BD Biosciences) was used to sort yeast cells based on their fluorescence
intensity. Five bins were defined along the distribution of log_{10} fluorescence intensities, numbered 0 to
4 from the lowest to the highest fluorescence levels. More cells were collected from the most populated

parts of the distribution (50 000 cells each from bins 2 and 3, versus 10 000 cells each from bins 0,1
and 4). Each bin was sorted twice, resulting in two technical replicates, in a 96 well plate. The resulting
fractions were each used to inoculate 3 ml of SC complete [pH = 6.0] media in a 24 deep-well plate and
grown for 24h at 30°C with shaking (400 rpm). Each culture was spun down at 3000 rpm (1000 g) for
5 min and all pellets were washed with sterile water and transferred to a 1.5 ml microtube. All
microtubes were centrifuged at 15 000 rpm (21 000 g) for 1 min, supernatants were removed and pellets
were frozen at -80°C. This process was repeated two times for the two other biological replicates, such
that six replicate samples were obtained for each bin.

779
DNA library preparation and sequencing
The same approach was used for all sequencing libraries, for both the bulk competition and sort-seq
experiments. Genomic DNA was first extracted from each cell pellet using a standard phenol-
chloroform extraction protocol [87], including RNAse treatment. For the resequencing of bulk competition
samples in PE300, an additional RNAse treatment (10 minutes, 37°C) was performed on previously
purified DNA samples, followed by precipitation with sodium acetate and cold ethanol. A first PCR
was then performed using Q5 High-Fidelity DNA Polymerase (New England Biolabs) to amplify the
genotyping region chosen for each type of samples, which included the F3-F4 promoter regions and
flanking sequences of varying lengths. This PCR step added Illumina primer-binding sites at both ends,
separated from the fragment of interest by small numbers of N bases (between 2 and 8; 10 total per PCR
amplicon). For the initial (PE150) sequencing of bulk competition samples, primers CLOP311-C6 to
CLOP314-H9 were used (as ordered in **Supplementary Table 2**). Two reactions were made from each
DNA extraction from cultures with added coding sequence mutants. One was specific to the F3-F4 pool
while the other targeted only the added *FCY1* variants. This was possible because the pool of coding
sequence mutations was generated from a codon-optimized sequence, harboring many nucleotide
changes from the WT [6]. For the PE300 resequencing of the same samples, only F3-F4 PCR were
performed, using oligos CLOP328-A6 to CLOP328-H7. Finally, for the sequencing of sort-seq bins,
also in PE300, oligos CLOP328-A6 to CLOP341-A10 were employed. In this case, 7 N bases were
added per amplicon.

799
In all cases, amplicons from this first PCR step were purified with AMPure XP magnetic beads
(Beckman Coulter Life Sciences) and diluted to 0.01 ng/µl. These dilutions were used as template for
a second set of PCR reactions using Q5 polymerase, which added Nextera barcodes and Illumina
adapters i5 and i7. A unique combination of i5 and i7 indexes was used for each DNA sample. The final
PCR products were also purified with AMPure XP beads, before quantification using a NanoDrop
spectrophotometer (Thermo Fisher Scientific). Amplicons from the same sequencing round (PE150
competition, PE300 competition or PE300 sort-seq) were mixed in equal proportions based on the
NanoDrop quantification and the resulting pool was sent for sequencing. PE150 sequencing was
performed on a NovaSeq Illumina instrument (sequencing platform of Centre de recherche du CHU de
Québec-Université Laval, Québec), while PE300 sequencing was performed on an AVITI instrument
from Element Biosciences (Genomics Analysis Platform [PAG] of Institut de Biologie Intégrative et
des Systèmes, Université Laval, Québec).

812
Analysis of sequencing reads
Following each sequencing run, reads were received already demultiplexed, according to the sample id
associated with each combination of i5 and i7 adaptors used. Read quality was first assessed using
FastQC (v 0.12.1) and MultiQC (v 1.12) [88]. Paired-end reads were next merged using PANDAseq (v
2.11) [89], with the '-k' parameter set to 4. The N positions which had been added at both ends during the

first PCR step of library construction were trimmed using command "fastx_filter" of VSEARCH (v
2.28.1) [90], according to a table listing the number of N added for each sample. The first and last 20 bp
of the resulting sequences, which were the binding sites for the oligonucleotide primers used in the first
PCR step, were also trimmed using the same command. Following these trimming steps, all identical
sequences were aggregated with the "derep_fulllength" command of VSEARCH. Each unique sequence
was aligned to the corresponding reference (depending on the type of sample and the length of the
sequenced region) using the needle aligner from the EMBOSS suite (v 6.5.7) [91]. For promoter (F3-F4)
libraries, the opening and extension of gaps was equally penalized (4,4), while penalties of 50 and 0.5
were respectively used for opening and extension for *FCY1* coding sequence libraries [6]. Custom Python
functions were then used to call variants from each alignment. In addition to WT reads, only sequences
harboring single-nucleotide mutations (F3-F4 libraries) or single-codon substitutions (*FCY1* libraries)
were kept for downstream analysis. These variants were further filtered by length, to remove those that
differed too much from the expected length of the PCR fragment. For bulk competition samples, read
counts of each single mutant were converted to WT-relative frequencies. These read counts were instead
used as is in the downstream analysis of sort-seq samples.

Estimation of fitness effects

Calculation of selection coefficients

For each bulk competition culture, time series of WT-relative genotype frequencies were reconstituted
from variant frequencies obtained at the four timepoints (T0 to T3). The number of generations
associated with each timepoint was obtained from the corresponding durations in hours, which varied
between cultures, and the doubling time of the WT strain in the same conditions (SC media with or
without 5-FC [12 μM]), as estimated from growth curve experiments. For each single-mutant genotype,
the selection coefficient s was obtained by fitting a linear model using the number of generations and
the log-transformed WT-relative frequencies as dependent and independent variables, respectively. This
was performed through ordinary least squares regression, using the "ols" implementation of the
statsmodels (v 0.12.2) [92] Python module, with an intercept added to the model. Regressions were
performed iteratively over all time intervals which included the T0 (T0-T1, T0-T2, T0-T3; with
intermediate data points when applicable). Because selection coefficients were inferred separately
within each culture replicate, as many replicate values as there were cultures were obtained for each of
the intervals mentioned above.

To identify the selection coefficients which should be used in further analysis, two diagnostics were
performed. First, the agreement between the initial (T0) log-scaled WT-relative frequencies and the
fitted intercept from each regression model was assessed. Second, the linearity of log-scaled frequencies
through time, as expected for cultures growing exponentially, was validated graphically. This revealed
that no interval was reliable for the initial 150 paired-end genotyping of promoter (F3-F4) libraries,
while T0-T2 selection coefficients should be used for the 300 paired-end resequencing of the promoter
fragments and the 150 paired-end sequence of *FCY1* coding sequence (see Supplementary Note). The
chosen selection coefficients were then used to identify a T0 abundance threshold for variant filtering,
such that inferred s would not be correlated with initial read counts. This threshold was set at 85 reads,
both for the promoter (300 paired-end) and coding sequence (150 paired-end) sequencing libraries. To
ensure that low frequency but beneficial secondary mutations could be assessed, any variant which was
below the threshold at T0 but over it at the end of the experiment was kept.

Removal of outliers and aggregation of replicates

In 5-FC samples, mutations sometimes appeared beneficial in only one pair of cultures (coming from the same transformation replicate), likely due to secondary resistance-conferring mutations (See Supplementary Note). Outlier removal was thus performed, separately for two types of libraries (promoter or coding sequence). The promoter (F3-F4) libraries were further separated depending on whether coding-sequence mutants had been added in the corresponding cultures. Within each group of samples, the median absolute deviation (MAD) of selection coefficients was first computed for each mutant. This was then used to compute robust z-scores for each replicate measurement within each genotype. The resulting distribution of robust z-scores calculated across all mutant genotypes was compared and its 95th percentile was defined as the threshold to identify outlier replicates. The filtering of outliers was however restricted to a subset of mutants which displayed a higher variance, to ensure that small absolute variations between replicates were not identified as outliers. This was assessed from each genotype's standard deviation, with the 95th percentile of the overall distribution used as threshold. For F3-F4 sequencing libraries obtained from cultures with added coding sequence mutations, the 85th percentile was instead used.

To obtain final estimates of the fitness effects of promoter mutations, only F3-F4 libraries without added coding sequence mutants were taken into account. Four of the eight replicate cultures, corresponding to two of the four transformation replicates, were additionally ignored. In these four samples, selection coefficients displayed a temporal shift towards negative values which appeared consistent with some undetected resistance-conferring mutations being genotyped as WT (see Supplementary Note). For each of the 1020 assayed promoter mutations, the corresponding fitness effect was estimated as the median of the selection coefficients observed across the four remaining replicates, which were considered unbiased. The statistical significance of selection coefficients was assessed from the same four replicate measurements for each mutation, by comparing their mean to 0 using the SciPy (v 1.15.2) [93] implementation of the one-sample t-test, followed by Benjamini-Hochberg correction controlling the false discovery rate at 5%. For the coding sequence mutations added as controls, fitness effects were instead estimated as the median of the eight replicate measurements, without ignoring any of the corresponding samples.

Comparison with published effects of amino acid substitutions in Fcy1p

The selection coefficients of added coding sequence mutations were first aggregated into grand medians, according to the corresponding amino acid substitutions. They were then compared with previously published variant effect scores obtained in the same conditions (12 μM 5-FC) [6]. The agreement between the two experiments was further assessed by classifying variants as resistant and susceptible in both cases. To this end, Gaussian mixture models with two components and a "tied" covariance matrix were fitted on each of the two datasets, using the scikit-learn (v 1.6.1) [94] implementation.

For the comparison of promoter and coding sequence mutations, the full dataset from [6] was used. All single-nucleotide substitutions in the reference sequence of the *FCY1* gene, obtained from the Saccharomyces Genome Database [86], were first performed computationally. The corresponding variant effects were retrieved from the Després *et al.* (2022) data [6], according to the resulting amino acid substitutions. Of 1431 possible single-nucleotide substitutions in *FCY1*, only 20 were not covered, mostly in the first and last codons. The Gaussian mixture model previously fitted on the full set of amino acid changes was then used to classify all single mutants as either resistant or susceptible. From this

911 classification, the fraction of adaptive (resistance-conferring) mutations in the coding sequence was
912 obtained.
913

Calculation of sort-seq expression scores

Within each of the six replicate sort-seq experiments, all variants which reached a total abundance of at
least 85 reads across the five sampled bins were kept for downstream analysis. Although this included
a small number of secondary mutations (outside of the F3-F4 mutagenized region), any variant which
occurred in the mEGFP sequence was filtered out. For each of the remaining genotypes i, which includes
the WT, fluorescence intensity was quantified from a weighted average B_i of bins where the
corresponding reads were detected:

$$B_i = \frac{\sum_{j=0}^{4} j \cdot \frac{r_{i,j}}{r_{tot,j}} \frac{n_j}{n_{tot}}}{\sum_{j=0}^{4} \frac{r_{i,j}}{r_{tot,j}} \frac{n_j}{n_{tot}}}$$

921
922 In the above formula, j is the bin number, where 0 and 4 are associated with the lowest and highest
923 fluorescence intensities, respectively. Variables $r_{i,j}$ and $r_{tot,j}$ are the read count for genotype i, as well
924 as the total number of reads, in bin j. Finally, n_j and n_{tot} are respectively the number of cells collected
925 from bin j and the total amount of cells collected across the five bins. The n_j/n_{tot} ratio was introduced
926 because the number of yeast cells collected was not uniform across all bins.
927

928 Separately for each replicate experiment, the difference was computed between the B_m weighted
929 average of each mutant m and the B_{WT} weighted average obtained for the WT genotype:

$$\Delta_{Ref,m} = B_m - B_{WT}$$

930
931 For each mutant, up to six replicate Δ_{Ref} scores were obtained (one for each sort-seq experiment). The
932 median of these values was used as an estimate of each mutation's effect on fluorescence intensity.
933 Positive values denote an increase relative to the WT, while negative values instead indicate a decrease.
934

Selection and construction of validation mutants

The sort-seq data was used to select 30 mutant genotypes for validation. These mutations were chosen
to cover the whole range of scores, as well as all regions of the promoter where single-nucleotide
changes appeared impactful. Substitutions were favored to ensure a high success rate to the downstream
reconstruction of genotypes. To ensure that low-confidence mutations were not selected, a one-sample
t-test (with Benjamini-Hochberg correction at FDR=5%) was first performed on the replicate sort-seq
scores to identify Δ_{Ref} values which were significantly different from 0. This was also repeated with
Δ_{mod} scores computed from B_{mod} weighted averages obtained from reads of bin 1 to 4, excluding the
0th one displaying the lowest fluorescence intensities. Because the Δ_{Ref} scores appeared biased
towards positive values compared with Δ_{mod}, the latter were used to select significant mutations to be
reconstructed, except when annotated *. This ensured a more stringent selection of genotypes displaying
altered fluorescence intensity compared with the WT reference. Within each of the mutant groups
described below, the specified number of sequences were chosen randomly, unless stated otherwise.
Among mutations which significantly decreased Δ_{Mod} below zero, there were seven groups: 1) indels
before the second methionine of *FCY1* (n=2); 2) indels after the second methionine of *FCY1* (n=2)*; 3)
top 3 lowest scores between positions -50 and 0 of the promoter*; 4) substitutions between positions -
133 and -127 (n=6); 5) substitutions between position -45 and -11 (n=2); 6) substitutions and insertions
near the -75 position (one each) and 7) two chosen substitutions outside of regions with many significant
effects. Among mutations which significantly increased Δ_{Mod} above zero, a similar approach was
employed. The four categories were: 1) substitutions between position -125 and -118 (n=4); 2)

substitutions between positions -112 to -109 (n=2); 3) substitutions between positions -63 to -57 (n=2)
and 4) variants among the 25 highest scores (n=4).
957
The corresponding validation mutants were individually reconstructed using the CRISPR-Cas9 yeast
transformation protocol described above. For each genotype, this was performed in two genetic
backgrounds: fluorescent strain SA0247 (ΔF3-F4 with 42bp-mEGFP construct) and *FCY1* deletion
(ΔF3-F4) mutant SA0159 (see **Supplementary Table 1**). A fusion PCR product harboring the single
mutation of interest was used as donor DNA. This amplicon was generated by two consecutive rounds
of PCR. In the first round, two reactions were performed with WT genomic DNA as template. The first
one used a generic forward primer hybridizing upstream of F3 (CLOP326-A11) with a mutant-specific
reverse primer harboring the mutation of interest (**Supplementary Table 2**). In the second reaction, a
forward primer with the mutation of interest (**Supplementary Table 2**) was used alongside a generic
reverse primer binding either in the mEGFP sequence (CLOP326-G10; for background SA0247) or in
the *FCY1* coding sequence (CLOP326-E10; for background SA0159). The two mutant-specific
mutagenesis primers are complementary to each other, such that there is homology between the two
amplicons. The second PCR round used the two PCR products from the first round diluted 1/40 as
template and the same two generic primers used in the first pair of PCR (CLOP326-A11 with either
CLOP326-G10 or CLOP326-E10, depending on the mutagenized background), which allowed fusion
of both PCR parts. This approach thus allows to amplify the full F3-F4 promoter region (with a short
segment of *FCY1*) while introducing a single mutation within it and adding homology regions on both
ends for subsequent genomic insertion. It was additionally used to perform deletion of short segments
of the promoter (-133 to -127 and -125 to -118), similarly in both genetic backgrounds. After
transformation of each fusion PCR product, clones from each transformation were sent for Sanger
sequencing to validate proper genomic integration of the mutation. All 66 resulting strains, including
reconstructions of the WT promoter, are listed in **Supplementary Table 1**.

Expression measurements by flow cytometry
Measurement of promoter activity using fluorescent strains was performed with a Guava EasyCyte HT
cytometer (CytekBio). Each strain of interest was grown overnight in three replicates in SC complete
media at 30°C with shaking, along with similar replicate cultures of a reference strain with the WT
promoter (SA0247 or AKD1462) and of a non-fluorescent control (AKD1123 or AKD1429). The
overnight cultures were diluted at 0.15 OD_{595} in fresh SC-Trp (MSG) media and grown at 30°C with
shaking until ~0.5 OD_{595}. Exponentially growing cells were then diluted to 0.05 OD_{595} in fresh media
and single-cell fluorescence intensity measurements were performed with the Guava EasyCyte HT
system. For each replicate culture of each strain, 5000 events were acquired (488 nm excitation; 525
nm detection). The medians of the corresponding distributions of fluorescence intensities were used to
estimate relative expression levels.
991
Within each cytometry experiment, events were first filtered out according to forward scatter and side
scatter values, with the 99th percentile of each distribution used as thresholds. All fluorescence
intensities were then normalized by the corresponding forward scatter measurement. From these
normalized fluorescence intensities, a median was computed for each replicate culture assayed. These
medians were then normalized by the grand median of measurements made for the reference strain,
such that they were converted into WT-relative fluorescence intensities. The resulting values, and
especially their medians, were used as estimates of relative expression levels. Accordingly, three
replicate measurements of WT-relative fluorescence intensity were obtained for each tested strain, with
a value of 1 being equal to the WT level.
1001

Calibration of sort-seq scores and bootstrapping of relative expression levels

WT-relative fluorescence intensities were first measured for all validation mutants reconstructed in the SA0247 (42bp-mEGFP) background, as described above. These measurements were then compared with the corresponding sort-seq expression scores and a LOWESS regression was fitted. This approach allowed to simultaneously validate the sort-seq experiments and convert the resulting scores into more interpretable WT-relative fluorescence intensities. The Δ_{Ref} scores were used, since some genotypes with very low fluorescence intensities were absent from the Δ_{mod} dataset. In addition, because fluorescence intensity measurements from reconstructed mutants were used to fit the relationship, any bias in Δ_{Ref} was unlikely to affect the downstream inference of relative expression levels. So that uncertainties could also be obtained for each of the >1000 mutant expression levels, both the LOWESS fitting and the conversion of sort-seq scores into relative fluorescence intensities were performed within a bootstrapping routine.

At the start of each iteration, Δ_{Ref} scores were first resampled with replacement. For each genotype which had been included in the sort-seq analysis, as many values were sampled as there were replicate measurements among the six experiments. Per-genotype medians were then computed on the newly sampled data. For each reconstructed mutant for which fluorescence had been measured directly, three median WT-relative fluorescence intensities were sampled with replacement among the corresponding replicate measurements. Grand medians were computed from each trio of observations, which were combined with the newly recomputed median Δ_{Ref} scores of the corresponding validation genotypes to obtain 30 pairs. From these data points, LOWESS regression was performed with median Δ_{Ref} as independent variable, using the "nonparametric.lowess" implementation from statsmodels (v 0.14.4) [92] with default parameters. This regression was then evaluated at each of the >1000 median Δ_{Ref} values obtained from the first resampling steps. This generated an estimate of WT-relative fluorescence intensity for each of the genotypes which had been assayed in the sort-seq screen. The regression was also evaluated at 100 equally spaced values from the minimal to the maximal median Δ_{Ref} observed experimentally. This process was repeated 5000 times. After all iterations had been completed, medians and corresponding 95% prediction intervals were computed from the distribution of WT-relative fluorescence intensities obtained for each mutant genotype. These medians were used as WT-relative expression levels associated with the corresponding F3-F4 mutations. The same calculations were performed for each of the 100 equally spaced values at which the LOWESS had been evaluated, thus allowing to plot the relationship between Δ_{Ref} and relative fluorescence intensity.

Experimental measurements of the expression-fitness function

Construction of strains with distinct *FCY1* expression levels

Six yeast promoters covering a wide range of expression levels were used to construct two sets of strains, respectively from *FCY1* background AKD1123 and from fluorescent background SA0228. The six promoter cassettes, each including a G418 resistance marker, were PCR-amplified from the corresponding plasmids [84] (**Supplementary Table 3**) using oligonucleotides CLOP338-G6 and CLOP338-H6 (**Supplementary Table 2**). The two primers provided homology for genomic insertion immediately upstream of *FCY1* (or of its first 42 bp for the SA0228 background), between the gene and its native promoter. Yeast transformations were performed in the two genetic backgrounds as previously described, using each of the corresponding PCR products. Transformants were selected on YPD+HYG+G418 media. Correct integration of the promoter cassette was validated by PCR in four clones from each transformation, using oligonucleotides CLOP310-D1 and CLOP291-A3 (**Supplementary Table 2**). Two to three validated clones were obtained in most cases, such that

1048 biological replicates could be used in downstream assays for the majority of genotypes. All 25
1049 corresponding yeast strains are listed in **Supplementary Table 1**.

Measurements of growth rates and fluorescence intensities
The set of *FCY1* strains (background AKD1123) was used to measure growth rates in liquid media
containing different concentrations of 5-FC (3 μM, 12 μM, 24 μM and 48 μM) or cytosine (47 μM, 63
μM, 84 μM and 105 μM), following the growth assays protocol described above. The set of fluorescent
strains (background SA0228) was instead used in flow cytometry experiments to measure expression
levels (see Expression measurements by flow cytometry). In both cases, all available biological
replicates were used for each genotype. When less than three strains were available, three replicate
cultures were nonetheless inoculated.

The growth assays in 5-FC media were performed in two separate experiments. More than three
replicate measurements were thus available for two concentrations (12 μM and 3 μM). Growth rates
from the first experiment were adjusted to have the same grand medians as those of the second one, by
the addition of the difference between the two experiments' grand medians. This was performed
separately for each concentration and allowed to combine the two sets of growth rates, in spite of
potential systematic differences between them – for instance if the true concentration of 5-FC in the
media varied. The cytometry measurements, which were done in parallel with the growth assays, were
also performed in two separate experiments. In this case, potential systematic differences were
controlled for by computing WT-relative fluorescence intensities within each experiment and only
combining measurements after this step.

Fitting of expression-fitness functions
Fluorescence measurements associated with each promoter were combined with the corresponding
growth rates. An exponential function was fitted for each of the eight conditions, using WT-relative
fluorescence as the independent variable and growth rate as the dependent variable. This could model
the sharp decline observed in 5-FC as well as the steep increase seen in cytosine conditions. The fitting
of all expression-fitness curves was performed within a bootstrapping routine. For each promoter, two
WT-relative fluorescence intensity measurements were first sampled with replacement from each
cytometry experiment. Then, the four resulting values were resampled with replacement. Corresponding
growth rates were sampled similarly. If a given condition had been tested in two experiments, two
replicate growth rates were sampled with replacement from each experiment for each promoter, and the
four measurements were afterwards resampled with replacement. This approach was employed to avoid
favoring one experiment over the other, since the number of replicates sometimes varied between them.
When only one experiment had been performed, replicate growth rates from this assay were instead
resampled with replacement. Once relative fluorescence intensities and growth rates had been sampled,
values obtained for the same promoter were paired randomly into data points. The exponential function
was then fitted using the "optimize.curve_fit" implementation of the SciPy module (v 1.15.2) [93] module
and evaluated over 250 equally spaced values covering the full range of WT-relative fluorescence
intensities. After 5000 iterations, medians and corresponding 95% prediction intervals were computed
for each of these 250 fluorescence levels, thus generating the expression-fitness function. This was
repeated for each of the 5-FC and cytosine concentrations tested.

Validation of *FCY1* promoter variants fitness effects
1093 The previously described validation mutants reconstructed in the SA0159 *FCY1* background
1094 (**Supplementary Table 1**) were grown in SC complete (MSG) media with 12 μM 5-FC and without,

according to the growth assays protocol described above. One genotype (AKD1417; C to G substitution at position -123) displayed both an elevated fitness in SC media and an outlier low growth rate in 5-FC conditions and was not included in downstream analyses. For all other mutants, growth rates measured in 12 μM 5-FC were converted into selection coefficients as described in the growth assays section, using growth rates measured in the same experiment for WT strain AKD1123. The expression-fitness function obtained in the same conditions and the bounds of its 95% prediction interval were also converted into selection coefficients using the same approach. In this case, the WT growth rate interpolated from the curve was instead used as comparison.

Assessment of 5-FC resistance arising from mutant library construction

To test whether mutations conferring 5-FC resistance were introduced during mutant library construction, the process described in *Construction of mutant libraries* was repeated, but using WT sequences of the F3 and F4 regions. Briefly, the same two oligonucleotide pairs (CLOP299-C3/CLOP299-C4 and CLOP299-D3/CLOP299-D4) were first used to amplify F3 and F4 from genomic DNA of the AKD1123 strain, using KAPA HiFi HotStart DNA polymerase (Roche Molecular Systems). Competent cells from strains SA0129 and SA0130 (prepared as described by [85]) were then transformed with the F3 and F4 inserts, respectively. Two biological replicates of each transformation were made. This was performed following the CRISPR-Cas9 approach described above for the pools of single mutations. After 48 hours of incubation at 30°C, transformants from each Petri dish were collected in 5 ml YPD by scraping, as previously described. From each of the resulting cell suspension, cultures were started in 5 ml of SC complete [pH=6] + HYG, from 100 μl of cells.

After overnight incubation at 30°C with shaking, OD_{595} was measured for each culture and dilutions at $OD_{595} = 1$ were made in 200 μl of sterile water. Additional dilutions were afterwards made for OD_{595} values from 0.2 to 0.002. From each dilution, 150 μl were plated on SC complete (MSG) solid media with and without 5-FC [100 μg/ml]. Petri dishes were incubated at 30°C for 48 hours and colonies were counted. While colonies were too numerous to be counted on SC complete controls, some dilutions plated on 5-FC media resulted in countable colony numbers (between 20 and 340). For each plating (one per CRISPR-Cas9 transformation), colony counting was performed for two dilution levels, which acted as technical replicates. Since the SA0130 background was initially resistant to 5-FC due to the insertion of a natNT2 in place of F4, all associated platings were replica plated on YPD+NAT media. Nourseothricin-resistant colonies were subtracted from the total number of 5-FC-resistant colonies. The frequency of 5-FC resistance was then estimated as follows, separately for each plating on media containing 5-FC:

$$F_{LOF} = \frac{n_{FC}}{OD_d \, V_{plate} \, c_{ml}}$$

In the equation above, n_{FC} is the number of colonies on 5-FC media (after subtracting NAT resistant ones, when applicable), while OD_d is the OD_{595} of the plated dilution, V_{plate} is the volume deposited on each plate (0.15 ml) and c_{ml} is the number of yeast cells in 1 ml of media at $OD_{595} = 1$, for which a value of 10 million cells/ml was used. The resulting F_{LOF} was considered as a putative frequency of *FCY1* LOF in the corresponding transformations. To validate whether that was the case, 16 colonies were selected for Sanger sequencing, using primer CLOP326-F10, which covered nearly 70% of the *FCY1* coding sequence **(Supplementary Table 2)**. All colony counts, resistance rates and relevant calculations are listed in **Supplementary Table 7**.

Simulations with *invisible* resistance-conferring mutations

A simulation framework replicating the bulk competition experiment was designed. It involved a library of neutral variants within which a small number of invisible resistance-conferring mutations was

introduced, thus mimicking secondary mutations occurring outside of the genotyping region in the experiment. A set of variants of interest (hereafter named genotype labels) was first generated by sampling 1000 neutral mutations ($s \sim \mathcal{N}(0, 0.005)$) four times. The standard deviation used was set according to what was observed in sequencing libraries from cultures without 5-FC. Four transformations were then obtained by replicate random samplings of 10 000 clones, among which 17% were defined as WT ($s = 0$) and the remaining were drawn from the set of genotype labels. Within each such transformation replicate, a small fraction of clones was defined as harboring an invisible resistance-conferring mutation (or resistance flag), which increased its s (between 0.4 to 0.6, depending on simulations) without changing its genotype label. From each transformation, two replicate cultures were generated by randomly sampling two sets of 2.5 million cells, resulting in eight cultures, as in the bulk competition experiment (5-FC condition without added coding sequence variants). From each culture, 1 million of sequencing reads (genotype labels) were sampled randomly, to simulate sequencing of the initial T0 samples. Simulated cells in each culture were next aggregated according to genotype labels and presence/absence of a resistance flag, prior to calculation of WT-relative frequencies for each group (called true genotypes). Growth over the course of a bulk competition experiment was simulated by modeling exponential changes of WT-relative frequencies through time for each true genotype i, as follows:

$$F_{i,T+1} = F_{i,T}\, e^{s_i n_{gen}}$$

The WT-relative frequency F of true genotype i at timepoint $T + 1$ is obtained from the corresponding WT-relative frequency at previous timepoint T according to the genotype's true selection coefficient (increased by a resistance-conferring mutation when applicable) and the time interval n_{gen} between the two timepoints, in WT generations. Three such timepoints, from T1 to T3, were simulated and their respective durations were set in accordance with the bulk competition experiment. At each timepoint, 1 million reads as well as a new initial population of 2.5 million cells were sampled from each culture, as described above. Selection coefficients for all variants were then computed from time-series of simulated read counts, as described for experimental variant frequencies. This was performed from genotype labels only as well as with knowledge of true genotypes (resistance flags). Such simulations of one full experiment were repeated while varying both the initial frequency and the fitness effect of resistance-conferring mutations. For each combination, multiple replicate simulated experiments (each with eight simulated cultures) were performed.

Data availability

All sequencing data from bulk competition and sort-seq experiments has been deposited on the NCBI SRA (https://www.ncbi.nlm.nih.gov/sra/PRJNA1321317). All other experimental data and files needed to replicate the analysis have been deposited on Zenodo: https://doi.org/10.5281/zenodo.17088129.

Code availability

All analyses were performed using custom Python scripts and notebooks, which are available at: https://github.com/Landrylab/Aube_et_al_2025_FCY1promoter. A custom Python pipeline was deployed on a computing cluster (Institut de Biologie Intégrative et des Systèmes, Université Laval) to compute variant counts and selection coefficients from raw sequencing data. As described above, this used the following software: FastQC (v 0.12.1), MultiQC (v 1.12) [88], PANDAseq (v 2.11) [89], VSEARCH (v 2.28.1) [90] and EMBOSS suite (v 6.5.7) [91]. The versions of

all Python modules used are included as requirements files, separately for the analysis pipeline and for the notebooks, which were executed locally.

Acknowledgements

We thank Philippe Després, Isabelle Hatin, Olivier Namy and all Landry lab members, especially Soham Dibyachintan, David Jordan, Pascale Lemieux and Camille Bédard, for helpful discussions. We are grateful to Pavithra Venkataraman, Angel Cisneros, Soham Dibyachintan and Philippe Després, who provided comments on this manuscript. We additionally thank Angel Cisneros for helpful suggestions on the simulation framework. This work was supported by a grant from the Natural Sciences and Engineering Research Council of Canada (NSERC RGPIN-2020-04844) for CRL. CRL holds the Canada Research Chair in Cellular Systems and Synthetic Biology. SA was supported by doctoral research fellowships from Natural Sciences and Engineering Research Council of Canada and Fonds de recherche du Québec - Nature et technologies. The F65V-34a-mEGFP plasmid used in this study was a kind gift from Dr Michael Springer.

Author contributions

CRL, AKD and SA designed the study. SA and AKD performed the experiments as well as the data analysis. SA, AKD and CRL interpreted the results. SA, AKD and CRL wrote the manuscript.

References

1.   Wright, S. The Roles of Mutation, Inbreeding, crossbreeding and Selection in Evolution. *Proceedings of the XI International Congress of Genetics* **8**, 209–222 (1932).

2.   Smith, J. M. Natural selection and the concept of a protein space. *Nature* **225**, 563–564 (1970).

3.   Weinreich, D. M., Delaney, N. F., Depristo, M. A. & Hartl, D. L. Darwinian evolution can follow only very few mutational paths to fitter proteins. *Science* **312**, 111–114 (2006).

4.   Aguilar-Rodríguez, J., Payne, J. L. & Wagner, A. A thousand empirical adaptive landscapes and their navigability. *Nat. Ecol. Evol.* **1**, 45 (2017).

5.   Fowler, D. M. & Fields, S. Deep mutational scanning: a new style of protein science. *Nat. Methods* **11**, 801–807 (2014).

6.   Després, P. C. *et al.* Asymmetrical dose responses shape the evolutionary trade-off between antifungal resistance and nutrient use. *Nat. Ecol. Evol.* **6**, 1501–1515 (2022).

7.   Bédard, C. *et al.* Most azole resistance mutations in the Candida albicans drug target confer cross-resistance without intrinsic fitness cost. *Nat. Microbiol.* **9**, 3025–3040 (2024).

8.   Rouleau, F. D. *et al.* Deep mutational scanning of Pneumocystis jirovecii dihydrofolate reductase reveals allosteric mechanism of resistance to an antifolate. *PLoS Genet.* **20**, e1011252 (2024).

9.   Jiang, L. *et al.* A Balance between Inhibitor Binding and Substrate Processing Confers

Influenza Drug Resistance. *Journal of molecular biology* **428**, (2016).

10.   Flynn, J., Samant, N., Schneider-Nachum, G., Tenzin, T. & Bolon, D. N. A. Mutational

fitness landscape and drug resistance. *Curr. Opin. Struct. Biol.* **78**, 102525 (2023).

11.   Estevam, G. O. *et al.* Mapping kinase domain resistance mechanisms for the MET receptor

tyrosine kinase via deep mutational scanning. *Elife* **13**, (2025).

12.   Gresham, D. *et al.* The repertoire and dynamics of evolutionary adaptations to controlled

nutrient-limited environments in yeast. *PLoS Genet.* **4**, e1000303 (2008).

13.   Payen, C. *et al.* The dynamics of diverse segmental amplifications in populations of

Saccharomyces cerevisiae adapting to strong selection. *G3 (Bethesda)* **4**, 399–409 (2014).

14.   Rich, M. S. *et al.* Comprehensive Analysis of the SUL1 Promoter of Saccharomyces

cerevisiae. *Genetics* **203**, 191–202 (2016).

15.   Ascencio, D. *et al.* Expression attenuation as a mechanism of robustness against gene

duplication. *Proc. Natl. Acad. Sci. U. S. A.* **118**, e2014345118 (2021).

16.   Siddiq, M. A., Duveau, F. & Wittkopp, P. J. Plasticity and environment-specific relationships

between gene expression and fitness in Saccharomyces cerevisiae. *Nat. Ecol. Evol.* **8**, 2184–2194

(2024).

17.   Siu, L. K., Ho, P. L., Yuen, K. Y., Wong, S. S. & Chau, P. Y. Transferable hyperproduction

of TEM-1 beta-lactamase in Shigella flexneri due to a point mutation in the pribnow box. *Antimicrob.*

Agents Chemother. **41**, 468–470 (1997).

18.   Jay, A., Jordan, D. F., Gerstein, A. & Landry, C. R. The role of gene copy number variation

in antimicrobial resistance in human fungal pathogens. *NPJ Antimicrob Resist* **3**, 1 (2025).

19.   Dekel, E. & Alon, U. Optimality and evolutionary tuning of the expression level of a protein.

Nature **436**, 588–592 (2005).

20.   Jiang, L., Mishra, P., Hietpas, R. T., Zeldovich, K. B. & Bolon, D. N. A. Latent effects of

Hsp90 mutants revealed at reduced expression levels. *PLoS Genet.* **9**, e1003600 (2013).

21.   Rest, J. S. *et al.* Nonlinear fitness consequences of variation in expression level of a

eukaryotic gene. *Mol. Biol. Evol.* **30**, 448–456 (2013).

22.   Keren, L. *et al.* Massively Parallel Interrogation of the Effects of Gene Expression Levels on

Fitness. *Cell* **166**, 1282–1294.e18 (2016).

23.   Arita, Y. *et al.* A genome-scale yeast library with inducible expression of individual genes.

Mol. Syst. Biol. **17**, e10207 (2021).

24.   Cisneros, A. F. *et al.* Epistasis between promoter activity and coding mutations shapes gene

evolvability. *Sci. Adv.* **9**, eadd9109 (2023).

25.   Tsuru, S., Hatanaka, N. & Furusawa, C. Promoters constrain evolution of expression levels of

essential genes in Escherichia coli. *Mol. Biol. Evol.* **41**, msae185 (2024).

26.   Metzger, B. P. H. *et al.* Contrasting Frequencies and Effects of cis- and trans-Regulatory

Mutations Affecting Gene Expression. *Mol. Biol. Evol.* **33**, 1131–1146 (2016).

27.   Hodgins-Davis, A., Duveau, F., Walker, E. A. & Wittkopp, P. J. Empirical measures of mutational effects define neutral models of regulatory evolution in Saccharomyces cerevisiae. *Proceedings of the National Academy of Sciences* **116**, 21085–21093 (2019).

28.   Vaishnav, E. D. *et al.* The evolution, evolvability and engineering of gene regulatory DNA. *Nature* **603**, 455–463 (2022).

29.   Li, X. C., Fuqua, T., van Breugel, M. E. & Crocker, J. Mutational scans reveal differential evolvability of Drosophila promoters and enhancers. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **378**, 20220054 (2023).

30.   Patwardhan, R. P. *et al.* High-resolution analysis of DNA regulatory elements by synthetic saturation mutagenesis. *Nat. Biotechnol.* **27**, 1173–1175 (2009).

31.   Patwardhan, R. P. *et al.* Massively parallel functional dissection of mammalian enhancers in vivo. *Nat. Biotechnol.* **30**, 265–270 (2012).

32.   Melnikov, A. *et al.* Systematic dissection and optimization of inducible enhancers in human cells using a massively parallel reporter assay. *Nat. Biotechnol.* **30**, 271–277 (2012).

33.   Kircher, M. *et al.* Saturation mutagenesis of twenty disease-associated regulatory elements at single base-pair resolution. *Nat. Commun.* **10**, 3583 (2019).

34.   Duveau, F. *et al.* Fitness effects of altering gene expression noise in Saccharomyces cerevisiae. *Elife* **7**, e37272 (2018).

35.   Ringel, A. E. *et al.* Yeast Tdh3 (glyceraldehyde 3-phosphate dehydrogenase) is a Sir2-interacting factor that regulates transcriptional silencing and rDNA recombination. *PLoS Genet.* **9**, e1003871 (2013).

36.   Kankipati, H. N., Rubio-Texeira, M., Castermans, D., Diallinas, G. & Thevelein, J. M. Sul1 and Sul2 sulfate transceptors signal to protein kinase A upon exit of sulfur starvation. *J. Biol. Chem.* **290**, 10430–10446 (2015).

37.   Vermes, A., Guchelaar, H. J. & Dankert, J. Flucytosine: a review of its pharmacology, clinical indications, pharmacokinetics, toxicity and drug interactions. *J. Antimicrob. Chemother.* **46**, 171–179 (2000).

38.   Lin, Z., Wu, W.-S., Liang, H., Woo, Y. & Li, W.-H. The spatial distribution of cis regulatory elements in yeast promoters and its implications for transcriptional regulation. *BMC Genomics* **11**, 581 (2010).

39.   Lemieux, P., Bradley, D., Dubé, A. K., Dionne, U. & Landry, C. R. Dissection of the role of a Src homology 3 domain in the evolution of binding preference of paralogous proteins. *Genetics* **226**, iyad175 (2024).

40.   Dionne, U. *et al.* Protein context shapes the specificity of SH3 domain-mediated interactions in vivo. *Nat. Commun.* **12**, 1597 (2021).

41.   Matuszewski, S., Hildebrandt, M. E., Ghenu, A.-H., Jensen, J. D. & Bank, C. A statistical

guide to the design of deep mutational scanning experiments. *Genetics* **204**, 77–87 (2016).

42.   de Boer, C. G. *et al.* Deciphering eukaryotic gene-regulatory logic with 100 million random

promoters. *Nat. Biotechnol.* **38**, 56–65 (2020).

43.   Peterman, N. & Levine, E. Sort-seq under the hood: implications of design choices on large-

scale characterization of sequence-function relations. *BMC Genomics* **17**, 206 (2016).

44.   Eisenberg, A. R. *et al.* Translation initiation site profiling reveals widespread synthesis of

non-AUG-initiated protein isoforms in yeast. *Cell Syst.* **11**, 145–160.e5 (2020).

45.   de Boer, C. G. & Hughes, T. R. YeTFaSCo: a database of evaluated yeast transcription factor

sequence specificities. *Nucleic Acids Res.* **40**, D169–79 (2012).

46.   Rhee, H. S. & Pugh, B. F. Genome-wide structure and organization of eukaryotic pre-

initiation complexes. *Nature* **483**, 295–301 (2012).

47.   McMillan, J., Lu, Z., Rodriguez, J. S., Ahn, T.-H. & Lin, Z. YeasTSS: an integrative web

database of yeast transcription start sites. *Database (Oxford)* **2019**, baz048 (2019).

48.   Huang, Q., Szklarczyk, D., Wang, M., Simonovic, M. & von Mering, C. PaxDb 5.0: Curated

protein quantification data suggests adaptive proteome changes in yeasts. *Mol. Cell. Proteomics* **22**,

100640 (2023).

49.   Erb, I. & van Nimwegen, E. Transcription factor binding site positioning in yeast: proximal

promoter motifs characterize TATA-less promoters. *PLoS One* **6**, e24279 (2011).

50.   Pelechano, V., Wei, W. & Steinmetz, L. M. Extensive transcriptional heterogeneity revealed

by isoform profiling. *Nature* **497**, 127–131 (2013).

51.   Struhl, K. Naturally occurring poly(dA-dT) sequences are upstream promoter elements for

constitutive transcription in yeast. *Proc. Natl. Acad. Sci. U. S. A.* **82**, 8419–8423 (1985).

52.   de Boer, C. G. & Hughes, T. R. Poly-dA:dT tracts form an in vivo nucleosomal turnstile.

PLoS One **9**, e110479 (2014).

53.   Ichikawa, Y. *et al.* Sequence-directed nucleosome-depletion is sufficient to activate

transcription from a yeast core promoter in vivo. *Biochem. Biophys. Res. Commun.* **476**, 57–62

(2016).

54.   Jansen, A., van der Zande, E., Meert, W., Fink, G. R. & Verstrepen, K. J. Distal chromatin

structure influences local nucleosome positions and gene expression. *Nucleic Acids Res.* **40**, 3870–

3885 (2012).

55.   Chevin, L.-M. On measuring selection in experimental evolution. *Biol. Lett.* **7**, 210–213

(2011).

56.   Hoekstra, H. E. & Coyne, J. A. The locus of evolution: evo devo and the genetics of

adaptation: The locus of evolution. *Evolution* **61**, 995–1016 (2007).

57.   Delma, F. Z. *et al.* Genetic mutations in Cryptococcus neoformans pyrimidine salvage

pathway enzymes contribute to reduced susceptibility against 5-fluorocytosine. *NPJ Antimicrob.*

Resist. **2**, 24 (2024).

58.   Kruglyak, L. *et al.* Insufficient evidence for non-neutrality of synonymous mutations. *Nature* **616**, E8–E9 (2023).

59.   Wyman, S. K. *et al.* Synthesis errors in oligonucleotides are propagated into the genome by homology-directed repair. *bioRxiv* 2025.07.08.663753 (2025) doi:10.1101/2025.07.08.663753.

60.   Basehoar, A. D., Zanton, S. J. & Pugh, B. F. Identification and distinct regulation of yeast TATA box-containing genes. *Cell* **116**, 699–709 (2004).

61.   Jin, Z. *et al.* Unraveling the regulatory dynamics of bidirectional promoters for modulating gene co-expression and metabolic flux in Saccharomyces cerevisiae. *Nucleic Acids Res.* **53**, gkaf511 (2025).

62.   Gruber, J. D., Vogel, K., Kalay, G. & Wittkopp, P. J. Contrasting properties of gene-specific regulatory, coding, and copy number mutations in Saccharomyces cerevisiae: frequency, effects, and dominance. *PLoS Genet.* **8**, e1002497 (2012).

63.   Bédard, C. *et al.* FungAMR: a comprehensive database for investigating fungal mutations associated with antimicrobial resistance. *Nat. Microbiol.* 1–15 (2025) doi:10.1038/s41564-025-02084-7.

64.   Kemble, H., Nghe, P. & Tenaillon, O. Recent insights into the genotype-phenotype relationship from massively parallel genetic assays. *Evol. Appl.* **12**, 1721–1742 (2019).

65.   Vanella, R. *et al.* Understanding activity-stability tradeoffs in biocatalysts by enzyme proximity sequencing. *Nat. Commun.* **15**, 1807 (2024).

66.   Amorosi, C. J. *et al.* Massively parallel characterization of CYP2C9 variant enzyme activity and abundance. *Am. J. Hum. Genet.* **108**, 1735–1751 (2021).

67.   Prywes, N. *et al.* A map of the rubisco biochemical landscape. *Nature* **638**, 823–828 (2025).

68.   Grønbæk-Thygesen, M. *et al.* Deep mutational scanning reveals a correlation between degradation and toxicity of thousands of aspartoacylase variants. *Nat. Commun.* **15**, 4026 (2024).

69.   Choudhury, A. *et al.* Deep mutational scanning reveals the molecular determinants of RNA polymerase-mediated adaptation and tradeoffs. *Nat. Commun.* **14**, 6319 (2023).

70.   Beltran, A., Jiang, X. 'er, Shen, Y. & Lehner, B. Site-saturation mutagenesis of 500 human protein domains. *Nature* **637**, 885–894 (2025).

71.   Otwinowski, J., McCandlish, D. M. & Plotkin, J. B. Inferring the shape of global epistasis. *Proc. Natl. Acad. Sci. U. S. A.* **115**, E7550–E7558 (2018).

72.   Wylie, C. S. & Shakhnovich, E. I. A biophysical protein folding model accounts for most mutational fitness effects in viruses. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 9916–9921 (2011).

73.   Serohijos, A. W. R. & Shakhnovich, E. I. Merging molecular mechanism and evolution: theory and computation at the interface of biophysics and evolutionary population genetics. *Curr. Opin. Struct. Biol.* **26**, 84–91 (2014).

74.   Tokuriki, N. & Tawfik, D. S. Stability effects of mutations and protein evolvability. *Curr.*
Opin. Struct. Biol. **19**, 596–604 (2009).
75.   Després, P. C. *et al.* Compensatory mutations potentiate constructive neutral evolution by
gene duplication. *Science* **385**, 770–775 (2024).
76.   Lagator, M., Paixão, T., Barton, N. H., Bollback, J. P. & Guet, C. C. On the mechanistic
nature of epistasis in a canonical cis-regulatory element. *Elife* **6**, (2017).
77.   Nagalakshmi, U. *et al.* The transcriptional landscape of the yeast genome defined by RNA
sequencing. *Science* **320**, 1344–1349 (2008).
78.   Luthra, I. *et al.* Regulatory activity is the default DNA state in eukaryotes. *Nat. Struct. Mol.*
Biol. **31**, 559–567 (2024).
79.   Meneu, L. *et al.* Sequence-dependent activity and compartmentalization of foreign DNA in a
eukaryotic nucleus. *Science* **387**, eadm9466 (2025).
80.   Cools, H. J., Hawkins, N. J. & Fraaije, B. A. Constraints on the evolution of azole resistance
in plant pathogenic fungi. *Plant Pathol.* **62**, 36–42 (2013).
81.   Todd, R. T. & Selmecki, A. Expandable and reversible copy number amplification drives
rapid adaptation to antifungal drugs. *Elife* **9**, e58349 (2020).
82.   Brachmann, C. B. *et al.* Designer deletion strains derived from Saccharomyces cerevisiae
S288C: a useful set of strains and plasmids for PCR-mediated gene disruption and other applications.
Yeast **14**, 115–132 (1998).
83.   Gietz, R. D. & Schiestl, R. H. High-efficiency yeast transformation using the LiAc/SS carrier
DNA/PEG method. *Nat. Protoc.* **2**, 31–34 (2007).
84.   Janke, C. *et al.* A versatile toolbox for PCR-based tagging of yeast genes: new fluorescent
proteins, more markers and promoter substitution cassettes. *Yeast* **21**, 947–962 (2004).
85.   Ryan, O. W., Poddar, S. & Cate, J. H. D. CRISPR-Cas9 Genome Engineering in
Saccharomyces cerevisiae Cells. *Cold Spring Harb. Protoc.* **2016**, db.prot086827 (2016).
86.   Engel, S. R. *et al.* Saccharomyces Genome Database: advances in genome annotation,
expanded biochemical pathways, and other key enhancements. *Genetics* **229**, iyae185 (2025).
87.   Amberg, D. C., Burke, D. J. & Strathern, J. N. *Methods in Yeast Genetics 2005: A Cold*
Spring Harbor Laboratory Course Manual. (Cold Spring Harbor Laboratory Press, New York, NY,
2005).
88.   Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: summarize analysis results for
multiple tools and samples in a single report. *Bioinformatics* **32**, 3047–3048 (2016).
89.   Masella, A. P., Bartram, A. K., Truszkowski, J. M., Brown, D. G. & Neufeld, J. D.
PANDAseq: paired-end assembler for illumina sequences. *BMC Bioinformatics* **13**, 31 (2012).
90.   Rognes, T., Flouri, T., Nichols, B., Quince, C. & Mahé, F. VSEARCH: a versatile open
source tool for metagenomics. *PeerJ* **4**, e2584 (2016).
91.   Rice, P., Longden, I. & Bleasby, A. EMBOSS: The European molecular biology open

software suite. *Trends Genet.* **16**, 276–277 (2000).

92.	Seabold, S. & Perktold, J. statsmodels: Econometric and statistical modeling with python. in *9th Python in Science Conference* (2010).

93.	Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat. Methods* **17**, 261–272 (2020).

94.	Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *Journal of Machine Learning Research* **12**, 2825–2830 (2011).

95.	Chereji, R. V., Ramachandran, S., Bryson, T. D. & Henikoff, S. Precise genome-wide mapping of single nucleosomes and linkers in vivo. *Genome Biol.* **19**, 19 (2018).

 Extended Data Figures:

Extended Data Fig. 1: **The deletion of the F4 promoter fragment is clearly adaptive in 5-FC conditions. a,** Schematics of the four 77 bp fragments into which the intergenic region between *JID1* (YPR061C) and *FCY1* (YPR062W) was divided. There is a 26 bp overlap between F2 and F3. The nucleosome-depleted region (NDR) of the corresponding bidirectional promoter [61,95] as well as the TATA-like element, involved in transcription initiation at *FCY1* [46,47], are shown. **b,** Fitness of deletion mutants in [12 μM] 5-FC. The ΔF1-F4 mutant refers to deletion of the complete intergenic region, while Δregion instead represents the simultaneous deletion of the intergenic region and the *FCY1* gene. Exponential growth rates and the corresponding selection coefficients (*Methods*) show that F4 is the only promoter fragment for which deletion confers a clear growth advantage (and thus 5-FC resistance). **c,** Fitness of reconstructed WT strains in [12 μM] 5-FC. Here, F2, F4 and F1-F4 refer to CRISPR reconstruction of the WT intergenic region from the corresponding deletion mutant (*Methods*). In all three cases, the WT 5-FC sensitivity is recovered. All growth measurements were made from three replicate cultures of each strain.

1448
Extended Data Fig. 2: Complete overview of the bulk competition experiment. a, Full experimental
design showing all 32 cultures and their relationships as transformation and/or culture replicates. Each
pool of substitutions (Sub), insertions (Ins) or deletions (Del) was constructed four times and
corresponding replicates were mixed to obtain each of the four *transformation replicates* (grey
rectangles). Each transformation replicate was used to inoculate two culture replicates, with or without
the addition of a set of previously characterized coding sequence mutations (*FCY1*; n=960). Created in
BioRender. Aubé, S. (2025) https://BioRender.com/acpk3ks. **b,** The selected coding sequence variants
cover the full range of variant effects reported in [6], both in 5-FC and cytosine (without uracil) conditions,
where Fcy1 activity is respectively toxic and essential.

1458

Extended Data Fig. 3: **Coding sequence mutations included as positive controls are in agreement**
with previously reported variant effects. All mutants which were added to the "+*FCY1*" samples were
classified into resistant (R), and sensitive (S) variants according to their effect in 5-FC conditions, using
Gaussian mixture models (*Methods*). This was done using either the FC score reported by [6] or the
selection coefficients inferred in the current study (left). The confusion matrix (right) shows that, among
all variants which were classified as resistant using data from [6], none appear sensitive to 5-FC in the
current study. Mutated sequences were collapsed into amino acid substitutions (as in [6]), resulting in 311
variants from 960 nucleotide mutations.

Extended Data Fig. 4: **Mixed cultures of promoter and coding sequence variants were quickly taken over by the _FCY1_ mutants. a,** Gel electrophoresis image of the first PCR amplification step to prepare sequencing libraries of the F3-F4 promoter region (_Methods_), for the eight +_FCY1_ cultures. Primers were designed to be specific to the pool of promoter mutants, to avoid amplifying the WT promoter sequence of the added _FCY1_ variants. On the left, T0 and T1 PCR reactions have been loaded in alternance (from sample 1A to sample 4B), while the same was done on the right for T2 and T3 PCR reactions. Almost no band is visible for the T2 and T3 samples, although the same volume of PCR reaction was loaded in all cases. This suggests a much lower quantity of template DNA, and thus a smaller population size for promoter mutants. **b,** Comparisons of inferred selection coefficients in [12 µM] 5-FC for variants in the pool of promoter mutations, across samples with and without added coding sequence mutants. Variants which are identified as "CDS" are secondary mutations in the coding sequence which occurred during construction of the F3-F4 library (see **Supplementary Fig. 1**). Median selection coefficients display high agreement between the two types of samples (left). Using the minimal _s_ observed among replicate cultures (right) however reveals outlier negative values for samples with added _FCY1_ mutants. This is consistent with the stochastic disappearance of some genotypes due to the decreased population size of F3-F4 promoter mutants.

Extended Data Fig. 5: **Validation of the 42bp-mEGFP construction. a,** Medians of normalized fluorescence intensity, as measured by flow cytometry (*Methods*), for three mEGFP insertions at the genomic *FCY1* locus. Each point is a distinct clone of the corresponding strain, except for the WT. In the 42bp-mEGFP construct, which was used in the current study to measure promoter activity, the mEGFP green fluorescent protein is fused in C-terminal of the first 42 bp of the *FCY1* coding sequence. In contrast, FCY1-mEGFP designates a C-terminal fusion with full-length *FCY1,* while mEGFP stands for the insertion of the fluorescent protein in place of the *FCY1* coding sequence. **b,** As on the left, but showing the full distribution of log-scaled normalized fluorescence intensities for each clone (5000 events).

1494

Extended Data Fig. 6: **Indels before and after the second methionine codon of *FCY1* have**
contrasting effects on expression level. a, Comparison of relative expression levels for the four indel
mutations which were included among the set of validation mutations, classified according to whether
they occur before or after the second methionine codon of *FCY1*. Cytometry measurements from
reconstructed mutants as well as estimates obtained from the large-scale sort-seq experiment are shown.
b, Relative expression levels, as estimated from the sort-seq experiment, for all detected indel mutations
in the *FCY1* coding sequence. Those occurring after the second methionine are clearly more impactful.
c, Median effect of insertion and deletion variants detected in the sort-seq (top) and bulk competition
(bottom) experiments. The shaded area encompasses the three bases of the second methionine codon.
Mutations occurring before this codon were recovered at a much lower frequency from the bulk
competition, which is consistent with the observation that they do not completely abolish *FCY1*
expression and thus do not confer resistance.

1507

Extended Data Fig. 7: More detailed look at the calibration curve of sort-seq scores. a, LOWESS
fit as in **Fig. 4c**, but performed separately for true single mutants and for the minority of validation
mutants (11/30) which featured an additional T insertion within a poly-T region (positions -110 to -99)
in the F3 promoter fragment. The two fits result in almost the same curve, thus showing that these
unwanted double mutants do not have a systematic effect on the conversion of Δ_{Ref} scores into relative
expression levels. **b,** Sort-seq scores observed across all six replicates for the corresponding insertion
(I) of a single T at position -110. All values are below the median Δ_{Ref} observed across the six sort-seq
experiments (dashed line). **c,** Relative expression level inferred from the sort-seq scores shown in **b**
(**Fig. 4c**), with error bars displaying the corresponding 95% prediction interval. Panels **b** and **c** show
that the insertion alone has a weak to nonexistent impact on *FCY1* expression, thus confirming the
robustness of relative expression estimates to the (accidental) inclusion of double mutants among the
validation set.

Extended Data Fig. 8: Deletion of putative TF binding sites is not more impactful than single-nucleotide mutations in the same regions. a, Deletion of positions -133 to -127 of the *FCY1* promoter, corresponding to site 1 (positions -132 to -127) of **Fig. 4e**. The WT-relative fluorescence of the deletion mutant, from three replicate measurements in flow cytometry, is shown as an horizontal line whose width displays the minimum and maximum values among the three corresponding medians. Among reconstructed validation mutants (left), one single-nucleotide substitution decreases *FCY1* expression as much as the deletion, as shown by its median WT-relative fluorescence intensities (dots; grand median of three replicate cytometry measurements). Among all mutations assayed by sort-seq (right), the most impactful single-nucleotide changes decrease expression as much as the deletion, as shown by the corresponding relative expression level estimates (dots; median of the 95% prediction interval obtained from sort-seq calibration). **b,** Deletion of positions -125 to -118, corresponding to site 2 (-125 to -120) of **Fig. 4e**. The plots are the same as in **a**, but this time for a deletion which increases *FCY1* expression. Cytometry measurements of validation mutants (left) and sort-seq estimates (right) both show that many single-nucleotide changes have an effect that is comparable to that of the deletion – or even larger.

1536

Extended Data Fig. 9: The insensitivity of fitness to changes in *FCY1* expression level is constant
across many concentrations of 5-FC and cytosine. a, Fitness functions in 5-FC. In [12 μM] 5-FC
(left), a constant plateau of low fitness is reached starting from ~0.18 WT-relative expression (same as
Fig. 5b). Across three additional concentrations of 5-FC (right), the boundary of the plateau does not
change. Its height however varies, such that lower growth rates are observed for increasing 5-FC
concentrations. **b,** Fitness functions in media lacking uracil, where deamination of cytosine by Fcy1p is
essential. When cytosine is added at [84 μM] (left), a similar plateau is observed from ~0.18 WT-relative
expression. Any further increase in *FCY1* expression does not increase growth rate, such expression
level variations around the WT have no fitness effect. When three additional concentrations of cytosine
are tested (right), the boundary of the fitness plateau also remains constant. Error bars around each dot
display minima and maxima of three replicate measurements for growth rate and relative fluorescence
intensity for each combination of promoter and 5-FC/cytosine concentration. In addition, dashed
vertical lines identify the WT expression level (relative fluorescence intensity of 1.0).