The shape of fitness functions and the distribution of mutational effect sizes jointly limit adaptation by regulatory mutations

Simon Aubé[1,2,3,4,5*]**, Alexandre K. Dubé**[1,2,3,4,5] **& Christian R. Landry**[1,2,3,4,5*]

1. Département de Biochimie, microbiologie et bio-informatique, Université Laval, 1045 Avenue de la Médecine, Québec, QC, Canada, G1V 0A6
2. Institut de Biologie Intégrative et des Systèmes (IBIS), 1030, Avenue de la Médecine, Université Laval, Québec, QC, Canada, G1V 0A6
3. PROTEO-Regroupement Québécois de Recherche sur la Fonction, l'Ingénierie et les Applications des Protéines, Université Laval, 1030, Avenue de la Médecine Québec, QC, Canada, G1V 0A6
4. Centre de Recherche en Données Massives de l'Université Laval, Université Laval, 1065, Avenue de la Médecine, Québec, QC, Canada, G1V 0A6
5. Centre de recherche en infectiologie, Université Laval, 2705 boul. Laurier, Québec, QC, G1V 4G2

Correspondence: simon.aube.2@ulaval.ca and christian.landry@bio.ulaval.ca

Abstract

Mutations in gene regulatory regions have been shown to play a role in rapid adaptation but the factors determining their contribution are largely unknown. Using the yeast metabolic enzyme cytosine deaminase, we examine if adaptation to 5-fluorocytosine (5-FC), which requires reduced cytosine deamination and can readily arise from amino acid substitutions, may be reached by single promoter mutations. We generated all single-nucleotide substitutions and indels in the *FCY1* promoter and assayed the resulting mutants in presence of 5-FC. This revealed that no promoter mutation is sufficient for adaptation to occur. We next investigated how this inaccessibility of adaptation arises by combining large-scale expression measurements with the experimental characterization of the corresponding expression-fitness function. These experiments showed that the shape of the corresponding fitness function precludes single promoter mutations from being adaptive. Although 24% of mutations significantly affect expression, the fitness curve is flat around wild-type level. As such, adaptation can only emerge from a severe reduction of expression, which cannot occur from a single mutation in the promoter. Our results show that the contribution of regulatory mutations to rapid adaptation not only depends on the distribution of mutational effect sizes on expression level but also on the shape of the function linking fitness to expression levels.

Introduction

For rapid adaptation to occur, advantageous phenotypes must be immediately available through mutation. Whether a given mutation is adaptive depends on how the corresponding molecular change translates into a fitness effect. For expression changes, this is defined by the affected gene's fitness function (**Fig. 1a**) – describing the relationship linking expression level to organismal fitness –, of which a diversity of shapes have been reported[1]. As such, the corresponding distribution of mutational effects (DME) on expression level and the shape of the relevant fitness function jointly dictate if adaptive changes can arise within a given cis-regulatory region. These two factors thus have a major impact on the available pathways to adaptation, as well as on the rate and mode of regulatory evolution. In spite of this importance, the interplay between them remains unknown for most genes[2]. Adaptation would for instance be unlikely to arise from the combination of a narrow DME, such that mutations have small effects on expression, with a step-like fitness function where only large expression changes are beneficial. Conversely, single-nucleotide mutations sampled from the same narrow DME could be adaptive if the fitness function instead displayed more gradual variation, allowing small expression changes to provide a benefit (**Fig. 1b**). Fully understanding the evolution of cis-regulatory sequences and its contribution to adaptation hence requires assaying mutational effects on expression level and characterizing their conversion through fitness functions[2].

Across organisms, a major engine of rapid adaptation is loss-of-function (LOF) mutations, both in coding and regulatory sequences. In humans, the best known mechanisms of resistance to malaria are the loss of expression of specific genes in red blood cells[3, 4] and the partial loss of a key metabolic enzyme activity[5]. Among the best documented adaptive morphological changes in vertebrates in natural populations arose through the repeated loss of expression of a developmental gene[6]. LOF mutations in coding sequences are also a major pathway to antimicrobial resistance, for instance in fungal pathogens evolving in response to antifungal drugs[7]. Expression-decreasing epigenetic changes are involved in the adaptive evolution of cancer cells, by reducing expression of tumor suppressor genes[8]. Finally, laboratory microbial evolution has shown that adaptation occurs through LOF at least as frequently as through gain of function(s)[9–16]. LOF therefore provides a powerful model to investigate questions regarding the contributions of mutations to rapid adaptation.

Adaptive LOF sometimes occurs in regulatory elements of genes and sometimes in coding sequences. While this relative contribution could be caused by chance only, some gene properties could make one type of change more likely to occur than another. However, past experiments are insufficient to answer this question. Large-scale studies which investigated the effect of mutations within promoters and cis-regulatory sequences[17–23] mostly focused on expression level, without directly measuring fitness effects. In few cases where the fitness consequences of promoter mutations are well known, for instance in *TDH3*[24–26] and *SUL1*[27] in yeast, only a

93 handful of coding sequence mutations have comparatively been assayed for the same
94 genes[28,29]. The contribution of regulatory mutations to rapid adaptation, relative to that
95 of coding changes, thus remains unclear.
96

Fig. 1: The DME and the shape of the fitness function jointly define the
accessibility of adaptation. a, Fitness functions describe the relationship between
the expression level of a gene and organismal fitness. Hypothetical fitness functions
are shown for two genes. In each case, the shape of the curve displays how fitness
varies with increasing expression level of the corresponding gene. **b,** The accessibility
of adaptation through expression level changes depends on mutational effect sizes
and fitness function shape. For the two fitness functions, adaptation from the same
low-fitness wild-type (WT) expression level is shown. When the function is flat around
WT expression (left), only large expression increases from the high-variance DME can
be adaptive. In contrast, when the fitness function is curved near WT expression (right),
small expression increases are adaptive, such that adaptation is accessible from the
low-variance DME as well as the high-variance one. Created in BioRender. Aubé, S.
(2025) https://BioRender.com/rw8qljy.

We focused on the adaptive loss of Fcy1 activity in yeast exposed to prodrug 5-
fluorocytosine (5-FC). While this enzyme's canonical activity is cytosine deamination,
it also converts 5-FC into 5-fluorouracil, which leads to growth arrest and cell death[30].
Accordingly, complete or partial LOF mutations in the *FCY1* coding sequence confer
resistance and are adaptive[31]. To assess whether regulatory mutations can also
contribute to rapid adaptation to this condition, we performed saturation mutagenesis
of the *FCY1* promoter. We measured the effect of all possible single nucleotide
substitutions and indels in this region on fitness and expression level. These
experiments revealed no measurable fitness gain in medium with 5-FC, although a
quarter of single-nucleotide changes significantly affect gene expression level.
Through the characterization of the underlying fitness function, we show that this
discrepancy emerges jointly from the molecular effect of mutations and from the shape

of this relationship. Because the fitness function is flat around the endogenous expression level, adaptation can only occur through large expression decreases, which cannot arise from a single-nucleotide change in the promoter. These findings highlight how the availability of adaptative phenotypes through mutation can be restricted by the combined action of the DME and of the relevant fitness function. In *FCY1*, this results in rapid adaptation being only accessible through mutations in the coding sequence. This observation may be generalizable to other enzymes which exert low control over metabolic fluxes.

Results

Measuring the effects of mutations within the *FCY1* promoter

To identify promoter regions to mutate, we performed deletions of successive 77 bp fragments of the intergenic region upstream of *FCY1*. This revealed that only the most proximal fragment, among four, is essential for gene function (**Extended Data Fig. 1**). For this reason, and because most regulatory elements in yeast promoters are found within the 150 bp closest to the transcribed gene[32,33], we focused on the 154 bp immediately upstream of *FCY1*. This corresponds to two fragments hereafter named F3 and F4 (**Fig. 2a**). Within this region, we generated all single-nucleotide mutations – substitutions, insertions and deletions. The corresponding variants of F3 and F4 were inserted at the endogenous *FCY1* locus using a CRISPR-Cas9-based pooled approach (**Fig. 2b**, *Methods*[34,35]). This resulted in 1021 unique mutated sequences (462 substitutions, 463 insertions and 96 deletions), of which 1020 were successfully recovered by sequencing of the mutant pools (**Supplementary Fig. 2**). This process was repeated four times for each of the three classes of mutations (**Fig. 2c**). The corresponding pools of substitutions, insertions and deletions were then combined, resulting in four independent transformation replicates, each containing all mutations. Because the set of substitutions was obtained by successively degenerating each nucleotide position, the WT *FCY1* promoter was also (re)constructed along with the desired mutants. Within each combined transformation replicate, its initial frequency was ~17%.

From each of the four transformation replicates, two culture replicates were made, which were both grown in liquid media with [12 µM] 5-FC and without, resulting in 16 cultures. This concentration has been shown to inhibit the growth of *Saccharomyces cerevisiae* by ~50% and was previously used to assay all single-codon substitutions in the *FCY1* coding sequence[31] (**Extended Data Fig. 2**). Additional replicates of all 16 cultures were also done by combining the promoter mutants with 960 representative coding sequence variants[31], to ensure that the fitness effects of promoter mutations were directly comparable to those of coding mutations(**Extended Data Fig. 3**).

Fig. 2: Experimental design of the *FCY1* promoter deep mutational scanning.
a, 5-FC is converted into 5-FU, a toxic compound, by the cytosine deaminase (Fcy1p), such that *FCY1* expression can have a strong effect on fitness. Fragments 3 (F3) and 4 (F4) of the *FCY1* promoter were replaced with a library of synthetic sequences corresponding to all possible single-nucleotide substitutions (Sub), insertions (Ins) and deletions (Del). The oligonucleotides used to sequence the region and genotype mutants are marked in red (short fragment) and black (extended fragment, including the recombination breakpoints). **b,** Mutated F3 and F4 sequences were inserted at the endogenous yeast *FCY1* locus by CRISPR-Cas9 genome editing. A NAT resistance cassette was first introduced in F3 or F4 by homologous recombination. Using a specific guide RNA, Cas9 could be targeted to the cassette, such that synthetic DNA fragments would be inserted at F3 or F4. This step was repeated four independent times. **c,** The three types of pools (Sub, Ins, Del) were combined and grown in synthetic complete media (SC) and then transferred to media with 5-FC to allow for competition based on the abundance of Fcy1p. Loss of function of *FCY1* has no detectable effect in SC media without 5-FC. At every timepoint, DNA was extracted and the edited locus was sequenced to follow the frequency of each mutant in the culture, allowing to estimate selection coefficient for each mutation from their enrichment or depletion through time. Schematics created in BioRender. Aubé, S. (2025) https://BioRender.com/frtbee2.

All 32 cultures were grown for ~20 generations in liquid media, with dilution every ~7
generations to maintain exponential growth (*Methods*). Samples were taken at each
dilution step, as well as before the start of the experiment, resulting in a series of
timepoints from T0 to T3 (**Fig. 2c**). The relative frequencies of all mutations were
monitored by sequencing of the F3-F4 region (**Fig. 2a**). Due to the toxicity of Fcy1
activity in the presence of 5-FC, we expected that mutations reducing gene expression
would prove adaptive and increase in frequency, outcompeting those resulting in a
higher expression level. Accordingly, we estimated the selection coefficient (s)
associated with each promoter sequence from the slope of its log-scaled frequencies
relative to the WT[36] (**Fig. 2c**).

Mutant pools sampled from the competition experiment were first genotyped by
sequencing a 174 bp fragment, which covered the full F3-F4 region and the first 20 bp
of *FCY1* (red oligonucleotide pair; **Fig. 2a**). There were however major reproducibility
issues. While estimated selection coefficients were consistent between culture
replicates of the same transformation, they diverged substantially across
transformation replicates (**Supplementary Fig. 1a**). In addition, comparisons of the
distributions of selection coefficients across the different time intervals revealed a
gradual shift of the mean. From ~0 for the T0-T1 interval, it decreased below -0.15 for
the full competition experiment (**Supplementary Fig. 1b**). Through further
experiments, we found that these effects were caused by secondary mutations
introduced during library construction (**Supplementary Fig. 1c-d**). We modified our
sequencing protocol to genotype a longer region (black oligonucleotide pair; **Fig. 2a**)
and repeated the analyses to eliminate these effects. We explain the observations in
detail in the supplementary material (**Supplementary Note**) as we suspect these
could be common in genome editing studies and thus be of broad interest.

Fitness is robust to *FCY1* promoter mutations

After correcting for the impact of secondary mutations, the inferred selection
coefficients for the F3-F4 mutants reveal only small fitness effects, which vary mostly
uniformly along the mutagenized region (**Fig. 3a**). Across all assayed promoter
mutations, median s estimates obtained from replicate experiments range from ~-
0.037 to ~0.035 gen^{-1}, for a grand median of ~-0.002 gen^{-1}. Experimental noise
moreover appears of a similar magnitude as these deviations from 0, as shown by an
average difference of ~0.028 gen^{-1} between the highest and lowest s among replicate
estimates for the same genotype, up to ~0.110 gen^{-1} (**Extended Data Fig. 4a**). A
small number of secondary mutations remain undetected (not found within the longer
sequencing fragment), resulting in outlier selection coefficients for one pair of culture
replicates within some F3-F4 genotypes. Following stringent removal of these rare
outliers (**Supplementary Fig. 8**), none of the 1020 s estimates for *FCY1* promoter
mutations appear significantly different from 0 ($p > 0.05$; one-sample t-test with
Benjamini-Hochberg correction at FDR=5%). Identifying significant effects using $|s|$
thresholds on the most extreme replicates instead results in small numbers of

significant changes (12 at $|s| > 0.01$), which are consistent with the null expectation from random permutations of the data (**Extended Data Fig. 4**). Overall, we thus conclude that there were no measurable fitness effects among the assayed single-nucleotide mutations.

Fig. 3: Mutations in the *FCY1* promoter do not have a measurable impact on fitness in 5-FC conditions. a, Fitness effects of single-nucleotide mutations in the *FCY1* promoter. All substitutions (Sub; n=462), insertions (Ins; n=462) and deletions (Del; n=96) are shown. Points display the median selection coefficient for each mutant genotype, while error bars show the minimal and maximal values observed at each position along the mutated region. Nucleotide positions which appear to be missing are due to mutations which result in the same genotype (**Supplementary Fig. 2**), for instance deletion of any single base in a homopolymer region. Only data from two unbiased transformation replicates (each duplicated into two culture replicates; four samples in total) are shown (**Supplementary Note**, **Supplementary Fig. 6**). **b,** Distribution of mutational effects on fitness for single-nucleotide mutations in the *FCY1* promoter. The median selection coefficient obtained for each single mutant is shown, either as separate distributions for each of three classes of mutation (top) or as the full distribution (bottom; n=1020 observations, one per mutant).

This absence of mutational effects is not due to a failure of the competition experiment, as confirmed by two layers of controls. First, when considering all mutations (F3-F4 and secondary), the variance of estimated selection coefficients is indeed much

narrower for cultures without 5-FC (**Supplementary Fig. 9**). Second, selection coefficients inferred for the added coding sequence mutants correlate well with previously published variant effects[31] ($\rho = 0.780$ with $p < 10^{-64}$; **Extended Data Fig. 5a**). There is moreover almost perfect agreement between the two experiments regarding the number and identity of resistance-conferring mutations (**Extended Data Fig. 5b**). Together, these results confirm that adaptive changes could have been observed in our assay, if they could occur through single-nucleotide promoter mutations. We note that, whereas samples which contained coding sequence variants also included the complete pool of promoter mutations (**Extended Data Fig. 3**), the resulting cultures were not used to precisely estimate F3-F4 mutational effects. Although noisier, the corresponding selection coefficients were nonetheless qualitatively similar to those reported in **Fig. 3a** (**Supplementary Fig. 10**).

This lack of adaptive mutations within the *FCY1* promoter contrasts markedly with the high accessibility of adaptation through changes in the corresponding coding sequence. Within the *FCY1* gene, ~29% (408/1411) of single-nucleotide substitutions indeed result in codon changes which confer resistance to 5-FC in the same conditions (**Extended Data Fig. 5c**)[31]. Among these, only 67 introduce a premature stop codon. As such, although adaptation to 5-FC is readily available through *FCY1* mutations, it cannot be conferred by single-nucleotide changes in the promoter. In order to understand how this inaccessibility of adaptation arises, we first measured the effect of promoter mutations on *FCY1* expression.

Many mutations in the *FCY1* promoter substantially affect expression level

To obtain expression level measurements for all promoter mutants, we used a sort-seq approach[37,38]. The complete pool of F3-F4 single mutants was reconstructed in a genetic background where the distal part of the *FCY1* gene – starting from its 43rd base – was replaced with a mEGFP sequence (*Methods*, **Extended Data Fig. 6**). As such, the translation initiation context could be maintained while adding a fluorescent reporter to measure promoter activity. Yeast cells of the corresponding mutant library were sorted into five bins (numbered 0 to 4) according to their log-scaled fluorescence intensity (**Fig. 4a, Supplementary Fig. 11**). All the resulting samples were sequenced and read counts were used to compute a weighted average of bins for each genotype (*Methods*). The effect of each mutation was quantified using a Δ_{Ref} score equal to the difference between the corresponding mean bin and the average obtained for the WT. This was performed separately within each of six replicates (three transformation replicates, each subdivided into two culture replicates).

Fig. 4: Mutations in the *FYC1* promoter impact expression level. a, Schematics of the sort-seq experiment. Yeast cells from pools of promoter mutants were sorted into five bins according to their fluorescence intensity (representative distribution shown). From the sequencing of each bin, WT-relative fluorescence intensity could be estimated as a Δ_{Ref} score. Created in BioRender. Aubé,S. (2025) https://BioRender.com/uzzv5xy. **b,** Median sort-seq scores obtained for all variants detected across six replicate experiments. In addition to 1019 F3-F4 variants ("Promoter"), 28 single mutations in the first 42 bp of *FCY1* were identified ("CDS"), of which the vast majority are indels. The full distribution of scores ("All") as well as the 30 mutants selected for reconstruction ("Validation") are shown, split between the two categories in each case. **c,** Calibration curve of sort-seq scores**.** Points display the medians of sort-seq scores (n=6) and WT-relative fluorescence intensities (n=3) for all 30 validation genotypes. Horizontal and vertical error bars show the minimal and

maximal observations associated with each median. The high correlation between
medians is shown with Spearman's rho ($p < 10^{-23}$). The line and the corresponding
shaded area display a LOWESS fit with its 95% confidence interval (*Methods*). **d,**
Distributions of WT-relative expression levels for all single-nucleotide promoter
mutations, as inferred from the calibration of sort-seq scores. At the top, distributions
are shown separately for substitutions (Sub; n=461), insertions (Ins; n=462) and
deletions (Del; n=96), with a dashed line indicating WT expression level (1.0). At the
bottom, the full distribution, combining all three types of single mutants, is shown
(n=1019). **e,** Distribution of promoter mutations with a significant effect on expression
(n=245), along the mutagenized region. Separately for Sub (n=113), Ins (n=111) and
Del (n=21) variants, the estimated WT-relative expression (point) and its
corresponding 95% prediction interval (error bar) are shown for each mutant affecting
FCY1 expression by 5% or more (minimum or maximum of the corresponding 95%
prediction interval; *Methods*). Shaded rectangles display the minima and maxima
across all prediction intervals for each position along the sequence, including variants
which are not shown. In the upper plot, vertical lines show the midpoint of hits (n=41)
to known TF recognition motifs, identified using YeTFaSCo[39]. In the lower plot, the
black and white rectangles respectively display A/T and G/C positions. Promoter
positions corresponding with the two main clusters (sites "1" and "2"), the approximate
AT-rich region and the TATA-like element ("TA")[40,41] are also identified.

A narrow range of scores was obtained, with the median Δ_{Ref} of most promoter
mutations falling between -1 and 0.5 (**Fig. 4b**). Of the 1020 F3-F4 mutants, 1019 were
covered in at least two replicates after abundance-based filtering (*Methods*). Many
secondary mutations were also recovered (**Fig. 4b**), including indels in the first 42 bp
of *FCY1*. Interestingly, while most such mutations were associated with a low
fluorescence intensity and thus a negative Δ_{Ref}, some did not completely abolish
expression. We specifically found out that indels which occurred before the second
methionine codon of *FCY1* maintained ~10% of the WT fluorescence level (**Extended**
Data Fig. 7 a,b). This is consistent with a previous report of a minor translation
initiation site at this codon[42].

To validate the sort-seq assays, we selected 30 mutants for independent
reconstruction (**Fig. 4b**). For each newly constructed strain as well as for the WT
reference, fluorescence intensity was measured by flow cytometry (*Methods*). The
resulting WT-relative intensities are in strong agreement with the corresponding
median Δ_{Ref} scores (**Fig. 4c, Supplementary Fig. 12**). We additionally used this
validation to draw up a calibration curve, enabling the conversion of sort-seq scores
into more easily interpretable relative fluorescence intensities. To this end, we fit a
LOWESS regression (**Fig. 4c**), to avoid making specific assumptions about the
relationship between the two variables. This was performed within a bootstrapping
routine which involved repeated resampling of Δ_{Ref} values as well as validation
fluorescence measurements, thus generating 95% prediction intervals for the curve
itself and for the relative fluorescence intensities of all genotypes (*Methods*,
Supplementary Table 9). We hereafter refer to the medians of the latter intervals as

WT-relative expression levels for the corresponding mutations, where a value of 1
denotes the WT level.
349
Examining the inferred relative expression levels for all F3-F4 mutants reveals a slight
bias towards expression increases, with medians lying between 1.04 and 1.06 (**Fig.**
4d). That mutations are more likely to increase expression than to decrease it may
appear surprising, since Fcy1p is already highly expressed and is among the 10%
most abundant proteins in *S. cerevisiae*[43]. While this bias could be artefactual, the y-
intercept of the calibration curve suggests otherwise (**Fig. 4c**). It is very close to 1
(~1.00, with 95% prediction interval from 0.98 to 1.02) – as expected for a Δ_{Ref} of 0 –
, although no WT measurements were used in the LOWESS fitting.
358
The overall distribution of effects highlights that most single mutations in the *FCY1*
promoter impact expression levels by less than 15%, with 75% of variants lying
between values of 0.98 and 1.11 (**Fig. 4d**). Yet, we also observe a small number of
mutations with larger effects, decreasing expression to ~50% of its WT level, or even
lower. This tail appears more pronounced than previously reported in yeast[17,24],
potentially because we specifically mutated the most proximal part of the *FCY1*
promoter, where mutations are more likely to disrupt processes such as transcription
initiation. Among the 1019 F3-F4 mutations, 245 significantly affect *FCY1* expression,
by 5% or more (95% prediction intervals). In agreement with a previous study of a
yeast promoter[27], these significant mutations are not distributed uniformly along the
mutagenized promoter region (**Fig. 4e**). This distribution reveals important
determinants of *FCY1* expression level.
371
The fact that significant mutations form two main clusters, from positions -132 to -127
(site 1) and from -125 to -120 (site 2), first highlights the importance of transcription
factor (TF) binding. Within these two sites, 92.4 % (73/79) of all variants change
expression by 5% or more, in a consistent direction within each cluster. Both sites are
within a 80 bp segment known to be critical for TF binding in yeast promoters[18,37,44],
while the whole -132 to -120 region is a hotspot of predicted TF binding sites[39] (**Fig.**
4e top; Supplementary Table 10). As such, the clusters likely reveal TF binding sites.
Interestingly, the deletion of either of the two sites is not more impactful than the
strongest mutations within the corresponding cluster (**Extended Data Fig. 8**),
suggesting that single mutations are sufficient to disrupt them.
382
The position of the two most impactful mutations additionally highlights how alterations
of transcription initiation may severely reduce *FCY1* expression. These variants, a
substitution and an insertion which respectively decrease expression to ~16% and
~13% of its WT level (cytometry; **Fig. 4c**), both occur at position -9 (**Fig. 4e**). This
position is immediately downstream of the previously identified TATA-like element of
the *FCY1* promoter[40,41] – from -17 to -10 –, where the transcription pre-initiation
complex assembles. Additionally, the transcription of *FCY1* has been shown to initiate

at position -7[33,45]. The corresponding mutational effects thus likely reflect the disruption
of transcription initiation. The conflicting report of a most abundant transcript isoform
beginning further upstream, at -29[46], however adds uncertainty.
393
Finally, the complete set of significant mutations underscores that most single-
nucleotide changes within the poly(dA:dT) region of the *FCY1* promoter weakly affect
its transcriptional activity. The corresponding AT-rich sequence, approximately
between positions -118 and -73 (17.39% GC; **Fig. 4e**, bottom), is indeed comparatively
depleted in strong mutational effects. Only 17.6 % (52/295) of variants within it are
significant, compared to 26.7 % (193/724) in the remainder of the mutagenized region.
This may seem contradictory, considering that such regions are important for the
establishment of nucleosome-depleted regions within promoters[47,48], which favor
transcription and can be sufficient to induce expression from a downstream core
promoter[49]. It is however consistent with previous work on the *URA3* promoter, which
showed that deletion of more than half of the corresponding 50 bp poly(dA:dT) was
necessary to abolish expression[50].
406
407   Overall, we show that 24% (245/1019) of single-nucleotide mutations in the *FCY1*
408   promoter have a significant effect on expression level, many of which appear to disrupt
409   the binding of TFs at two putative sites. As such, *FCY1* expression is sensitive to
410   mutations, meaning that the DME on expression level is not sufficient to explain the
411   absence of fitness changes in 5-FC conditions.
412

The expression-fitness relationship for *FCY1* is mostly flat
To investigate the mechanism underlying the absence of measurable fitness effects,
we set out to characterize experimentally the expression-fitness relationship for *FCY1*
(**Fig. 5a**). Six yeast promoters covering a wide range of expression levels were used
to construct two sets of strains, in which each promoter was inserted upstream of either
the native *FCY1* sequence or the previously described 42bp-mEGFP construct
(**Extended Data Fig. 6c**). The *FCY1* strains were grown in media with 5-FC (12 µM;
like in the competition experiment) – to obtain growth rate measurements –, while
mEGFP strains were used to measure expression levels (*Methods*). For each
promoter, the measurements from the two strains were combined to draw the
corresponding expression-fitness curve (**Fig. 5a**).
424

This experiment revealed a very steep fitness function, where growth rate decreases
sharply until *FCY1* expression reaches ~18% of its WT level and then plateaus (**Fig.**
5b). Consequently, this fitness curve is flat around the WT, such that small expression
changes have no fitness effect. The shape of this function is thus consistent with the
fact that no single-nucleotide promoter mutation conferred resistance to 5-FC in our
bulk competition experiment. To further validate this result, we leveraged previously
characterized validation mutants. The set of alleles used in the calibration of sort-seq
measurements, which also includes a small number of indels in the coding sequence,

was reconstructed in the native *FCY1* background and the resulting strains were phenotyped in 5-FC conditions (*Methods*). In combination with the corresponding relative fluorescence intensities, these growth measurements confirm the flatness of the curve around WT expression level. Almost all promoter mutants display WT-like growth rates, with no clear trend along the complete range of effects (**Fig. 5c**). Among three assayed coding sequence mutations, the only one which reached the high fitness region is associated with a ~98% decrease in fluorescence intensity (**Supplementary Table 11**), caused by a frameshift-inducing indel. This highlights how adaptation is only possible through very large reductions in Fcy1 abundance, which cannot be achieved by single promoter mutations. Zooming in on the promoter mutants further illustrates this point. Among them, the two single mutations with the largest effect barely reach the threshold below which fitness rapidly increases, although they respectively decrease *FCY1* expression to ~16% and ~13% of its WT level (**Fig. 5d**). As such, the shape of the fitness function, jointly with the corresponding DME, renders adaptation to 5-FC inaccessible from single changes in the *FCY1* promoter.

The major contribution of the fitness function is made clearer by comparing the selection coefficients inferred from the competition experiment to the distribution expected from the relationship between 5-FC concentration and growth rate (**Extended Data Fig. 2a**). Under the assumption that varying the abundance of Fcy1p would have the same effect as changing the concentration of 5-FC (*Methods*), the most impactful promoter mutation would have been associated with a *s* of 0.19, while seven other mutations would have resulted in $s > 0.1$ (**Fig. 5e**). Adaptation would then have been accessible through regulatory changes, albeit much less readily than from mutations in the coding sequence.

Whereas growth rate varies according to 5-FC concentration, the shape of the fitness function remains mostly constant. Across concentrations resulting in ~10% to ~70% inhibition for the WT, growth rate variation is restricted to the same narrow range of relative expression levels and only the height of the corresponding plateau varies (**Extended Data Fig. 9a**). A similar phenomenon is observed in media lacking uracil, where the deamination of cytosine into uracil by Fcy1p is essential for growth. With increasing concentrations of cytosine, the height of the plateau of high fitness increases, while the region of varying fitness remains mostly the same (**Extended Data Fig. 9b**). At 5-FC and cytosine concentrations resulting in similar levels of relative inhibition or growth, the fitness functions obtained in the two conditions are very similar, with a sign inversion (**Extended Data Fig. 10ab**). This suggests that our results also extend to the canonical cytosine deaminase function of the Fcy1 enzyme. Growth measurements for a set of promoter mutants covering the full range of mutational effects confirm that increasing *FCY1* expression, as allowed by the DME of single mutations, is not adaptive under uracil starvation (**Extended Data Fig. 10c**). Overall, these experiments demonstrate that adaptation is not accessible from single-nucleotide changes in the *FCY1* promoter, in LOF (5-FC) as well as in gain-of-function

477 (cytosine) contexts. This is due in part to the small number of large mutational effects
478 on gene expression, but more importantly to the shape of the fitness function, which
479 is flat around WT expression level.

480
Fig. 5: The *FCY1* expression-fitness function is flat around WT expression level,
making fitness robust to promoter mutations. a, Strains with varying expression
levels were constructed to measure the fitness function. A set of yeast promoters
covering a wide range of activities were introduced in the genome in place of the
endogenous *FCY1* promoter, either upstream of *FCY1* or of a fluorescent reporter.
Created in BioRender. Aubé, S. (2025) https://BioRender.com/1p60mp7. **b,**
Expression-fitness relationship for *FCY1* in 5-FC conditions. All strains with *FCY1*
expression equal to or higher than ~18% of its WT level display a growth rate which is
indistinguishable from the WT. For each promoter, medians of growth rate and WT-
relative fluorescence (n=3) are shown along with the corresponding minima and
maxima (error bars). The fitness function is displayed as a fitted exponential decay
function, with its corresponding 95% confidence interval (*Methods*). The WT strain,

which lacks a resistance cassette and is not directly comparable to the other ones (*Methods*), was not used in the fit. **c,** Growth rates of reconstructed mutants are consistent with the fitness function. For validation mutants (**Fig 4c**) which could also be reconstructed in the *FCY1* background, medians (n=3) are shown for growth rates in 5-FC [12µM] media and WT-relative fluorescence intensities. This includes single promoter mutations ("Promoter"; n=26) and indels in the first 42 bp of the gene ("CDS"; n=3). The dashed line identifies the WT expression level (1.0), while the shaded area displays the range of growth rates measured for the WT background (panel **a**), into which the mutations have been reconstructed. **d,** Reconstructed promoter mutations are mostly neutral. For the same set of single mutations in the promoter (n=26), medians (n=3) are shown for selection coefficients and relative fluorescence intensities, with error bars displaying the corresponding minimal and maximal replicates. Selection coefficients are computed from growth measurements (*Methods*) shown in the previous panel, relative to the WT strain. The previously fitted fitness function (panels **b** and **c**) is displayed with its corresponding 95% confidence interval, after conversion into selection coefficients. Only the region of the expression-fitness curve where promoter mutants lie is shown. The outlier mutant which is hidden is associated with a median s of ~-0.28. **e,** The shape of the fitness function prevents expression changes from affecting fitness. The DME on fitness observed in the competition experiment is much narrower than the one expected from the relationship between the concentration of 5-FC and growth rate (**Extended Data Fig. 2a**).

Discussion

The rapid accessibility of adaptive changes through regulatory mutations depends on their effect sizes on gene expression and on the relationship between fitness and expression level. One powerful approach to investigate these determinants is to focus on a gene where adaptation is known to arise and systematically assay all single regulatory mutations. To this end, we studied the promoter of the gene *FCY1*, a protein-coding region within which single-nucleotide changes have been shown to confer resistance to the drug 5-FC[31,51]. We show that no promoter mutation has a measurable fitness effect in yeast exposed to this molecule, although large changes in expression level (reaching a reduction of 87%) are observed. In this adaptive landscape, beneficial phenotypes are thus inaccessible from single nucleotide variation in the promoter, although they arise readily from mutations in the corresponding coding sequence[7,31].

Our initial data instead suggested that most promoter mutations were highly deleterious. Further investigations revealed that this result was due to secondary resistance-conferring mutations introduced during library construction, outside of the sequenced region (**Supplementary Note; Supplementary Fig. 1,4-8**). These beneficial mutations largely occurred in the WT promoter background, since this genotype was generated at high frequency within the mutant pool. This comparatively made all true promoter single mutants appear less fit. Interestingly, bias thus arose from the reconstruction of the WT within the CRISPR library, which is widely regarded as a best practice to ensure the accuracy of mutant fitness estimates[52]. Such artifacts

may be common in large-scale genome editing experiments, due to the increased frequency of single-nucleotide mutations – and especially indels – in synthetic DNA oligonucleotides[53]. This highlights the importance of designing library construction workflows which minimize the use of genic regions as homology arms to guide recombination. Alternatively, these segments should at least be fully included in the genotyping region, as we did to correct our experiment.

Within our model system, the inaccessibility of adaptation through promoter mutations emerges jointly from the DME on expression level and from the shape of the fitness function. Since these two parameters vary across genes[1,17], the lower adaptive potential of promoter mutations is only generalizable to those which are similar to *FCY1* in this regard. There may be a substantial number of such genes, as the *FCY1* promoter is a typical yeast promoter. It is TATA-less like the majority of such regions [40,54], features the most common poly-T/poly-A configuration[48] and belongs to the most abundant class of yeast promoters (2474/5378)[55]. Although this promoter is bidirectional, hundreds of such regions have been identified in the *S. cerevisiae* genome and it drives much higher expression of *FCY1* than of the opposite gene[56]. The endogenous expression level of Fcy1 is in addition not exceptional for an enzyme. Among proteins included in the recent yeast9 genome-scale metabolic model of *S. cerevisiae*[57], more than 16% (196/1161) are more abundant than Fcy1p[43]. Because of these typical features, our results likely reflect general properties.

The generality of our findings is first shown by the distribution of expression changes we report, which is consistent with previous work. The large-scale mutagenesis of promoters and regulatory elements in yeast[17,24] and mammals[20–23] indeed showed that only a minority of single mutations strongly affect expression. In the yeast *TDH3* promoter, for instance, only 7.7% (18/235) of mutations were reported to decrease expression by more than 6%[24]. Within the *FCY1* promoter, we instead observe that 17.2% of single-nucleotide changes (175/1019) reduce expression level by at least this magnitude. As such, the rarity of strong expression-decreasing mutations is likely not unique to our model. This is further underscored by genome-scale computational predictions of cis-regulatory mutations in yeast, according to which 10% of mutations account for 50% of expression variation[18]. These observations suggest that, rather than being an exception, the DME on expression level which we report is probably representative of many yeast genes. It is however worth noting that it is likely incomplete, since only single-nucleotide indels were included. If indels of variable lengths[58] had instead been considered, larger expression changes could potentially have been observed.

Second, the shape of the fitness function, which severely restricts the adaptive potential of promoter mutations, is not atypical. Saturating abundance-fitness curves with a fitness plateau, like observed for Fcy1 (**Extended Data Figs 9, 10**), have generally been reported for enzymes, across organisms[1,25,26,59–63]. They are additionally expected from enzyme kinetics and metabolic control theory[60,64]. Whereas

the WT expression level can sometimes be close to the steep region where fitness
varies[1,63], it is more often away from this cliff, as we report here. In the case of *FCY1*,
the flatness of the fitness function around WT expression may arise from permeases
exerting control over metabolic flux, rather than the enzyme. This is suggested by the
observation that variations in 5-FC and cytosine concentrations have large effects on
fitness, contrary to *FCY1* expression changes (**Extended Data Figs 2, 9, 10**). A similar
phenomenon has been reported in bacteria, with the β-galactosidase enzyme and the
corresponding permease. While the former displays a step-like fitness function as
FCY1, the latter is associated with a much more curved fitness function, such that
small activity or abundance changes may have selective consequences[61]. More
generally, recent proteomics and metabolomics studies have shown that, in most
cases, changing the abundance of a single yeast enzyme has a weak impact on the
corresponding metabolic flux[57,65,66]. Overall, because the shape of the expression-
fitness relationship for *FCY1* likely emerges from fundamental principles, the
inaccessibility of adaptation through single promoter mutations may be generalizable
to a substantial number of genes, and especially enzymes.
598
Rapid adaptation through cis-regulatory changes may generally be rare, which would
be consistent with it being less accessible through mutation. For instance, increasing
the expression of drug target *ERG11* is a common mechanism by which fungi acquire
resistance to antifungals. Yet, it does not occur through single promoter mutations, as
only copy number variations (CNV) have been reported[67,68]. Similarly, whereas CNVs
of the sulfate transporter *SUL1* consistently arise in yeast evolved in sulfate-limited
conditions[69,70], single mutations in the corresponding promoter have been shown to
provide much less adaptive benefit[27]. There are however reports of single promoter
mutations conferring adaptation, especially in bacteria exposed to antibiotics[71–73].
Additionally, the analysis of natural variation between two yeast strains revealed an
enrichment of promoter mutations among variants with fitness effects[74]. In contrast,
mutations in noncoding regions have been found to be a very minor contributor to
adaptation in laboratory-evolved yeast[16]. Our results will help better understand in the
future why some genes are more likely to contribute to adaptation by regulatory
mutations by providing an integrative framework that connects mutations to phenotype
and fitness.
615
Contrary to promoter mutations, single-nucleotide substitutions in the *FCY1* coding
sequence readily confer adaptation to 5-FC[31] (**Extended Data Fig. 5**). In accordance
with the fitness function we report (**Fig. 5b**), this would mean that 29% (408/1411) of
coding mutations decrease the abundance or activity of the Fcy1 enzyme below 5%
of its WT level. As the strongest promoter mutation only decreases expression to 13%
of its WT level, this shows that, in *FCY1*, coding changes have larger molecular effects
than cis-regulatory ones. Whether this is generally the case across genes is unclear,
but saturation mutagenesis studies have shown that a substantial fraction of missense
mutations in protein sequences induce LOF[59,75–81]. Almost abolishing gene expression
may be comparatively difficult to achieve, as illustrated in yeast by genome-wide

spurious transcription[82] and by the expression of introduced exogenous and random
DNA[83,84]. As such, it is plausible that adaptation by LOF mostly occurs through coding
sequence mutations, as for *FCY1*. This may even be the case for rapid adaptation in
general, including gain-of-function. Two recent surveys of adaptive mutations, in yeast
evolved across different environments[16] and in pathogenic fungi exposed to diverse
antifungal drugs[7], have collectively identified only tens of adaptive noncoding changes,
compared with thousands of coding mutations. Yet, a large fraction of these mutations
are LOF-inducing. Elucidating whether coding or regulatory mutations generally
contribute more to adaptation will likely require additional experiments as well as a
systematic literature comparison similar to that of Hoekstra & Coyne[85].

These investigations should first focus on characterizing fitness functions. An
important role for fitness function shape in defining whether cis-regulatory or coding
mutations contribute more to adaptation was previously theorized from data on a
handful of mutations conferring β-lactam resistance in bacteria[86]. When the WT
resides on a fitness plateau, as is the case for *FCY1*, expression changes are unlikely
to provide a large fitness benefit (**Fig. 6a**). Coding mutations, on the other hand, may
cause complete LOF – like in our *FCY1* model – or bypass the fitness plateau by
modifying the curvature of the fitness function (in gain-of-function contexts), for
instance by changing the affinity of an enzyme for its substrate[86]. When the WT instead
resides in a region of the curve where fitness varies rapidly, expression variation
provided by regulatory mutations could be equally beneficial as coding changes (**Fig.**
6b), or even more if the fitness function is peaked and mutations may overshoot its
optimum (**Fig. 6c**). Second, a complementary avenue of investigation would be to
assess if the respective DME of promoter and coding regions are fundamentally
different at the molecular level. Within coding sequences, DME are often bimodal and
display peaks corresponding to WT-like and LOF mutations[59]. This is observed for
fitness effects[77,79–81] – as for the *FCY1* gene –, but also when other phenotypes like
fluorescence, protein activity and binding are measured[75,76,78,81,87,88]. Although such
multimodality can emerge from saturating abundance-fitness functions (like in **Fig.**
5b), it may also be a more general feature of mutations within coding sequences. This
could arise from nonlinear relationships between the biophysical properties of proteins
and fitness-relevant traits like protein abundance[59,87]. One example may be protein
stability, which dictates the fraction of correctly folded (functional) proteins through a
sigmoid function[79,89–91]. By potentially increasing the mutational accessibility of LOF,
this relationship could for instance underlie the major contribution of such mutations
to rapid adaptation[12,14,16]. Other properties of coding and cis-regulatory sequences
could make other categories of mutational effects more or less likely in one type of
region or the other.

Altogether, previous studies and our results demonstrate the importance of
considering the shape and curvature of fitness functions to understand the effects of
mutations and the molecular bases of adaptation. This goes beyond the relationship
between fitness and expression levels in microbes as it appears key for phenotypic

traits in general. For instance, it was recently shown that considering the full shape gene dosage response curves significantly enhances our understanding of genotype-phenotype relationships in humans[92].

Fig. 6: Different shapes of fitness function may favor mutational effects of different magnitudes. Assuming that mutations in coding sequences have larger effects on molecular traits than changes within promoters, as is the case for *FCY1*, this can modulate whether adaptation is accessible through one type of mutation or the other. Although we only measured *FCY1* expression, we generalize expression-fitness relationships to total enzymatic activity. **a,** A step-like function, which is flat around WT, favors mutations with large effects on total activity. Accordingly, adaptation is accessible through coding sequence changes (arrow), while promoter mutations do not increase fitness and are not selected for (blunt arrow). This is what we report for *FCY1*. **b,** A fitness function with high curvature around WT allows mutations with small and large effects to be adaptive. Here, less impactful promoter mutations are also beneficial, such that adaptation can occur through one large-effect coding sequence mutation or through successive promoter mutations. **c,** A non-monotonic fitness function may favor small changes in total activity. Because they overshoot the curve's optimum, coding mutations with large effects are deleterious and thus not selected for (blunt arrow), while promoter changes with smaller effects can be adaptive (arrow). Created in BioRender. Aubé, S. (2025) https://BioRender.com/tj8z7wz.

Methods

Strains, oligonucleotides, plasmids and culture media

All *S. cerevisiae* strains constructed in this study were generated from a BY4742 (*his3Δ leu2Δ lys2Δ ura3Δ*) background[93] harboring a hygromycin resistance marker at the *HO* locus (*hoΔ::HPHMX*). This AKD1123 parental background and all other strains are listed in **Supplementary Table 1**, along with the identity of the corresponding genomic inserts and PCR primers as well as the selective media used. The sequences and references for all oligonucleotides used in this study can be found in **Supplementary Table 2**, while references for all plasmids are listed in **Supplementary Table 3**. Recipes for culture media are detailed

703 in **Supplementary Table 4**. When specified, selective media was prepared by adding one or
704 multiple of the following agents: geneticin [200 µg/mL] (G418; Bioshop Canada),
705 nourseothricin [100 µg/mL] (NAT; Jena Bioscience) and hygromycin B [250 µg/mL] (HYG;
706 Bioshop Canada). All Sanger sequencing was performed at the sequencing platform of Centre
707 de recherche du CHU de Québec-Université Laval (Université Laval, Québec, Canada).
708

Yeast transformation

Whenever inserts which included a selection marker had to be inserted in the yeast genome,
a standard lithium acetate yeast transformation protocol was used[94]. After selection of
transformants on the appropriate selective media, four to eight clones were selected for
validation PCR, depending on the strain. Unless stated otherwise, colony PCR was performed
on each side of the integration site to confirm proper integration, using Taq polymerase
(Bioshop Canada). All genomic inserts were amplified using the KAPA HiFi HotStart DNA
polymerase (Roche Molecular Systems).
717

This approach was for instance used to delete each fragment F1 to F4 of the intergenic region
between *JID1* (*YPR061C*) and *FCY1* (*YPR062W*), as well as the full intergenic segment and
the complete *FCY1* region (including the corresponding gene), by the insertion of a *natNT2*
cassette[95]. This resulted in a set of six strains harboring a different deletion of the *FCY1* region
(SA0127, SA0128, SA0129, SA0130, SA0131, SA0132). Similar deletions of regions F3 and
F4 were also performed in the 42bp-mEGFP construct obtained by CRISPR-Cas9 yeast
transformation (see below), generating fluorescent strains SA0246, SA0247 and SA0254
725

CRISPR-Cas9 yeast transformation

When sequences needed to be inserted in the yeast genome without any flanking selection
marker, a CRISPR-Cas9 approach was used. This required the prior generation of a strain
harboring a *natNT2* cassette at the chosen insertion site, which was done through standard
yeast transformation as described above. Competent cells from this genetic background could
then be cotransformed with the pCAS-NAT plasmid (**Supplementary Table 3**) – encoding a
guide RNA engineered to target the Cas9 endonuclease to the *natNT2* cassette[34] – and the
desired repair DNA, amplified with KAPA HiFi HotStart DNA polymerase (Roche Molecular
Systems). A previously published CRISPR-Cas9 transformation protocol was followed[96].
Briefly, competent cells were prepared as described, and transformed with unpurified PCR
amplification of the insert, DNA carrier [10 mg/ml], pCAS-NAT and PLATE solution
(polyethylene glycol 2000 [40%], lithium acetate [0.1 M], Tris-EDTA [10 mM Tris-Cl, 1 mM
EDTA] buffer [pH=8.0]). After mixing, cells were incubated for 30 minutes at 30°C, heat
shocked for 15 minutes at 42°C, resuspended in YPD media and incubated for 4-6 hours at
30°C. Transformation reactions were plated on YPD media containing G418, to select for
uptake of the pCas plasmid, and incubated at 30°C for 48 hours. Eight colonies from each
transformation were then grown overnight (30°C with shaking) in liquid YPD and streaked on
solid YPD. After 48 hours of incubation at 30°C, one isolated clone from each streaking was
grown overnight in YPD (30°C with shaking) and spotted on both YPD and YPD+G418 solid
media, to confirm the loss of the pCAS plasmid. Because the AKD1123 parental background
is resistant to hygromycin, HYG was added to the YPD throughout. Correct integration was
then validated by colony PCR and Sanger sequencing.
748

749   Using this protocol, the reinsertion of deleted regions F2, F4 and F1-F4 (full intergenic) was
750   performed, resulting in strains SA0162, SA0166 and SA0170. It was also used to generate
751   the 42bp-mEGFP construct (SA0228) to measure *FCY1* expression level and all validation
752   clones expressing either *FCY1* or mEGFP from the *FCY1* locus.
753

Growth assays

Unless stated otherwise, the following protocol was used each time the growth of *FCY1* strains
was assessed in media containing 5-FC (Alfa Aesar) or cytosine (Alfa Aesar). Each yeast
strain of interest was first inoculated in three replicates in SC [pH=6] liquid media – potentially
with HYG, NAT and/or G418, depending on each strain's resistance marker(s) – and grown
overnight at 30°C with shaking. After ~16 hours of incubation, a sample (100 µl) of each culture
was centrifuged and resuspended in the same volume of sterile water, before being aliquoted
and diluted again in sterile water to obtain up to 250 µl of cells at 1 optical density (OD) unit/ml.
Growth assays were then assembled in Greiner flat-bottom transparent 96-well plates as
follows: 160 µl of media (SC complete for 5-FC or SC-Ura for cytosine); 20 µl of diluted cells
[OD=1]; 20 µl of 5-FC or cytosine diluted to obtain the desired final concentration in a volume
of 200 µl. Each plate was incubated for 48 hours at 30°C without shaking in a Tecan Infinite
M Nano (Tecan Life Sciences) and OD_{595} was measured at 15 minutes intervals.
767

To obtain exponential growth rates, the exponential part of each growth curve was first
identified. The end T_{max} of the exponential phase was defined as the time where the derivative
of OD, estimated as the moving median of differences between consecutive OD
measurements, reached its maximum. To limit experimental noise, the first two hours of OD
measurements were discarded, while any T_{max} above 30 hours or below 3 hours was ignored.
In such cases, the first ten hours of the growth curve were instead defined as the exponential
phase. Exponential growth rates were then estimated as the slope of the linear regression of
$log(OD)$ over time. When converting the growth rate of any strain i into selection coefficient
s_i, the following formula was used[97]:

$$s_i = \frac{\mu_i - \mu_{WT}}{\mu_{WT}} \log 2$$

778   Variables μ_i and μ_{WT} are the growth rates for the strain of interest and the corresponding WT
779   reference, respectively. This comparison was always made using WT measurements obtained
780   in the same experiment as the growth rate of strain i.
781

Construction of the mutant libraries

Amplification of mutated F3 and F4 promoter fragments

All single-nucleotide substitutions, insertions and deletions in the F3 and F4 regions were
ordered as six separate oPools Oligo Pools (Integrated DNA Technologies Inc.) containing
sequences of 117 bp, 118 bp and 116 bp, respectively. Within the different pools, each
oligonucleotide is a variant of the full-length F3 or F4 region (77 bp) with flanking 20 bp
homology regions at both ends. The latter sequences are respectively the 20 bp immediately
upstream or downstream of the corresponding fragment in the reference genome of *S.*
cerevisiae S288C, as retrieved from the Saccharomyces Genome Database[98]. For
substitutions, each position of each fragment was iteratively degenerated into all four possible
nucleotides by specifying a N base, while for insertions, the same mixed base was inserted in
3' of each position in the reference F3 and F4 sequences. When such an N is encountered,
an equal ratio of each of the four standard nucleotides (A, T, G, C) is dispensed during

synthesis. For the two pools of deletions, single-nucleotide removal was performed iteratively
at each position of F3 and F4 and duplicates (occurring whenever two or more consecutive
nucleotides are identical in the reference) were removed to obtain non-redundant sets. All
sequences ordered as part of the six oPools are listed in **Supplementary Table 5**.

Upon reception, each oPool was spun down and resuspended in fresh PCR-grade water, to
obtain a final concentration of 50 µM. Aliquots of 100 µl at 0.05 µM were then obtained by
serial dilution from 2 µl of the initial resuspension. From these aliquots, PCR amplification of
each oPool was performed in four replicates using KAPA HiFi HotStart DNA polymerase
(Roche Molecular Systems) and the following cycle: 98°C for 3 min; {98°C for 30 s; 60°C for
15 s; 72°C for 10 s} x 22; 72°C for 20 s; Hold at 10°C. Each reaction was done in 50 µl:

Volume (µl)	Reagent
33	PCR-grade water
10	KAPA HF buffer [5X]
1.5	dNTPs [10 mM]
1.5	Forward oligo [10 µM]
1.5	Reverse oligo [10 µM]
1.0	KAPA Hi-Fi HotStart DNA polymerase
1.5*	oPool [0.05 µM] (or PCR-grade water)
50	**Total**

In each case, the diluted oPool resuspension was not added to the reaction mix and was
instead added individually in each reaction tube. In addition, one control without the oPool
dilution was performed for each primer combination. Both primer pairs were designed to
extend the 20 bp homology regions included in the oPool, such that mutated F3 and F4
featured 60 bp homology arms upstream and downstream of the corresponding genomic
region. The two pairs were CLOP299-C3 / CLOP299-C4 and CLOP299-D3 / CLOP299-D4,
respectively to amplify F3 and F4 oPools.

Genomic insertion of mutated fragments in *S. cerevisiae*
To insert promoter mutations upstream of *FCY1*, competent yeast cells harboring a *natNT2*
deletion of the F3 or F4 region (strains SA0129 and SA0130) were transformed with the
corresponding PCR-amplified oPools. For each mutation category and each region, four
replicate transformations were performed from the corresponding four PCR reactions
described above – for a total of 24 transformations. The protocol described by Ryan *et al.*
(2016)[96] was followed (see *CRISPR-Cas9 yeast transformation*, above). After incubation,
G418-positive colonies from each agar plate were collected separately. Liquid YPD (5 ml) was
added to the plate, cells were resuspended by scraping with a glass rake and the suspension
was transferred to a 15 ml Falcon tube. From this suspension, glycerol stocks (687 µl cells
mixed with 313 µl sterile [80%] glycerol) were made for each transformation and stored at -
80°C. Two 500 µl aliquots from each tube were spun down and stored at -20°C after aspiration
of the supernatant. From these samples, mutant libraries were validated by PE150 sequencing
on an Illumina MiSeq instrument (Genomics Analysis Platform [PAG] of Institut de Biologie

828   Intégrative et des Systèmes, Université Laval, Québec). This approach was also employed to
829   insert the same set of promoter mutations upstream of the 42bp-mEGFP construct, using
830   strains SA0246 and SA0254. In this case, three replicate transformations were instead
831   performed per amplified oPool.
832
Selection of coding sequence mutations
Fifteen codon positions of the *FCY1* gene, for which all possible states had previously been
generated by DMS[31], were selected to obtain a diversity similar to that of the complete
mutational scanning of the F3-F4 region (15 positions x 64 codons = 960 mutants). Among
these positions, six were chosen for their wide range of variant effects and nine others were
selected randomly, to fully cover the previously reported range of mutational effects. The 15
corresponding pools of mutants were each grown for ~24 hours in 4 ml SC complete media
(30°C with shaking), from 25 µl of glycerol stock. Optical densities were measured and all
cultures were mixed in equal amounts to make four replicate 1 ml glycerol stocks (687 µl mixed
cultures and 313 µl sterile [80%] glycerol) of this pool of Fcy1p mutants, which were stored at
-80°C.
844
Pooling of mutants
The different sets of promoter mutations (substitutions, insertions and deletions, separately
for F3 and F4) were grown separately in 4 ml SC complete media, from 50-100 µl of the
corresponding glycerol stock. After ~16 hours of incubation at 30°C with shaking, OD_{595} was
measured for all cultures. Subpools were then mixed to obtain equal proportions of all single
mutants, according to their OD_{595} and how many unique genotypes they each contained.
851
For mutations in the *FCY1* background, the four replicate pools of coding sequence mutations
were also grown from 75 µl of glycerol stock. All subpools were then combined as described,
according to their replicate number (i.e. replicate 1 for F3 substitutions was mixed with
replicate 1 of all other subpools). Each subpool of promoter mutations was thus included in
only two final pools: one with added coding sequence mutations and the other without. For
mutations in the 42bp-mEGFP background, all subpools of promoter variants were combined
as described without the addition of other mutants, except that there were only three replicates
of each.
860

Bulk competition experiment in 5-FC media
Each mutant pool was grown in two replicates in 4 ml SC complete media (30°C, 200 rpm) for
~20 hours, from 111 µl or 65 µl of the corresponding glycerol stock. These inoculum volumes
were chosen to have at least 2000x cells for each mutant, respectively for pools with and
without added coding sequence mutations. The competition experiment was initiated by
diluting each preculture at final OD_{595} of 0.01 in 25 ml of SC complete media (with and without
5-FC [12 µM]) in 50 ml Falcon tubes. T0 samples were also prepared by centrifuging two
microtubes containing 5 OD_{595} units for each preculture (850 g for 5 minutes; 4°C). After
aspiration of supernatant, cell pellets were stored at -20°C. Newly inoculated cultures were
incubated (30°C with shaking) in the dark until OD_{595} reached ~1.0. At this first timepoint T1,
each culture was diluted in at final OD_{595} of 0.01 in 25 ml of fresh media. Additional cell pellets
were prepared from the remaining cultures. The full volume was first centrifuged at 4°C (1000
g; 5 minutes). Cell pellets were resuspended in ~5 ml of supernatants and two samples of ~5
OD_{595} units were prepared as previously. This process was repeated at timepoint T2 when the

new cultures reached OD_{595} ~1.0 and final T3 samples were prepared when the newly
inoculated T2-T3 cultures reached the same OD_{595}. The timings of each timepoint and the
corresponding OD_{595} values for all mutant pools are listed in **Supplementary Table 6**.
878

Sort-seq experiment

One final pool of F3-F4 mutants in the 42bp-mEGFP background was grown overnight from
100 µl glycerol stock in 5 ml of SC complete [pH = 6.0] media, at 30°C with shaking. After
measurement of its OD$_{595}$, the culture was diluted to 0.15 OD/ml in 20 ml of fresh SC complete
[pH = 6.0] and incubated for 5 hours at 30°C with shaking, to reach exponential phase (~0.5
OD/mL). From this, 4.5 ml were transferred to a 5 ml polystyrene round-bottom Falcon tube.
The remaining 15.5 ml were centrifuged to collect cells, so that genotype frequencies before
sorting could also be measured by sequencing.
887

A FACSMelody cell sorter (BD Biosciences) was used to sort yeast cells based on their
fluorescence intensity. Five bins were defined along the distribution of log_{10} fluorescence
intensities, numbered 0 to 4 from the lowest to the highest fluorescence levels (Supplementary
Methods). More cells were collected from the most populated parts of the distribution (50 000
cells each from bins 2 and 3, versus 10 000 cells each from bins 0,1 and 4). Each bin was
sorted twice, resulting in two technical replicates, in a 96 well plate. The resulting fractions
were each used to inoculate 3 ml of SC complete [pH = 6.0] media in a 24 deep-well plate and
grown for 24h at 30°C with shaking (400 rpm). Each culture was spun down at 3000 rpm (1000
g) for 5 min and all pellets were washed with sterile water and transferred to a 1.5 ml
microtube. All microtubes were centrifuged at 15 000 rpm (21 000 g) for 1 min, supernatants
were removed and pellets were frozen at -80°C. This process was repeated two times for the
two other biological replicates, such that six replicate samples were obtained for each bin.
900

DNA library preparation and sequencing

The same approach was used for all sequencing libraries, for both the bulk competition and
sort-seq experiments. Genomic DNA was first extracted from each cell pellet using a standard
phenol-chloroform extraction protocol[99], including RNAse treatment. For the resequencing of
bulk competition samples in PE300, an additional RNAse treatment (10 minutes, 37°C) was
performed on previously purified DNA samples, followed by precipitation with sodium acetate
and cold ethanol. A first PCR was then performed using Q5 High-Fidelity DNA Polymerase
(New England Biolabs) to amplify the genotyping region chosen for each type of samples,
which included the F3-F4 promoter regions and flanking sequences of varying lengths. This
PCR step added Illumina primer-binding sites at both ends, separated from the fragment of
interest by small numbers of N bases (between 2 and 8; 10 total per PCR amplicon). For the
initial (PE150) sequencing of bulk competition samples, primers CLOP311-C6 to CLOP314-
H9 were used (as ordered in **Supplementary Table 2**). Two reactions were made from each
DNA extraction from cultures with added coding sequence mutants. One was specific to the
F3-F4 pool while the other targeted only the added *FCY1* variants. This was possible because
the pool of coding sequence mutations was generated from a codon-optimized sequence,
harboring many nucleotide changes from the WT [31]. For the PE300 resequencing of the same
samples, only F3-F4 PCR were performed, using oligos CLOP328-A6 to CLOP328-H7.
Finally, for the sequencing of sort-seq bins, also in PE300, oligos CLOP328-A6 to CLOP341-
A10 were employed. In this case, 7 N bases were added per amplicon.
921

In all cases, amplicons from this first PCR step were purified with AMPure XP magnetic beads (Beckman Coulter Life Sciences) and diluted to 0.01 ng/μl. These dilutions were used as template for a second set of PCR reactions using Q5 polymerase, which added Nextera barcodes and Illumina adapters i5 and i7. A unique combination of i5 and i7 indexes was used for each DNA sample. The final PCR products were also purified with AMPure XP beads, before quantification using a NanoDrop spectrophotometer (Thermo Fisher Scientific). Amplicons from the same sequencing round (PE150 competition, PE300 competition or PE300 sort-seq) were mixed in equal proportions based on the NanoDrop quantification and the resulting pool was sent for sequencing. PE150 sequencing was performed on a NovaSeq Illumina instrument (sequencing platform of Centre de recherche du CHU de Québec-Université Laval, Québec), while PE300 sequencing was performed on an AVITI instrument from Element Biosciences (Genomics Analysis Platform [PAG] of Institut de Biologie Intégrative et des Systèmes, Université Laval, Québec).

Analysis of sequencing reads

Following each sequencing run, reads were received already demultiplexed, according to the sample id associated with each combination of i5 and i7 adaptors used. Read quality was first assessed using FastQC (v 0.12.1) and MultiQC (v 1.12)[100]. Paired-end reads were next merged using PANDAseq (v 2.11)[101], with the '-k' parameter set to 4. The N positions which had been added at both ends during the first PCR step of library construction were trimmed using command "fastx_filter" of VSEARCH (v 2.28.1)[102], according to a table listing the number of N added for each sample. The first and last 20 bp of the resulting sequences, which were the binding sites for the oligonucleotide primers used in the first PCR step, were also trimmed using the same command. Following these trimming steps, all identical sequences were aggregated with the "derep_fulllength" command of VSEARCH. Each unique sequence was aligned to the corresponding reference (depending on the type of sample and the length of the sequenced region) using the needle aligner from the EMBOSS suite (v 6.5.7)[103]. For promoter (F3-F4) libraries, the opening and extension of gaps was equally penalized (4,4), while penalties of 50 and 0.5 were respectively used for opening and extension for *FCY1* coding sequence libraries[31]. Custom Python functions were then used to call variants from each alignment. In addition to WT reads, only sequences harboring single-nucleotide mutations (F3-F4 libraries) or single-codon substitutions (*FCY1* libraries) were kept for downstream analysis. These variants were further filtered by length, to remove those that differed too much from the expected length of the PCR fragment. For bulk competition samples, read counts of each single mutant were converted to WT-relative frequencies. These read counts were instead used as is in the downstream analysis of sort-seq samples.

Estimation of fitness effects
Calculation of selection coefficients

For each bulk competition culture, time series of WT-relative genotype frequencies were reconstituted from variant frequencies obtained at the four timepoints (T0 to T3). The number of generations associated with each timepoint was obtained from the corresponding durations in hours, which varied between cultures, and the doubling time of the WT strain in the same conditions (SC media with or without 5-FC [12 μM]), as estimated from growth curve experiments. For each single-mutant genotype, the selection coefficient s was obtained by fitting a linear model using the number of generations and the log-transformed WT-relative frequencies as dependent and independent variables, respectively. This was performed

through ordinary least squares regression, using the "ols" implementation of the statsmodels (v 0.12.2)[104] Python module, with an intercept added to the model. Regressions were performed iteratively over all time intervals which included the T0 (T0-T1, T0-T2, T0-T3; with intermediate data points when applicable). Because selection coefficients were inferred separately within each culture replicate, as many replicate values as there were cultures were obtained for each of the intervals mentioned above.

To identify the selection coefficients which should be used in further analysis, two diagnostics were performed. First, the agreement between the initial (T0) log-scaled WT-relative frequencies and the fitted intercept from each regression model was assessed. Second, the linearity of log-scaled frequencies through time, as expected for cultures growing exponentially, was validated graphically. This revealed that no interval was reliable for the initial 150 paired-end genotyping of promoter (F3-F4) libraries, while T0-T2 selection coefficients should be used for the 300 paired-end resequencing of the promoter fragments and the 150 paired-end sequence of *FCY1* coding sequence (see Supplementary Note). The chosen selection coefficients were then used to identify a T0 abundance threshold for variant filtering, such that inferred s would not be correlated with initial read counts. This threshold was set at 85 reads, both for the promoter (300 paired-end) and coding sequence (150 paired-end) sequencing libraries. To ensure that low frequency but beneficial secondary mutations could be assessed, any variant which was below the threshold at T0 but over it at the end of the experiment was kept.

Removal of outliers and aggregation of replicates
In 5-FC samples, mutations sometimes appeared beneficial in only one pair of cultures (coming from the same transformation replicate), likely due to secondary resistance-conferring mutations (See Supplementary Note). Outlier removal was thus performed, separately for two types of libraries (promoter or coding sequence). The promoter (F3-F4) libraries were further separated depending on whether coding-sequence mutants had been added in the corresponding cultures. Within each group of samples, the median absolute deviation (MAD) of selection coefficients was first computed for each mutant. This was then used to compute robust z-scores for each replicate measurement within each genotype. The resulting distribution of robust z-scores calculated across all mutant genotypes was compared and its 95th percentile was defined as the threshold to identify outlier replicates. The filtering of outliers was however restricted to a subset of mutants which displayed a higher variance, to ensure that small absolute variations between replicates were not identified as outliers. This was assessed from each genotype's standard deviation, with the 95th percentile of the overall distribution used as threshold. For F3-F4 sequencing libraries obtained from cultures with added coding sequence mutations, the 85th percentile was instead used.

To obtain final estimates of the fitness effects of promoter mutations, only F3-F4 libraries without added coding sequence mutants were taken into account. Four of the eight replicate cultures, corresponding to two of the four transformation replicates, were additionally ignored. In these four samples, selection coefficients displayed a temporal shift towards negative values which appeared consistent with some undetected resistance-conferring mutations being genotyped as WT (see Supplementary Note). For each of the 1020 assayed promoter mutations, the corresponding fitness effect was estimated as the median of the selection coefficients observed across the four remaining replicates, which were considered unbiased.

The statistical significance of selection coefficients was assessed from the same four replicate
measurements for each mutation, by comparing their mean to 0 using the SciPy (v 1.15.2)[105]
implementation of the one-sample t-test, followed by Benjamini-Hochberg correction
controlling the false discovery rate at 5%. For the coding sequence mutations added as
controls, fitness effects were instead estimated as the median of the eight replicate
measurements, without ignoring any of the corresponding samples. Alternatively, significant
selection coefficients were also defined according to thresholds of minimal absolute deviations
from 0, using the most extreme positive and negative replicates for each genotype. In this
case, the number of fitness changes identified was compared with a null expectation derived
from the observed experimental noise. To this end, the complete dataset of final selection
coefficients (from the four culture replicates mentioned above) was shuffled into 1020 random
"genotypes". From each culture replicate, 1020 selection coefficients were first sampled with
replacement. The resulting four sets of values were then combined randomly into the 1020
genotypes, such that each was associated with four replicate measurements. The number of
random genotypes exceeding the chosen threshold of absolute deviation from 0 was then
compiled. This was repeated 10 000 times.
1033
Comparison with published effects of amino acid substitutions in Fcy1p
The selection coefficients of added coding sequence mutations were first aggregated into
grand medians, according to the corresponding amino acid substitutions. They were then
compared with previously published variant effect scores obtained in the same conditions (12
µM 5-FC)[31]. The agreement between the two experiments was further assessed by classifying
variants as resistant and susceptible in both cases. To this end, Gaussian mixture models with
two components and a "tied" covariance matrix were fitted on each of the two datasets, using
the scikit-learn (v 1.6.1)[106] implementation.
1042
For the comparison of promoter and coding sequence mutations, the full dataset from[31] was
used. All single-nucleotide substitutions in the reference sequence of the *FCY1* gene, obtained
from the Saccharomyces Genome Database[98], were first performed computationally. The
corresponding variant effects were retrieved from the Després *et al.* (2022) data[31], according
to the resulting amino acid substitutions. Of 1431 possible single-nucleotide substitutions in
FCY1, only 20 were not covered, mostly in the first and last codons. The Gaussian mixture
model previously fitted on the full set of amino acid changes was then used to classify all single
mutants as either resistant or susceptible. From this classification, the fraction of adaptive
(resistance-conferring) mutations in the coding sequence was obtained.
1052
Calculation of sort-seq expression scores
Within each of the six replicate sort-seq experiments, all variants which reached a total
abundance of at least 85 reads across the five sampled bins were kept for downstream
analysis. Although this included a small number of secondary mutations (outside of the F3-F4
mutagenized region), any variant which occurred in the mEGFP sequence was filtered out.
For each of the remaining genotypes i, which includes the WT, fluorescence intensity was
quantified from a weighted average B_i of bins where the corresponding reads were detected:

$$B_i = \frac{\sum_{j=0}^{4} j \frac{r_{i,j}}{r_{tot,j}} \frac{n_j}{n_{tot}}}{\sum_{j=0}^{4} \frac{r_{i,j}}{r_{tot,j}} \frac{n_j}{n_{tot}}}$$

1060

In the above formula, j is the bin number, where 0 and 4 are associated with the lowest and
highest fluorescence intensities, respectively. Variables $r_{i,j}$ and $r_{tot,j}$ are the read count for
genotype i, as well as the total number of reads, in bin j. Finally, n_j and n_{tot} are respectively
the number of cells collected from bin j and the total amount of cells collected across the five
bins. The n_j/n_{tot} ratio was introduced because the number of yeast cells collected was not
uniform across all bins.
1067
1068 Separately for each replicate experiment, the difference was computed between the B_m
1069 weighted average of each mutant m and the B_{WT} weighted average obtained for the WT
1070 genotype:

$$\Delta_{Ref,m} = B_m - B_{WT}$$

1071

1072 For each mutant, up to six replicate Δ_{Ref} scores were obtained (one for each sort-seq
1073 experiment). The median of these values was used as an estimate of each mutation's effect
1074 on fluorescence intensity. Positive values denote an increase relative to the WT, while
1075 negative values instead indicate a decrease.
1076

Selection and construction of validation mutants
The sort-seq data was used to select 30 mutant genotypes for validation. These mutations
were chosen to cover the whole range of scores, as well as all regions of the promoter where
single-nucleotide changes appeared impactful. Substitutions were favored to ensure a high
success rate to the downstream reconstruction of genotypes. To ensure that low-confidence
mutations were not selected, a one-sample t-test (with Benjamini-Hochberg correction at
FDR=5%) was first performed on the replicate sort-seq scores to identify Δ_{Ref} values which
were significantly different from 0. This was also repeated with Δ_{mod} scores computed from
B_{mod} weighted averages obtained from reads of bin 1 to 4, excluding the 0th one displaying
the lowest fluorescence intensities. Because the Δ_{Ref} scores appeared biased towards
positive values compared with Δ_{mod}, the latter were used to select significant mutations to be
reconstructed, except when annotated *. This ensured a more stringent selection of genotypes
displaying altered fluorescence intensity compared with the WT reference. Within each of the
mutant groups described below, the specified number of sequences were chosen randomly,
unless stated otherwise. Among mutations which significantly decreased Δ_{Mod} below zero,
there were seven groups: 1) indels before the second methionine of *FCY1* (n=2); 2) indels
after the second methionine of *FCY1* (n=2)*; 3) top 3 lowest scores between positions -50 and
0 of the promoter*; 4) substitutions between positions -133 and -127 (n=6); 5) substitutions
between position -45 and -11 (n=2); 6) substitutions and insertions near the -75 position (one
each) and 7) two chosen substitutions outside of regions with many significant effects. Among
mutations which significantly increased Δ_{Mod} above zero, a similar approach was employed.
The four categories were: 1) substitutions between position -125 and -118 (n=4); 2)
substitutions between positions -112 to -109 (n=2); 3) substitutions between positions -63 to -
57 (n=2) and 4) variants among the 25 highest scores (n=4).
1101

The corresponding validation mutants were individually reconstructed using the CRISPR-
Cas9 yeast transformation protocol described above. For each genotype, this was performed
in two genetic backgrounds: fluorescent strain SA0247 (ΔF3-F4 with 42bp-mEGFP construct)
and *FCY1* deletion (ΔF3-F4) mutant SA0159 (see **Supplementary Table 1**). A fusion PCR
product harboring the single mutation of interest was used as donor DNA. This amplicon was
generated by two consecutive rounds of PCR. In the first round, two reactions were performed

with WT genomic DNA as template. The first one used a generic forward primer hybridizing
upstream of F3 (CLOP326-A11) with a mutant-specific reverse primer harboring the mutation
of interest (**Supplementary Table 2**). In the second reaction, a forward primer with the
mutation of interest (**Supplementary Table 2**) was used alongside a generic reverse primer
binding either in the mEGFP sequence (CLOP326-G10; for background SA0247) or in the
FCY1 coding sequence (CLOP326-E10; for background SA0159). The two mutant-specific
mutagenesis primers are complementary to each other, such that there is homology between
the two amplicons. The second PCR round used the two PCR products from the first round
diluted 1/40 as template and the same two generic primers used in the first pair of PCR
(CLOP326-A11 with either CLOP326-G10 or CLOP326-E10, depending on the mutagenized
background), which allowed fusion of both PCR parts. This approach thus allows to amplify
the full F3-F4 promoter region (with a short segment of *FCY1*) while introducing a single
mutation within it and adding homology regions on both ends for subsequent genomic
insertion. It was additionally used to perform deletion of short segments of the promoter (-133
to -127 and -125 to -118), similarly in both genetic backgrounds. After transformation of each
fusion PCR product, clones from each transformation were sent for Sanger sequencing to
validate proper genomic integration of the mutation. All 66 resulting strains, including
reconstructions of the WT promoter, are listed in **Supplementary Table 1**.
1126

Expression measurements by flow cytometry
Measurement of promoter activity using fluorescent strains was performed with a Guava
EasyCyte HT cytometer (CytekBio). Each strain of interest was grown overnight in three
replicates in SC complete media at 30°C with shaking, along with similar replicate cultures of
a reference strain with the WT promoter (SA0247 or AKD1462) and of a non-fluorescent
control (AKD1123 or AKD1429). The overnight cultures were diluted at 0.15 OD_{595} in fresh
SC-Trp (MSG) media and grown at 30°C with shaking until ~0.5 OD_{595}. Exponentially growing
cells were then diluted to 0.05 OD_{595} in fresh media and single-cell fluorescence intensity
measurements were performed with the Guava EasyCyte HT system. For each replicate
culture of each strain, 5000 events were acquired (488 nm excitation; 525 nm detection). The
medians of the corresponding distributions of fluorescence intensities were used to estimate
relative expression levels.
1139

Within each cytometry experiment, events were first filtered out according to forward scatter
and side scatter values, with the 99th percentile of each distribution used as thresholds. All
fluorescence intensities were then normalized by the corresponding forward scatter
measurement. From these normalized fluorescence intensities, a median was computed for
each replicate culture assayed. These medians were then normalized by the grand median of
measurements made for the reference strain, such that they were converted into WT-relative
fluorescence intensities. The resulting values, and especially their medians, were used as
estimates of relative expression levels. Accordingly, three replicate measurements of WT-
relative fluorescence intensity were obtained for each tested strain, with a value of 1 being
equal to the WT level.
1150

Calibration of sort-seq scores and bootstrapping of relative expression
levels
WT-relative fluorescence intensities were first measured for all validation mutants
reconstructed in the SA0247 (42bp-mEGFP) background, as described above. These

measurements were then compared with the corresponding sort-seq expression scores and
a LOWESS regression was fitted. This approach allowed to simultaneously validate the sort-
seq experiments and convert the resulting scores into more interpretable WT-relative
fluorescence intensities. The Δ_{Ref} scores were used, since some genotypes with very low
fluorescence intensities were absent from the Δ_{mod} dataset. In addition, because
fluorescence intensity measurements from reconstructed mutants were used to fit the
relationship, any bias in Δ_{Ref} was unlikely to affect the downstream inference of relative
expression levels. So that uncertainties could also be obtained for each of the >1000 mutant
expression levels, both the LOWESS fitting and the conversion of sort-seq scores into relative
fluorescence intensities were performed within a bootstrapping routine.
1165
At the start of each iteration, Δ_{Ref} scores were first resampled with replacement. For each
genotype which had been included in the sort-seq analysis, as many values were sampled as
there were replicate measurements among the six experiments. Per-genotype medians were
then computed on the newly sampled data. For each reconstructed mutant for which
fluorescence had been measured directly, three median WT-relative fluorescence intensities
were sampled with replacement among the corresponding replicate measurements. Grand
medians were computed from each trio of observations, which were combined with the newly
recomputed median Δ_{Ref} scores of the corresponding validation genotypes to obtain 30 pairs.
From these data points, LOWESS regression was performed with median Δ_{Ref} as
independent variable, using the "nonparametric.lowess" implementation from statsmodels (v
0.14.4)[104] with default parameters. This regression was then evaluated at each of the >1000
median Δ_{Ref} values obtained from the first resampling steps. This generated an estimate of
WT-relative fluorescence intensity for each of the genotypes which had been assayed in the
sort-seq screen. The regression was also evaluated at 100 equally spaced values from the
minimal to the maximal median Δ_{Ref} observed experimentally. This process was repeated
5000 times. After all iterations had been completed, medians and corresponding 95%
prediction intervals were computed from the distribution of WT-relative fluorescence intensities
obtained for each mutant genotype. These medians were used as WT-relative expression
levels associated with the corresponding F3-F4 mutations. The same calculations were
performed for each of the 100 equally spaced values at which the LOWESS had been
evaluated, thus allowing to plot the relationship between Δ_{Ref} and relative fluorescence
intensity.
1188

Experimental measurements of the expression-fitness function

Fitting of dose-response curves

The WT background AKD1123 was phenotyped in liquid media containing various
concentrations of 5-FC and cytosine, according to the growth assays protocol described
above. From two to three replicates were made for each concentration of 5-FC and cytosine.
A five-parameter logistic model was then fitted, using the exponential growth rate
measurements and the concentrations of 5-FC / cytosine as dependent and independent
variables, respectively. The resulting curves were converted into relative inhibition or relative
growth, using measured growth rates in absence of 5-FC or at the highest cytosine
concentration as references.
1199
1200

Construction of strains with distinct *FCY1* expression levels

Six yeast promoters covering a wide range of expression levels were used to construct two sets of strains, respectively from *FCY1* background AKD1123 and from fluorescent background SA0228. The six promoter cassettes, each including a G418 resistance marker, were PCR-amplified from the corresponding plasmids[95] (**Supplementary Table 3**) using oligonucleotides CLOP338-G6 and CLOP338-H6 (**Supplementary Table 2**). The two primers provided homology for genomic insertion immediately upstream of *FCY1* (or of its first 42 bp for the SA0228 background), between the gene and its native promoter. Yeast transformations were performed in the two genetic backgrounds as previously described, using each of the corresponding PCR products. Transformants were selected on YPD+HYG+G418 media. Correct integration of the promoter cassette was validated by PCR in four clones from each transformation, using oligonucleotides CLOP310-D1 and CLOP291-A3 (**Supplementary Table 2**). Two to three positive clones were obtained in most cases and validated by Sanger sequencing, such that biological replicates could be used in downstream assays for the majority of genotypes. All 25 corresponding yeast strains are listed in **Supplementary Table 1**.

Measurements of growth rates and fluorescence intensities

The set of *FCY1* strains (background AKD1123) was used to measure growth rates in liquid media containing different concentrations of 5-FC or cytosine, following the growth assays protocol described above. The set of fluorescent strains (background SA0228) was instead used in flow cytometry experiments to measure expression levels (see Expression measurements by flow cytometry). In both cases, all available biological replicates were used for each genotype. When less than three strains were available, three replicate cultures were nonetheless inoculated.

The growth assays in 5-FC media were performed in more than one experiment. To combine measurements obtained for concentrations of 12 µM and 3 µM (**Extended Data Fig. 9**), a normalization step was first performed.Growth rates from the first experiment were adjusted to have the same grand medians as those of the second one, by the addition of the difference between the two experiments' grand medians. This was performed separately for each concentration and allowed to combine the two sets of growth rates, in spite of potential systematic differences between them – for instance if the true concentration of 5-FC in the media varied. The cytometry measurements, which were done in parallel with the growth assays, were also performed in two separate experiments. In this case, potential systematic differences were controlled for by computing WT-relative fluorescence intensities within each experiment and only combining measurements after this step.

Fitting of expression-fitness functions

Fluorescence measurements associated with each promoter were combined with the corresponding growth rates. Measurements made from the AKD1123 WT strain were however not used, because this strain lacks the G418 resistance marker included in promoter cassettes and is thus not directly comparable. An exponential function was fitted for each condition, using WT-relative fluorescence as the independent variable and growth rate as the dependent variable. This could model the sharp decline observed in 5-FC as well as the steep increase seen in cytosine conditions. The fitting of all expression-fitness curves was performed within a bootstrapping routine. For each promoter, two WT-relative fluorescence intensity measurements were first sampled with replacement from each cytometry experiment. Then,

the four resulting values were resampled with replacement. Corresponding growth rates were
sampled similarly. If a given condition had been tested in two experiments, two replicate
growth rates were sampled with replacement from each experiment for each promoter, and
the four measurements were afterwards resampled with replacement. This approach was
employed to avoid favoring one experiment over the other, since the number of replicates
sometimes varied between them. When only one experiment had been performed, replicate
growth rates from this assay were instead resampled with replacement. Once relative
fluorescence intensities and growth rates had been sampled, values obtained for the same
promoter were paired randomly into data points. The exponential function was then fitted using
the "optimize.curve_fit" implementation of the SciPy module (v 1.15.2)[105] module and
evaluated over 250 equally spaced values covering the full range of WT-relative fluorescence
intensities. After 5000 iterations, medians and corresponding 95% prediction intervals were
computed for each of these 250 fluorescence levels, thus generating the expression-fitness
function. This was repeated for each of the 5-FC and cytosine concentrations tested.
1263

Inference of the expected distribution of effects on fitness for promoter mutations
Changes in *FCY1* expression were assumed to have the same effect on fitness as variations
in 5-FC concentration. The dose-response curve (**Extended Data Fig. 2**) was converted such
that concentrations were expressed as relative changes from 12 µM (used in the bulk
competition experiment). From this new curve, corresponding exponential growth rates were
inferred for the 1019 relative expression estimates obtained from the sort-seq experiment. All
growth rates were then converted into selection coefficients, as described above.
1271

Validation of *FCY1* promoter variants fitness effects
The previously described validation mutants reconstructed in the SA0159 *FCY1* background
(**Supplementary Table 1**) were grown in SC complete (MSG) media with 12 µM 5-FC and
without, according to the growth assays protocol described above. For all other mutants,
growth rates measured in 12 µM 5-FC were converted into selection coefficients as described
in the growth assays section, using growth rates measured in the same experiment for
CRISPR reconstructions of the WT genotype in the SA0159 background (generated at the
same time as the validation mutants). The expression-fitness function obtained in the same
conditions and the bounds of its 95% prediction interval were also converted into selection
coefficients using the same approach. In this case, the WT growth rate interpolated from the
curve was instead used as comparison.
1283

Assessment of 5-FC resistance arising from mutant library construction
To test whether mutations conferring 5-FC resistance were introduced during mutant library
construction, the process described in *Construction of mutant libraries* was repeated, but using
WT sequences of the F3 and F4 regions. Briefly, the same two oligonucleotide pairs
(CLOP299-C3/CLOP299-C4 and CLOP299-D3/CLOP299-D4) were first used to amplify F3
and F4 from genomic DNA of the AKD1123 strain, using KAPA HiFi HotStart DNA polymerase
(Roche Molecular Systems). Competent cells from strains SA0129 and SA0130 (prepared as
described by [96]) were then transformed with the F3 and F4 inserts, respectively. Two biological
replicates of each transformation were made. This was performed following the CRISPR-Cas9
approach described above for the pools of single mutations. After 48 hours of incubation at
30°C, transformants from each Petri dish were collected in 5 ml YPD by scraping, as

previously described. From each of the resulting cell suspension, cultures were started in 5 ml
of SC complete [pH=6] + HYG, from 100 µl of cells.

1297

After overnight incubation at 30°C with shaking, OD_{595} was measured for each culture and
dilutions at $OD_{595} = 1$ were made in 200 µl of sterile water. Additional dilutions were
afterwards made for OD_{595} values from 0.2 to 0.002. From each dilution, 150 µl were plated
on SC complete (MSG) solid media with and without 5-FC [100 µg/ml]. Petri dishes were
incubated at 30°C for 48 hours and colonies were counted. While colonies were too numerous
to be counted on SC complete controls, some dilutions plated on 5-FC media resulted in
countable colony numbers (between 20 and 340). For each plating (one per CRISPR-Cas9
transformation), colony counting was performed for two dilution levels, which acted as
technical replicates. Since the SA0130 background was initially resistant to 5-FC due to the
insertion of a natNT2 in place of F4, all associated platings were replica plated on YPD+NAT
media. Nourseothricin-resistant colonies were subtracted from the total number of 5-FC-
resistant colonies. The frequency of 5-FC resistance was then estimated as follows, separately
for each plating on media containing 5-FC:

$$F_{LOF} = \frac{n_{FC}}{OD_d \, V_{plate} \, c_{ml}}$$

1311
In the equation above, n_{FC} is the number of colonies on 5-FC media (after subtracting NAT
resistant ones, when applicable), while OD_d is the OD_{595} of the plated dilution, V_{plate} is the
volume deposited on each plate (0.15 ml) and c_{ml} is the number of yeast cells in 1 ml of media
at $OD_{595} = 1$, for which a value of 10 million cells/ml was used. The resulting F_{LOF} was
considered as a putative frequency of *FCY1* LOF in the corresponding transformations. To
validate whether that was the case, 16 colonies were selected for Sanger sequencing, using
primer CLOP326-F10, which covered nearly 70% of the *FCY1* coding sequence
(Supplementary Table 2). All colony counts, resistance rates and relevant calculations are
listed in **Supplementary Table 7**.

1321

Simulations with *invisible* resistance-conferring mutations
A simulation framework replicating the bulk competition experiment was designed. It involved
a library of neutral variants within which a small number of invisible resistance-conferring
mutations was introduced, thus mimicking secondary mutations occurring outside of the
genotyping region in the experiment. A set of variants of interest (hereafter named genotype
labels) was first generated by sampling 1000 neutral mutations from $s \sim \mathcal{N}(0, 0.004)$ four
times. The standard deviation used was set according to what was observed in sequencing
libraries from cultures without 5-FC. Four transformations were then obtained by replicate
random samplings of 10 000 clones, among which 17% were defined as WT ($s = 0$) and the
remaining were drawn from the set of genotype labels. Within each such transformation
replicate, a small fraction of clones was defined as harboring an invisible resistance-conferring
mutation (or resistance flag), which increased its s (between 0.2 to 0.4, depending on
simulations) without changing its genotype label. From each transformation, two replicate
cultures were generated by randomly sampling two sets of 2.5 million cells, resulting in eight
cultures, as in the bulk competition experiment (5-FC condition without added coding
sequence variants). From each culture, 1 million of sequencing reads (genotype labels) were
sampled randomly, to simulate sequencing of the initial T0 samples. Simulated cells in each
culture were next aggregated according to genotype labels and presence/absence of a
resistance flag, prior to calculation of WT-relative frequencies for each group (called true

genotypes). Growth over the course of a bulk competition experiment was simulated by modeling exponential changes of WT-relative frequencies through time for each true genotype i, as follows:

$$F_{i,T+1} = F_{i,T}\, e^{s_i n_{gen}}$$

The WT-relative frequency F of true genotype i at timepoint $T+1$ is obtained from the corresponding WT-relative frequency at previous timepoint T according to the genotype's true selection coefficient (increased by a resistance-conferring mutation when applicable) and the time interval n_{gen} between the two timepoints, in WT generations. Three such timepoints, from T1 to T3, were simulated and their respective durations were set in accordance with the bulk competition experiment. At each timepoint, 1 million reads as well as a new initial population of 2.5 million cells were sampled from each culture, as described above. Selection coefficients for all variants were then computed from time-series of simulated read counts, as described for experimental variant frequencies. This was performed from genotype labels only as well as with knowledge of true genotypes (resistance flags). Such simulations of one full experiment were repeated while varying both the initial frequency and the fitness effect of resistance-conferring mutations. For each combination, multiple replicate simulated experiments (each with eight simulated cultures) were performed.

Data availability

All sequencing data from bulk competition and sort-seq experiments has been deposited on the NCBI SRA (https://www.ncbi.nlm.nih.gov/sra/PRJNA1321317). All other experimental data and files needed to replicate the analysis have been deposited on Zenodo: https://doi.org/10.5281/zenodo.17088129.

Code availability

All analyses were performed using custom Python scripts and notebooks, which are available at: https://github.com/Landrylab/Aube_et_al_2025_FCY1promoter. A custom Python pipeline was deployed on a computing cluster (Institut de Biologie Intégrative et des Systèmes, Université Laval) to compute variant counts and selection coefficients from raw sequencing data. As described above, this used the following software: FastQC (v 0.12.1), MultiQC (v 1.12)[100], PANDAseq (v 2.11)[101], VSEARCH (v 2.28.1)[102] and EMBOSS suite (v 6.5.7)[103]. The versions of all Python modules used are included as requirements files, separately for the analysis pipeline and for the notebooks, which were executed locally.

Acknowledgements

We thank Philippe Després, Isabelle Hatin, Olivier Namy and all Landry lab members, especially Soham Dibyachintan, David Jordan, Pascale Lemieux and Camille Bédard, for helpful discussions. We are grateful to Pavithra Venkataraman, Angel Cisneros, Soham Dibyachintan and Philippe Després, who provided comments on this manuscript. We additionally thank Angel Cisneros for helpful suggestions on the simulation framework. This work was supported by a grant from the Natural Sciences and Engineering Research Council of Canada (NSERC RGPIN-2020-04844) for CRL. CRL holds the Canada Research Chair in Cellular Systems and Synthetic Biology. SA

was supported by doctoral research fellowships from Natural Sciences and Engineering Research Council of Canada and Fonds de recherche du Québec - Nature et technologies. The F65V-34a-mEGFP plasmid used in this study was a kind gift from Dr Michael Springer.

Author contributions

C.R.L., A.K.D and S.A. designed the study. S.A. and A.K.D. performed the experiments as well as the data analysis. S.A., A.K.D. and C.R.L. interpreted the results. S.A., A.K.D and C.R.L. wrote the manuscript.

References

1.   Keren, L. *et al.* Massively Parallel Interrogation of the Effects of Gene Expression Levels on Fitness. *Cell* **166**, 1282–1294.e18 (2016).

2.   Hill, M. S., Vande Zande, P. & Wittkopp, P. J. Molecular and evolutionary processes generating variation in gene expression. *Nat. Rev. Genet.* **22**, 203–215 (2021).

3.   McManus, K. F. *et al.* Population genetic analysis of the DARC locus (Duffy) reveals adaptation from standing variation associated with malaria resistance in humans. *PLoS Genet.* **13**, e1006560 (2017).

4.   Tournamille, C., Colin, Y., Cartron, J. P. & Le Van Kim, C. Disruption of a GATA motif in the Duffy gene promoter abolishes erythroid gene expression in Duffy-negative individuals. *Nat. Genet.* **10**, 224–228 (1995).

5.   Breeyear, J. H. *et al.* Adaptive selection at G6PD and disparities in diabetes complications. *Nat. Med.* **30**, 2480–2488 (2024).

6.   Chan, Y. F. *et al.* Adaptive evolution of pelvic reduction in sticklebacks by recurrent deletion of a Pitx1 enhancer. *Science* **327**, 302–305 (2010).

7.   Bédard, C. *et al.* FungAMR: a comprehensive database for investigating fungal mutations associated with antimicrobial resistance. *Nat. Microbiol.* (2025) doi:10.1038/s41564-025-02084-7.

8.   Guerrero-Preston, R. *et al.* Key tumor suppressor genes inactivated by 'greater promoter' methylation and somatic mutations in head and neck cancer. *Epigenetics* **9**, 1031–1046 (2014).

9.   Behe, M. J. Experimental evolution, loss-of-function mutations, and 'the first rule of

adaptive evolution'. *Q. Rev. Biol.* **85**, 419–445 (2010).

10. Gerstein, A. C., Lo, D. S. & Otto, S. P. Parallel Genetic Changes and Nonparallel Gene–

Environment Interactions Characterize the Evolution of Drug Resistance in Yeast.

Genetics **192**, 241–252 (2012).

11. Kvitek, D. J. & Sherlock, G. Whole genome, whole population sequencing reveals that

loss of signaling networks is the major adaptive strategy in a constant environment.

PLoS Genet. **9**, e1003972 (2013).

12. Lang, G. I. & Desai, M. M. The spectrum of adaptive mutations in experimental

evolution. *Genomics* **104**, 412–416 (2014).

13. Lind, P. A., Libby, E., Herzog, J. & Rainey, P. B. Predicting mutational routes to new

adaptive phenotypes. *Elife* **8**, e38822 (2019).

14. Murray, A. W. Can gene-inactivating mutations lead to evolutionary novelty? *Curr. Biol.*

30, R465–R471 (2020).

15. Durand, R., Jalbert-Ross, J., Fijarczyk, A., Dubé, A. K. & Landry, C. R. Cross-feeding

affects the target of resistance evolution to an antifungal drug. *PLoS Genet.* **19**,

e1011002 (2023).

16. Chen, P. & Zhang, J. The loci of environmental adaptation in a model eukaryote. *Nat.*

Commun. **15**, 5672 (2024).

17. Hodgins-Davis, A., Duveau, F., Walker, E. A. & Wittkopp, P. J. Empirical measures of

mutational effects define neutral models of regulatory evolution in Saccharomyces

cerevisiae. *Proceedings of the National Academy of Sciences* **116**, 21085–21093

(2019).

18. Vaishnav, E. D. *et al.* The evolution, evolvability and engineering of gene regulatory

DNA. *Nature* **603**, 455–463 (2022).

19. Tsuru, S., Hatanaka, N. & Furusawa, C. Promoters constrain evolution of expression

levels of essential genes in Escherichia coli. *Mol. Biol. Evol.* **41**, msae185 (2024).

20. Patwardhan, R. P. *et al.* High-resolution analysis of DNA regulatory elements by

synthetic saturation mutagenesis. *Nat. Biotechnol.* **27**, 1173–1175 (2009).

21. Patwardhan, R. P. *et al.* Massively parallel functional dissection of mammalian

enhancers in vivo. *Nat. Biotechnol.* **30**, 265–270 (2012).

22. Melnikov, A. *et al.* Systematic dissection and optimization of inducible enhancers in

human cells using a massively parallel reporter assay. *Nat. Biotechnol.* **30**, 271–277

(2012).

23. Kircher, M. *et al.* Saturation mutagenesis of twenty disease-associated regulatory

elements at single base-pair resolution. *Nat. Commun.* **10**, 3583 (2019).

24. Metzger, B. P. H. *et al.* Contrasting Frequencies and Effects of cis- and trans-

Regulatory Mutations Affecting Gene Expression. *Mol. Biol. Evol.* **33**, 1131–1146

(2016).

25. Duveau, F., Toubiana, W. & Wittkopp, P. J. Fitness effects of Cis-regulatory variants in

the Saccharomyces cerevisiae TDH3 promoter. *Mol. Biol. Evol.* **34**, 2908–2912 (2017).

26. Siddiq, M. A., Duveau, F. & Wittkopp, P. J. Plasticity and environment-specific

relationships between gene expression and fitness in Saccharomyces cerevisiae. *Nat.*

Ecol. Evol. **8**, 2184–2194 (2024).

27. Rich, M. S. *et al.* Comprehensive Analysis of the SUL1 Promoter of Saccharomyces

cerevisiae. *Genetics* **203**, 191–202 (2016).

28. Ringel, A. E. *et al.* Yeast Tdh3 (glyceraldehyde 3-phosphate dehydrogenase) is a Sir2-

interacting factor that regulates transcriptional silencing and rDNA recombination. *PLoS*

Genet. **9**, e1003871 (2013).

29. Kankipati, H. N., Rubio-Texeira, M., Castermans, D., Diallinas, G. & Thevelein, J. M.

Sul1 and Sul2 sulfate transceptors signal to protein kinase A upon exit of sulfur

starvation. *J. Biol. Chem.* **290**, 10430–10446 (2015).

30. Vermes, A., Guchelaar, H. J. & Dankert, J. Flucytosine: a review of its pharmacology,

clinical indications, pharmacokinetics, toxicity and drug interactions. *J. Antimicrob.*

Chemother. **46**, 171–179 (2000).

31. Després, P. C. *et al.* Asymmetrical dose responses shape the evolutionary trade-off

between antifungal resistance and nutrient use. *Nat. Ecol. Evol.* **6**, 1501–1515 (2022).

32. Lin, Z., Wu, W.-S., Liang, H., Woo, Y. & Li, W.-H. The spatial distribution of cis

regulatory elements in yeast promoters and its implications for transcriptional regulation.

BMC Genomics **11**, 581 (2010).

33. Mahendrawada, L., Warfield, L., Donczew, R. & Hahn, S. Low overlap of transcription

factor DNA binding and regulatory targets. *Nature* **642**, 796–804 (2025).

34. Lemieux, P., Bradley, D., Dubé, A. K., Dionne, U. & Landry, C. R. Dissection of the role

of a Src homology 3 domain in the evolution of binding preference of paralogous

proteins. *Genetics* **226**, iyad175 (2024).

35. Dionne, U. *et al.* Protein context shapes the specificity of SH3 domain-mediated

interactions in vivo. *Nat. Commun.* **12**, 1597 (2021).

36. Matuszewski, S., Hildebrandt, M. E., Ghenu, A.-H., Jensen, J. D. & Bank, C. A statistical

guide to the design of deep mutational scanning experiments. *Genetics* **204**, 77–87

(2016).

37. de Boer, C. G. *et al.* Deciphering eukaryotic gene-regulatory logic with 100 million

random promoters. *Nat. Biotechnol.* **38**, 56–65 (2020).

38. Peterman, N. & Levine, E. Sort-seq under the hood: implications of design choices on

large-scale characterization of sequence-function relations. *BMC Genomics* **17**, 206

(2016).

39. de Boer, C. G. & Hughes, T. R. YeTFaSCo: a database of evaluated yeast transcription

factor sequence specificities. *Nucleic Acids Res.* **40**, D169–79 (2012).

40. Rhee, H. S. & Pugh, B. F. Genome-wide structure and organization of eukaryotic pre-

initiation complexes. *Nature* **483**, 295–301 (2012).

41. McMillan, J., Lu, Z., Rodriguez, J. S., Ahn, T.-H. & Lin, Z. YeasTSS: an integrative web

database of yeast transcription start sites. *Database (Oxford)* **2019**, baz048 (2019).

42. Eisenberg, A. R. *et al.* Translation initiation site profiling reveals widespread synthesis of

non-AUG-initiated protein isoforms in yeast. *Cell Syst.* **11**, 145–160.e5 (2020).

43. Huang, Q., Szklarczyk, D., Wang, M., Simonovic, M. & von Mering, C. PaxDb 5.0:

Curated protein quantification data suggests adaptive proteome changes in yeasts. *Mol. Cell. Proteomics* **22**, 100640 (2023).

44. Erb, I. & van Nimwegen, E. Transcription factor binding site positioning in yeast: proximal promoter motifs characterize TATA-less promoters. *PLoS One* **6**, e24279 (2011).

45. Park, D., Morris, A. R., Battenhouse, A. & Iyer, V. R. Simultaneous mapping of transcript ends at single-nucleotide resolution and identification of widespread promoter-associated non-coding RNA governed by TATA elements. *Nucleic Acids Res.* **42**, 3736–3749 (2014).

46. Pelechano, V., Wei, W. & Steinmetz, L. M. Extensive transcriptional heterogeneity revealed by isoform profiling. *Nature* **497**, 127–131 (2013).

47. Struhl, K. Naturally occurring poly(dA-dT) sequences are upstream promoter elements for constitutive transcription in yeast. *Proc. Natl. Acad. Sci. U. S. A.* **82**, 8419–8423 (1985).

48. de Boer, C. G. & Hughes, T. R. Poly-dA:dT tracts form an in vivo nucleosomal turnstile. *PLoS One* **9**, e110479 (2014).

49. Ichikawa, Y. *et al.* Sequence-directed nucleosome-depletion is sufficient to activate transcription from a yeast core promoter in vivo. *Biochem. Biophys. Res. Commun.* **476**, 57–62 (2016).

50. Jansen, A., van der Zande, E., Meert, W., Fink, G. R. & Verstrepen, K. J. Distal chromatin structure influences local nucleosome positions and gene expression. *Nucleic Acids Res.* **40**, 3870–3885 (2012).

51. Delma, F. Z. *et al.* Genetic mutations in Cryptococcus neoformans pyrimidine salvage pathway enzymes contribute to reduced susceptibility against 5-fluorocytosine. *NPJ Antimicrob. Resist.* **2**, 24 (2024).

52. Kruglyak, L. *et al.* Insufficient evidence for non-neutrality of synonymous mutations. *Nature* **616**, E8–E9 (2023).

53. Wyman, S. K. *et al.* Synthesis errors in oligonucleotides are propagated into the

genome by homology-directed repair. *bioRxiv* 2025.07.08.663753 (2025) doi:10.1101/2025.07.08.663753.

54. Basehoar, A. D., Zanton, S. J. & Pugh, B. F. Identification and distinct regulation of yeast TATA box-containing genes. *Cell* **116**, 699–709 (2004).

55. Rossi, M. J. *et al.* A high-resolution protein architecture of the budding yeast genome. *Nature* **592**, 309–314 (2021).

56. Jin, Z. *et al.* Unraveling the regulatory dynamics of bidirectional promoters for modulating gene co-expression and metabolic flux in Saccharomyces cerevisiae. *Nucleic Acids Res.* **53**, gkaf511 (2025).

57. Zhang, C. *et al.* Yeast9: a consensus genome-scale metabolic model for S. cerevisiae curated by the community. *Mol. Syst. Biol.* **20**, 1134–1150 (2024).

58. Zhu, Y. O., Siegal, M. L., Hall, D. W. & Petrov, D. A. Precise estimates of mutation rate and spectrum in yeast. *Proc. Natl. Acad. Sci. U. S. A.* **111**, E2310–8 (2014).

59. Kemble, H., Nghe, P. & Tenaillon, O. Recent insights into the genotype-phenotype relationship from massively parallel genetic assays. *Evol. Appl.* **12**, 1721–1742 (2019).

60. Kacser, H. & Burns, J. A. The molecular basis of dominance. *Genetics* **97**, 639–666 (1981).

61. Dykhuizen, D. E., Dean, A. M. & Hartl, D. L. Metabolic flux and fitness. *Genetics* **115**, 25–31 (1987).

62. Jiang, L., Mishra, P., Hietpas, R. T., Zeldovich, K. B. & Bolon, D. N. A. Latent effects of Hsp90 mutants revealed at reduced expression levels. *PLoS Genet.* **9**, e1003600 (2013).

63. Rest, J. S. *et al.* Nonlinear fitness consequences of variation in expression level of a eukaryotic gene. *Mol. Biol. Evol.* **30**, 448–456 (2013).

64. Hartl, D. L., Dykhuizen, D. E. & Dean, A. M. Limits of adaptation: the evolution of selective neutrality. *Genetics* **111**, 655–674 (1985).

65. Hackett, S. R. *et al.* Systems-level analysis of mechanisms regulating yeast metabolic flux. *Science* **354**, aaf2786–aaf2786 (2016).

66. Li, X., Walhout, A. J. M. & Yilmaz, L. S. Enhanced flux potential analysis links changes

in enzyme expression to metabolic flux. *Mol. Syst. Biol.* **21**, 413–445 (2025).

67. Cools, H. J., Hawkins, N. J. & Fraaije, B. A. Constraints on the evolution of azole

resistance in plant pathogenic fungi. *Plant Pathol.* **62**, 36–42 (2013).

68. Todd, R. T. & Selmecki, A. Expandable and reversible copy number amplification drives

rapid adaptation to antifungal drugs. *Elife* **9**, e58349 (2020).

69. Gresham, D. *et al.* The repertoire and dynamics of evolutionary adaptations to

controlled nutrient-limited environments in yeast. *PLoS Genet.* **4**, e1000303 (2008).

70. Payen, C. *et al.* The dynamics of diverse segmental amplifications in populations of

Saccharomyces cerevisiae adapting to strong selection. *G3 (Bethesda)* **4**, 399–409

(2014).

71. Risch, T. *et al.* YgiV promoter mutations cause resistance to cystobactamids and

reduced virulence factor expression in Escherichia coli. *NPJ Antimicrob. Resist.* **2**, 33

(2024).

72. Händel, N., Schuurmans, J. M., Feng, Y., Brul, S. & ter Kuile, B. H. Interaction between

mutations and regulation of gene expression during development of de novo antibiotic

resistance. *Antimicrob. Agents Chemother.* **58**, 4371–4379 (2014).

73. Siu, L. K., Ho, P. L., Yuen, K. Y., Wong, S. S. & Chau, P. Y. Transferable

hyperproduction of TEM-1 beta-lactamase in Shigella flexneri due to a point mutation in

the pribnow box. *Antimicrob. Agents Chemother.* **41**, 468–470 (1997).

74. Sharon, E. *et al.* Functional genetic variants revealed by massively parallel precise

genome editing. *Cell* **175**, 544–557.e16 (2018).

75. Vanella, R. *et al.* Understanding activity-stability tradeoffs in biocatalysts by enzyme

proximity sequencing. *Nat. Commun.* **15**, 1807 (2024).

76. Amorosi, C. J. *et al.* Massively parallel characterization of CYP2C9 variant enzyme

activity and abundance. *Am. J. Hum. Genet.* **108**, 1735–1751 (2021).

77. Prywes, N. *et al.* A map of the rubisco biochemical landscape. *Nature* **638**, 823–828

(2025).

78. Grønbæk-Thygesen, M. *et al.* Deep mutational scanning reveals a correlation between

degradation and toxicity of thousands of aspartoacylase variants. *Nat. Commun.* **15**,

4026 (2024).

79. Beltran, A., Jiang, X. 'er, Shen, Y. & Lehner, B. Site-saturation mutagenesis of 500

human protein domains. *Nature* **637**, 885–894 (2025).

80. Bédard, C. *et al.* Most azole resistance mutations in the Candida albicans drug target

confer cross-resistance without intrinsic fitness cost. *Nat. Microbiol.* **9**, 3025–3040

(2024).

81. Tabet, D. R. *et al.* The functional landscape of coding variation in the familial

hypercholesterolemia gene LDLR. *Science* eady7186 (2025).

82. Nagalakshmi, U. *et al.* The transcriptional landscape of the yeast genome defined by

RNA sequencing. *Science* **320**, 1344–1349 (2008).

83. Luthra, I. *et al.* Regulatory activity is the default DNA state in eukaryotes. *Nat. Struct.*

Mol. Biol. **31**, 559–567 (2024).

84. Meneu, L. *et al.* Sequence-dependent activity and compartmentalization of foreign DNA

in a eukaryotic nucleus. *Science* **387**, eadm9466 (2025).

85. Hoekstra, H. E. & Coyne, J. A. The locus of evolution: evo devo and the genetics of

adaptation: The locus of evolution. *Evolution* **61**, 995–1016 (2007).

86. Brown, K. M., Depristo, M. A., Weinreich, D. M. & Hartl, D. L. Temporal constraints on

the incorporation of regulatory mutants in evolutionary pathways. *Mol Biol Evol* **26**,

2455–2462 (2009).

87. Otwinowski, J., McCandlish, D. M. & Plotkin, J. B. Inferring the shape of global epistasis.

Proc. Natl. Acad. Sci. U. S. A. **115**, E7550–E7558 (2018).

88. Sarkisyan, K. S. *et al.* Local fitness landscape of the green fluorescent protein. *Nature*

533, 397–401 (2016).

89. Wylie, C. S. & Shakhnovich, E. I. A biophysical protein folding model accounts for most

mutational fitness effects in viruses. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 9916–9921

(2011).

90. Serohijos, A. W. R. & Shakhnovich, E. I. Merging molecular mechanism and evolution:

theory and computation at the interface of biophysics and evolutionary population

genetics. *Curr. Opin. Struct. Biol.* **26**, 84–91 (2014).

91. Tokuriki, N. & Tawfik, D. S. Stability effects of mutations and protein evolvability. *Curr.*

Opin. Struct. Biol. **19**, 596–604 (2009).

92. Milind, N. *et al.* Buffering and non-monotonic behavior of gene dosage response curves

for human complex traits. *medRxiv* 2024.11.11.24317065 (2025)

doi:10.1101/2024.11.11.24317065.

93. Brachmann, C. B. *et al.* Designer deletion strains derived from Saccharomyces

cerevisiae S288C: a useful set of strains and plasmids for PCR-mediated gene

disruption and other applications. *Yeast* **14**, 115–132 (1998).

94. Gietz, R. D. & Schiestl, R. H. High-efficiency yeast transformation using the LiAc/SS

carrier DNA/PEG method. *Nat. Protoc.* **2**, 31–34 (2007).

95. Janke, C. *et al.* A versatile toolbox for PCR-based tagging of yeast genes: new

fluorescent proteins, more markers and promoter substitution cassettes. *Yeast* **21**, 947–

962 (2004).

96. Ryan, O. W., Poddar, S. & Cate, J. H. D. CRISPR-Cas9 Genome Engineering in

Saccharomyces cerevisiae Cells. *Cold Spring Harb. Protoc.* **2016**, db.prot086827

(2016).

97. Chevin, L.-M. On measuring selection in experimental evolution. *Biol. Lett.* **7**, 210–213

(2011).

98. Engel, S. R. *et al.* Saccharomyces Genome Database: advances in genome annotation,

expanded biochemical pathways, and other key enhancements. *Genetics* **229**, iyae185

(2025).

99. Amberg, D. C., Burke, D. J. & Strathern, J. N. *Methods in Yeast Genetics 2005: A Cold*

Spring Harbor Laboratory Course Manual. (Cold Spring Harbor Laboratory Press, New

York, NY, 2005).

100. Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: summarize analysis results

for multiple tools and samples in a single report. *Bioinformatics* **32**, 3047–3048 (2016).

101. Masella, A. P., Bartram, A. K., Truszkowski, J. M., Brown, D. G. & Neufeld, J. D. PANDAseq: paired-end assembler for illumina sequences. *BMC Bioinformatics* **13**, 31 (2012).

102. Rognes, T., Flouri, T., Nichols, B., Quince, C. & Mahé, F. VSEARCH: a versatile open source tool for metagenomics. *PeerJ* **4**, e2584 (2016).

103. Rice, P., Longden, I. & Bleasby, A. EMBOSS: The European molecular biology open software suite. *Trends Genet.* **16**, 276–277 (2000).

104. Seabold, S. & Perktold, J. statsmodels: Econometric and statistical modeling with python. in *9th Python in Science Conference* (2010).

105. Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat. Methods* **17**, 261–272 (2020).

106. Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *Journal of Machine Learning Research* **12**, 2825–2830 (2011).

107. Chereji, R. V., Ramachandran, S., Bryson, T. D. & Henikoff, S. Precise genome-wide mapping of single nucleosomes and linkers in vivo. *Genome Biol.* **19**, 19 (2018).

Extended Data Figures:

Extended Data Fig. 1: **The deletion of the F4 promoter fragment is clearly adaptive in 5-FC**
conditions. a, Schematics of the four 77 bp fragments into which the intergenic region between *JID1*
(YPR061C) and *FCY1* (YPR062W) was divided. There is a 26 bp overlap between F2 and F3. The
nucleosome-depleted region (NDR) of the corresponding bidirectional promoter[56,107] as well as the
TATA-like element, involved in transcription initiation at *FCY1*[40,41], are shown. **b,** Fitness of deletion
mutants in [12 µM] 5-FC. The ΔF1-F4 mutant refers to deletion of the complete intergenic region, while
Δregion instead represents the simultaneous deletion of the intergenic region and the *FCY1* gene.
Exponential growth rates and the corresponding selection coefficients (*Methods*) show that F4 is the
only promoter fragment for which deletion confers a clear growth advantage (and thus 5-FC resistance).
c, Fitness of reconstructed WT strains in [12 µM] 5-FC. Here, F2, F4 and F1-F4 refer to CRISPR
reconstruction of the WT intergenic region from the corresponding deletion mutant (*Methods*). In all
three cases, the WT 5-FC sensitivity is recovered. All growth measurements were made from three
replicate cultures of each strain.
1680

1681
Extended Data Fig. 2: The growth rate of wild-type yeast varies sharply across a range of 5-FC
and cytosine concentrations. a, Exponential growth rates for the wild-type reference strain AKD1123
across 5-FC concentrations. Two replicate cultures were assayed for each tested concentration (dots).
The line displays a fitted five-parameter logistic function (Methods), while the vertical dashed line
identifies the 12 µM concentration used in the bulk competition experiment. b, Same as panel a, but for
a range of cytosine concentrations in media lacking uracil. c, The dose-response curve obtained for 5-
FC is consistent with a previously published relationship. The curve from a is shown after conversion
into relative inhibition levels, using growth rates measured in absence of 5-FC within the same
experiment as comparison (0 relative inhibition). This curve, labeled "Exponential - This paper", was
used in downstream analyses to obtain the relative inhibition level associated with a given 5-FC
concentration. When the same growth curve data is used to compute maximum linear growth rates
(rather than exponential growth rates) and then fit a dose-response curve as in Després et al.[31], the
published relationship is replicated. d, Same as panel c, but using the cytosine growth data shown in
panel b. Here, the published relationship cannot be replicated by using the same analysis approach.
The curve, labeled "Exponential - This paper", was nonetheless used in downstream analyses to obtain
relative growth levels.

1698
Extended Data Fig. 3: Complete overview of the bulk competition experiment. a, Full experimental
design showing all 32 cultures and their relationships as transformation and/or culture replicates. Each
pool of substitutions (Sub), insertions (Ins) or deletions (Del) was constructed four times and
corresponding replicates were mixed to obtain each of the four *transformation replicates* (grey
rectangles). Each transformation replicate was used to inoculate two culture replicates, with or without
the addition of a set of previously characterized coding sequence mutations (*FCY1*; n=960)[31]. Created
in BioRender. Aubé, S. (2025) https://BioRender.com/acpk3ks. **b,** The selected coding sequence
variants cover the full range of variant effects reported in[31], both in 5-FC and cytosine (without uracil)
conditions, where Fcy1 activity is respectively toxic and essential.
1708

1709
Extended Data Fig. 4: The number and magnitude of small fitness effects observed among
promoter single-mutants are consistent with the level of noise in the bulk competition
experiment. a, Selection coefficient measurements for the same genotype typically cover an interval
of ~0.028 gen^{-1}. The distributions of maximal $|\Delta s|$, the difference between the two most extreme
selection coefficient estimates obtained across replicate cultures (n=4), are shown for all assayed
single-nucleotide substitutions (Sub; n=462), insertions (Ins; n=462) and deletions (Del; n=96) in the
FCY1 promoter. **b,** The number of mutations exceeding different thresholds of minimal fitness effect is
consistent with the expectation derived from random shuffling of the data. Across a range of minimal
$|s|$, all genotypes for which the lowest fitness estimate is above $+s$ (significant beneficial effects) or the
highest estimate is below $-s$ (significant deleterious effects) are shown (dots). The solid line and shaded
area display the median and associated 95% confidence interval for the number of significant effects
(using the same definition) obtained from random permutations of the data (n=10 000). Within each
iteration, 1020 mock single mutants were assembled by randomly combining selection coefficients from
each of the four replicate cultures (Methods). **c,** Twelve promoter single-mutants display $|s| > 0.01$ in
the bulk competition experiment. For each genotype, all replicate s estimates are shown (dots). The
identifiers of the mutations (x axis) inform on their type (deletion, insertion or substitution), position
relative to the *FCY1* coding sequence (where 0 is the base immediately upstream of the gene), and
nucleotide change. **d,** The small fitness effects measured are not clearly related to corresponding
changes in *FCY1* expression level. For mutants included in panel **c,** the median selection coefficient
estimate is compared with WT-relative expression level estimated from sort-seq (see **Fig. 4e**). Whereas
the correlation is negative as expected, the second highest fitness effect is observed for a mutation
increasing expression (>1) and the most negative selection coefficient is associated with a moderate
decrease of *FCY1* expression.

1733
1734

1735
Extended Data Fig. 5: Single-nucleotide substitutions in the *FCY1* coding sequence readily confer
resistance to 5-FC, across experiments. **a,** Comparison of median selection coefficients for coding
sequence mutants (n=960) added to the bulk competition experiment with previously published variant
effect scores measured in the same conditions[31]. Values have been aggregated into medians for all
genotypes which result in the same amino acid sequence, such that n=311 points are shown. A
LOWESS fit highlights a mostly monotonically increasing trend, as shown by a high Spearman
correlation ($p < 10^{-64}$). **b,** The set of coding variants which confer resistance to 5-FC is consistent
across the two experiments. All mutants which were added to the "+*FCY1*" samples (**Extended Data**
Fig. 3a) were classified into resistant (R), and sensitive (S) variants according to their effect in 5-FC
conditions, using Gaussian mixture models (*Methods*). This was done using either the FC score
reported by[31] (top) or the selection coefficients inferred in the current study (left). The confusion matrix
(right) shows that, among all variants which were classified as resistant using data from[31], none appear
sensitive to 5-FC in the current study. As in the previous panel, mutated sequences were collapsed into
amino acid substitutions resulting in 311 variants. **c,** Almost a third of single-nucleotide substitutions in
FCY1 confer resistance to 5-FC. The distribution of variant effect scores from[31] for all single-nucleotide
substitutions in *the coding sequence*. For each genotype (n=1411), the score reported for the
corresponding amino acid variant was used. Only 20 of the possible nucleotide substitutions are
missing, mostly in the first and last codons. Variants are identified as resistant (R) or susceptible (S)
according to the classification shown in panel **b**. The "R" class includes ~29% (408/1411) of the single
substitutions shown.
1756

1757
Extended Data Fig. 6: **Validation of the 42bp-mEGFP construction. a,** Medians of normalized
fluorescence intensity, as measured by flow cytometry (*Methods*), for three mEGFP insertions at the
genomic *FCY1* locus. Each point is a distinct clone of the corresponding strain, except for the WT. In
the 42bp-mEGFP construct, which was used in the current study to measure promoter activity, the
mEGFP green fluorescent protein is fused in C-terminal of the first 42 bp of the *FCY1* coding sequence.
In contrast, FCY1-mEGFP designates a C-terminal fusion with full-length *FCY1,* while mEGFP stands
for the insertion of the fluorescent protein in place of the *FCY1* coding sequence. **b,** As on the left, but
showing the full distribution of log-scaled normalized fluorescence intensities for each clone (5000
events). **c,** Comparisons of median WT-relative fluorescence intensities for the different promoter
constructs, in the 42bp-mEGFP or FCY1-mEGFP backgrounds. In each case, the WT-relative value
has been computed relative to the corresponding reference (WT *FCY1* promoter sequence, upstream
of either the 42bp-mEGFP or FCY1-mEGFP construct). The ranks of the promoters are unchanged
between the two backgrounds, although WT-relative fluorescence intensities are consistently higher
when the fluorescent protein is fused to full-length *FCY1*. Values shown for the 42bp-mEGFP are the
same ones which were used throughout the paper to draw expression-fitness curves, although replicate
measurements obtained from the same clone were aggregated into grand medians. The FCY1-mEGFP
strains had not been validated prior to cytometry. All clones which displayed a fluorescence intensity
clearly different from the corresponding WT are shown. Because the promoter cassettes were directly
inserted in place of the endogenous promoter in the WT background, the WT expression level (and thus
fluorescence intensity) is expected in cases where integration failed (*Methods*).

1778
Extended Data Fig. 7: **Indels before and after the second methionine codon of *FCY1* have**
contrasting effects on expression level. a, Comparison of relative expression levels for the four indel
mutations which were included among the set of validation mutations, classified according to whether
they occur before or after the second methionine codon of *FCY1*. Cytometry measurements from
reconstructed mutants as well as estimates obtained from the large-scale sort-seq experiment are
shown. **b,** Relative expression levels, as estimated from the sort-seq experiment, for all detected indel
mutations in the *FCY1* coding sequence. Those occurring after the second methionine are clearly more
impactful. **c,** Median effect of insertion and deletion variants detected in the sort-seq (top) and bulk
competition (bottom) experiments. The shaded area encompasses the three bases of the second
methionine codon. Mutations occurring before this codon were recovered at a much lower frequency
from the bulk competition, which is consistent with the observation that they do not completely abolish
FCY1 expression and thus do not confer resistance.
1791

1792

Extended Data Fig. 8: Deletion of putative TF binding sites is not more impactful than single-nucleotide mutations in the same regions. a, Deletion of positions -133 to -127 of the *FCY1* promoter, corresponding to site 1 (positions -132 to -127) of **Fig. 4e**. The WT-relative fluorescence of the deletion mutant, from three replicate measurements in flow cytometry, is shown as an horizontal line whose width displays the minimum and maximum values among the three corresponding medians. Among reconstructed validation mutants (left), one single-nucleotide substitution decreases *FCY1* expression as much as the deletion, as shown by its median WT-relative fluorescence intensities (dots; grand median of three replicate cytometry measurements). Among all mutations assayed by sort-seq (right), the most impactful single-nucleotide changes decrease expression as much as the deletion, as shown by the corresponding relative expression level estimates (dots; median of the 95% prediction interval obtained from sort-seq calibration). **b,** Deletion of positions -125 to -118, corresponding to site 2 (-125 to -120) of **Fig. 4e**. The plots are the same as in **a**, but this time for a deletion which increases *FCY1* expression. Cytometry measurements of validation mutants (left) and sort-seq estimates (right) both show that many single-nucleotide changes have an effect that is comparable to that of the deletion – or even larger.

1808

Extended Data Fig. 9: The insensitivity of fitness to changes in *FCY1* expression level is constant
across many concentrations of 5-FC and cytosine. a, Fitness functions in 5-FC. In [12 μM] 5-FC
(left), a constant plateau of low fitness is reached starting from ~0.18 WT-relative expression (same as
Fig. 5b). Across three additional concentrations of 5-FC (right), the boundary of the plateau does not
change. Its height however varies, such that lower growth rates are observed for increasing 5-FC
concentrations. **b,** Fitness functions in media lacking uracil, where deamination of cytosine by Fcy1p is
essential. When cytosine is added at [63 μM] (left), a similar plateau is observed from ~0.18 WT-relative
expression, such that any further increase in *FCY1* expression does not increase growth rate. When
three additional concentrations of cytosine are tested (right), the boundary of the fitness plateau also
remains constant. Error bars around each dot display minima and maxima of three replicate
measurements for growth rate and relative fluorescence intensity for each combination of promoter and
5-FC/cytosine concentration. In addition, dashed vertical lines identify the WT expression level (relative
fluorescence intensity of 1.0).

1822

Extended Data Fig. 10: The expression-fitness function of *FCY1* displays a sign inversion between 5-FC and cytosine conditions. a, Fitness functions across 5-FC and cytosine concentrations covering the full ranges of the corresponding dose-response curves (**Extended Data Fig. 2**). Exponential growth rate measurements were performed in duplicates for each of the promoter constructs which define the elbow of the fitness function: pGAL-L, pCYC1, pCUP1 and pTEF1. In 5-FC (left), the conditions shown, from the lowest to the highest concentration, respectively result in relative inhibition levels of 0.20, 0.29, 0.40 and 0.69. In cytosine (right), the concentrations shown are respectively associated with the following relative growth levels: 0.19, 0.64, 0.93 and 0.97. **b,** Comparison of fitness functions obtained in 5-FC (left) and cytosine (right**)** concentrations resulting in matching levels of relative inhibition or growth for the WT reference strain (**Extended Data Fig. 2**). Although growth does not reach 0 in cytosine conditions when *FCY1* expression is decreased to very low levels, the two pairs of curves display very similar shapes. Same data as in panel **a**, but converted into relative inhibition or growth values using the relevant dose-response curve. **c,** Increasing *FCY1* expression through single mutations in the promoter does not provide a benefit in cytosine conditions,

where Fcy1 activity is essential for growth. Exponential growth rate measurements were performed in
triplicates for eight reconstructed promoter mutants covering the full range of expression changes
available from single-nucleotide changes (**Fig. 5d**). From the lowest to the highest, the three cytosine
concentrations used respectively result in relative growth levels of 0.42, 0.64 and 0.97. The dashed
horizontal lines display the medians of growth rates obtained for the two mutants closest to the WT
expression level. Across all three concentrations, the mutant associated with the highest expression
level is not clearly fitter than this baseline. This confirms that, just like they are not deleterious in 5-FC
conditions, promoter mutations increasing *FCY1* expression are not beneficial in cytosine conditions.