Choroid Plexus Hosts the Earliest Detectable Circadian Clock in the Brain

Hélène Vitet [1], Vuong Hung Truong [1], Jihwan Myung [1,2,*]

[1] Laboratory of Braintime, Graduate Institute of Mind, Brain and Consciousness (GIMBC), Taipei Medical University, New Taipei City 235, Taiwan
[2] Graduate Institute of Medical Sciences (GIMS), College of Medicine, Taipei Medical University, Taipei 110, Taiwan

* Correspondence: Jihwan Myung (jihwan@tmu.edu.tw)

Abstract

Embryonic circadian rhythms support normal development, and maternal chronodisruption is linked to miscarriages and neurodevelopmental risk, yet the identity of the earliest brain circadian oscillator and the mechanisms governing its emergence remain unknown. Using bioluminescent reporters for *Per2* and *Bmal1*, we identified the developmental window in which autonomous ~24 h oscillations first appear *ex vivo*. Contrary to the prevailing view that places the suprachiasmatic nucleus (SCN) as the earliest circadian structure at E14.5–E15.5, we found that the fourth ventricle choroid plexus (4VCP) exhibits PER2::LUC oscillations by E11.5–E12.5. Transcriptional profiling revealed significant phase shifts in multiple clock genes between E11.5 and E13.5, indicating a transition away from maternal synchrony and the onset of a differentiation program. Detailed characterization of these early oscillations revealed hallmarks consistent with a saddle-node on an invariant circle (SNIC) bifurcation. Our findings establish the 4VCP as the earliest detectable brain circadian oscillator and clarify the process leading to the onset of the embryonic circadian clock in the brain. These results provide insights into maternal and fetal clock interactions during development, with potential implications for neonatal care.

Keywords: circadian clocks, choroid plexus, embryonic development, SNIC bifurcation, clock emergence, maternal synchrony

30
31 **Graphical Abstract:**
32

33
34
35 **Timeline of 4VCP clock emergence before the SCN.** In the embryonic 4VCP, PER2::LUC oscillations appear
36 at E11.5–E12.5, and cellular synchrony increases abruptly around E14.5. *Bmal1-ELuc* oscillations appear at
37 E13.5. The 4VCP entrains to a 0.5 °C temperature cycle through E16 but not at E18. The lower pie charts and
38 gradients summarize the fraction of embryonic clock genes that are circadian and their correlation with
39 maternal *Bmal1*, *Dbp*, and *Per2* expression.
40

41   **Introduction**
42
Circadian clocks are fundamental pacemakers that align physiology with the 24 h rhythms of Earth.
Physiological processes occur within a circadian timescale autonomously maintained by the
transcription-translation feedback loop (TTFL). In this loop, the positive feedback component
BMAL1:CLOCK activates transcription of the core negative feedback components such as *Per* and *Cry*,
and the resulting PER:CRY complexes accumulate with a delay before repressing their own
transcription upon translocation. While most mammalian tissues express circadian clocks, the
suprachiasmatic nucleus (SCN) in the hypothalamus acts as the central clock, synchronizing other
clocks throughout the body. Although mature circadian systems are increasingly well characterized, the
timing and locus of the emergence of circadian rhythms during development remain poorly defined.
52
When? Circadian rhythms have been observed in prenatal stages in humans, non-human primates, and
rodents (Maeda et al., 2006; Serón-Ferré et al., 2012; Landgraf et al., 2014). However, detecting
embryonic clocks presents technical challenges, as experimental findings are often inconsistent due to
low-amplitude oscillations and tissue-specific differences in clock gene expression. Although *Per* and
Bmal1 did not show clear rhythms when measured in whole-embryo *ex vivo* preparations (E10 to E17)
(Cao et al., 2022; Dolatshad et al., 2010; Serón-Ferré et al., 2012), the circadian rhythmicity of PER2 has
been detected in the SCN *ex vivo* around E14.5 to E15.5 (Wreschnig et al., 2014; Carmona-Alcocer et al.,
2018) and around E18 in peripheral organs such as the kidney (Dan et al., 2020), the heart (Umemura et
al., 2017), and the liver (Dolatshad et al., 2010).
62
Embryonic brain morphogenesis depends on a complex interplay between the neural tube and the
cerebrospinal fluid pressure gradients. The dynamic homeostasis of cerebrospinal fluid (CSF), produced
primarily by the choroid plexus (CP), must be tightly regulated during this process. This non-neuronal
tissue performs two roles: the CP filters the blood through its epithelial cells to produce CSF, acting as a
gatekeeper at the fluid boundary of the brain, and it releases signaling molecules (5-HT, miRNAs) as
well as peptides and proteins (TTR, cytokines, IGF2) into the CSF to support processes such as
metabolism and signal transduction (Saunders et al., 2023; Chau et al., 2015; Courtney et al., 2025;
Edelbo et al., 2023). In the embryo, the onset timing of CP epithelial CSF production remains unclear, in
part because early in development the CP is not the only structure that modulates CSF composition
(Bueno et al., 2020; Parvas et al., 2008). CSF is present as early as neural tube closure at E9.5, initially
derived from the enclosed amniotic fluid. Its composition then transitions to CP-produced fluid
between E10.5 and E14.5, coinciding with CP formation (Chau et al., 2015).
75
Where? Little is known about the exact locus in the brain where the earliest circadian clock emerges.
Using bioluminescent reporters of circadian gene expression, PER2::LUC and *Bmal1-ELuc*, we scanned
ex vivo brain slices for circadian oscillations and monitored tissue explants collected from E9.5 onward
throughout embryogenesis. This screening identified the 4VCP as the earliest locus with detectable
circadian clock oscillations. We recently found that, in adult mice, the CP acts as a strong clock, as
indicated by the persistence and amplitude of PER2::LUC reporter oscillations and by its ability to
entrain the SCN (Myung et al., 2018). The CP is present in each of the brain ventricles, which are filled

with the CSF. The lateral ventricle CP (LVCP) is contiguous with the third ventricle CP (3VCP) and
separate from the fourth ventricle CP (4VCP) in the hindbrain. Differences among CPs include their
developmental timelines and transcriptomes (Saunders et al., 2023; Lun et al., 2015a; Dziegielewska et
al., 2001). In mice, the 4VCP develops first, followed by the LVCP (Lun et al., 2015b). Arising from the
lower rhombic lip, between the roof plate and dorsal neuroepithelium, 4VCP development begins as
early as E9.5, when it expresses *Wnt1*, transthyretin (*Ttr*), and potassium voltage-gated channel
subfamily E member 2 (*Kcne2*). Development peaks between E11 and E12.5 (Lun et al., 2015b; Hunter
and Dymecki, 2007), after which the 4VCP becomes anatomically apparent and is thought to be
complete by E14, although proliferation continues postnatally (Lun et al., 2015b). At this stage, 4VCP
cells are mature and express *Ttr, Otx2,* and *Wnt5a*, among other genes.

Embryonic CSF is crucial for the survival, proliferation, and differentiation of neural progenitors (Miyan
et al., 2003; Martin et al., 2009; Gato et al., 2005; Parada et al., 2008; Alonso et al., 2011 and 2017;
Huang et al., 2009) by delivering peptides and molecules, such as morphogens (retinoic acid, Sonic
Hedgehog) and growth or trophic factors, such as FGF2 (Martin et al., 2006), EGF (Birecree et al., 1991),
and IGF (Zappaterra and Lehtinen, 2012). CSF is also known for its role in shaping the brain through
pressure against neuroepithelial walls (Gato and Desmond, 2009). The strong clock in the CP is thought
to regulate this CSF secretome, which is sensitive to time of the day (Edelbo et al., 2023). Very recently,
apocrine secretion from the CP to the CSF has been identified as a cause of neuronal progenitor
stimulation and offspring behavior modification (Courtney et al., 2025). However, the embryonic CP
clock remains poorly characterized, although it could modulate the temporal composition of CSF and,
consequently, neurogenesis.

How?   Circadian clocks are present across many mammalian cell types (Balsalobre et al., 1998;
Yamazaki et al., 2000; Yoo et al., 2004), yet the locus-specific sequence of its emergence in the brain
suggests a programmed onset of circadian oscillations (Wreschnig et al., 2014; Carmona-Alcocer et al.,
2018). Developmentally, the early embryo operates on distinct timescales, with cell divisions occurring
on the order of hours before lengthening during neurogenesis, and somitogenesis proceeding on a
short timescale of hours (Caviness et al., 1995; Oates et al., 2012). These rhythms may not synchronize
well with the circadian clock, especially as the cell division timescale changes gradually. An abrupt
transition between the somitogenesis era and the circadian era has been proposed (Yagita *et al.*, 2024),
making the sudden emergence of the circadian clock *in vivo* plausible.

In nonlinear dynamics, a stable equilibrium can transition to a stable oscillation through mechanisms
called bifurcations (Strogatz, 2024). Several types of bifurcation are distinguished by how amplitude
and period behave near threshold and by their robustness to perturbations. Many oscillating biological
systems, including circadian oscillators, are described by a (supercritical) Hopf bifurcation,
characterized by a gradual increase in oscillation amplitude while the period stays near 24 h. The Hopf-
based oscillator often fits dynamics of circadian clocks, consistent with experimental and modeling
work (Murayama et al., 2017; Battogtokh and Tyson, 2018). Similarly, adult sinoatrial node pacemaking
is typically governed by Hopf-type mechanisms, whereas the developmental onset of the first
heartbeat in zebrafish arises via a noisy SNIC bifurcation (Jia et al., 2023). The saddle-node on an

invariant circle (SNIC), also called SNIPER, is characterized by the abrupt appearance of nonzero
amplitude and an initially long period that shortens toward ~24 h. At the single-cell level in the 4VCP,
we observe a near-step increase in amplitude at onset and a progressive shortening of period over
development, hallmarks consistent with SNIC-type dynamics. This bifurcation has specific implications
in maternal-fetal coupling. Near a SNIC threshold, slow dynamics make oscillation frequency highly
sensitive to weak biases. Consequently, weak maternal cues can efficiently entrain the nascent CP clock
and synchronize weakly coupled CP cells by common drive. We observed such behavior at E14 in *ex vivo*
4VCP explants subjected to low-amplitude rhythmic temperature input.
133
In this study, we pinpoint when circadian rhythms first emerge in the 4VCP, identifying it as the earliest
detectable clock in the embryonic mouse brain. We find hallmarks consistent with discrete appearance
of the circadian clock, including SNIC-like bifurcation behaviors and abrupt synchronization over
development. These results clarify the timing, locus, and onset dynamics of the brain's first clock that
potentially governs fluid homeostasis and suggest implications for maternal chronodisruption,
including shift work, during early pregnancy that can elevate risks of miscarriages and preterm birth
(Cai et al., 2019; Davari et al., 2018), and for circadian support in neonatal care settings such as the
Neonatal Intensive Care Unit (NICU) where preterm infants often show impaired rhythmicity (Govindan
et al., 2024).
143

144 **Material and Methods**
145
Mice

Homozygous PER2::LUC mice (B6.129S6-*Per2*^{tm1Jt}/J, RRID: IMSR_JAX:006852) and heterozygous
Bmal1-ELuc mice (Nakajima et al., 2010; a gift from Daisuke Ono, Nagoya University), maintained on
the C57BL/6J background, were crossed with wild-type C57BL/6J animals to generate heterozygous
PER2::LUC$^{+/-}$ or *Bmal1-ELuc*$^{+/-}$ embryos. The PER2::LUC fusion protein reporter tracks PER2 abundance
through a luciferase reaction (Yoo et al., 2004). *Bmal1* transcriptional activity was monitored with the
Bmal1-ELuc reporter, with high baseline removed through Hodrick-Prescott filter-based detrending
(Myung et al., 2012). In this study, we treat *Bmal1-ELuc* as a transcript-level proxy and PER2::LUC as a
protein-level proxy, as its luciferase activity peaks several hours after *Per2* transcription (Myung et al.,
2025). Both females and males were maintained under equinox light-dark (LD) cycles (12 h light:12 h
dark; lights on at 07:00 and off at 19:00), with light-dark schedules controlled by LocoBox software
(Truong and Myung, 2023) and constant ambient temperature and humidity conditions (24.0 ± 1.8°C;
50.2 ± 8.7% RH). For sampling at ZT14 and ZT20, we used a reverse dark-light (DL) cycle (lights on at
22:00 and off at 10:00) and validated that gene expression levels matched those under LD cycles at
E13.5 (**Fig. S1**). All animal protocols were reviewed and approved by the Institutional Animal Care and
Use Committee (IACUC) of Taipei Medical University Laboratory Animal Center (protocols LAC-2019-
0430, LAC2022-0381, LAC2022-0477, LAC2023-0037, and LAC2023-0038).
163
Tissue isolation and explant culture

Direct dissection. Tissue culture and bioluminescence recording protocols were identical to those used
previously (Myung et al., 2018). Pregnant female mice were anesthetized with isoflurane and
euthanized at defined Zeitgeber times, and brains and embryos were rapidly removed into ice-cold
Hank's balanced salt solution (HBSS; Sigma-Aldrich), supplemented with penicillin-streptomycin
(10,000 U/mL; Gibco) and HEPES (1 M, pH 7.4). Embryonic tissues were dissected after decapitation
using forceps and a scalpel under a stereomicroscope. For E9.5, the prosencephalon (containing both
the SCN and LVCP progenitors) and the rhombencephalon (containing the 4VCP progenitors) were
collected. For E10.5–E12.5, the 4VCP was extracted from the inferior part of the hindbrain roof plate
epithelium, i.e., the lower rhombic lip containing rhombomere 2-8, located between the otic vesicles
and the posterior part of rhombomere 1 (Hunter and Dymecki, 2007). For the LVCP, the cortical midline
was considered to host the LVCP progenitors. After E12.5, the CP becomes clearly distinguishable, and
its dissection is similar to that of the adult mouse. The precision of 4VCP isolation throughout
development was validated by RT-qPCR (**Fig. S2**). The SCN from E10.5 to E13.5 was dissected from the
ventral hypothalamus between the two optic cups (developing eyes). After E13.5, the SCN and the optic
chiasm become more distinct, and the dissection procedure is the same as in the adult. This protocol
was also used to isolate maternal and embryonic tissues in ice-cold PBS for biomolecular experiments.

Placenta dissection. We separated the light pink outer layer of the placenta (maternal decidua,
considered here as the maternal placenta) from the darker pink inner layer (junctional zone and
labyrinth, considered here as the embryonic placenta). The weak junction between these two sides of
the placenta allows dissection with tweezers.

Tissue slicing. Heads or brains of mouse embryos at E11.5, E13.5, or E15.5 were sliced *ex vivo* on a
vibratome (VT1000S, Leica Biosystems). Tissue was briefly dipped in two successive baths of low-
melting point agarose (2% w/v; CAS 9012-36-6) at 37 °C, then embedded in an agarose block, glued to
the vibratome stage and immersed in ice-cold HBSS. Using a diamond blade (Delaware Diamond
Knives) advancing at ~0.025 mm/s and maximum frequency, 300 μm-thick slices were made and
transferred to a culture membrane.

Explant culture. Each explant was cultured on a culture membrane insert (PICM0RG50, Merck
Millipore) in a 35-mm dish containing 1 mL DMEM (Sigma-Aldrich) supplemented with B-27 (Gibco) and
200 μM beetle luciferin (Promega). The dish lid was sealed with high-vacuum grease (Molykote,
DuPont) to prevent evaporation and maintained at a constant temperature of 37.00 ± 0.01°C in a
hybrid refrigerator-heater incubator (MIR-154-PA, PHCbi). Temperature was monitored by placing an
iButton (DS1921L, Maxim Integrated) adjacent to the dishes.

iDISCO tissue clearing and immunofluorescence labeling

After culture, the membrane supporting the brain slices was cut out and cleared using the iDISCO
protocol (https://idisco.info/idisco-protocol). Briefly, after fixation in 4% PFA, the slices were rinsed and
incubated in PBS for 3 days, then dehydrated in MeOH/H_2O in micro glass vials with PTFE caps,
followed by an overnight incubation in 66% DCM/33% MeOH. Samples were then washed and bleached

in 5% H_2O_2 overnight, before being rehydrated in MeOH/H_2O. Immunolabelling was conducted in a 96-
well plate, starting with permeabilization (1 day) and blocking (1 day) at 37°C, followed by incubation
with primary antibody TTR (AB_10723946; GTX85112, chicken, 1:500, Genetex) for 2 days at 37°C. Slices
were then washed and incubated with secondary antibody (anti-chicken Alexa Fluor 568; AB_175477;
ab175477, goat, 1:1000, Abcam) for 2 days at 37°C. Similar steps of wash and blocking were applied
before consecutive incubations with LHX1 (AB_531784; 4F2-c, mouse, 1:500, DSHB) and anti-mouse
CF350 (SAB4600222, goat, 1:1000, Sigma) as before. Slices were then transferred back to the micro
glass vials to undergo final clearing, beginning with dehydration in MeOH/H_2O followed by a 3-h
incubation in 66% DCM/33% MeOH at room temperature. Finally, two 15 min incubations in 100% DCM
were performed before mounting the slices on glass slides. DBE was added directly over the slices,
confined under a coverslip, and sealed with nail polish. Slices were then imaged using a DM IL
fluorescence microscope (Leica Microsystems).
218
Time-lapse bioluminescence luminometry and imaging

Bioluminescence luminometry. For luminometry, the overall bioluminescence from each sample was
recorded every 15 min with an integration time window of 108 s per dish, using a 32-dish rotating wheel
with four photon-counting photomultiplier tubes (PMTs) (LumiCycle 32, Actimetrics), housed in a
refrigerator-heater incubator for precise temperature control (MIR-154-PA, PHCbi). Temperature was
recorded continuously throughout the experiment.
225
Bioluminescence imaging. Single-cell imaging was performed on a custom-built incubator-microscope
system equipped with the Orca R2 cooled CCD camera (Hamamatsu) with water cooling, combined
with a 4x, NA 0.16 objective (UPLXAPO, Olympus) coupled to a 0.35x reduction lens (U-TV0.35XC-2,
Olympus) to increase photon collection from bioluminescence (Myung et al., 2024), with a stage-top
incubator (INUG2-WELS, Tokai Hit). Binning was set to 4×4, and exposure time was 60 min. Images
were acquired using Micro-Manager 1.4 (Vale Lab, UCSF) via the DCAM-API (Hamamatsu) driver.
232
Real-time quantitative qPCR (RT-qPCR)

RNA Isolation, cDNA synthesis, and RT-qPCR. After dissection of the embryonic or maternal 4VCPs at
ZT2, ZT8, ZT14, and ZT20, the samples were stored at -80°C. Embryonic 4VCPs from one dam (3-11
embryos per litter) were pooled, and at least 2-3 litters per time point were used. RNA sampling and RT-
qPCR methods followed our previous protocol (Myung et al., 2015). 300 µL of TRK lysis buffer
containing 2% β-mercaptoethanol was added to each sample, followed by an equal volume of
chloroform after intensive homogenization. After phase separation and ethanol addition,
microcolumns from the ENZA MicroElute Total RNA Kit (Omega Bio-Tek) were used to collect total
RNA, with a final elution volume of 21 µL in DEPC-treated water. After evaluating the RNA
concentration, cDNA was synthesized from 0.5–1 µg of total RNA using SuperScript IV Reverse
Transcriptase (Thermo Fisher Scientific) with oligo(dT)$_{20}$ primers. PowerUp SYBR Green Master Mix
(Applied Biosystems) was added to the cDNA samples, and RT-qPCR was performed in technical

triplicate on a QuantStudio 3 Real-Time PCR System (Thermo Fisher Scientific). Primer sequences
tested and used are listed in **Table S1**.

Normalization. *Gapdh*, *Actb* (*β-actin*), and *Rpl13a* served as internal controls for embryonic samples
(Hildyard et al., 2022). G*apdh* and *Actb* were used for maternal samples. For embryonic samples,
differentiation status of each gene was assessed as the relative abundance of mRNA at a given
embryonic stage (*n* > 8) normalized to the maximum abundance observed between E9.5 and E16.5. For
example, at E9.5:

$$RNA_{E9.5} \, (\%) = \frac{mean \, (RNA_{E9.5})}{max \, (RNA_{E9.5-E16.5})} \times 100 \qquad (1)$$

Spectral analysis of bioluminescent time series and time of emergence quantification

Preprocessing. Visualization and preprocessing, including periodogram and fast Fourier transform
(FFT)-based spectral analysis, were performed on Mathematica 13 (Wolfram Research) using the
custom PMTAnalysis package (http://sourceforge.net/projects/imaginganalysis). Before analysis, the
first 6 h of each recording were truncated. All circadian periods of the bioluminescence oscillations were
estimated using FFT, and amplitudes were quantified as the root mean square (RMS) of detrended
oscillations over fixed analysis windows: nominally 7 days for luminometry and 4VCP imaging, and 4
days for whole brain slice imaging. Because the first 6 h were removed, some recordings were slightly
shorter than 7 days. For image analysis of brain slices or the 4VCP, a 2×2-pixel grid was applied to the
time-lapse images after removal of outlier pixels, and each ROI's standard deviation, center of mass,
and mean gray value were measured using ImageJ (NIH).

Circadian power. Using a sliding-window FFT (2-day for luminometry data and 3-day for imaging data),
the time-lapse bioluminescence signals from PER2::LUC or *Bmal1-ELuc* were decomposed into spectral
power and visualized as spectrograms over time. After integrating the power within specific period
bands, the circadian power was calculated as the ratio of the total power in the circadian range (24 ± 4
h) to the total power across the full period range (3–58 h) (**Fig. S3**). This yielded a time series of percent
circadian power, which quantifies how strongly a gene oscillates within the circadian band relative to
other period bands. Finally, the overall circadian power was computed as the average of these window-
wise circadian power values across the entire recording. Circadian power thus provides a metric that
captures the robustness of the circadian rhythm and accounts for damping, period variability, and
noise. For a perfect sinusoid in a 2-day window, the maximal circadian power reaches ~60%, whereas
for transient, rapidly damped oscillations, it falls below 20% (**Fig. S3**). The blank control (luciferase
medium without tissue explant) showed a maximal circadian power of ~20% (*n*=10) (**Fig. 2D**). For SCN
and 4VCP explants, the half-maximal level of the fitted circadian power curve across development was
~30% (30.83 ± 2.36% and 31.28 ± 1.70%, respectively).

Emergence Time. For rigorous identification of the emergence time points of the PER2::LUC and
Bmal1-ELuc circadian rhythms, we performed nonlinear curve fitting using a sigmoid function. Blank

recordings were used to estimate baseline circadian power and to provide null data points prior to E9.5.
We then defined the emergence time (T_e) as the time when one-half of the maximum circadian power
was reached (eq. 2).

287
$$P(t) = \frac{A}{1+e^{-(t-T_e)/\tau}} + C_0 \qquad\qquad (2)$$

288   where $P(t)$ is the circadian power at time t (time of sampling for explant), A is the amplitude above
289   baseline, T_e is the half-maximal emergence time, τ is the time constant (slope parameter), and C_0 is the
290   baseline determined from blank recordings, with maximal power given by $A + C_0$.
291

Hierarchical clustering and spectral analysis of qPCR gene expression profile

Hierarchical clustering. To ensure comparability across genes, expression levels were first normalized
using the internal control genes, then standardized using Z-scores: for each gene, the mean normalized
expression was subtracted from each time point and the result was divided by the standard deviation
across all time points. To explore temporal relationships between genes, we used hierarchical clustering
as implemented in the seaborn.clustermap function (v0.11.2) in Python (v3.6.13). This function
computes pairwise distances between gene expression profiles using the Pearson correlation r_{ij} as the
distance metric between genes i and j. Clustering was performed using average linkage, and results
were visualized as a cluster heat map with a dendrogram that represents the similarity relationships
among genes based on their temporal patterns. Average linkage distance $D(A, B)$ was calculated as:
302
$$D(A, B) = \frac{1}{m\,n}\sum_{i\in A}\ \ \sum_{j\in B}\quad d(i, j) \qquad\qquad (3)$$

where m and n are the numbers of genes in clusters A and B, respectively, and $d(i, j)$ is the distance (1 -
r_{ij}) between genes i and j. To summarize clusters, we identified representative genes for each cluster.
For a given cluster, we computed its centroid by averaging the Z-scored expression profiles of all
member genes across time points. We then calculated the Pearson correlation between each gene in
the cluster and the cluster centroid. The gene with the highest correlation to the centroid was
designated as the representative gene for that cluster, as it best reflected the average temporal pattern
of its cluster.
310

Synchrosqueezed continuous wavelet transform (SST-CWT). Circadian gene expression profiles in
embryos are sparsely sampled, and conventional Fourier or continuous wavelet transforms (CWT) often
yield broad, smeared power spectra when signals have non-stationary components. SST-CWT sharpens
time-frequency estimates by reassigning CWT coefficients from a range of supporting frequencies to a
more precise instantaneous frequency (Daubechies et al., 2011). We implemented SST-CWT using the
cwt_sst   function   from   the   squeezepy   library   (v0.6.5;
https://github.com/OverLordGoldDragon/ssqueezepy). Prior to analysis, normalized gene expression
time series were detrended with a moving average filter to reduce baseline drift. The signal was then
mirrored and extended to improve edge behavior during the wavelet transform. We used a Morlet
wavelet and 90 voices per octave. Although SST-CWT yields an amplitude spectrum, we refer to it here
as a "power spectrum" to align with common usage in circadian biology (true power can be obtained by

squaring the amplitude). From the SST-CWT output, we retained periods from 3 to 58 h and calculated
circadian power at each time point using the same 24 ± 4 h band as in the PER2::LUC and *Bmal1-ELuc*
spectral analysis. This definition enables direct comparison between bioluminescence and qPCR-
derived rhythmicity.
326
Phase dynamics and visualization. We estimated instantaneous phase (radians) along ridges of the
time-frequency representation. Instead of plotting phase alone, we generated phase-weighted
spectrograms, multiplying SST-CWT amplitude by its corresponding phase angle to highlight dominant
oscillatory components with phase information. To improve visual clarity, colormap normalization and
adaptive contrast scaling were applied using the maximum absolute value of each map. An illustration
of this approach using simulated data is shown in **Fig. S6A**.
333
Circular phase plots. To track phase across development (E10.5, E12.5, E14.5, E16.5), we constructed
circular phase plots based on the estimated instantaneous phase within the circadian range (20–28 h).
For each embryonic cluster, the representative gene was plotted clockwise in a phase plot resembling a
conventional clock, with 0 rad positioned at 12 o'clock. Non-representative genes from the same cluster
were plotted with smaller, semi-transparent markers for comparison. The left semicircle was shaded to
distinguish early vs late phases. Marker shape and color indicated cluster identity, and genes with
circadian power ≥ 20% were outlined in black. To assess potential maternal-embryonic coordination,
dual-radius phase plots were used: the inner ring showed embryonic cluster representatives, and the
outer ring showed selected maternal genes. Maternal-embryonic pairs were connected with dotted
lines when both exhibited circadian power ≥ 20% and their phase difference was $< \pi/2$ rad (≤ 6 h),
indicating possible phase alignment.
345
Sliding-window correlation and network construction. To assess temporal coordination among
maternal and embryonic genes, time series were padded by duplicating the first and last two points,
and 5-timepoint sliding windows were applied. Pearson correlation matrices were computed for each
window, generating a time-resolved sequence of correlations. Examples of dynamic correlation are
shown in **Fig. S6B**. Networks were built in NetworkX (v2.5.1), with genes as nodes and edges weighted
by average correlation within a developmental period. Edge consistency across periods was shown with
solid lines, and changes in correlation sign with dashed lines. For maternal-embryo networks, only the
top 30% strongest correlations were highlighted; the remainder were rendered in semi-transparent
gray.
355

356   **Results**
357
PER2::LUC circadian oscillations emerge in the 4VCP before the SCN
The embryonic SCN is widely recognized as the earliest reported brain region to exhibit a detectable
circadian rhythm, which emerges between E14.5 and E15.5 (Wreschnig et al., 2014; Carmona-Alcocer et
al., 2018). To determine whether the SCN clock is unique at this developmental stage, we prepared

brain slices from E15.5 PER2::LUC$^{+/-}$ embryos and performed bioluminescence imaging for 5–7 days
(**Fig. 1Aa**). Spectral analysis revealed two consistent loci with ~24 h periods, circadian power exceeding
30% (see Materials and Methods), and high RMS amplitude (**Fig. S4**). The ventral locus (white arrow)
was LHX1-positive, confirming its identity as the SCN (VanDunk et al., 2011), whereas the posterior
structure (red arrow) was TTR-positive, a known marker for the CP (Stauder et al., 1986) (**Fig. 1Cc**).
When analysis was restricted to the SCN and 4VCP, a strong rhythm with a period between 20 and 28 h
was detected (**Fig. 1Cd**). Notably, the pituitary gland also showed rhythmicity, but with a markedly
lower RMS amplitude (**Fig. S4**).

To test whether the 4VCP begins to oscillate before the SCN, we repeated the experiment at E13.5 (**Fig.
1D**). We did not observe any circadian oscillation from ventral structures corresponding to SCN
progenitors, consistent with previous reports (Wreschnig et al., 2014; Carmona-Alcocer et al., 2018)
(**Fig. 1Db,Dd**). However, only the 4VCP (red arrow) exhibited a near-24 h rhythm with strong circadian
power and high RMS amplitude (**Fig. 1Db** and **S4**), which we confirmed after restricting the analysis to
the 4VCP (**Fig. 1Dd**). At E11.5, no significant circadian rhythm was detected in the 4VCP region in brain
slices (**Fig. 1E**). Because the 4VCP in a slice could still be influenced by neighboring tissues, we asked
whether the rhythm observed in E13.5 could originate from an endogenous 4VCP clock. Therefore, we
dissected E13.5 4VCPs and recorded their bioluminescence *ex vivo* for 7 days (**Fig. 1F**). We found that
the isolated 4VCP explant exhibited robust circadian rhythmicity at E13.5, with a mean circadian power
of ~40%, well above the detection floor determined from blank luciferase medium (Blk, < 20%) (**Fig.
1Fb**).

To pinpoint the time of the clock emergence, we dissected 4VCPs from PER2::LUC embryos at
embryonic stages beginning at E9.5, when the CSF compartment forms during neural tube closure
(Chau et al., 2015), to postnatal stages. We recorded the 4VCP PER2::LUC bioluminescence from 110
embryos, with 3–11 embryos from at least 2–3 litters per stage, for 7 days with PMTs. The period,
initially long and dispersed (37.08 ± 6.53 h at E10.5; *n*=17), converged toward 24 h by E11.5 (25.08 ± 0.88
h; *n*=15), when the first detectable oscillations appeared and circadian power exceeded 20% (31.21 ±
2.36% at E11.5; *n*=15; vs. 17.7 ± 1.69% at E10.5; *n*=17) (**Fig. 2A,C**). However, as seen in the head slice
(**Fig. 1Ec**), these oscillations were weakly sustained, with low RMS amplitude. At E12.5, the oscillations
persisted for the full 7-day recording, and their amplitude increased further at E13.5. The rhythm
became fully circadian at E15.5 (24.47 ± 0.60 h; *n*=5), with high circadian power (50.96 ± 3.42%; *n*=5)
relative to the 2-day window ceiling (~60%) (**Fig. 2A,C,D**).

We also used the *Bmal1-ELuc* reporter strain to characterize the molecular assembly of the embryonic
circadian clock. A 24 h period in *Bmal1-ELuc* oscillation was observed from E13.5 to E14.5,
approximately 2 days after the emergence of PER2 oscillations. From the sigmoidal fits to circadian
power trajectories (see Materials and Methods), we estimated the emergence time T_e and found that
PER2::LUC oscillations started to be circadian at E11.7 ± 0.14 *ex vivo*, whereas *Bmal1-ELuc* oscillations
emerged at E13.52 ± 0.70 *ex vivo* (**Fig. 2D**). This result aligns with the sudden shift of the circadian
power of PER2::LUC above 20% observed at E12.5 (**Fig. S5**). Because the circadian clock and
developmental programs are intertwined, we mapped the embryonic 4VCP's developmental trajectory

by quantifying mRNA levels for stage-associated genes. For progenitor or stem cell markers, we
measured *Nanog* (stem cells), *Gli3* (neuroepithelium), *Wnt1* (roof-plate and CP epithelium progenitors)
(Awatramani et al., 2003; Hunter and Dymecki, 2007), and *Ki67* (proliferation). We compared these with
differentiation markers *Aqp1*, *Wnt5a*, *Ttr*, and *Otx2*, which are upregulated as the 4VCP matures
(Johansson et al., 2013; Lun et al., 2015b; Langford et al., 2020) (**Fig. 2E**). After normalizing each
transcript to its peak abundance between E9.5 and E16.5, we observed that progenitor markers
predominate from E9.5 to E12.5, whereas differentiation genes dominate thereafter (**Fig. 2E**).
Sigmoidal fits estimated the progenitor half-minimum at E13.63 ± 0.79 and the differentiation half-
maximum at E13.52 ± 0.20, marking a sharp transition that coincided with the *Bmal1-ELuc* T_e in the
4VCP (E13.52 ± 0.70), as anticipated from developmental maturation (Yagita, *et al.*, 2024). By contrast,
PER2::LUC oscillations appear earlier and autonomously (E11.70 ± 0.14), and precede *Bmal1* and the
differentiation switch, an order that is atypical for a canonical TTFL (**Fig. 2G**).

Finally, we compared the emergence time of PER2::LUC rhythms across the brain and peripheral
organs in isolated explant culture. Among all tissues tested, the 4VCP was the first to become circadian,
preceding the LVCP (E12.37 ± 0.19), which develops later than the 4VCP (Lun et al., 2015b), and the
SCN (E14.21 ± 0.41), consistent with previous reports (Carmona-Alcocer et al., 2018). In the SCN, the
period approached 24 h at E14.5 (24.47 ± 2.35 h, *n*=7) and became fully circadian at E15.5 with high
circadian power (44.56 ± 2.48%, *n*=3). Peripheral organs emerged later in gestation and after the SCN
(liver: E14.53 ± 0.61; kidney: E15.08 ± 0.33) (**Fig. 2F,G**).

Early-to-late circadian gene expression in the embryonic 4VCP

Because mRNA and protein levels can be uncoupled during development (Umemura et al., 2017), we
assessed the emergence of circadian mRNA expression rhythms in the 4VCP. To capture circadian
variation, we sampled embryonic 4VCPs from 104 litters across E9–E16 at four circadian timepoints
(ZT2, ZT8, ZT14, ZT20) each embryonic day and performed RT-qPCR. We first compared the overall
mRNA levels of a specific gene set throughout development, and observed that at E9, the highest
transcript levels were for *Rora*, *Clock*, *Cry1*, and *Chrono*, rather than *Bmal1* and *Per1/2/3* (**Fig. 3A**).

We then clustered the transcripts according to their full temporal expression profiles across
development (**Fig. 3B**). Pairwise distances between expression profiles were computed using the
Pearson correlation, followed by hierarchical clustering. Four clusters emerged with distinct onset
patterns: early and sustained from E9.5 (Cluster 1: *Cry2*); inverted U-shaped, rising at ~E12.5 and
declining after ~E15.5 (Cluster 2: *Per2*, *Chrono*, *Rora*, *Rorc*); late-emerging at ~E13.5 (Cluster 3: *Per1*,
Per3, *Bmal1*, *Dbp*, *Clock*, *Rev-Erbβ*); and U-shaped, with an early onset at ~E9.5, a transient silence after
E11.5, and a rebound from ~E13.5 onward (Cluster 4: *Cry1*, *Rev-Erbα*). These clusters do not strictly align
with promoter motif architecture underlying the circadian transcriptional logic of the TTFL (Ukai-
Tadenuma et al., 2008; Takahashi, 2017). Notably, *Bmal1* and *Clock*, which act as drivers of the E-box
loop, belong to Cluster 3, and their baseline expression increases later than *Cry2* and *Per2* (**Fig. 3A**),
mirroring the onset-timing difference noted in *Bmal1-ELuc* and PER2::LUC reporter oscillations (**Fig.
2G**). In addition, *Per1* and *Per2*, both with glucocorticoid response elements (GREs) in their promoters

that can reset the clock in the CP (Liška et al., 2021), fall into two different clusters. These point to a
possibility that embryonic circadian gene expression in the 4VCP diverges from a simple TTFL-based
expectation.
448
We evaluated how circadian rhythms of each gene compared with the clusters, which are more
influenced by baseline changes across development, by quantifying circadian power using SST-CWT
analysis (Materials and Methods; **Fig. S7, S8A**). This analysis revealed that the strength of circadian
rhythmicity is unrelated to baseline expression patterns. Despite sustained expression, the Cluster 1
gene *Cry2* is circadian from E9 but rapidly loses rhythmicity; Cluster 2 shows heterogeneous behavior,
with *Per2* remaining below 20% circadian power; despite late increases in baseline expression, most
Cluster 3 genes become circadian around E11.5; and Cluster 4 genes remain circadian from E9 to E16
(**Fig. 3C**). To further validate our findings, we created phase-weighted power spectrograms of
representative transcripts (red), which visualize changes in dominant period and power together with
phase across cycles (**Fig. 3C**, lower panels). For several genes, including *Cry2, Chrono, Rora, Clock, Per1*,
and *Rev-erbβ*, we confirmed an initial circadian rhythm at E9–E10, which became indistinct around E13–
E14 and re-emerged at E15–E16 with an almost inverted phase (**Fig. S7, S8A**). Overall, most clock
genes in the embryonic 4VCP are transiently circadian before E16.5.
462
To better understand the establishment of the molecular clock and the dynamics between clusters, we
next compared the expression of the embryonic clock genes across development with the
representative gene of each cluster (**Fig. 3D**) and summarized the phase correlations in **Fig. 3E**. From
these correlative heat maps, we identified clusters whose correlations remained stable across time
(Cluster 2, 4 and claudin genes *Cldn2* and *Cldn11*) (**Fig. 3E**), with Cluster 2 and the claudins were
strongly positively correlated (**Fig. S8C**), and other clusters showing correlations that varied over
development. For the latter, three eras were apparent: E9–E11, E11–E13, and E13–E16. Although this
division suits most genes, it is especially clear for the dynamics between Cluster 3 (containing *Bmal1*)
and Cluster 4 (containing *Rev-Erbα*); while they are generally antiphasic (negatively correlated), they
become in-phase (positively correlated) between E11.5–E13.5 (**Fig. 3D, S8B**). The opposite trend can be
seen with Cluster 2 (*Rora*) and Cluster 3 (*Clock*); while generally positively correlated, they become
uncorrelated or negatively correlated around E11.5–E13.5 (**Fig. 3D, S8B, S8C**). Finally, we studied the
phase of expression to understand the origin of the change in correlation. Not only did the phase of
Cluster 3 and Cluster 4 change between E10.5 and E12.5, but the phases within each cluster also
became more homogeneous (**Fig. 3F**). By E16.5, we observed cluster polarization, with most positively
correlated and in-phase, whereas Cluster 4 was negatively correlated and antiphase to the others (**Fig.**
3E, 3F).
480
The absence of sustained PER2::LUC or *Bmal1-ELuc* rhythms in the embryonic 4VCP before E11.5–
E12.5 *ex vivo*, despite *Rev-Erbα* and *Cry2* being circadian *in vivo*, suggests that the early circadian gene
expression may depend on external inputs, such as maternal cues (Wong et al., 2022). We therefore
compared the expression profiles of selected clock genes in the 4VCP of 91 dams with the 4VCP of
PER2::LUC embryos. From hierarchical clustering, we identified 3 clusters of maternal clock genes,
which largely align with the gene classes, except for Cluster 1 which contains both E-box- and RRE-

driven genes: *Per2*, *Bmal1*, and *Cry1* (**Fig. 4A**). Surprisingly, in the maternal 4VCP, *Bmal1* and *Per2*
expression, initially antiphasic, became in-phase around E11.5–E13.5 (**Fig. 4A, 4C, S10A**). SST-CWT
analysis and circadian power estimation for each maternal gene (**Fig. S9**) showed all genes exhibited
strong circadian power (**Fig. 4B, S9**).

To investigate how maternal signals might influence embryonic clock development, we examined
temporal correlations between maternal and embryonic gene expression (**Fig. 4D, 4E**). We compared
maternal *Per2*, *Bmal1*, and *Dbp* with representative genes from each embryonic cluster (**Fig. S10C**). We
found that these correlations were not static but dynamically reorganized during development,
revealing three distinct epochs that mirror the embryonic expression timeline, although less sharply
defined. Before E11.5, when the embryonic clock is still immature and only Cluster 1 and 4 show
circadian rhythmicity, maternal and embryonic *Bmal1* were largely uncorrelated. Cluster-specific
patterns emerged nonetheless: Cluster 1 correlated positively with maternal *Bmal1*, whereas Cluster 4
correlated negatively with maternal *Dbp*, which are two maternal genes that are negatively correlated
with each other (**Fig. 4C, 4E, S10B**). Between E11.5 and E13.5, as Clusters 2 and 3 acquire circadian
rhythmicity, maternal-embryonic relations reorganized: correlation shifted away from maternal *Bmal1*
and *Dbp* toward maternal *Per2*, suggesting that maternal *Per2* or associated pathway may help
synchronize newly rhythmic components of the embryonic clock. This period might signify a major
developmental phase transition, where the embryo's internal clock becomes more autonomous but still
responsive to maternal signals. From E13.5 to E16.5, as most embryonic clusters acquired circadian
rhythmicity (except Cluster 4), embryonic genes became negatively correlated with maternal *Bmal1*.
Toward the end of this phase, correlation weakened substantially, consistent with increasing embryonic
autonomy. Dual-radius phase plots further revealed a shift in maternal *Per2* and *Rora* expression around
E12.5, while maternal *Bmal1* and *Dbp* remained relatively stable (**Fig. 4F**). This shift occurred precisely
when maternal-embryonic correlation patterns reorganized, raising the possibility that some of
maternal clock genes might undergo their own adjustments during this development window. Whether
this reflects intrinsic maternal dynamics or embryo-evoked changes remains to be determined.

Taken together, these results indicate that maternal and embryonic clocks are coordinated, especially
at E9.5–E11.5 when embryonic Clusters 1 and 4 are strongly correlated, and that the maternal clock
appears to undergo a transient reorganization during the E11.5–E13.5 epoch. Notably, this epoch
coincides with the functional maturation of the placenta, when maternal nutrient and oxygen supply to
the embryonic circulation begins (Elmore et al., 2022). To test whether placental maturation
contributes to this change in clock dynamics, we separated embryonic and maternal placentas and
recorded PER2::LUC and *Bmal1-ELuc* bioluminescence *ex vivo*. As reported previously (Čečmanová et
al., 2019), the embryonic placenta lacked autonomous circadian oscillations, whereas the maternal
placenta exhibited robust rhythms with a transient mid-gestational attenuation (**Fig. 4G**). In the
maternal placenta, circadian power of PER2::LUC and *Bmal1-ELuc* oscillations showed a mid-
gestational nadir of ~20–30% at E15.3 ± 0.4 and E15.6 ± 0.6, respectively, estimated from inverted
Gaussian fits. This dip occurs during E13–E15 and coincides with timing of increased activity of 11β-
hydroxysteroid dehydrogenase type 2 (11β-HSD2) (E12.5-E16.5) (Salvante et al., 2017; Brown RW et al.,
1996; Thompson *et al.*, 2002), which inactivates glucocorticoids and is mainly active in the maternal

placenta, suggesting a potential endocrine gating mechanism that shields the embryo from the
maternal influence (Astiz and Oster, 2021), although causality remains to be tested.
531
Temperature cycles can drive the embryonic 4VCP rhythmicity until E16.5

Because the circadian maternal core body temperature changes around E11.5 (Smarr et al., 2016) and
because nutrition supply shifts from histiotrophic (yolk sac-based) to hemotrophic (blood-based),
beginning at ~E10 and becoming fully functional by E12.5–E13.5 (Elmore et al., 2022), we hypothesized
that small intrauterine temperature cycles could serve as a non-bloodborne driver of the nascent
embryonic clock. During pregnancy, the maternal core-temperature rhythm amplitude dampens to ~1
°C, and to ~0.5 °C from around mid-gestation (Smarr et al., 2016). By contrast, many peripheral clocks
in vitro require larger temperature cycles for entrainment (~4 °C, Brown et al., 2002; ~2.5 °C, Buhr et al.,
2010; ~1.5 °C, Abraham et al., 2010), consistent with heat shock factor activation thresholds near ~2.5
°C (Brown and Rush, 1996; Buhr et al., 2010). If the embryonic 4VCP entrains to a 0.5 °C cycle, this
would imply heightened phase sensitivity near onset, as expected for a system close to a SNIC
threshold, rather than the higher-amplitude requirements typical of mature Hopf-type limit cycle
oscillators.
545
We therefore tested temperature entrainment directly by recording PER2::LUC activity in cultured
embryonic 4VCP explants. Since maternal body temperature at E6 varies between 37.5°C and 37°C
(Wharfe et al., 2016), we applied 2 days at 37°C followed by 12 days of a 37–37.5°C (± 0.1 °C) cycle with a
26-h period to E14, E16, and E18 4VCP and lung explants (**Fig. 5A**). At E14, 4VCP PER2 rhythms
entrained to the 26 h period temperature cycle, and at E16 they entrained after a brief transient,
whereas at E18 the 0.5 °C temperature gradient did not entrain the 4VCP (**Fig. 5B**). Across all
embryonic stages, the lung was entrained by the temperature cycle, possibly reflecting its prolonged
structural maturation that continues postnatally. We applied the same protocol to the LVCP and SCN
and found that the SCN remained entrainable at E18, which is plausible given that the SCN network is
still immature at this stage (Bedont and Blackshaw, 2015), but the LVCP, like the 4VCP, was not (**Fig.
S11A**). Together, these results indicate that small temperature cycles can influence the embryonic
4VCP through E16, whereas by E18, the oscillator exhibits reduced sensitivity to small temperature
gradients and behaves more like a robust, limit cycle oscillator that maintains a stable rhythm under
constant conditions.
560
Abrupt, SNIC-like onset and synchronization of clock emergence

A transition from a stable equilibrium to self-sustained oscillation is called a bifurcation. Single-cell
circadian clocks are well described by robust limit cycles often modeled near a supercritical Hopf
bifurcation, with amplitude that increases smoothly while the period remains near 24 h (Leloup and
Goldbeter, 1999). By contrast, systems near a SNIC threshold show abrupt appearance of finite-
amplitude oscillations, with very long initial periods that shorten toward ~24 h and can also display
pronounced phase sensitivity to weak perturbations (Ermentrout, 1996). The bifurcation type can be
probed by tracking how amplitude and period change over development. In our single-cell-resolution

PER2::LUC imaging data, we observe a sharply defined onset, with a long initial period and stepwise
increase in single-cell amplitude (**Fig. 5C**), consistent with a SNIC-like onset near the birth of the
circadian clock. Whole-tissue PER2::LUC luminometry likewise shows a long initial period that
converges to ~24 h through development (**Fig. 2B**). The apparent linear increase in whole-tissue
amplitude may partly reflect growth in 4VCP cell number across development (**Fig. S11B**).
574
At the population level, we observe an abrupt increase in synchronization at E14.5 (**Fig. 5D**), quantified
by the Kuramoto order parameter, where 0 indicates the absence of synchronization and 1 indicates
complete synchronization. The abrupt developmental shift is visible also in spatial phase distributions
at day 4 in culture (**Fig. 5E**), reminiscent of abrupt synchronization emergence in the SCN (Carmona-
Alcocer et al., 2018). These let us conclude that the clock emergence process at both single-cell and
population levels are not gradual but discretely staged in developmental epochs.
581

582 **Discussion**
583
Our study identifies the embryonic choroid plexus as an early brain circadian clock that emerges before
the SCN. Through PER2::LUC time lapse recording in 4VCP explant cultures, we found that circadian
PER2 rhythms appear from E12.5, 2–3 days earlier than the central clock. We also observed staged
assembly of the transcriptional clock circuit, with a delayed rise of *Bmal1* transcripts, coincident with
the expression of the differentiation markers. Accordingly, we detected the postponed onset of *Bmal1-*
Eluc oscillations *ex vivo* around E13.5.
590
When we performed circadian profiling of select transcripts across embryonic stages, we observed that
12 out of the 13 clock genes showed at least transient circadian expression patterns before E16.5 and
displayed strong, developmentally shifting correlations with the maternal clock gene expression.
Within a specific developmental window between E11.5 and E13.5, correlations among embryonic clock
gene clusters were largely disrupted. This window coincides with placental maturation and the onset of
maternal-fetal circulation. Before hemotrophic nutrition is established, the embryonic clock may be
susceptible to blood-independent cues such as temperature, consistent with SNIC-like sensitivity near
onset. After placental maturation, endocrine gating at the placenta may reduce maternal influence,
enabling a developmental window for autonomous emergence and organization of circadian clocks in
the embryo.
601

The CP, the first clock in the brain?
Contrary to the prevailing assumption that the SCN is the primary structure to be circadian in the
mouse embryo because of its crucial role in adulthood, we found that the 4VCP clock emerges earlier
than the SCN. In our study, the 4VCP is the earliest in the brain structure to display clear circadian
rhythmicity, though our search is limited to select sagittal sections rather than across the entire 3D
aspect of the embryonic brain. The 4VCP nevertheless has strong functional implications as the first
clock, given our earlier demonstration that the adult 4VCP acts as a robust and influential oscillator

capable of entraining the SCN both *ex vivo* and *in vivo* (Myung et al., 2018). It is also notable that the
first three structures we identified as circadian before or around E15.5 are the CP, SCN, and pituitary
gland, which are all strategically positioned to regulate brain homeostasis, process external light cues,
and propagate outputs through endocrine pathways. Strikingly, these three structures also show the
strongest oscillations within the postnatal mouse brain (Myung et al., 2018).

The CSF-containing ventricles play a substantial role in early development; by E10.5–E11.5 the brain
ventricle volume is ≥10% of the head volume, substantially bathing the developing brain (Aristizábal et
al., 2013). This embryonic CSF is filled with morphogens and growth factors (Kaiser and Bryja, 2020;
Saunders et al., 2023; Lun et al., 2015a,b), mostly secreted from the CP. If the embryonic CP secretome
is released in a circadian manner, the neuroepithelium lining the large ventricles would receive
temporally patterned cues. Some of these inputs modulate transcription factors in neural progenitors
and influence fate decisions (Lamus et al., 2020; Lehtinen et al., 2011; Subashini et al., 2017). Although
not circadian, oscillations of basic-helix-loop-helix (bHLH) factors such as *Hes1*, *Ascl1*, and *Olig2*
promote cell proliferation, whereas sustained expression promotes cell differentiation (Imayoshi et al.,
2013). By inhibiting differentiation, the progenitor pool can expand before neurogenesis begins, a
prerequisite for reaching an appropriate number of both neurons and neural progenitors in the adult
stage (Caviness et al., 1995; Kriegstein and Alvarez-Buylla, 2009; Imayoshi and Kageyama, 2014). Other
factor-inducing pathways such as *Wnt*, *Hedgehog*, and *Notch* induce oscillatory patterns of target-gene
expression to regulate neural progenitor development (Imayoshi et al., 2014). This volume-transmission
from the CP is particularly important early in development, when synaptic circuits have yet to mature.
Synaptogenesis in the visual cortex, for instance, begins late in gestation and matures postnatally (Li et
al., 2010).

Clock and CP differentiation

The temporal convergence between 4VCP clock onset in culture and the start of the developmental
program, as well as the delay in clock onset between the 4VCP and the LVCP, illustrate the requirement
of differentiation for the clock to begin (Dan et al., 2020). These results agree with previous studies
showing that induced pluripotent stem cells (iPCSs), embryonic stem cells (ESCs), and pluripotent stem
cells (PSCs) lack a circadian clock, which emerges during differentiation (Yagita et al., 2010; Kowalska
et al., 2010; Kaneko et al., 2023; Umemura et al., 2017).

However, despite a progenitor profile marked by predominant markers such as *Gli3* and *Ki67* (**Fig. 2**),
we observed early circadian expression *in vivo* as early as E9.5 for *Cry2* when tissue is processed
immediately after sampling (**Fig. S7, S8**). This differs from whole-embryo analyses (Cao et al., 2022;
Dolatshad et al., 2010) and underscores the need to evaluate clock gene rhythmicity in a tissue-specific
manner. It is possible that some early rhythmic transcripts in the 4VCPs reflect maternally driven
activity. Challenges remain in determining the autonomous nature of such early gene oscillations as the
developing clock system may not conform to the adult model. For example, neonatal circadian clocks,
including the $Cry1^{-/-};Cry2^{-/-}$ SCN, deviate from the behavior expected under the canonical TTFL model
(Ono et al., 2013).

Expression of *Per2* and *Bmal1* is typically antiphasic in the mature clock, since BMAL1-complexes drive
Per2 transcription, and after a delay due to translational and post-translational modifications, PER2-
complexes repress transcription driven by BMAL1-complexes. In contrast, in our mRNA dataset, the
two components do not co-emerge in the 4VCP, and in cultured explant recordings, *Bmal1-ELuc* starts
circadian expression about ~2 days after PER2::LUC. This highlights the possibility that the very first
oscillation of the embryonic clock may be independent of circadian *Bmal1* expression. It is plausible that
other E-box activators operate during the early developmental stage, which do not need to be circadian
themselves, as non-rhythmic *Bmal1* overexpression can lead to circadian expression of *Per2* (Liu et al.,
2008) and *Bmal1* itself (Padlom et al., 2022). These features may relate to SNIC-like behavior, which
most TTFL models do not predict. Regardless of the exact mechanism, this early clock could be
physiologically important; for example, metabolic rhythms in glucose uptake have been observed in
undifferentiated stem cells even when core clock genes appear arrhythmic (Paulose et al., 2012).

Entrainment of the late embryonic clock

Multiple lines of evidence indicate that a relatively autonomous clock operates by late gestation. In
preterm infants, circadian-like rhythms in body temperature, heart rate, and rest-activity can be
detectable in the NICU (Bueno et al., 2016; Mirmiran et al., 1990; Mirmiran and Kok, 1991; Govindan et
al., 2024), consistent with a late-developing endogenous embryonic clock. Pups born from dams with
late SCN lesions do not show inter-pup desynchrony, unlike litters from early SCN-lesioned dams (Davis
and Gorski, 1988; Jud and Albrecht, 2006). In preterm infants, one study even demonstrated that the 24
h sleep-wake rhythm appears earlier in preterm than in full term babies (Guyer et al., 2015), although
amplitude may be lower and phase may differ (e.g., inverted heart-rate rhythm; Govindan et al., 2024).
Such phase differences could arise from changes in external zeitgebers such as feeding schedules. In
mice, food restriction during and after pregnancy can shift offspring rhythms (Canaple et al., 2018).
While ultradian temperature variation in the NICU generally does not seem to disrupt circadian
development in infants born after mid-gestation (Thomas, 2001; Bueno et al., 2016; Mirmiran et al.,
1990; Mirmiran and Kok, 1991), our data show that even when the 4VCP exhibits a strong rhythm at
E16, it remains sensitive to weak temperature changes (0.5 °C), with reduced sensitivity by E18. These
observations may help frame hypotheses about temperature-based care protocols customized to
different stages of preterms.

Conclusion

The early emergence of the circadian clock in the 4VCP from E11.5 to E12.5 challenges the long-held
view of the SCN as the central player of the circadian pacemaking in the mammalian brain and, by
extension, as the locus of developmental onset. Our findings also reveal that the onset is a discrete,
multi-stage process. We show that the *ex vivo* rhythm of the canonical transcriptional activator *Bmal1*
appears later than that of *Per2*, suggesting that the embryonic clock is not established by the classic
BMAL1:CLOCK feedback. Instead, qPCR analysis of the embryonic 4VCP uncovered non-canonical
rhythmic expressions of *Chrono* and *Rev-Erbα* as early as E9.5. At the point of emergence, while mature

circadian oscillators are often described by a Hopf bifurcation, we find that this transition shows hallmarks of a SNIC bifurcation. Implications of these findings are twofold. First, the early clock in the 4VCP is best poised to guide the rhythms in other loci of the brain through the rhythmic secretion of factors into the CSF. Second, the internally multi-stage process and externally sudden emergence align with the idea that the circadian clock onset is a programmed developmental event, which provides a new framework for understanding the maternal-fetal clock interactions and the potential impact of their disruption. By characterizing the precise timing, locus, and mechanism of the brain's first clock, our work provides insights into early circadian homeostasis that are critically relevant to preterm and neonatal settings.

Acknowledgements

We thank Lilian Byer for critical reading of the manuscript. This work was supported by the National Science and Technology Council (NSTC) (109-2314-B-038-071, 109-2320-B-038-020, 110-2314-B-038-162, 110-2311-B-038-003, 111-2314-B-038-008, 112-2314-B-038-063, 113-2314-B-038-121, 114-2320-B-038-052-MY3, 114-2811-B-038-015) and the Higher Education Sprout Project of the Ministry of Education (MOE) in Taiwan.

References

Abraham U, Granada AE, Westermark PO, Heine M, Kramer A, Herzel H (2010). Coupling governs entrainment range of circadian clocks. *Mol Syst Biol*, 6:438.

Alonso MI, Martín C, Carnicero E, Bueno D, Gato A (2011). Cerebrospinal fluid control of neurogenesis induced by retinoic acid during early brain development. *Dev Dyn*, 240:1650–1659.

Alonso MI, Lamus F, Carnicero E, Moro JA, de la Mano A, Fernández JMF, Desmond ME, Gato A (2017). Embryonic cerebrospinal fluid increases neurogenic activity in the brain ventricular-subventricular zone of adult mice. *Front Neuroanat*, 11:124.

Aristizábal O, Mamou J, Ketterling JA, Turnbull DH (2013). High-Throughput, High-Frequency 3-D Ultrasound for in Utero Analysis of Embryonic Mouse Brain Development. *Ultrasound Med Biol*, 39:2321–2332.

Astiz M, Oster H (2021). Feto-maternal crosstalk in the development of the circadian clock system. *Front Neurosci*, 14:631687.

Awatramani R, Soriano P, Rodriguez C, Mai JJ, Dymecki SM (2003). Cryptic boundaries in roof plate and choroid plexus identified by intersectional gene activation. *Nat Genet*, 35:70–75.

Balsalobre A, Damiola F, Schibler U (1998). A serum shock induces circadian gene expression in mammalian tissue culture cells. *Cell*, 93:929–937.

Battogtokh D, Tyson JJ (2018). Deciphering the dynamics of interlocked feedback loops in a model of
the mammalian circadian clock. *Biophys J*, 115:2055–2066.

Bedont JL, Blackshaw S (2015). Constructing the suprachiasmatic nucleus: a watchmaker's perspective
on the central clockworks. *Front Syst Neurosci*, 9:74.

Birecree E, King LE, Nanney LB (1991). Epidermal growth factor and its receptor in the developing
human nervous system. *Dev Brain Res*, 60:145–154.

Brown IR, Rush SJ (1996). In vivo activation of neural heat shock transcription factor HSF1 by a
physiologically relevant increase in body temperature. *J Neurosci Res*, 44:52-57.

Brown RW, Diaz R, Robson AC, Kotelevtsev YV, Mullins JJ, Kaufman MH, Seckl JR (1996). The ontogeny
of 11 beta-hydroxysteroid dehydrogenase type 2 and mineralocorticoid receptor gene expression reveal
intricate control of glucocorticoid action in development. *Endocrinology*, 137:794–797.

Brown SA, Zumbrunn G, Fleury-Olela F, Preitner N, Schibler U (2002). Rhythms of mammalian body
temperature can sustain peripheral circadian clocks. *Curr Biol*, 12:1574–1583.

Bueno C, Menna-Barreto L (2016). Development of sleep/wake, activity and temperature rhythms in
newborns maintained in a neonatal intensive care unit and the impact of feeding schedules. *Infant
Behav Dev*, 44:21–28.

Bueno D, Parvas M, Nabiuni M, Miyan J (2020). Embryonic cerebrospinal fluid formation and regulation.
Semin Cell Dev Biol, 102:3–12.

Buhr ED, Yoo SH, Takahashi JS (2010). Temperature as a universal resetting cue for mammalian
circadian oscillators. *Science*, 330:379–385.

Cai C, Vandermeer B, Khurana R, Nerenberg K, Featherstone R, Sebastianski M, Davenport MH (2019).
The impact of occupational shift work and working hours during pregnancy on health outcomes: a
systematic review and meta-analysis. Am J Obstet Gynecol, 221:563–576.

Canaple L, Gréchez-Cassiau A, Delaunay F, Dkhissi-Benyahya O, Samarut J (2018). Maternal eating
behavior is a major synchronizer of fetal and postnatal peripheral clocks in mice. *Cell Mol Life Sci*,
75:3991–4005.

Cao X, Yan Y, Luo X, Yang X, Cui H, Yang Y, Li H (2022). Analyses of the circadian clock genes
expression in whole embryos and maternal major tissues of mice. *J Mol Histol*, 53:473–482.

Carmona-Alcocer V, Abel JH, Sun TC, Petzold LR, Doyle FJ, Simms CL, Herzog ED (2018). Ontogeny of
circadian rhythms and synchrony in the suprachiasmatic nucleus. *J Neurosci*, 38:1326–1334.

Caviness VS, Takahashi T, Nowakowski RS (1995). Numbers, time and neocortical neuronogenesis: a
general developmental and evolutionary model. *Trends Neurosci*, 18:379–383.

Čečmanová V, Houdek P, Šuchmanová K, Sládek M, Sumová A (2019). Development and entrainment
of the fetal clock in the suprachiasmatic nuclei: the role of glucocorticoids. *J Biol Rhythms*, 34:307–322.

Chau KF, Springel MW, Broadbelt KG, Park HY, Topal S, Lun MP, Mullan H, Maynard T, Steen H,
LaMantia AS, Lehtinen MK (2015). Progressive differentiation and instructive capacities of amniotic
fluid and cerebrospinal fluid proteomes following neural tube closure. *Dev Cell*, 35:789–802.

Courtney Y, Head JP, Dani N, Chechneva OV, Shipley FB, Zhang Y, Holtzman MJ, Sadegh C, Libermann
TA, Lehtinen MK (2025). Choroid plexus apocrine secretion shapes CSF proteome during mouse brain
development. *Nat Neurosci*, 28:1446–1459.

Dan H, Ruan T, Sampogna RV (2020). Circadian clock regulation of developmental time in the kidney.
Cell Rep, 31:107661.

Daubechies I, Lu J, Wu HT (2011). Synchrosqueezed wavelet transforms: An empirical mode
decomposition-like tool. *Appl Comput Harmon Anal*, 30:243–261.

Davari MH, Naghshineh E, Mostaghaci M, Mirmohammadi SJ, Bahaloo M, Jafari A, Mehrparvar AH
(2018). Shift work effects and pregnancy outcome: A historical cohort study. *J Family Reprod Health*,
12:84–88.

Davis FC, Gorski RA (1988). Development of hamster circadian rhythms: role of the maternal
suprachiasmatic nucleus. *J Comp Physiol A*, 162:601–610.

Dolatshad H, Cary AJ, Davis FC (2010). Differential expression of the circadian clock in maternal and
embryonic tissues of mice. *PLoS ONE*, 5:e9855.

Dziegielewska KM, Ek J, Habgood MD, Saunders NR (2001). Development of the choroid plexus.
Microsc Res Tech, 52:5–20.

Edelbo BL, Andreassen SN, Steffensen AB, MacAulay N (2023). Day-night fluctuations in choroid plexus
transcriptomics and cerebrospinal fluid metabolomics. *PNAS Nexus*, 2:pgad262.

Elmore SA, Cochran RZ, Bolon B, Lubeck B, Mahler B, Sabio D, Ward JM (2022). Histology atlas of the
developing mouse placenta. *Toxicol Pathol*, 50:60–117.

Ermentrout B (1996). Type I membranes, phase resetting curves, and synchrony. *Neural Comput*, 8:979–
1001.

Gato A, Moro JA, Alonso MI, Bueno D, de La Mano A, Martín C (2005). Embryonic cerebrospinal fluid
regulates neuroepithelial survival, proliferation, and neurogenesis in chick embryos. *Anat Rec A Discov
Mol Cell Evol Biol*, 284:475–484.

Gato A, Desmond ME (2009). Why the embryo still matters: CSF and the neuroepithelium as
interdependent regulators of embryonic brain growth, morphogenesis and histiogenesis. *Dev Biol*,
327:263–272.

Govindan RB, Andescavage NN, Basu S, Murnick J, Ngwa J, Galla JT, Kapse K, Limperopoulos C, du
Plessis A (2024). Circadian rhythm development in preterm infants: The role of postnatal versus
postmenstrual age. *Early Hum Dev*, 196:106084.

Guyer C, Huber R, Fontijn J, Bucher HU, Nicolai H, Werner H, Molinari L, Latal B, Jenni OG (2015). Very
preterm infants show earlier emergence of 24-hour sleep-wake rhythms compared to term infants.
Early Hum Dev, 91:37–42.

Hildyard JCW, Wells DJ, Piercy RJ (2022). Identification of qPCR reference genes suitable for
normalising gene expression in the developing mouse embryo. *Wellcome Open Res*, 6:197.

Huang X, Ketova T, Fleming JT, Wang H, Dey SK, Litingtung Y, Chiang C (2009). Sonic hedgehog
signaling regulates a novel epithelial progenitor domain of the hindbrain choroid plexus. *Development*,
136:2535–2543.

Hunter NL, Dymecki SM (2007). Molecularly and temporally separable lineages form the hindbrain roof
plate and contribute differentially to the choroid plexus. *Development*, 134:3449–3460.

Imayoshi I, Isomura A, Harima Y, Kawaguchi K, Kori H, Miyachi H, Fujiwara T, Ishidate F, Kageyama R
(2013). Oscillatory control of factors determining multipotency and fate in mouse neural progenitors.
Science, 342:1203–1208.

Imayoshi I, Kageyama R (2014). bHLH factors in self-renewal, multipotency, and fate choice of neural
progenitor cells. *Neuron*, 82:9–23.

Jia BZ, Qi Y, Wong-Campos JD, Megason SG, Cohen AE (2023). A bioelectrical phase transition patterns
the first vertebrate heartbeats. *Nature*, 622:149–155.

Johansson PA, Irmler M, Acampora D, Beckers J, Simeone A, Götz M (2013). The transcription factor
Otx2 regulates choroid plexus development and function. *Development*, 140:1055–1066.

Jud C, Albrecht U (2006). Circadian rhythms in murine pups develop in absence of a functional maternal
circadian clock. *J Biol Rhythms*, 21:149–154.

Kaiser K, Bryja V (2020). Choroid plexus: the orchestrator of long-range signalling within the CNS. *Int J
Mol Sci*, 21:4760.

Kaneko H, Kaitsuka T, Tomizawa K (2023). Artificial induction of circadian rhythm by combining
exogenous BMAL1 expression and polycomb repressive complex 2 inhibition in human induced
pluripotent stem cells. *Cell Mol Life Sci*, 80:200.

Kowalska E, Moriggi E, Bauer C, Dibner C, Brown SA (2010). The circadian clock starts ticking at a
developmentally early stage. *J Biol Rhythms*, 25:442–449.

Kriegstein A, Alvarez-Buylla A (2009). The glial nature of embryonic and adult neural stem cells. *Annu
Rev Neurosci*, 32:149–184.

Lamus F, Martín C, Carnicero E, Moro JA, Fernández JMF, Mano A, Gato Á, Alonso MI (2020). FGF2/EGF
contributes to brain neuroepithelial precursor proliferation and neurogenesis in rat embryos: the
involvement of embryonic cerebrospinal fluid. *Dev Dyn*, 249:141–153.

Landgraf D, Koch CE, Oster H, Alvarez-Bolado G, Pombal MA (2014). Embryonic development of
circadian clocks in the mammalian suprachiasmatic nuclei. *Front Neuroanat*, 8:143.

Langford MB, O'Leary CJ, Veeraval L, White A, Lanoue V, Cooper HM (2020). WNT5a regulates
epithelial morphogenesis in the developing choroid plexus. *Cereb Cortex*, 30:3617–3631.

Lehtinen MK, Zappaterra MW, Chen X, Yang YJ, Hill AD, Lun M, Maynard T, Gonzalez D, Kim S, Ye P,
D'Ercole AJ, Wong ET, LaMantia AS, Walsh CA (2011). The cerebrospinal fluid provides a proliferative
niche for neural progenitor cells. *Neuron*, 69:893–905.

Li M, Cui Z, Niu Y, Liu B, Fan W, Yu D, Deng J (2010). Synaptogenesis in the developing mouse visual
cortex. *Brain Res Bull*, 81:107–113.

Liu AC, Tran HG, Zhang EE, Priest AA, Welsh DK, Kay SA (2008). Redundant function of REV-ERBα and
β and non-essential role for Bmal1 cycling in transcriptional regulation of intracellular circadian
rhythms. *PLoS Genet*, 4:e1000023.

Liška K, Sládek M, Čečmanová V, Sumová A (2021). Glucocorticoids reset circadian clock in choroid
plexus via period genes. *J Endocrinol*, 248:155–166.

Lun MP, Johnson MB, Broadbelt KG, Watanabe M, Kang YJ, Chau KF, Springel MW, Malesz A, Sousa
AMM, Pletikos M, Adelita T, Calicchio ML, Zhang Y, Holtzman MJ, Lidov HGW, Sestan N, Steen H,
Monuki ES, Lehtinen MK (2015a). Spatially heterogeneous choroid plexus transcriptomes encode
positional identity and contribute to regional CSF production. *J Neurosci*, 35:4903–4916.

Lun MP, Monuki ES, Lehtinen MK (2015b). Development and functions of the choroid plexus-
cerebrospinal fluid system. *Nat Rev Neurosci*, 16:445–457.

Maeda Y, Muro M, Shono M, Shono H, Iwasaka T (2006). Diurnal rhythms in fetal heart rate baseline
and sustained fetal tachycardia in twin pregnancy. *Early Hum Dev*, 82:637–644.

Martín C, Bueno D, Alonso MI, Moro JA, Callejo S, Parada C, Martín P, Carnicero E, Gato A (2006). FGF2
plays a key role in embryonic cerebrospinal fluid trophic properties over chick embryo neuroepithelial
stem cells. *Dev Biol*, 297:402–416.

Martin C, Alonso MI, Santiago C, Moro JA, de la Mano A, Carretero R, Gato A (2009). Early embryonic
brain development in rats requires the trophic influence of cerebrospinal fluid. *Int J Dev Neurosci*,
27:733–740.

Mirmiran M, Kok JH, de Kleine MJK, Koppe JG, Overdijk J, Witting W (1990). Circadian rhythms in
preterm infants: a preliminary study. *Early Hum Dev*, 23:139–146.

Mirmiran M, Kok JH (1991). Circadian rhythms in early human development. *Early Hum Dev*, 26:121–
128.

Miyan JA, Nabiyouni M, Zendah M (2003). Development of the brain: a vital role for cerebrospinal fluid.
Can J Physiol Pharmacol, 81:317–328.

Murayama Y, Kori H, Oshima C, Kondo T, Iwasaki H, Ito H (2017). Low temperature nullifies the
circadian clock in cyanobacteria through Hopf bifurcation. *Proc Natl Acad Sci USA*, 114:5641–5646.

Myung J, Hong S, Hatanaka F, Nakajima Y, De Schutter E, Takumi T (2012). Period coding of Bmal1
oscillators in the suprachiasmatic nucleus. *J Neurosci*, 32:8900–8918.

Myung J, Hong S, DeWoskin D, De Schutter E, Forger DB, Takumi T (2015). GABA-mediated repulsive
coupling between circadian clock neurons in the SCN encodes seasonal time. *Proc Natl Acad Sci USA*,
112:E3920–E3929.

Myung J, Schmal C, Hong S, Tsukizawa Y, Rose P, Zhang Y, Holtzman MJ, De Schutter E, Herzel H,
Takumi T (2018). The choroid plexus is an important circadian clock component. *Nat Commun*, 9:1–13.

Myung J (2024). Enhancing bioluminescence imaging of cultured tissue explants using optical
telecompression. *Sensors*, 24:6041.

Myung J, Vitet H, Truong VH, Ananthasubramaniam B (2025). The role of the multiplicity of circadian
clocks in mammalian systems. *Sleep Med*, 131:106518.

Nakajima Y, Yamazaki T, Nishii S, Noguchi T, Hoshino H, Niwa K, Viviani VR, Ohmiya Y (2010).
Enhanced beetle luciferase for high-resolution bioluminescence imaging. *PLoS ONE*, 5:e10011.

Oates AC, Morelli LG, Ares S (2012). Patterning embryos with oscillations: structure, function and
dynamics of the vertebrate segmentation clock. *Development*, 139:625–639.

Ono D, Honma S, Honma KI (2013). Cryptochromes are critical for the development of coherent
circadian rhythms in the mouse suprachiasmatic nucleus. *Nat Commun*, 4:1666.

Padlom A, Ono D, Hamashima R, Furukawa Y, Yoshimura T, Nishiwaki-Ohkawa T (2022). Level of
constitutively expressed BMAL1 affects the robustness of circadian oscillations. *Sci Rep*, 12:19519.

Parada C, Escolà-Gil JC, Bueno D (2008). Low-density lipoproteins from embryonic cerebrospinal fluid
are required for neural differentiation. *J Neurosci Res*, 86:2674–2684.

Parvas M, Parada C, Bueno D (2008). A blood-CSF barrier function controls embryonic CSF protein
composition and homeostasis during early CNS development. *Dev Biol*, 321:51–63.

Paulose JK, Rucker EB, Cassone VM (2012). Toward the beginning of time: Circadian rhythms in
metabolism precede rhythms in clock gene expression in mouse embryonic stem cells. *PLoS ONE*,
7:e49555.

Salvante KG, Milano K, Kliman HJ, Nepomnaschy PA (2017). Placental 11 β-hydroxysteroid
dehydrogenase type 2 (11β-HSD2) expression very early during human pregnancy. *J Dev Orig Health*
Dis, 8:149–154

Saunders NR, Dziegielewska KM, Fame RM, Lehtinen MK, Liddelow SA (2023). The choroid plexus: A
missing link in our understanding of brain development and function. *Physiol Rev*, 103:919–956.

Serón-Ferré M, Mendez N, Abarzua-Catalan L, Vilches N, Valenzuela FJ, Reynolds HE, Llanos AJ, Rojas
A, Valenzuela GJ, Torres-Farfan C (2012). Circadian rhythms in the fetus. *Mol Cell Endocrinol*, 349:68–75.

Smarr BL, Zucker I, Kriegsfeld LJ (2016). Detection of successful and unsuccessful pregnancies in mice
within hours of pairing through frequency analysis of high temporal resolution core body temperature
data. *PLoS ONE*, 11:e0160127.

Stauder AJ, Dickson PW, Aldred AR, Schreiber G, Mendelsohn FA, Hudson P (1986). Synthesis of
transthyretin (pre-albumin) mRNA in choroid plexus epithelial cells, localized by in situ hybridization in
rat brain. *J Histochem Cytochem*, 34:949–952.

Strogatz SH (2024). *Nonlinear Dynamics and Chaos: With Applications to Physics, Biology, Chemistry,*
and Engineering. 3rd ed. New York: Chapman & Hall/CRC.

Subashini C, Dhanesh SB, Chen CM, Riya PA, Meera V, Divya TS, Kuruvilla R, Buttler K, James J (2017).
Wnt5a is a crucial regulator of neurogenesis during cerebellum development. *Sci Rep*, 7:42523.

Takahashi JS (2017). Transcriptional architecture of the mammalian circadian clock. *Nat Rev Genet*,
18:164–179.

Thomas KA (2001). Biological rhythm development in preterm infants: Does health status influence
body temperature circadian rhythm? *Res Nurs Health*, 24:170–180.

Thompson, A., Han, V. K. M., & Yang, K. (2002). Spatial and temporal patterns of expression of 11β-
hydroxysteroid dehydrogenase types 1 and 2 messenger RNA and glucocorticoid receptor protein in the
murine placenta and uterus during late pregnancy. *Biol Repro*, 67:1708–1718.

Truong VH, Myung J (2023). LocoBox: Modular Hardware and Open-Source Software for Circadian
Entrainment and Behavioral Monitoring in Home Cages. *Sensors*, 23:9469.

Ukai-Tadenuma M, Kasukawa T, Ueda HR (2008). Proof-by-synthesis of the transcriptional logic of
mammalian circadian clocks. *Nat Cell Biol*, 10:1154–1163.

Umemura Y, Koike N, Ohashi M, Tsuchiya Y, Meng QJ, Minami Y, Hara M, Hisatomi M, Yagita K (2017).
Involvement of posttranscriptional regulation of Clock in the emergence of circadian clock oscillation
during mouse development. *Proc Natl Acad Sci USA*, 114:E7479–E7488.

VanDunk C, Hunter LA, Gray PA (2011). Development, maturation, and necessity of transcription
factors in the mouse suprachiasmatic nucleus. *J Neurosci*, 31:6457–6467.

Wharfe MD, Mark PJ, Wyrwoll CS, Smith JT, Yap C, Clarke MW, Waddell BJ (2016). Pregnancy-induced
adaptations of the central circadian clock and maternal glucocorticoids. *J Endocrinol*, 228:135–147.

Wong SD, Wright KP, Spencer RL, Vetter C, Hicks LM, Jenni OG, LeBourgeois MK (2022). Development
of the circadian system in early life: maternal and environmental factors. *J Physiol Anthropol*, 41:22.

Wreschnig D, Dolatshad H, Davis FC (2014). Embryonic development of circadian oscillations in the
mouse hypothalamus. *J Biol Rhythms*, 29:299–310.

Yagita K (2024). Emergence of the circadian clock oscillation during the developmental process in
mammals. *Curr Opin Genet Dev*, 84:102152.

Yagita K, Horie K, Koinuma S, Nakamura W, Yamanaka I, Urasaki A, Shigeyoshi Y, Kawakami K,
Shimada S, Takeda J, Uchiyama Y (2010). Development of the circadian oscillator during differentiation
of mouse embryonic stem cells in vitro. *Proc Natl Acad Sci USA*, 107:3846–3851.

Yamazaki S, Numano R, Abe M, Hida A, Takahashi RI, Ueda M, Block GD, Sakaki Y, Menaker M, Tei H
(2000). Resetting central and peripheral circadian oscillators in transgenic rats. *Science*, 288:682–685.

Yoo SH, Yamazaki S, Lowrey PL, Shimomura K, Ko CH, Buhr ED, Siepka SM, Hong HK, Oh WJ, Yoo OJ,
Menaker M, Takahashi JS (2004). PERIOD2::LUCIFERASE real-time reporting of circadian dynamics
reveals persistent circadian oscillations in mouse peripheral tissues. *Proc Natl Acad Sci USA*, 101:5339–
5346.

Zappaterra MW, Lehtinen MK (2012). The cerebrospinal fluid: Regulator of neurogenesis, behavior, and
beyond. *Cell Mol Life Sci*, 69:2863–2878.

A brain slice *C, D, E* | 4VCP isolation *F*

B SCN — Circadian PER2::LUC
E11.5   E13.5   E15.5   E18.5
4VCP — Circadian PER2::LUC

C a Bright field — E15.5 | PER2::LUC
c LHX1 | TTR
b Period | Circadian power
d PER2::LUC | PER2::LUC

D a Bright field — E13.5 | PER2::LUC
c PER2::LUC Hypothalamus | PER2::LUC 4VCP
b Period | Circadian power
d PER2::LUC | PER2::LUC

E a Bright field — E11.5 | PER2::LUC
c PER2::LUC Hypothalamus | PER2::LUC 4VCP
b Period | Circadian power
d PER2::LUC | PER2::LUC

F a PER2::LUC — E13.5 | b Period | Circadian power | c PER2::LUC

Figure 1. The 4VCP is circadian before the SCN in the mouse embryonic brain. (A) Schematic of the protocol used for recording PER2::LUC bioluminescence from embryonic brain or head slices (left) or from the isolated 4VCP (right). (**B**) Illustration summarizing the finding that the 4VCP is circadian earlier than the SCN in the mouse embryo. (**C**) **a**: Bright field and PER2::LUC images of the bioluminescence obtained from a PER2::LUC E15.5 brain slice. **b**: Analysis of the recording showing the period (left) and the circadian power (right). **c**: Images of immunofluorescence of LHX1 (left) and TTR (right) after brain clearing **d**: Detrended time series of PER2::LUC bioluminescence from the areas indicated by the white boxes in **c.** (**D**) **a**: Bright field and PER2::LUC images of the bioluminescence obtained from a PER2::LUC E13.5 head slice. **b**: Analysis of the recording showing the period (left) and the circadian power (right). **c**: PER2::LUC images with white boxes indicating the putative hypothalamus (left) and 4VCP (right). **d**: Detrended time series of PER2::LUC bioluminescence from the areas indicated by the white boxes in **c.** (**E**) **a**: Bright field and PER2::LUC images of the bioluminescence obtained from a PER2::LUC E11.5 head slice. **b**: Analysis of the recording showing the period (left) and the circadian power (right). **c**: PER2::LUC images with white boxes indicating the putative hypothalamus (left) and 4VCP (right). **d**: Detrended time series of PER2::LUC bioluminescence from the areas indicated by the white boxes in **c.** (**F**) **a**: PER2::LUC image of the bioluminescence obtained from a PER2::LUC E13.5 isolated 4VCP. **b**: Analysis of the recording showing the period (left) and the circadian power (right). **c**: Detrended time series of the recording. Scale bar: 500 μm. Red arrows indicate 4VCP and white arrows indicate SCN.

Figure 2. The 4VCP clock emerges at E11.5–E12.5, preceding the LVCP and the SCN and coinciding with the onset of 4VCP differentiation. (**A**) Individual (thin colored) and mean (black) detrended traces (top) and spectrograms (bottom) of PER2::LUC bioluminescence recorded from embryonic 4VCP from E9.5 to E18.5. Zoom-in insets highlight low amplitude rhythms. (**B**) Individual (thin colored) and mean (black) detrended traces (top) and spectrograms (bottom) of *Bmal1-ELuc* bioluminescence recorded from embryonic 4VCP from E10.5 to E18.5. Zoom-in insets highlight low amplitude rhythms. (**C**) Period scatter plots for PER2::LUC (left) and *Bmal1-ELuc* (right) with $1/\sqrt{(t - t_0)}$ fits (consistent with SNIC). (**D**) Circadian power scatter plots for PER2::LUC (left) and *Bmal1-ELuc* (right) with sigmoid fits versus embryonic age; emergence times (T_e) were E11.70 ± 0.14 and E13.52 ± 0.70, respectively (best fit estimate ± SE). (**E**) Heat map of the clustering of genes characterizing the developmental state (top) and comparison of transcript levels of stem cell and progenitor markers

(blue), differentiation markers (red) in the 4VCP across embryonic stages, normalized to the highest
expression between E9.5 and E16.5 (down). Each data point represents a mean of 5–15 litters (3–11
embryos per litter). Whiskers plot displays the minimum to maximum values, the 25[th] and 75[th]
percentile and the median. (**F**) Developmental trajectories of circadian power for PER2::LUC oscillation
in the LVCP and the SCN (upper panels) and liver and kidney (lower panels), overlaid with sigmoidal
fits. (**G**) Comparison of the half-maximal times for progenitor markers (E13.63 ± 0.79), differentiation
markers (E13.52 ± 0.20), emergence time of *Bmal1-ELuc* and PER2::LUC oscillation in the 4VCP, and
of the PER2::LUC emergence time for each tissue.

Figure 3. Expression patterns of clock genes in the embryonic 4VCP. (**A**) Absolute mRNA levels

of clock gene transcripts across embryonic stages E9 to E16. Bars show the mean ± SEM of 8–20

litters, each litter containing 3–11 embryos. (**B**) Hierarchical clustering and heat map of mRNA Z-scores
across embryonic stages and ZT for the clock genes. Each data point represents the mean of 2–5
litters, each with 3–11 embryos. (**C**) Circadian power for each embryonic clock gene, grouped by
cluster (top); the representative gene of each cluster is plotted in red. Circadian phase-weighted power
spectrogram across embryonic days for the representative gene of each cluster (bottom); the left white
box marks the initial phase and the right white box marks the shifted phase in the end. (**D**) Heat maps
of pairwise correlations between each embryonic gene and *Cry2* (left), *Clock* (middle) and *Rev-Erba*
(right), the representative genes. (**E**) Temporal correlation networks of embryonic genes. Nodes denote
gene clusters. Edge color and width encode the sign and absolute value of the average Pearson within
the specified developmental window, respectively (blue negative, red positive, greater width indicates
larger magnitude). Solid edges indicate correlations with consistent signs across stages; dashed
edges indicate a sign change across stages. Black outlines indicate genes with circadian power ≥
20%. (**F**) Circular phase plots of embryonic gene expression across developmental stages. Each
phase plot has 0 rad at 12 o'clock and increases clockwise. Representative genes are shown with
larger opaque markers; non-representative genes are smaller and semi-transparent. The left-half
shading indicates earlier phases. Marker shape and color indicate cluster membership; bold outlines
highlight genes with circadian power ≥ 20%.

Figure 4. Correlations between maternal and embryonic gene expression vary with embryonic development. (**A**) Hierarchical clustering and heat map of the mRNA Z-scores across embryonic

stages and ZT for maternal clock genes. Each data point represents the mean of 2-3 dams. (**B**)
Circadian power for each maternal clock gene per cluster (top). The representative gene of each
cluster is shown in red. Circadian phase-weighted power spectrogram across embryonic days for the
representative gene of each maternal gene cluster (bottom). (**C**) Heat map of the correlation of maternal
Per2 and *Dbp* with *Bmal1*. (**D**) Heat map of the correlation of each embryonic gene with maternal *Per2*
(left), *Bmal1* (middle), and *Dbp* (right). (**E**) Temporal correlation networks of maternal-embryonic gene
expression correlations. Nodes colored by embryonic clusters; edges show average Pearson
correlation (red positive, blue negative, thickness magnitude). Solid edges indicate consistent sign,
dashed indicate changing sign. Black outlines mark genes with circadian power ≥ 20%. Only the top
30% strongest correlations are highlighted; others are semi-transparent gray. (**F**) Dual-radius phase
plots of maternal and embryonic gene expression. Inner ring, embryonic cluster representatives; outer
ring, selected maternal genes. Maternal-embryonic pairs are connected with dotted lines when both
exhibit circadian power ≥ 20% and their phase difference is < $\pi/2$ rad (≤ 6 h). (**G**) PER2::LUC and
Bmal1-ELuc circadian power in the embryonic (top, dark pink) and maternal (down, light pink) placenta
from 97 embryonic and 47 maternal placentas from E10.5 to E18.5.

Figure 5. E14 and E16 4VCP, but not E18 4VCP, are driven to a 26-h period temperature cycle.
(**A**) Raw PER2::LUC time series from 4VCP (black) and lung (orange) explanted at E14 (left) and E18
(right) for 12 days with a 0.5°C square-wave cycle (26 h period) after 2 days at 37 °C. Red bars indicate
the 37.5 °C epochs. (**B**) Double-plots of the peak phase of PER2::LUC time series from 1–4 embryonic
4VCPs and lungs at E14 (left), E16 (middle), and E18 (right). Red bars indicate the 37.5°C epochs. (**C**)

Violin plots showing amplitude (left) and period (right) per ROI in embryonic 4VCP across embryonic stages from E11.5 to E18.5 (579–2,787 cells from 3–4 4VCPs from 2–3 litters). The dotted line represents the median and the black lines represent the quartiles. Representative maps showing the single-cell level amplitude (top left) and period (top right) across stages. (**D**) Bar plot showing the mean ± SEM synchronization index in embryonic 4VCP overlaid with scatter plots showing individual 4VCP data across stages (579–2,787 cells from 3–4 4VCPs from 2–3 litters). (**E**) Representative 4VCP maps showing the normalized phase after 4 days in culture at E12.5, E13.5 and E14.5. Scale bar: 2 mm.