Endothelial Type I Interferon signaling modulates the vascular response to ischemic brain injury

Mary Claire Tuohy[1,2,3], Ping-Chang Kuo[4], Adrian Chelminski[3], Eti Muharremi[3], Claudia De Sanctis[5], Aomeng Cui[1,3], Alicia Russo[3], Danny Jamoul[3], Elizabeth Hillman[6,7], John F. Crary[5], Jui-Hung Jimmy Yen[4] and Dritan Agalliu [3,8,*]

[1] Vagelos College of Physicians and Surgeons, [2] Department of Neuroscience, [3] Department of Neurology, Columbia University Irving Medical Center, New York, NY USA

[4] Department of Microbiology and Immunology, Indiana University School of Medicine, Fort Wayne, IN, USA

[5] Neuropathology Brain Bank & Research CoRE, Mount Sinai School of Medicine, New York, NY, USA

[6] Mortimer B. Zuckerman Mind Brain Behavior Institute, [7] Department of Biomedical Engineering and Radiology, Columbia University, New York, NY, USA

[8] Department of Pathology and Cell Biology, Columbia University Irving Medical Center, New York, NY, USA

*Corresponding author: E-mail: da191@cumc.columbia.edu

Mailing address: Columbia University Irving Medical Center

650 West 168th Street, Black Building Room 310, New York, NY 10032

Telephone: 212-305-0323

ORCID: 0000-0002-5375-4143

Abstract

Vascular normalization [stabilization of aberrant angiogenesis and restoration of blood-brain barrier (BBB)] is critical for reducing long-term secondary sequelae after ischemic stroke. How immune and developmental signaling pathways coordinate these processes is poorly understood. Here we identify a unique brain endothelial cell (BEC) type one interferon (IFN1) signature in human and mouse ischemic stroke tissue. By leveraging two clinically-relevant murine ischemic stroke models, single-cell transcriptomics, and BBB functional assays, we find that deletion of endothelial IFN1 receptor (Ifnar1) exacerbates post-stroke BBB disruption and expands a BEC population expressing angiogenic and immature BBB markers. Conversely, IFNβ administration immediately after stroke reduces acute BBB disruption. Activation of IFN1 signaling in mouse BECs *in vitro* promotes junctional stabilization and reduces vascular endothelial growth factor (VEGF) signaling to enhance barrier properties, but suppress angiogenic features. Thus, endogenous endothelial IFN1 signaling modulates BBB dysfunction and angiogenesis to promote vascular normalization after ischemic brain injury.

MAIN TEXT

Ischemic stroke is a leading cause of mortality and morbidity [1]. Persistent cerebrovascular dysfunction has been proposed as a covert pathophysiological mechanism driving post-stroke morbidity [2-5]. Yet, our understanding of the underlying molecular mechanisms that reduce acute vascular dysfunction and promote long-term vascular normalization after ischemic brain

injury is limited. To restore central nervous system (CNS) homeostasis, surviving brain endothelial cells (BECs) must reacquire blood-brain barrier (BBB) properties through restoration of mature tight junctions (TJs), suppression of caveolae-mediated transcytosis, and expression of distinct transporters [6]. Studies also suggest that peri-infarct angiogenesis, triggered after ischemic stroke, positively correlates with functional outcomes [7, 8]. However, it is unclear how angiogenesis is temporally coordinated with BBB re-establishment, and whether deviations in this process may contribute to variability in acute brain injury and post-stroke cognitive impairment.

Developmental CNS angiogenesis and barriergenesis are tightly coordinated by Wnt/β-catenin signaling [9-11]. Endothelial Wnt/β-catenin signaling is repurposed in the adult brain to accelerate BBB repair and improve cerebral blood flow perfusion after ischemic stroke [12, 13]. In contrast to development, Wnt signaling in the ischemic brain occurs on a backdrop of acute inflammation [14-17]. Yet how immune and developmental programs cooperate to regulate post-stroke vascular remodeling and normalization remains poorly understood. The inflammatory response to danger associated molecular patterns (DAMPs) released during tissue injury in many ways parallels that observed during microbial infection. Amongst these shared signaling pathways is the type 1 interferon (IFN1) family [18]. IFN1 signaling is classically activated by microbial products that bind pattern recognition receptors and cytosolic nucleic acid sensors. Binding of IFN1s to the IFN1 receptors (Ifnar1 & Ifnar2) triggers their dimerization, autophosphorylation of tyrosine kinase 2 (Tyk2) and Janus kinase 1 (Jak1) and activation of signal transducer and activator of transcription (Stat1) family to induce transcription of interferon stimulated genes (ISGs). Although best understood for coordinating anti-microbial responses, ISGs are a highly conserved transcriptional signature observed across many brain diseases characterized by acute (i.e. ischemic

stroke, traumatic brain injury) and chronic (i.e. multiple sclerosis, neurodegeneration, aging) sterile inflammation [19-24].

Acute administration of exogenous IFNβ attenuates ischemic brain injury [25, 26]. In line with mounting evidence that IFN1 signaling regulates microglia function, this rescue is partially mediated via modulation of microglia polarization [27]. However, the IFN1 receptor is expressed on all nucleated cells. Single-cell RNA sequencing (scRNA-seq) studies have found an IFN responsive BEC population after stroke, highlighting a potential role for endothelial IFN1 signaling both in response to the primary ischemic insult and exogenous IFNβ treatment [28]. Acute endothelial IFN1 signaling promotes peripheral tumor vascular normalization and BBB stabilization following neurotrophic viral infection [29-31]. In contrast, chronic high levels of IFN1 stimulated either by autoimmunity or comorbid systemic infection impede vascular repair in systemic lupus erythematosus (SLE) and traumatic brain injury (TBI) [32, 33]. These findings demonstrate that the biological impact of IFN signaling depends on IFN type, cellular identity, signaling duration, and physiological context.

Here we show endogenous IFN1 signaling is upregulated in BECs after ischemic stroke. Amplification of this signaling by exogenous IFNβ administration ameliorates acute BBB disruption after stroke. *In vitro* activation of the IFN1 pathway in primary BECs suppresses inflammation-induced barrier disruption and angiogenic responses. To isolate the functional significance of endogenous IFN1 signaling on vascular function following ischemic brain injury, we generated IFN1 receptor (Ifnar1)-inducible EC knockout mice. Ablation of post-ischemic BEC

IFN1 signaling worsens acute BBB disruption and expands angiogenic BECs with immature BBB markers. These findings demonstrate that endogenous BEC IFN1 signaling modulates BBB dysfunction and angiogenesis to promote vascular normalization after ischemic brain injury.

RESULTS

Ischemic brain upregulates endogenous IFN1 and shows conserved BEC IFN signature

Although the IFN1 receptor is expressed on all nucleated cells, the role of IFN1 signaling has primarily been studied in myeloid cells [34]. To investigate the potential contribution of endothelial IFN1 signaling after ischemic brain injury, we analyzed initially several published murine single-cell RNA-sequencing (scRNA-seq) datasets in homeostasis and ischemic stroke [28, 35]. In accordance with organ specific EC transcriptional heterogeneity, the IFN1 receptor, *Ifnar1*, is enriched on BECs compared to ECs in other tissues (Figure 1a). This pattern is not observed with the IFN2 receptor, *Ifngr1*, which is uniformly enriched in ECs from all organs (brain, heart, large intestine, and lung; Extended Data Figure 1a). Analysis of a stroke dataset revealed that border associated macrophages (BAMs) constitutively make IFNβ and contribute to its upregulation after ischemic brain injury (Extended Data Figure 1b). In contrast, the aged brain loses this IFN1 signature, preferentially tipping toward IFN2 (IFNγ) signaling by T cells (Extended Data Figure 1b) [28]. To extend and validate this analysis, we examined the spatial distribution and temporal kinetics of IFNβ and several downstream ISGs (Irf9, Isg20, Cxcl10, and Stat1) by RNA *in situ* hybridizations (RNA-ISH) in two distinct murine models of ischemic stroke, photothrombotic ischemia (RBPT) and transient-middle cerebral artery occlusion (t-MCAO) [25, 36]. Within both models, *IFNβ* mRNA was upregulated within the ischemic territory as early as

24 hours after brain injury, with its expression peaking at 72 hours and reduced at post-stroke day 7 (Figure 1b). In contrast, the analyzed ISGs exhibited a broader spatial distribution in the adjacent non-ischemic parenchyma with the highest expression observed closest to the stroke tissue border (Extended Data Figure 1c, d; data not shown).

Do BECs respond to this endogenous upregulation of IFNβ? To examine this question, we performed scRNA-seq on fluorescence-activated cell sorting (FACS)-purified CD31+CD11b-BECs isolated from the ipsilateral cortex at 72 hours after RBPT and from age-matched sham controls (healthy) (Figure 1c-d, Supplementary Table 3a). Consistent with previous findings [28], we observed differential expression of several ISGs across BEC clusters under homeostatic conditions, but this signature was significantly upregulated following ischemic injury (Figure 1e, f, cluster 3). Immunofluorescence staining for one of the ISGs, Ifitm3, confirmed low expression in the normal brain vasculature, which was significantly increased within the ischemic territory by 72 hours in both t-MCAO and RBPT models (Figure 1g, h).

To determine the relevance of these findings for human stroke, we performed immunostaining for CD31, IFITM3, and STAT1 in postmortem brain specimens from a small cohort of individuals with ischemic stroke and age-matched controls (Figure 2a, Extended Data Table 1). Images were acquired from ischemic stroke lesions, adjacent normal-appearing tissue (NAT) confirmed by hematoxylin and eosin staining of a serial section, and brain region-matched controls. As observed in murine brain tissue (Figure 1g, h), baseline homeostatic expression of IFITM3 was present in human brain vasculature in regions unaffected by ischemia (Figure 2a).

However, vascular IFITM3 coverage and intensity were significantly increased in ischemic lesions, accompanied with changes in vessel morphology, compared to adjacent NAT and age-/region-matched control brains (Figure 2b-d). Thus, upregulation of endogenous IFN1 signaling and the responsiveness of a subset of BECs to this pathway are conserved features of both murine and human ischemic stroke.

Endothelial IFN1 signaling mitigates inflammation-mediated barrier disruption

We next sought to determine the impact of acute activation of IFN1 signaling on the functional barrier properties of BECs. TNFα and IL1β have been shown to mediate BBB disruption after ischemic stroke and in primary BECs [37]. To model the inflammatory microenvironment of the ischemic brain, primary murine BECs (mBECs) were grown to confluency and transendothelial electrical resistance (TEER) was measured upon treatment with TNFα/IL1β ("Cyt", each used at 10ng/mL) in the presence or absence of IFNβ (250U/mL). The addition of TNFα/IL1β led to a significant reduction in mBEC TEER (TEER = -1041 Ω*cm^2/hrs +/- 7.406; Figure 3a). In contrast, mBECs treated concurrently with TNFα/IL1β and IFNβ showed a unique rebound effect, characterized by a significant, albeit transient, recovery in TEER values (397 Ω*cm^2/hrs +/- 4.808; Figure 3a). Western blotting revealed that the recovery observed in IFNβ-treated mBECs was not due to an increase in total protein levels of Claudin-5 (Cldn5), a key BBB tight junction protein (Figure 3d). Instead, the recovery specifically correlated with increased junctional Cldn5 localization and a reduction in intracellular Cldn5 aggregates, as seen by immunofluorescent staining 24 hours post-IFNβ treatment (Figure 3b) Similarly, treatment of human brain microvascular endothelial cells (HBMECs) with IFNβ ameliorated TNFα/IL1β-induced reduction in TEER (90 Ω*cm^2/hrs +/- 5.773; Figure 3c). This rescue was not observed in human umbilical

vein endothelial cells (HUVECs; Figure 3c). Pure IFNβ treatment also increased TEER values in both mBECs (32.30 Ω^*cm^2/hrs +/- 3.852) and HBMECs (846 Ω^*cm^2/hrs +/- 7.951), but not in HUVECs, underscoring that IFN1-mediated BEC barrier modulation occurs independently of inflammatory cytokines and barrier disruption (Extended Data Figure 2a-b). In contrast to IFNβ, IFNγ treatment resulted in a significant decrease in BEC TEER (-63.10 Ω^*cm^2/hrs +/- 3.406), indicating that specifically IFN1 signaling, but not IFN2, strengthens BEC barrier function (Extended Data Figure 2a). Although ISGs are a common transcriptional signature of IFN signaling in RNA sequencing datasets, the identification of ISGs is not unique to IFN signaling (e.g. NF-κB pathway, Extended Data Figure 2c) and does not distinguish the class of IFN signaling (IFN1, IFN2, or IFN3) [38]. This is an important distinction as IFN1 and IFN2 signaling have an opposite effect on barrier function in primary mBECs (Extended Data Figure 2a).

To examine potential downstream genes underlying IFNβ-mediated modulation of BEC barrier function, we collected control, TNFα/IL1β, and TNFα/IL1β + IFNβ-treated mBECs for bulk RNA sequencing and performed differential gene expression (DGE) analysis followed by gene set enrichment analysis (GSEA) (Supplementary Table 1a-d). As expected, the addition of IFNβ resulted in a significant upregulation of ISGs compared to mBECs treated with TNFα/IL1β alone (Extended Data Figure 2c, Supplementary Table 1d). Activation of IFN1 signaling also upregulated genes involved in antigen presentation and chemokine signaling more broadly, which resulted in an enhanced inflammatory transcriptional signature (Figure 3e). Beyond immunomodulation, genes associated with Wnt signaling were significantly upregulated in BECs treated with IFNβ (Figure 3f). A similar transcriptional signature was observed in mBECs treated with pure IFNβ (Extended Data Figure 2d-e, Supplementary Table2a-b). Interestingly, the

enrichment of Wnt signaling was primarily driven by genes related to β-catenin independent (non-canonical) planar cell polarity (PCP) signaling (e.g. *Cdc42, Daam1, Rhoa, Rock1*), rather than canonical Wnt/β-catenin signaling which promotes BBB acquisition and maintenance (Figure 3f, Extended Data Fig 2e) [39]. This led us to hypothesize that IFN1 signaling may strengthen BEC junctions via regulation of local actin cytoskeleton interactions. To test this hypothesis, we assessed whether antagonism of Rho GTPases, central coordinators of actin cytoskeleton dynamics and cell junctions, modify the barrier response to IFNβ treatment. Pharmacological inhibition of Rho (RhoI), but not Rac1 (Rac1I), attenuated the IFNβ-induced increase in TEER, while preserving the overall shape of the response curve (Figure 3g). Altogether, these observations suggest that BEC IFN1 signaling may promote barrier function through Rho-dependent actin remodeling which reinforces tight junction proteins at the cell membrane (Figure 3h).

IFNβ treatment may reduce acute BBB dysfunction after ischemic brain injury

Administration of exogenous IFNβ has been shown to modulate brain injury in the t-MCAO model of ischemic stroke [25]. To understand how IFNβ overexpression alters BBB dynamics after stroke *in vivo*, we administered IFNβ 3hrs post t-MCAO and examined the biodistribution of 5-(and-6-) tetramethylrhodamine biocytin (biocytin-TMR, MW = 869 Da) tracer injected intravenously at 72 hours post-t-MCAO (Figure 4a). IFNβ administration significantly improved the neuroscore at 72 hours post t-MCAO (Figure 4b), consistent with prior findings at the 48-hour timepoint [25]. Additionally, IFNβ-treated mice showed a trend toward decreased biocytin-TMR leakage area along the rostral-caudal brain axis compared to vehicle-treated mice, with the difference approaching significance (p-value = 0.0533; Figure 4c-e). Brain sections at three antero-posterior positions relative to bregma showed a significant reduction in the area of tracer leakage across the

BBB (Figure 4c, d) To characterize the effects of exogenous IFNβ treatment on ischemia mediated tight junction (TJ) remodeling, we quantified the fraction of vessel segments with either intact Cldn5+ junction strands ("intact"), Cldn5+ junction strands with large gaps (> 2.5 μm, "gap"), or absent junctions ("absent") in both the ischemic ipsilateral cortex and the contralateral cortex as previously described [6] (Figure 4f). Intact Cldn5+ strands, representing mature and stable vascular junctions, covered the majority of vessels in the contralateral cortex of both vehicle and IFNβ-treated mice (Figure 4g). In contrast, the ischemic ipsilateral cortex showed an increased percentage of vessels with gaps in Cldn5+ junctional immunostaining (Figure 4g-h''). Compared to controls, IFNβ-treated mice showed an increased percentage of vessels with intact Cldn5+ strands and a decreased percentage of vessels with Cldn5+ gaps in the ipsilateral cortex (Figure 4g-h''). Therefore, IFNβ treatment reduces acute structural and functional BBB damage after ischemic injury.

Given the well-established role of IFN1 signaling in myeloid cells, we further assessed myeloid cell activation in vehicle- and IFNβ-treated mice. CD68 immunoreactivity within Iba1+ cells was significantly increased in the ipsilateral compared to contralateral cortex of both vehicle- and IFNβ-treated mice at 72 hours post t-MCAO (Extended Data Figure 3a-b). However, this myeloid cell activation was significantly reduced in IFNβ-treated mice compared to controls and corresponded with decreased biocytin-TMR leakage (Extended Data Figure 3b-c). Despite the systemic response to ischemic stroke and the peripheral administration of IFNβ, contralateral cortices showed no significant differences in CD68 immunoreactivity across treatments (Extended Data Figure 3a, b). Concurrent changes in TJ morphology and myeloid cell reactivity suggest that

BBB rescue observed with IFNβ treatment may be mediated either *directly* through endothelial IFN1 signaling or *indirectly* through myeloid IFN1 signaling (Extended Data Figure 3d).

Ablation of endogenous BEC IFN1 signaling modestly exacerbates acute post-stroke BBB dysfunction

Having established that the ischemic brain upregulates IFNβ and that activation of endothelial IFN1 signaling modulates BEC barrier properties *in vitro* and *in vivo*, we next sought to determine the impact of *endogenous* BEC IFN1 signaling on acute BBB dysfunction after ischemic stroke. To isolate the biological impact of endogenous IFN1 specific signaling on BECs *in vivo*, we generated *Ifnar1*-inducible EC knockout (*Ifnar1^{iECKO}: Ifnar1$^{fl/fl}$; VECad-CreERT2$^{+/-}$*) mice and control (*Ifnar1$^{fl/fl}$*) mice (Figure 5a) . We confirmed that a course of consecutive intraperitoneal injections of 3mg of tamoxifen over five days (15mg in total) induced robust Cre-mediated recombination in BECs (Extended Data Figure 4a-b). Following a tamoxifen washout period of at least 7days, t-MCAO was then induced, and as in previous experiments biocytin-TMR was injected via tail vein to assess paracellular BBB permeability in *Ifnar1^{iECKO}* and *Ifnar1$^{fl/fl}$* mice 72 hours after reperfusion. In contrast to exogenous IFNβ treatment, the ablation of endothelial IFN1 signaling did not significantly affect the neuroscore during the first 72 hours after t-MCAO (Figure 5b). *Ifnar1^{iECKO}* mice showed a trend, albeit not significant, toward increased biocytin-TMR leakage area along the rostral-caudal brain axis compared to *Ifnar1$^{fl/fl}$* control mice (p=0.0525; Figure 5d-e). This effect size was comparable to the barrier rescue observed with IFNβ treatment (Figure 4e). Together, these data suggest that activation of endothelial IFN1 signaling either by an

endogenous or exogenous source of IFNβ has a relatively modest effect on BBB function 72 hours after ischemic brain injury.

IFN1 signaling suppresses angiogenic features of BECs

Activation of IFN1 signaling in mBECs strengthened their barrier integrity *in vitro* in the presence of inflammatory cytokines (IL1β/TNFα), despite amplifying endothelial transcriptional inflammatory signatures (Figure 3e). To gain insight into pathways regulated by IFN1 signaling in BECs, we returned to the GSEA of cytokine-treated mBECs *in vitro*. Inflammatory signaling and vascular remodeling are intimately intertwined to preserve homeostasis [40] Correspondingly, genes involved in cell growth and proliferation (e.g. *Erh, Gspt1, Kpnb1, Pwp1, Rack1*) were upregulated in mBECs treated with TNFα/IL1β (Figure 6a). This Myc activity signature was reversed in mBECs concurrently treated with IFNβ (Figure 6a), consistent with IFN-regulated antiproliferative responses reported in other cell types [41]. Unique to BECs, marker genes known to be enriched in tip cells (*Apln, Dll4, Lama4, Ecscr*), as well as downstream targets of vascular endothelial growth factor A (VEGF-A) were highly downregulated in response to IFNβ (Figure 6a-c). A similar antiproliferative transcriptional response was observed in mBECs treated with IFNβ alone, and this response persisted at the 24 hour timepoint even as ISG expression began to normalize (Extended Data Figure 2d, f). By western blotting, IFNβ-treated mBECs show decreased levels of VEGF receptor 2 (Vegfr2) (Figure 6d). To understand the significance of this angiogenic switch, we performed functional assays to assess migration and proliferation of mBECs, and VEGF-A-mediated vascular permeability. Priming of mBECs with IFNβ significantly delayed wound closure (Figure 6e) and suppressed VEGF-A-induced acute permeability (Figure 6f).

Notably, the rescue in permeability was observed with IFNβ priming but not with simultaneous VEGF-A and IFNβ application (Figure 6f), potentially due to temporal lag of IFNβ-mediated Vegfr2 degradation. To understand the implications of this signaling for sprouting behavior in human cells, we used a fibrin bead angiogenesis assay. HBMECs treated with IFNβ had a reduced sprout length and network length compared to untreated HBMECs in the fibrin bead assay (Figure 6g-i). Together, these findings suggest that IFN1 signaling modulates angiogenic features of BECs potentially through crosstalk with VEGF-A and Myc signaling pathways.

Ablation of BEC IFN1 signaling results in expansion of angiogenic BECs after ischemic brain injury

Next, we evaluated the molecular impact of endogenous BEC IFN1 signaling on the vascular response after ischemic brain injury. Unlike t-MCAO, RBPT induced a robust peri-infarct BEC proliferation by 72 hours after ischemia (Extended Data Figure 5a), providing an opportunity to examine how IFN1 signaling modulates angiogenic and barrier properties of BECs in this model of ischemic stroke. We performed scRNA-seq of FACs-sorted BECs (CD31+ CD11b-) isolated from dissected ipsilateral cortices (2-3 mice pooled/sample; 4 samples/ genotype) of *Ifnar1*^{iECKO} and *Ifnar1*$^{fl/fl}$ mice 72 hours after photothrombosis (Figure 7a, Extended Data Figure 5b, Supplementary Table 3b). Following evaluation of gene number and mitochondrial read counts, duplicates and low-quality cells were removed. Unique transcripts were normalized for total read depth and clustering was visualized with uniform manifold approximation and projection (UMAP). Contaminating smooth muscle cells (*Acta2, Ccl6*), pericytes (*Pdgfrb, Atp13a5*), immune cells (*Hexb, Tmem119, P2ry12, Csf1r, Trem2, Ly6c2, Ccr2, CD3a, CD4, CD8a*), and astrocytes (*S100b, Gfap, Aldh1l1*) were removed to obtain a total of 30,975 high-quality BECs. The dataset

was then corrected for batch effects with Harmony (Extended Data Figure 5c, [42]), and isolated BECs were re-clustered, which revealed 10 distinct clusters detected in each sequenced sample, though not at the same abundance (Figure 7b). Loss of IFN1 signaling induced a marked transcriptional shift in BECs (Fig 7c-d). This shift could not be explained by differences in EC subtype proportion. A similar proportion of BEC subtypes along the artery-capillary-venous tree (Arterial: *Bmx, Efnb1, Hey1, Amd1, Eln, Clu*; Capillary: *Slc1a1, Mfsd2a*; Veinous: *Icam1, Nr2f2, Lcn2, Prcp, Bst1, Il1r1, Lyve1*) was obtained from both genotypes (Extended Data Fig 5d-h). Subclustering and visualization of BECs by vascular subtype (3,690 arterial BECs; 26,437 capillary BECs; 848 venous BECs) revealed that the capillary compartment was the most sensitive to changes in IFN1 signaling and mainly contributed to the transcriptional shift at the BEC population level (Extended Data Figure 6a-c).

To gain insight into IFN1-regulated BEC biological processes after ischemic brain injury, we performed differential gene expression analysis of BECs isolated from *Ifnar1*[iECKO] and *Ifnar1*[fl/fl] mice for each vessel subtype and conducted GSEA (Supplementary Table 1a). Gene sets associated with cell growth ("Myc targets"), angiogenesis ("Tip Cell", "Angiogenesis", "Notch Signaling"), vascular remodeling related processes ("Non-Canonical Wnt Signaling", "Endothelial Cell Migration"), and inflammation ("Chemokine/Cytokine Signaling", "Antigen Processing & Presentation") were significantly upregulated in BECs derived from *Ifnar1*[iECKO] samples (Figure 7e, Supplementary Table 3c-e). Conversely, gene sets associated with BBB properties ("BBB", "Transporter", "Receptor-Mediated Transcytosis Receptors") were downregulated in *Ifnar1*[iECKO] samples, presenting an inverse correlation between angiogenesis and barrier function (Figure 7e, Supplementary Table 3c-e). The large number of capillary BECs in our scRNA-seq dataset

allowed more detailed characterization of capillary-specific heterogeneity following ischemic brain injury. Genes associated with angiogenesis were specifically upregulated in a subset of capillary BEC clusters (cBEC5, cBEC6, cBEC7, and cBEC8; Figure 7f, Supplementary Table 3f-i). cBEC7 was enriched in *Ifnar1*$^{fl/fl}$ samples and maintained high tight junction and specialized transporter gene expression (e.g. *Cldn5, Glut1*; Figure 7f). In contrast, cBEC5, cBEC6, and cBEC8 enriched in *Ifnar1*^{iECKO} samples showed a further upregulation of distinct angiogenic genes (i.e. *Angpt2, Rhoc, Ecscr, Hbegf, Apln*) and a decreased expression of genes related to BBB function (Figure 7f). We further compared BEC angiogenic signatures in the ischemic brain with those reported in experimental autoimmune encephalomyelitis (EAE, ECs isolated from spinal cord) and CNS developmental angiogenesis (BECs isolated from embryonic day 11.5 - 17.5) (Supplementary Table 4a-b) [43, 44]. Congruency analysis revealed a substantial transcriptome overlap of angiogenic BECs, with a greater number of upregulated genes shared between stroke and CNS development (108), than stroke and EAE (53) (Extended Data Figure 7a-d, Supplementary Table 4a-e). Moreover, spinal cord ECs in EAE showed a subtype-specific upregulation of ISGs in arterial and capillary BECs during acute, and to a lesser extent, chronic EAE (Extended Data Figure 7e) [43]. Consistent with the upregulation of the ISG signature in EAE spinal cord ECs, Ifitm3 was highly upregulated in blood vessels in acute, but not chronic EAE (Extended Data Figure 7f). These data suggest that while there may be a core transcriptional module underlying the angiogenic switch in BECs, different microenvironmental triggers likely regulate distinct BEC angiogenic processes and distinct CNS insults may elicit a convergent BEC IFN signaling response.

To validate scRNAseq findings *in vivo*, we used fluorescent in situ hybridization (FISH) and immunofluorescence to assess expression of three angiogenic markers (*Apln, Vegfr2, Angpt2*) highly upregulated in *Ifnar1^{iECKO}* BECs (Figure 7f, 8b-d). *Ifnar1$^{fl/fl}$* and *Ifnar1^{iECKO}* brain tissue were collected 72 hours after RPBT and images were acquired from the contralateral cortex (#1), peri-infarct cortex (#2), and ischemic lesion (#3) (Figure 8a). Basal expression of Vegfr2 is observed across vasculature in the contralateral cortex, while Apln and Angpt2 vascular expression were low-to-absent (Figure 8e-g). Peri-infarct vessels in *Ifnar1^{iECKO}* mice showed a significant upregulation of Vegfr2 relative to contralateral vessels that was not observed uniformly in *Ifnar1$^{fl/fl}$* mice (Figure 8e). Within this same peri-infarct region, a subset of BECs upregulated the tip cell marker, *Apln* (Figure 8f). In line with scRNAseq findings, *Ifnar1^{iECKO}* peri-infarct BECs showed significantly increased *Apln* mean fluorescence intensity (MFI) and vascular surface area (SA) compared to *Ifnar1$^{fl/fl}$* peri-infarct BECs (Fig 8f). In contrast to peri-infarct Vegfr2 protein and *Apln* mRNA localization, Angpt2 protein was upregulated on vessels within the ischemic lesion (Figure 8g). This highlighted two molecularly distinct compartments of vascular remodeling after photo-thrombosis, suggesting region specific vascular responses to ischemic brain injury not captured by scRNAseq. Analysis of vascular Angpt2 in the ischemic lesion, however, showed no significant difference in expression across *Ifnar1$^{fl/fl}$* and *Ifnar1^{iECKO}* mice (Figure 8f). Taken together, these results show that endogenous BEC IFN1 signaling fine-tunes the peri-infarct angiogenic response to brain ischemia (Figure 8g).

DISCUSSION

ISGs have emerged as a highly conserved transcriptional signature upregulated in many brain diseases [19-24, 45, 46]. Most studies investigating the functional significance of this IFN-

dependent signaling have been limited to the myeloid cell compartment. In this study, we report that ischemic brain injury upregulates endogenous IFN1 signaling and robust ISG expression in BECs. Ablation of post-ischemic BEC IFN1 signaling results in the expansion of angiogenic BECs and an exacerbation of barrier function. Conversely, exogenous IFNβ treatment partially rescues in-vivo barrier dysfunction and suppresses inflammatory-induced BEC barrier disruption and angiogenic responses in vitro. These results indicate that IFN1 signaling regulates vascular remodeling in the acute period after ischemic brain injury.

First discovered for their crucial role in antiviral immunity, IFN1s are best known for their immunomodulatory properties [47]. Our work adds to a growing body of literature which further supports the role of IFN1-dependent regulation of vascular homeostasis. In a model of West Nile virus encephalitis, IFN1 signaling stabilizes tight junctions and reduces viral neuroinvasion [30]. However, excessive or sustained IFN1 signaling can result in endothelial dysfunction, vascular leakage, and septic shock [48, 49]. In cancer, endothelial IFN signaling coordinates anti-tumor activity through promoting vascular normalization and tumor immunogenicity [29, 50, 51]. Yet in a rodent model of Aicardi Goutieres syndrome (AGS), chronic BEC IFNα signaling provokes pathological vascular inflammation, microaneurysm formation, BBB disruption, and neurotoxicity [52]. Thus, distinct cell types and microenvironments differentially modulate IFN1 signaling intensity, duration, and crosstalk to generate context-specific responses [53]. Here we find that IFNβ treatment of mBECs and HBMECs, but not HUVECs, promotes barrier function *in vitro*. As BECs are the primary interface between the CNS parenchyma and blood, it is possible that enrichment of the IFN1 receptor promotes tonic BEC IFN1 signaling to coordinate unique CNS immune surveillance. Indeed, microbiota induced IFN1 production metabolically instructs

immune populations to be poised for future pathogen contact [54]. It will be interesting to understand whether endothelial cell heterogeneity alters IFN1 signaling to mediate tissue-specific function.

Due to overlapping cytokine-induced effectors, the detection of ISGs cannot distinguish between IFN families or even IFN-independent pathways, making it challenging to attribute downstream effects to a specific cytokine [55]. In this study, we use *Ifnar1*^{iECKO} mice to specifically isolate IFN1-mediated BEC function and demonstrate that mice deficient in post-stroke BEC IFN1 signaling upregulate angiogenic markers. These markers are suppressed when BECs are treated *in vitro* with IFNβ. Our data suggests that IFN1 may modulate angiogenic processes through regulation of VEGF-A/VEGFR2 signaling (Figure 8h). Previous studies report that post-stroke VEGF-A treatment enhances peri-infarct angiogenesis, reduces ischemic brain damage, and improves functional outcomes [56-58]. However, others studies have found that antagonism of endogenous VEGF-A reduces infarct volume and administration of VEGF-A significantly increases BBB leakage and hemorrhagic transformation [59, 60]. These contradictory findings highlight the multi-step nature of angiogenesis which inextricably couples changes in basement membrane composition with EC adhesion, proliferation, migration and maturation through VEGF-A signaling [61]. Critically, endogenous IFNβ peaks within the lesion core at 72 hours after ischemic brain injury and decreases by 7 days. We postulate that this tight temporal and spatial control may allow for suppression of VEGF-A mediated vascular permeability during the acute period (24 – 48 hours post-stroke) when hemorrhagic risk is high, but keep intact the delayed VEGF-A signaling to support vascular growth and functional recovery in the subacute phase of ischemic stroke (Figure 8g). BEC IFN1 signaling may further regulate angiogenic and

barrier properties through direct crosstalk with Rho and/or Myc signaling to induce cytoskeletal reorganization, critical for junctional stability and / or cell migration, and metabolic reprograming respectively (Figure 8h). Ablation of BEC IFN1 signaling also resulted in upregulation of inflammatory transcripts, including ISGs, after ischemic injury. This aligns with previous studies showing that ISGs are downregulated only when multiple IFN families are deficient, highlighting the transcriptional redundancy of ISGs and potential cross-regulation between IFN signaling [62].

Our scRNA-seq analyses further provides greater molecular insight into post-stroke angiogenesis. In particular, the expression of tip-cell enriched genes (i.e. *Dll4, Apln, Angpt2, Pgf)* are highly prevalent in the ischemic brain. During developmental angiogenesis, tip cells at the leading front of the vascular sprout direct migration along a VEGF-A gradient, while stalk cells behind proliferate to propel the plexus forward. Binding of VEGF-A to tip cell enriched Vegfr2 activates both phosphatidylinositiol 3-kinase (PI3K)/AKT signaling as well as Rac/Rho/Cdc42 to stimulate actin reorganization, filopodia formation, and endothelial cell migration [63-66]. Transcripts encoding for proteases involved in ECM remodeling (*Sparc, Adamts9, Adamts1, Mmp15*), chemotactic receptors (*Nrp1, Ecscr, Plxnd1*), basement membrane components (*Col4a1, Col4a2, Lama4*), and cytoskeletal related proteins (*Tuba1a, Dynll1, Arpc3, Vcl*) are amongst tip-cell enriched genes after ischemic brain injury. Our congruency analysis shows that this BEC angiogenic transcriptional signature is more similar to that seen in development than a model of neuroinflammation, EAE [43, 44]. This may be because tissue ischemia is the primary stimulus for angiogenesis during development and stroke, whereas inflammation drives it in EAE. Consequently, the molecular mechanisms regulating CNS angiogenesis may be distinct across neurological diseases.

Our work shows that BEC IFN1 signaling modulates the vascular responses during the acute period after ischemic brain injury. Future studies will determine how ischemia-induced BEC IFN1 signaling may shape vascular trajectories in long term and post-stroke neurological recovery. Although older age and larger infarct volume are well-established predictors of poor functional outcomes after ischemic stroke, there is considerable variability among patients with comparable infarct burden. This heterogeneity likely reflects differences in baseline vascular integrity, cerebral autoregulation, and inflammatory processes, including IFN1 signaling. Understanding how these factors interact to affect longitudinal stroke outcomes will be an important point of future investigation. Moreover, given the prevalence of CNS IFN signatures, our findings may have broader implications for understanding conserved mechanisms that regulate vascular pathology and normalization across various neuroinflammatory brain states.

AVAILABILITY OF DATA AND MATERIALS

All data are available in the manuscript and the supplementary information. Any additional information required to reanalyze the data reported in this study will be available upon request.

COMPETING INTEREST

The authors declare no competing or financial interests relevant to the subject matter in this contribution.

FUNDING

M.C.T was supported by T32 GM007367 and a grant from the National Institute of Neurological Disorders and Stroke (F31NS13093). A.C, E.M., D.J., A.C., and D.A. were supported by grants from the National Eye Institute (R01EY033994), National Heart Lung and Blood Institute (R61/R33 HL159949), National Institute on Aging (RF1AG078352), and National Institute of Neurological Disorders and Stroke (R21NS130265).

AUTHOR CONTRIBUTIONS

M.C.T and D.A conceived the study and designed the experiments. M.C.T, P.C.K, A.C, E.M, C.D, A.C, A.R, D.J, J.Y, and D.A performed the experiments. M.C.T, A.C, E.M, A.C, and D.J analyzed the data. D.A supervised the project. J.Y, J.F.C, and E.M.C.H provided resources. M.C.T wrote the manuscript and D.A provided revisions. Project was administered and funding was acquired by D.A.

ACKNOWLEDGEMENTS

We thank Michael Kissner and staff from the Columbia Stem Cell Initiative Flow Cytometry Core, Erin Bush and staff from the Columbia Single Cell Analysis Core and JP Sulzberger Columbia Genome Center, and Theresa Swayne and Haojie Ji from the Herbert Irving Comprehensive Cancer Center Confocal and Specialized Microscopy Shared Resource (funded in part through NIH/NCI Cancer Center Support Grant P30CA013696) for their technical support. We also thank Sukanya Sarkar for critical reading of this manuscript.

METHODS

Mice: 4- to 6-month-old male age-matched mice were used for all experiments. C57BL/6J (B6) (JAX stock #000664), B6(Cg)-Ifnar1[tm1.1Ees]/J (Ifnar1[fl/fl]; JAX stock #028256) [67], and B6.Cg-Gt(ROSA)26Sor[tm14(CAG-tdTomato)Hze]/J (Ai14; JAX stock #007914) [68] mice were obtained from Jackson Laboratories. Cdh5(PAC)-Cre[ERT] (VEC-PAC) mice were kindly provided by Ralf Adams and have been previously described [69]. Ifnar1[iECKO] and Ai14 VEC-PAC reporter mice were generated by crossing VEC-PAC to Ifnar1[fl/fl] and Ai14 mice, respectively. Cre activity was induced in adult mice by consecutive intraperitoneal injections with 3mg of Tamoxifen (Millipore Sigma, T5648) dissolved in 150uL of corn oil (Millipore Sigma, C8267) over the course of five days (15mg of Tamoxifen in total). Cre negative tamoxifen injected Ifnar1[fl/fl] and Ai14 littermates were used for controls. Mice that received tamoxifen were allowed a one-week washout period prior to rose bengal photothrombosis (RBPT) and a three-week washout period prior to t-MCAO surgery. Mice were bred and maintained under specific pathogen-free conditions at CUIMC or Indiana University School of Medicine, under 12-hour light/12-hour dark cycle.

Ethics approval: All experimental procedures in mice were approved by the Columbia University and Purdue Institutional Animal Care and Use Committee.

Rose Bengal Photothrombosis (RBPT): Mice were anesthetized with isoflurane (3% induction, 1.5% maintenance) and affixed to a stereotaxic frame. Temperature was maintained at 37°C with a feedback temperature control system (Harvard Apparatus 55-7030). Following scalp retraction and skull exposure, a mask with a small aperture was placed on the somatosensory cortex: -2 mm anterior-posterior (AP), 3mm medial-lateral (ML). Rose Bengal (dissolved in 0.9% saline solution

at 15mg/mL) was injected intraperitoneally at 10uL/g body weight and allowed to circulate for 5 minutes. For sham surgeries, pure saline was injected. An optical fiber attached to a cold light source (150W) was aligned above the mask aperture and turned on for 15 minutes to achieve complete photothrombotic occlusion of surface vessels in region of interest [36]. The scalp incision was closed with vetbond tissue adhesive and mice were placed in 37°C recovery cage until anesthesia recovery.

Transient Middle Cerebral Occlusion (t-MCAO): Cerebral ischemia was induced as previously described [25]. Under anesthesia, mice were subjected to 40-minute intraluminal occlusion of the right middle cerebral artery through insertion of a silicone-coated nylon monofilament (Doccol Corp, Sharon, MA). Cerebral blood flow (CBF) was measured before, during and after ischemia using laser Doppler flowmetry. Mice with a total CBF reduction of less than 80% during surgery were excluded from the study as they did not meet the surgical criteria. The animal's body temperature was maintained at 37°C throughout the surgery by a warming lamp and heating pad. After surgery, mice were placed in a 37°C recovery cage for 1 hour. For IFNβ treatment studies, 10,000U of recombinant murine IFNβ (PBL Assay Science) suspended in 100μL of PBS was administered via intravenous injection 3hrs post reperfusion.

Neurological Assessment: Neurological function was examined directly after t-MCAO and again 24hrs, 48hrs, and 72 hrs later. Neurological deficits were scored as follows: 0, normal; 1, mild turning behavior with or without inconsistent rotation when picked up by tail with <50% attempts to curl to the contralateral side; 2, mild consistent curling with >50% attempts to curl to

contralateral side; 3, consistent strong and immediate curling behavior that is held for more than 1-2 secs; 4, severe curling progressing into barreling with loss of walking or righting reflex; 5, comatose or moribund [70].

Analysis of publicly available scRNA-seq data: For analysis of endothelial *Ifnar1* and *Ifngr1* expression, publicly available processed EC data and associated tissue metadata from Kalucka et al., 2020 was used [35]. The percentage of ECs within a given tissue-specific cluster and their corresponding gene expression values (*Ifnar1* or *Ifngr1*) were visualized via a histogram. For analysis of *Ifnb1* and *Ifng* at resting state and after ischemic stroke in young and aged mice, publicly available data from Garcia-Bonilla et al., 2024 was used [28]. The Seurat R package was used to import paired count data and metadata, followed by quality control [71]. Raw count matrices were filtered to remove cells that expressed fewer than 200 unique genes (low quality), greater than 2,500 unique genes (potentially doublets), or greater than 5% mitochondrial genes (potentially dying). Remaining data was normalized and used to visualize changes in *Ifnb1* and *Ifng* by treatment and age.

Isolation of BECs for scRNA-seq: Mice were anesthetized with isoflurane and perfused with PBS for 5 minutes. Ischemic cortices were dissected, chopped up with a sterile scalpel blade, and placed in Hanks' Balanced Salt Solution (HBSS) without Ca^{2+} and Mg^{2+}. Two or three animals were pooled per sample. Cells were dissociated with the Neural Tissue Dissociation Kit (P) (Miltenyi #130-092-628) and automated gentleMACs (Miltenyi #130-096-427; 37C_NTDK_1 program). Following dissociation, myelin debris was removed using magnetic bead separation (Miltenyi

#130-096-733) in LS columns (Miltenyi #130-042-401), according to manufacturer instructions. Eluent was washed twice and incubated with DRAQ5 (BioLegend #424101, 1:1000) and CD16/CD32 Fc block (BD Biosciences #553141, 1:200) for 15 minutes on ice. Cells were washed and incubated with antibodies against CD31 (PE, BD Biosciences Cat# 553373, 1:200) and CD11b (BV421, BioLegend #101235, 1:200) for 30 minutes on ice. Cells were washed and resuspended in FACS buffer with propidium iodide (1:10,000). Live, nucleated, endothelial cells (DRAQ5^{hi} PI^{lo} CD31+ CD11b-) were sorted on a Sony MA900 (Columbia Stem Cell Initiative Flow Cytometry Core) with 100 um chip. Sorted ECs were resuspended in 50uL PBS with 0.04% BSA and submitted to Columbia Sulzberger Genome Center for scRNA-seq.

scRNA-seq and analysis: Sequencing was performed with 10x Genomics Chromium Single Cell 3' technology and data was processed using the CellRanger v3.0 analysis pipeline to align reads and generate feature-barcode matrices. The Seurat R package was used to read the output of the Cell Ranger pipeline, aggregate data, perform quality control, and conduct downstream analyses [71]. Cells that expressed fewer than 200 unique genes (low quality), greater than 2,500 unique genes (potentially doublets), or greater than 20% mitochondrial genes (potentially dying) were removed. Data was normalized and highly variable genes were identified. Data was then scaled and summarized by principal component analysis (PCA) using previously determined highly variable features (2,000 genes). Dimensionality of the dataset was determined using an Elbow plot and cells were clustered by applying the weighted shared nearest-neighbor graph-based clustering method using the first 20 principal components and resolution value 0.5, followed by visualization using UMAP. Cluster identities were assigned using known cell type markers: ECs (Cldn5, Pecam1), smooth muscle cells (*Acta2*, *Ccl6*), pericytes (*Pdgfrb*, *Atp13a5*), immune cells (*Hexb*,

Tmem119, P2ry12, Csf1r, Trem2, Ly6c2, Ccr2, CD3a, CD4, CD8a), and astrocytes (*S100b, Gfap, Aldh1l1*). Non-ECs were removed and the Harmony package v1.0 was used for batch correction [42]. Batch corrected ECs were then reclustered to refine resolution and EC subtype-specific genes were used to assign EC clusters to a major vessel subtype (Arterial: *Bmx, Efnb1, Hey1, Amd1, Eln, Clu*; Capillary: *Slc1a1, Mfsd2a*; Veinous: *Icam1, Nr2f2, Lcn2, Prcp, Bst1, Il1r1, Lyve1*). This same preprocessing was performed on murine stroke and EAE scRNA-seq dataset. All downstream analyses were performed on ECs only.

For the murine stroke dataset, Seurat *FindMarkers* function was used to compare gene expression across IFNAR1[iECKO] and IFNAR1[fl/fl] Ctrl samples with minimum percent of 0.25, fold-change threshold of 0.25 and p-value cutoff of less than 0.05. Following differential expression (DE) analysis, a score was calculated for each gene using the formula: gene score = $-\log_{10}(pval) \times sign(\log_2 fc)$. This score was used to rank each gene in the DE gene list. GSEA (Broad Institute, v4.3.2) was performed on the ranked list using database-derived (mh.all.v2024.1.Mm.symbols.gmt) and curated gene lists for VEGF signaling, Canonical Wnt signaling, Non-canonical Wnt signaling, Chemokine/cytokine signaling, Tgfb signaling, ECM, Cell-cell adhesion, Transporters, Notch signaling, Angiogenesis, EC proliferation, EC migration, Apoptosis, Antigen processing & presentation, BBB, Inflammation, EndoMT, and Tip cells as previously reported [43]

Congruency Analysis: DEGs (adjusted p-value < 0.001) related to angiogenesis (from curated lists [43]) that were significantly upregulated in the following comparisons were extracted: 1) acute EAE vs CFA spinal cord venous BECs 2) embryonic vs healthy adult BECs and 3) RBPT72 angio capillary clusters (C6, C5, C7, C8) vs other capillary clusters. A Python script was then used to

find the intersection of upregulated angiogenesis-related genes in stroke vs. EAE and stroke vs development. Shared (congruent) genes and unique (non-congruent) genes were counted and used to construct Venn diagrams.

RNA *in situ* hybridization: Digoxigenin-labeled probes were synthesized from plasmids obtained from TransOMIC using the Roche Applied Science in vitro transcription kit (Sigma #11175025910). Mice were anesthetized with isoflurane and perfused with PBS for 5 minutes. The brain was dissected, embedded in Tissue-Plus OCT compound (Fisher 4585), flash frozen on dry ice, and stored at -80C. Coronal sections were cut at 12um on a Leica CM3050 S cryostat and slides were stored at -80C. Alkaline phosphatase (AP) and fluorescent in situ hybridization (FISH) were performed as previously described using anti-sense mRNA probes for full length mouse *IFNβ*, *Irf9*, *Stat1*, *Cxcl10*, and *Apln* [6, 72]. Fluorescent in situ hybridization for *Apln* was followed by immunostaining for Glut1 (Millipore Sigma Cat# 400060, Lot# 2762506). Tissue used for AP in situ hybridization was tiled at 10x magnification using a Zeiss AxioImager. Tissue used for FISH was imaged using a Zeiss LSM700 confocal microscope at 20x magnification.

Immunofluorescence: Mice were anesthetized with isoflurane and perfused with PBS for 4 minutes, followed by 4% paraformaldehyde (PFA) for 6 minutes. The brain was dissected and postfixed in 4% PFA for five hours. After three 30-minute PBS washes, the brain was transferred to 30% sucrose overnight and embedded in OCT. Immunofluorescence for Vegfr2 and Angpt2 was performed in fresh frozen tissue prepared as described for RNA in situ hybridization. Coronal sections were cut at 12um on a Leica CM3050 S cryostat and slides were stored at -80C. Tissue

was washed with PBS for 10 minutes to remove excess OCT and then incubated with blocking buffer (PBS containing 0.1% Triton X-100 (PBST) and 10% BSA) for 1 hour at room temperature. Fresh frozen tissue (Vegfr2, Angpt2) was fixed for 10mins with 4% PFA prior to PBS washes and block. For Cldn5 immunostaining, tissue underwent antigen retrieval (Citrate Buffer, 100C, 35mins) prior to cooling, PBS washes, and block. Primary antibodies were diluted in PBST containing 1% BSA and incubated overnight at 4C.

Antibody	Catalog	Lot	Dilution
Ifitm3	Proteintech #11714-1-AP	#00132970	1:500
Podocalyxin	R&D Systems #AF1556	#JPC0121081	1:300
NeuN	Millipore Sigma #MAB377	#4109965	1:500
Iba1	Wako #019-19741	#5KN4887	1:500
CD68	Abcam #ab53444	#1046837-24	1:2000
Cldn5	Invitrogen #34-1600	#SJ259075	1:200
Ki67	Invitrogen #14-5698-82	#2496198	1:200
Vegfr2	R&D Systems #AF644	#COA0423051	1:300
Angpt2	R&D Systems #MAB098	#FBB0623091	1:200
Glut1	Millipore Sigma #400060	#2762506	1:1000

The following day, slides were washed three times for 10 min with PBST and then incubated for 2 hours at room temperature with fluorescently conjugated secondary antibodies diluted in PBST with 1% BSA (AlexaFluor 488 and Alexa Fluor 594 used at 1:1000; AlexaFluor 647 used at 1:500). Following three washes with PBST and two washes with PBS, slides were coverslipped with Vectashield containing DAPI and then sealed with clear nail polish to store at -20C.

Immunostaining was imaged using a Zeiss LSM700 confocal microscope at 20x magnification.

Biocytin-TMR BBB permeability assay: Mice were injected with 100uL of 1% Biocytin-TMR (Thermo # T12921) dissolved in DPBS with CaCl and MgCl by tail vein. Tracer was allowed to circulate for 30 minutes prior to mice being perfused with PBS for 4 minutes and then PFA for 6 minutes. Tissue was collected and handled as described above for immunofluorescence.

Immunohistochemistry of human tissue: De-identified case and control postmortem brain tissue were provided by the Mount Sinai Neuropathology Brain Bank. Case inclusion criteria were individuals with a neuropathological diagnosis of acute or subacute ischemic stroke with no concurrent clinical diagnosis of active infection or history of autoimmune disease. Controls were sex, age, and brain region matched with no stroke-related neuropathological changes. Microtome serial sections of formalin-fixed and paraffin-embedded tissues were processed for hematoxylin and eosin (H&E) staining as well as immunohistochemistry. Chromogenic Multiplex IHC was performed on Ventana Discovery Ultra according to manufacturer's directions (Neuropathology Brain Bank and Research CoRe, Mt. Sinai). Tissue sections were incubated in three rounds of staining with primary antibodies against CD31 (1:200, Dako #M0823, Clone JC70A), Ifitm3 (1:2000, Proteintech #11714-1-AP), Stat1 (1:250, Cell Signaling Technology #14994), and secondary antibodies Multimer HRP OminiMAP-Anti Rabbit (760-4311, Roche Diagnostics) and OminiMAP-Anti Mouse (760-4310, Roche Diagnostics). The detection was performed using DISCOVERY Yellow HRP Kit, DISCOVERY Green HRP Kit, DISCOVERY Purple Kit (760-250, 760-271, 760-229, Roche Diagnostics). Sodium citrate antigen retrieval (pH 6.0) was used in

between rounds to remove the antibody from the previous round, to avoid any cross-reactivity. Hematoxylin and Bluing reagent (760-2021, 760-2037, Roche Diagnostics) were used as a nuclear counter stain. Slides were digitally scanned (Leica, AperioGT 450) at 40x magnification. H&E staining was used to annotate normal appearing (NA) tissue and ischemic lesions. Corresponding regions on IHC were captured in ImageScope software for downstream analysis.

Cell culture: Primary mBECs (Cell Biologics #C57-6023; Lot M120118W5D17AA.JSS) were grown in mouse endothelial cell medium (Cell Biologics #M1168) containing 10% fetal bovine serum (FBS) and supplements (0.1% VEGF, 0.1% ECGS, 0.1% Heparin, 0.1% EGF, 0.1% Hydrocortisone, 1% L-glutamine, 1.1% Penicillin/Streptomycin) on poly D-Lysine (Sigma #P6407) and collagen IV (Sigma #C5533) coated plates. HBMECs (Cell System #ACBRI 376, Lot 376.03.02.01.2F) were stained with CD31 (FITC, BioLegend #303103, 1:200). The CD31+ HBMEC population was sorted on a Sony MA900 (Columbia Stem Cell Initiative Flow Cytometry Core) and subsequently grown in endothelial cell growth medium (Promocell #C-39220) on collagen IV and fibronectin (Corning #35600) coated plates. Once mBECs and HBMECs were confluent, growth factors were removed and FBS was dropped to 1% for experimental medium. 24hrs after switching to experimental medium, mBECs were treated with 10ng/mL TNFα (R&D Systems #410-MT, Lot CS1420021), 10ng/mL IL1β (R&D Systems #401-ML, Lot BN0720031), 250U/mL IFNβ (PBL Assay #124001-1, Lot 7414), 250U/mL IFNα (PBL Assay #12100-1, Lot 7239), 50ng/mL IFNγ (R&D Systems #485-MI-100, Lot CFP2922051), 1 ug/mL Rho Inhibitor I (Cytoskeleton #CT04-A, Lot 119), or 100 ng/mL VEGF 164 (R&D Systems #493-MV, Lot RQ0321011). HBMECs were treated with 20ng/mL TNFα (R&D Systems #210-TA), 20ng/mL IL1β (R&D Systems #201-LB), or 250U IFNβ (PBL Assay #114201-1, Lot 7333).

TEER measurements: mBECs were plated on 96-well plates containing electrode arrays (Applied BioPhysics, 96W20idf) and TEER was measured in real time using an electric cell-substrate impedance sensing (ECIS) instrument (Applied BioPhysics, ZTheta 96 Well Array Station). Resistance was measured during growth phase and cytokine treatment. Resistance values over time were plotted and the AUC was calculated for each condition.

In vitro immunostaining: For Cldn5 immunostaining, mBECs were fixed with ice cold 95% ethanol for 30 minutes, followed by 1 minute with room temperature acetone. To probe all other proteins of interest, mBECs were fixed with 4% PFA for 10 min. Fixed cells were then washed twice with PBS, blocked for 1 hour in PBST containing 10% BSA, and incubated at 4C overnight with primary antibodies diluted in PBST containing 1% BSA.

Antibody	Catalogue #	Lot #	Dilution
Cldn5	Invitrogen #34-1600	#SJ259075	1:500
ZO1	Invitrogen #33-9100	#ZG392213	1:500
VE-Cadherin	Abcam #ab33168	#GR3452507-1	1:250
Ctnna1	Abcam #ab51032	#1010196-1	1:250

The following day, cells were washed two times for 10 min with PBST and then incubated for 2 hours at room temperature with fluorescently conjugated secondary antibodies diluted in PBST with 1% BSA (AlexaFluor 488 and Alexa Fluor 594 used at 1:1000; AlexaFluor 647 used at 1:500). In some experiments, phalloidin (Abcam #ab176753, 1:1000) was added during secondary antibody incubation. To remove excess secondary antibodies, cells were then washed two times

with PBST and two times with PBS. Prior to imaging on a Zeiss LSM700 confocal microscope, cells were incubated with DAPI (1:5000).

Western blotting: For western blotting, the following primary antibodies were used: rabbit anti-Claudin5 (Invitrogen #34-1600, 1:500), rabbit anti-VE-cadherin (Abcam #ab33168, 1:500), rabbit anti-VEGFR2 (Cell Signaling Technology #2479S, 1:500), and mouse anti-β-actin (Novus #NB600-501, 1:10,000). IR-Dyes 680 (LI-COR #926-68070, 1:20,000) and 800 (LI-COR #926-32211, 1:20,000) were used as secondary antibodies and membranes were imaged using the Odyssey Sa infrared imaging system (LI-COR). Protein levels were quantified using the Empiria Studio software (LI-COR) with β-actin as an internal loading control, and data was presented as normalized to control mean.

Bulk RNA sequencing: RNA was extracted using RNeasy Mini Kit (QIAGEN). RNA sequencing was performed by JP Sulzberger Columbia Genome Center. mRNA was enriched from total RNA samples using poly-A pull down and cDNA libraries were prepared using Illumina Truseq chemistry, followed by sequencing using Illumina NovaSeq 6000. Pseudoalignment of RNAseq reads was performed to a kallisto index created from transcriptome Ensembl v96, Mouse:GRCm38.p6 using kallisto (0.44.0). DESeq2 was used to identify differential gene expression across mBEC treatment conditions and GSEA was performed as described above. For heat map visualization, size factors were calculated using a median ratio method to normalize raw gene counts across samples and z-scores were calculated for each gene's normalized counts.

Scratch Assay: mBECs were treated with experimental medium +/- 250U/mL IFNβ. 24 hours later, scratch wounds were created by the BioTek AutoScratch wound making tool and cells were imaged every 3 hours with the BioTek Cytation 5 Cell Imaging MultiMode Reader (Columbia Confocal and Specialized Microscopy). Wound area fraction was quantified at each time point using the Wound_healing_size_tool Fiji plugin [73].

Bead fibrin angiogenesis assay: Cytodex3 beads were coated with HBMECs and cultured overnight as previously described [74, 75]. The following day (Day 0), HBMEC-coated beads were embedded in fibrin gels consisting of 2.5 mg/mL fibrinogen, 50 U/mL thrombin, and 4 U/mL aprotinin. Fibrin gels with HBMEC-coated beads were treated with standard EC growth media (Promocell) with or without addition of 500 U IFNb, and media was replenished with or without IFNb treatment on Day 2. Gels with beads were fixed on Day 3 for staining and imaging. Data were analyzed using the Sprout Morphology plugin on Fiji [74].

Image analysis

Vascular expression: In Fiji, the vessel marker (Podocalyxin, Cav1, or Glut1) was thresholded and converted into a vascular mask. Median fluorescent intensity (Ifitm3, Vegfr2, Apln, Angpt2) and surface area (Apln, Angpt2) was measured within the vascular mask. Statistical analysis was performed on the mean value calculated across all images for each animal, with a minimum of 15 images per animal per analysis. All analyses were performed blinded to animal genotype and unblinding was performed after completion of statistical analysis.

Human stroke tissue: Using a Python script, image pixels were mapped to 3D RGB color space and an RGB value was assigned as a green 'Ifitm3' reference point. The Euclidian distance in RGB space between each image pixel and the reference point was calculated and visualized as a heatmap. Ifitm3 percent coverage was assessed by applying a Euclidian distance threshold and quantifying the percentage of pixels below that threshold. Statistical analysis was performed on the mean Ifitm3 percent coverage value calculated across 15 images for each region of interest (ROI: control, normal appearing tissue, ischemic lesion). Vascular median Euclidian distance was used as a proxy for Ifitm3 intensity and the median Euclidian distance was calculated for 45 vessel segments sampled across each ROI. Statistical analysis was conducted on the mean value calculated across the 45 vessels for each ROI.

BBB leakage: Tiled bregma sections were loaded into Fiji and uniformly thresholded to measure the area of fluorescence (Biocytin-TMR) that exceeded the threshold, normalized to the total area of each bregma section. This Biocytin-TMR percent leakage area was plotted across specific bregmas (2.34, 1.34, 0.26, -0.22, -0.7, -1.06) and the AUC was calculated to assess the leakage along the rostral-caudal axis of the brain. All analyses were performed blinded to animal treatment/genotype, with unblinding occurring only after the statistical analysis was completed.

Myeloid cell activation: In Fiji, the surface area of thresholded CD68 signal was extracted and normalized to the surface area of Iba1 across anatomically matched ipsilateral and contralateral cortical ROIs at bregma 1.34. Statistical analysis was performed on the mean values calculated from 10 images for both ipsilateral and contralateral ROIs for each animal. All analyses were conducted in a blinded manner regarding animal treatment, with unblinding occurring after the completion of statistical analysis.

Tight junction morphology: To quantify junctional abnormalities, intact junctions, junctional gaps, and absent junctions were identified manually as described before [6]. 8 images/ animal were acquired from both ipsilateral (~114 vessels) and contralateral cortical ROIs (~114 vessels) at bregma -0.22 and the percentage of vessel segments with either intact TJ strands, TJ strands with gaps or absent junctional strands over the total number of vessel segments was calculated per image in ImageJ. We defined junctional status along a Podocalyxin-positive endothelial cell mask as follows: a) A "gap" required a continuous Podocalyxin -positive segment in which ZO-1 intensity fell below a pre-specified threshold while Podocalyxin staining remained intact. Quantification of the TJ strand gaps was restricted to the Podocalyxin membrane mask and puncta off the membrane did not influence the scoring. The vessels coverage of junctional strands per each ROI was calculated by thresholding the junctional marker within the vascular marker after masking in ImageJ.

REFERENCES

1.   Collaborators, G.B.D.S., Global, regional, and national burden of stroke and its risk factors, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. Lancet Neurol, 2021. **20**(10): p. 795-820.
2.   Hankey, G.J., et al., Long-term disability after first-ever stroke and related prognostic factors in the Perth Community Stroke Study, 1989-1990. Stroke, 2002. **33**(4): p. 1034-40.
3.   El Husseini, N., et al., Cognitive Impairment After Ischemic and Hemorrhagic Stroke: A Scientific Statement From the American Heart Association/American Stroke Association. Stroke, 2023. **54**(6): p. e272-e291.
4.   Lo, J.W., et al., Long-Term Cognitive Decline After Stroke: An Individual Participant Data Meta-Analysis. Stroke, 2022. **53**(4): p. 1318-1327.
5.   Jiang, X., et al., Blood-brain barrier dysfunction and recovery after ischemic stroke. Prog Neurobiol, 2018. **163-164**: p. 144-171.
6.   Knowland, D., et al., Stepwise recruitment of transcellular and paracellular pathways underlies blood-brain barrier breakdown in stroke. Neuron, 2014. **82**(3): p. 603-17.
7.   Rust, R., et al., Nogo-A targeted therapy promotes vascular repair and functional recovery following stroke. Proc Natl Acad Sci U S A, 2019. **116**(28): p. 14270-14279.
8.   Williamson, M.R., et al., Reactive astrocytes facilitate vascular repair and remodeling after stroke. Cell Rep, 2021. **35**(4): p. 109048.
9.   Stenman, J.M., et al., Canonical Wnt signaling regulates organ-specific assembly and differentiation of CNS vasculature. Science, 2008. **322**(5905): p. 1247-50.
10.   Liebner, S., et al., Wnt/beta-catenin signaling controls development of the blood-brain barrier. J Cell Biol, 2008. **183**(3): p. 409-17.
11.   Daneman, R., et al., Wnt/beta-catenin signaling is required for CNS, but not non-CNS, angiogenesis. Proc Natl Acad Sci U S A, 2009. **106**(2): p. 641-6.
12.   Chang, J., et al., Gpr124 is essential for blood-brain barrier integrity in central nervous system disease. Nat Med, 2017. **23**(4): p. 450-460.
13.   Song, S., et al., Activation of endothelial Wnt/beta-catenin signaling by protective astrocytes repairs BBB damage in ischemic stroke. Prog Neurobiol, 2021. **199**: p. 101963.
14.   Iadecola, C., M.S. Buckwalter, and J. Anrather, Immune responses to stroke: mechanisms, modulation, and therapeutic potential. J Clin Invest, 2020. **130**(6): p. 2777-2788.
15.   Pedragosa, J., et al., CCR2 deficiency in monocytes impairs angiogenesis and functional recovery after ischemic stroke in mice. J Cereb Blood Flow Metab, 2020. **40**(1_suppl): p. S98-S116.
16.   Beuker, C., et al., Stroke induces disease-specific myeloid cells in the brain parenchyma and pia. Nat Commun, 2022. **13**(1): p. 945.
17.   Mastorakos, P., et al., Temporally distinct myeloid cell responses mediate damage and repair after cerebrovascular injury. Nat Neurosci, 2021. **24**(2): p. 245-258.
18.   Lukhele, S., G.M. Boukhaled, and D.G. Brooks, Type I interferon signaling, regulation and gene stimulation in chronic virus infection. Semin Immunol, 2019. **43**: p. 101277.
19.   Androvic, P., et al., Decoding the Transcriptional Response to Ischemic Stroke in Young and Aged Mouse Brain. Cell Rep, 2020. **31**(11): p. 107777.
20.   Barrett, J.P., et al., Interferon-beta Plays a Detrimental Role in Experimental Traumatic Brain Injury by Enhancing Neuroinflammation That Drives Chronic Neurodegeneration. J Neurosci, 2020. **40**(11): p. 2357-2370.

21. Prinz, M., et al., Distinct and nonredundant in vivo functions of IFNAR on myeloid cells limit autoimmunity in the central nervous system. Immunity, 2008. **28**(5): p. 675-86.

22. Roy, E.R., et al., Type I interferon response drives neuroinflammation and synapse loss in Alzheimer disease. J Clin Invest, 2020. **130**(4): p. 1912-1930.

23. Baruch, K., et al., Aging. Aging-induced type I interferon response at the choroid plexus negatively affects brain function. Science, 2014. **346**(6205): p. 89-93.

24. Roy, E.R., et al., Fate-mapping and functional dissection reveal perilous influence of type I interferon signaling in mouse brain aging. Mol Neurodegener, 2024. **19**(1): p. 48.

25. Kuo, P.C., et al., Interferon-beta Modulates Inflammatory Response in Cerebral Ischemia. J Am Heart Assoc, 2016. **5**(1).

26. Veldhuis, W.B., et al., Interferon-beta blocks infiltration of inflammatory cells and reduces infarct volume after ischemic stroke in the rat. J Cereb Blood Flow Metab, 2003. **23**(9): p. 1029-39.

27. Kuo, P.C., et al., Interferon-beta alleviates delayed tPA-induced adverse effects via modulation of MMP3/9 production in ischemic stroke. Blood Adv, 2020. **4**(18): p. 4366-4381.

28. Garcia-Bonilla, L., et al., Analysis of brain and blood single-cell transcriptomics in acute and subacute phases after experimental stroke. Nat Immunol, 2024. **25**(2): p. 357-370.

29. Yang, H., et al., STING activation reprograms tumor vasculatures and synergizes with VEGFR2 blockade. J Clin Invest, 2019. **129**(10): p. 4350-4364.

30. Daniels, B.P., et al., Viral pathogen-associated molecular patterns regulate blood-brain barrier integrity via competing innate cytokine signals. mBio, 2014. **5**(5): p. e01476-14.

31. Daniels, B.P., et al., Regional astrocyte IFN signaling restricts pathogenesis during neurotropic viral infection. J Clin Invest, 2017. **127**(3): p. 843-856.

32. Thacker, S.G., et al., The detrimental effects of IFN-alpha on vasculogenesis in lupus are mediated by repression of IL-1 pathways: potential role in atherogenesis and renal vascular rarefaction. J Immunol, 2010. **185**(7): p. 4457-69.

33. Mastorakos, P., et al., Antimicrobial immunity impedes CNS vascular repair following brain injury. Nat Immunol, 2021. **22**(10): p. 1280-1293.

34. Goldmann, T., T. Blank, and M. Prinz, Fine-tuning of type I IFN-signaling in microglia--implications for homeostasis, CNS autoimmunity and interferonopathies. Curr Opin Neurobiol, 2016. **36**: p. 38-42.

35. Kalucka, J., et al., Single-Cell Transcriptome Atlas of Murine Endothelial Cells. Cell, 2020. **180**(4): p. 764-779 e20.

36. Labat-gest, V. and S. Tomasi, Photothrombotic ischemia: a minimally invasive and reproducible photochemical cortical lesion model for mouse stroke studies. J Vis Exp, 2013(76).

37. Doll, D.N., T.L. Barr, and J.W. Simpkins, Cytokines: their role in stroke and potential use as biomarkers and therapeutic targets. Aging Dis, 2014. **5**(5): p. 294-306.

38. Forero, A., et al., Differential Activation of the Transcription Factor IRF1 Underlies the Distinct Immune Responses Elicited by Type I and Type III Interferons. Immunity, 2019. **51**(3): p. 451-464 e6.

39. Tapon, N. and A. Hall, Rho, Rac and Cdc42 GTPases regulate the organization of the actin cytoskeleton. Curr Opin Cell Biol, 1997. **9**(1): p. 86-92.

40. Aguilar-Cazares, D., et al., Contribution of Angiogenesis to Inflammation and Cancer. Front Oncol, 2019. **9**: p. 1399.

41. Bekisz, J., et al., Antiproliferative Properties of Type I and Type II Interferon. Pharmaceuticals (Basel), 2010. **3**(4): p. 994-1015.

42. Korsunsky, I., et al., Fast, sensitive and accurate integration of single-cell data with Harmony. Nat Methods, 2019. **16**(12): p. 1289-1296.

43. Shahriar, S., et al., VEGF-A-mediated venous endothelial cell proliferation results in neoangiogenesis during neuroinflammation. Nat Neurosci, 2024. **27**(10): p. 1904-1917.

44. Hupe, M., et al., Gene expression profiles of brain endothelial cells during embryonic development at bulk and single-cell levels. Sci Signal, 2017. **10**(487).

45. Escoubas, C.C., et al., Type-I-interferon-responsive microglia shape cortical development and behavior. Cell, 2024. **187**(8): p. 1936-1954 e24.

46. Jin, M., et al., Type-I-interferon signaling drives microglial dysfunction and senescence in human iPSC models of Down syndrome and Alzheimer's disease. Cell Stem Cell, 2022. **29**(7): p. 1135-1153 e8.

47. Gonzalez-Navajas, J.M., et al., Immunomodulatory functions of type I interferons. Nat Rev Immunol, 2012. **12**(2): p. 125-35.

48. Lee, P.Y., et al., Type I interferon as a novel risk factor for endothelial progenitor cell depletion and endothelial dysfunction in systemic lupus erythematosus. Arthritis Rheum, 2007. **56**(11): p. 3759-69.

49. Baccala, R., et al., Type I interferon is a therapeutic target for virus-induced lethal vascular damage. Proc Natl Acad Sci U S A, 2014. **111**(24): p. 8925-30.

50. Fuertes, M.B., et al., Host type I IFN signals are required for antitumor CD8+ T cell responses through CD8alpha+ dendritic cells. J Exp Med, 2011. **208**(10): p. 2005-16.

51. Demaria, O., et al., STING activation of tumor endothelial cells initiates spontaneous and therapeutic antitumor immunity. Proc Natl Acad Sci U S A, 2015. **112**(50): p. 15408-13.

52. Viengkhou, B., et al., The brain microvasculature is a primary mediator of interferon-alpha neurotoxicity in human cerebral interferonopathies. Immunity, 2024. **57**(7): p. 1696-1709 e10.

53. van Boxel-Dezaire, A.H., M.R. Rani, and G.R. Stark, Complex modulation of cell type-specific signaling in response to type I interferons. Immunity, 2006. **25**(3): p. 361-72.

54. Schaupp, L., et al., Microbiota-Induced Type I Interferons Instruct a Poised Basal State of Dendritic Cells. Cell, 2020. **181**(5): p. 1080-1096 e19.

55. Schoggins, J.W. and C.M. Rice, Interferon-stimulated genes and their antiviral effector functions. Curr Opin Virol, 2011. **1**(6): p. 519-25.

56. Hayashi, T., K. Abe, and Y. Itoyama, Reduction of ischemic damage by application of vascular endothelial growth factor in rat brain after transient ischemia. J Cereb Blood Flow Metab, 1998. **18**(8): p. 887-95.

57. Gu, W., et al., Vascular endothelial growth factor-A and -C protein up-regulation and early angiogenesis in a rat photothrombotic ring stroke model with spontaneous reperfusion. Acta Neuropathol, 2001. **102**(3): p. 216-26.

58. Dzietko, M., et al., Delayed VEGF treatment enhances angiogenesis and recovery after neonatal focal rodent stroke. Transl Stroke Res, 2013. **4**(2): p. 189-200.

59. van Bruggen, N., et al., VEGF antagonism reduces edema formation and tissue damage after ischemia/reperfusion injury in the mouse brain. J Clin Invest, 1999. **104**(11): p. 1613-20.

60. Zhang, Z.G., et al., VEGF enhances angiogenesis and promotes blood-brain barrier leakage in the ischemic brain. J Clin Invest, 2000. **106**(7): p. 829-38.

61. Stupack, D.G. and D.A. Cheresh, ECM remodeling regulates angiogenesis: endothelial integrins look for new ligands. Sci STKE, 2002. **2002**(119): p. pe7.

62. Ashby, K.M., et al., Sterile production of interferons in the thymus affects T cell repertoire selection. Sci Immunol, 2024. **9**(97): p. eadp1139.

63. del Toro, R., et al., Identification and functional analysis of endothelial tip cell-enriched genes. Blood, 2010. **116**(19): p. 4025-33.

64. Gerhardt, H., et al., VEGF guides angiogenic sprouting utilizing endothelial tip cell filopodia. J Cell Biol, 2003. **161**(6): p. 1163-77.

65. Abid, M.R., et al., Vascular endothelial growth factor activates PI3K/Akt/forkhead signaling in endothelial cells. Arterioscler Thromb Vasc Biol, 2004. **24**(2): p. 294-300.

66. Soga, N., et al., Rho family GTPases regulate VEGF-stimulated endothelial cell motility. Exp Cell Res, 2001. **269**(1): p. 73-87.

67. Prigge, J.R., et al., Type I IFNs Act upon Hematopoietic Progenitors To Protect and Maintain Hematopoiesis during Pneumocystis Lung Infection in Mice. J Immunol, 2015. **195**(11): p. 5347-57.

68. Madisen, L., et al., A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. Nat Neurosci, 2010. **13**(1): p. 133-40.

69. Wang, Y., et al., Ephrin-B2 controls VEGF-induced angiogenesis and lymphangiogenesis. Nature, 2010. **465**(7297): p. 483-6.

70. Rousselet, E., J. Kriz, and N.G. Seidah, Mouse model of intraluminal MCAO: cerebral infarct evaluation by cresyl violet staining. J Vis Exp, 2012(69).

71. Hao, Y., et al., Integrated analysis of multimodal single-cell data. Cell, 2021. **184**(13): p. 3573-3587 e29.

72. Lengfeld, J.E., et al., Endothelial Wnt/beta-catenin signaling reduces immune cell infiltration in multiple sclerosis. Proc Natl Acad Sci U S A, 2017. **114**(7): p. E1168-E1177.

73. Suarez-Arnedo, A., et al., An image J plugin for the high throughput image analysis of in vitro scratch wound healing assays. PLoS One, 2020. **15**(7): p. e0232565.

74. Eglinger, J., H. Karsjens, and E. Lammert, Quantitative assessment of angiogenesis and pericyte coverage in human cell-derived vascular sprouts. Inflamm Regen, 2017. **37**: p. 2.

75. Nakatsu, M.N., J. Davis, and C.C. Hughes, Optimized fibrin gel bead assay for the study of angiogenesis. J Vis Exp, 2007(3): p. 186.

FIGURES AND FIGURE LEGENDS

Figure 1. Ischemic brain upregulates endogenous IFN1 and shows expansion of IFN responsive BECs. a, Distribution of *Ifnar1* transcript across endothelial cells (ECs) in four distinct tissues (brain, liver, lung, spleen). X-axis shows normalized *Ifnar1* mRNA expression values and Y axis shows the percentage of ECs in organ-specific clusters that express *Ifnar1* mRNA at the respective expression level. **b,** Brightfield images of Dig-labelled RNA *in situ* hybridizations showing expression of *IFNβ* mRNA in brain coronal sections at 24 hours, 72 hours, and 7 days after RBPT and t-MCAO (n= 3 mice per timepoint). The dotted red curve demarcates the stroke lesion and the black rectangle shows the area of the zoomed image. Scale bars = 100 μm and 25 μm (insets). **c,** UMAP clustering of mBECs from sham ("resting state", n = 2 mice) and dissected ischemic ("RBPT72", n = 3 mice) cerebral cortex. Dotted line highlights highly IFN responsive cluster. **d,** Clusters colored by condition ("resting state" grey vs "RBPT72" red). **e,** Cluster 3 abundance in resting state and RBPT72 conditions. Dots represent independent samples and data shown as percentage of total cells in sample (two-tailed unpaired t-test). **f,** Dot plot of ISG expression showing upregulation in cluster 3. **g,** Representative immunostaining and median fluorescent intensity (MFI) quantification of vascular Ifitm3 (green) in ischemic ipsilateral and contralateral cortex 72 hours after t-MCAO (n = 6 mice; two-tailed paired t-test). **h,** Representative immunostaining of vascular Ifitm3 in ischemic territory and contralateral cortex 72 hours after RBPT. Podocalyxin was used as vascular marker. All bars indicate mean ± SEM, *p<0.05.

Figure 2. BEC IFN signature is conserved in human ischemic stroke. a, Immunostaining of IFN-related proteins (IFITM3, STAT1) and vascular marker (CD31) in the ischemic lesion and normal appearing tissue (NAT) in postmortem ischemic stroke brain tissue (N = 6 cases) and age- and brain region-matched control tissue (Ctrl, N = 6 cases). **b,** Example of image pixels projected onto 3D RGB color space. **c,** Quantification of IFITM3 coverage following thresholding, normalized to total number of image pixels. **d,** Quantification of vascular median Euclidian distance as a proxy for vascular IFITM3 intensity. Bars represent mean ± SEM, *p<0.05, ****p<0.0001, two-tailed paired t-test (NAT versus Ischemic Lesion), two tailed unpaired t-test (comparisons with controls). Closed data points with the same color indicate the same case, while open data points with same color indicate corresponding matched control.

Figure 3. Activation of IFN1 signaling in primary BECs mitigates inflammation-mediated barrier disruption. a, Transendothelial electrical resistance (TEER) measurements and area under the curve (AUC) quantifications for mouse brain endothelial cells (mBECs) treated with two inflammatory cytokines (Cyt: IL-1β (10 ng/mL) and TNFα (10ng/mL)) +/- IFNβ (250 U/mL). Curves represent the mean TEER values from 8 independent wells per treatment group (technical replicates). Grey shaded region marks the time window used to calculate AUC. The bar graphs

indicate mean ± SEM, ****p<0.0001, two-tailed unpaired t-test from n=3 biological replicates. **b,** Representative Western blot and quantification of Claudin-5 (normalized to β-actin, relative to control) 24 hours after treatment (Control n = 6 replicates, Cyt n= 6 replicates, Cyt + IFNβ = 5 replicates). **c,** Representative immunofluorescence images of mBECs stained for Claudin-5 (red), ZO-1 (green), and DAPI (blue) 24 hours after treatment. White arrows indicate colocalization of Claudin-5 and ZO-1 tight junctions whereas yellow arrows indicate lack of colocalization. White dashed circles indicate intracellular Claudin-5 aggregates. Scale bar = 25um. **d,** TEER measurements and AUC quantifications of HBMECs and HUVECs treated with TNFα/IL1β (Cyt; each used at 20ng/mL) ± IFNβ (250U/mL). Curves represent the mean TEER values from 8 independent wells per treatment group. Grey shaded region marks the time window used to calculate AUC. The bars in bar graphs indicate mean ± SEM, ****p<0.0001, two-tailed unpaired t-test from n=3 biological replicates. **e, f,** Heatmap visualization of z-scores for differentially expressed genes (DEGs) associated with **e,** Inflammation and **f,** Wnt Signaling that are significantly upregulated in mBECs treated with Cyt + IFNβ for 24 hours compared to treatment with Cyt alone, as identified by GSEA (n=3 for each treatment group, with each column representing an independent bulk-RNAseq sample). **g,** TEER measurements and AUC quantifications of mBECs treated with TNFα/IL1β ± IFNβ ± Rho Inhibitor (RhoI, 1ug/mL) vs Rac1 Inhibitor (Rac1I, 5uM). Curves represent the mean TEER values from 8 independent wells per treatment group Grey shaded region marks the time window used to calculate AUC. The bars in bar graphs indicate mean ± SEM, ****p<0.0001, two-tailed unpaired t-test from n=3 biological replicates. **h,** Putative mechanism by which IFN1 may stabilize BEC junctional integrity through local Rho-dependent actin remodeling.

Figure 4. IFNβ treatment reduces acute BBB disruption after ischemic brain injury. a, Experimental timeline for the t-MCAO and IFNβ treatment. Mice were treated with vehicle or IFNβ (10,000 U) 3 hours post reperfusion and analyses were performed at 72 hours. **b,** Neurological scores of vehicle- (n=9) and IFNβ-treated (n=10) mice. **c,** Biocytin-TMR tracer extravasation in seven brain sections located at different distances from the bregma in vehicle and IFNβ-treated mice 72 hours after t-MCAO (biocytin-TMR was injected 30-45 minutes before analysis). Solid line demarcates the border of the brain section and dotted line demarcates leakage area. **d,** Quantification of biocytin-TMR leakage area analyzed across seven distinct brain sections at 72 hours after t-MCAO (Vehicle n=10 mice, IFNβ-treated n =9 mice). **e,** Quantification of area under curve (AUC) as measure of cumulative biocytin-TMR leakage along rostral-caudal brain axis. **f,** Immunofluorescence staining of anatomically matched ROIs in contralateral and ipsilateral ischemic cortices of vehicle- (n=10) and IFNβ- treated (n=9) mice at 72 hours after t-MCAO for Claudin-5 (Cldn-5; green) and Podocalyxin (purple). Blue arrowheads indicate intact junctions. Yellow arrowheads indicate junctions with gaps. Red arrowheads indicate absent junctions. **g,** Pie charts showing the average proportion of vessels with intact Cldn-5+ strands, gaps in Cldn-5+ strands, and absent Cldn-5+ strands in the contralateral and ipsilateral ischemic cortices of vehicle- and IFNβ- treated mice. Percentage of vessels with **h,** intact Cldn-5+ strands, **h',** gaps in Cldn-5+ strands, and **h'',** absent Cldn-5+ strands in contralateral and reperfusion cortices of vehicle- and IFNβ- treated mice. Bars indicate mean ± SEM, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, two-tailed unpaired t-test between treatments, two-tailed paired t-test between cortices within treatment.

Figure 5: *In vivo* **ablation of BEC IFN1 signaling exacerbates acute barrier disruption after ischemic brain injury. a,** Generation of inducible endothelial cell Ifnar1knockout (*Ifnar1[iECKO]*) and the experimental t-MCAO timeline. **b,** Neurological scores of Ifnar1[iECKO] (n = 6) and Ifnar1[fl/fl] Ctrl (n = 7) mice. **c,** Biocytin-TMR tracer extravasation in seven brain sections located at different distances from the bregma landmark in *Ifnar1[iECKO]* (n = 6) and *Ifnar1[fl/fl]* (n = 7) mice 72 hours after t-MCAO (biocytin-TMR was injected 30-45 minutes before analysis). Solid line demarcates the border of the brain section and dotted line demarcates leakage area. **d,** Quantification of biocytin-TMR leakage area analyzed across seven distinct brain regions at 72 hours after t-MCAO. **e,**

Quantification of area under curve (AUC) as a measure of cumulative biocytin-TMR leakage along rostral-caudal brain axis. Bars indicate mean ± SEM, *p<0.05, two-tailed unpaired t-test.

a Myc Targets

Ctrl   Cyt   Cyt + IFNβ

b VEGF Signaling

Ctrl   Cyt   Cyt + IFNβ

c Tip Cells

Ctrl   Cyt   Cyt + IFNβ

g Phalloidin DAPI

Control

IFNβ

d

Ctrl   Cyt   Cyt + IFNβ

Vegfr2, 210, 230kDa

Bactin, 42kDa

VEGFR2 (fold change of control)

Ctrl   Cyt   Cyt + IFNβ

**

**

h

Average sprout length (μm)

control   +IFNβ

**

e

Ctrl

IFNβ priming

0hr 3hr 6hr 9hr 12hr 15hr 18hr 21hr 24hr

Wound area fraction

Time post scratch

00:00 6:00 12:00 18:00 24:00

Control
IFNβ priming

*   **   ***   ****

i

Total network length (μm)

control   +IFNβ

****

f

± IFNβ ± VEGFa

Resistance (ohms*cm2)

Time (hrs)

Ctrl --> Ctrl
IFNβ --> Ctrl
IFNβ --> VEGFa
Ctrl --> VEGFa
Ctrl --> IFNβ + VEGFa

Resistance (ohms*cm²)

Ctrl/Ctrl   IFNβ/Ctrl   IFNβ/VEGFa   Ctrl/VEGFa   Ctrl/IFNβ+VEGFa

**   ****   ****   ns

Figure 6. IFN1 signaling suppresses angiogenic features of BECs *in vitro*

a-c, Heatmap visualization of z-scores for differentially expressed genes (DEGs) associated with **a,** Myc Targets **b,** VEGF Signaling and **c,** Tip Cells that are significantly downregulated in mBECs treated with Cyt + IFNβ for 24hrs compared to treatment with Cyt alone, as identified by GSEA (n=3 for each treatment group, with each column representing an independent bulk-RNAseq sample) **d,** Representative western blot and quantification of VEGFR2 (normalized to β-actin, relative to control) 24 hours after indicated treatment (Ctrl n = 5 replicates, Cyt n= 6 replicates, Cyt + IFNβ = 5 replicates) **e,** Representative migration images and quantification of mBECs in wound healing assay ± 24 hour IFNβ (250U/mL) pre-treatment. Curves represent the mean of two technical replicates averaged across three independent experiments; wound area fraction quantified using Wound_healing_size_tool Fiji plugin [73]. **f,** TEER measurement in mBECs following VEGF-A (100ng/mL) ± 24 hour IFNβ (250U/mL) pre-treatment. Grey shaded region indicates time-period used to quantify acute VEGF-A-mediated permeability. Curves represent the mean TEER values from 8 independent wells per treatment group. The bars in bar graphs indicate mean ± SEM, ****$p<0.0001$, two-tailed unpaired t-test from n=3 biological replicates. **g,** Representative bead fibrin assay images for HBMECs ± 72 hours IFNβ (500U/mL) stained for Phalloidin (green) and DAPI (blue) (n=15 beads / condition with data representative of 3 independent experiments) and **h,** quantification for sprout length and **i,** network length. Bars indicated mean ± SEM, **$p<0.01$, ****$p<0.0001$, two-tailed unpaired t-test.

Figure 7. Ablation of endogenous BEC IFN1 signaling leads to expansion of angiogenic BECs following ischemic brain injury. a, Schematic of the workflow for scRNAseq experiments. **b,** UMAP visualization of 10 distinct BEC clusters isolated from the dissected ischemic cortex of *Ifnar1^{fl/fl}* and *Ifnar1^{iECKO}* mice (2-3 mice pooled/sample; 4 samples/genotype) 72 hours after rose bengal photothrombosis (RBPT). **c,** Clusters colored by genotype [*Ifnar1^{fl/fl}* (grey) versus *Ifnar1^{iECKO}* (red)]. **d,** Quantification of cluster composition by genotype. Y axis represents percent

of BECs in each cluster from *Ifnar1^{fl/fl}* and *Ifnar^{iECKO}* ischemic cortex normalized for total number of cells per sample. **e,** Processes significantly upregulated and downregulated in arterial (red), capillary (grey), and venule (blue) BECs isolated from ischemic cortex of *Ifnar^{iECKO}* compared to *Ifnar1^{fl/fl}* mice, as identified by GSEA. Bars indicate significance of enrichment (-log$_2$FDR) and black dots indicate the number of genes in each gene set. **f,** Gene-expression heatmap showing the log fold-change of the average expression (avglog2FC) of BBB and angiogenesis related genes of each capillary cluster (C0-C8, Extended Data Figure 6). Red text highlights clusters enriched (>75%) in BECs derived from *Ifnar^{iECKO}* mice.

Figure 8: *In vivo* **validation of angiogenesis-related transcripts enriched in *Ifnar1iECKO* mice after ischemic brain injury. a,** Schematic illustrating regions of image acquisition in the RBPT

model: contralateral cortex (1), peri-infarct cortex (2), and ischemic lesion (3). Violin plots for **b,** *Kdr*, **c,** *Apln*, and **d,** *Angpt2* mRNA expression in Ifnar1$^{fl/fl}$ Ctrl and Ifnar1^{iECKO} BECs generated from scRNA-seq dataset. Brain sections from Ifnar1$^{fl/fl}$ Ctrl and Ifnar1^{iECKO} mice 72 hours after RBPT were immunostained for **e,** Vegfr2 protein (contralateral and peri-infarct cortex) and **f,** Angpt2 protein (contralateral cortex and cortical ischemic lesion) or **g,** probed with FISH for *Apln* mRNA (contralateral and peri-infarct cortex), and combined with immunostaining for Caveolin-1 protein or Glut1 (vascular markers) to quantify vascular expression levels. Bars indicate mean ± SEM, *p<0.05, **p<0.01, two-tailed unpaired t-test. **h,** Model for role of IFN1 signaling in the vasculature after ischemic stroke. Ischemia-induced cell death triggers local upregulation of IFNβ, which is temporally aligned with the period of vascular remodeling that occurs following ischemic stroke. Upregulation of endothelial IFN1 signaling suppresses angiogenic processes and acutely strengthens BEC junctions, potentially through reorganization of the actin cytoskeleton and degradation of VEGFR2, thereby promoting BEC quiescence and acquisition of barrier function. In the absence of BEC IFN1 signaling, there is a potential expansion of angiogenic BECs that lack barrier properties, exacerbating vascular permeability and preventing restoration of CNS vascular homeostatic function after ischemic stroke.