Title

Sex-biased expression of enteroendocrine cell-derived hormones contributes to higher

fat storage in *Drosophila* females

Authors

Puja Biswas[1,2], Elizabeth J. Rideout[1,*]

Affiliations

[1] Department of Cellular and Physiological Sciences, Life Sciences Institute, The

University of British Columbia, Vancouver, Canada

[2] Present Address: Department of Pediatrics, BC Children's Hospital Research Institute,

The University of British Columbia, Vancouver, Canada

* Corresponding author: elizabeth.rideout@ubc.ca

Abstract

Enteroendocrine (EE) cells in the *Drosophila* gut produce and release multiple factors,

including Allatostatin A (AstA), Allatostatin C (AstC), neuropeptide F (NPF), tachykinin

(Tk), Diuretic hormone 31 (Dh31), Bursicon, CCHamide 1 (CCHa1) and CCHamide 2

(CCHa2), and short neuropeptide F (sNPF). Collectively, these peptides ensure that

physiology (e.g., fat storage, fluid balance) and behavior (e.g., feeding, sleep) are

coordinated with environmental factors such as nutrient quantity and quality. Despite

notable sex differences in physiology and behavior, it remains unclear whether the

regulation and function of these EE cell-derived factors is shared between males and

females. Given that recent data identified sex-biased physiological effects of two EE

cell-derived hormones on *Drosophila* food intake and energy mobilization, we performed

a detailed characterization of these hormones in male and female flies. Despite an

overall male bias in mRNA levels of *AstA*, *AstC*, *Tk*, *NPF*, *Dh31* in whole-body and head

samples, we observed a strong female bias in mRNA levels of *AstC*, *Tk*, and *NPF* in the

gut. To determine whether this sex-biased regulation was physiologically significant, we

monitored triglyceride levels in flies with gut-specific loss of EE cell-derived hormones.

In 5-day-old flies, loss of either EE cell-derived AstC or Tk reduced fat storage in

females with no effect in males. These female-specific effects on fat storage were

reproduced in flies with neuron-specific loss of the AstC (AstC-R2) and Tk receptors

(TkR99D). Together, these data uncover strongly sex-biased regulation of EE cell-

derived hormones, and show that gut-specific loss of two of these hormones had a

female-specific effect on body fat.

Keywords

Sex difference, gut, enteroendocrine cell, physiology, triglyceride, hormones, fat storage

Highlights:

• Enteroendocrine cell-expressed hormones show strongly sex-biased expression

• Loss of enteroendocrine cell-derived AstC and Tk reduced body fat only in females

• Neuronal loss of AstC or Tk receptors reduced stored fat only in females

1. Introduction

In *Drosophila*, females store more fat than males [1–7]. Greater female fat storage has been observed in both mated and unmated females compared with age-matched males [1,3–7]. In flies, as in other animals, triglyceride is the main form of stored fat. Although triglyceride is present in many cell types and organs (*e.g.*, oenocytes, glia, neurons, gut) [5,6,8–14], the majority of triglyceride is stored in an organ called the fat body [3,12,15].

In females, high levels of triglyceride in the fat body play a key role in supporting reproduction and physiology [5,16]. Indeed, triglyceride from the fat body is a key energy source for the developing embryo [17]. Increased fat storage in adult females also supports prolonged survival during nutrient deprivation compared with males [4,6]. Despite these clear benefits of greater fat storage for female fertility, excess fat accumulation in males adversely affects their reproductive output. Specifically, males carrying a mutation that promotes excess triglyceride accumulation show reduced testis size, defects consistent with delays in spermatogenesis, and ultimately a reduction in sperm number [8]. The sex difference in fat storage therefore likely reflects the fact that males and females differ in how much whole-body triglyceride storage supports optimal fertility.

Recent studies have identified genes and pathways that contribute to sex differences in fat storage [5–7,18]. In mated females, the steroid hormone ecdysone acts on neurons to promote food intake, which is associated with increased body fat [5]. In unmated adult females, the insulin/insulin-like growth factor signaling pathway (IIS) plays a key role in maintaining an elevated level of triglyceride storage compared with adult males [18]. Specifically, adult females have higher production of *Drosophila*

insulin-like peptide 3 (Dilp3) and greater insulin sensitivity, leading to higher IIS activity.

This elevated IIS activity is important for females to store more triglyceride than males,

as adult-specific ablation of the insulin-producing cells reduces body fat in females but

not males [18].

In males, body fat levels are maintained by higher expression and activity of two

catabolic pathways that promote fat breakdown. One pathway is regulated by

triglyceride lipase *brummer* (*bmm*), where males show higher *bmm* mRNA levels

compared with females [6]. This elevated *bmm* expression contributes to the sex

difference in fat storage by restricting triglyceride accumulation in males. Similarly,

males have higher production and secretion of Adipokinetic hormone (Akh), a key

lipolytic hormone in insects [7]. As with *bmm*, high levels of Akh in males contribute to

the sex difference in fat storage by limiting fat accumulation in males [7]. Together,

these studies have defined a model of the sex difference in fat storage in which females

maintain higher levels of fat storage in part due to a higher relative activity level for

anabolic pathway IIS, whereas males have lower fat storage due to higher relative

activity of catabolic effectors such as *bmm* and Akh. While some progress has been

made in revealing the mechanisms underlying the sex-specific regulation of Akh and IIS

[7,18], we do not have a complete understanding of the factors that determine the sex-

biased regulation and function of these key metabolic factors.

Recent clues into the regulation of Akh, *bmm*, and IIS have emerged from

studies on peptide hormone function [19–30], as several hormones influence physiology

via effects on Akh- and Dilp-producing cells [19–22,24–33]. In particular, recent studies

have illuminated an important role for hormones produced by the enteroendocrine (EE)

cells of the gut [32,34–36]. Adult *Drosophila* EE cells produce and release hormones

such as Allatostatin A (AstA), Allatostatin C (AstC), neuropeptide F (NPF), tachykinin

(Tk), Diuretic hormone 31 (Dh31), Bursicon, CCHamide 1 (CCHa1) and CCHamide 2

(CCHa2), and short neuropeptide F (sNPF) [37–41].

EE cells are identified by expression of the homeodomain protein Prospero in

adults [42,43], where EE cells that produce distinct hormones are present in

anatomically defined regions of the adult gut [43]. For example, AstA- and Dh31-

producing EE cells are located in the posterior midgut, whereas EE cells that produce

AstC and Tk are found along the entire length of the midgut [37]. NPF-producing cells

are found in the anterior and middle midgut [41]. Supporting a role for EE cell-derived

hormones in regulating Akh/IIS, in fed conditions studies show EE cell-derived

hormones such as Bursicon inhibit Akh secretion [28], whereas NPF enhances Dilp

secretion from the insulin-producing cells [29]. During starvation, AstC promotes Akh

release from Akh-producing cells to enable lipid mobilization [22]. While some EE cell-

derived hormones have been shown to have sex-biased effects on food intake and

energy mobilization [22,44,45], sex differences in the regulation and function of most of

these hormones remain unclear. Defining potential differences in EE cells is an

important task, as prior studies have revealed profound differences in gut biology

between males and females.

For example, males and females differ in overall gut size and shape [46,47] and

in the number of intestinal stem cell divisions [46,48–50]. The absorptive lining of the

gut also shows sex differences during aging [49,51]. After mating, females show

pronounced changes to gut size, function, and gene expression [46,52–55]. While sex

differences in intestinal stem cells and enterocytes play a key role in mediating these

differences in gut biology, we know less about male-female differences in EE cells. We

therefore aimed to perform a detailed characterization of EE cell-derived hormones in

males and females, and to determine the contribution of these hormones to physiology

in each sex. We reveal profound sex-biased regulation of EE cell-expressed hormones

AstC, Tk, and NPF: females show higher mRNA levels of these hormones in the gut

than males. For at least two EE cell-derived hormones this sex-biased regulation was

physiologically significant, as loss of AstC and Tk in the gut reduced fat storage in

females with no effect in males. These female-specific fat storage defects were

reproduced by loss of AstC and Tk receptors in neurons and/or neuropeptide-producing

cells. While the specific cell type targeted by these EE cell-derived hormones to

influence fat storage remains unclear, this reveals a female-specific contribution of EE

cell-derived hormones in regulating body fat.

2. RESULTS

2.1. Sex differences in expression of gut-derived peptide hormones

Given that several EE cell-derived hormones such as AstA (FBgn0015591), AstC

(FBgn0032336), Tk (FBgn0037976), NPF (FBgn0027109), Dh31 (FBgn0032048) are

expressed in cells outside the gut [20,56–70], we used quantitative real-time PCR

(qPCR) to assess whole-body mRNA levels of genes encoding EE cell-expressed

hormones. In particular, we focused on hormones known to influence whole-body fat

metabolism [20,22,29,44,71,72], though an important future direction of this work will be

to assess sex differences in all EE cell-expressed hormones. In 5-day-old w^{1118}

unmated adult males and females, we found that whole-body mRNA levels of *AstA*,

AstC, *Tk*, *NPF*, and *Dh31* were significantly higher in males than in females (Figure 1A-

E). To gain further insight into this sex-biased expression, we analyzed mRNA levels of

these factors from isolated heads and intestines, as these are the main sites of AstA,

AstC, Tk, NPF, and Dh31 production [35,37]. A significant male bias in mRNA levels

was found in the head for *AstA*, *AstC*, *Tk*, *NPF,* and *Dh31* (Figure 1F-1J). In contrast,

mRNA levels of *AstC*, *Tk*, and *NPF* in isolated intestines showed a strong female bias

(Figure 1L-1N). No sex bias in the expression of *AstA* or *Dh31* was observed in the gut

(Figure 1K, 1O).

Building on the sex bias in mRNA levels, we next examined mRNA levels of

receptors that correspond to EE cell-expressed hormones with sex-biased expression in

whole-body, fat body, and head samples. Whole-body mRNA levels of the receptors for

AstA (*AstA-R2*), AstC (*AstC-R2*), Tk (*TkR99D*), NPF (*NPFR*), and Dh31 (*Dh31-R*) were

significantly higher in 5-day-old w^{1118} males compared with age-matched females

(Figure S1A-E). For most peptides, the male bias was due to a higher mRNA level in

the head and not the fat body (Figure S1A-E); however, *TkR99D* mRNA levels were

higher in male fat bodies with no difference in head mRNA levels (Figure S1C). We

therefore cannot rule out a contribution of additional anatomical sites to the male bias in

expression of EE cell-expressed hormones, which is an interesting area for future

investigation. Taken together with our data on peptide mRNA levels, our data suggest

sex differences exist in both the expression of EE cell-derived hormones and in the

ability of tissues to respond to available peptide.

158 **2.2. Sex determination gene *transformer* does not regulate sex differences in EE**

159 **cell-derived peptide mRNA levels**

160 To determine the mechanism by which these differences in mRNA levels are

161 established, we tested a role for sex determination gene *transformer* (*tra*). Normally, a

162 functional Tra protein is only expressed in females, where Tra specifies most aspects of

163 female sexual development and behavior [73–76]. Indeed, ectopic Tra expression in

164 males is sufficient to specify many aspects of female sexual development and

165 physiology [7,48,49,74,77,78]. Because *tra* mRNA is detected in the gut, specifically in

166 ISC and EE cells [48,79], we asked whether broad overexpression of Tra in neurons

167 and/or EE cells contributes to the sex difference in mRNA levels of EE cell-expressed

168 hormones. For these data, cell type-specific Tra overexpression was considered to have

169 a significant effect on EE cell-expressed hormones only if the experimental genotype

170 (e.g., *tissue-GAL4>UAS-tra^F^*) significantly differed from both parental strains (e.g.,

171 *tissue-GAL4>+* and *+>UAS-tra^F^*) with the same direction of effect. We found that sex

172 differences in mRNA levels of *AstA*, *AstC*, *Tk*, *NPF,* and *Dh31* were unaffected when we

173 used either *voila-GAL4* (Figure 2A-2J) which expresses in EE and sensory cells, or

174 *elav-GAL4* (Figure 2K-2T) which expresses in neurons and neuropeptide-producing

175 cells, to drive Tra expression in these cells. However, we note that Tra expression in EE

176 cells further augments the male bias in head *Tk* mRNA levels (Figure 2H), whereas Tra

177 expression in female neurons paradoxically decreases *NPF* mRNA levels in the head

178 (Figure 2S). Tra expression in neurons similarly had no effect on mRNA levels of *AstC-*

179 *R2*, *TkR99D*, or *Dh31-R* in the head (Figure S2A-C). Thus, sex determination *tra* does

not establish sex differences in levels of the mRNAs that encode either EE cell-derived

hormones or their receptors.

2.3. Gut-derived Tachykinin and Allatostatin C promote female fat storage

Given that EE cell-derived AstC, NPF, and Tk regulate fat storage and phenotypes

associated with fat storage (e.g., starvation resistance) in single- and mixed-sex animal

groups [22,29,44,72], we wanted to assess whether these hormones contribute to the

sex difference in fat storage. We used RNAi to knock down levels of *AstC, Tk* and *NPF*

with GAL4 drivers targeting these specific EE populations (*AstC-GAL4, Tk-GAL4,* and

NPF-GAL4, respectively). Importantly, GAL4 activity for each driver line was restricted

to the gut using *R57C10-GAL80*, a validated approach to target only the gut cells that

produce these hormones [22,44,45]. For all fat storage data, cell type-specific RNAi was

considered to have a significant effect on fat storage only if the experimental genotype

(e.g., *tissue-GAL4>UAS-RNAi*) significantly differed from both parental strains (e.g.,

tissue-GAL4>+ and *+>UAS-RNAi*) with the same direction of effect. We found gut-

specific loss of *AstC* (genotype *AstC-GAL4>UAS-AstC-RNAi, R57C10-GAL80*) caused

a significant reduction in body fat in females with no effect in males (Figure 3A). Gut-

specific knockdown of *Tk* (genotype *Tk-GAL4>UAS-Tk-RNAi, R57C10-GAL80*) similarly

showed a trend toward a female-specific decrease in fat storage (p^{GAL4}=0.1109;

p^{UAS}=0.0118), with no significant effect on male body fat (Figure 3B). This suggests a

role for gut-derived AstC and a potential role for gut-derived Tk in regulating female

body fat, whereas gut-derived AstC or Tk do not play a role in regulating male body fat.

In contrast, gut-specific loss of *NPF* (*NPF-GAL4>UAS-NPF-RNAi, R57C10-GAL80*) did

not significantly alter whole-body fat storage in either males or females (Figure 3C).

Together, these data suggest that the female-biased expression of AstC and Tk in the

gut play physiologically significant roles in regulating fat storage in females.

2.4. Allatostatin C receptor and Tachykinin receptor in neurons promote fat

storage in females but not males

Neurons and neuropeptide-producing cells are key cell types upon which AstC, Tk, and

NPF act to influence physiology [22,29,32,72,80,81]. We therefore predicted that loss of

AstC-R2 and *TkR99D* in these cells would reproduce the reduced fat storage we

observed in females with loss of EE cell-derived AstC and Tk, though we cannot fully

rule out effects of Tk and AstC mediated by other receptors as we did not test these

additional receptors [82,83]. To test this, we used *elav-GAL4* to knock down *AstC-R2*

and *TkR99D* in post-mitotic neurons and neuropeptide-producing cells. In females, loss

of *AstC-R2* in neurons and neuropeptide-producing cells caused a significant decrease

in body fat, with no effect in males (Figure 4A). This reproduced the body fat phenotype

caused by loss of gut AstC. A similar female-specific reduction in fat storage was

observed with loss of *TkR99D* in neurons and neuropeptide-producing cells (Figure 4B),

reproducing the potential fat storage phenotype of females with loss of gut-derived Tk.

In line with the lack of body fat effect due to loss of gut-derived NPF, we saw no

significant change in fat storage in either males or females with loss of *NPFR* in neurons

and neuropeptide-producing cells (Figure 4C). Together, these data suggest that AstC

and Tk may promote whole-body fat storage in females via effects on neurons and

neuropeptide-producing cells.

To narrow down the neurons and neuropeptide-producing cells in which *AstC-R2*

and *TkR99D* act to mediate their effects on female fat storage, we used cell-type-

specific GAL4 drivers to overexpress RNAi transgenes directed at these genes. Given

that these gut-derived peptides have been shown to influence metabolic homeostasis

and feeding via effects on the insulin-producing cells (IPCs) and the Akh-producing cells

(APCs) [19,22,68], we first knocked down *AstC-R2* and *TkR99D* in these cells. We used

dilp2-GAL4 to drive expression of *UAS-Astc-R2-RNAi* and *UAS-TkR99D-RNAi* in the

IPCs, and *Akh-GAL4* to drive expression of these transgenes in the APCs. Loss of

AstC-R2 in the IPC had no significant effect on fat storage in either males or females

compared with sex-matched controls (Figure 4D). IPC-specific loss of *TkR99D*, on the

other hand, caused a significant increase in whole-body fat storage in males (Figure 4E)

with no change in females. In the APC, loss of neither receptor altered fat storage in

males or females (Figure 4F, 4G). These findings are interesting for several reasons.

For example, in males, loss of EE cell-derived *Tk* and loss of *TkR99D* across neurons

had no effect on fat storage, in contrast to the greater fat storage observed with IPC-

specific *TkR99D* loss. This suggests that Tk derived from outside of the gut, and likely in

the head, regulates fat storage via effects on *TkR99D* in the IPC. Future experiments

will be needed to test this model, and to determine how Tk affects IPC biology in males.

Further studies will also be needed to understand why IPC but not pan-neuronal loss of

TkR99D causes an effect on male body fat. Possible explanations include greater

knockdown in the IPC using *Dilp2-GAL4,* or that Tk mediates opposing effects on body

fat via effects on *TkR99D* in multiple neuron groups. In females, more work will be

needed to identify the neurons upon which Tk acts to regulate body fat, and to test the

relative contributions of EE cell- and brain-derived Tk in regulating body fat.

3. DISCUSSION

EE cell-derived hormones regulate body fat in single- and mixed-sex animal groups;

however, it has been unclear whether the regulation and function of these peptides

differ between the sexes. The goal of our study was to perform a detailed comparison of

EE cell-derived hormones between the sexes, and to test if these hormones contribute

to the sex difference in fat storage. Our assessment revealed profound female-biased

expression of EE cell-expressed hormones within the gut. This differential expression

was physiologically significant, as we showed that EE cell-derived Tk and AstC promote

female fat storage. Interestingly, these effects were not mediated by the IPC or APC,

cells that we have previously shown contribute to the sex difference in fat storage.

Taken together, our data provide additional insight into the highly complex

mechanism(s) by which unmated female flies achieve higher fat storage than male flies

(Figure 5).

While it was not the main goal of our study, our survey of EE cell-expressed

hormones in *Drosophila* revealed that the sex bias in expression was not uniform across

tissues. In the gut, mRNA levels of *AstC*, *Tk*, and *NPF* were higher in females than in

males. In the brain, mRNA levels of these hormones showed a significant male bias, in

line with data from previous reports on Tk [57] and NPF [64]. While it remains unclear

whether the tissue-specific sex bias in expression is physiologically significant, peptides

derived from the gut and the brain have been shown to mediate distinct effects on

physiology and/or behavior. For example, EE cell-derived AstC regulates energy

homeostasis and food-seeking behaviors in adult females [22], whereas neuron-derived

AstC is involved in regulating locomotion in adult males [60] and the circadian regulation

of oogenesis in adult females [70]. Neuron-derived Tk similarly regulates locomotion

[65,69], food consumption [68], Dilp secretion [19], and aggression [57], whereas gut-

derived Tk regulates intestinal lipogenesis [72] and stem cell divisions in the midgut

[84–86]. Supporting a potential sex-specific role for peptides derived from different

anatomical sites in regulating physiology, gut-derived AstC stimulates fat breakdown

during starvation through the Akh pathway in mated females with no effect in males

[22]. Future studies are therefore needed to determine whether there are sex

differences in whether the effects of EE cell-derived hormones are primarily mediated

by local or systemic mechanisms.

Another important task for future studies will be to elucidate how sex differences

in neuropeptide expression are established. The first step in understanding these

mechanisms will be to determine which factors specify the sex bias in neuropeptide

mRNA levels. Because our data shows that sex determination gene *tra* does not

regulate the sex bias in neuropeptide expression in either the brain or the gut, the role

of other factors that influence sexual identity and sexual differentiation must be

assessed. One strong candidate is the steroid hormone ecdysone, as virgin females

have higher ecdysone titers than males [87–89]. Ecdysone plays a role in regulating

sexual differentiation and development [5,90,91], and contributes to male-female

differences in multiple aspects of intestinal physiology (e.g., intestinal stem cell

proliferation) [46,50] and brain development [92,93]. Another candidate is juvenile

hormone, which has been shown to regulate sexual maturation in *Drosophila* and other

insects [94–100]. While it remains unclear whether juvenile hormone titers differ

between virgin males and females, juvenile hormone regulates many aspects of gut

physiology in mated females (e.g., intestinal lipid accumulation, intestinal stem cell

proliferation) [54,101] and influences brain development [102]. Other than hormones, it

is possible that sex determination gene *Sex-lethal* plays a role in regulating the sex

difference in mRNA levels of EE cell-derived hormones, as *tra*-independent effects of

Sex-lethal have been described in the brain [103]. Future studies will also need to test

additional members of the sex determination pathway. While sex differences in

expression of EE cell-derived hormones does not involve *tra*, and is therefore unlikely to

involve known *tra* targets such as *fruitless* [104,105], without further experiments we

cannot fully rule out these additional sex determination pathway members.

In parallel to identifying the factor(s) responsible for establishing the sex

difference in EE cell-expressed hormones, it will be important to reveal the cellular basis

for this differential expression. For example, a sex difference in the number of EE cells

and neuropeptide-expressing cells in the brain could explain the differences in

expression. Supporting this, gut length, overall brain size, and neuron number have

been shown to differ between males and females [34,48,106–113]. In the gut, the

difference in length is at least partially due to a sex difference in the proliferation of

intestinal stem cells [46,48,49], which undergo asymmetric divisions and subsequent

differentiation to generate all gut cell types including EE cells [34,42,114–118].

In the brain, males and females differ in the number of neurons found within

many identified clusters [106,107,109–113,119], including the cells that produce NPF

[64] and Tk [57]. Differences in neuron number have been primarily attributed to sex-

specific programmed cell death [112,119–122]; however, sex differences in neuroblast

cell death and/or proliferation may also play a role [111,123–125].

Beyond the effects of cell number, sex differences in EE cell-derived hormone

mRNA levels may also be due to differential gene and/or protein expression of these

factors, which have been reported for other peptide hormones [7,18,64]. Because sex

differences in the activity of peptide hormone-producing cells have also been previously

described [7], it is clear that a detailed examination of sex differences in EE cells, and

more generally in neuropeptide-producing cells, is needed to gain a comprehensive

picture of how these cells differ between males and females. Benefits of such a detailed

study include gaining insight into potential mechanisms underlying sex differences in

other aspects of physiology and behavior. For example, EE cells regulate ISC

homeostasis [84,126], and EE cell-derived hormones act locally and systemically to

regulate appetite, food ingestion, food digestion, gut motility, and immune responses

[35,81,127,128]. Importantly, male-female differences in many of these phenotypes

have been reported [52,53,129–132].

Overall, our findings identify EE cell-derived hormones AstC and Tk as important

factors that promote higher fat storage in *Drosophila* adult virgin females. This builds on

a recent paper identifying a key role for IIS in promoting higher levels of fat storage in

unmated females but not males [18], advancing knowledge of the factors that establish

an optimal level of stored fat in each sex.

4. MATERIALS AND METHODS

4.1. Fly strains

The following fly strains from the Bloomington *Drosophila* Stock Center were used: w^{1118} (RRID:BDSC_3605), *voila-GAL4* (RRID:BDSC_80572), *dilp2-GAL4* (IPCs) (RRID:BDSC_37516), *UAS-tra^F* (RRID:BDSC_4590), *UAS-NPF-RNAi* (RRID:BDSC_27237), *UAS-TkR99D-RNAi* (RRID:BDSC_27513), *UAS-AstC-RNAi* (RRID:BDSC_25868), *UAS-NPFR-RNAi* (RRID:BDSC_25939), *UAS-AstC-R2-RNAi* (RRID:BDSC_36888), *UAS-Tk-RNAi* (RRID:BDSC_25800), *elav-GAL4* (RRID:BDSC_458). We obtained *R57C10-GAL80; NPF-GAL4, R57C10-GAL80; Tk-GAL4 and R57C10-GAL80; AstC-GAL4* as kind gifts from Dr. Kim Rewitz at the University of Copenhagen, and *Akh-GAL4* was a kind gift from Dr. Mike Gordon at The University of British Columbia. We acknowledge FlyBase as an essential resource providing genetic, genomic, and functional data and tools that supported this study [133].

4.2. Fly husbandry

Fly media was prepared with the following ingredients: 20.5 g/L sucrose, 70.9 g/L D-glucose, 48.5 g/L cornmeal, 45.3 g/L yeast, 4.55 g/L agar, 0.5 g/L $CaCl_2•2H_2O$, 0.5 g/L $MgSO_4•7H_2O$, 11.77 mL/L acid mix (propionic acid/phosphoric acid). For all experiments, we allowed female flies to lay eggs on grape juice agar plates for 12 hr. At 24 hr after egg laying, 50 larvae were picked into vials containing 10 mL of food and reared at 22°C. Males and females were distinguished by the presence of sex combs in the late pupal period and placed into single-sex vials to eclose. After eclosion, adult flies were maintained at a density of twenty flies per vial in single-sex groups. Unless

otherwise stated, all experiments used 5- to 7-day-old unmated flies. We used unmated

flies to identify genetic factors that regulate the sex difference in body fat; mated

females were not used to avoid mating-induced changes in physiology mediated by

additional factors (*e.g.*, Sex-peptide) and behavioral changes due to altered food

preferences.

4.3.   Adult weight

Groups of 10 flies were placed in pre-weighed 1.5 ml microcentrifuge tubes (Diamed

Lab Supplies, DIATEC610-2550) and weighed on an analytical balance (Mettler-Toledo,

ME104).

4.4.   Whole-body triglyceride measurements

Triglyceride is the main form of stored fat in the body, with very little in the circulation

[134]. We therefore refer to whole-body triglyceride as 'fat storage' or 'body fat'. One

biological replicate consisted of five flies. Flies were collected in a 1.5 ml tube and

homogenized in 350 µl of 0.1% Tween (Amresco, 0777-1L) in 1X phosphate-buffered

saline (PBS; Sigma-Aldrich, P5493) using 50 µl of glass beads (Sigma-Aldrich,

Z250473) that were agitated at 8 m/s for 5 s (OMNI International BeadRuptor 24).

Triglyceride concentration was measured using the Stanbio Triglyceride Liquid Reagent

(FT7610, BD386a/d, BD386b) according to the manufacturer's instructions and as

described previously [13,18] with minor modifications. Briefly, 10 µl of either

homogenate or triglyceride standard (FT7610) was added to 190 µl of activated

triglyceride reagent (Enzymatic Triglyceride Reagent, BD386a/d; Triglyceride Activator,

Cat. No. BD386b) in a 96-well plate. After a 15 min incubation at room temperature, the

absorbance was read at 540 nm (Thermo Scientific – Multiskan FC Microplate

Photometer).

4.5.   RNA extraction, cDNA synthesis, and Quantitative real-time PCR (qPCR)

One biological replicate consisted of 3-5 adult fly guts, or 10 adult fly heads, or 5 whole-

body adult flies. Samples were homogenized in 500 µl Trizol (Thermo Fisher Scientific;

15596018). Chloroform was added to Trizol to separate the mixture into aqueous and

organic phases, and isopropanol was added to the aqueous phase (in a fresh tube) to

precipitate the RNA. RNA was resuspended in either 20-25 µl (for guts and heads) or

200 µl (for whole-body) of molecular biology grade water (Corning, 46-000-CV). RNA

was stored at -80°C until use. Each experiment contained 5-10 biological replicates per

sex and per genotype; each experiment was repeated twice.

For genomic DNA elimination and cDNA synthesis, an equal amount of RNA per

reaction was DNase-treated and reverse transcribed according to manufacturer's

instructions using the QuantiTect Reverse Transcription Kit (Qiagen, 205314). Relative

mRNA transcript levels were quantified using qPCR as described previously [6]. Data

were normalized to the average fold change of *Actin5C* and *β-tubulin*. For a full primer

list, refer to Document S1.

4.6.   Statistical analysis

Statistical analyses and data presentation were completed using GraphPad Prism 9

(GraphPad Software, San Diego, CA, USA). All data were tested for normality using the

Shapiro-Wilk test. Normally-distributed data were subjected to parametric tests as

appropriate, including Student's *t*-test and two-way ANOVA followed by Bonferroni post-

hoc test. For non-normally distributed data, we used the Mann-Whitney test. For two-

way ANOVA involving data that do not satisfy the normality assumption, aligned rank

transformation was first applied using the art() function from the ARTool R package

[135]. Then, ANOVA was performed on the transformed data with the base R anova()

function. Finally, the art.con() function from the ARTool package was used to extract the

main as well as the interaction effects. Default parameters were used in each step of

the analysis. For all statistical analyses, differences were considered significant if

$p < 0.05$.

CRediT authorship contribution statement

Puja Biswas: Conceptualization, Data curation, Formal analysis, Investigation,

Methodology, Visualization, Writing – original draft and editing.

Elizabeth J. Rideout: Conceptualization, Funding acquisition, Methodology,

Resources, Supervision, Writing – original draft, Writing – review and editing.

Funding

This study was supported by operating grants to EJR from the Canadian Institutes for

Health Research (PJT-153072 and PJT-183786), CIHR Sex and Gender Science Chair

program (GS4-171365), Michael Smith Foundation for Health Research (16876), and

the Canada Foundation for Innovation (JELF-34879). PB was supported by a 4-year

CELL fellowship from UBC.

Acknowledgments

The authors thank FlyBase, which is supported by a grant from the National Human

Genome Research Institute at the U.S. National Institutes of Health (U41 HG000739)

and by the British Medical Research Council (MR/N030117/1). Stocks obtained from the

Bloomington *Drosophila* Stock Center (NIHP40OD018537) were used in this study. The

authors thank the TRiP at Harvard Medical School (NIH/NIGMS R01-GM084947) for

providing transgenic RNAi fly stocks and/or plasmid vectors used in this study. The

funders had no role in study design, data collection and analysis, decision to publish, or

preparation of the manuscript. We thank members of the Rideout lab for valuable

feedback. We acknowledge that our research takes place on the traditional, ancestral,

and unceded territory of the Musqueam people; a privilege for which we are grateful.

Declaration of competing interests

The authors declare no competing interests.

Appendix A. Supplementary data

Document S1. Supplemental Figures S1-S2

Data availability

Table S1. Excel file containing raw data with calculations

Table S2. Excel file containing statistics for all data

References

[1]   De Groef, S., Wilms, T., Balmand, S., Calevro, F., Callaerts, P., 2021. Sexual
Dimorphism in Metabolic Responses to Western Diet in Drosophila melanogaster.
Biomolecules 12(1): 33, Doi: 10.3390/biom12010033.

[2]   Lease, H.M., Wolf, B.O., 2011. Lipid content of terrestrial arthropods in relation to
body size, phylogeny, ontogeny and sex. Physiological Entomology 36(1): 29–38,
Doi: 10.1111/j.1365-3032.2010.00767.x.

[3]   Parisi, M., Li, R., Oliver, B., 2011. Lipid profiles of female and male Drosophila.
BMC Research Notes 4(1): 198, Doi: 10.1186/1756-0500-4-198.

[4]   Schwasinger-Schmidt, T.E., Kachman, S.D., Harshman, L.G., 2012. Evolution of
starvation resistance in Drosophila melanogaster: measurement of direct and
correlated responses to artificial selection. Journal of Evolutionary Biology 25(2):
378–87, Doi: 10.1111/j.1420-9101.2011.02428.x.

[5]   Sieber, M.H., Spradling, A.C., 2015. Steroid Signaling Establishes a Female
Metabolic State and Regulates SREBP to Control Oocyte Lipid Accumulation.
Current Biology: CB 25(8): 993–1004, Doi: 10.1016/j.cub.2015.02.019.

[6]   Wat, L.W., Chao, C., Bartlett, R., Buchanan, J.L., Millington, J.W., Chih, H.J., et
al., 2020. A role for triglyceride lipase brummer in the regulation of sex differences
in Drosophila fat storage and breakdown. PLoS Biology 18(1): e3000595, Doi:
10.1371/journal.pbio.3000595.

[7]   Wat, L.W., Chowdhury, Z.S., Millington, J.W., Biswas, P., Rideout, E.J., 2021. Sex
determination gene transformer regulates the male-female difference in
Drosophila fat storage via the adipokinetic hormone pathway. eLife 10: e72350,
Doi: 10.7554/eLife.72350.

[8]   Chao, C.F., Pesch, Y.-Y., Yu, H., Wang, C., Aristizabal, M.J., Huan, T., et al.,
2024. An important role for triglyceride in regulating spermatogenesis. eLife 12,
Doi: 10.7554/eLife.87523.3.

[9]   Gutierrez, E., Wiggins, D., Fielding, B., Gould, A.P., 2007. Specialized
hepatocyte-like cells regulate Drosophila lipid metabolism. Nature 445(7125):
275–80, Doi: 10.1038/nature05382.

[10]   Kis, V., Barti, B., Lippai, M., Sass, M., 2015. Specialized Cortex Glial Cells
Accumulate Lipid Droplets in Drosophila melanogaster. PloS One 10(7):
e0131250, Doi: 10.1371/journal.pone.0131250.

[11]   Kühnlein, R.P., 2011. The contribution of the Drosophila model to lipid droplet
research. Progress in Lipid Research 50(4): 348–56, Doi:
10.1016/j.plipres.2011.04.001.

[12]   Kühnlein, R.P., 2012. Thematic review series: Lipid droplet synthesis and
metabolism: from yeast to man. Lipid droplet-based storage fat metabolism in
Drosophila. Journal of Lipid Research 53(8): 1430–6, Doi: 10.1194/jlr.R024299.

[13]   Sieber, M.H., Thummel, C.S., 2009. The DHR96 nuclear receptor controls
triacylglycerol homeostasis in Drosophila. Cell Metabolism 10(6): 481–90, Doi:
10.1016/j.cmet.2009.10.010.

[14]   Villanueva, J.E., Livelo, C., Trujillo, A.S., Chandran, S., Woodworth, B., Andrade,
L., et al., 2019. Time-restricted feeding restores muscle function in Drosophila

models of obesity and circadian-rhythm disruption. Nature Communications 10(1):
2700, Doi: 10.1038/s41467-019-10563-9.

[15]   Arrese, E.L., Soulages, J.L., 2010. INSECT FAT BODY: ENERGY,
       METABOLISM, AND REGULATION. Annual Review of Entomology 55: 207–25,
       Doi: 10.1146/annurev-ento-112408-085356.

[16]   Buszczak, M., Lu, X., Segraves, W.A., Chang, T.Y., Cooley, L., 2002. Mutations in
       the midway gene disrupt a Drosophila acyl coenzyme A: diacylglycerol
       acyltransferase. Genetics 160(4): 1511–8, Doi: 10.1093/genetics/160.4.1511.

[17]   Matsuoka, S., Armstrong, A.R., Sampson, L.L., Laws, K.M., Drummond-Barbosa,
       D., 2017. Adipocyte Metabolic Pathways Regulated by Diet Control the Female
       Germline Stem Cell Lineage in Drosophila melanogaster. Genetics 206(2): 953–
       71, Doi: 10.1534/genetics.117.201921.

[18]   Biswas, P., Bako, J.A., Liston, J.B., Yu, H., Wat, L.W., Miller, C.J., et al., 2025.
       Insulin/insulin-like growth factor signaling pathway promotes higher fat storage in
       Drosophila females. Cell Reports 44(7), Doi: 10.1016/j.celrep.2025.115915.

[19]   Birse, R.T., Söderberg, J.A.E., Luo, J., Winther, A.M.E., Nässel, D.R., 2011.
       Regulation of insulin-producing cells in the adult Drosophila brain via the
       tachykinin peptide receptor DTKR. The Journal of Experimental Biology 214(Pt
       24): 4201–8, Doi: 10.1242/jeb.062091.

[20]   Hentze, J.L., Carlsson, M.A., Kondo, S., Nässel, D.R., Rewitz, K.F., 2015. The
       Neuropeptide Allatostatin A Regulates Metabolism and Feeding Decisions in
       Drosophila. Scientific Reports 5: 11680, Doi: 10.1038/srep11680.

[21]   Kapan, N., Lushchak, O.V., Luo, J., Nässel, D.R., 2012. Identified peptidergic
       neurons in the Drosophila brain regulate insulin-producing cells, stress responses
       and metabolism by coexpressed short neuropeptide F and corazonin. Cellular and
       Molecular Life Sciences 69(23): 4051–66, Doi: 10.1007/s00018-012-1097-z.

[22]   Kubrak, O., Koyama, T., Ahrentløv, N., Jensen, L., Malita, A., Naseem, M.T., et
       al., 2022. The gut hormone Allatostatin C/Somatostatin regulates food intake and
       metabolic homeostasis under nutrient stress. Nature Communications 13(1): 692,
       Doi: 10.1038/s41467-022-28268-x.

[23]   Kubrak, O., Jørgensen, A.F., Koyama, T., Lassen, M., Nagy, S., Hald, J., et al.,
       2024. LGR signaling mediates muscle-adipose tissue crosstalk and protects
       against diet-induced insulin resistance. Nature Communications 15(1): 6126, Doi:
       10.1038/s41467-024-50468-w.

[24]   Kubrak, O.I., Lushchak, O.V., Zandawala, M., Nässel, D.R., 2016. Systemic
       corazonin signalling modulates stress responses and metabolism in Drosophila.
       Open Biology 6(11): 160152, Doi: 10.1098/rsob.160152.

[25]   Nässel, D.R., Enell, L.E., Santos, J.G., Wegener, C., Johard, H.A., 2008. A large
       population of diverse neurons in the Drosophilacentral nervous system expresses
       short neuropeptide F, suggesting multiple distributed peptide functions. BMC
       Neuroscience 9(1): 90, Doi: 10.1186/1471-2202-9-90.

[26]   Oh, Y., Lai, J.S.-Y., Mills, H.J., Erdjument-Bromage, H., Giammarinaro, B.,
       Saadipour, K., et al., 2019. A glucose-sensing neuron pair regulates insulin and
       glucagon in Drosophila. Nature 574(7779): 559–64, Doi: 10.1038/s41586-019-
       1675-4.

[27] Sano, H., Nakamura, A., Texada, M.J., Truman, J.W., Ishimoto, H., Kamikouchi,
A., et al., 2015. The Nutrient-Responsive Hormone CCHamide-2 Controls Growth
by Regulating Insulin-like Peptides in the Brain of Drosophila melanogaster. PLOS
Genetics 11(5): e1005209, Doi: 10.1371/journal.pgen.1005209.
[28] Scopelliti, A., Bauer, C., Yu, Y., Zhang, T., Kruspig, B., Murphy, D.J., et al., 2019.
A Neuronal Relay Mediates a Nutrient Responsive Gut/Fat Body Axis Regulating
Energy Homeostasis in Adult Drosophila. Cell Metabolism 29(2): 269-284.e10,
Doi: 10.1016/j.cmet.2018.09.021.
[29] Yoshinari, Y., Kosakamoto, H., Kamiyama, T., Hoshino, R., Matsuoka, R., Kondo,
S., et al., 2021. The sugar-responsive enteroendocrine neuropeptide F regulates
lipid metabolism through glucagon-like and insulin-like hormones in Drosophila
melanogaster. Nature Communications 12(1): 4818, Doi: 10.1038/s41467-021-
25146-w.
[30] Zandawala, M., Yurgel, M.E., Texada, M.J., Liao, S., Rewitz, K.F., Keene, A.C., et
al., 2018. Modulation of Drosophila post-feeding physiology and behavior by the
neuropeptide leucokinin. PLOS Genetics 14(11): e1007767, Doi:
10.1371/journal.pgen.1007767.
[31] Nässel, D.R., Broeck, J.V., 2016. Insulin/IGF signaling in Drosophila and other
insects: factors that regulate production, release and post-release action of the
insulin-like peptides. Cellular and Molecular Life Sciences 73(2): 271–90, Doi:
10.1007/s00018-015-2063-3.
[32] Nässel, D.R., Zandawala, M., 2019. Recent advances in neuropeptide signaling in
Drosophila, from genes to physiology and behavior. Progress in Neurobiology
179: 101607, Doi: 10.1016/j.pneurobio.2019.02.003.
[33] Toprak, U., 2020. The Role of Peptide Hormones in Insect Lipid Metabolism.
Frontiers in Physiology 11, Doi: 10.3389/fphys.2020.00434.
[34] Miguel-Aliaga, I., Jasper, H., Lemaitre, B., 2018. Anatomy and Physiology of the
Digestive Tract of Drosophila melanogaster. Genetics 210(2): 357–96, Doi:
10.1534/genetics.118.300224.
[35] Wegener, C., Veenstra, J.A., 2015. Chemical identity, function and regulation of
enteroendocrine peptides in insects. Current Opinion in Insect Science 11: 8–13,
Doi: 10.1016/j.cois.2015.07.003.
[36] Zhou, X., Ding, G., Li, J., Xiang, X., Rushworth, E., Song, W., 2020. Physiological
and Pathological Regulation of Peripheral Metabolism by Gut-Peptide Hormones
in Drosophila. Frontiers in Physiology 11: 577717, Doi:
10.3389/fphys.2020.577717.
[37] Chen, J., Kim, S., Kwon, J.Y., 2016. A Systematic Analysis of *Drosophila*
Regulatory Peptide Expression in Enteroendocrine Cells. Molecules and Cells
39(4): 358–66, Doi: 10.14348/molcells.2016.0014.
[38] Hung, R.-J., Hu, Y., Kirchner, R., Liu, Y., Xu, C., Comjean, A., et al., 2020. A cell
atlas of the adult Drosophila midgut. Proceedings of the National Academy of
Sciences of the United States of America 117(3): 1514–23, Doi:
10.1073/pnas.1916820117.
[39] Reiher, W., Shirras, C., Kahnt, J., Baumeister, S., Isaac, R.E., Wegener, C., 2011.
Peptidomics and peptide hormone processing in the Drosophila midgut. Journal of
Proteome Research 10(4): 1881–92, Doi: 10.1021/pr101116g.

[40] Veenstra, J.A., 2009. Peptidergic paracrine and endocrine cells in the midgut of
the fruit fly maggot. Cell and Tissue Research 336(2): 309–23, Doi:
10.1007/s00441-009-0769-y.
[41] Veenstra, J.A., Agricola, H.-J., Sellami, A., 2008. Regulatory peptides in fruit fly
midgut. Cell and Tissue Research 334(3): 499–516, Doi: 10.1007/s00441-008-
0708-3.
[42] Micchelli, C.A., Perrimon, N., 2006. Evidence that stem cells reside in the adult
Drosophila midgut epithelium. Nature 439(7075): 475–9, Doi:
10.1038/nature04371.
[43] Ohlstein, B., Spradling, A., 2006. The adult Drosophila posterior midgut is
maintained by pluripotent stem cells. Nature 439(7075): 470–4, Doi:
10.1038/nature04333.
[44] Malita, A., Kubrak, O., Koyama, T., Ahrentløv, N., Texada, M.J., Nagy, S., et al.,
2022. A gut-derived hormone suppresses sugar appetite and regulates food
choice in Drosophila. Nature Metabolism 4(11): 1532–50, Doi: 10.1038/s42255-
022-00672-z.
[45] Kubrak, O., Malita, A., Ahrentløv, N., Nagy, S., Texada, M.J., Rewitz, K., 2026.
Gut hormone signaling drives sex differences in metabolism and behavior.
Molecular Metabolism 103: 102312, Doi: 10.1016/j.molmet.2025.102312.
[46] Ahmed, S.M.H., Maldera, J.A., Krunic, D., Paiva-Silva, G.O., Pénalva, C.,
Teleman, A.A., et al., 2020. Fitness trade-offs incurred by ovary-to-gut steroid
signalling in Drosophila. Nature 584(7821): 415–9, Doi: 10.1038/s41586-020-
2462-y.
[47] Blackie, L., Gaspar, P., Mosleh, S., Lushchak, O., Kong, L., Jin, Y., et al., 2024.
The sex of organ geometry. Nature 630(8016): 392–400, Doi: 10.1038/s41586-
024-07463-4.
[48] Hudry, B., Khadayate, S., Miguel-Aliaga, I., 2016. The sexual identity of adult
intestinal stem cells controls organ size and plasticity. Nature 530(7590): 344–8,
Doi: 10.1038/nature16953.
[49] Regan, J.C., Khericha, M., Dobson, A.J., Bolukbasi, E., Rattanavirotkul, N.,
Partridge, L., 2016. Sex difference in pathology of the ageing gut mediates the
greater response of female lifespan to dietary restriction. eLife 5: e10956, Doi:
10.7554/eLife.10956.
[50] Zipper, L., Jassmann, D., Burgmer, S., Görlich, B., Reiff, T., 2020. Ecdysone
steroid hormone remote controls intestinal stem cell fate decisions via the PPARγ-
homolog Eip75B in Drosophila. eLife 9: e55795, Doi: 10.7554/eLife.55795.
[51] Regan, J.C., Lu, Y.-X., Ureña, E., Meilenbrock, R.L., Catterson, J.H., Kißler, D., et
al., 2022. Sexual identity of enterocytes regulates autophagy to determine
intestinal health, lifespan and responses to rapamycin. Nature Aging 2(12): 1145–
58, Doi: 10.1038/s43587-022-00308-7.
[52] Cognigni, P., Bailey, A.P., Miguel-Aliaga, I., 2011. Enteric neurons and systemic
signals couple nutritional and reproductive status with intestinal homeostasis. Cell
Metabolism 13(1): 92–104, Doi: 10.1016/j.cmet.2010.12.010.
[53] Hadjieconomou, D., King, G., Gaspar, P., Mineo, A., Blackie, L., Ameku, T., et al.,
2020. Enteric neurons increase maternal food intake during reproduction. Nature
587(7834): 455–9, Doi: 10.1038/s41586-020-2866-8.

[54] Reiff, T., Jacobson, J., Cognigni, P., Antonello, Z., Ballesta, E., Tan, K.J., et al.,
2015. Endocrine remodelling of the adult intestine sustains reproduction in
Drosophila. eLife 4: e06930, Doi: 10.7554/eLife.06930.
[55] White, M.A., Bonfini, A., Wolfner, M.F., Buchon, N., 2021. Drosophila
melanogaster sex peptide regulates mated female midgut morphology and
physiology. Proceedings of the National Academy of Sciences 118(1):
e2018112118, Doi: 10.1073/pnas.2018112118.
[56] Abruzzi, K.C., Zadina, A., Luo, W., Wiyanto, E., Rahman, R., Guo, F., et al., 2017.
RNA-seq analysis of Drosophila clock and non-clock neurons reveals neuron-
specific cycling and novel candidate neuropeptides. PLOS Genetics 13(2):
e1006613, Doi: 10.1371/journal.pgen.1006613.
[57] Asahina, K., Watanabe, K., Duistermars, B.J., Hoopfer, E., González, C.R.,
Eyjólfsdóttir, E.A., et al., 2014. Tachykinin-Expressing Neurons Control Male-
Specific Aggressive Arousal in Drosophila. Cell 156(1): 221–35, Doi:
10.1016/j.cell.2013.11.045.
[58] Beshel, J., Dubnau, J., Zhong, Y., 2017. A Leptin analog locally produced in the
brain acts via a conserved neural circuit to modulate obesity-linked behaviors in
Drosophila. Cell Metabolism 25(1): 208, Doi: 10.1016/j.cmet.2016.12.013.
[59] Chung, B.Y., Ro, J., Hutter, S.A., Miller, K.M., Guduguntla, L.S., Kondo, S., et al.,
2017. Drosophila Neuropeptide F Signaling Independently Regulates Feeding and
Sleep-Wake Behavior. Cell Reports 19(12): 2441–50, Doi:
10.1016/j.celrep.2017.05.085.
[60] Díaz, M.M., Schlichting, M., Abruzzi, K.C., Long, X., Rosbash, M., 2019.
Allatostatin-C/AstC-R2 Is a Novel Pathway to Modulate the Circadian Activity
Pattern in Drosophila. Current Biology: CB 29(1): 13-22.e3, Doi:
10.1016/j.cub.2018.11.005.
[61] Hergarden, A.C., Tayler, T.D., Anderson, D.J., 2012. Allatostatin-A neurons inhibit
feeding behavior in adult Drosophila. Proceedings of the National Academy of
Sciences 109(10): 3967–72, Doi: 10.1073/pnas.1200778109.
[62] Kunst, M., Hughes, M.E., Raccuglia, D., Felix, M., Li, M., Barnett, G., et al., 2014.
Calcitonin gene-related peptide neurons mediate sleep-specific circadian output in
Drosophila. Current Biology: CB 24(22): 2652–64, Doi:
10.1016/j.cub.2014.09.077.
[63] Kurogi, Y., Imura, E., Mizuno, Y., Hoshino, R., Nouzova, M., Matsuyama, S., et
al., 2023. Female reproductive dormancy in Drosophila is regulated by DH31-
producing neurons projecting into the corpus allatum. Development (Cambridge,
England) 150(10): dev201186, Doi: 10.1242/dev.201186.
[64] Lee, G., Bahn, J.H., Park, J.H., 2006. Sex- and clock-controlled expression of the
neuropeptide F gene in Drosophila. Proceedings of the National Academy of
Sciences 103(33): 12580–5, Doi: 10.1073/pnas.0601171103.
[65] Lee, S.H., Cho, E., Yoon, S.-E., Kim, Y., Kim, E.Y., 2021. Metabolic control of
daily locomotor activity mediated by tachykinin in Drosophila. Communications
Biology 4(1): 1–14, Doi: 10.1038/s42003-021-02219-6.
[66] Luan, H., Lemon, W.C., Peabody, N.C., Pohl, J.B., Zelensky, P.K., Wang, D., et
al., 2006. Functional Dissection of a Neuronal Network Required for Cuticle

Tanning and Wing Expansion in Drosophila. Journal of Neuroscience 26(2): 573–84, Doi: 10.1523/JNEUROSCI.3916-05.2006.

[67] Mendive, F.M., Van Loy, T., Claeysen, S., Poels, J., Williamson, M., Hauser, F., et al., 2005. Drosophila molting neurohormone bursicon is a heterodimer and the natural agonist of the orphan receptor DLGR2. FEBS Letters 579(10): 2171–6, Doi: 10.1016/j.febslet.2005.03.006.

[68] Qi, W., Wang, G., Wang, L., 2021. A novel satiety sensor detects circulating glucose and suppresses food consumption via insulin-producing cells in Drosophila. Cell Research 31(5): 580–8, Doi: 10.1038/s41422-020-00449-7.

[69] Winther, Å.M.E., Acebes, A., Ferrús, A., 2006. Tachykinin-related peptides modulate odor perception and locomotor activity in *Drosophila*. Molecular and Cellular Neuroscience 31(3): 399–406, Doi: 10.1016/j.mcn.2005.10.010.

[70] Zhang, C., Daubnerova, I., Jang, Y.-H., Kondo, S., Žitňan, D., Kim, Y.-J., 2021. The neuropeptide allatostatin C from clock-associated DN1p neurons generates the circadian rhythm for oogenesis. Proceedings of the National Academy of Sciences 118(4): e2016878118, Doi: 10.1073/pnas.2016878118.

[71] Song, T., Qin, W., Lai, Z., Li, H., Li, D., Wang, B., et al., 2023. Dietary cysteine drives body fat loss via FMRFamide signaling in Drosophila and mouse. Cell Research 33(6): 434–47, Doi: 10.1038/s41422-023-00800-8.

[72] Song, W., Veenstra, J.A., Perrimon, N., 2014. Control of lipid metabolism by tachykinin in Drosophila. Cell Reports 9(1): 40–7, Doi: 10.1016/j.celrep.2014.08.060.

[73] Camara, N., Whitworth, C., Van Doren, M., 2008. The creation of sexual dimorphism in the Drosophila soma. Current Topics in Developmental Biology 83: 65–107, Doi: 10.1016/S0070-2153(08)00403-1.

[74] McKeown, M., Belote, J.M., Boggs, R.T., 1988. Ectopic expression of the female transformer gene product leads to female differentiation of chromosomally male Drosophila. Cell 53(6): 887–95, Doi: 10.1016/s0092-8674(88)90369-8.

[75] Oliver, B., 2002. Genetic control of germline sexual dimorphism in Drosophila. International Review of Cytology 219: 1–60, Doi: 10.1016/s0074-7696(02)19010-3.

[76] Sturtevant, A.H., 1945. A Gene in Drosophila Melanogaster That Transforms Females into Males. Genetics 30(3): 297–9, Doi: 10.1093/genetics/30.3.297.

[77] Billeter, J.-C., Rideout, E.J., Dornan, A.J., Goodwin, S.F., 2006. Control of male sexual behavior in Drosophila by the sex determination pathway. Current Biology: CB 16(17): R766-776, Doi: 10.1016/j.cub.2006.08.025.

[78] Rideout, E.J., Narsaiya, M.S., Grewal, S.S., 2015. The Sex Determination Gene transformer Regulates Male-Female Differences in Drosophila Body Size. PLoS Genetics 11(12): e1005683, Doi: 10.1371/journal.pgen.1005683.

[79] Hérault, C., Pihl, T., Hudry, B., 2024. Cellular sex throughout the organism underlies somatic sexual differentiation. Nature Communications 15(1): 6925, Doi: 10.1038/s41467-024-51228-6.

[80] Chopra, G., Kaushik, S., Kain, P., 2022. Nutrient Sensing via Gut in Drosophila melanogaster. International Journal of Molecular Sciences 23(5): 2694, Doi: 10.3390/ijms23052694.

[81] Guo, X., Lv, J., Xi, R., 2022. The specification and function of enteroendocrine
cells in Drosophila and mammals: a comparative review. The FEBS Journal
289(16): 4773–96, Doi: 10.1111/febs.16067.
[82] Kreienkamp, H.-J., Larusson, H.J., Witte, I., Roeder, T., Birgul, N., Honck, H.-H.,
et al., 2002. Functional annotation of two orphan G-protein-coupled receptors,
Drostar1 and -2, from Drosophila melanogaster and their ligands by reverse
pharmacology. The Journal of Biological Chemistry 277(42): 39937–43, Doi:
10.1074/jbc.M206931200.
[83] Poels, J., Birse, R.T., Nachman, R.J., Fichna, J., Janecka, A., Vanden Broeck, J.,
et al., 2009. Characterization and distribution of NKD, a receptor for Drosophila
tachykinin-related peptide 6. Peptides 30(3): 545–56, Doi:
10.1016/j.peptides.2008.10.012.
[84] Amcheslavsky, A., Song, W., Li, Q., Nie, Y., Bragatto, I., Ferrandon, D., et al.,
2014. Enteroendocrine cells support intestinal stem-cell-mediated homeostasis in
Drosophila. Cell Reports 9(1): 32–9, Doi: 10.1016/j.celrep.2014.08.052.
[85] Foronda, D., Weng, R., Verma, P., Chen, Y.-W., Cohen, S.M., 2014. Coordination
of insulin and Notch pathway activities by microRNA miR-305 mediates adaptive
homeostasis in the intestinal stem cells of the Drosophila gut. Genes &
Development 28(21): 2421–31, Doi: 10.1101/gad.241588.114.
[86] O'Brien, L.E., Soliman, S.S., Li, X., Bilder, D., 2011. Altered Modes of Stem Cell
Division Drive Adaptive Intestinal Growth. Cell 147(3): 603–14, Doi:
10.1016/j.cell.2011.08.048.
[87] Bownes, M., Dübendorfer, A., Smith, T., 1984. Ecdysteroids in adult males and
females of *Drosophila melanogaster*. Journal of Insect Physiology 30(10): 823–30,
Doi: 10.1016/0022-1910(84)90019-2.
[88] Harshman, L.G., Loeb, A.M., Johnson, B.A., 1999. Ecdysteroid titers in mated and
unmated *Drosophila melanogaster* females. Journal of Insect Physiology 45(6):
571–7, Doi: 10.1016/S0022-1910(99)00038-4.
[89] Schwedes, C.C., Carney, G.E., 2012. Ecdysone signaling in adult *Drosophila
melanogaster*. Journal of Insect Physiology 58(3): 293–302, Doi:
10.1016/j.jinsphys.2012.01.013.
[90] Li, Y., Ma, Q., Cherry, C.M., Matunis, E.L., 2014. Steroid signaling promotes stem
cell maintenance in the *Drosophila* testis. Developmental Biology 394(1): 129–41,
Doi: 10.1016/j.ydbio.2014.07.016.
[91] Riddiford, L.M., Cherbas, P., Truman, J.W., 2000. Ecdysone receptors and their
biological actions. Vitamins and Hormones 60: 1–73, Doi: 10.1016/s0083-
6729(00)60016-x.
[92] Dalton, J.E., Lebo, M.S., Sanders, L.E., Sun, F., Arbeitman, M.N., 2009.
Ecdysone receptor acts in fruitless- expressing neurons to mediate drosophila
courtship behaviors. Current Biology: CB 19(17): 1447–52, Doi:
10.1016/j.cub.2009.06.063.
[93] Zhang, B., Sato, K., Yamamoto, D., 2018. Ecdysone signaling regulates
specification of neurons with a male-specific neurite in Drosophila. Biology Open
7(2): bio029744, Doi: 10.1242/bio.029744.
[94] Argue, K.J., Yun, A.J., Neckameyer, W.S., 2013. Early manipulation of juvenile
hormone has sexually dimorphic effects on mature adult behavior in Drosophila

melanogaster. Hormones and Behavior 64(4): 589–97, Doi:
10.1016/j.yhbeh.2013.08.018.

[95] Barth, R.H., Lester, L.J., 1973. Neuro-hormonal control of sexual behavior in
insects. Annual Review of Entomology 18: 445–72, Doi:
10.1146/annurev.en.18.010173.002305.

[96] Bilen, J., Atallah, J., Azanchi, R., Levine, J.D., Riddiford, L.M., 2013. Regulation of
onset of female mating and sex pheromone production by juvenile hormone in
Drosophila melanogaster. Proceedings of the National Academy of Sciences
110(45): 18321–6, Doi: 10.1073/pnas.1318119110.

[97] Flatt, T., Tu, M.-P., Tatar, M., 2005. Hormonal pleiotropy and the juvenile
hormone regulation of Drosophila development and life history. BioEssays: News
and Reviews in Molecular, Cellular and Developmental Biology 27(10): 999–1010,
Doi: 10.1002/bies.20290.

[98] Raikhel, A.S., Brown, M.R., Belles, X., 2005. 3.9 - Hormonal Control of
Reproductive Processes. In: Gilbert, L.I., editor. Comprehensive Molecular Insect
Science, Amsterdam: Elsevier p. 433–91.

[99] Wijesekera, T.P., Saurabh, S., Dauwalder, B., 2016. Juvenile Hormone Is
Required in Adult Males for Drosophila Courtship. PloS One 11(3): e0151912,
Doi: 10.1371/journal.pone.0151912.

[100] Wyatt, G.R., Davey, K.G., 1996. Cellular and Molecular Actions of Juvenile
Hormone. II. Roles of Juvenile Hormone in Adult Insects. In: Evans, P.D., editor.
Advances in Insect Physiology, vol. 26. Academic Press p. 1–155.

[101] Rahman, M.M., Franch-Marro, X., Maestro, J.L., Martin, D., Casali, A., 2017.
Local Juvenile Hormone activity regulates gut homeostasis and tumor growth in
adult Drosophila. Scientific Reports 7(1): 11677, Doi: 10.1038/s41598-017-11199-
9.

[102] Liu, Z.-H., Zhai, Y., Xia, Y., Liao, Q., 2024. Mating modifies oxidative stress in the
brain and confers protection against Parkinson's Disease in a *Drosophila* model.
Biochemical and Biophysical Research Communications 737: 150911, Doi:
10.1016/j.bbrc.2024.150911.

[103] Evans, D.S., Cline, T.W., 2013. Drosophila switch gene Sex-lethal can bypass its
switch-gene target transformer to regulate aspects of female behavior.
Proceedings of the National Academy of Sciences of the United States of America
110(47): E4474-4481, Doi: 10.1073/pnas.1319063110.

[104] Heinrichs, V., Ryner, L.C., Baker, B.S., 1998. Regulation of Sex-Specific Selection
of fruitless 5′ Splice Sites by transformerand transformer-2. Molecular and Cellular
Biology 18(1): 450–8, Doi: 10.1128/MCB.18.1.450.

[105] Ryner, L.C., Goodwin, S.F., Castrillon, D.H., Anand, A., Villella, A., Baker, B.S., et
al., 1996. Control of Male Sexual Behavior and Sexual Orientation in Drosophila
by the fruitless Gene. Cell 87(6): 1079–89, Doi: 10.1016/S0092-8674(00)81802-4.

[106] Asahina, K., 2018. Sex differences in Drosophila behavior: Qualitative and
Quantitative Dimorphism. Current Opinion in Physiology 6: 35–45, Doi:
10.1016/j.cophys.2018.04.004.

[107] Cachero, S., Ostrovsky, A.D., Yu, J.Y., Dickson, B.J., Jefferis, G.S.X.E., 2010.
Sexual Dimorphism in the Fly Brain. Current Biology 20(18): 1589–601, Doi:
10.1016/j.cub.2010.07.045.

[108] Jiao, W., Spreemann, G., Ruchti, E., Banerjee, S., Vernon, S., Shi, Y., et al.,
2022. Intact Drosophila central nervous system cellular quantitation reveals
sexual dimorphism. eLife 11: e74968, Doi: 10.7554/eLife.74968.

[109] Kimura, K.-I., Ote, M., Tazawa, T., Yamamoto, D., 2005. Fruitless specifies
sexually dimorphic neural circuitry in the Drosophila brain. Nature 438(7065):
229–33, Doi: 10.1038/nature04229.

[110] Nojima, T., Rings, A., Allen, A.M., Otto, N., Verschut, T.A., Billeter, J.-C., et al.,
2021. A sex-specific switch between visual and olfactory inputs underlies adaptive
sex differences in behavior. Current Biology 31(6): 1175-1191.e6, Doi:
10.1016/j.cub.2020.12.047.

[111] Ren, Q., Awasaki, T., Huang, Y.-F., Liu, Z., Lee, T., 2016. Cell Class-Lineage
Analysis Reveals Sexually Dimorphic Lineage Compositions in the Drosophila
Brain. Current Biology 26(19): 2583–93, Doi: 10.1016/j.cub.2016.07.086.

[112] Sanders, L.E., Arbeitman, M.N., 2008. Doublesex establishes sexual dimorphism
in the *Drosophila* central nervous system in an isoform-dependent manner by
directing cell number. Developmental Biology 320(2): 378–90, Doi:
10.1016/j.ydbio.2008.05.543.

[113] Sato, K., Yamamoto, D., 2023. Molecular and cellular origins of behavioral sex
differences: a tiny little fly tells a lot. Frontiers in Molecular Neuroscience 16:
1284367, Doi: 10.3389/fnmol.2023.1284367.

[114] Buchon, N., Osman, D., 2015. All for one and one for all: Regionalization of the
Drosophila intestine. Insect Biochemistry and Molecular Biology 67: 2–8, Doi:
10.1016/j.ibmb.2015.05.015.

[115] Driver, I., Ohlstein, B., 2014. Specification of regional intestinal stem cell identity
during Drosophila metamorphosis. Development 141(9): 1848–56, Doi:
10.1242/dev.104018.

[116] Jiang, H., Edgar, B.A., 2011. Intestinal stem cells in the adult *Drosophila* midgut.
Experimental Cell Research 317(19): 2780–8, Doi: 10.1016/j.yexcr.2011.07.020.

[117] Lemaitre, B., Miguel-Aliaga, I., 2013. The digestive tract of Drosophila
melanogaster. Annual Review of Genetics 47: 377–404, Doi: 10.1146/annurev-
genet-111212-133343.

[118] Schonhoff, S.E., Giel-Moloney, M., Leiter, A.B., 2004. Minireview: Development
and Differentiation of Gut Endocrine Cells. Endocrinology 145(6): 2639–44, Doi:
10.1210/en.2004-0051.

[119] Kimura, K.-I., 2011. Role of cell death in the formation of sexual dimorphism in the
Drosophila central nervous system. Development, Growth & Differentiation 53(2):
236–44, Doi: 10.1111/j.1440-169X.2010.01223.x.

[120] Garner, S.R.C., Castellanos, M.C., Baillie, K.E., Lian, T., Allan, D.W., 2018.
Drosophila female-specific Ilp7 motoneurons are generated by Fruitless-
dependent cell death in males and by a double-assurance survival role for
Transformer in females. Development (Cambridge, England) 145(1): dev150821,
Doi: 10.1242/dev.150821.

[121] Kimura, K., Hachiya, T., Koganezawa, M., Tazawa, T., Yamamoto, D., 2008.
Fruitless and Doublesex Coordinate to Generate Male-Specific Neurons that Can
Initiate Courtship. Neuron 59(5): 759–69, Doi: 10.1016/j.neuron.2008.06.007.

[122] Nojima, T., Kimura, K., Koganezawa, M., Yamamoto, D., 2010. Neuronal synaptic
outputs determine the sexual fate of postsynaptic targets. Current Biology: CB
20(9): 836–40, Doi: 10.1016/j.cub.2010.02.064.
[123] Ghosh, N., Bakshi, A., Khandelwal, R., Rajan, S.G., Joshi, R., 2019. The Hox
gene Abdominal-B uses DoublesexF as a cofactor to promote neuroblast
apoptosis in the Drosophila central nervous system. Development (Cambridge,
England) 146(16): dev175158, Doi: 10.1242/dev.175158.
[124] Taylor, B.J., Truman, J.W., 1992. Commitment of abdominal neuroblasts in
Drosophila to a male or female fate is dependent on genes of the sex-determining
hierarchy. Development (Cambridge, England) 114(3): 625–42, Doi:
10.1242/dev.114.3.625.
[125] Truman, J.W., Bate, M., 1988. Spatial and temporal patterns of neurogenesis in
the central nervous system of Drosophila melanogaster. Developmental Biology
125(1): 145–57, Doi: 10.1016/0012-1606(88)90067-x.
[126] Scopelliti, A., Cordero, J.B., Diao, F., Strathdee, K., White, B.H., Sansom, O.J., et
al., 2014. Local Control of Intestinal Stem Cell Homeostasis by Enteroendocrine
Cells in the Adult Drosophila Midgut. Current Biology 24(11): 1199–211, Doi:
10.1016/j.cub.2014.04.007.
[127] Kamareddine, L., Robins, W.P., Berkey, C.D., Mekalanos, J.J., Watnick, P.I.,
2018. The Drosophila immune deficiency pathway modulates enteroendocrine
function and host metabolism. Cell Metabolism 28(3): 449-462.e5, Doi:
10.1016/j.cmet.2018.05.026.
[128] Ren, G.R., Hauser, F., Rewitz, K.F., Kondo, S., Engelbrecht, A.F., Didriksen, A.K.,
et al., 2015. CCHamide-2 Is an Orexigenic Brain-Gut Peptide in Drosophila. PloS
One 10(7): e0133017, Doi: 10.1371/journal.pone.0133017.
[129] Belmonte, R.L., Corbally, M.-K., Duneau, D.F., Regan, J.C., 2020. Sexual
Dimorphisms in Innate Immunity and Responses to Infection in Drosophila
melanogaster. Frontiers in Immunology 10, Doi: 10.3389/fimmu.2019.03075.
[130] Duneau, D.F., Kondolf, H.C., Im, J.H., Ortiz, G.A., Chow, C., Fox, M.A., et al.,
2017. The Toll pathway underlies host sexual dimorphism in resistance to both
Gram-negative and Gram-positive bacteria in mated Drosophila. BMC Biology
15(1): 124, Doi: 10.1186/s12915-017-0466-3.
[131] Peng, Y., Lyu, J., Li, Q., Ligoxygakis, P., Xia, Y., Xiao, Q., 2025. Sexual
dimorphism in the immune response of *Drosophila melanogaster* to the
entomopathogenic fungus *Metarhizium anisopliae*. Journal of Invertebrate
Pathology 213: 108422, Doi: 10.1016/j.jip.2025.108422.
[132] Shahrestani, P., Chambers, M., Vandenberg, J., Garcia, K., Malaret, G.,
Chowdhury, P., et al., 2018. Sexual dimorphism in Drosophila melanogaster
survival of Beauveria bassiana infection depends on core immune signaling.
Scientific Reports 8(1): 12501, Doi: 10.1038/s41598-018-30527-1.
[133] Öztürk-Çolak, A., Marygold, S.J., Antonazzo, G., Attrill, H., Goutte-Gattat, D.,
Jenkins, V.K., et al., 2024. FlyBase: updates to the Drosophila genes and
genomes database. Genetics 227(1): iyad211, Doi: 10.1093/genetics/iyad211.
[134] Palm, W., Sampaio, J.L., Brankatschk, M., Carvalho, M., Mahmoud, A.,
Shevchenko, A., et al., 2012. Lipoproteins in Drosophila melanogaster—

Assembly, Function, and Influence on Tissue Lipid Composition. PLoS Genetics
8(7): e1002828, Doi: 10.1371/journal.pgen.1002828.
[135] Wobbrock, J.O., Findlater, L., Gergle, D., Higgins, J.J., 2011. The Aligned Rank
Transform for nonparametric factorial analyses using only ANOVA procedures.
CHI 2011 - 29th Annual CHI Conference on Human Factors in Computing
Systems, Conference Proceedings and Extended Abstracts: 143–6, Doi:
10.1145/1978942.1978963.

FIGURE LEGENDS

Figure 1 – Sex differences in expression of gut-derived peptide hormones.

(A-E) mRNA levels of *AstA* ($p<0.0001$; Student's *t*-test) (A), *AstC* ($p=0.0002$; Mann-

Whitney test) (B), *Tk* ($p<0.0001$; Student's *t*-test) (C), *NPF* ($p=0.0001$; Student's *t*-test)

(D), *Dh31* ($p=0.0002$; Mann-Whitney test) (E) in whole-body were significantly higher in

5-day-old w^{1118} males compared to females. $n=7-8$ biological replicates.

(F-J) mRNA levels of *AstA* ($p<0.0001$; Student's *t*-test) (F), *AstC* ($p=0.001$; Student's *t*-

test) (G), *Tk* ($p<0.0001$; Student's *t*-test) (H), *NPF* ($p=0.0015$; Student's *t*-test) (I), *Dh31*

($p<0.0001$; Student's *t*-test) (J) in heads were significantly higher in 5-day-old w^{1118}

males compared to females. $n=8-10$ biological replicates.

(K) mRNA levels of *AstA* ($p=0.5039$; Student's *t*-test) in guts were not significantly

different between 5-day-old w^{1118} females and males. $n=7$ biological replicates.

(L-N) mRNA levels of *AstC* ($p=0.0002$; Student's *t*-test) (L), *Tk* ($p<0.0001$; Student's *t*-

test) (M), *NPF* ($p=0.0006$; Mann-Whitney test) (N) in guts were significantly higher in 5-

day-old w^{1118} females compared to males. $n=7$ biological replicates.

(O) mRNA levels of *Dh31* ($p=0.7517$; Student's *t*-test) in guts were not significantly

different between 5-day-old w^{1118} females and males. $n=7$ biological replicates.

All data plotted as mean ± SEM. ns indicates not significant with $p > 0.05$; ** $p < 0.01$, ***

$p < 0.001$, **** $p < 0.0001$.

See also Figure S1.

Figure 2 – Sex determination gene *transformer* does not regulate sex differences

in EE cell-derived peptide mRNA levels.

For all data, cell type-specific RNAi was considered to have a significant effect only if

the experimental genotype (e.g., *tissue-GAL4>UAS-RNAi*) significantly differed from

both parental strains (e.g., *tissue-GAL4>+* and *+>UAS-RNAi*) with the same direction of

effect. (A-E) mRNA levels of *AstA* (A), *AstC* (B), *Tk* (C), *NPF* (D), and *Dh31* (E) in the

gut were measured in *voila-GAL4>UAS-tra^F* flies and respective genetic controls (*voila-*

GAL4>+ and *+>UAS-tra^F*) in females and males. Tra expression did not alter *AstA*

(female: $p^{GAL4} > 0.9999$ and $p^{UAS} > 0.9999$; male: $p^{GAL4} > 0.9999$ and $p^{UAS} = 0.0084$), *AstC*

(female: $p^{GAL4} = 0.6814$ and $p^{UAS} = 1.0$; male: $p^{GAL4} = 0.9965$ and $p^{UAS} = 0.0463$), *Tk* (female:

$p^{GAL4} = 0.2258$ and $p^{UAS} > 0.9999$; male: $p^{GAL4} = 0.0006$ and $p^{UAS} = 0.0004$), *NPF* (female:

$p^{GAL4} = 0.1579$ and $p^{UAS} = 0.3389$; male: $p^{GAL4} = 0.6639$ and $p^{UAS} = 0.9043$) or *Dh31* (female:

$p^{GAL4} = 0.0439$ and $p^{UAS} = 0.9745$; male: $p^{GAL4} = 0.9953$ and $p^{UAS} = 0.1370$) levels in either

sex. Sex:genotype interaction: *AstA* ($p < 0.0001$), *AstC* ($p = 0.1078$), *Tk* ($p = 0.0004$), *NPF*

($p = 0.1655$), *Dh31* ($p = 0.1945$). Data were analyzed by two-way ANOVA with Bonferroni

or Tukey's HSD post-hoc tests as appropriate (aligned rank transform applied for non-

parametric data in B and E); $n = 5$ biological replicates.

(F–J) mRNA levels of *AstA* (F), *AstC* (G), *Tk* (H), *NPF* (I), and *Dh31* (J) in the head

were measured in *voila-GAL4>UAS-tra^F* flies and respective genetic controls (*voila-*

GAL4>+ and *+>UAS-tra^F*) in females and males. Tra expression did not alter *AstA*

(female: p^{GAL4}>0.9999 and p^{UAS}=0.3344; male: p^{GAL4}>0.9999 and p^{UAS}>0.9999), *AstC*

(female: p^{GAL4}>0.9999 and p^{UAS}<0.0001; male: p^{GAL4}=0.6687 and p^{UAS}=0.0236), *NPF*

(female: p^{GAL4}>0.9999 and p^{UAS}=0.0006; male: p^{GAL4}=0.5030 and p^{UAS}=0.6158) or *Dh31*

(female: p^{GAL4}>0.9999 and p^{UAS}=0.0003; male: p^{GAL4}>0.9999 and p^{UAS}>0.9999) levels in

either sex. In females, *Tk* (p^{GAL4}>0.9999 and p^{UAS}=0.0044) levels did not alter, but in

males, *Tk* was increased compared with both controls (p^{GAL4}=0.0120 and p^{UAS}=0.0156).

Sex:genotype interaction: *AstA* (*p*=0.2471), *AstC* (*p*=0.0189), *Tk* (*p*=0.0003), *NPF*

(*p*=0.1408). *Dh31* (*p*=0.0403). Data were analyzed by two-way ANOVA with Bonferroni

post-hoc tests; *n*=8 biological replicates.

(K–O) mRNA levels of *AstA* (K), *AstC* (L), *Tk* (M), *NPF* (N), and *Dh31*(O) in the gut were

measured in *elav-GAL4>UAS-tra^F* flies and respective genetic controls (*elav-GAL4>+*

and *+>UAS-tra^F*) in females and males. Neuronal Tra expression did not alter *AstA*

(female: p^{GAL4}>0.9999 and p^{UAS}=0.0534; male: p^{GAL4}=0.5496 and p^{UAS}=0.1858), *AstC*

(female: p^{GAL4}=0.5948 and p^{UAS}=0.0878; male: p^{GAL4}=0.1745 and p^{UAS}=0.1745), *Tk*

(female: p^{GAL4}>0.9999 and p^{UAS}=0.0269; male: p^{GAL4}>0.9999 and p^{UAS}=0.2110), *NPF*

(female: p^{GAL4}>0.9999 and p^{UAS}=0.1158; male: p^{GAL4}>0.9999 and p^{UAS}=0.6652), or

Dh31 (female: p^{GAL4}=0.5442 and p^{UAS}=0.8086; male: p^{GAL4}=0.1650 and p^{UAS}=0.5258)

levels in either sex. Sex:genotype interaction: *AstA* (*p*=0.6125), *AstC* (*p*=0.4992), *Tk*

(*p*=0.5212), *NPF* (*p*=0.6546), *Dh31* (*p*=0.9566). Data were analyzed by two-way

ANOVA with Bonferroni or Tukey's HSD post-hoc tests as appropriate (aligned rank

transform applied for non-parametric data in L and O); *n*=6 biological replicates.

(P–T) mRNA levels of *AstA* (P), *AstC* (Q), *Tk* (R), *NPF* (S), and *Dh31* (T) in the head

were measured in *elav-GAL4>UAS-tra^F* flies and respective genetic controls (*elav-*

GAL4>+ and *+>UAS-tra^F*) in females and males. Neuronal Tra expression did not alter

AstA (female: p^{GAL4}=0.9843 and p^{UAS}=0.7086; male: p^{GAL4}=0.0628 and p^{UAS}=0.9936),

AstC (female: p^{GAL4}=0.1253 and p^{UAS}=0.8540; male: p^{GAL4}=0.9086 and p^{UAS}=0.0188),

Tk (female: p^{GAL4}=0.6051 and p^{UAS}=0.9999; male: p^{GAL4}=0.3600 and p^{UAS}=0.2760), or

Dh31 (female: p^{GAL4}>0.9999 and p^{UAS}>0.9999; male: p^{GAL4}=0.2918 and p^{UAS}=0.5990)

levels in either sex. *NPF* was reduced in females compared with both controls

(p^{GAL4}=0.0347 and p^{UAS}=0.0273) but was unchanged in males (p^{GAL4}=0.0656 and

p^{UAS}=0.6253). Sex:genotype interaction: *AstA* (p=0.0171), *AstC* (p=0.0198), *Tk*

(p=0.2324), *NPF* (p=0.6872), *Dh31* (p=0.4360). Data were analyzed by two-way

ANOVA with Tukey's HSD or Bonferroni post-hoc tests as appropriate (aligned rank

transform applied for non-parametric data in P, Q, R, S); *n*=5–6 biological replicates.

All data plotted as mean ± SEM. ns indicates not significant with p>0.05; * p<0.05, **

p<0.01, *** p<0.001, **** p<0.0001.

See also Figure S2.

Figure 3 – Gut-derived Tachykinin and Allatostatin C promote female fat storage.

For all data, cell type-specific RNAi was considered to have a significant effect only if

the experimental genotype (e.g., *tissue-GAL4>UAS-RNAi*) significantly differed from

both parental strains (e.g., *tissue-GAL4>+* and *+>UAS-RNAi*) with the same direction of

effect. (A) Whole-body triglyceride levels were significantly lower in *AstC-GAL4>UAS-*

AstC-RNAi,R57C10-GAL80 females compared with *AstC-GAL4>+,R57C10-GAL80 and*

+>UAS-AstC-RNAi control females (p^{GAL4}<0.0001 and p^{UAS}<0.0001), an effect that was

not observed in males (p^{GAL4}=0.4527 and p^{UAS}=0.1056) (sex:genotype interaction

$p<0.0001$). Two-way ANOVA followed by Bonferroni post-hoc test; n=8 biological

replicates.

(B) Whole-body triglyceride levels were not significantly different in *Tk-GAL4>UAS-Tk-*

RNAi,R57C10-GAL80 females and males compared with controls (female:

p^{GAL4}=0.1109 and p^{UAS}=0.0118; male: p^{GAL4}<0.0001 and p^{UAS}=0.5704) (sex:genotype

interaction $p<0.0001$) though we note a trend toward lower body fat in females. Two-

way ANOVA followed by Bonferroni post-hoc test; n=8 biological replicates.

(C) Whole-body triglyceride levels were not significantly different in *NPF-GAL4>UAS-*

NPF-RNAi,R57C10-GAL80 females and males compared with controls (female:

p^{GAL4}>0.9999 and p^{UAS}>0.9999; male: p^{GAL4}>0.9999 and p^{UAS}=0.4134) (sex:genotype

interaction p=0.2890). Two-way ANOVA followed by Bonferroni post-hoc test; n=8

biological replicates.

All data plotted as mean ± SEM. ns indicates not significant with $p>0.05$; ** $p<0.01$, ****

$p<0.0001$.

1016

Figure 4 – Allatostatin C receptor and Tachykinin receptor in neurons promote fat

storage in females but not males.

For all data, cell type-specific RNAi was considered to have a significant effect only if

the experimental genotype (e.g., *tissue-GAL4>UAS-RNAi*) significantly differed from

both parental strains (e.g., *tissue-GAL4>+* and *+>UAS-RNAi*) with the same direction of

effect. (A–C) Whole-body triglyceride levels were measured in *elav-GAL4>UAS-RNAi*

flies targeting *AstC-R2* (A), *TkR99D* (B), and *NPFR* (C), along with respective genetic

controls (*elav-GAL4>+* and *+>UAS-RNAi*) in females and males. Knockdown of *AstC-*

R2 (female: p^{GAL4}=0.0001 and p^{UAS}=0.0013; male: p^{GAL4}=0.0564 and p^{UAS}>0.9999;

sex:genotype interaction p=0.0631) and *TkR99D* (female: p^{GAL4}<0.0001 and

p^{UAS}<0.0001; male: p^{GAL4}>0.9999 and p^{UAS}>0.9999; sex:genotype interaction p<0.0001)

significantly reduced triglyceride levels in females compared with no significant

differences in males. Knockdown of *NPFR* did not change triglyceride levels in either

females or males compared with controls (female: p^{GAL4}<0.0001 and p^{UAS}>0.9999;

male: p^{GAL4}<0.0001 and p^{UAS}>0.9999; sex:genotype interaction p=0.2470). Data were

analyzed by two-way ANOVA followed by Bonferroni post-hoc tests; n=8 biological

replicates.

(D) Whole-body triglyceride levels were not significantly different in *dilp2-GAL4>UAS-*

AstC-R2-RNAi females and males compared with controls (female: p^{GAL4}<0.0001 and

p^{UAS}>0.9999; male: p^{GAL4}=0.0156 and p^{UAS}=0.3419; sex:genotype interaction

p<0.0001). Two-way ANOVA followed by Bonferroni post-hoc test; n=8 biological

replicates.

(E) Whole-body triglyceride levels were not significantly different in *dilp2-GAL4>UAS-*

TkR99D-RNAi females compared with controls (p^{GAL4}=0.0321 and p^{UAS}=0.0724).

Whole-body triglyceride levels were significantly higher in *dilp2-GAL4>UAS-TkR99D-*

RNAi males compared with *dilp2-GAL4>+* and *+>UAS-TkR99D-RNAi* control males

(p^{GAL4}<0.0001 and p^{UAS}=0.0003; sex:genotype interaction p<0.0001). Two-way ANOVA

followed by Bonferroni post-hoc test; n=8 biological replicates.

(F) Whole-body triglyceride levels were not significantly different in *Akh-GAL4>UAS-*

AstC-R2-RNAi females and males compared with controls (female: p^{GAL4}=0.3817 and

p^{UAS}=0.0181; male: p^{GAL4}=0.1229 and p^{UAS}>0.9999; sex:genotype interaction

p=0.0241). Two-way ANOVA followed by Bonferroni post-hoc test; n=8 biological

replicates.

(G) Whole-body triglyceride levels were not significantly different in *Akh-GAL4>UAS-*

TkR99D-RNAi females and males compared with controls (female: p^{GAL4}=0.4601 and

p^{UAS}>0.9999; male: p^{GAL4}>0.9999 and p^{UAS}=0.8744; sex:genotype interaction

p=0.0595). Two-way ANOVA followed by Bonferroni post-hoc test; n=8 biological

replicates.

All data plotted as mean ± SEM. ns indicates not significant with p>0.05; * p<0.05, **

p<0.01, *** p<0.001, **** p<0.0001.

1057

Figure 5 – Model of sex differences in fat storage.

Schematic representation of complex mechanisms that promote higher fat storage in

Drosophila females. Profound female-biased EE cell-derived Tk and AstC promote

female fat storage, an effect mediated by neurons but independent of the IPC or APC.

EE cell-derived factors therefore act alongside a greater insulin/insulin-like growth factor

signaling pathway activity, lower Adipokinetic hormone (Akh) signaling, and lower

expression of triglyceride lipase *brummer* (*bmm*) to specify higher adiposity in female

flies. In males, we detected no contribution of gut-derived factors to fat storage, which

we previously showed is kept at a lower level than in females by higher *bmm* expression

and Akh signaling. Thus, multiple complex mechanism specify higher fat storage in

unmated female flies compared with males. IPCs: insulin-producing cells; APCs:

adipokinetic hormone-producing cells; CNS: central nervous system; *bmm*: *brummer*;

EC: enterocyte; EB: enteroblast; EE: enteroendocrine; ISC: intestinal stem cells; AstC:

Allatostatin C; Tk: Tachykinin; Dilps: *Drosophila* insulin-like peptides; Akh: Adipokinetic

hormones. Created in BioRender. Rideout, E. (https://BioRender.com/9z1gko5) is

licensed under CC BY 4.0.

Figure 1

Figure 2: Sex determination gene transformer does not regulate sex differences in EE cell-derived peptide mRNA levels

Figure 3: Gut-derived Tachykinin and Allatostatin C promote female fat storage.

Figure 4   Allatostatin C receptor and Tachykinin receptor in neurons promote fat storage in females but not males.

Figure 5