Title

Sex-biased expression of enteroendocrine cell-derived hormones contributes to higher

fat storage in *Drosophila* females

Authors

Puja Biswas[1,2], Elizabeth J. Rideout[1,*]

Affiliations

[1] Department of Cellular and Physiological Sciences, Life Sciences Institute, The

University of British Columbia, Vancouver, Canada

[2] Present Address: Department of Pediatrics, BC Children's Hospital Research Institute,

The University of British Columbia, Vancouver, Canada

* Corresponding author: elizabeth.rideout@ubc.ca

Abstract

Enteroendocrine (EE) cells in the *Drosophila* gut produce and release multiple factors,

including Allatostatin A (AstA), Allatostatin C (AstC), neuropeptide F (NPF), tachykinin

(Tk), Diuretic hormone 31 (Dh31), Bursicon, CCHamide 1 (CCHa1) and CCHamide 2

(CCHa2), and short neuropeptide F (sNPF). Collectively, these peptides ensure that

physiology (e.g., fat storage, fluid balance) and behavior (e.g., feeding, sleep) are

coordinated with environmental factors such as nutrient quantity and quality. Despite

notable sex differences in physiology and behavior, it remains unclear whether the

regulation and function of these EE cell-derived factors is shared between males and

females. Given that recent data identified sex-biased physiological effects of two EE

cell-derived hormones on *Drosophila* food intake and energy mobilization, we performed

a detailed characterization of these hormones in male and female flies. Despite an

overall male bias in mRNA levels of *AstA*, *AstC*, *Tk*, *NPF*, *Dh31* in the whole body and

head, we observed a strong female bias in mRNA levels of *AstC*, *Tk*, and *NPF* in the

gut. To determine whether this sex-biased regulation was physiologically significant, we

monitored triglyceride levels in flies with gut-specific loss of EE cell-derived hormones.

In 5-day-old flies, loss of either EE cell-derived AstC or Tk reduced fat storage in

females with no effect in males. These female-specific effects on fat storage were

reproduced in flies with neuron-specific loss of the AstC (AstC-R2) and Tk receptors

(TkR99D). Together, these data uncover strongly sex-biased regulation of EE cell-

derived hormones, and show that gut-specific loss of two of these hormones had a

female-specific effect on body fat.

Keywords

Sex difference, gut, enteroendocrine cell, physiology, triglyceride, hormones, fat storage

Highlights:

• Enteroendocrine cell-expressed hormones show strongly sex-biased expression

• Loss of enteroendocrine cell-derived AstC and Tk reduced body fat only in females

• Neuronal loss of AstC or Tk receptors reduced stored fat only in females

1. Introduction

In *Drosophila*, females store more fat than males [1–7]. Greater female fat storage has been observed in both mated and unmated females compared with age-matched males [1,3–7]. In flies, as in other animals, triglyceride is the main form of stored fat. Although triglyceride is present in many cell types and organs (*e.g.*, oenocytes, glia, neurons, gut) [5,6,8–14], the majority of triglyceride is stored in an organ called the fat body [3,12,15].

In females, high levels of triglyceride in the fat body play a key role in supporting reproduction and physiology [5,16]. Indeed, triglyceride from the fat body is a key energy source for the developing embryo [17]. Increased fat storage in adult females also supports prolonged survival during nutrient deprivation compared with males [4,6]. Despite these clear benefits of greater fat storage for female fertility, excess fat accumulation in males adversely affects their reproductive output. Specifically, males carrying a mutation that promotes excess triglyceride accumulation show reduced testis size, defects consistent with delays in spermatogenesis, and ultimately a reduction in sperm number [8]. The sex difference in fat storage therefore likely reflects the fact that males and females differ in how much whole-body triglyceride storage supports optimal fertility.

Recent studies have identified genes and pathways that contribute to sex differences in fat storage [5–7,18]. In mated females, the steroid hormone ecdysone acts on neurons to promote food intake, which is associated with increased body fat [5]. In unmated adult females, the insulin/insulin-like growth factor signaling pathway (IIS) plays a key role in maintaining an elevated level of triglyceride storage compared with adult males [18]. Specifically, adult females have higher production of *Drosophila*

insulin-like peptide 3 (Dilp3) and greater insulin sensitivity, leading to higher IIS activity.

This elevated IIS activity is important for females to store more triglyceride than males,

as adult-specific ablation of the insulin-producing cells reduces body fat in females but

not males [18].

In males, body fat is maintained via alternative mechanisms. For example, males

show higher expression and activity of two catabolic pathways that promote fat

breakdown. One pathway is regulated by triglyceride lipase *brummer* (*bmm*), where

males show higher *bmm* mRNA levels compared with females [6]. This elevated *bmm*

expression contributes to the sex difference in fat storage by restricting triglyceride

accumulation in males. Similarly, males have higher production and secretion of

Adipokinetic hormone (Akh), a key lipolytic hormone in insects [7]. As with *bmm*, high

levels of Akh in males contribute to the sex difference in fat storage by limiting fat

accumulation in males [7]. Together, these studies have defined a model of the sex

difference in fat storage in which females maintain higher levels of fat storage due to the

anabolic action of IIS, whereas males have lower fat storage due to the catabolic effects

of *bmm* and Akh. While some progress has been made in revealing the mechanisms

underlying the sex-specific regulation of Akh and IIS [7,18], we do not have a complete

understanding of the factors that determine the sex-biased regulation and function of

these key metabolic factors.

Recent clues into the regulation of Akh, *bmm*, and IIS have emerged from

studies on peptide hormone function [19–30], as several hormones influence physiology

via effects on Akh- and Dilp-producing cells [19–22,24–33]. In particular, recent studies

have illuminated an important role for hormones produced by the enteroendocrine (EE)

cells of the gut [32,34–36]. Adult *Drosophila* EE cells produce and release hormones

such as Allatostatin A (AstA), Allatostatin C (AstC), neuropeptide F (NPF), tachykinin

(Tk), Diuretic hormone 31 (Dh31), Bursicon, CCHamide 1 (CCHa1) and CCHamide 2

(CCHa2), and short neuropeptide F (sNPF) [37–41].

EE cells are identified by expression of the homeodomain protein Prospero in

adults [42,43], where EE cells that produce distinct hormones are present in

anatomically defined regions of the adult gut [43]. For example, AstA- and Dh31-

producing EE cells are located in the posterior midgut, whereas EE cells that produce

AstC and Tk are found along the entire length of the midgut [37]. NPF-producing cells

are found in the anterior and middle midgut [41]. Supporting a role for EE cell-derived

hormones in regulating Akh/IIS, in fed conditions studies show EE cell-derived

hormones such as Bursicon inhibit Akh secretion [28], whereas NPF enhances Dilp

secretion from the insulin-producing cells [29]. During starvation, AstC promotes Akh

release from Akh-producing cells to enable lipid mobilization [22]. While at least two EE

cell-derived hormones have been shown to have sex-biased effects on food intake and

energy mobilization [22,44], sex differences in the regulation and function of most of

these hormones remain unclear. Defining potential differences in EE cells is an

important task, as prior studies have revealed profound differences in gut biology

between males and females.

For example, males and females differ in overall gut size and shape [45,46] and

in the number of intestinal stem cell divisions [45,47–49]. The absorptive lining of the

gut also shows sex differences during aging [48,50]. After mating, females show

pronounced changes to gut size, function, and gene expression [45,51–54]. While sex

differences in intestinal stem cells and enterocytes play a key role in mediating these

differences in gut biology, we know less about male-female differences in EE cells. We

therefore aimed to perform a detailed characterization of EE cell-derived hormones in

males and females, and to determine the contribution of these hormones to physiology

in each sex. We reveal profound sex-biased regulation of EE cell-expressed hormones

AstC, Tk, and NPF: females show higher mRNA levels of these hormones in the gut

than males. For at least two EE cell-derived hormones this sex-biased regulation was

physiologically significant, as loss of AstC and Tk in the gut reduced fat storage in

females with no effect in males. These female-specific fat storage defects were

reproduced by loss of AstC and Tk receptors in neurons and/or neuropeptide-producing

cells. While the specific cell type targeted by these EE cell-derived hormones to

influence fat storage remains unclear, this reveals a female-specific contribution of EE

cell-derived hormones in regulating body fat.

2. RESULTS

2.1. Sex differences in expression of gut-derived peptide hormones

Given that several EE cell-derived hormones such as AstA (FBgn0015591), AstC

(FBgn0032336), Tk (FBgn0037976), NPF (FBgn0027109), Dh31 (FBgn0032048) are

expressed in cells outside the gut [20,55–69], we used quantitative real-time PCR

(qPCR) to assess whole-body mRNA levels of genes encoding EE cell-expressed

hormones. In particular, we focused on hormones known to influence whole-body fat

metabolism [20,22,29,44,70,71]. In 5-day-old w^{1118} unmated adult males and females,

we found that whole-body mRNA levels of *AstA*, *AstC*, *Tk*, *NPF*, and *Dh31* were

significantly higher in males than in females (Figure 1A-E). To gain further insight into

this sex-biased expression, we analyzed mRNA levels of these factors from isolated

heads and intestines, as these are the main sites of AstA, AstC, Tk, NPF, and Dh31

production [35,37]. A significant male bias in mRNA levels was found in the head for

AstA, *AstC*, *Tk*, *NPF,* and *Dh31* (Figure 1F-1J). In contrast, mRNA levels of *AstC*, *Tk*,

and *NPF* in isolated intestines showed a strong female bias (Figure 1K-1M). No sex

bias in the expression of *AstA* or *Dh31* was observed in the gut (Figure 1N, 1O).

Building on the sex bias in mRNA levels, we next examined mRNA levels of

receptors that correspond to EE cell-expressed hormones with sex-biased expression in

whole-body, fat body, and head samples. Whole-body mRNA levels of the receptors for

AstA (*AstA-R2*), AstC (*AstC-R2*), Tk (*TkR99D*), NPF (*NPFR*), and Dh31 (*Dh31-R*) were

significantly higher in 5-day-old w^{1118} males compared with age-matched females

(Figure S1A-E). For most peptides, the male bias was due to a higher mRNA level in

the head and not the fat body (Figure S1A-E); however, *TkR99D* mRNA levels were

higher in male fat bodies with no difference in head mRNA levels (Figure S1C). Taken

together with our data on peptide mRNA levels, our data suggest sex differences exist

in both the expression of EE cell-derived hormones and in the ability of tissues to

respond to available peptide.

2.2. Sex determination gene *transformer* does not regulate sex differences in EE

cell-derived peptide mRNA levels

To determine the mechanism by which these differences in mRNA levels are

established, we tested a role for sex determination gene *transformer* (*tra*). Normally, a

functional Tra protein is only expressed in females, where Tra specifies most aspects of

female sexual development and behavior [72–75]. Indeed, ectopic Tra expression in

males is sufficient to specify many aspects of female sexual development and

physiology [7,47,48,73,76,77]. Because *tra* mRNA is detected in the gut, specifically in

ISC and EE cells [47,78], we asked whether broad overexpression of Tra in neurons

and/or EE cells contributes to the sex difference in mRNA levels of EE cell-expressed

hormones. We found that sex differences in mRNA levels of *AstA*, *AstC*, *Tk*, *NPF,* and

Dh31 were unaffected when we used either *voila-GAL4* (Figure 2A-2J) or *elav-GAL4*

(Figure 2K-2T) to drive Tra expression in these cells. Tra expression in neurons

similarly had no effect on mRNA levels of *AstC-R2*, *TkR99D*, or *Dh31-R* in the head

(Figure S2A-C). Thus, sex determination *tra* does not establish sex differences in levels

of the mRNAs that encode either EE cell-derived hormones or their receptors.

2.3. Gut-derived Tachykinin and Allatostatin C promote female fat storage

Given that EE cell-derived AstC, NPF, and Tk regulate fat storage and phenotypes

associated with fat storage (e.g., starvation resistance) in single- and mixed-sex animal

groups [22,29,44,71], we wanted to assess whether these hormones contribute to the

sex difference in fat storage. We used RNAi to knock down levels of *AstC, Tk* and *NPF*

with GAL4 drivers targeting these specific EE populations (*AstC-GAL4, Tk-GAL4,* and

NPF-GAL4, respectively). Importantly, GAL4 activity for each driver line was restricted

to the gut using *R57C10-GAL80*, a validated approach to target only the gut cells that

produce these hormones [22,44,71]. We found gut-specific loss of *AstC* (genotype

AstC-GAL4>UAS-AstC-RNAi, R57C10-GAL80) caused a significant reduction in body

fat in females with no effect in males (Figure 3A). Gut-specific knockdown of *Tk*

(genotype *Tk-GAL4>UAS-Tk-RNAi, R57C10-GAL80*) similarly showed a trend toward a

female-specific decrease in fat storage (GAL4 control p=0.1109; UAS control

p=0.0118), with no significant effect on male body fat (Figure 3B). In contrast, gut-

specific loss of *NPF* (*NPF-GAL4>UAS-NPF-RNAi, R57C10-GAL80*) did not significantly

alter whole-body fat storage in either males or females (Figure 3C). Together, these

data suggest that the female-biased expression of AstC and Tk in the gut play

physiologically significant roles in regulating fat storage in females.

2.4. Allatostatin C receptor and Tachykinin receptor in neurons promote fat

storage in females but not males

Neurons and neuropeptide-producing cells are key cell types upon which AstC, Tk, and

NPF act to influence physiology [22,29,32,71,79,80]. We therefore predicted that loss of

AstC-R2 and *TkR99D* in these cells would reproduce the reduced fat storage we

observed in females with loss of EE cell-derived AstC and Tk. To test this, we used

elav-GAL4 to knock down *AstC-R2* and *TkR99D* in post-mitotic neurons and

neuropeptide-producing cells. In females, loss of *AstC-R2* in neurons and neuropeptide-

producing cells caused a significant decrease in body fat, with no effect in males (Figure

4A). This reproduced the body fat phenotype caused by loss of gut AstC. A similar

female-specific reduction in fat storage was observed with loss of *TkR99D* in neurons

and neuropeptide-producing cells (Figure 4B), reproducing the fat storage phenotype of

females with loss of gut-derived Tk. In line with the lack of body fat effect due to loss of

gut-derived NPF, we saw no significant change in fat storage in either males or females

with loss of *NPFR* in neurons and neuropeptide-producing cells (Figure 4C). Together,

these data suggest that AstC and Tk may promote whole-body fat storage in females

via effects on neurons and neuropeptide-producing cells.

To narrow down the neurons and neuropeptide-producing cells in which *AstC-R2*

and *TkR99D* act to mediate their effects on female fat storage, we used cell-type-

specific GAL4 drivers to overexpress RNAi transgenes directed at these genes. Given

that these gut-derived peptides have been shown to influence metabolic homeostasis

and feeding via effects on the insulin-producing cells (IPCs) and the Akh-producing cells

(APCs) [19,22,67], we first knocked down *AstC-R2* and *TkR99D* in these cells. We used

dilp2-GAL4 to drive expression of *UAS-Astc-R2-RNAi* and *UAS-TkR99D-RNAi* in the

IPCs, and *Akh-GAL4* to drive expression of these transgenes in the APCs. Loss of

AstC-R2 in the IPC had no significant effect on fat storage in either males or females

compared with sex-matched controls (Figure 4D). IPC-specific loss of *TkR99D*, on the

other hand, caused a significant increase in whole-body fat storage in males (Figure 4E)

with no change in females. In the APC, loss of neither receptor altered fat storage in

males or females (Figure 4F, 4G). Thus, while Tk normally restricts male body fat via

effects in the IPC, AstC and Tk promote female body fat via neurons other than the IPC

and APC.

3. DISCUSSION

EE cell-derived hormones regulate body fat in single- and mixed-sex animal groups;

however, it has been unclear whether the regulation and function of these peptides

differ between the sexes. The goal of our study was to perform a detailed comparison of

EE cell-derived hormones between the sexes, and to test if these hormones contribute

to the sex difference in fat storage. Our assessment revealed profound female-biased

expression of EE cell-expressed hormones within the gut. This differential expression

was physiologically significant, as we showed that EE cell-derived Tk and AstC promote

female fat storage. Taken together, our data provide additional insight into the

mechanism(s) by which unmated female flies achieve higher fat storage than male flies.

While it was not the main goal of our study, our survey of EE cell-expressed

hormones in *Drosophila* revealed that the sex bias in expression was not uniform across

tissues. In the gut, mRNA levels of *AstC*, *Tk*, and *NPF* were higher in females than in

males. In the brain, mRNA levels of these hormones showed a significant male bias, in

line with data from previous reports on Tk [56] and NPF [63]. While it remains unclear

whether the tissue-specific sex bias in expression is physiologically significant, peptides

derived from the gut and the brain have been shown to mediate distinct effects on

physiology and/or behavior. For example, EE cell-derived AstC regulates energy

homeostasis and food-seeking behaviors in adult females [22], whereas neuron-derived

AstC is involved in regulating locomotion in adult males [59] and the circadian regulation

of oogenesis in adult females [69]. Neuron-derived Tk similarly regulates locomotion

[64,68], food consumption [67], Dilp secretion [19], and aggression [56], whereas gut-

derived Tk regulates intestinal lipogenesis [71] and stem cell divisions in the midgut

[81–83]. Supporting a potential sex-specific role for peptides derived from different

anatomical sites in regulating physiology, gut-derived AstC stimulates fat breakdown

during starvation through the Akh pathway in mated females with no effect in males

[22]. Future studies are therefore needed to determine whether there are sex

differences in whether the effects of EE cell-derived hormones are primarily mediated

by local or systemic mechanisms.

Another important task for future studies will be to elucidate how sex differences

in neuropeptide expression are established. The first step in understanding these

mechanisms will be to determine which factors specify the sex bias in neuropeptide

mRNA levels. Because our data shows that sex determination gene *tra* does not

regulate the sex bias in neuropeptide expression in either the brain or the gut, the role

of other factors that influence sexual identity and sexual differentiation must be

assessed. One strong candidate is the steroid hormone ecdysone, as virgin females

have higher ecdysone titers than males [84–86]. Ecdysone plays a role in regulating

sexual differentiation and development [5,87,88], and contributes to male-female

differences in multiple aspects of intestinal physiology (e.g., intestinal stem cell

proliferation) [45,49] and brain development [89,90]. Another candidate is juvenile

hormone, which has been shown to regulate sexual maturation in *Drosophila* and other

insects [91–97]. While it remains unclear whether juvenile hormone titers differ between

virgin males and females, juvenile hormone regulates many aspects of gut physiology in

mated females (e.g., intestinal lipid accumulation, ISC proliferation) [53,98] and

influences brain development [99]. Other than hormones, it is possible that sex

determination gene *Sex-lethal* plays a role in regulating the sex difference in mRNA

levels of EE cell-derived hormones, as *tra*-independent effects of *Sex-lethal* have been

described in the brain [100].

In parallel to identifying the factor(s) responsible for establishing the sex

difference in EE cell-expressed hormones, it will be important to reveal the cellular basis

for this differential expression. For example, a sex difference in the number of EE cells

and neuropeptide-expressing cells in the brain could explain the differences in

expression. Supporting this, gut length, overall brain size, and neuron number have

been shown to differ between males and females [34,47,101–108]. In the gut, the

difference in length is at least partially due to a sex difference in the proliferation of

intestinal stem cells [45,47,48], which undergo asymmetric divisions and subsequent

differentiation to generate all gut cell types including EE cells [34,42,109–113].

In the brain, males and females differ in the number of neurons found within

many identified clusters [101,102,104–108,114], including the cells that produce NPF

[63] and Tk [56]. Differences in neuron number have been primarily attributed to sex-

specific programmed cell death [107,114–117]; however, sex differences in neuroblast

cell death and/or proliferation may also play a role [106,118–120].

Beyond the effects of cell number, sex differences in EE cell-derived hormone

mRNA levels may also be due to differential gene and/or protein expression of these

factors, which have been reported for other peptide hormones [7,18,63]. Because sex

differences in the activity of peptide hormone-producing cells have also been previously

described [7], it is clear that a detailed examination of sex differences in EE cells, and

more generally in neuropeptide-producing cells, is needed to gain a comprehensive

picture of how these cells differ between males and females. Benefits of such a detailed

study include gaining insight into potential mechanisms underlying sex differences in

other aspects of physiology and behavior. For example, EE cells regulate ISC

homeostasis [81,121], and EE cell-derived hormones act locally and systemically to

regulate appetite, food ingestion, food digestion, gut motility, and immune responses

[35,80,122,123]. Importantly, male-female differences in many of these phenotypes

have been reported [51,52,124–127].

Overall, our findings identify EE cell-derived hormones AstC and Tk as important

factors that promote higher fat storage in *Drosophila* adult virgin females. This builds on

a recent paper identifying a key role for IIS in promoting higher levels of fat storage in

unmated females but not males [18], advancing knowledge of the factors that establish

an optimal level of stored fat in each sex.

4. MATERIALS AND METHODS

4.1.   Fly strains

The following fly strains from the Bloomington *Drosophila* Stock Center were used: *w*^{1118}

(#3605), *voila-GAL4* (#80572), *dilp2-GAL4* (IPCs) (#37516), *UAS-tra*^{F} (#4590), *UAS-*

NPF-RNAi (#27237), *UAS-TkR99D-RNAi* (#27513), *UAS-AstC-RNAi* (#25868), *UAS-*

NPFR-RNAi (#25939), *UAS-AstC-R2-RNAi* (#36888), *UAS-Tk-RNAi* (#25800), *elav-*

GAL4 (#458). We obtained *R57C10-GAL80; NPF-GAL4, R57C10-GAL80; Tk-GAL4 and*

R57C10-GAL80; AstC-GAL4 as kind gifts from Dr. Kim Rewitz at the University of

Copenhagen, and *Akh-GAL4* was a kind gift from Dr. Mike Gordon at The University of

British Columbia.

4.2.   Fly husbandry

Fly media was prepared with the following ingredients: 20.5 g/L sucrose, 70.9 g/L D-

glucose, 48.5 g/L cornmeal, 45.3 g/L yeast, 4.55 g/L agar, 0.5 g/L CaCl$_2$•2H$_2$O, 0.5 g/L

MgSO$_4$•7H$_2$O, 11.77 mL/L acid mix (propionic acid/phosphoric acid). For all

experiments, we allowed female flies to lay eggs on grape juice agar plates for 12 hr. At

24 hr after egg laying, 50 larvae were picked into vials containing 10 mL of food and

reared at 22°C. Males and females were distinguished by the presence of sex combs in

the late pupal period and placed into single-sex vials to eclose. After eclosion, adult flies

were maintained at a density of twenty flies per vial in single-sex groups. Unless

otherwise stated, all experiments used 5- to 7-day-old unmated flies.

4.3.   Adult weight

Groups of 10 flies were placed in pre-weighed 1.5 ml microcentrifuge tubes (Diamed

Lab Supplies, DIATEC610-2550) and weighed on an analytical balance (Mettler-Toledo,

ME104).

4.4.   Whole-body triglyceride measurements

One biological replicate consisted of five flies. Flies were collected in a 1.5 ml tube and

homogenized in 350 μl of 0.1% Tween (Amresco, 0777-1L) in 1X phosphate-buffered

saline (PBS; Sigma-Aldrich, P5493) using 50 μl of glass beads (Sigma-Aldrich,

Z250473) that were agitated at 8 m/s for 5 s (OMNI International BeadRuptor 24).

Triglyceride concentration was measured using the Stanbio Triglyceride Liquid Reagent

(FT7610, BD386a/d, BD386b) according to the manufacturer's instructions and as

described previously [13,18] with minor modifications. Briefly, 10 μl of either

homogenate or triglyceride standard (FT7610) was added to 190 μl of activated

triglyceride reagent (Enzymatic Triglyceride Reagent, BD386a/d; Triglyceride Activator,

Cat. No. BD386b) in a 96-well plate. After a 15 min incubation at room temperature, the

absorbance was read at 540 nm (Thermo Scientific – Multiskan FC Microplate

Photometer).

4.5. RNA extraction, cDNA synthesis, and Quantitative real-time PCR (qPCR)

One biological replicate consisted of 3-5 adult fly guts, or 10 adult fly heads, or 5 whole-

body adult flies. Samples were homogenized in 500 µl Trizol (Thermo Fisher Scientific;

15596018). Chloroform was added to Trizol to separate the mixture into aqueous and

organic phases, and isopropanol was added to the aqueous phase (in a fresh tube) to

precipitate the RNA. RNA was resuspended in either 20-25 µl (for guts and heads) or

200 µl (for whole-body) of molecular biology grade water (Corning, 46-000-CV). RNA

was stored at -80°C until use. Each experiment contained 5-10 biological replicates per

sex and per genotype; each experiment was repeated twice.

For genomic DNA elimination and cDNA synthesis, an equal amount of RNA per

reaction was DNase-treated and reverse transcribed according to manufacturer's

instructions using the QuantiTect Reverse Transcription Kit (Qiagen, 205314). Relative

mRNA transcript levels were quantified using qPCR as described previously [6]. Data

were normalized to the average fold change of *Actin5C* and *β-tubulin*. For a full primer

list, refer to Document S1.

4.6. Statistical analysis

Statistical analyses and data presentation were completed using GraphPad Prism 9

(GraphPad Software, San Diego, CA, USA). All data were tested for normality using the

Shapiro-Wilk test. Normally-distributed data were subjected to parametric tests as

appropriate, including Student's *t*-test and two-way ANOVA followed by Bonferroni post-

hoc test. For non-normally distributed data, we used the Mann-Whitney test. For two-

way ANOVA involving data that do not satisfy the normality assumption, aligned rank

transformation was first applied using the art() function from the ARTool R package

[128]. Then, ANOVA was performed on the transformed data with the base R anova()

function. Finally, the art.con() function from the ARTool package was used to extract the

main as well as the interaction effects. Default parameters were used in each step of

the analysis. For all statistical analyses, differences were considered significant if

$p < 0.05$.

CRediT authorship contribution statement

Puja Biswas: Conceptualization, Data curation, Formal analysis, Investigation,

Methodology, Visualization, Writing – original draft.

Elizabeth J. Rideout: Conceptualization, Funding acquisition, Methodology,

Resources, Supervision, Writing – original draft, Writing – review & editing.

Funding

This study was supported by operating grants to EJR from the Canadian Institutes for

Health Research (PJT-153072 and PJT-183786), CIHR Sex and Gender Science Chair

program (GS4-171365), Michael Smith Foundation for Health Research (16876), and

the Canada Foundation for Innovation (JELF-34879). PB was supported by a 4-year

CELL fellowship from UBC.

Acknowledgments

The authors thank FlyBase, which is supported by a grant from the National Human

Genome Research Institute at the U.S. National Institutes of Health (U41 HG000739)

and by the British Medical Research Council (MR/N030117/1). Stocks obtained from the

Bloomington *Drosophila* Stock Center (NIHP40OD018537) were used in this study. The

authors thank the TRiP at Harvard Medical School (NIH/NIGMS R01-GM084947) for

providing transgenic RNAi fly stocks and/or plasmid vectors used in this study. The

funders had no role in study design, data collection and analysis, decision to publish, or

preparation of the manuscript. We thank members of the Rideout lab for valuable

feedback. We acknowledge that our research takes place on the traditional, ancestral,

and unceded territory of the Musqueam people; a privilege for which we are grateful.

Declaration of competing interests

The authors declare no competing interests.

Appendix A. Supplementary data

Document S1. Supplemental Figures S1-S4

Data availability

Table S1. Excel file containing raw data with calculations

Table S2. Excel file containing statistics for all data

References

[1]   De Groef, S., Wilms, T., Balmand, S., Calevro, F., Callaerts, P., 2021. Sexual Dimorphism in Metabolic Responses to Western Diet in Drosophila melanogaster. Biomolecules 12(1): 33, Doi: 10.3390/biom12010033.

[2]   Lease, H.M., Wolf, B.O., 2011. Lipid content of terrestrial arthropods in relation to body size, phylogeny, ontogeny and sex. Physiological Entomology 36(1): 29–38, Doi: 10.1111/j.1365-3032.2010.00767.x.

[3]   Parisi, M., Li, R., Oliver, B., 2011. Lipid profiles of female and male Drosophila. BMC Research Notes 4(1): 198, Doi: 10.1186/1756-0500-4-198.

[4]   Schwasinger-Schmidt, T.E., Kachman, S.D., Harshman, L.G., 2012. Evolution of starvation resistance in Drosophila melanogaster: measurement of direct and correlated responses to artificial selection. Journal of Evolutionary Biology 25(2): 378–87, Doi: 10.1111/j.1420-9101.2011.02428.x.

[5]   Sieber, M.H., Spradling, A.C., 2015. Steroid Signaling Establishes a Female Metabolic State and Regulates SREBP to Control Oocyte Lipid Accumulation. Current Biology: CB 25(8): 993–1004, Doi: 10.1016/j.cub.2015.02.019.

[6]   Wat, L.W., Chao, C., Bartlett, R., Buchanan, J.L., Millington, J.W., Chih, H.J., et al., 2020. A role for triglyceride lipase brummer in the regulation of sex differences in Drosophila fat storage and breakdown. PLoS Biology 18(1): e3000595, Doi: 10.1371/journal.pbio.3000595.

[7]   Wat, L.W., Chowdhury, Z.S., Millington, J.W., Biswas, P., Rideout, E.J., 2021. Sex determination gene transformer regulates the male-female difference in Drosophila fat storage via the adipokinetic hormone pathway. eLife 10: e72350, Doi: 10.7554/eLife.72350.

[8]   Chao, C.F., Pesch, Y.-Y., Yu, H., Wang, C., Aristizabal, M.J., Huan, T., et al., 2024. An important role for triglyceride in regulating spermatogenesis. eLife 12, Doi: 10.7554/eLife.87523.3.

[9]   Gutierrez, E., Wiggins, D., Fielding, B., Gould, A.P., 2007. Specialized hepatocyte-like cells regulate Drosophila lipid metabolism. Nature 445(7125): 275–80, Doi: 10.1038/nature05382.

[10]   Kis, V., Barti, B., Lippai, M., Sass, M., 2015. Specialized Cortex Glial Cells Accumulate Lipid Droplets in Drosophila melanogaster. PloS One 10(7): e0131250, Doi: 10.1371/journal.pone.0131250.

[11]   Kühnlein, R.P., 2011. The contribution of the Drosophila model to lipid droplet research. Progress in Lipid Research 50(4): 348–56, Doi: 10.1016/j.plipres.2011.04.001.

[12]   Kühnlein, R.P., 2012. Thematic review series: Lipid droplet synthesis and metabolism: from yeast to man. Lipid droplet-based storage fat metabolism in Drosophila. Journal of Lipid Research 53(8): 1430–6, Doi: 10.1194/jlr.R024299.

[13]   Sieber, M.H., Thummel, C.S., 2009. The DHR96 nuclear receptor controls triacylglycerol homeostasis in Drosophila. Cell Metabolism 10(6): 481–90, Doi: 10.1016/j.cmet.2009.10.010.

[14]   Villanueva, J.E., Livelo, C., Trujillo, A.S., Chandran, S., Woodworth, B., Andrade, L., et al., 2019. Time-restricted feeding restores muscle function in Drosophila

models of obesity and circadian-rhythm disruption. Nature Communications 10(1):
2700, Doi: 10.1038/s41467-019-10563-9.

[15] Arrese, E.L., Soulages, J.L., 2010. INSECT FAT BODY: ENERGY, METABOLISM,
AND REGULATION. Annual Review of Entomology 55: 207–25, Doi:
10.1146/annurev-ento-112408-085356.

[16] Buszczak, M., Lu, X., Segraves, W.A., Chang, T.Y., Cooley, L., 2002. Mutations in
the midway gene disrupt a Drosophila acyl coenzyme A: diacylglycerol
acyltransferase. Genetics 160(4): 1511–8, Doi: 10.1093/genetics/160.4.1511.

[17] Matsuoka, S., Armstrong, A.R., Sampson, L.L., Laws, K.M., Drummond-Barbosa,
D., 2017. Adipocyte Metabolic Pathways Regulated by Diet Control the Female
Germline Stem Cell Lineage in Drosophila melanogaster. Genetics 206(2): 953–71,
Doi: 10.1534/genetics.117.201921.

[18] Biswas, P., Bako, J.A., Liston, J.B., Yu, H., Wat, L.W., Miller, C.J., et al., 2025.
Insulin/insulin-like growth factor signaling pathway promotes higher fat storage in
Drosophila females. Cell Reports 44(7), Doi: 10.1016/j.celrep.2025.115915.

[19] Birse, R.T., Söderberg, J.A.E., Luo, J., Winther, A.M.E., Nässel, D.R., 2011.
Regulation of insulin-producing cells in the adult Drosophila brain via the tachykinin
peptide receptor DTKR. The Journal of Experimental Biology 214(Pt 24): 4201–8,
Doi: 10.1242/jeb.062091.

[20] Hentze, J.L., Carlsson, M.A., Kondo, S., Nässel, D.R., Rewitz, K.F., 2015. The
Neuropeptide Allatostatin A Regulates Metabolism and Feeding Decisions in
Drosophila. Scientific Reports 5: 11680, Doi: 10.1038/srep11680.

[21] Kapan, N., Lushchak, O.V., Luo, J., Nässel, D.R., 2012. Identified peptidergic
neurons in the Drosophila brain regulate insulin-producing cells, stress responses
and metabolism by coexpressed short neuropeptide F and corazonin. Cellular and
Molecular Life Sciences 69(23): 4051–66, Doi: 10.1007/s00018-012-1097-z.

[22] Kubrak, O., Koyama, T., Ahrentløv, N., Jensen, L., Malita, A., Naseem, M.T., et al.,
2022. The gut hormone Allatostatin C/Somatostatin regulates food intake and
metabolic homeostasis under nutrient stress. Nature Communications 13(1): 692,
Doi: 10.1038/s41467-022-28268-x.

[23] Kubrak, O., Jørgensen, A.F., Koyama, T., Lassen, M., Nagy, S., Hald, J., et al.,
2024. LGR signaling mediates muscle-adipose tissue crosstalk and protects
against diet-induced insulin resistance. Nature Communications 15(1): 6126, Doi:
10.1038/s41467-024-50468-w.

[24] Kubrak, O.I., Lushchak, O.V., Zandawala, M., Nässel, D.R., 2016. Systemic
corazonin signalling modulates stress responses and metabolism in Drosophila.
Open Biology 6(11): 160152, Doi: 10.1098/rsob.160152.

[25] Nässel, D.R., Enell, L.E., Santos, J.G., Wegener, C., Johard, H.A., 2008. A large
population of diverse neurons in the Drosophilacentral nervous system expresses
short neuropeptide F, suggesting multiple distributed peptide functions. BMC
Neuroscience 9(1): 90, Doi: 10.1186/1471-2202-9-90.

[26] Oh, Y., Lai, J.S.-Y., Mills, H.J., Erdjument-Bromage, H., Giammarinaro, B.,
Saadipour, K., et al., 2019. A glucose-sensing neuron pair regulates insulin and
glucagon in Drosophila. Nature 574(7779): 559–64, Doi: 10.1038/s41586-019-
1675-4.

[27] Sano, H., Nakamura, A., Texada, M.J., Truman, J.W., Ishimoto, H., Kamikouchi, A.,
et al., 2015. The Nutrient-Responsive Hormone CCHamide-2 Controls Growth by
Regulating Insulin-like Peptides in the Brain of Drosophila melanogaster. PLOS
Genetics 11(5): e1005209, Doi: 10.1371/journal.pgen.1005209.
[28] Scopelliti, A., Bauer, C., Yu, Y., Zhang, T., Kruspig, B., Murphy, D.J., et al., 2019.
A Neuronal Relay Mediates a Nutrient Responsive Gut/Fat Body Axis Regulating
Energy Homeostasis in Adult Drosophila. Cell Metabolism 29(2): 269-284.e10, Doi:
10.1016/j.cmet.2018.09.021.
[29] Yoshinari, Y., Kosakamoto, H., Kamiyama, T., Hoshino, R., Matsuoka, R., Kondo,
S., et al., 2021. The sugar-responsive enteroendocrine neuropeptide F regulates
lipid metabolism through glucagon-like and insulin-like hormones in Drosophila
melanogaster. Nature Communications 12(1): 4818, Doi: 10.1038/s41467-021-
25146-w.
[30] Zandawala, M., Yurgel, M.E., Texada, M.J., Liao, S., Rewitz, K.F., Keene, A.C., et
al., 2018. Modulation of Drosophila post-feeding physiology and behavior by the
neuropeptide leucokinin. PLOS Genetics 14(11): e1007767, Doi:
10.1371/journal.pgen.1007767.
[31] Nässel, D.R., Broeck, J.V., 2016. Insulin/IGF signaling in Drosophila and other
insects: factors that regulate production, release and post-release action of the
insulin-like peptides. Cellular and Molecular Life Sciences 73(2): 271–90, Doi:
10.1007/s00018-015-2063-3.
[32] Nässel, D.R., Zandawala, M., 2019. Recent advances in neuropeptide signaling in
Drosophila, from genes to physiology and behavior. Progress in Neurobiology 179:
101607, Doi: 10.1016/j.pneurobio.2019.02.003.
[33] Toprak, U., 2020. The Role of Peptide Hormones in Insect Lipid Metabolism.
Frontiers in Physiology 11, Doi: 10.3389/fphys.2020.00434.
[34] Miguel-Aliaga, I., Jasper, H., Lemaitre, B., 2018. Anatomy and Physiology of the
Digestive Tract of Drosophila melanogaster. Genetics 210(2): 357–96, Doi:
10.1534/genetics.118.300224.
[35] Wegener, C., Veenstra, J.A., 2015. Chemical identity, function and regulation of
enteroendocrine peptides in insects. Current Opinion in Insect Science 11: 8–13,
Doi: 10.1016/j.cois.2015.07.003.
[36] Zhou, X., Ding, G., Li, J., Xiang, X., Rushworth, E., Song, W., 2020. Physiological
and Pathological Regulation of Peripheral Metabolism by Gut-Peptide Hormones in
Drosophila. Frontiers in Physiology 11: 577717, Doi: 10.3389/fphys.2020.577717.
[37] Chen, J., Kim, S., Kwon, J.Y., 2016. A Systematic Analysis of Drosophila
Regulatory Peptide Expression in Enteroendocrine Cells. Molecules and Cells
39(4): 358–66, Doi: 10.14348/molcells.2016.0014.
[38] Hung, R.-J., Hu, Y., Kirchner, R., Liu, Y., Xu, C., Comjean, A., et al., 2020. A cell
atlas of the adult Drosophila midgut. Proceedings of the National Academy of
Sciences of the United States of America 117(3): 1514–23, Doi:
10.1073/pnas.1916820117.
[39] Reiher, W., Shirras, C., Kahnt, J., Baumeister, S., Isaac, R.E., Wegener, C., 2011.
Peptidomics and peptide hormone processing in the Drosophila midgut. Journal of
Proteome Research 10(4): 1881–92, Doi: 10.1021/pr101116g.

[40] Veenstra, J.A., 2009. Peptidergic paracrine and endocrine cells in the midgut of the
fruit fly maggot. Cell and Tissue Research 336(2): 309–23, Doi: 10.1007/s00441-
009-0769-y.
[41] Veenstra, J.A., Agricola, H.-J., Sellami, A., 2008. Regulatory peptides in fruit fly
midgut. Cell and Tissue Research 334(3): 499–516, Doi: 10.1007/s00441-008-
0708-3.
[42] Micchelli, C.A., Perrimon, N., 2006. Evidence that stem cells reside in the adult
Drosophila midgut epithelium. Nature 439(7075): 475–9, Doi:
10.1038/nature04371.
[43] Ohlstein, B., Spradling, A., 2006. The adult Drosophila posterior midgut is
maintained by pluripotent stem cells. Nature 439(7075): 470–4, Doi:
10.1038/nature04333.
[44] Malita, A., Kubrak, O., Koyama, T., Ahrentløv, N., Texada, M.J., Nagy, S., et al.,
2022. A gut-derived hormone suppresses sugar appetite and regulates food choice
in Drosophila. Nature Metabolism 4(11): 1532–50, Doi: 10.1038/s42255-022-
00672-z.
[45] Ahmed, S.M.H., Maldera, J.A., Krunic, D., Paiva-Silva, G.O., Pénalva, C., Teleman,
A.A., et al., 2020. Fitness trade-offs incurred by ovary-to-gut steroid signalling in
Drosophila. Nature 584(7821): 415–9, Doi: 10.1038/s41586-020-2462-y.
[46] Blackie, L., Gaspar, P., Mosleh, S., Lushchak, O., Kong, L., Jin, Y., et al., 2024. The
sex of organ geometry. Nature 630(8016): 392–400, Doi: 10.1038/s41586-024-
07463-4.
[47] Hudry, B., Khadayate, S., Miguel-Aliaga, I., 2016. The sexual identity of adult
intestinal stem cells controls organ size and plasticity. Nature 530(7590): 344–8,
Doi: 10.1038/nature16953.
[48] Regan, J.C., Khericha, M., Dobson, A.J., Bolukbasi, E., Rattanavirotkul, N.,
Partridge, L., 2016. Sex difference in pathology of the ageing gut mediates the
greater response of female lifespan to dietary restriction. eLife 5: e10956, Doi:
10.7554/eLife.10956.
[49] Zipper, L., Jassmann, D., Burgmer, S., Görlich, B., Reiff, T., 2020. Ecdysone steroid
hormone remote controls intestinal stem cell fate decisions via the PPARγ-homolog
Eip75B in Drosophila. eLife 9: e55795, Doi: 10.7554/eLife.55795.
[50] Regan, J.C., Lu, Y.-X., Ureña, E., Meilenbrock, R.L., Catterson, J.H., Kißler, D., et
al., 2022. Sexual identity of enterocytes regulates autophagy to determine intestinal
health, lifespan and responses to rapamycin. Nature Aging 2(12): 1145–58, Doi:
10.1038/s43587-022-00308-7.
[51] Cognigni, P., Bailey, A.P., Miguel-Aliaga, I., 2011. Enteric neurons and systemic
signals couple nutritional and reproductive status with intestinal homeostasis. Cell
Metabolism 13(1): 92–104, Doi: 10.1016/j.cmet.2010.12.010.
[52] Hadjieconomou, D., King, G., Gaspar, P., Mineo, A., Blackie, L., Ameku, T., et al.,
2020. Enteric neurons increase maternal food intake during reproduction. Nature
587(7834): 455–9, Doi: 10.1038/s41586-020-2866-8.
[53] Reiff, T., Jacobson, J., Cognigni, P., Antonello, Z., Ballesta, E., Tan, K.J., et al.,
2015. Endocrine remodelling of the adult intestine sustains reproduction in
Drosophila. eLife 4: e06930, Doi: 10.7554/eLife.06930.

[54] White, M.A., Bonfini, A., Wolfner, M.F., Buchon, N., 2021. Drosophila melanogaster
sex peptide regulates mated female midgut morphology and physiology.
Proceedings of the National Academy of Sciences 118(1): e2018112118, Doi:
10.1073/pnas.2018112118.
[55] Abruzzi, K.C., Zadina, A., Luo, W., Wiyanto, E., Rahman, R., Guo, F., et al., 2017.
RNA-seq analysis of Drosophila clock and non-clock neurons reveals neuron-
specific cycling and novel candidate neuropeptides. PLOS Genetics 13(2):
e1006613, Doi: 10.1371/journal.pgen.1006613.
[56] Asahina, K., Watanabe, K., Duistermars, B.J., Hoopfer, E., González, C.R.,
Eyjólfsdóttir, E.A., et al., 2014. Tachykinin-Expressing Neurons Control Male-
Specific Aggressive Arousal in Drosophila. Cell 156(1): 221–35, Doi:
10.1016/j.cell.2013.11.045.
[57] Beshel, J., Dubnau, J., Zhong, Y., 2017. A Leptin analog locally produced in the
brain acts via a conserved neural circuit to modulate obesity-linked behaviors in
Drosophila. Cell Metabolism 25(1): 208, Doi: 10.1016/j.cmet.2016.12.013.
[58] Chung, B.Y., Ro, J., Hutter, S.A., Miller, K.M., Guduguntla, L.S., Kondo, S., et al.,
2017. Drosophila Neuropeptide F Signaling Independently Regulates Feeding and
Sleep-Wake Behavior. Cell Reports 19(12): 2441–50, Doi:
10.1016/j.celrep.2017.05.085.
[59] Díaz, M.M., Schlichting, M., Abruzzi, K.C., Long, X., Rosbash, M., 2019.
Allatostatin-C/AstC-R2 Is a Novel Pathway to Modulate the Circadian Activity
Pattern in Drosophila. Current Biology: CB 29(1): 13-22.e3, Doi:
10.1016/j.cub.2018.11.005.
[60] Hergarden, A.C., Tayler, T.D., Anderson, D.J., 2012. Allatostatin-A neurons inhibit
feeding behavior in adult Drosophila. Proceedings of the National Academy of
Sciences 109(10): 3967–72, Doi: 10.1073/pnas.1200778109.
[61] Kunst, M., Hughes, M.E., Raccuglia, D., Felix, M., Li, M., Barnett, G., et al., 2014.
Calcitonin gene-related peptide neurons mediate sleep-specific circadian output in
Drosophila. Current Biology: CB 24(22): 2652–64, Doi: 10.1016/j.cub.2014.09.077.
[62] Kurogi, Y., Imura, E., Mizuno, Y., Hoshino, R., Nouzova, M., Matsuyama, S., et al.,
2023. Female reproductive dormancy in Drosophila is regulated by DH31-producing
neurons projecting into the corpus allatum. Development (Cambridge, England)
150(10): dev201186, Doi: 10.1242/dev.201186.
[63] Lee, G., Bahn, J.H., Park, J.H., 2006. Sex- and clock-controlled expression of the
neuropeptide F gene in Drosophila. Proceedings of the National Academy of
Sciences 103(33): 12580–5, Doi: 10.1073/pnas.0601171103.
[64] Lee, S.H., Cho, E., Yoon, S.-E., Kim, Y., Kim, E.Y., 2021. Metabolic control of daily
locomotor activity mediated by tachykinin in Drosophila. Communications Biology
4(1): 1–14, Doi: 10.1038/s42003-021-02219-6.
[65] Luan, H., Lemon, W.C., Peabody, N.C., Pohl, J.B., Zelensky, P.K., Wang, D., et al.,
2006. Functional Dissection of a Neuronal Network Required for Cuticle Tanning
and Wing Expansion in Drosophila. Journal of Neuroscience 26(2): 573–84, Doi:
10.1523/JNEUROSCI.3916-05.2006.
[66] Mendive, F.M., Van Loy, T., Claeysen, S., Poels, J., Williamson, M., Hauser, F., et
al., 2005. Drosophila molting neurohormone bursicon is a heterodimer and the

natural agonist of the orphan receptor DLGR2. FEBS Letters 579(10): 2171–6, Doi:
10.1016/j.febslet.2005.03.006.

[67] Qi, W., Wang, G., Wang, L., 2021. A novel satiety sensor detects circulating glucose
and suppresses food consumption via insulin-producing cells in Drosophila. Cell
Research 31(5): 580–8, Doi: 10.1038/s41422-020-00449-7.

[68] Winther, Å.M.E., Acebes, A., Ferrús, A., 2006. Tachykinin-related peptides
modulate odor perception and locomotor activity in *Drosophila*. Molecular and
Cellular Neuroscience 31(3): 399–406, Doi: 10.1016/j.mcn.2005.10.010.

[69] Zhang, C., Daubnerova, I., Jang, Y.-H., Kondo, S., Žitňan, D., Kim, Y.-J., 2021. The
neuropeptide allatostatin C from clock-associated DN1p neurons generates the
circadian rhythm for oogenesis. Proceedings of the National Academy of Sciences
118(4): e2016878118, Doi: 10.1073/pnas.2016878118.

[70] Song, T., Qin, W., Lai, Z., Li, H., Li, D., Wang, B., et al., 2023. Dietary cysteine
drives body fat loss via FMRFamide signaling in Drosophila and mouse. Cell
Research 33(6): 434–47, Doi: 10.1038/s41422-023-00800-8.

[71] Song, W., Veenstra, J.A., Perrimon, N., 2014. Control of lipid metabolism by
tachykinin in Drosophila. Cell Reports 9(1): 40–7, Doi:
10.1016/j.celrep.2014.08.060.

[72] Camara, N., Whitworth, C., Van Doren, M., 2008. The creation of sexual
dimorphism in the Drosophila soma. Current Topics in Developmental Biology 83:
65–107, Doi: 10.1016/S0070-2153(08)00403-1.

[73] McKeown, M., Belote, J.M., Boggs, R.T., 1988. Ectopic expression of the female
transformer gene product leads to female differentiation of chromosomally male
Drosophila. Cell 53(6): 887–95, Doi: 10.1016/s0092-8674(88)90369-8.

[74] Oliver, B., 2002. Genetic control of germline sexual dimorphism in Drosophila.
International Review of Cytology 219: 1–60, Doi: 10.1016/s0074-7696(02)19010-3.

[75] Sturtevant, A.H., 1945. A Gene in Drosophila Melanogaster That Transforms
Females into Males. Genetics 30(3): 297–9, Doi: 10.1093/genetics/30.3.297.

[76] Billeter, J.-C., Rideout, E.J., Dornan, A.J., Goodwin, S.F., 2006. Control of male
sexual behavior in Drosophila by the sex determination pathway. Current Biology:
CB 16(17): R766-776, Doi: 10.1016/j.cub.2006.08.025.

[77] Rideout, E.J., Narsaiya, M.S., Grewal, S.S., 2015. The Sex Determination Gene
transformer Regulates Male-Female Differences in Drosophila Body Size. PLoS
Genetics 11(12): e1005683, Doi: 10.1371/journal.pgen.1005683.

[78] Hérault, C., Pihl, T., Hudry, B., 2024. Cellular sex throughout the organism underlies
somatic sexual differentiation. Nature Communications 15(1): 6925, Doi:
10.1038/s41467-024-51228-6.

[79] Chopra, G., Kaushik, S., Kain, P., 2022. Nutrient Sensing via Gut in Drosophila
melanogaster. International Journal of Molecular Sciences 23(5): 2694, Doi:
10.3390/ijms23052694.

[80] Guo, X., Lv, J., Xi, R., 2022. The specification and function of enteroendocrine cells
in Drosophila and mammals: a comparative review. The FEBS Journal 289(16):
4773–96, Doi: 10.1111/febs.16067.

[81] Amcheslavsky, A., Song, W., Li, Q., Nie, Y., Bragatto, I., Ferrandon, D., et al., 2014.
Enteroendocrine cells support intestinal stem-cell-mediated homeostasis in
Drosophila. Cell Reports 9(1): 32–9, Doi: 10.1016/j.celrep.2014.08.052.

[82] Foronda, D., Weng, R., Verma, P., Chen, Y.-W., Cohen, S.M., 2014. Coordination
of insulin and Notch pathway activities by microRNA miR-305 mediates adaptive
homeostasis in the intestinal stem cells of the Drosophila gut. Genes &
Development 28(21): 2421–31, Doi: 10.1101/gad.241588.114.

[83] O'Brien, L.E., Soliman, S.S., Li, X., Bilder, D., 2011. Altered Modes of Stem Cell
Division Drive Adaptive Intestinal Growth. Cell 147(3): 603–14, Doi:
10.1016/j.cell.2011.08.048.

[84] Bownes, M., Dübendorfer, A., Smith, T., 1984. Ecdysteroids in adult males and
females of *Drosophila melanogaster*. Journal of Insect Physiology 30(10): 823–30,
Doi: 10.1016/0022-1910(84)90019-2.

[85] Harshman, L.G., Loeb, A.M., Johnson, B.A., 1999. Ecdysteroid titers in mated and
unmated *Drosophila melanogaster* females. Journal of Insect Physiology 45(6):
571–7, Doi: 10.1016/S0022-1910(99)00038-4.

[86] Schwedes, C.C., Carney, G.E., 2012. Ecdysone signaling in adult *Drosophila*
melanogaster. Journal of Insect Physiology 58(3): 293–302, Doi:
10.1016/j.jinsphys.2012.01.013.

[87] Li, Y., Ma, Q., Cherry, C.M., Matunis, E.L., 2014. Steroid signaling promotes stem
cell maintenance in the *Drosophila* testis. Developmental Biology 394(1): 129–41,
Doi: 10.1016/j.ydbio.2014.07.016.

[88] Riddiford, L.M., Cherbas, P., Truman, J.W., 2000. Ecdysone receptors and their
biological actions. Vitamins and Hormones 60: 1–73, Doi: 10.1016/s0083-
6729(00)60016-x.

[89] Dalton, J.E., Lebo, M.S., Sanders, L.E., Sun, F., Arbeitman, M.N., 2009. Ecdysone
receptor acts in fruitless- expressing neurons to mediate drosophila courtship
behaviors. Current Biology: CB 19(17): 1447–52, Doi: 10.1016/j.cub.2009.06.063.

[90] Zhang, B., Sato, K., Yamamoto, D., 2018. Ecdysone signaling regulates
specification of neurons with a male-specific neurite in Drosophila. Biology Open
7(2): bio029744, Doi: 10.1242/bio.029744.

[91] Argue, K.J., Yun, A.J., Neckameyer, W.S., 2013. Early manipulation of juvenile
hormone has sexually dimorphic effects on mature adult behavior in Drosophila
melanogaster. Hormones and Behavior 64(4): 589–97, Doi:
10.1016/j.yhbeh.2013.08.018.

[92] Barth, R.H., Lester, L.J., 1973. Neuro-hormonal control of sexual behavior in
insects. Annual Review of Entomology 18: 445–72, Doi:
10.1146/annurev.en.18.010173.002305.

[93] Bilen, J., Atallah, J., Azanchi, R., Levine, J.D., Riddiford, L.M., 2013. Regulation of
onset of female mating and sex pheromone production by juvenile hormone in
Drosophila melanogaster. Proceedings of the National Academy of Sciences
110(45): 18321–6, Doi: 10.1073/pnas.1318119110.

[94] Flatt, T., Tu, M.-P., Tatar, M., 2005. Hormonal pleiotropy and the juvenile hormone
regulation of Drosophila development and life history. BioEssays: News and
Reviews in Molecular, Cellular and Developmental Biology 27(10): 999–1010, Doi:
10.1002/bies.20290.

[95] Raikhel, A.S., Brown, M.R., Belles, X., 2005. 3.9 - Hormonal Control of
Reproductive Processes. In: Gilbert, L.I., editor. Comprehensive Molecular Insect
Science, Amsterdam: Elsevier p. 433–91.

[96] Wijesekera, T.P., Saurabh, S., Dauwalder, B., 2016. Juvenile Hormone Is Required in Adult Males for Drosophila Courtship. PloS One 11(3): e0151912, Doi: 10.1371/journal.pone.0151912.

[97] Wyatt, G.R., Davey, K.G., 1996. Cellular and Molecular Actions of Juvenile Hormone. II. Roles of Juvenile Hormone in Adult Insects. In: Evans, P.D., editor. Advances in Insect Physiology, vol. 26. Academic Press p. 1–155.

[98] Rahman, M.M., Franch-Marro, X., Maestro, J.L., Martin, D., Casali, A., 2017. Local Juvenile Hormone activity regulates gut homeostasis and tumor growth in adult Drosophila. Scientific Reports 7(1): 11677, Doi: 10.1038/s41598-017-11199-9.

[99] Liu, Z.-H., Zhai, Y., Xia, Y., Liao, Q., 2024. Mating modifies oxidative stress in the brain and confers protection against Parkinson's Disease in a *Drosophila* model. Biochemical and Biophysical Research Communications 737: 150911, Doi: 10.1016/j.bbrc.2024.150911.

[100] Evans, D.S., Cline, T.W., 2013. Drosophila switch gene Sex-lethal can bypass its switch-gene target transformer to regulate aspects of female behavior. Proceedings of the National Academy of Sciences of the United States of America 110(47): E4474-4481, Doi: 10.1073/pnas.1319063110.

[101] Asahina, K., 2018. Sex differences in Drosophila behavior: Qualitative and Quantitative Dimorphism. Current Opinion in Physiology 6: 35–45, Doi: 10.1016/j.cophys.2018.04.004.

[102] Cachero, S., Ostrovsky, A.D., Yu, J.Y., Dickson, B.J., Jefferis, G.S.X.E., 2010. Sexual Dimorphism in the Fly Brain. Current Biology 20(18): 1589–601, Doi: 10.1016/j.cub.2010.07.045.

[103] Jiao, W., Spreemann, G., Ruchti, E., Banerjee, S., Vernon, S., Shi, Y., et al., 2022. Intact Drosophila central nervous system cellular quantitation reveals sexual dimorphism. eLife 11: e74968, Doi: 10.7554/eLife.74968.

[104] Kimura, K.-I., Ote, M., Tazawa, T., Yamamoto, D., 2005. Fruitless specifies sexually dimorphic neural circuitry in the Drosophila brain. Nature 438(7065): 229–33, Doi: 10.1038/nature04229.

[105] Nojima, T., Rings, A., Allen, A.M., Otto, N., Verschut, T.A., Billeter, J.-C., et al., 2021. A sex-specific switch between visual and olfactory inputs underlies adaptive sex differences in behavior. Current Biology 31(6): 1175-1191.e6, Doi: 10.1016/j.cub.2020.12.047.

[106] Ren, Q., Awasaki, T., Huang, Y.-F., Liu, Z., Lee, T., 2016. Cell Class-Lineage Analysis Reveals Sexually Dimorphic Lineage Compositions in the Drosophila Brain. Current Biology 26(19): 2583–93, Doi: 10.1016/j.cub.2016.07.086.

[107] Sanders, L.E., Arbeitman, M.N., 2008. Doublesex establishes sexual dimorphism in the *Drosophila* central nervous system in an isoform-dependent manner by directing cell number. Developmental Biology 320(2): 378–90, Doi: 10.1016/j.ydbio.2008.05.543.

[108] Sato, K., Yamamoto, D., 2023. Molecular and cellular origins of behavioral sex differences: a tiny little fly tells a lot. Frontiers in Molecular Neuroscience 16: 1284367, Doi: 10.3389/fnmol.2023.1284367.

[109] Buchon, N., Osman, D., 2015. All for one and one for all: Regionalization of the *Drosophila* intestine. Insect Biochemistry and Molecular Biology 67: 2–8, Doi: 10.1016/j.ibmb.2015.05.015.

[110] Driver, I., Ohlstein, B., 2014. Specification of regional intestinal stem cell identity
during Drosophila metamorphosis. Development 141(9): 1848–56, Doi:
10.1242/dev.104018.
[111] Jiang, H., Edgar, B.A., 2011. Intestinal stem cells in the adult *Drosophila* midgut.
Experimental Cell Research 317(19): 2780–8, Doi: 10.1016/j.yexcr.2011.07.020.
[112] Lemaitre, B., Miguel-Aliaga, I., 2013. The digestive tract of Drosophila
melanogaster. Annual Review of Genetics 47: 377–404, Doi: 10.1146/annurev-
genet-111212-133343.
[113] Schonhoff, S.E., Giel-Moloney, M., Leiter, A.B., 2004. Minireview: Development and
Differentiation of Gut Endocrine Cells. Endocrinology 145(6): 2639–44, Doi:
10.1210/en.2004-0051.
[114] Kimura, K.-I., 2011. Role of cell death in the formation of sexual dimorphism in the
Drosophila central nervous system. Development, Growth & Differentiation 53(2):
236–44, Doi: 10.1111/j.1440-169X.2010.01223.x.
[115] Garner, S.R.C., Castellanos, M.C., Baillie, K.E., Lian, T., Allan, D.W., 2018.
Drosophila female-specific Ilp7 motoneurons are generated by Fruitless-dependent
cell death in males and by a double-assurance survival role for Transformer in
females. Development (Cambridge, England) 145(1): dev150821, Doi:
10.1242/dev.150821.
[116] Kimura, K., Hachiya, T., Koganezawa, M., Tazawa, T., Yamamoto, D., 2008.
Fruitless and Doublesex Coordinate to Generate Male-Specific Neurons that Can
Initiate Courtship. Neuron 59(5): 759–69, Doi: 10.1016/j.neuron.2008.06.007.
[117] Nojima, T., Kimura, K., Koganezawa, M., Yamamoto, D., 2010. Neuronal synaptic
outputs determine the sexual fate of postsynaptic targets. Current Biology: CB
20(9): 836–40, Doi: 10.1016/j.cub.2010.02.064.
[118] Ghosh, N., Bakshi, A., Khandelwal, R., Rajan, S.G., Joshi, R., 2019. The Hox gene
Abdominal-B uses DoublesexF as a cofactor to promote neuroblast apoptosis in
the Drosophila central nervous system. Development (Cambridge, England)
146(16): dev175158, Doi: 10.1242/dev.175158.
[119] Taylor, B.J., Truman, J.W., 1992. Commitment of abdominal neuroblasts in
Drosophila to a male or female fate is dependent on genes of the sex-determining
hierarchy. Development (Cambridge, England) 114(3): 625–42, Doi:
10.1242/dev.114.3.625.
[120] Truman, J.W., Bate, M., 1988. Spatial and temporal patterns of neurogenesis in the
central nervous system of Drosophila melanogaster. Developmental Biology
125(1): 145–57, Doi: 10.1016/0012-1606(88)90067-x.
[121] Scopelliti, A., Cordero, J.B., Diao, F., Strathdee, K., White, B.H., Sansom, O.J., et
al., 2014. Local Control of Intestinal Stem Cell Homeostasis by Enteroendocrine
Cells in the Adult Drosophila Midgut. Current Biology 24(11): 1199–211, Doi:
10.1016/j.cub.2014.04.007.
[122] Kamareddine, L., Robins, W.P., Berkey, C.D., Mekalanos, J.J., Watnick, P.I., 2018.
The Drosophila immune deficiency pathway modulates enteroendocrine function
and host metabolism. Cell Metabolism 28(3): 449-462.e5, Doi:
10.1016/j.cmet.2018.05.026.

[123] Ren, G.R., Hauser, F., Rewitz, K.F., Kondo, S., Engelbrecht, A.F., Didriksen, A.K.,
et al., 2015. CCHamide-2 Is an Orexigenic Brain-Gut Peptide in Drosophila. PloS
One 10(7): e0133017, Doi: 10.1371/journal.pone.0133017.
[124] Belmonte, R.L., Corbally, M.-K., Duneau, D.F., Regan, J.C., 2020. Sexual
Dimorphisms in Innate Immunity and Responses to Infection in Drosophila
melanogaster. Frontiers in Immunology 10, Doi: 10.3389/fimmu.2019.03075.
[125] Duneau, D.F., Kondolf, H.C., Im, J.H., Ortiz, G.A., Chow, C., Fox, M.A., et al., 2017.
The Toll pathway underlies host sexual dimorphism in resistance to both Gram-
negative and Gram-positive bacteria in mated Drosophila. BMC Biology 15(1): 124,
Doi: 10.1186/s12915-017-0466-3.
[126] Peng, Y., Lyu, J., Li, Q., Ligoxygakis, P., Xia, Y., Xiao, Q., 2025. Sexual dimorphism
in the immune response of *Drosophila melanogaster* to the entomopathogenic
fungus *Metarhizium anisopliae*. Journal of Invertebrate Pathology 213: 108422, Doi:
10.1016/j.jip.2025.108422.
[127] Shahrestani, P., Chambers, M., Vandenberg, J., Garcia, K., Malaret, G.,
Chowdhury, P., et al., 2018. Sexual dimorphism in Drosophila melanogaster
survival of Beauveria bassiana infection depends on core immune signaling.
Scientific Reports 8(1): 12501, Doi: 10.1038/s41598-018-30527-1.
[128] Wobbrock, J.O., Findlater, L., Gergle, D., Higgins, J.J., 2011. The Aligned Rank
Transform for nonparametric factorial analyses using only ANOVA procedures. CHI
2011 - 29th Annual CHI Conference on Human Factors in Computing Systems,
Conference Proceedings and Extended Abstracts: 143–6, Doi:
10.1145/1978942.1978963.

FIGURE LEGENDS

Figure 1 – Sex differences in expression of gut-derived peptide hormones.

(A-E) mRNA levels of *AstA* ($p<0.0001$; Student's *t*-test) (A), *AstC* ($p=0.0002$; Mann-

Whitney test) (B), *Tk* ($p<0.0001$; Student's *t*-test) (C), *NPF* ($p=0.0001$; Student's *t*-test)

(D), *Dh31* ($p=0.0002$; Mann-Whitney test) (E) in whole-body were significantly higher in

5-day-old w^{1118} males compared to females. n=7-8 biological replicates.

(F-J) mRNA levels of *AstA* ($p<0.0001$; Student's *t*-test) (F), *AstC* ($p=0.001$; Student's *t*-

test) (G), *Tk* ($p<0.0001$; Student's *t*-test) (H), *NPF* ($p=0.0015$; Student's *t*-test) (I), *Dh31*

($p<0.0001$; Student's *t*-test) (J) in heads were significantly higher in 5-day-old w^{1118}

males compared to females. n=8-10 biological replicates.

(K-O) mRNA levels of *AstC* (*p*=0.0002; Student's *t*-test) (K), *Tk* (*p*<0.0001; Student's *t*-

test) (L), *NPF* (*p*=0.0006; Mann-Whitney test) (M) in guts were significantly higher in 5-

day-old w^{1118} females compared to males. n=7 biological replicates.

(N-O) mRNA levels of *AstA* (*p*=0.5039; Student's *t*-test) (N), *Dh31* (*p*=0.7517; Student's

t-test) (O) in guts were not significantly different between 5-day-old w^{1118} females and

males. n=7 biological replicates.

All data plotted as mean ± SEM. ns indicates not significant with *p*>0.05; ** *p*<0.01, ***

p<0.001, **** *p*<0.0001.

See also Figure S1.

Figure 2 – Sex determination gene *transformer* does not regulate sex differences

in EE cell-derived peptide mRNA levels.

(A) mRNA levels of *AstA* in gut were not significantly different between *voila-*

GAL4>UAS-tra^F females and *voila-GAL4>+* and *+>UAS-tra^F* control females (*p*>0.9999

and *p*>0.9999, respectively). mRNA levels of *AstA* in gut were significantly higher in

voila-GAL4>UAS-tra^F males compared to *+>UAS-tra^F* control males, but were not

significantly different from *voila-GAL4>+* control males (*p*=0.0084 and *p*>0.9999,

respectively) (sex:genotype interaction *p*<0.0001). Two-way ANOVA followed by

Bonferroni post-hoc test; n=5 biological replicates.

(B) mRNA levels of *AstC* in gut were not significantly different between *voila-*

GAL4>UAS-tra^F females and *voila-GAL4>+* and *+>UAS-tra^F* control females (*p*=0.6814

and *p*=1.0, respectively). mRNA levels of *AstC* in gut were significantly higher in *voila-*

GAL4>UAS-tra^F males compared to *+>UAS-tra^F* control males, but were not

significantly different from *voila-GAL4>+* control males (*p*=0.0463 and *p*=0.9965,

respectively) (sex:genotype interaction *p*=0.1078). Two-way ANOVA followed by Tukey

HSD on data processed using the aligned rank transform for non-parametric data; n=5

biological replicates.

(C) mRNA levels of *Tk* in gut were not significantly different between *voila-GAL4>UAS-*

tra^F females and *voila-GAL4>+* and *+>UAS-tra^F* control females (*p*=0.2258 and

p>0.9999, respectively). mRNA levels of *Tk* in gut were significantly higher in *voila-*

GAL4>UAS-tra^F males compared to *+>UAS-tra^F* control males, but were significantly

lower compared to *voila-GAL4>+* control males (*p*=0.0004 and *p*=0.0006, respectively)

(sex:genotype interaction *p*=0.0004). Two-way ANOVA followed by Bonferroni post-hoc

test; n=5 biological replicates.

(D) mRNA levels of *NPF* in gut were not significantly different between *voila-*

GAL4>UAS-tra^F females and *voila-GAL4>+* and *+>UAS-tra^F* control females (*p*=0.1579

and *p*=0.3389, respectively). mRNA levels of *NPF* in gut were not significantly different

between *voila-GAL4>UAS-tra^F* males and *voila-GAL4>+* and *+>UAS-tra^F* control males

(*p*=0.6639 and *p*=0.9043, respectively) (sex:genotype interaction *p*=0.1655). Two-way

ANOVA followed by Bonferroni post-hoc test; n=5 biological replicates.

(E) mRNA levels of *Dh31* in gut were significantly lower in *voila-GAL4>UAS-tra^F*

females compared to *voila-GAL4>+* control females, but were not significantly different

from *+>UAS-tra^F* control females (*p*=0.0439 and *p*=0.9745, respectively). mRNA levels

of *Dh31* in gut were not significantly different between *voila-GAL4>UAS-tra^F* males and

voila-GAL4>+ and *+>UAS-tra^F* control males (*p*=0.9953 and *p*=0.1370, respectively)

(sex:genotype interaction *p*=0.1945). Two-way ANOVA followed by Tukey HSD on data

processed using the aligned rank transform for non-parametric data; n=5 biological

replicates.

(F) mRNA levels of *AstA* in head were not significantly different between *voila-*

GAL4>UAS-tra^F females and *voila-GAL4>+* and *+>UAS-tra^F* control females ($p>0.9999$

and $p=0.3344$, respectively). mRNA levels of *AstA* in head were not significantly

different between *voila-GAL4>UAS-tra^F* males and *voila-GAL4>+* and *+>UAS-tra^F*

control males ($p>0.9999$ and $p>0.9999$, respectively) (sex:genotype interaction

$p=0.2471$). Two-way ANOVA followed by Bonferroni post-hoc test; n=8 biological

replicates.

(G) mRNA levels of *AstC* in head were significantly lower in *voila-GAL4>UAS-tra^F*

females compared to *+>UAS-tra^F* control females, but were not significantly different

from *voila-GAL4>+* control females ($p<0.0001$ and $p>0.9999$, respectively). mRNA

levels of *AstC* in head were significantly lower in *voila-GAL4>UAS-tra^F* males compared

to *+>UAS-tra^F* control males, but were not significantly different from *voila-GAL4>+*

control males ($p=0.0236$ and $p=0.6687$, respectively) (sex:genotype interaction

$p=0.0189$). Two-way ANOVA followed by Bonferroni post-hoc test; n=8 biological

replicates.

(H) mRNA levels of *Tk* in head were significantly lower in *voila-GAL4>UAS-tra^F* females

compared to *+>UAS-tra^F* control females, but were not significantly different from *voila-*

GAL4>+ control females ($p=0.0044$ and $p>0.9999$, respectively). mRNA levels of *Tk* in

head were significantly higher in *voila-GAL4>UAS-tra^F* males compared to both *voila-*

GAL4>+ and *+>UAS-traF* control males ($p=0.0120$ and $p=0.0156$, respectively)

(sex:genotype interaction p=0.0003). Two-way ANOVA followed by Bonferroni post-hoc

test; n=8 biological replicates.

(I) mRNA levels of *NPF* in head were significantly lower in *voila-GAL4>UAS-tra^F^*

females compared to *+>UAS-tra^F^* control females, but were not significantly different

from *voila-GAL4>+* control females (p=0.0006 and p>0.9999, respectively). mRNA

levels of *NPF* in head were not significantly different between *voila-GAL4>UAS-tra^F^*

males and *voila-GAL4>+* and *+>UAS-tra^F^* control males (p=0.5030 and p=0.6158,

respectively) (sex:genotype interaction p=0.1408). Two-way ANOVA followed by

Bonferroni post-hoc test; n=8 biological replicates.

(J) mRNA levels of *Dh31* in head were significantly lower in *voila-GAL4>UAS-tra^F^*

females compared to *+>UAS-tra^F^* control females, but were not significantly different

from *voila-GAL4>+* control females (p=0.0003 and p>0.9999, respectively). mRNA

levels of *Dh31* in head were not significantly different between *voila-GAL4>UAS-tra^F^*

males and *voila-GAL4>+* and *+>UAS-tra^F^* control males (p>0.9999 and p>0.9999,

respectively) (sex:genotype interaction p=0.0403). Two-way ANOVA followed by

Bonferroni post-hoc test; n=8 biological replicates.

(K) mRNA levels of *AstA* in gut were not significantly different between *elav-*

*GAL4>UAS-tra^F^* females and *elav-GAL4>+* and *+>UAS-tra^F^* control females (p>0.9999

and p=0.0534, respectively). mRNA levels of *AstA* in gut were not significantly different

between *elav-GAL4>UAS-tra^F^* males and *elav-GAL4>+* and *+>UAS-tra^F^* control males

(p=0.5496 and p=0.1858, respectively) (sex:genotype interaction p=0.6125). Two-way

ANOVA followed by Bonferroni post-hoc test; n=6 biological replicates.

(L) mRNA levels of *AstC* in gut were not significantly different between *elav-*

GAL4>UAS-tra^F females and *elav-GAL4>+* and *+>UAS-tra^F* control females (p=0.5948

and p=0.0878, respectively). mRNA levels of *AstC* in gut were not significantly different

between *elav-GAL4>UAS-tra^F* males and *elav-GAL4>+* and *+>UAS-tra^F* control males

(p=0.1745 and p=0.1745, respectively) (sex:genotype interaction p=0.4992). Two-way

ANOVA followed by Tukey HSD on data processed using the aligned rank transform for

non-parametric data; n=6 biological replicates.

(M) mRNA levels of *Tk* in gut were significantly lower in *elav-GAL4>UAS-tra^F* females

compared to *+>UAS-tra^F* control females, but were not significantly different from *elav-*

GAL4>+ control females (p=0.0269 and p>0.9999, respectively). mRNA levels of *Tk* in

gut were not significantly different between *elav-GAL4>UAS-tra^F* males and *elav-*

GAL4>+ and *+>UAS-tra^F* control males (p>0.9999 and p=0.2110, respectively)

(sex:genotype interaction p=0.5212). Two-way ANOVA followed by Bonferroni post-hoc

test; n=6 biological replicates.

(N) mRNA levels of *NPF* in gut were not significantly different between *elav-*

GAL4>UAS-tra^F females and *elav-GAL4>+* and *+>UAS-tra^F* control females (p>0.9999

and p=0.1158, respectively). mRNA levels of *NPF* in gut were not significantly different

between *elav-GAL4>UAS-tra^F* males and *elav-GAL4>+* and *+>UAS-tra^F* control males

(p>0.9999 and p=0.6652, respectively) (sex:genotype interaction p=0.6546). Two-way

ANOVA followed by Bonferroni post-hoc test; n=6 biological replicates.

(O) mRNA levels of *Dh31* in gut were not significantly different between *elav-*

GAL4>UAS-tra^F females and *elav-GAL4>+* and *+>UAS-tra^F* control females (p=0.5442

and p=0.8086, respectively). mRNA levels of *Dh31* in gut were not significantly different

between *elav-GAL4>UAS-tra^F* males and *elav-GAL4>+* and *+>UAS-tra^F* control males

(*p*=0.1650 and *p*=0.5258, respectively) (sex:genotype interaction *p*=0.9566). Two-way

ANOVA followed by Tukey HSD on data processed using the aligned rank transform for

non-parametric data; n=6 biological replicates.

(P) mRNA levels of *AstA* in head were not significantly different between *elav-*

GAL4>UAS-tra^F females and *elav-GAL4>+* and *+>UAS-tra^F* control females (*p*=0.9843

and *p*=0.7086, respectively). mRNA levels of *AstA* in head were not significantly

different between *elav-GAL4>UAS-tra^F* males and *elav-GAL4>+* and *+>UAS-tra^F* control

males (*p*=0.0628 and *p*=0.9936, respectively) (sex:genotype interaction *p*=0.0171).

Two-way ANOVA followed by Tukey HSD on data processed using the aligned rank

transform for non-parametric data; n=5-6 biological replicates.

(Q) mRNA levels of *AstC* in head were not significantly different between *elav-*

GAL4>UAS-tra^F females and *elav-GAL4>+* and *+>UAS-tra^F* control females (*p*=0.1253

and *p*=0.8540, respectively). mRNA levels of *AstC* in head were significantly lower in

elav-GAL4>UAS-tra^F males compared to *+>UAS-tra^F* control males, but were not

significantly different from *elav-GAL4>+* control males (*p*=0.0188 and *p*=0.9086,

respectively) (sex:genotype interaction *p*=0.0198). Two-way ANOVA followed by Tukey

HSD on data processed using the aligned rank transform for non-parametric data; n=5-6

biological replicates.

(R) mRNA levels of *Tk* in head were not significantly different between *elav-GAL4>UAS-*

tra^F females and *elav-GAL4>+* and *+>UAS-tra^F* control females (*p*=0.6051 and

p=0.9999, respectively). mRNA levels of *Tk* in head were not significantly different

between *elav-GAL4>UAS-tra^F* males and *elav-GAL4>+* and *+>UAS-tra^F* control males

(p=0.3600 and p=0.2760, respectively) (sex:genotype interaction p=0.2324). Two-way

ANOVA followed by Tukey HSD on data processed using the aligned rank transform for

non-parametric data; n=5-6 biological replicates.

(S) mRNA levels of *NPF* in head were significantly lower in *elav-GAL4>UAS-tra^F*

females compared to both *elav-GAL4>+* and *+>UAS-tra^F* control females (p=0.0347 and

p=0.0273, respectively). mRNA levels of *NPF* in head were not significantly different

between *elav-GAL4>UAS-tra^F* males and *elav-GAL4>+* and *+>UAS-tra^F* control males

(p=0.0656 and p=0.6253, respectively) (sex:genotype interaction p=0.6872). Two-way

ANOVA followed by Tukey HSD on data processed using the aligned rank transform for

non-parametric data; n=5-6 biological replicates.

(T) mRNA levels of *Dh31* in head were not significantly different between *elav-*

GAL4>UAS-tra^F females and *elav-GAL4>+* and *+>UAS-tra^F* control females (p>0.9999

and p>0.9999, respectively). mRNA levels of *Dh31* in head were not significantly

different between *elav-GAL4>UAS-tra^F* males and *elav-GAL4>+* and *+>UAS-tra^F* control

males (p=0.2918 and p=0.5990, respectively) (sex:genotype interaction p=0.4360).

Two-way ANOVA followed by Bonferroni post-hoc test; n=5-6 biological replicates.

All data plotted as mean ± SEM. ns indicates not significant with p>0.05; * p<0.05, **

p<0.01, *** p<0.001, **** p<0.0001.

See also Figure S2.

1006

Figure 3 – Gut-derived Tachykinin and Allatostatin C promote female fat storage.

(A) Whole-body triglyceride levels were significantly lower in *AstC-GAL4>UAS-AstC-*

RNAi,R57C10-GAL80 females compared to *AstC-GAL4>+,R57C10-GAL80 and*

+>UAS-AstC-RNAi control females (*p*<0.0001 and *p*<0.0001, respectively). Whole-body

triglyceride levels were not significantly different between *AstC-GAL4>UAS-AstC-*

RNAi,R57C10-GAL80 males and *AstC-GAL4>+,R57C10-GAL80* and *+>UAS-AstC-*

RNAi control males (*p*=0.4527 and *p*=0.1056, respectively) (sex:genotype interaction

p<0.0001). Two-way ANOVA followed by Bonferroni post-hoc test; n=8 biological

replicates.

(B) Whole-body triglyceride levels were significantly lower in *Tk-GAL4>UAS-Tk-*

RNAi,R57C10-GAL80 females compared to *+>UAS-Tk-RNAi* control females

(*p*=0.0118) but were not significantly different from *Tk-GAL4>+,R57C10-GAL80* control

females (*p*=0.1109). Whole-body triglyceride levels were significantly higher in *Tk-*

GAL4>UAS-Tk-RNAi,R57C10-GAL80 males compared to *Tk-GAL4>+,R57C10-GAL80*

control males (*p*<0.0001) but were not significantly different from *+>UAS-Tk-RNAi*

control males (*p*=0.5704) (sex:genotype interaction *p*<0.0001). Two-way ANOVA

followed by Bonferroni post-hoc test; n=8 biological replicates.

(C) Whole-body triglyceride levels were not significantly different between *NPF-*

GAL4>UAS-NPF-RNAi,R57C10-GAL80 females and *NPF-GAL4>+,R57C10-GAL80* and

+>UAS-NPF-RNAi control females (*p*>0.9999 and *p*>0.9999, respectively). Whole-body

triglyceride levels were not significantly different between *NPF-GAL4>UAS-NPF-*

RNAi,R57C10-GAL80 males and *NPF-GAL4>+,R57C10-GAL80* and *+>UAS-NPF-RNAi*

control males (*p*>0.9999 and *p*=0.4134, respectively) (sex:genotype interaction

p=0.2890). Two-way ANOVA followed by Bonferroni post-hoc test; n=8 biological

replicates.

All data plotted as mean ± SEM. ns indicates not significant with $p>0.05$; ** $p<0.01$, ****

$p<0.0001$.

1034

Figure 4 – Allatostatin C receptor and Tachykinin receptor in neurons promote fat

storage in females but not males.

(A) Whole-body triglyceride levels were significantly lower in *elav-GAL4>UAS-AstC-R2-*

RNAi females compared to *elav-GAL4>+* and *+>UAS-AstC-R2-RNAi* control females

($p=0.0001$ and $p=0.0013$, respectively). Whole-body triglyceride levels were not

significantly different between *elav-GAL4>UAS-AstC-R2-RNAi* males and *elav-GAL4>+*

and *+>UAS-AstC-R2-RNAi* control males ($p=0.0564$ and $p>0.9999$, respectively)

(sex:genotype interaction $p=0.0631$). Two-way ANOVA followed by Bonferroni post-hoc

test; n=8 biological replicates.

(B) Whole-body triglyceride levels were significantly lower in *elav-GAL4>UAS-TkR99D-*

RNAi females compared to *elav-GAL4>+* and *+>UAS-TkR99D-RNAi* control females

($p<0.0001$ and $p<0.0001$, respectively). Whole-body triglyceride levels were not

significantly different between *elav-GAL4>UAS-TkR99D-RNAi* males and *elav-GAL4>+*

and *+>UAS-TkR99D-RNAi* control males ($p>0.9999$ and $p>0.9999$, respectively)

(sex:genotype interaction $p<0.0001$). Two-way ANOVA followed by Bonferroni post-hoc

test; n=8 biological replicates.

(C) Whole-body triglyceride levels were significantly lower in *elav-GAL4>UAS-NPFR-*

RNAi females compared to *elav-GAL4>+* control females ($p<0.0001$) but were not

significantly different from *+>UAS-NPFR-RNAi* control females ($p>0.9999$). Whole-body

triglyceride levels were significantly lower in *elav-GAL4>UAS-NPFR-RNAi* males

compared to *elav-GAL4>+* control males ($p<0.0001$) but were not significantly different

from *+>UAS-NPFR-RNAi* control males ($p>0.9999$) (sex:genotype interaction

$p=0.2470$). Two-way ANOVA followed by Bonferroni post-hoc test; n=8 biological

replicates.

(D) Whole-body triglyceride levels were significantly lower in *dilp2-GAL4>UAS-AstC-R2-*

RNAi females compared to *dilp2-GAL4>+* control females ($p<0.0001$) but were not

significantly different from *+>UAS-AstC-R2-RNAi* control females ($p>0.9999$). Whole-

body triglyceride levels were significantly higher in *dilp2-GAL4>UAS-AstC-R2-RNAi*

males compared to *dilp2-GAL4>+* control males ($p=0.0156$) but were not significantly

different from *+>UAS-AstC-R2-RNAi* control males ($p=0.3419$) (sex:genotype interaction

$p<0.0001$). Two-way ANOVA followed by Bonferroni post-hoc test; n=8 biological

replicates.

(E) Whole-body triglyceride levels were significantly lower in *dilp2-GAL4>UAS-TkR99D-*

RNAi females compared to *dilp2-GAL4>+* control females ($p=0.0321$) but were not

significantly different from *+>UAS-TkR99D-RNAi* control females ($p=0.0724$). Whole-

body triglyceride levels were significantly higher in *dilp2-GAL4>UAS-TkR99D-RNAi*

males compared to *dilp2-GAL4>+* and *+>UAS-TkR99D-RNAi* control males ($p<0.0001$

and $p=0.0003$, respectively) (sex:genotype interaction $p<0.0001$). Two-way ANOVA

followed by Bonferroni post-hoc test; n=8 biological replicates.

(F) Whole-body triglyceride levels were not significantly different in *Akh-GAL4>UAS-*

AstC-R2-RNAi females compared to *Akh-GAL4>+* control females ($p=0.3817$) but were

significantly lower than *+>UAS-AstC-R2-RNAi* control females ($p=0.0181$). Whole-body

triglyceride levels were not significantly different in *Akh-GAL4>UAS-AstC-R2-RNAi*

males compared to *Akh-GAL4>+* and *+>UAS-AstC-R2-RNAi* control males (*p*=0.1229

and *p*>0.9999, respectively) (sex:genotype interaction *p*=0.0241). Two-way ANOVA

followed by Bonferroni post-hoc test; n=8 biological replicates.

(G) Whole-body triglyceride levels were not significantly different in *Akh-GAL4>UAS-*

TkR99D-RNAi females compared to *Akh-GAL4>+* and *+>UAS-TkR99D-RNAi* control

females (*p*=0.4601 and *p*>0.9999, respectively). Whole-body triglyceride levels were not

significantly different in *Akh-GAL4>UAS-TkR99D-RNAi* males compared to *Akh-*

GAL4>+ and *+>UAS-TkR99D-RNAi* control males (*p*>0.9999 and *p*=0.8744,

respectively) (sex:genotype interaction *p*=0.0595). Two-way ANOVA followed by

Bonferroni post-hoc test; n=8 biological replicates.

All data plotted as mean ± SEM. ns indicates not significant with *p*>0.05; * *p*<0.05, **

p<0.01, *** *p*<0.001, **** *p*<0.0001.

Figure 1

Figure 2

Figure 3

Figure 4