Polyelectrolyte domains in intrinsically disordered regions differentially regulate intracellular biomolecular condensation

Yoon Jeong Choi[1,2,3], Yujin Lee[1,6], Soojin Cho[1,6], Yaqi Pan [1,6], Changi Pack[4,5], Yemuk Kang[1], Youngtae Jeong[2,3], and Kiwon Song[1,*]

[1]Department of Biochemistry, College of Life Science and Biotechnology, Yonsei University, Seoul 03722, Republic of Korea

[2]Department of New Biology, DGIST, Daegu, 43014, Korea

[3]New Biology Research Center, DGIST, Daegu, 43014, Korea

[4]Department of Biomedical Engineering, University of Ulsan College of Medicine, Seoul 05505, South Korea

[5]Convergence Medicine Research Center, Asan Institute for Life Sciences, Asan Medical Center, Seoul 05505, South Korea

[6]These authors contributed equally.

*Correspondence: bc5012@yonsei.ac.kr

SUMMARY

Intracellular condensation is a hallmark of many proteins containing intrinsically disordered regions (IDRs). Decoding the regulatory mechanisms of condensation encoded within the amino acid sequences of condensate-forming proteins is essential for understanding both how intracellular condensates assemble and how they contribute to various disease pathologies. Here, we investigated the budding yeast P-body proteins Nst1 and Edc3, which exhibit pronounced intracellular self-condensation, and identified charged amino acid clusters in the low complexity regions (LCRs) that modulate their phase behavior. Guided by the stickers-and-spacers framework, we hypothesized that clusters of charged

residues within IDRs function in promoting intracellular condensation process. Using a sliding-window

approach, we designated and dissected the polyelectrolyte (PE) domains of Nst1 and Edc3 and

established their essential contributions to condensate coalescence. In Nst1, the polyanionic (PA)

subdomain enhances intermolecular attraction and promotes coalescence, whereas the polyampholyte

(PAM) subdomain imparts rigidity to the condensate as a distinct modulatory element and recruits

additional components into the condensate. PA of Edc3 also regulates coalescence. Extending these

findings to the neurodegenerative disease-associated protein TDP-43, we show that its PA regions are

critical for coalescence, consistent with the presence of negatively charged amino acid clusters as potent

molecular stickers functioning in condensate coalescence in cells. These results highlight charged-

residue clusters as versatile sequence determinants of phase behavior and indicate that their systematic

identification will be pivotal to defining the molecular grammar of protein condensation underlying the

condensation of disease-related proteins.

KEYWORDS

biomolecular condensates, intrinsically disordered region (IDR), stickers and spacers model,

polyelectrolyte, coalescence, polyampholyte, polyanion, P-body, Nst1, Edc3, TDP-43

INTRODUCTION

Biomolecular condensation observed in cells not only deepens our understanding of molecular dynamics

in response to various physiological conditions, but also shed light on the mechanisms underlying protein

amyloid formation (Alberti and Hyman, 2021; Alshareedah et al., 2024; Banani et al., 2017; Niu et al.,

2023). These condensations occur within the crowded and complex cellular environment, exhibiting

diverse physical properties depending on the degree of condensation.

Liquid-liquid phase separation (LLPS) represents one form of condensation, characterized by partitioned

liquid-like droplets distinct from solidified protein amyloids (Ranganathan et al., 2022).

Neurodegeneration-associated proteins usually act as scaffolds that drive the formation of liquid-like

condensates with self-condensation ability and high multivalency. Recent studies have increasingly

reported the co-occurrence of amyloid fibrils and liquid-like condensates of pathogenic proteins, such as

TDP-43, both in the brain tissues of patients with neurodegenerative diseases and in cellular models of

these diseases (Ahmad et al., 2022; Alshareedah *et al.*, 2024; Gruijs da Silva et al., 2022; Sneideriene et

al., 2025). This observation strongly suggests that the aggregation of pathogenic proteins does not

proceed through direct solid aggregation, but rather through prior enrichment into liquid-like condensates,

followed by a structural conversion into amyloid assemblies. That is, dysregulation of these liquid-like

condensates is expected to promote the formation of protein amyloids implicated in neurodegenerative

disorders (Hurtle et al., 2023; Nam and Gwon, 2023; Visser et al., 2024). Therefore, elucidating the

underlying mechanisms of liquid-like condensation is critical both for understanding the molecular

dynamics underlying physiological condensate formation, and for identifying therapeutic strategies to

regulate pathological condensate formation (Alberti and Hyman, 2021; Choi et al., 2024; Gao et al., 2023;

Lu et al., 2024; Mehta and Zhang, 2022).

Stickers-and-spacers model is a leading framework of a molecular mechanism for biomolecular liquid-

like condensation (Banjade and Rosen, 2014; Choi et al., 2020; Sood and Zhang, 2024). This model

defines "stickers" as molecular elements that mediate attractive interactions, and "spacers" as flexible

linkers that separate stickers and modulate condensate properties (Choi *et al.*, 2020). Amino acids in

intrinsically disordered regions (IDRs), capable of functioning as stickers, were precisely identified through

in vitro studies. The cation-π interaction between tyrosine and arginine, as well as the interaction between

lysine and RNA, were revealed to act as stickers (Ukmar-Godec et al., 2019; Wang et al., 2018b). The

sequential pattern of these components significantly influences the dynamic characteristics of the

condensate (Alshareedah *et al.*, 2024; Alshareedah et al., 2021; Wang *et al.*, 2018b). Notably, when

sticker amino acids are clustered within the sequence, rather than dispersed, the resultant condensate

tends towards solidification (Alshareedah *et al.*, 2024; Alshareedah *et al.*, 2021; Wang *et al.*, 2018b).

Thus, the stickers-and-spacers model offers a solid basis for elucidating the condensation characteristics

of molecules *in vitro*, offering valuable insights into the behavior of biomolecular condensates.

Despite the explanatory power of the stickers-and-spacers model, the molecular grammar underlying

intracellular biomolecular condensation remains enigmatic. First, the morphology and physical properties

of condensates depend on the characteristics of condensate-forming proteins and the intracellular

environment, and these properties can change over time as condensation progresses. In other words, the

stickers that function may vary depending on the phase of condensation. Second, although the molecular

grammar of condensation has been extensively characterized for prion-like low-complexity domains

(PLCDs) - such as poly Q/N tracts in yeast amyloidogenic proteins and the RGG motifs in disease-related

proteins like FUS and TDP-43 (Patel et al., 2015; Sanders et al., 2020; Wang et al., 2018), other

intrinsically disordered regions (IDRs) with distinct combinations of sticker amino acids remain largely

unexplored. Critically, many IDRs lack canonical PLCD features but still drive phase separation,

suggesting the existence of alternative molecular grammars.

Recent *in vitro* and simulation studies have reported that the charge distribution within amino acid

sequences strongly affects the viscosity, surface tension, and dynamic properties of condensates (Biswas

and Potoyan, 2024; Linsenmeier et al., 2023; Sundaravadivelu Devarajan et al., 2024). These reports

strongly indicate the need for research on the regulation of cellular condensation process by novel low

complexity regions (LCRs) composed of sticker amino acids such as charged amino acid clusters, in

addition to the previously studied PLCDs.

In this study, we aimed to elucidate the role of the polyelectrolyte region in the cellular condensation of

biomolecules via Nst1. Both Nst1 and Edc3, established P-body components in budding yeast, possess

intrinsic self-condensation capabilities. Upon Nst1 and Edc3 overexpression, cells initially exhibit

numerous small puncta of Nst1 and Edc3, which progressively fuse into larger structures over time,

thereby reducing the overall number of observed puncta (Fig. 1A). Utilizing this model, we examined how

the polyelectrolyte region affects both the molecular dynamics of NST1 condensates and the condensate

formation of P-body components mediated by NST1. Furthermore, we investigated whether the function

of the polyelectrolyte region in Nst1 condensation could be extrapolated to TDP-43, a stress granule

constituent that plays a pivotal role in stress granule dynamics and is closely linked to the pathology of

neurodegenerative diseases.

RESULTS

Overexpression of Nst1 and Edc3 leads to mature condensate formation, whereas overexpression

of their polyelectrolyte (PE) domain deletion mutants within the IDR maintains a stabilized

microphase-separated state.

In our previous study, we reported that GFP-tagged Nst1 and Edc3, when overexpressed under the

control of the GAL promoter, formed condensates that were larger and denser than those formed by Dcp2

or Dhh1 (Choi et al., 2022). Here, we monitored the formation dynamics of these condensates from the

early phase of Nst1 and Edc3 overexpression until their intracellular expression levels reached saturation

(Fig. 1A). Numerous small condensates—here referred to as microphase-separated structures—began to

appear simultaneously throughout the cell as early as 1 hour after galactose induction and were

observed in nearly all cells by 2 hours. When we analyze the mean integrated density and the number of

puncta of each cell at 4 hours post-induction, the number of condensates per cell had decreased, while

their internal density had increased markedly, indicative of a transition to a mature condensate state

(Fig. 1B). These observations suggest that microphase Nst1 condensates progressively undergo

coalescence as the intracellular concentration of Nst1 rises.

Next, we confirmed this distinctive self-condensation potential of Nst1 and Edc3 through analysis of

puncta formation patterns in correlation with cytoplasmic protein concentration (Fig. 1C-D). Every

molecule has its own critical concentration at which it forms a condensate, and a lower critical

concentration indicates a higher potential for molecular condensation. By measuring the cytoplasmic

concentration of proteins and correlating it with the degree of puncta formation (microphase separation,

mature condensates), we can compare condensation patterns between proteins under cellular conditions.

Using a galactose promoter, GFP-tagged P-body components such as Nst1, Edc3, Dcp2, and Dhh1,

along with the stress granule component Pab1, were overexpressed for 4 hours. Afterward, the

cytoplasmic concentration of the overexpressed proteins and the degree of self-condensation were

measured using fluorescence correlation spectroscopy (FCS) (Fig.1 A, C-D). We classified the patterns of

puncta formation into three categories: diffuse state, microphase separation, and mature condensates

(Fig. 1A). GFP control did not exhibit puncta formation at any measured cytoplasmic concentration,

whereas Nst1 and Edc3 demonstrated mature condensates at remarkably low concentrations (Fig. 1C-D).

Another P-body component, Dcp2 and Dhh1, exhibited microphase separation at higher cytoplasmic

concentrations than the concentrations required for mature condensate formation by Nst1 and Edc3.

However, mature condensates similar to those observed with Nst1 and Edc3 were not detected for Dcp2

and Dhh1 (Fig. 1C-D). These results confirm that, compared to Dcp2 and Dhh1, Nst1 and Edc3 have a

greater intrinsic propensity for self-condensation in cells, which indicates their intrinsic potential for

condensation.

Next, we hypothesized that a particular charged-amino acid cluster within intrinsically disordered regions

(IDRs) in Nst1 protein sequence may function as a "sticker" domain during condensate formation. In our

previous study, we identified that PD (polyampholyte domain), a segment of a charged-amino acid cluster

in Nst1, functions in the condensation of other P-body components in cells overexpressing Nst1 (Choi *et*

al., 2022). However, Nst1 has extended charged amino acid regions beyond PD. We systematically

searched for polyelectrolyte-like clusters of charged amino acids in the IDRs of Nst1 and Edc3 using a

sliding window approach, as described in Supplementary Figure 1E (Fig. 1E). Charged amino acids (K, R,

H, D, E) were classified, and their proportion within each window was calculated to generate a scoring

system. Subsequently, a density plot of charged amino acids was created based on their positions within

the sequence (Fig. 1E, S1, S2). Using this method, we segmented polyelectrolyte (PE) amino acid

sequence regions spanning residues 65–205 in Edc3 and 460–752 in Nst1 (Fig. 1F, S1, S2).

Then, we generated deletion mutant clones of Nst1ΔPE and Edc3ΔPE, overexpressed each protein,

and observed the cells after 4 hours of induction. Consistent with previous findings, overexpression of

Nst1 deleted with the aggregation prone domain (APD), Nst1ΔAPD, did not result in condensate

formation (Choi *et al.*, 2022) even at higher cytoplasmic concentrations (24 h induction), while those

overexpressing wild-type Nst1 and Edc3 formed mature condensates (Fig. 1G-J). Interestingly, cells

overexpressing the PE deletion mutants of both Nst1 and Edc3 exhibited microphase separation, even

when their cytoplasmic concentrations were comparable with those of wild type Nst1 and Edc3 (Fig. 1G-J).

To rule out insufficient expression or time as the cause of microphase separation, we extended the

culture to 24 hours in galactose medium. However, even after prolonged induction, both Nst1ΔPE and

Edc3ΔPE failed to form mature condensates and continued to exhibit microphase separation (Fig. 1G, H).

These results indicate that each polyelectrolyte (PE) is necessary for the coalescence of both Nst1 and

Edc3 condensates, thereby facilitating the formation of mature condensates of Nst1 and Edc3.

Polyanion (PA) domain of the Polyelectrolyte (PE) region specifically promotes the coalescence of

Nst1 condensates, whereas the polyampholyte (PAM) domain in the PE region does not.

Since the PE region of Nst1 exhibits a unique pattern of charged amino acid arrangement compared to

the PE region of Edc3, the PE region of Nst1 was further characterized. Analysis of the Nst1 PE region

using FELLS (http://protein.bio.unipd.it/fells/ BioComputing UP) revealed two distinct segments: a

polyanion (PA) region spanning residues 460–631, a cluster consisting of remarkably high proportion of

negatively charged amino acids, and a polyampholyte (PAM) domain spanning residues 632–752 (Fig. 2

A, B). In our previous study, we reported that the PAM domain (PD in the previous report) is required for

promoting the condensation of other P-body components such as Dcp2 and Dhh1, which is induced by

Nst1 condensation, but does not affect the self-condensation of Nst1 (Choi *et al.*, 2022).

To validate whether the effects of PE on Nst1 condensation were specifically due to the PA domain, we

observed the time-dependent pattern of puncta formation of each GFP-tagged Nst1 domain deletion

mutant in *Δnst1* cells. When overexpressed for 1 hour using the galactose induction system, the wild-type

Nst1 (Nst1 full-length, F.L.), Nst1ΔPA, Nst1ΔPAM, and Nst1ΔPE all formed microphase separation (Fig.

2C). After 4 hours of induction, wild-type Nst1 and Nst1ΔPAM exhibited mature condensates with a

reduced number of puncta per cell, indicating that the numerous small puncta observed after 1 hour of

induction coalesced into fewer, larger puncta as the cytoplasmic concentration increased over time. (Fig.

2C). However, cells overexpressing Nst1ΔPA and Nst1ΔPE failed to undergo coalescence, and showed

an increased number of puncta and decreased signal intensities compared to cells overexpressing

wildtype Nst1 and Nst1ΔPAM (Fig. 2C). Even after 24 hours of induction, wild-type Nst1 and Nst1ΔPAM

formed mature condensates with intense fluorescence signals (Fig. 2C). However, cells expressing

Nst1ΔPA and Nst1ΔPE did not show a reduction in puncta number, although some puncta increased in

size and intensity compared to those observed after 4 hours of induction (Fig. 2C). This result indicates

that PA domain is essential for coalescence of Nst1 condensates. Since several mutations in a protein

can alter the physical property of condensates (Banani et al., 2022; Sanfeliu-Cerdan et al., 2023), we also

determined whether these condensates exhibit LLPS or amyloid-like characteristics. Cells overexpressing

each domain deletion mutant for 4 hours were treated with 1,6-hexanediol for 1 hour, then observed.

Most puncta generated by each mutant dispersed upon treatment, indicating that none of the domains

significantly alters the liquid-like properties of these condensates (Fig. 2D).

To evaluate the condensation potential of each domain deletion mutant, we measured the protein

concentrations associated with each intracellular puncta formation pattern for each Nst1 variant after 4

hours of induction by FCS (Fig. 2E). Cells expressing GFP or Nst1ΔAPD remained diffuse, whereas cells

expressing Nst1ΔPA or Nst1ΔPE exhibited microphase separation without forming mature condensates,

despite having cytoplasmic protein concentrations comparable to those of wild-type Nst1 and Nst1ΔPAM

(Fig. 2E).

When we initially constructed the Nst1ΔPA mutant, positively charged amino acids near the C-terminal

portion of the defined PA region were included in the deletion. To better distinguish the function of

negatively charged clusters within this region, we generated an additional mutant called Nst1ΔPA-1 that

excluded positively charged amino acids from deletion while maintaining the integrity of the negatively-

charged domain within PA. Consistently, cells overexpressing Nst1ΔPA-1 displayed numerous

microphase separations from 1hour to 24 hours induction, similar to cells overexpressing Nst1ΔPA (Fig.

S4). These findings indicate that the overexpression phenotype observed in Nst1ΔPA is due to the

polyanion located within the PA region.

Next, we investigated whether the results from the Nst1 domain deletion mutants could be applied in

condensate formation of endogenous Nst1. Since endogenously expressed Nst1 forms condensates from

diauxic shift to stationary phase (Fig. 2F), we examined the puncta generation patterns by condensation

of endogenously expressed full-length Nst1, the Nst1ΔPA, the Nst1ΔPAM, and the Nst1ΔAPD from early

log phase to stationary phase (Fig. 2F).

We cultured cells expressing the chromosome-integrated GFP-tagged versions of each domain deletion

mutant under the control of the endogenous Nst1 promoter in a *Δnst1* strain and observed them at 16, 30,

48, and 80 hours after seeding at an initial OD_{600} = 0.1 (Fig. 2G). These mutants exhibited similar puncta

generation patterns to those observed in cells overexpressing these deletion mutants. Consistent with

previous reports (Choi *et al.*, 2022), Nst1ΔAPD did not form puncta, confirming that the aggregation-

prone region critical for hydrophobic core assembly is essential for condensate formation (Fig. 2H-I).

Among the tested mutants, Nst1ΔPA was the first to initiate condensation, starting at 10 hours during the

early log phase (Fig. 2H-I), and notably during the mid-log phase (14–18 hours). While the wild-type Nst1

and other mutants had not yet begun puncta formation, Nst1ΔPA condensates increased significantly

(Fig. 2H-I). After 18 hours, condensation began for the full-length Nst1 and Nst1ΔPAM. Interestingly,

starting at 22 hours, as cells entered the stationary phase, the intensity of Nst1ΔPAM condensates were

markedly higher than those of the wild-type Nst1 and Nst1ΔPA condensates (Fig. 2I). At 46 hours, during

the stationary phase, we observed that the intensities of Nst1ΔPA condensates were lower than those of

wild-type Nst1 and Nst1ΔPAM condensates (Fig. 2J). Together, these findings demonstrate that each

domain of IDR differentially contributes to Nst1 condensation even under physiological conditions.

Polyanion (PA) domain exclusively functions in coalescence of Nst1 condensates.

Next, we investigated whether the contribution of the PA domain to Nst1 condensation is unique to the

PA region, or whether other sequence regions of similar size can also play a similar role, thereby

assessing potential non-specific effects arising from deletions of comparable regions. To clarify this, we

generated five additional domain deletion mutants of similar size as the PA domain in Nst1: Nst1ΔD1[2-122],

Nst1ΔD2[123-293], Nst1ΔD3[294-459], Nst1ΔD4[753-844], Nst1ΔD5[845-1015] (Fig. 3A). The *Δnst1* w303a cells were

transformed with plasmids that can overexpress each GFP-tagged Nst1 domain deletion mutant (D1~D5).

All strains were induced with galactose and observed over time using a fluorescence microscope at

specified time points after the induction (Fig. 3B).

Except the Nst1ΔAPD deletion mutant, the overexpression of D1~D5 deletion mutants began to induce

condensates approximately 1 hour after galactose induction (Fig. 3B). Similar to the Nst1ΔPAM (Fig. 2C,

3B), the condensates of the deletion mutants from D1 to D5, increased in size as the number of

condensates per cell decreased, except for the Nst1ΔPA (Fig. 3B). When cytoplasmic protein

concentrations were measured by FCS after 4 hours of induction, Nst1ΔPA exhibited microphase

separation at all measured concentrations, whereas all other mutants formed mature condensates at

similar cytoplasmic concentrations (Fig. 3C). Even after 24 hours of induction, Nst1ΔPA failed to generate

mature condensates, while all the other mutants, ΔD1 to ΔD5, successfully formed mature condensates

(Fig. 3B). These results further support that the function of the negatively charged amino acid cluster in

the IDR of Nst1 is unique and critical for coalescence of condensates in the process of biomolecular

condensation of Nst1.

The polyanion (PA) domain of Nst1 enhances intermolecular interaction, whereas the

polyampholyte (PAM) domain of Nst1 increases the rigidity of Nst1 condensates.

Next, we examined whether the PA and PAM domains influence the dynamics of Nst1 condensates by

employing fluorescence recovery after photobleaching (FRAP). Since the condensates of the

overexpressed Nst1ΔPA did not coalescence and retained minimal puncta size, the abnormal function in

the coalescence process is likely related to the molecular dynamics affected by the PA domain deletion.

We examined GFP-tagged the full-length Nst1, as well as the Nst1ΔPA, Nst1ΔPAM, and Nst1ΔD1

mutants, with GFP control after 4 hours of induction (Fig. 4A). Puncta from each mutant were imaged,

photobleached, and fluorescence recovery was monitored over time (Fig. 4B). Prior to be analyzed,

recovery curves from each mutant FRAP analysis were fitted and normalized (Fig. 4C-D, S6 A-E).

Because molecular dynamics can be affected by molecular size (Pack, 2021), the Nst1ΔD1 mutant was

also used as a size control for the analyses of Nst1ΔPA and Nst1ΔPAM.

We measured the maximal recovery velocity and recovery half-life for each Nst1 domain deletion mutant

(Fig. 4C, F-G). The maximal recovery velocity represents the rate at which unbleached fluorescent

molecules exchange with bleached molecules within puncta; a lower maximal recovery velocity indicates

stronger intermolecular interactions within condensates, while a higher velocity suggests weaker

interactions and increased molecular dynamics. The recovery half-life reflects the time required for

fluorescence in the bleached region to reach half of its maximum equilibrium intensity, serving as an

additional measure of molecular dynamics. The full-length Nst1 exhibited the slowest and lowest maximal

recovery velocity among all samples, whereas the PAM-, PA-, and D1-deleted mutants recovered more

rapidly (Fig. 4D-F), indicating that molecules with smaller molecular weight are generally more dynamic in

the condensates. Notably, Nst1ΔPA puncta showed the fastest maximal recovery velocity among all

deletion mutants (Fig. 4F). In contrast, the maximal recovery velocity and half-life of Nst1ΔPAM were

comparable to those of Nst1ΔD1 (Fig. 4F-G). These results suggest that Nst1ΔPA exhibits particularly

increased molecular dynamics due to weaker and less stable intermolecular interactions compared to

Nst1ΔPAM and Nst1ΔD1.

Regarding mobile fraction as area under curve (AUC, %), the full-length Nst1 had the lowest AUC

compared with these Nst1 deletion mutant proteins (Fig. 4E). The AUC of the Nst1Δ PAM was

significantly higher than those of the Nst1ΔPA and Nst1ΔD1 (Fig. 4H). This indicates that puncta formed

by Nst1ΔPAM have a higher proportion of mobile fractions compared to those formed by Nst1ΔPA and

Nst1ΔD1. This suggests that the PAM domain of Nst1 enhances the rigidity of Nst1 condensates.

Overall, our results strongly suggest that polyampholyte (PAM) deletion weakens cohesion of internal

networks, increasing overall molecular mobility within the condensate, while the polyanion (PA) deletion

specifically enhances single-molecule dynamics by reducing electrostatic molecular affinity. That is,

polyanions might participate in intermolecular interactions, enforcing molecular affinity, while

polyampholytes contribute to restricting the movement of molecules in the confined state within

condensates. These results highlight how different charged domains contribute uniquely to condensate

properties and molecular behavior.

The PAM domain is an RNA-independent multivalent interaction domain for interaction with a P-body component, Dcp2.

In our previous study, we found that overexpression of Nst1 induces the formation of Nst1 condensates, which simultaneously promote the condensation of other P-body components, including Dcp2 (Choi *et al.*, 2022). We suggested that the PAM domain, previously described as PD, is a multivalent interaction region responsible for the condensation of P-body components such as Dcp2 (Choi *et al.*, 2022).

Since the function of PAM domain was resolved by measuring endogenously expressing GFP-tagged Dcp2, Dhh1, and Xrn1 in cells overexpressing non-tagged Nst1 variants (Choi *et al.*, 2022), we need to confirm the function of PAM domain using a direct approach that can elucidate its role in inducing Dcp2 condensation. We utilized the chromosomally mKate2-tagged Dcp2 strain overexpressing GFP-tagged Nst1 variants, including Nst1ΔPAM, Nst1ΔPA, and Nst1ΔD1. The cells were induced to generate condensates of each Nst1 variant by 4 hour-overexpression, then treated with cycloheximide which binds to the ribosomal A site and inhibits translation elongation, resulting in decrease of free RNA level (Schneider-Poetsch et al., 2010). The reduction of the cytoplasmic concentration of free RNA, thereby eliminates Dcp2 condensates that are formed by free RNA (Teixeira et al., 2005; Van Treeck and Parker, 2018). With this strategy, the majority of Dcp2 condensates formed by Nst1 variants could be observed under these conditions (Fig. 5A).

Using high-resolution confocal microscopy, we clearly observed the formation of mixed or multiphasic condensates composed of Nst1 and Dcp2 (Fig. 5B). Most wild-type Nst1 condensates were co-localized with mKate2-tagged Dcp2, suggesting that Nst1 condensates promote the formation of Dcp2 condensates (Fig. 5C). Interestingly, the fluorescence signals from Nst1 and Dcp2 condensates did not completely overlap; instead, the two signals appeared as if distinct condensates were docked together (Fig. 5B).

To further investigate potential intermolecular interactions between Nst1 and Dcp2, we performed fluorescence cross-correlation spectroscopy (FCCS). However, no correlation between the two signals

was detected, indicating that individual Nst1 and Dcp2 molecules do not directly interact with each other

(data not shown). These findings suggest that Dcp2 condensation induced by Nst1 overexpression is not

mediated by direct intermolecular interactions but rather occurs as a consequence of the Nst1

condensation process itself.

Both Nst1ΔPA and Nst1ΔD1 condensates also exhibited co-localization with Dcp2-mKate2 condensates

(Fig. 5B-C). Although Nst1ΔPA condensates were able to induce Dcp2 condensation, their size was

relatively smaller compared to other Nst1 variant condensates, reminding the unique role of PA in

coalescence of microphase condensates (Fig. 5B). In contrast, Nst1ΔPAM condensates rarely promoted

the formation of Dcp2 condensates, highlighting the critical role of the PAM domain in facilitating Dcp2

condensation upon Nst1 overexpression (Fig. 5B-C).

These observations suggest that if the PAM domain induces Dcp2 condensation directly, rather than via

RNA, then the number of Dcp2 condensates should not be reduced in Nst1ΔPA cells even after

cycloheximide treatment, because Nst1ΔPA condensates, which contain the PAM domain, fail to

coalesce and instead increase in number. In contrast, Nst1ΔPAM should lead to a significant reduction in

the number of Dcp2 condensates (Fig. 5D). Condensates formed by overexpression of GFP-tagged Nst1

and its mutants demonstrated no change in the integrated density of puncta before and after CHX

treatment (S7. A). This indicates that a decrease in cytoplasmic concentration of ribosomal free RNA

induced by CHX treatment does not affect Nst1 condensation.

To further verify the role of PAM, we quantitatively measured Dcp2 condensation induced by

overexpression of Nst1ΔPA and Nst1ΔPAM in each cell. For the quantitative analysis of Dcp2

condensates induced by condensation of wild-type Nst1 and its domain deletion mutants, instead of

relying on the weak signals from mKate2-tagged Dcp2, we employed a GFP-tagged Dcp2 strain that

allowed more robust quantification to further validate the function of the PAM domain (Fig. 5D). Cells

overexpressing either non-tagged full-length Nst1 or the deletion mutants (Nst1ΔPA, Nst1ΔPAM) for 4

hours were treated with cycloheximide for 10 minutes prior to imaging. Subsequently, the puncta formed

by chromosomally EGFP-tagged Dcp2 in these overexpression conditions were segmented and

quantified (Fig. 5D-E). Notably, cells overexpressing Nst1ΔPA exhibited a substantially higher number of

Dcp2 puncta per cell compared to cells overexpressing wild-type Nst1 (Fig. 5E), suggesting that the

increased number of Nst1ΔPA condensates containing the PAM domain may promote the formation of a

greater number of Dcp2 condensates than wild-type Nst1. In contrast, the number of Dcp2 puncta per cell

was reduced in Nst1ΔPAM-overexpressing cells relative to the wild-type (Fig. 5E), indicating that the PAM

domain provides a physical platform necessary for efficient Dcp2 condensation. Collectively, these results

demonstrate that the patterns of Dcp2 puncta formation differ between the PA and PAM deletion mutants,

as depicted in Fig. 5D.

To further elucidate how the PA and PAM domains of Nst1 influence condensate formation in relation to

free-RNA, we analyzed GFP-Dcp2, GFP-Edc3, and GFP-Nst1 condensates in lysates derived from

stationary-phase cells endogenously expressing each of these GFP-tagged proteins (Fig. 5F).

Observations were conducted both before and after the addition of an RNase inhibitor, which prevents

RNA degradation and increases the concentration of pre-initiation RNA complexes (Fig. 5F). Previous

studies have demonstrated that P-bodies extracted from Xrn1-deleted cells disassemble upon RNase

treatment *in vitro* (Teixeira *et al.*, 2005), suggesting that RNA serves as a scaffold for P-body formation.

Based on this, we hypothesized that the condensation of specific P-body components would either

increase or decrease depending on their reliance on RNA as a scaffold. If the condensation of a particular

P-body component requires pre-initiation complex RNA, its condensation is expected to be promoted

when the concentration of pre-initiation complex RNA increases. Conversely, if condensation is

independent of pre-initiation complex RNA, an increase in pre-initiation complex RNA concentration

would either reduce condensation or leave it unaffected.

As previously reported, we observed that puncta of Dcp2, which lacks self-condensation potential,

became intensified after RNase inhibitor treatment, supporting free RNA dependency in Dcp2 condensate

formation (Fig. 5G-H, and(Cohen et al., 2005; Tibble et al., 2021; Van Treeck and Parker, 2018). In

contrast, we observed that Edc3 condensates decreased upon RNase inhibitor treatment (Fig. 5G-H).

Since both Edc3 and pre-initiation complex RNA can function as scaffolds for P-body generation, this

observation likely reflects the competition between Edc3 and RNA for inducing condensation.

To investigate the role of RNA in condensate formation mediated by Nst1 and its domains, we analyzed

the puncta of endogenously expressed GFP-Nst1, GFP-Nst1ΔPA, and GFP-Nst1ΔPAM in stationary-

phase cells, with or without RNase inhibitor treatment. Similar to Edc3, Nst1 condensates were reduced

in cells treated with the RNase inhibitor (Fig. 5I-J). The condensates formed by the Nst1 ΔPA mutant

displayed comparable intensity regardless of RNase inhibitor treatment (Fig. 5I-J). In contrast, the Nst1

ΔPAM mutant exhibited a significant increase in condensate intensity following RNase inhibitor treatment,

suggesting an enhanced dependency of Nst1ΔPAM on pre-initiation RNA complexes (Fig. 5I-J). These

findings strongly support that self-condensation of both Nst1 and Edc3 occurs independently of pre-

initiation RNA complexes. Furthermore, the PAM domain appears to serve as an interaction platform that

promotes RNA-independent condensation of other P-body components.

The Polyanion (PA) domain of human TDP-43 *promotes* coalescence of condensates, similar to

Nst1.

We expanded the functional analysis of PA and PAM subdomains to the scaffold protein Edc3 focusing

on their roles in biomolecular condensation. Since Edc3 also has a distinctive negative charged amino

acid cluster in PE domain, we divided the PE domain of Edc3 into two parts: the PA region, characterized

by a cluster of negatively charged amino acids, and the PAM region, where positively and negatively

charged amino acids are more evenly distributed (Fig. S2, S8). We constructed plasmids for the

overexpression of GFP-tagged Edc3ΔPAM and Edc3ΔPA mutants, and overexpressed them together

with GFP-tagged Edc3, Edc3ΔPE, and Edc3ΔAPD in the *Δedc3* strain to examine their condensate

phenotypes. Remarkably, as observed with Nst1, after 4 hours of overexpression, Edc3 formed mature

condensates, whereas Edc3ΔPA, similar to Edc3ΔPE, exhibited microphase separation in all cells (Fig.

S8). Even after 24 hours of induction, Edc3ΔPA failed to form mature condensates in almost all cells, and

only microphase-separated condensates were observed, demonstrating the role of PA in coalescence of

Edc3 condensates. Edc3ΔAPD, consistent with the result of Nst1, did not form condensates (Fig. S8).

We then investigated whether the phenotype of PA deletion mutants observed in the budding yeast

proteins Nst1 and Edc3 is also present in proteins with self-condensation potential that are implicated in

neurodegenerative diseases. Since TDP-43 are known to undergo self-condensation upon

overexpression in budding yeast (Fushimi et al., 2011; Johnson et al., 2008; Outeiro and Lindquist, 2003;

Sun et al., 2011), we first confirmed TDP-43 aggregation upon overexpression in budding yeast as been

previously reported. Under the same overexpression conditions used for Nst1 and Edc3, TDP-43 also

underwent self-condensation in budding yeast (Fig. 6B). Two PA domains (PA1, PA2) in the TDP-43

sequence were designated with using the sliding window approach (Fig. 6A, S3). The low complexity

region in the C-terminal domain (CTD) has been known to be critical for TDP-43 amyloid aggregation

(Chien et al., 2021; Li et al., 2021; Shenoy et al., 2023). The CTD has a relatively low charged amino acid

cluster score, but within the CTD, residues 255-276, which have a relatively high density of charged

amino acid, were designated as the PAM domain. To identify intrinsic sequence determinants of TDP-43

aggregation, we generated and overexpressed those deletion mutants of TDP-43 in the budding yeast

system and monitored them over time. Strikingly, the PA1 and PA2 deletion mutants failed to form mature

condensates at 4 hours of overexpression compared to the full-length TDP-43, suggesting their roles in

coalescence of condensates as observed role of PA in Nst1 and Edc3 (Fig. 6B). Notably, the previously

characterized CTD deletion mutant also formed condensates when overexpressed, similar to the reported

study (Johnson *et al.*, 2008; Wang et al., 2018a). The morphology of these condensates differed from that

of the full-length TDP-43: while the full-length TDP-43 formed both large and small, irregularly shaped

condensates, the CTD deletion mutant produced very large, round condensates in much lower numbers

per cell (Fig. 6B). Interestingly, the condensate morphology of the TDP-43ΔCTD mutant was

recapitulated in the TDP-43ΔPAM mutant, suggesting that the phenotype might be attributable to the loss

of the PAM domain within the CTD rather than the entire NTD existence or CTD deletion. This implies that

the charged amino acid cluster within the CTD may play a key role in determining the physical properties

and dynamics of TDP-43 condensates. Since these TDP-43 mutants generated retain the nuclear

localization signal (NLS), all overexpressed mutants predominantly localize to and form condensates in

the nucleus. To observe the condensates of mutants in the cytoplasm, we additionally generated NLS-

mutated TDP-43 variants. These variants exhibited phenotypes similar to those observed in the nucleus,

except for the CTD-deleted TDP-43 variant: the PA1 and PA2 deletion mutants failed to form mature

condensates (Fig 6B). Notably, while overexpressed TDP-43ΔCTD forms condensates in the nucleus, it

remains diffusely distributed in the cytoplasm (Fig. 6B). The phenotypic pattern observed for each mutant

after 4 hours of induction remained unchanged in cells induced for 8, 10, or 12 hours (S9).

We next investigated whether the TDP-43 coalescence tendency observed for the PA1 and PA2

mutants, when expressed in yeast, was attributable to the negative charges of the clustered amino acids.

To this end, we generated *PA1-1*, *PA1-2*, *PA2-1*, and *PA2-2* mutants in which the glutamate (E) and

aspartate (D) residues in the partial N- and C-terminal regions of PA1 and PA2 were individually

substituted with glutamine (Q) and asparagine (N), respectively, as well as *PA1-1PA1-2* and *PA2-1PA2-2*

mutants in which all glutamate and aspartate residues were replaced with glutamine and asparagine (Fig.

6C).

Strikingly, upon galactose induction in cells, the PA1 and PA2 mutants with all negatively charged

amino acids replaced by uncharged glutamine and asparagine exhibited completely different profile in

puncta formation of the wildtype TDP-43 protein (Fig. 6D). Unlike the ΔPA1 mutant, the *PA1-1PA1-2*

mutant exhibited a distinct phenotype: while the overexpressed ΔPA1 mutant displayed microphase

separation, the overexpressed *PA1-1PA1-2* showed a pronounced inhibition of condensation itself (Fig.

6D). This mutation appears to confer a gain of function, since substituting the negatively charged aspartic

acid (D) and glutamic acid (E) residues in PA1 with uncharged but chemically similar glutamine (Q) and

asparagine (N) not only abolishes coalescence, but also impairs the initial seeding of puncta formation—a

property not observed in either the wild-type protein or the ΔPA1 mutant. Furthermore, the *PA1-1* mutant,

with partial N-terminal substitution, exhibited morphology identical to wildtype condensates, whereas

PA1-2 resembled the *PA1-1PA1-2* mutant, suggesting that the substitution of E and D residues with Q

and N within the [13]DEPIEIPSEDD[23] region of PA1 is critical in driving this diffused phenotype. These

results indicate that the gain of function induced by the specific substitutions in PA1 includes both the

inhibition of coalescence and the blocking of seeding during puncta formation.

Notably, the results for PA2 domain are remarkable. The *PA2-1PA2-2* exhibited puncta morphology

identical to that of ΔPA2 and, unlike the wildtype TDP-43, displayed a pronounced tendency toward

microphase separation (Fig. 6D). Unlike the PA1 mutants, both *PA2-1, PA2-2* mutant having partial

substitution of negatively charged amino acids in either the N-terminal or C-terminal regions failed to fully

recapitulate the puncta generation patterns of ΔPA2 (Fig. 6D). These results indicate that it is not specific

negatively charged residues within PA2 that are critical, but rather the overall clusters of negative charged

amino acids in PA2 contribute to TDP-43 coalescence.

To determine whether the condensate phenotypes of PA1, PA2, PAM, and CTD deletion mutants of

TDP-43 observed in budding yeast are also present in mammalian cells, we overexpressed GFP-fused

TDP-43 and TDP-43 mutants in U2OS cells. Consistent with our findings in budding yeast, the PA1 and

PA2 deletion mutants did not form mature condensates in mammalian cells (Fig. 6E). In particular, the

PA2 deletion mutant failed to form condensates in all cells expressing the protein, as indicated by the

presence of GFP signal (Fig. 6E). Interestingly, the PAM domain deletion mutant exhibited a markedly

larger size and stronger intensity of puncta formation compared to other mutants. To observe the

condensation of TDP-43 derivatives in the cytoplasm, we introduced an additional ΔNLS (K82A, R83A,

K84A) mutation into each TDP-43 domain deletion mutant and examined their expression in U2OS cells

(Lu et al., 2022). The TDP-43 derivatives in the nucleus displayed a similar pattern of puncta formation.

Notably, in the case of the ΔPA1ΔPA2 double deletion, puncta formation was further inhibited. These

data suggest that PA domains could be a general key contributor to condensation of TDP-43.

DISCUSSION

Condensates by LLPS are known to undergo formation, maturation, and subsequent transition into

aggregates within the cellular environment (Shen et al., 2023; Wang et al., 2021). Through *in vitro* studies,

it has been found that the maturation of liquid-like condensates proceeds via three major principal

mechanisms: coalescence, whereby individual condensates fuse to form larger structures; Ostwald

ripening, in which constituent molecules from smaller condensates are transferred to larger ones,

resulting in the growth of the latter and shrinkage of the former; and diffusion-limited growth,

characterized by the incorporation of freely diffusing molecules into existing condensates (Banerjee et al.,

2024; Lee et al., 2021; Linsenmeier et al., 2021). The intrinsic sequence properties of proteins are

believed to play critical roles in regulating each of these processes; however, *in vivo* studies elucidating

their molecular grammar remain relatively limited.

Low complexity regions (LCRs) are particularly intriguing segments within protein sequences. They are

composed of highly repetitive stretches consisting of just one or a few types of amino acids. It is known

that, compared to prokaryotes, LCRs are more than 200 times more prevalent in eukaryotes (Dickson and

Golding, 2022; Sim and Creamer, 2002). LCRs are closely associated with phenotypic variation (Fondon

and Garner, 2004) and function as multivalent interaction hubs, playing a critical role in protein

condensation and amyloidogenesis (Banjade and Rosen, 2014; Linsenmeier *et al.*, 2023). Collectively,

these findings strongly suggest that LCRs are physiologically and pathologically important components in

complex biological systems.

Charged amino acid clusters which is included in LCRs represent one of the most frequently observed

sequence motifs within proteins consisting of RNP granules (Niedner-Boblenz et al., 2024; Regy et al.,

2020; Saitoh et al., 2004). Within intrinsically disordered regions (IDRs), there have been numerous clues

that certain charged amino acids may contribute to the regulation of condensation, either by promoting or

inhibiting condensation (Saar et al., 2021). Both negative and positive charged amino acids are thought to

modulate intermolecular affinity via electrostatic interactions, where the pattern of charged residues along

the sequence plays a pivotal role. As mentioned previously, when charged residues are uniformly

distributed, the driving force for phase separation is attenuated; in contrast, when charged residues are

clustered or segregated from other residues, the propensity for phase separation is enhanced (Borcherds

et al., 2021; Chen et al., 2021; Miyagi et al., 2022). Therefore, the charged amino acid clusters within

condensate-forming proteins likely play critical roles in condensation.

This study presents the first report on the role of the low-complexity domains (LCDs) consisting of the

negative charged amino acids, as sequence elements required for the maturation of protein condensates

in cells. Deletion of the PA region in Nst1, Edc3, and TDP-43 allowed condensate formation but impaired

their coalescence and maturation, suggesting that the PA domain may be generally required for the

formation of mature condensates in other proteins as well.

Substituting the negatively charged aspartate (D) and glutamate (E) in the PA1 of TDP-43 with

uncharged but chemically similar glutamine (Q) and asparagine (N) abolishes puncta formation or

maturation, demonstrating the critical role of the negative charge cluster in PA1 for condensate seeding

and coalescence (Fig. 6). This strongly suggests that the polyanion region plays a critical role in proteins

with high condensation potential and underscores the necessity to further validate its function in TDP-43

condensation within mammalian models. Notably, in motor neurons of ALS-FTD patients, the reduction of

intracellular TDP-43 monomer concentration due to aberrant condensation and aggregation is a major

driver of disease progression (Kellett et al., 2025; Oiwa et al., 2023). Therefore, identifying domains

capable of modulating TDP-43 condensation could represent an initial step toward developing novel

therapeutic strategies. The PA1 and PA2 domains of TDP-43 identified in this study are short regions,

each comprising approximately 20 amino acids. Subsequent studies should focus on pinpointing specific

residues within PA1 and PA2 which function in TDP-43 coalescence.

Bioinformatics studies revealed that D/E repeats are widespread across eukaryotic proteomes (Wang et

al., 2023), with nearly half of the proteins containing these negatively charged clusters being DNA/RNA-

binding proteins (Wang *et al.*, 2023). In DNA-binding transcription factors such as HMGB1, RFX1, Sox11,

and UBF1, these negative charge clusters are thought to interact with their own positive-charged DNA

binding domains, mediating autoinhibition and thereby modulating DNA binding affinity. The recent

research on HMGB1 is particularly noteworthy (Mensah et al., 2023). HMGB1 functions as a chromatin-

binding architectural protein that regulates DNA accessibility, gene expression, and genome stability

within the cell (Lange and Vasquez, 2009). The C-terminal domain of HMGB1, which consists of a

negative charge amino acid cluster, when replaced by positively charged amino acids in a mutant,

exhibits amorphous and rigid physical properties *in vitro*, contrasting with the wild-type protein. In addition,

its mutants in which the negative charge amino acid cluster is deleted, fail to localize properly in the

nucleolus, thereby influencing physiological function of HMGB1 (Mensah *et al.*, 2023). This suggests that

a broader study is warranted on the correlation between the condensation behavior of proteins containing

sequence-specific charged amino acid clusters and the effect of such condensation on their physiological

functions.

Another noteworthy observation regarding the PAM domain deletion mutant of TDP-43 is that

condensates formed by this mutant become enlarged compared to the condensates of wild-type (Fig. 6).

A similar, though less pronounced, trend was also observed in Nst1 (Fig. 2). It is possible that deletion of

the PAM domain reduces the local charge surrounding the PA domain, thereby amplifying the functional

contribution of the PA domain. Furthermore, because loss of the PAM domain itself increases the fluidity

of condensates, the combined effect of enhanced affinity and elevated fluidity may lead to the observed

enlargement of condensates.

Altogether, our study provides compelling evidence that the arrangement of low complexity regions

(LCRs) and clustered charged amino acids within proteins serves as a key regulator in the formation and

maturation of biomolecular condensates associated with liquid-liquid phase separation (LLPS) in cell.

These findings underscore the significance of correlating the patterns and organization of LCRs, such as

charged clusters, with the material state, subcellular localization, and functional outcomes of condensates.

Further investigation into the biochemical characteristics of condensates, which vary according to the

molecular grammar of proteins, will illuminate the diverse physiological functions of condensates within

cells, and may reveal a link between excessive condensate or aggregate formation and the onset of

pathological states.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT OR RESOURCE	SOURCE	IDENTIFIER
Antibodies		
goat anti-GFP	ROCKLAND, USA	Cat# 600-101-215
mouse anti-Tub1	Sigma, USA	Cat# T5168
anti-goat (secondary antibody)	Jackson ImmunoResearch Laboratories, Inc, USA	Cat# 705-035-003

anti-Mouse (secondary antibody)	Santa Cruz Biotechnology, USA	Cat# sc-2005
Bacterial and virus strains		
Escherichia coli DH5α cells One Shot™ TOP10 Chemically Competent *E. coli*	Invitrogen, USA	Cat# C404003
Biological samples		
Salmon sperm DNA (Deoxyribonucleic acid sodium salt from salmon testes)	Sigma, USA	Cat# D1626
Chemicals, peptides, and recombinant proteins		
1,6-Hexanediol	DAEJUNG CHEMICAL & METALS CO., LTD, South Korea	Cat# 629-11-8
4% paraformaldehyde	T&I, South Korea	Cat# BPP-9004
	Cytiva,USA,	Cat# SH30088.03
WestGlowTM PICO PLUS Chemiluminescent Substrate	BIOMAX, South Korea	Cat# BWP0200
20X TBS-T (20X Tris-Buffered Saline/1% Tween-20)	T&I, South Korea	Cat# BTT-9120
Antibiotic-antimycotic solution (Anti-Anti)	Gibco, USA,	Cat# 15240-062
Experimental models: Cell lines		
U2OS cells Human osteosarcoma cell line	ATCC	HTB-96
Experimental models: Organisms/strains		
Saccharomyces cerevisiae W303a *leu2-3,112 trp1-1 can1-100 ura3-1 ade2-1 his3-11,15*	S. Elledge	YSK3549
Saccharomyces cerevisiae W303a except as *Δedc3::KANMX*	this study	YSK3707
Saccharomyces cerevisiae W303a except as *Δnst1::KANMX*	this study	YSK3490
Saccharomyces cerevisiae W303a except as *NST1-EGFP::KANMX*	K.Song	YSK3498
Saccharomyces cerevisiae W303a except as *DCP2-EGFP::HIS3MX6*	K.Song	YSK3527
Saccharomyces cerevisiae W303a except as *EDC3-EGFP::HIS3MX6*	K.Song	YSK3528

Saccharomyces cerevisiae W303a except as *Δnst1::KANMX pNST1-GFP-NST1::URA3*	this study	YSK3700
Saccharomyces cerevisiae W303a except as *Δnst1::KANMX pNST1-GFP-NST1ΔPA::URA3*	this study	YSK3701
Saccharomyces cerevisiae W303a except as *Δnst1::KANMX pNST1-GFP-NST1ΔPAM::URA3*	this study	YSK3702
Saccharomyces cerevisiae W303a except as *Δnst1::KANMX pNST1-GFP-NST1ΔAPD::URA3*	this study	YSK3714
Saccharomyces cerevisiae W303a except as *Δnst1::KANMX PAB1-EGFP::HIS3MX6*	this study	YSK3704
Saccharomyces cerevisiae W303a expect as *Δnst1::KANMX EDC3-EGFP::HIS3MX6*	this study	YSK3694
Saccharomyces cerevisiae W303a expect as *Δnst1::KANMX DHH1-EGFP::HIS3MX6*	this study	YSK3695
Saccharomyces cerevisiae W303a expect as *Δnst1::KANMX DCP2-EGFP::HIS3MX6*	this study	YSK3709
Saccharomyces cerevisiae BY4741expect as *Δnst1::KANMX DCP2-mKate2-sphis5*	this study	YSK3663
Oligonucleotides		
pMW20(U)-pGAL-GFP-NST1Δ2-122 deletion top 5' ATTACACATGGCATGGATGAACGAGCTCAGTTTAATGTCATAAGGAATTACCAGGACGAT 3'	this study	KS2262
pMW20(U)-pGAL-GFP-NST1Δ2-122 deletion bottom 5' ATCGTCCTGGTAATTCCTTATGACATTAAACTGAGCTCGTTCATCCATGCCATGTGTAAT 3'	this study	KS2263
pMW20(U)-pGAL-GFP-NST1Δ123-293 deletion top 5' ACTATTCTACGAATCGAAAAGGAGGAAGTACAAAAAATGATGATGTATCCGAATATTAGA 3'	this study	KS2384
pMW20(U)-pGAL-GFP-NST1Δ123-293 deletion bottom 5' TCTAATATTCGGATACATCATCATTTTTTGTACTTCCTCCTTTTCGATTCGTAGAATAGT 3'	this study	KS2385

pMW20(U)-pGAL-GFP-NST1Δ294-459 deletion top 5' TCACATCCCAAAATAGCAGAGGAGTATGTAGAAGATGAAGATGAGGAAGAC TATGATGAC 3'	this study	KS2386
pMW20(U)-pGAL-GFP-NST1Δ294-459 deletion bottom 5' GTCATCATAGTCTTCCTCATCTTCATCTTCTACATACTCCTCTGCTATTTTGG GATGTGA 3'	this study	KS2387
pMW20(U)-pGAL-GFP-NST1Δ460-630 deletion top 5' GATACTTCGTTAGATACTGATTCTATATATCTAGAAGAAGAAAAAAGAAAGAA AAGAGAG 3'	this study	KS2280
pMW20(U)-pGAL-GFP-NST1Δ460-630 deletion bottom 5' CTCTCTTTTCTTTCTTTTTTCTTCTTCTAGATATATAGAATCAGTATCTAACGA AGTATC 3'	this study	KS2281
pMW20(U)-pGAL-GFP-NST1Δ755-844 deletion top 5' AGAGAACAGAAAAGGCTAGAACAAGAAAAGGATCTACTTCTTCCTTCAACTA ACAACCAG 3'	this study	KS2397
pMW20(U)-pGAL-GFP-NST1Δ755-844 deletion bottom 5' CTGGTTGTTAGTTGAAGGAAGAAGTAGATCCTTTTCTTGTTCTAGCCTTTTCT GTTCTCT 3'	this study	KS2398
pMW20(U)-pGAL-GFP-NST1Δ845-1015 deletion top 5' CCAGTTTCACCCACAGGCTTTAACGTTCATGATCAAAAAAGCTCATTTTCGG GACCGTCG 3'	this study	KS2388
pMW20(U)-pGAL-GFP-NST1Δ845-1015 deletion bottom 5' CGACGGTCCCGAAAATGAGCTTTTTTGATCATGAACGTTAAAGCCTGTGGG TGAAACTGG 3'	this study	KS2389
pMW20(U)-pGAL-GFP-NST1Δ460-596 deletion top 5' GATACTTCGTTAGATACTGATTCTATATATCGGAAATTGATTCAGATAGCCAT TACAAAG 3'	this study	KS2437
pMW20(U)-pGAL-GFP-NST1Δ460-596 deletion bottom 5'GATACTTCGTTAGATACTGATTCTATATATCGGAAATTGATTCAGATAGCC ATTACAAAG 3'	this study	KS2438
pMW20(U)-pGAL-GFP-EDC3Δ65-205 deletion top 5' AGGTTCTAACTGTTGCCTCTCCCAAATACCTACCAATAAC 3'	this study	KS2360

pMW20(U)-pGAL-GFP-EDC3Δ65-205 deletion bottom 5' GTTATTGGTAGGTATTTGGGAGAGGCAACAGTTAGAACCT 3'	this study	KS2361
pMW20(U)-pGAL-GFP-EDC3Δ65-180 deletion top 5' AAGGACTTAAAGGTTCTAACTGTTGCCTCTCCGCAACAAGATGCTCAAGATG ATCTGGTT 3'	this study	KS2446
pMW20(U)-pGAL-GFP-EDC3Δ65-180 deletion bottom 5' AACCAGATCATCTTGAGCATCTTGTTGCGGAGAGGCAACAGTTAGAACCTTT AAGTCCTT 3'	this study	KS2447
pMW20(U)-pGAL-GFP-EDC3Δ181-205 deletion top 5' AGAAATGAGCAAAGCACACACCAATCTCAGCCCAAATACCTACCAATAACTC AGTCTTTG 3'	this study	KS2444
pMW20(U)-pGAL-GFP-EDC3Δ181-205 deletion bottom 5' CAAAGACTGAGTTATTGGTAGGTATTTGGGCTGAGATTGGTGTGTGCTTTG CTCATTTCT 3'	this study	KS2445
pMW20(U)-pGAL-GFP-EDC3Δ330-380 deletion top 5' AACTAGGTGGCCGTGCACGTCTACAGGATTCCATGGTCAA 3'	this study	KS2362
pMW20(U)-pGAL-GFP-EDC3Δ330-380 deletion bottom 5' TTGACCATGGAATCCTGTAGACGTGCACGGCCACCTAGTT 3'	this study	KS2363
pMW20(U)-pGAL-GFP-TARDBP infusion cloning top 5' GCATGGATGAACGAGCTTCTGAATATATTCGGGTAAC 3'	this study	KS2401
pMW20(U)-pGAL-GFP-TARDBP infusion cloning bottom 5' GTCGACTCTAGAGGATCCCTACATTCCCCAGCCAGAAG 3'	this study	KS2402
pMW20(U)-pGAL-GFP-TARDBPΔ2-29 deletion top 5' GGGATTACACATGGCATGGATGAACGAGCTACGGTTACAGCCCAGTTTCCA GGGGCGTGT 3'	this study	KS2429
pMW20(U)-pGAL-GFP-TARDBPΔ2-29 deletion bottom 5' ACACGCCCCTGGAAACTGGGCTGTAACCGTAGCTCGTTCATCCATGCCATG TGTAATCCC 3'	this study	KS2430
pMW20(U)-pGAL-GFP-TARDBPΔ197-220 deletion top 5' CCTTTGAGAAGCAGAAAAGTGTTTGTGGGGTTCATCCCCAAGCCATTCAGG GCCTTTGCC 3'	this study	KS2448

pMW20(U)-pGAL-GFP-TARDBPΔ197-220 deletion bottom 5' GGCAAAGGCCCTGAATGGCTTGGGGATGAACCCCACAAACACTTTTCTGCT TCTCAAAGG 3'	this study	KS2449
pMW20(U)-pGAL-GFP-TARDBPΔ255-276 deletion top 5' GGAGAGGACTTGATCATTAAAGGAATCAGCGGTGGTAATCCAGGTGGCTTT GGGAATCAG 3'	this study	KS2431
pMW20(U)-pGAL-GFP-TARDBPΔ255-276 deletion bottom 5' CTGATTCCCAAAGCCACCTGGATTACCACCGCTGATTCCTTTAATGATCAAG TCCTCTCC 3'	this study	KS2432
pMW20(U)-pGAL-GFP-TARDBPΔ255-414 deletion top 5' GGACTTGATCATTAAAGGAATCAGCTAGGGATCCTCTAGAGTCGACCTGC 3'	this study	KS2439
pMW20(U)-pGAL-GFP-TARDBPΔ255-414 deletion bottom 5' GCAGGTCGACTCTAGAGGATCCCTAGCTGATTCCTTTAATGATCAAGTCC 3'	this study	KS2440
TARDBPΔNLS (K82A, R83A, K84A) top 5' GTGTATGTTGTCAACTATCCAAAAGATAACGCAGCAGCAATGGATGAGACA GATGCTTCATCAGCAGTG 3'	this study	KS2472
TARDBPΔNLS (K82A, R83A, K84A) bottom 5' CACTGCTGATGAAGCATCTGTCTCATCCATTGCTGCTGCGTTATCTTTTGGA TAGTTGACAACATACAC 3'	this study	KS2473
pRS306-pNST1-GFP-NST1 infusion cloning top 5' CGTCCCGGGATGAGTAAAGGAGAAGAACTTTTC 3'	this study	KS2238
pRS306-pNST1-GFP-NST1 infusion cloning bottom 5' CGTCTCGAGCAAATTTAAGACGTTTGCGTAG 3'	this study	KS2239
pRS306-pNST1-GFP-NST1Δ460-630 top 5' GATACTTCGTTAGATACTGATTCTATATATCTAGAAGAAGAAAAAAGAAAGAA AAGAGAG 3'	this study	Ks2280
pRS306-pNST1-GFP-NST1Δ460-630 bottom 5' CTCTCTTTTCTTTCTTTTTTCTTCTTCTAGATATATAGAATCAGTATCTAACGA AGTATC 3'	this study	Ks2281

pRS306-pNST1-GFP-NST1Δ631-752 top 5' GATAATAACAGACTAAAACTTCTACAAGAACTACAGAAAGAAAAAGAAGAAG AAGAGAGA 3'	this study	KS2176
pRS306-pNST1-GFP-NST1Δ631-752 bottom 5' TCTCTCTTCTTCTTCTTTTTCTTTCTGTAGTTCTTGTAGAAGTTTTAGTCTGTT ATTATC 3'	this study	KS2177
pRS306-pNST1-GFP-NST1Δ1016-1240 top 5' GGCATGGATGAACGAGCTCAGGATCAAAAAAGCTCATTTTCG 3'	this study	KS2194
pRS306-pNST1-GFP-NST1Δ1016-1240 bottom 5' CGAAAATGAGCTTTTTTGATCCTGAGCTCGTTCATCCATGCC 3'	this study	KS2195
pcDNA3.1myc-HisA-GFP-TARDBP infusion cloning top 5'AGCTGGCTAGTTAAGCTTGGTACCATGAGTAAAGGAGAAGAACTTTTC 3'	this study	KS2455
pcDNA3.1myc-HisA-GFP-TARDBP infusion cloning bottom 5'GCGGCCGCCACTGTGCTGGATATCCTACATTCCCCAGCCAGAAGACTT 3'	this study	KS2456
TARDBP *PA1-1* (E3Q, E9Q, D10N, E11Q) ΔNLS (K82A, R83A, K84A) top 5' GCATGGATGAACGAGCTTCTCAGTATATTCGGGTAACCCAGAACCAGAACG ATGAGCCCATTGAAAT 3'	this study	KS2492
TARDBP *PA1-1* (E3Q, E9Q, D10N, E11Q) ΔNLS (K82A, R83A, K84A) bottom 5' ATTTCAATGGGCTCATCGTTCTGGTTCTGGGTTACCCGAATATACTGAGAAG CTCGTTCATCCATGC 3'	this study	KS2493
TARDBP *PA1-2* (D13N, E14Q, E17Q, E21Q, D22N, D23N) ΔNLS (K82A, R83A, K84A) top 5' GGGTAACCGAAGATGAGAACAACCAGCCCATTCAGATACCATCGCAGAACA ACGGGACGGTGCTGCTCTCCAC 3'	this study	KS2494
TARDBP *PA1-2* (D13N, E14Q, E17Q, E21Q, D22N, D23N) ΔNLS (K82A, R83A, K84A) bottom 5' GTGGAGAGCAGCACCGTCCCGTTGTTCTGCGATGGTATCTGAATGGGCTG GTTGTTCTCATCTTCGGTTACCC 3'	this study	KS2495
TARDBP *PA1-1PA1-2* (E3Q, E9Q, D10N, E11Q, D13N, E14Q, E17Q, E21Q, D22N, D23N) ΔNLS (K82A, R83A, K84A) top 5' GCATGGATGAACGAGCTTCTCAGTATATTCGGGTAACCCAGAACCAGAACA ACCAGCCCATTCAGAT 3'	this study	KS2500
TARDBP *PA1-1PA1-2* (E3Q, E9Q, D10N, E11Q, D13N, E14Q, E17Q, E21Q, D22N, D23N) ΔNLS (K82A, R83A, K84A) bottom 5' ATCTGAATGGGCTGGTTGTTCTGGTTCTGGGTTACCCGAATATACTGAGAA GCTCGTTCATCCATGC 3'	this study	KS2501

TARDBP *PA2-1* (E200Q, D201N, E204Q, D205N, E206Q, E209Q) ΔNLS (K82A, R83A, K84A) top 5' TGTTTGTGGGGCGCTGTACACAGAACATGACTCAGAACCAGCTGCGGCAGT TCTTCTCTCAGTACGGGGA 3'	this study	KS2484
TARDBP *PA2-1* (E200Q, D201N, E204Q, D205N, E206Q, E209Q) ΔNLS (K82A, R83A, K84A) bottom 5' TCCCCGTACTGAGAGAAGAACTGCCGCAGCTGGTTCTGAGTCATGTTCTGT GTACAGCGCCCCACAAACA 3'	this study	KS2485
TARDBP *PA2-2* (D216N, D219N) ΔNLS (K82A, R83A, K84A) top 5' AGTTCTTCTCTCAGTACGGGAACGTGATGAACGTCTTCATCCCCAAGCCATT 3'	this study	KS2486
TARDBP *PA2-2* (D216N, D219N) ΔNLS (K82A, R83A, K84A) bottom 5' AATGGCTTGGGGATGAAGACGTTCATCACGTTCCCGTACTGAGAGAAGAAC T 3'	this study	KS2487
Recombinant DNA		
pMW20(U)-pGAL-GFP	This study	
pMW20(U)-pGAL-GFP-NST1	This study	
pMW20(U)-pGAL-GFP-NST1Δ2-122	This study	
pMW20(U)-pGAL-GFP-NST1Δ123-293	This study	
pMW20(U)-pGAL-GFP-NST1Δ294-459	This study	
pMW20(U)-pGAL-GFP-NST1Δ460-752	This study	
pMW20(U)-pGAL-GFP-NST1Δ460-630	This study	
pMW20(U)-pGAL-GFP-NST1Δ631-752	This study	
pMW20(U)-pGAL-GFP-NST1Δ753-844	This study	
pMW20(U)-pGAL-GFP-NST1Δ845-1015	This study	
pMW20(U)-pGAL-GFP-NST1Δ1016-1240	This study	
pMW20(U)-pGAL-GFP-NST1Δ460-596	This study	
pMW20(U)-pGAL-GFP-EDC3	This study	
pMW20(U)-pGAL-GFP-EDC3Δ65-205	This study	
pMW20(U)-pGAL-GFP-EDC3Δ65-180	This study	
pMW20(U)-pGAL-GFP-EDC3Δ181-205	This study	
pMW20(U)-pGAL-GFP-EDC3Δ330-380	This study	
pMW20(U)-pGAL-GFP-TARDBP	This study	

pMW20(U)-pGAL-GFP-TARDBPΔ2-29	This study	
pMW20(U)-pGAL-GFP-TARDBPΔ197-220	This study	
pMW20(U)-pGAL-GFP-TARDBPΔ255-276	This study	
pMW20(U)-pGAL-GFP-TARDBPΔ255-414	This study	
pMW20(U)-pGAL-GFP-TARDBPΔNLS (K82A, R83A, K84A)	This study	
pMW20(U)-pGAL-GFP-TARDBPΔ2-29ΔNLS (K82A, R83A, K84A)	This study	
pMW20(U)-pGAL-GFP-TARDBPΔ197-220ΔNLS (K82A, R83A, K84A)	This study	
pMW20(U)-pGAL-GFP-TARDBPΔ255-276ΔNLS (K82A, R83A, K84A)	This study	
pMW20(U)-pGAL-GFP-TARDBPΔ255-414ΔNLS (K82A, R83A, K84A)	This study	
pRS306-pNST1-GFP-NST1	This study	
pRS306-pNST1-GFP-NST1Δ460-630	This study	
pRS306-pNST1-GFP-NST1Δ631-752	This study	
pRS306-pNST1-GFP-NST1Δ1016-1240	This study	
pcDNA3.1myc-HisA-GFP-TARDBP	This study	
pcDNA3.1myc-HisA-GFP-TARDBPΔ2-29	This study	
pcDNA3.1myc-HisA-GFP-TARDBPΔ255-276	This study	
pcDNA3.1myc-HisA-GFP-TARDBPΔ255-414	This study	
pcDNA3.1myc-HisA-GFP-TARDBPΔ197-220	This study	
pcDNA3.1myc-HisA -GFP-TARDBPΔNLS(K82A, R83A, K84A)	This study	
pcDNA3.1myc-HisA -GFP-TARDBPΔ2-29ΔNLS(K82A, R83A, K84A)	This study	
pcDNA3.1myc-HisA -GFP-TARDBPΔ197-220ΔNLS(K82A, R83A, K84A)	This study	
pcDNA3.1myc-HisA -GFP-TARDBPΔ255-276ΔNLS(K82A, R83A, K84A)	This study	
pcDNA3.1myc-HisA -GFP-TARDBPΔ255-414ΔNLS(K82A, R83A, K84A)	This study	
pMW20(U)-pGAL-GFP-TARDBP *PA1-1*(E3Q, E9Q, D10N, E11Q) ΔNLS (K82A, R83A, K84A)	This study	
pMW20(U)-pGAL-GFP-TARDBP *PA1-2*(D13N, E14Q, E17Q, E21Q, D22N, D23N) ΔNLS (K82A, R83A, K84A)	This study	
pMW20(U)-pGAL-GFP-TARDBP *PA1-1PA1-2* (E3Q, E9Q, D10N, E11Q, D13N, E14Q, E17Q, E21Q, D22N, D23N) ΔNLS (K82A, R83A, K84A)	This study	
pMW20(U)-pGAL-GFP-TARDBP *PA2-1* (E200Q, D201N, E204Q, D205N, E206Q, E209Q) ΔNLS (K82A, R83A, K84A)	This study	
pMW20(U)-pGAL-GFP-TARDBP *PA2-2* (D216N, D219N) ΔNLS (K82A, R83A, K84A)	This study	

pMW20(U)-pGAL-GFP-TARDBP *PA2-1PA2-2* (E200Q, D201N, E204Q, D205N, E206Q, E209Q, D216N, D219N) ΔNLS (K82A, R83A, K84A)	This study	
Software and algorithms		
AIUPRed – Prediction of Intrinsically Unstructured Proteins	Eötvös Loránd University (ELTE), MTA-ELTE Momentum Bioinformatics Research Group, Hungary	AIUPred
Disopred3	University College London (UCL), Bioinformatics Group	PSIPRED Workbench
PrDOS – Protein Disorder Prediction System	Osaka University, Graduate School of Information Science and Technology	PrDOS - Protein disorder prediction server
PASTA 2.0 – prediction of amyloid structural aggregation	University of Padova, Department of Biomedical Sciences, INFN, Italy	Aggregation Prediction with PASTA
Aggrescan – prediction of "hot spot" of aggregation on polypeptides	Autònoma University of Barcelona, Institute of Biotechnology and Biomedicine, Department of Biochemistry and Molecular Biology, Spain	Aggrescan: Prediction of "hot spots" of aggregation in polypeptides
FELLS – Fast Estimator of Latent Local Structure	BioComputing Up, University of Padua, Italy	FELLS > Submit
GraphPad Prism 5	GraphPad	https://www.graphpad.com/
GraphPad Prism 10	GraphPad	https://www.graphpad.com/

ImageJ	Schindelin et al. Fiji: an open-source platform for biological-image analysis. *Nat. Methods.* 2012; 9: 676-682	https://imagej.nih.gov/ij/
Custom Python scripts	This paper	Available upon request
Other		
1.8 ml bead-beating tube	SIMPORT, Canada	Cat# T335-6SPR
Acid-washed 0.5 mm bead	Biospec, USA	Cat# 11079105
Polyvinylidene difluoride (PVDF) membrane	Merk Millipore, USA	Cat# IPVH00010

METHOD DETAILS

Yeast cell culture and transformation

Saccharomyces cerevisiae yeast cells were grown in YPAD medium until reaching the exponential growth phase (OD_{600} 0.5). Cells were harvested by centrifugation at 4,000 rpm for 5 min at room temperature (RT). The cell pellet was gently resuspended in 1 mL of 0.1 M LiOAc (Sigma-Aldrich, USA, L6883-250G) and incubated at RT for 5 min, followed by another centrifugation under the same condition. For transformation, cells were gently resuspended in 360 μL of yeast transformation mixture consisting of 50% polyethylene glycol (Sigma-Aldrich, USA, P4338-500G), 1 M LiOAc, 2 mg/mL salmon sperm DNA (Sigma, USA, D1626), and ddH$_2$O. 100 ng plasmid DNA was added to the cell suspension, and the mixture was incubated at RT for 30 min. Heat shock was performed by incubating the cells at 42°C for 30 min, followed by cooling at RT for 3 min. Cells were harvested again by centrifugation at 4,000 rpm for 5 min at RT, and plated onto SC-Ura plates. Plates were incubated at 25°C until colonies appeared typically 2–3 days.

Protein overexpression in yeast

The cells transformed with plasmids containing NST1, EDC3, *TARDBP* and their deletion mutants were

cultured in a SC-U + 2% glucose medium up to the exponential growth phase (OD$_{600}$ 0.5) and harvested.

The cells were washed with a SC-U + 2% raffinose + 0.1% glucose medium three times, diluted to half its

concentration, and cultured for an additional 3 hours in SC-U + 2% raffinose + 0.1% glucose. Thereafter,

galactose was added to a final 2% concentration and incubated for designated hours.

1,6-Hexanediol treatment of yeast

Yeast cells induced for protein overexpression in 2% galactose for 4 hours were harvested by

centrifugation at 4000 rpm for 5 minutes, and the resulting pellet was collected. The cell pellet was then

resuspended and incubated in SC-U medium supplemented with 2% galactose, 2% raffinose, 0.1%

glucose, and 10% 1,6-hexanediol (Daejung Chemical & Metals Co., LTD, South Korea, 629-11-8) for 1

hour.

Chromosomal integration of genes in yeast

Various NST1 deletion mutants were acquired by PCR-mediated deletion cloning with the full length

pRS306-pNST1-NST1 and verified by sequencing. Plasmids pRS306-pNST1-NST1 and its deletion

mutants were treated with EcoRV to make a single cut at the URA3 and were transformed into *Δnst1* cells

(YSK3490) by 42 °C heat shock. Integration into the chromosome occurred at low frequency

via homologous recombination at the URA3. The transformed cells were plated on the SC plate without

uracil and colony PCR was performed to confirm whether the fragment was successfully transformed.

Infusion cloning of genes

The PCR-amplified coding sequences were generated using primers containing 15-bp extensions

homologous to the vector ends. The destination vector pcDNA3.1-myc-His was linearized by restriction

enzyme digestion at 37°C for 3 h. The digested vector and PCR fragments were then subjected to

infusion cloning reaction (EZ-HT cloning kit, Enzynomics, Korea, EZ016TM), which utilizes single-strand

annealing activity to join DNA fragments with complementary ends, circumventing the need for

conventional ligation. The reaction products were transformed into chemically competent *E. coli* DH5α

cells, and transformants were selected on LB agar plates supplemented with ampicillin (100 μg/ml). The

integrity of the resulting constructs was verified by DNA sequencing.

TDP-43 PA1 and PA2 Domain Mutagenesis

PCR-based site-directed mutagenesis was performed using the plasmid pMW20(U)-pGAL-GFP-

TARDBPΔNLS (K82A, R83A, K84A) as a template to introduce targeted amino acid substitutions within

the TDP-43 PA1 and PA2 domain. The *PA1-1* mutant was generated with primers KS2492 and KS2493,

and the *PA1-2* mutant with KS2494 and KS2495. The *PA2-1* mutant was generated with primers KS2484

and KS2485, and the *PA2-2* mutant with primers KS2486 and KS2487. The *PA1-1PA1-2* and *PA2-*

1PA2-2 mutants, which harbor both the *PA1-1* and *PA1-2* mutations or *PA2-1* and *PA2-2* mutations,

were constructed using the *PA1-2* or *PA2-1* plasmid as a template with the *PA1-1PA1-2* primers (KS2500

and KS2501) or the *PA2-2* primers (KS2486 and KS2487), respectively. The sequences of all primers

were listed in the KEY RESOURCES TABLE. All mutant strains were verified by DNA sequencing.

Widefield microscopy of cells and image analysis

604 GFP-labeled proteins expressed in the cells were visualized using an Axioplan2 microscope (Carl

605 Zeiss,Jena, Germany) equipped with a 100x Plan-Neofluar oil immersion objective. Consistent culture

606 conditions, exposure times, and fluorescence intensities were maintained across all strains in this study to

607 ensure a comparative analysis of puncta intensity. Image analysis was performed using FIJI ImageJ

608 (https://imagej.nih.gov/ij/).

Confocal microscopy of cells and image analysis

GFP-labeled proteins expressed in the cells were imaged using both a ZEISS LSM900 laser scanning

confocal microscope (Carl Zeiss, Jena, Germany) equipped with high-sensitivity GaAsP-PMT detectors (2

standard and 1 additional GaAsP-PMT), and a ZEISS LSM980 confocal microscope with an Airyscan 2

detector (Carl Zeiss, Jena, Germany). For Airyscan super-resolution imaging with the LSM980, the

Airyscan Joint Deconvolution mode was utilized, achieving lateral resolution down to ~90 nm. Both

systems were fitted with a 63x/1.4 NA oil immersion objective, and all images were acquired at room

temperature. Microscope settings, including laser power, detector gain, and pinhole size, were kept

identical for all samples to ensure quantitative comparability. Image acquisition and processing were

performed using ZEISS Zen software (version 3.9, Carl Zeiss).

FCS measurements and data analysis

All FCS measurements were performed at 25°C on a confocal microscope (LSM780; Carl Zeiss, Inc.

Germany) combined with a ConfoCor3 and a 40×/1.2 NA water-immersion objective (Carl Zeiss, Inc.).

Details of the analysis of fluorescence auto-correlation functions (FAFs) were described in the previous

study (Kawai-Noma et al., 2006; Pack et al., 2014), FAFs were fitted to a two-component model with a

triplet term.

For each cell, the cytoplasmic region was identified by visual inspection of the fluorescence image,

avoiding the GFP visualized puncta. At least five independent measurements (each lasting 10 s) were

recorded per cell. Over 15 cells were analyzed per condition.

Autocorrelation curves were generated using ZEN black software (Carl Zeiss, Inc.) and fitted with two-

component 3D diffusion model including a triplet term:

$$G_x(\tau) = 1 + \frac{1}{N} \sum_i F_i (1 + \frac{\tau}{\tau_i})^{-1} (1 + \frac{\tau}{s^2 \tau_i})^{-1/2}$$

Where F_i and τ_i are the fraction and diffusion time of component I, respectively; N is the average number

of fluorescent molecules in the confocal volume defined by the radius w_0 and the length $2z_0$; and s is the

structure parameter representing the ratio $s = z_0/w_0$ (axial/lateral radius ratio of the confocal volume).

The structural parameter was calibrated using a known concentration of Rh6G standard solution at room

temperature. The relationship between absolute concentration and calculated N was calibrated using a

known concentration of Rh6G or recombinant mGFP solution. The effect of photobleaching on evaluation

of concentration was determined according to a previous study. Fluoprescence correlation functions

(FCFs) in live cells were measured sequentially 6 times with duration of 10s to minimize photobleaching

effect.

The cytoplasmic concentration of the target protein was calculated as

$$C = \frac{N}{N_A V_{eff}}$$

640 Where N_A is Avogadro's number and V_{eff} is the effective confocal volume determined from calibration

641 measurements with Rhodamine 6G solutdion.

Yeast cell lysis and RNase inhibitor treatment

For primary culture, yeast cells were inoculated and grown for 24 h at 25°C in 10 ml of SC media (w/ 2%

glucose). After 24 h, the primary cultured cells were diluted to 0.1 optical density (O.D.) value in 100 ml of

the same media and cultured for 30 h at 25°C. 50 ml of cells were then collected in 1.8 ml bead-beating

tubes (SIMPORT, Canada, #T335-6SPR). In each tube, we added 200 μl of acid-washed 0.5 mm bead

(Biospec, USA, #11079105) and 400 μl of pH 7.4 PBS lysis buffer with 0.1% NP40 (BIO BASIC, Canada,

#NDB0385), cOmplete^{TM} mini EDTA-free Protease Inhibitor Cocktail (Roche, Switzerland,

#11836170001), 1 mM DTT (Georgiachem, USA, #DT1050), 1 mM PMSF (Georgiachem, USA,

#PS2064), and 10 mM Ribonucleoside vanadyl complex (RVC, NEW ENGLAND Biolabs, USA,

#S1402S), a transition state analog acting as a non-covalently bound ribonuclease (RNase) inhibitor.

These tubes were incubated on ice for 15 min and then lysed by the bead beater for 4 times of a cycle set

of 30 sec beating and 8 min rest. The crude extracts were centrifuged at 2000 g for 5 min at 4°C. These

separated supernatants were centrifuged again at 10,000 g for 10 min at 4°C. To obtain the pellets, all the

supernatants were carefully removed. More than 5 fluorescent images of these pellets were acquired by

fluorescent microscopy.

Fluorescence recovery after photobleaching (FRAP) in living yeast cells

Fluorescence recovery after photobleaching (FRAP) was performed on puncta induced by GAL promoter

using a Zeiss LSM900 laser scanning confocal microscope with a 100×/1.40 NA oil immersion objective

(Carl Zeiss AG, Oberkochen, Germany). Cells were mounted on agarose pads, composed of SC 2%

raffinose/ 2% galactose media and 1~3% agarose, to minimize the delicate movements of cells. For the

confocal imaging, 0.6% 488 nm wavelength laser and GaAsP-PMT detector were used. Due to the low

GFP signals of proteins and the small size of yeast, the pinhole was opened to 3 Airy units, about 240

μm. For fast imaging, the acquisition mode was set to 512 px X 512 px frame size, 8 bits per pixel, 9

(about 1 µs of pixel times) scan speed, one-way, and none averaging scanning. For the FRAP

experiments, the region of interest (ROI) was selected, and 200 cycles with no interval of time series were

set. After snapping five images, bleaching on ROI was performed repeatedly three times with the power

of 10~100% 488 nm laser beam. Bleached signals on ROI were recovered, and these signals were

acquired for the left 194 cycles. All the experiment was repeated about 20~30 foci of each strain. If the

foci were out of focus for 200 cycles, these data were discarded.

Image analysis for FRAP

The mean intensity in ROI at each cycle was analyzed and drawn as graphs using Fiji with the Stowers

plugin. All the graphs were normalized to 0-1, and the average curves of these graphs were calculated

with GraphPad Prism 10 (GraphPad Software, CA, USA). These normalized curves were fitted with Eq.1.

$$y = y_{max} \times (1 - e^{-\frac{t}{\tau}}) \tag{1}$$

In Eq.1, y_{max} is the y value at infinite time ($t \approx \infty$), and τ (tau) is the time point when the y value is equal

to 63.2% of y_{max} value. The first derivative functions of the fitted curves were analyzed with GraphPad

Prism 10 to calculate the maximum velocity of the recovery curves.

Fixation and DAPI staining of yeast cells

Following 4 hours of galactose-mediated induction, yeast cells were harvested by centrifugation at 4,000

rpm for 5 minutes at RT. The supernatant was carefully removed, and cell pellets were resuspended in

4% paraformaldehyde (PFA) dissolved in Dulbecco's phosphate-buffered saline (DPBS) (T&I, South

Korea, BPP-9004) for fixation by gentle inversion mixing for 1 h at RT. To remove residual PFA, cells

were subjected to two successive wash cycles consisting of centrifugation (4,000 rpm, 2 minutes) and

resuspension in fresh DPBS (Welgene, South Korea, LB 001-02). The final cell pellet was reconstituted in

DPBS and evenly distributed onto microscope slides. Vectashield mounting medium for fluorescence with

DAPI (Vector Laboratories, Inc., USA, H-1200) was applied to each sample, followed by gentle mixing

using a pipette tip to ensure homogeneous distribution of cells. Cellular morphology and fluorescence

signals were subsequently analyzed using a Axioplan2 microscope (Carl Zeiss, Jena, Germany)

equipped with a DAPI-specific filter set.

U2OS cell culture and transfection

U2OS cells (ATCC, CRL-1427, HTB 96, Manassas, VA, USA) were cultured in Dulbecco's Modified Eagle Medium (DMEM with high glucose; Cytiva, USA SH30023.01) supplemented with 10% fetal bovine serum (FBS; Cytiva, USA, SH30088.03) and 1X antibiotic-antimycotic solution (Anti-Anti; Gibco, USA ,15240-062) at 37°C in a humidified incubator with 5% CO_2. Once the cells reached 70% confluence, they were transfected with pcDNA3.1-myc-hisA expression vectors encoding GFP-TDP-43 and its IDR deletion mutants using Lipofectamine 3000 (Invitrogen, USA, L300015), following the manufacturer's protocol. After transfection, the cells were selected with 600 ng/µL Geneticin™ Selective Antibiotic (G418 Sulfate)（Gibco, USA, 10131035）.

Fixation and DAPI staining of U2OS cells

Forty-eight hours after transfection, U2OS cells cultured on a cover slide were fixed with 4% paraformaldehyde (PFA, T&I, South Korea, BPP-9004) at RT for 1 h, washed three times with phosphate-buffered saline (PBS, WELGENE, South Korea, LB 001-02)., and mounted using a DAPI-containing mounting solution (Vector Laboratories, Inc., USA, H-1200). Fluorescence images were acquired using an Axioplan2 fluorescence microscope (Carl Zeiss, Jena, Germany) equipped with a 20x/0.5 Ph2 Plan-Neofluar objective.

Western blotting analysis

Protein extraction from yeast cells was obtained using the alkaline lysis as described in Choi and Song（*Int. J. Mol. Sci. 2022*, *23*, 2501）U2OS cell extract was obtained as previously described (PLOS ONE | https://doi.org/10.1371/journal.pone.0302936 May 7, 2024). Protein samples were separated by sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS-PAGE) and transferred onto polyvinylidene difluoride (PVDF) membranes (Merck Millipore, Burlington, MA, USA) using a semi-dry transfer system. Anti-GFP antibody (600-101-215 ROCKLAND, Limerick, PA, USA, 1:1000 dilution) was used for Western blotting. To confirm equal protein loading, the membranes were stripped of the anti-GFP antibody prior to re-probing with an anti-Tubulin antibody anti-Tubulin (Mouse, Sigma-Aldrich, Cat# T5168, 1:10000

716 dilution) and subsequently with HRP-conjugated anti-mouse secondary antibody (Santa Cruz

717 Biotechnology, Cat# sc-2005, 1:10000 dilution).

QUANTIFICATION AND STATISTICAL ANALYSIS

719 Only statistically analyzed data are shown as mean ± SEM. Statistical analyses were performed using

720 GraphPad Prism. Statistical significance was determined using unpaired two-tailed Student's t-test with

721 Welch's correction. $P < 0.05$ was considered statistically significant.

DATA AND CODE AVAILABILITY

723 Custom algorithms were implemented in Python 3.10 by the authors. The analysis pipeline included

724 modules written with NumPy and SciPy and was used to process experimental data shown in Figure 1.

725 The code used in this study was written by the authors and is available from the corresponding author

726 upon reasonable request.

ADDITIONAL RESOURCES

728 N/A

ACKNOWLEDGMENTS

We thank the Cellular Imaging Core Facility at the ConveRgence mEDIcine research cenTer (CREDIT),

Asan Medical Center, for technical support. The authors would like to thank Ewha Women's University

Fluorescence Core Imaging Center for valuable guidance in FRAP analysis. We specially appreciate the

technical assistance provided by professor Dongmin Kang, the Director, and Research Technician Sinae

Jang, of Ewha Fluorescence Core Imaging Center. TDP-43 gene clone was kindly provided by Dr.

Hyung-Jun Kim in Korea Brain Research Institute. This work was in part supported by the National

Research Foundation of Korea (NRF) grant provided to K. Song (RS-2023-00243284).

AUTHOR CONTRIBUTIONS

740   Conceptualization, K.S. and Y.J.C.; methodology, K.S., Y.J.C, and C.P.; Investigation, Y.J.C, Y.L., S.C.,

741   and Y.P.; writing—original draft, K.S. and Y.J.C.; writing—review & editing, K.S., Y.J.C, C.P, and Y.J.;

742   funding acquisition, K.S.; resources, K.S. and C.K.B.; supervision, K.S., and Y.J.C

DECLARATION OF INTERESTS

745   The authors (K.S., Y.J.C., S.C., and Y.P.) have applied for a patent entitled "Specific Amino Acid

746   Sequence Deletion Mutants for Inhibiting TDP-43 Aggregation Causing Degenerative Brain Diseases,

747   and Method for Identifying Mutant Sequences (DP 2025-0349)" with the Korean Intellectual Property

748   Office. The authors declare no other competing interests.

REFERENCES

Ahmad, A., Uversky, V.N., and Khan, R.H. (2022). Aberrant liquid-liquid phase separation and amyloid
aggregation of proteins related to neurodegenerative diseases. Int J Biol Macromol *220*, 703-720.
10.1016/j.ijbiomac.2022.08.132.
Alberti, S., and Hyman, A.A. (2021). Biomolecular condensates at the nexus of cellular stress, protein
aggregation disease and ageing. Nat Rev Mol Cell Biol *22*, 196-213. 10.1038/s41580-020-00326-6.
Alshareedah, I., Borcherds, W.M., Cohen, S.R., Singh, A., Posey, A.E., Farag, M., Bremer, A., Strout, G.W.,
Tomares, D.T., Pappu, R.V., et al. (2024). Sequence-specific interactions determine viscoelasticity and
aging dynamics of protein condensates. Nat Phys *20*, 1482-1491. 10.1038/s41567-024-02558-1.
Alshareedah, I., Moosa, M.M., Pham, M., Potoyan, D.A., and Banerjee, P.R. (2021). Programmable
viscoelasticity in protein-RNA condensates with disordered sticker-spacer polypeptides. Nat Commun
12, 6620. 10.1038/s41467-021-26733-7.
Banani, S.F., Afeyan, L.K., Hawken, S.W., Henninger, J.E., Dall'Agnese, A., Clark, V.E., Platt, J.M., Oksuz,
O., Hannett, N.M., Sagi, I., et al. (2022). Genetic variation associated with condensate dysregulation in
disease. Dev Cell *57*, 1776-1788 e1778. 10.1016/j.devcel.2022.06.010.
Banani, S.F., Lee, H.O., Hyman, A.A., and Rosen, M.K. (2017). Biomolecular condensates: organizers of
cellular biochemistry. Nat Rev Mol Cell Biol *18*, 285-298. 10.1038/nrm.2017.7.
Banerjee, D.S., Chigumira, T., Lackner, R.M., Kratz, J.C., Chenoweth, D.M., Banerjee, S., and Zhang, H.
(2024). Interplay of condensate material properties and chromatin heterogeneity governs nuclear
condensate ripening. bioRxiv. 10.1101/2024.05.07.593010.
Banjade, S., and Rosen, M.K. (2014). Phase Transitions of Multivalent Proteins Can Promote Clustering of
Membrane Receptors. Elife *3*. ARTN e04123

10.7554/eLife.04123.
Biswas, S., and Potoyan, D.A. (2024). Decoding Biomolecular Condensate Dynamics: An Energy
Landscape Approach. bioRxiv. 10.1101/2024.09.24.614805.

Borcherds, W., Bremer, A., Borgia, M.B., and Mittag, T. (2021). How do intrinsically disordered protein
regions encode a driving force for liquid-liquid phase separation? Curr Opin Struc Biol *67*, 41-50.
10.1016/j.sbi.2020.09.004.
Chen, C., Yamanaka, Y., Ueda, K., Li, P., Miyagi, T., Harada, Y., Tezuka, S., Narumi, S., Sugimoto, M.,
Kuroda, M., et al. (2021). Phase separation and toxicity of C9orf72 poly(PR) depends on alternate
distribution of arginine. J Cell Biol *220*. 10.1083/jcb.202103160.
Chien, H.M., Lee, C.C., and Huang, J.J. (2021). The Different Faces of the TDP-43 Low-Complexity
Domain: The Formation of Liquid Droplets and Amyloid Fibrils. Int J Mol Sci *22*. 10.3390/ijms22158213.
Choi, C.H., Lee, D.S.W., Sanders, D.W., and Brangwynne, C.P. (2024). Condensate interfaces can
accelerate protein aggregation. Biophysical Journal *123*, 1404-1413. 10.1016/j.bpj.2023.10.009.
Choi, J.M., Holehouse, A.S., and Pappu, R.V. (2020). Physical Principles Underlying the Complex Biology
of Intracellular Phase Transitions. Annu Rev Biophys *49*, 107-133. 10.1146/annurev-biophys-121219-
081629.
Choi, Y.J., Lee, Y., Lin, Y., Heo, Y., Lee, Y.H., and Song, K. (2022). The Multivalent Polyampholyte Domain
of Nst1, a P-Body-Associated Saccharomyces cerevisiae Protein, Provides a Platform for Interacting with
P-Body Components. Int J Mol Sci *23*. 10.3390/ijms23137380.
Cohen, L.S., Mikhli, C., Jiao, X., Kiledjian, M., Kunkel, G., and Davis, R.E. (2005). Dcp2 Decaps
m2,2,7GpppN-capped RNAs, and its activity is sequence and context dependent. Mol Cell Biol *25*, 8779-
8791. 10.1128/MCB.25.20.8779-8791.2005.
Dickson, Z.W., and Golding, G.B. (2022). Low Complexity Regions in Mammalian Proteins are Associated
with Low Protein Abundance and High Transcript Abundance. Mol Biol Evol *39*.
10.1093/molbev/msac087.
Fondon, J.W., 3rd, and Garner, H.R. (2004). Molecular origins of rapid and continuous morphological
evolution. Proc Natl Acad Sci U S A *101*, 18058-18063. 10.1073/pnas.0408118101.
Fushimi, K., Long, C., Jayaram, N., Chen, X., Li, L., and Wu, J.Y. (2011). Expression of human FUS/TLS in
yeast leads to protein aggregation and cytotoxicity, recapitulating key features of FUS proteinopathy.
Protein Cell *2*, 141-149. 10.1007/s13238-011-1014-5.
Gao, G., Sumrall, E.S., Pitchiaya, S., Bitzer, M., Alberti, S., and Walter, N.G. (2023). Biomolecular
condensates in kidney physiology and disease. Nat Rev Nephrol *19*, 756-770. 10.1038/s41581-023-
00767-0.
Gruijs da Silva, L.A., Simonetti, F., Hutten, S., Riemenschneider, H., Sternburg, E.L., Pietrek, L.M., Gebel,
J., Dotsch, V., Edbauer, D., Hummer, G., et al. (2022). Disease-linked TDP-43 hyperphosphorylation
suppresses TDP-43 condensation and aggregation. EMBO J *41*, e108443. 10.15252/embj.2021108443.
Hurtle, B.T., Xie, L., and Donnelly, C.J. (2023). Disrupting pathologic phase transitions in
neurodegeneration. J Clin Invest *133*. 10.1172/JCI168549.
Johnson, B.S., McCaffery, J.M., Lindquist, S., and Gitler, A.D. (2008). A yeast TDP-43 proteinopathy
model: Exploring the molecular determinants of TDP-43 aggregation and cellular toxicity. Proc Natl Acad
Sci U S A *105*, 6439-6444. 10.1073/pnas.0802082105.
Kawai-Noma, S., Ayano, S., Pack, C.G., Kinjo, M., Yoshida, M., Yasuda, K., and Taguchi, H. (2006).
Dynamics of yeast prion aggregates in single living cells. Genes Cells *11*, 1085-1096. 10.1111/j.1365-
2443.2006.01004.x.
Kellett, E.A., Bademosi, A.T., and Walker, A.K. (2025). Molecular mechanisms and consequences of TDP-
43 phosphorylation in neurodegeneration. Mol Neurodegener *20*, 53. 10.1186/s13024-025-00839-8.
Lange, S.S., and Vasquez, K.M. (2009). HMGB1: the jack-of-all-trades protein is a master DNA repair
mechanic. Mol Carcinog *48*, 571-580. 10.1002/mc.20544.
Lee, D.S.W., Wingreen, N.S., and Brangwynne, C.P. (2021). Chromatin Mechanics Dictates Subdiffusion
and Coarsening Dynamics of Embedded Condensates. Biophysical Journal *120*, 318a-318a.

Li, Q., Babinchak, W.M., and Surewicz, W.K. (2021). Cryo-EM structure of amyloid fibrils formed by the
entire low complexity domain of TDP-43. Nat Commun *12*, 1620. 10.1038/s41467-021-21912-y.
Linsenmeier, M., Faltova, L., Morelli, C., Capasso Palmiero, U., Seiffert, C., Kuffner, A.M., Pinotsi, D.,
Zhou, J., Mezzenga, R., and Arosio, P. (2023). The interface of condensates of the hnRNPA1 low-
complexity domain promotes formation of amyloid fibrils. Nat Chem *15*, 1340-1349. 10.1038/s41557-
023-01289-9.
Linsenmeier, M., Kopp, M.R.G., Stavrakis, S., de Mello, A., and Arosio, P. (2021). Analysis of biomolecular
condensates and protein phase separation with microfluidic technology. Biochim Biophys Acta Mol Cell
Res *1868*, 118823. 10.1016/j.bbamcr.2020.118823.
Lu, B., Qiu, R., Wei, J., Wang, L., Zhang, Q., Li, M., Zhan, X., Chen, J., Hsieh, I.Y., Yang, C., et al. (2024).
Phase separation of phospho-HDAC6 drives aberrant chromatin architecture in triple-negative breast
cancer. Nat Cancer. 10.1038/s43018-024-00816-y.
Lu, S., Hu, J., Arogundade, O.A., Goginashvili, A., Vazquez-Sanchez, S., Diedrich, J.K., Gu, J., Blum, J.,
Oung, S., Ye, Q., et al. (2022). Heat-shock chaperone HSPB1 regulates cytoplasmic TDP-43 phase
separation and liquid-to-gel transition. Nat Cell Biol *24*, 1378-1393. 10.1038/s41556-022-00988-8.
Mehta, S., and Zhang, J. (2022). Liquid-liquid phase separation drives cellular function and dysfunction in
cancer. Nat Rev Cancer *22*, 239-252. 10.1038/s41568-022-00444-7.
Mensah, M.A., Niskanen, H., Magalhaes, A.P., Basu, S., Kircher, M., Sczakiel, H.L., Reiter, A.M.V., Elsner,
J., Meinecke, P., Biskup, S., et al. (2023). Aberrant phase separation and nucleolar dysfunction in rare
genetic diseases. Nature *614*, 564-571. 10.1038/s41586-022-05682-1.
Miyagi, T., Yamazaki, R., Ueda, K., Narumi, S., Hayamizu, Y., Uji, I.H., Kuroda, M., and Kanekura, K.
(2022). The Patterning and Proportion of Charged Residues in the Arginine-Rich Mixed-Charge Domain
Determine the Membrane-Less Organelle Targeted by the Protein. Int J Mol Sci *23*.
10.3390/ijms23147658.
Nam, J., and Gwon, Y. (2023). Neuronal biomolecular condensates and their implications in
neurodegenerative diseases. Front Aging Neurosci *15*, 1145420. 10.3389/fnagi.2023.1145420.
Niedner-Boblenz, A., Monecke, T., Hennig, J., Klostermann, M., Hofweber, M., Davydova, E., Gerber,
A.P., Anosova, I., Mayer, W., Muller, M., et al. (2024). Intrinsically disordered RNA-binding motifs
cooperate to catalyze RNA folding and drive phase separation. Nucleic Acids Res *52*, 14205-14228.
10.1093/nar/gkae1107.
Niu, X., Zhang, L., Wu, Y., Zong, Z., Wang, B., Liu, J., Zhang, L., and Zhou, F. (2023). Biomolecular
condensates: Formation mechanisms, biological functions, and therapeutic targets. MedComm (2020) *4*,
e223. 10.1002/mco2.223.
Oiwa, K., Watanabe, S., Onodera, K., Iguchi, Y., Kinoshita, Y., Komine, O., Sobue, A., Okada, Y., Katsuno,
M., and Yamanaka, K. (2023). Monomerization of TDP-43 is a key determinant for inducing TDP-43
pathology in amyotrophic lateral sclerosis. Sci Adv *9*, eadf6895. 10.1126/sciadv.adf6895.
Outeiro, T.F., and Lindquist, S. (2003). Yeast cells provide insight into alpha-synuclein biology and
pathobiology. Science *302*, 1772-1775. 10.1126/science.1090439.
Pack, C.G. (2021). Application of quantitative cell imaging using label-free optical diffraction
tomography. Biophys Physicobiol *18*, 244-253. 10.2142/biophysico.bppb-v18.027.
Pack, C.G., Yukii, H., Toh-e, A., Kudo, T., Tsuchiya, H., Kaiho, A., Sakata, E., Murata, S., Yokosawa, H.,
Sako, Y., et al. (2014). Quantitative live-cell imaging reveals spatio-temporal dynamics and cytoplasmic
assembly of the 26S proteasome. Nat Commun *5*, 3396. 10.1038/ncomms4396.
Ranganathan, S., Liu, J., and Shakhnovich, E. (2022). Different states and the associated fates of
biomolecular condensates. Essays Biochem *66*, 849-862. 10.1042/EBC20220054.
Regy, R.M., Dignon, G.L., Zheng, W., Kim, Y.C., and Mittal, J. (2020). Sequence dependent phase
separation of protein-polynucleotide mixtures elucidated using molecular simulations. Nucleic Acids Res
48, 12593-12603. 10.1093/nar/gkaa1099.

Saar, K.L., Morgunov, A.S., Qi, R., Arter, W.E., Krainer, G., Lee, A.A., and Knowles, T.P.J. (2021). Learning
the molecular grammar of protein condensates from sequence determinants and embeddings. Proc Natl
Acad Sci U S A *118*. 10.1073/pnas.2019053118.
Saitoh, N., Spahr, C.S., Patterson, S.D., Bubulya, P., Neuwald, A.F., and Spector, D.L. (2004). Proteomic
analysis of interchromatin granule clusters. Mol Biol Cell *15*, 3876-3890. 10.1091/mbc.e04-03-0253.
Sanfeliu-Cerdan, N., Catala-Castro, F., Mateos, B., Garcia-Cabau, C., Ribera, M., Ruider, I., Porta-de-la-
Riva, M., Canals-Calderon, A., Wieser, S., Salvatella, X., and Krieg, M. (2023). A MEC-2/stomatin
condensate liquid-to-solid phase transition controls neuronal mechanotransduction during touch
sensing. Nat Cell Biol *25*, 1590-1599. 10.1038/s41556-023-01247-0.
Schneider-Poetsch, T., Ju, J., Eyler, D.E., Dang, Y., Bhat, S., Merrick, W.C., Green, R., Shen, B., and Liu, J.O.
(2010). Inhibition of eukaryotic translation elongation by cycloheximide and lactimidomycin. Nat Chem
Biol *6*, 209-217. 10.1038/nchembio.304.
Shen, Y., Chen, A., Wang, W., Shen, Y., Ruggeri, F.S., Aime, S., Wang, Z., Qamar, S., Espinosa, J.R.,
Garaizar, A., et al. (2023). The liquid-to-solid transition of FUS is promoted by the condensate surface.
Proc Natl Acad Sci U S A *120*, e2301366120. 10.1073/pnas.2301366120.
Shenoy, J., Lends, A., Berbon, M., Bilal, M., El Mammeri, N., Bertoni, M., Saad, A., Morvan, E., Grelard, A.,
Lecomte, S., et al. (2023). Structural polymorphism of the low-complexity C-terminal domain of TDP-43
amyloid aggregates revealed by solid-state NMR. Front Mol Biosci *10*, 1148302.
10.3389/fmolb.2023.1148302.
Sim, K.L., and Creamer, T.P. (2002). Abundance and distributions of eukaryote protein simple sequences.
Mol Cell Proteomics *1*, 983-995. 10.1074/mcp.m200032-mcp200.
Sneideriene, G., Gonzalez Diaz, A., Adhikari, S.D., Wei, J., Michaels, T., Sneideris, T., Linse, S.,
Vendruscolo, M., Garai, K., and Knowles, T.P.J. (2025). Lipid-induced condensate formation from the
Alzheimer's Abeta peptide triggers amyloid aggregation. Proc Natl Acad Sci U S A *122*, e2401307122.
10.1073/pnas.2401307122.
Sood, A., and Zhang, B. (2024). Preserving condensate structure and composition by lowering sequence
complexity. Biophys J *123*, 1815-1826. 10.1016/j.bpj.2024.05.026.
Sun, Z., Diaz, Z., Fang, X., Hart, M.P., Chesi, A., Shorter, J., and Gitler, A.D. (2011). Molecular
determinants and genetic modifiers of aggregation and toxicity for the ALS disease protein FUS/TLS.
PLoS Biol *9*, e1000614. 10.1371/journal.pbio.1000614.
Sundaravadivelu Devarajan, D., Wang, J., Szala-Mendyk, B., Rekhi, S., Nikoubashman, A., Kim, Y.C., and
Mittal, J. (2024). Sequence-dependent material properties of biomolecular condensates and their
relation to dilute phase conformations. Nat Commun *15*, 1912. 10.1038/s41467-024-46223-w.
Teixeira, D., Sheth, U., Valencia-Sanchez, M.A., Brengues, M., and Parker, R. (2005). Processing bodies
require RNA for assembly and contain nontranslating mRNAs. RNA *11*, 371-382. 10.1261/rna.7258505.
Tibble, R.W., Depaix, A., Kowalska, J., Jemielity, J., and Gross, J.D. (2021). Biomolecular condensates
amplify mRNA decapping by biasing enzyme conformation. Nat Chem Biol *17*, 615-623. 10.1038/s41589-
021-00774-x.
Ukmar-Godec, T., Hutten, S., Grieshop, M.P., Rezaei-Ghaleh, N., Cima-Omori, M.S., Biernat, J.,
Mandelkow, E., Soding, J., Dormann, D., and Zweckstetter, M. (2019). Lysine/RNA-interactions drive and
regulate biomolecular condensation. Nat Commun *10*, 2909. 10.1038/s41467-019-10792-y.
Van Treeck, B., and Parker, R. (2018). Emerging Roles for Intermolecular RNA-RNA Interactions in RNP
Assemblies. Cell *174*, 791-802. 10.1016/j.cell.2018.07.023.
Visser, B.S., Lipinski, W.P., and Spruijt, E. (2024). The role of biomolecular condensates in protein
aggregation. Nat Rev Chem *8*, 686-700. 10.1038/s41570-024-00635-w.
Wang, A., Conicella, A.E., Schmidt, H.B., Martin, E.W., Rhoads, S.N., Reeb, A.N., Nourse, A., Ramirez
Montero, D., Ryan, V.H., Rohatgi, R., et al. (2018a). A single N-terminal phosphomimic disrupts TDP-43
polymerization, phase separation, and RNA splicing. EMBO J *37*. 10.15252/embj.201797452.

Wang, B., Zhang, L., Dai, T., Qin, Z., Lu, H., Zhang, L., and Zhou, F. (2021). Liquid-liquid phase separation
in human health and diseases. Signal Transduct Target Ther *6*, 290. 10.1038/s41392-021-00678-1.
Wang, J., Choi, J.M., Holehouse, A.S., Lee, H.O., Zhang, X., Jahnel, M., Maharana, S., Lemaitre, R.,
Pozniakovsky, A., Drechsel, D., et al. (2018b). A Molecular Grammar Governing the Driving Forces for
Phase Separation of Prion-like RNA Binding Proteins. Cell *174*, 688-699 e616. 10.1016/j.cell.2018.06.006.
Wang, X., Bigman, L.S., Greenblatt, H.M., Yu, B., Levy, Y., and Iwahara, J. (2023). Negatively charged,
intrinsically disordered regions can accelerate target search by DNA-binding proteins. Nucleic Acids Res
51, 4701-4712. 10.1093/nar/gkad045.

Figure 1

Figure 1. *Overexpression of Nst1 and Edc3 leads to mature condensate formation, whereas*

overexpression of their polyelectrolyte (PE) domain deletion mutants within the IDR maintains a

stabilized microphase-separated state.

(A) Schematic representation of budding yeast overexpression system to observe intracellular

condensates.

(B) Diagram of puncta generation pattern of cells overexpressing GFP tagged Nst1 for 2 h (green) and 4

h (purple). *Δnst1* (YSK3490) cells transformed with pMW20-pGAL-GFP-NST1 were induced for 2 hours

(green) and 4 hours (purple) under the GAL promoter. Cells were imaged and analyzed; mean integrated

density and the number of puncta of each cell were measured and plotted.

(C-D) Representative fluorescence microscopy images and quantitative phase separation analysis of

cells overexpressing GFP and GFP-tagged Nst1, Edc3, Dcp3, Dhh1, and Pab1. (C) Scale bar: 10 μm. (D)

Cytoplasmic protein concentrations were measured via fluorescence correlation spectroscopy (FCS) as

described in Materials and Methods. The puncta formation patterns in cells were classified into three

categories: diffuse phase, microphase separation, and mature condensates. Box plot analysis of the

correlation between the pattern of protein puncta formation and cytoplasmic concentration was presented.

(E) Schematic representation of polyelectrolyte (PE) domain segmentation.

(F) Sequence diagram of Nst1 and Edc3. The red box indicates the polyelectrolyte (PE) region, while the

blue box represents the aggregation-prone domain (APD). Each sequence shown corresponds to the PE

region. Positively charged amino acids, lysine (K) and arginine (R),, are highlighted in red and marked

with a (+) sign below, whereas negatively charged amino acids, glutamic acid (E) and aspartic acid (D),

are highlighted in blue and marked with a (−) sign below.

(G) Fluorescence microscopy images showing puncta formation in cells overexpressing Edc3 and each

domain deletion mutant after 4 and 24 hours of galactose induction. In *Δedc3* cells (YSK3707), GFP-

tagged full-length Edc3 (pMW20-pGAL-GFP-EDC3), Edc3ΔPE (pMW20-pGAL-GFP-EDC3ΔPE), and

Edc3 Δ APD (pMW20-pGAL-GFP-EDC3ΔAPD) were overexpressed. Representative images were

selected from an average of more than 100 cells per strain at each time point. Scale bar: 5 μm.

(H) Fluorescence microscopy images showing puncta formation in cells expressing Nst1 and each Nst1

domain deletion mutant after 4 and 24 hours of induction as in (E). GFP-tagged full-length Nst1 (pMW20-

pGAL-GFP-NST1), Nst1ΔPE (pMW20-pGAL-GFP-NST1ΔPE), and Nst1ΔAPD (pMW20-pGAL-GFP-

NST1ΔAPD) were overexpressed in *Δnst1* (YSK3490) cells. Representative images were selected from

an average of more than 100 cells per strain at each time point. Scale bar: 5 μm.

(I) Box plot analysis of the correlation between the puncta formation pattern and cytoplasmic

concentration of the Edc3 and Edc3 derivatives. The puncta formation patterns in cells were classified

into three categories: diffuse phase, microphase separation, and mature condensates. Cytoplasmic

concentrations were measured in cells expressing each Edc3 domain deletion mutant after 4 hours of

induction by fluorescence correlation spectroscopy (FCS) as described in Materials and Methods. The

measured cytoplasmic concentrations of cells were analyzed with respect to puncta formation patterns.

(J) Box plot analysis of the correlation between the puncta formation pattern and cytoplasmic

concentration of the Nst1 and Nst1 derivatives. The cytoplasmic protein concentration measurement and

puncta formation analysis were performed same as (F).

Figure 2

A

Nst1

B

C

	GFP-Nst1	GFP-Nst1ΔPA	GFP-Nst1ΔPAM	GFP-Nst1ΔPE	GFP-Nst1ΔAPD
30 min					
45 min					
60 min					
75 min					
90 min					
110 min					
130 min					
4 h					
5 h					
24 h					

D

GFP-Nst1

+1,6-Hexanediol

F.L.

ΔPE

ΔPA

ΔPAM

E

F

G

H

I

J

Figure 2. *Polyanion (PA) domain of the Polyelectrolyte (PE) region specifically promotes the*

coalescence of Nst1 condensates, whereas the polyampholyte (PAM) domain in the PE region

does not.

(A) Schematic representation of Polyanion (PA) and Polyampholyte (PAM) domain segmentation with a

sliding window approach.

(B) Diagram of Nst1 with its domain architectures and sequences. The black line indicates the PE region,

while the green, red, and blue boxes denote the PA, PAM, and APD regions, respectively. The amino

acid sequences of the PA and PAM regions are shown below the diagram. Charges were annotated as

described in Figure 1D.

(C) Fluorescence microscopy images of condensate formation in cells overexpressing each GFP-tagged

Nst1 and its domain deletion mutants. Expression of GFP-tagged full-length Nst1 (pMW20-pGAL-GFP-

NST1) or NST1 lacking specific domains - Nst1ΔPE (pMW20-pGAL-GFP-NST1ΔPE), Nst1ΔPA (pMW20-

pGAL-GFP-NST1ΔPA), Nst1ΔPAM (pMW20-pGAL-GFP-NST1ΔPAM), and Nst1ΔAPD (pMW20-pGAL-

GFP-NST1ΔAPD) - was induced in *Δnst1* (YSK3490) cells. Each domain deletion mutant was

overexpressed by galactose induction for the indicated time periods. At each time point, fluorescence

images were collected from at least 100 cells to ensure representativeness. Scale bar: 5 μm.

(D) Fluorescence microscopy images of GFP-tagged Nst1 condensates induced for 4 hours, followed by

a 1-hour treatment with 1,6-hexanediol. Scale bar: 5 μm.

(E) Box plot analysis of the correlation between the puncta formation pattern and cytoplasmic

concentration of cells overexpressing GFP-tagged Nst1, Nst1ΔPA, Nst1ΔPAM, Nst1ΔPE, and Nst1ΔAPD.

Cytoplasmic protein concentrations were measured via fluorescence correlation spectroscopy (FCS). The

puncta formation patterns in cells were classified into three categories: diffuse phase, microphase

separation, and mature condensates.

(F-J) Biomolecular condensate formation of endogenously expressed GFP-tagged Nst1, Nst1ΔPA,

Nst1ΔPAM and Nst1ΔAPD during chronological aging in yeast. (F) The growth phase diagram of general

budding yeast with observation time points designated. The inverted triangles (▾) on the x-axis represent

the observed time points (10, 14, 18, 22, 26, and 46 hours). (G) Line chart showing the optical densities

(O.D.600) of cell cultures endogenously expressing GFP-tagged Nst1, Nst1ΔPA, Nst1ΔPAM and

Nst1ΔAPD under native NST1 promoter. The growth of *Δnst1* (YSK3490) cells integrated with pRS306-

pNST1-GFP-NST1 full-length and various NST1 deletion mutants (YSK3700, YSK3701, YSK3702,

YSK3714) was monitored over time. The 10 h,14 h and 18 h time points represent the log phase, 22 h the

diauxic shift, and 26 h and 46 h the stationary phase. (H) Condensates were segmented using Fiji

(ImageJ) and visualized in green. Shown are GFP images merged with corresponding phase-contrast

images. Scale bar: 10 μm. (I) Bar plot showing the average integrated GFP fluorescence density per cell

for each yeast strain expressing full-length Nst1 and its deletion mutants (Nst1ΔPA, Nst1ΔPAM,

Nst1ΔAPD), as shown in (H). For each condition, more than 300 cells were analyzed using Fiji (ImageJ).

Results are presented using GraphPad Prism 10. (J) Box plot showing the integrated GFP intensity of

condensates in stationary-phase (46 h) yeast cells expressing Nst1, Nst1ΔPA and Nst1ΔPAM. For each

condition, more than 300 cells were quantified using Fiji (ImageJ). Data from three independent biological

replicates were plotted with GraphPad Prism 10.

Figure 3

A

B

C

1012

1013

1014   **Figure 3. *Polyanion (PA) domain exclusively functions in coalescence of Nst1 condensates.***

1015

(A) A diagram of Nst1 with domain architectures. Residues 2-122 were designated as D1, 123-293 as D2,

294-459 as D3, 753-844 as D4, and 845-1015 as D5. The lengths of D1-D5 are similar to that of PA, and

all of the domains are comparable in size.

(B) Representative fluorescence microscopy images comparing time-dependent condensate formation in

the PA deletion mutant and other domain deletion mutants (ΔD1-D5, ΔAPD). In *Δnst1* (YSK3490) cells,

GFP-tagged Nst1ΔPA (pMW20-pGAL-GFP-NST1ΔPA), Nst1ΔD1 (pMW20-pGAL-GFP-NST1ΔD1),

Nst1ΔD2 (pMW20-pGAL-GFP-NST1ΔD2), Nst1ΔD3 (pMW20-pGAL-GFP-NST1ΔD3), Nst1ΔD4 (pMW20-

pGAL-GFP-NST1ΔD4), Nst1ΔD5 (pMW20-pGAL-GFP-NST1ΔD5), and Nst1ΔAPD (pMW20-pGAL-GFP-

NST1ΔAPD) were overexpressed. Overexpression of each Nst1 derivative was induced by galactose,

and images were obtained at the designated time point after the induction. Representative images were

selected among more than 100 cells. Scale bar: 5 μm.

(C) Box plot analysis showing the correlation between puncta formation patterns and cytoplasmic

concentrations in the cells overexpressing GFP-tagged Nst1, Nst1ΔPA, NstΔD1-ΔD5, and Nst1ΔAPD

after 4 hours of induction. Cytoplasmic protein concentrations were measured via fluorescence correlation

spectroscopy (FCS). The puncta formation patterns in cells were classified into three categories: diffuse

phase, microphase separation, and mature condensates.

1032

Figure 4

A

	Phase	GFP
GFP		
GFP-Nst1		
GFP-Nst1ΔPA		
GFP-Nst1ΔPAM		
GFP-Nst1ΔD1		

Δnst1

B

Pre-bleach   Post-bleach   Fluorescence recovery

Nst1

Nst1ΔPA

Nst1ΔPAM

Nst1ΔD1

C

$$y = y_{max} \times (1 - e^{-\frac{t}{\tau}})$$

D

Nst1

Nst1ΔPA

Nst1ΔPAM

Nst1ΔD1

E

Nst1
Nst1ΔPA
Nst1ΔPAM
Nst1ΔD1
GFP

F

Max velocity of the FRAP curve

Nst1, Nst1ΔPA, Nst1ΔPAM, Nst1ΔD1, GFP

**** ns

G

half-life

Nst1, Nst1ΔPA, Nst1ΔPAM, Nst1ΔD1, GFP

**** ns

H

AUC (%)

Nst1, Nst1ΔPA, Nst1ΔPAM, Nst1ΔD1, GFP

* ns

1033

1034 **Figure 4.** *The polyanion (PA) domain of Nst1 enhances intermolecular interaction, whereas the*

1035 *polyampholyte (PAM) domain increases the rigidity of Nst1 condensates.*

1036

(A-H) GAL promoter induction was performed for *Δnst1* (YSK3490) cells transformed with pMW20-pGAL -

GFP-NST1, pMW20-pGAL-GFP-NST1ΔPA*,* pMW20-pGAL-GFP-NST1ΔPAM*,* pMW20-pGAL-GFP-

NST1ΔD1*.* Fluorescence Recovery After Photobleaching (FRAP) experiments were performed on

induced condensates of these cells as described in Materials and Methods.

(A) Representative fluorescence microscopy images of cells overexpressing GFP-tagged full-length Nst1,

Nst1ΔPA, Nst1ΔPAM, Nst1ΔD1, and GFP alone (control) in the *Δnst1* strain. Scale bar: 10 μm.

(B) Representative fluorescence microscopy images at pre-beach point (cycle 1 of 200), post-bleach

(cycle 6 of 200), and fluorescence recovery (cycle 200 of 200) during FRAP analysis. The stitched lines

indicate the region of interest (ROI) selected for bleaching the puncta in cells. All experiments were

performed with more than 20 puncta each. Scale bar: 1 μm.

(C) Predicted FRAP recovery curve modeled as a single exponential function. t is time, y_{max} is the y

value at infinite time $(t \approx \infty)$, τ (tau) is the time point when the y value is equal to 63.2% of y_{max} value,

and c is half-life.

(D) The average FRAP recovery curves of Nst1 and its mutants shown in (B). Each graph for full-length

Nst1, Nst1ΔPA, Nst1ΔPAM, and Nst1ΔD1. The x-axis of these average graphs represents the time (sec)

taken in 200 cycles, and the y-axis indicates the relative fluorescence intensity.

(E) The average FRAP recovery curves for Nst1 and all Nst1 mutants in (D) were plotted together for

direct comparison. Full-length Nst1 (black line), Nst1ΔPA (pink dotted line), Nst1ΔPAM (blue dotted line),

Nst1ΔD1 (black dotted line), and GFP (grey line) as a control. The x-axis and y-axis represent the same

values as in (D).

(F) The maximum recovery rate of the FRAP curves for condensates of full-length Nst1, Nst1ΔPA,

Nst1ΔPAM, and Nst1ΔD1. Akima spline interpolation was applied to all FRAP curves. For each Nst1

mutant, the first derivative (slope) at the point of maximal recovery rate was calculated using GraphPad

Prism 10. The y-axis represents the maximum recovery rate (maximum slope) of the fitted FRAP curve.

Statistical significance was determined using unpaired two-tailed Student's t-test (**** $p<0.0001$) with

Welch's correction. $P < 0.05$ was considered statistically significant.

(G) The half-lives of FRAP curves for condensates of full-length Nst1, Nst1ΔPA, Nst1ΔPAM, and

Nst1ΔD1. The x-values corresponding to half-maximal recovery ($y = 0.5$) were calculated and averaged

using GraphPad Prism 10. Statistical significance was determined using unpaired two-tailed Student's t-

test (**** $p<0.0001$) with Welch's correction. $P < 0.05$ was considered statistically significant.

(H) The area under the curve (AUC) of the FRAP recovery curves for condensates of full-length Nst1,

Nst1ΔPA, Nst1ΔPAM, and Nst1ΔD1. The AUC was calculated for each normalized FRAP recovery curve

using GraphPad Prism 10. Statistical significance was determined using unpaired two-tailed Student's t-

test (* $p<0.01$) with Welch's correction. $P < 0.05$ was considered statistically significant.

1071

Figure 5

1073 **Figure 5.** *The PAM domain is an RNA-independent multivalent interaction domain for interaction*

1074 *with the P-body component Dcp2.*

1075

(A) Schematic representation of the experiments (B-C) designed to examine the effect of RNA on P-body

condensate formation of Nst1 and its PA and PAM deletion mutants.

(B) High-resolution confocal microscopy images showing the formation of mKate2-tagged Dcp2

condensates induced by condensates of GFP-tagged Nst1, Nst1ΔPA, Nst1ΔPAM, and Nst1ΔD1. W303a

Δnst1 cells endogenously expressing Dcp2-mKate2 (YSK3663) were transformed with pMW20-pGAL-

GFP-NST1*,* pMW20-pGAL-GFP-NST1ΔPA*,* pMW20-pGAL-GFP-NST1ΔPAM, or vector only (control).

Following galactose induction for 4 hours, cells were treated with cycloheximide (CHX, 100 µg/ml) for 10

minutes and imaged using a Zeiss LSM 980 confocal microscope equipped with an Airyscan 2 detector

(Carl Zeiss). The right panel shows a 2× magnified view of selected puncta in the merged images. GFP

and mKate2 channels are shown in green and red, respectively. Scale bar: 5 µm. Representative images

are shown.

(C) Quantitative analysis of the spatial co-localization between Dcp2 and Nst1 condensates shown in (B)

after CHX-treatment. The co-localization ratio was calculated as the proportion of Dcp2 condensates

overlapping with Nst1 and its mutants-derived puncta. Puncta were segmented and quantified using Fiji

(ImageJ) software. For each Nst1 and its mutants, at least 100 condensates were analyzed per

experiment across three independent replicates. Statistical significance was determined by unpaired t-test

(***p□<□0.001) using GraphPad Prism 10.

(D) Schematic representation of the proposed Dcp2 condensation model based on the functional roles of

the PA and PAM domains of Nst1.

(E) Quantification of Dcp2-EGFP puncta formation in yeast cells (YSK3709) imaged by widefield

epifluorescence microscopy. Cells endogenously expressing Dcp2-EGFP were transformed to

overexpress Nst1, Nst1ΔPA, or Nst1ΔPAM for 4 hours. Images were acquired using identical microscope

settings for all conditions. Dcp2-EGFP puncta were segmented in FIJI (ImageJ) by intensity thresholding.

The number of puncta per cell was quantified and normalized by cell number, with more than 300 cells

analyzed for each condition. Data are presented as bar graphs generated using GraphPad Prism 10.

(F) Schematic illustration of the experimental workflow for the experiments (G–J) designed to elucidate

the roles of the PA and PAM domains in condensate formation with respect to free-RNA.

(F-J) For endogenous-level expression, w303a, w303a expressing chromosomally EGFP-tagged Nst1,

Dcp2, or Edc3, and *Δnst1* strains expressing chromosomally integrated GFP-Nst1ΔPA or GFP-

Nst1ΔPAM were cultured for 30 hours. Cells were lysed with or without RNase inhibitor, and insoluble

pellets were collected. All experiments were performed at least three times.

(G, I) Phase-contrast and fluorescence images of cell pellets prepared with or without RNase inhibitor.

Condensates composed of GFP-tagged Dcp2 and Edc3 for (G), or Nst1, Nst1ΔPA, and Nst1ΔPAM for (I),

were formed in stationary phase cells (30 hour cultured), collected, and visualized. Each experiment was

repeated three times, with more than five images acquired per experiment. Scale bar: 10 μm.

(H, J) Quantification of the maximum intensity values of segmented puncta from the images in (G) was

shown in (H), and from (J) is shown in (I). For each punctum, the maximum intensity was measured by

segmenting puncta within the top 0.05% of intensity values using Fiji. Statistical significance was

assessed by unpaired t-test (****p < 0.0001) using GraphPad Prism 10.

1115

1116

Figure 6

1117

Figure 6. *The Polyanion (PA) domain of Edc3 and human TDP-43 promote coalescence of condensates, similar to Nst1.*

(A) Schematic representation of TDP-43 domain architecture. Residues 2–29 and 197–220 were identified as clusters of negatively charged amino acids, designated PA1 and PA2, based on a sliding window approach and prediction tools, as described in Supplementary Figure 3. The region spanning residue 255-276 was denoted as PAM, while residues 255-414 were characterized as the C-terminal domain. See also Supplementary figure 3.

(B) Representative widefield fluorescence microscopy images of budding yeast (W303a) cells after 4 hours of overexpression of TDP-43 and its domain deletion mutants (ΔPA1, ΔPA2, ΔPAM, ΔCTD and ΔPA1ΔPA2) with or without NLS (Nuclear localization signal) sequence under a galactose-inducible promoter. Cells were fixed and stained with DAPI to visualize nuclei. Scale bar: 5 µm.

(C) The amino acid sequence of the PA1 and PA2 domains of TDP-43, along with its *PA1-1, PA1-2, PA1-1PA1-2, PA2-1, PA2-2,* and *PA2-1PA2-2* mutant variants, is depicted at the top. In the *PA1-1, PA1-2, PA1-1PA1-2, PA2-1, PA2-2,* and *PA2-1PA2-2* mutants, the negatively charged glutamic acid (E) and aspartic acid (D) residues were individually substituted with glutamine (Q) and asparagine (N), respectively, as indicated.

(D) Representative widefield fluorescence microscopy images are shown for budding yeast (W303a) cells overexpressing TDP-43 full-length (F.L.), each PA domain deletion mutants (ΔPA1 and ΔPA2), as well as the *PA1-1PA1-2, PA1-1, PA1-2, PA2-1PA2-2, PA2-1*, and *PA2-2* variants, all of which lack the NLS. Images were taken following 4 and 12 hours of galactose induction. For each strain and time point, more than 100 cells were examined, and representative images are presented. Scale bar: 5 µm.

(E) Representative wildfield fluorescence microscopy images of cells expressing GFP-tagged TDP-43 and its variants (ΔPA1, ΔPA2, ΔPAM, ΔCTD and ΔPA1ΔPA2) in U2OS cells at 48 hours after transfection. U2OS cells were transfected with plasmids encoding GFP-tagged TDP-43 variants under

1143   CMV promoter as described in Materials and Methods. After 48 hours, cells were observed using

1144   fluorescence microscopy to assess condensation pattern of each TDP-43 variant. Scale bar: 20 μm.