CA2 neurons show abnormal responses to social stimuli in a rat model of Fragile X syndrome

Margaret M. Donahue[#,1,2,3], Emma Robson[#,2,3], Alyssa M. Marron[1,2,3,4], Edel J. Fernandez[2,3], Misty Hill[2,3], Alexandra J. Mably[2,3], John B. Trimper[2,3], Darrin H. Brager[2,4,5], Laura Lee Colgin[1,2,3]

[#]Contributed equally
[1]Interdisciplinary Neuroscience Program, The University of Texas at Austin, Austin, Texas, 78712
[2]Center for Learning and Memory, The University of Texas at Austin, Austin, Texas, 78712
[3]Department of Neuroscience, The University of Texas at Austin, Austin, Texas, 78712
[4]School of Life Sciences, University of Nevada at Las Vegas, Las Vegas, Nevada, 92154
[5]Interdisciplinary Neuroscience Program, University of Nevada at Las Vegas, Las Vegas, Nevada, 92154

Abbreviated title: CA2 neuronal responses in a Fragile X rat model

Corresponding author: Laura Lee Colgin, colgin@mail.clm.utexas.edu

Number of pages: 42; **Number of figures**: 8 main figures and 2 supplemental figures; **Number of tables**: 2

Number of words: Abstract: 138; Significance statement: 93; Introduction: 630; Discussion: 1350

Conflict of interest: The authors declare no competing financial interests.

Acknowledgments: This research was supported by the Department of Defense CDMRP award W81XWH1810314 (to L.L.C.), Brain and Behavior Research Foundation (BBRF) Distinguished Investigator Grant 32221 (to L.L.C.), and National Institutes of Health awards R56MH125655 (to D.H.B. and L.L.C), R01MH131317 (to D.H.B. and L.L.C.), and F31MH127933 (to M.M.D.). The authors thank Isabella Lee, Ayomide Akinsooto, and Sirisha Dhavala for technical assistance and Chenguang Zheng for providing MATLAB code for some of the analyses. We also thank Dr. Moses V. Chao and Dr. Robert C. Froemke from New York University School of Medicine for generously providing the OXTR-2 antibody used in this study. The authors acknowledge the Texas Advanced Computing Center (TACC) at The University of Texas at Austin for providing data storage resources that have contributed to the research described within this article. URL: http://www.tacc.utexas.edu

Author contributions:
M.M.D. performed research, analyzed data, and wrote the paper; E.R. designed research, performed research, analyzed data, and wrote the paper; A.M.M. performed research; E.J.F. analyzed data; M.H. performed research and analyzed data; A.J.M. designed research, performed research, and analyzed data; J.B.T. performed research; D.H.B. designed research, performed research, and wrote the paper; L.L.C. designed research, analyzed data, and wrote the paper.

Abstract

Fragile X Syndrome (FXS) is a neurodevelopmental disorder that is highly comorbid with autism

spectrum disorders and can cause abnormal social behaviors. The CA2 subregion of the

hippocampus is essential for social memory processing and social recognition. A social

interaction induces changes in CA2 neuronal firing; however, it is unknown whether these

changes are impaired in FXS models. Here, we examined CA2 activity in a rat model of Fragile

X Syndrome (*Fmr1* knockout rats). In *Fmr1* knockout rats, we observed impaired CA2 cell

responses to social stimuli, despite similar social behaviors. Further, in CA2 of *Fmr1* knockout

rats, we found reduced expression of oxytocin receptors and impaired whole cell responses to

oxytocin. Together, these results raise the possibility that abnormal CA2 activity contributes to

impaired social behavior in FXS and may suggest novel treatment targets for FXS patients.

Significance statement

Fragile X Syndrome (FXS) is a neurodevelopmental disorder that can result in abnormal social

behaviors, including social avoidance. Activity in the CA2 subregion of the hippocampus is

believed to support social recognition and social cognition. Yet, the extent to which the CA2

subregion of the hippocampus is affected by FXS is poorly understood. In this study, we

identified specific impairments in CA2 neuronal responses to social stimuli in a rat model of

FXS. Further, we provide evidence suggesting that CA2 responses to oxytocin, a neuropeptide

released during social interactions, are abnormal in FXS.

Introduction

Fragile X Syndrome (FXS) is a neurodevelopmental disorder that is highly co-morbid with

autism spectrum disorders. Patients with FXS often display abnormal social behaviors, including

social avoidance (Kau et al., 2004; Kaufmann et al., 2004; Lesniak-Karpiak et al., 2003;

Merenstein et al., 1996; Williams et al., 2014). FXS is caused by a deficit in the function of

Fragile X Messenger Ribonucleoprotein (FMRP). FMRP is the product of the *Fmr1* gene, which

is located on the X chromosome. FXS is caused by hypermethylation and transcriptional

silencing of the *Fmr1* gene (Pieretti et al., 1991; Verkerk et al., 1991), leading to a complete lack

of FMRP production. FMRP is an RNA-binding protein (Ashley et al., 1993; Siomi et al., 1993)

believed to regulate translation and protein synthesis in neurons (Jin & Warren, 2000; Richter &

Zhao, 2021). Mouse and rat models of FXS (hereafter termed "FXS mice" and "FXS rats",

respectively) have been developed by knocking out the homologous *fmr1* gene in these species

(Bakker et al., 1994; Hamilton et al., 2014; Mientjes et al., 2006; Tian et al., 2017).

Social behavior deficits have been observed in FXS rodent models. FXS mice display more

anxiety-like behaviors during social interactions (McNaughton et al., 2008; Spencer et al., 2005)

and lower levels of social interaction (Liu & Smith, 2009; McNaughton et al., 2008; Mineur et al.,

2006; Oddi et al., 2014) than wild-type (WT) mice. In addition, FXS mice and rats may be

deficient in distinguishing between positive and negative (McNaughton et al., 2008), or familiar

and novel (Tian et al., 2017), social interactions.

The CA2 region of the hippocampus is essential for social recognition (Oliva, 2022). Inactivating

CA2 pyramidal cells impairs social memory and recognition (Hitti & Siegelbaum, 2014; Meira et

al., 2018; Oliva et al., 2020; Stevenson & Caldwell, 2014). CA2, like all subregions of the rodent

hippocampus, contains place cells, neurons that have spatial receptive fields called place fields

(O'Keefe & Dostrovsky 1971). Place cells often alter their firing patterns in response to changes

in sensory or cognitive input in a process referred to as "remapping" (Colgin et al., 2008). During

"global" remapping, place fields may appear, disappear, or change location. During "rate"

remapping, place fields do not move, but cells' in-field firing rates change. Previous research

has shown that the introduction of a social stimulus to a familiar environment induces global

remapping in CA2 place cells (Alexander et al., 2016; Robson et al., 2025). Furthermore, this

process appears to be driven by the olfactory content of social experiences (Robson et al.,

2025). CA2 place cell responses to social stimuli may reflect a cellular mechanism underlying

CA2's contributions to social memory.

In this study, we recorded CA2 place cells in FXS rats and WT control rats freely exploring an

open field arena. We analyzed changes in CA2 place cell firing patterns when stimuli with a

social olfactory component were added to the arena. We found that the presentation of social

odors was associated with global remapping in CA2 place cells in WT rats but not FXS rats.

Receptors for oxytocin, a neuropeptide that regulates social behaviors and social memory

(Froemke & Young 2021), are prominently expressed in CA2 (Mitre et al. 2016), and

administration of an oxytocin receptor agonist potentiated excitatory postsynaptic currents

(EPSCs) in CA2 neurons (Pagani et al. 2015). Thus, we hypothesized that abnormal CA2

neuronal responses to oxytocin contribute to deficient CA2 place cell remapping to social stimuli

in FXS rats. In accord with this hypothesis, we observed reduced oxytocin receptor expression

and impaired depolarizing responses to oxytocin in CA2 neurons in FXS rats. Together, these

results suggest that CA2 cellular responses involved in social information processing may be

impaired in FXS. The results also point to potential mechanisms that may contribute to

abnormal social behaviors in FXS.

Materials & Methods

Subjects

The FXS rats used in this study were *fmr1*-knockout rats on a Sprague-Dawley background

(SD-*Fmr1*-/null[tm1Sage]). Control rats used in this study were WT Sprague-Dawley rats. Both WT

and FXS rats were obtained from Inotiv. As FXS is an X chromosome-linked disorder, the

prevalence of FXS is higher in males, and symptoms tend to be more severe in male patients

than female patients. Therefore, we chose to use male rats. All experiments were conducted

according to the guidelines of the United States National Institutes of Health Guide for the Care

and Use of Laboratory Animals. *In vivo* and *in vitro* studies were conducted under protocols

approved by the Institutional Animal Care and Use Committees at the University of Texas at

Austin and the University of Nevada at Las Vegas, respectively.

Subjects for CA2 place cell recordings and open field exploration. CA2 place cell recordings in

this study were collected from twelve male rats. Six were WT control rats, and six were FXS

rats. The six WT rats used in the *in vivo* portion of this study were included in one previously

published study (Robson et al., 2025); one of the WT rats and three of the FXS rats were

included in another previously published study (Donahue, Robson, & Colgin 2025). Rats were

between the ages of 3 and 10 months old at the time of surgery. Before surgery, rats were

double or tripled housed with littermates in same-genotype cages. Following surgery, rats were

housed singly in custom-built acrylic cages (40 cm x 40 cm x 40 cm) to prevent damage to

implants. Enrichment material (i.e., paper towel rolls, wooden blocks, etc.) was added to the

cages, and rats were housed directly next to the cage of their former cage mates throughout

behavioral neurophysiology data collection. Rats were maintained on a reverse light cycle (light:

8 p.m. – 8 a.m.), and all behavioral neurophysiology experiments were performed during the

dark cycle. When necessary to encourage open field exploration, rats were temporarily placed

on a food deprivation protocol. Five WT rats and one FXS rat experienced temporary food

deprivation. With the provided food and treats consumed during open field exploration (see

"*Open field behavioral task*" section below), rats on food deprivation maintained at least 97% of

their free-feeding body weight.

After experiments had concluded, tissue (~1-2 mm) was collected from anesthetized rats

immediately prior to terminal perfusion. Tissue samples were then stored at -20°C and sent to

Transnettyx Inc. to verify rats' genotypes.

Subjects for olfactory habituation/dishabituation testing. Thirty-six rats (18 WT rats and 18 FXS

rats) were included in the olfactory habituation/dishabituation behavioral task in this study. Two

of the WT rats and four of the FXS rats were the same rats used for recording CA2 place cells.

Rats who were included in both olfactory habituation/dishabituation tests and CA2 place cell

recording studies completed olfactory testing prior to undergoing implantation surgery and CA2

data collection. All olfactory habituation/dishabituation testing was performed during the dark

cycle. As with rats used for CA2 place cell recordings, tissue samples were collected from

anesthetized rats immediately prior to euthanasia. Tissue samples were then stored at -20°C

and sent to Transnettyx Inc. to verify rats' genotypes.

Subjects for analysis of oxytocin receptor expression. Tissue was collected from three WT rats

and nine FXS rats for oxytocin receptor immunostaining. Experimenters were blinded to rat

genotype during tissue collection, immunostaining, imaging, and image analysis. Tissue

samples were collected from anesthetized rats immediately prior to terminal perfusion. Tissue

samples were then sent to Transnettyx Inc. for genotyping to verify rats' genotyping. Following

genotype verification, it was discovered that one cage had been assigned with the incorrect

genotype. Estimates of oxytocin receptor levels for this cage were subsequently reassigned to

the other genotype group. The mislabeled cage was a cage of FXS rats, which is why the FXS

group had more rats than the WT group.

Subjects for in vitro recordings. CA2 intracellular recordings were made in acute hippocampal

slices from four WT and four FXS male rats. Rats had free access to food and water and were

housed on a reverse light-dark cycle (12 hours on/ 12 hours off). Rats were 4–5 months old at

the time of slice preparation.

Surgery and tetrode positioning for CA2 place cell recordings

"Hyperdrive" recording devices containing 14-21 independently movable tetrodes were

implanted above the right dorsal hippocampus (anterior-posterior -3.8 mm from bregma, medial-

lateral -3.0 mm from bregma). Tetrodes were constructed from 17 µm polymide-coated

platinum-iridium (90/10%) wire (California Fine Wire, Grover Beach, California). The tips of the

tetrodes designated for single unit recording were platinum-plated to reduce single-channel

impedances to ~150-300 kOhms. During surgery, eleven bone screws were placed in the skull

and covered with dental acrylic to anchor the hyperdrive to the skull. Two of the screws were

connected to the recording drive ground.

On the day of surgery, all tetrodes were lowered ~1 mm into the brain. Tetrodes were then

slowly lowered to the hippocampal pyramidal cell body layer over several weeks except for one

tetrode that was designated as a reference electrode for differential recording. This tetrode was

placed in an electrically quiet area of cortex at least 1 mm above the hippocampal cell body

layer. The reference signal was duplicated using an MDR-50 breakout board (Neuralynx,

Bozeman, Montana) and continuously recorded to ensure that unit activity or volume conducted

hippocampal signals of interest (e.g., theta rhythms, sharp wave-ripples) were not detected.

Another tetrode was placed in the apical dendritic layer of CA1 to monitor hippocampal local

field potential (LFP) signals; this tetrode helped to guide placement of the other tetrodes using

estimated depth and electrophysiological hallmarks of the hippocampus (e.g., sharp wave-

ripples).

Data acquisition for CA2 place cell experiments

Data were acquired using a Digital Lynx system and Cheetah recording software (Neuralynx,

Bozeman, Montana). The recording setup has been described in detail previously (Zheng et al.,

2016). Briefly, LFP signals from one randomly chosen channel within each tetrode were

continuously recorded at a 2000 Hz sampling rate and filtered in the 0.1–500 Hz band. Input

amplitude ranges were adjusted before each recording session to maximize resolution without

signal saturation. Input ranges for LFPs generally fell within ±2,000 to ±3,000 µV. To detect unit

activity, all four channels within each tetrode were bandpass filtered from 600 to 6,000 Hz.

Putative spikes were detected when the filtered continuous signal on one or more of the

channels exceeded 55 µV. Detected events were acquired with a 32,000 Hz sampling rate for 1

ms. For both LFPs and unit activity, signals were recorded differentially against a dedicated

reference channel (see "*Surgery and tetrode positioning for CA2 place cell recordings*" section

above).

Videos of rats' behavior were recorded through the Neuralynx system with a resolution of 720 ×

480 pixels and a frame rate of 29.97 frames/s. Rat position and head direction were tracked via

an array of red and green or red and blue light-emitting diodes (LEDs) on a HS-54 or HS-27

headstage (Neuralynx, Bozeman, Montana), respectively.

Open field behavioral task

Rats were familiarized to an open field arena (1 m x 1 m with 0.5 m wall height) for a minimum

of three days before data collection began. Rats freely explored the open field arena for four 20-

minute sessions per day, with 10-minute rest sessions preceding and following each active

exploration session. During active exploration, small pieces of sweetened cereal or cookies

were randomly scattered to encourage rats to explore the entirety of the arena. Rats had to

cover at least 60% of the arena across each of the four sessions for a day to be included for

further analysis. During each rest session, rats rested in a towel-lined, elevated flowerpot

outside of the arena. A plexiglass standard rat housing cage was placed in one corner of the

arena for all recording sessions. In the first and fourth sessions (A and A'), this cage contained

only clean bedding. In the middle two sessions (B and B') of the different experimental

conditions, the cage contained social stimuli (Figure 1). In the Odor condition, a cage that had

formerly housed familiar rat(s) and contained soiled bedding was used. In the Visual + Odor

condition, a familiar rat was presented in its home cage containing soiled bedding. In the Control

condition, a clean cage containing only clean bedding was presented in all four sessions.

Histology for tetrode localization for CA2 place cell recordings

After place cell recordings were completed, rats were perfused with 4% paraformaldehyde

solution in phosphate-buffered saline. Brains were cut coronally in 30 μm sections using a

cryostat. For four of the rats, brain slices were stained with CA2-expressed protein striatum-

enriched protein-tyrosine phosphatase (STEP; Cell Signaling, 4817, 1:1000 dilution, raised in

mouse). Slices were then incubated in an anti-mouse secondary antibody and developed with

3,3-diaminobenzidine (brown) (as in Alexander et al., 2016). Slices were then counter-stained

with cresyl-violet (purple/blue) to indicate cell bodies. Both the presence of STEP (Figure S1A)

and the enlargement of cell bodies indicative of the CA2 region of the hippocampus (Lu et al.,

2015) (Figure S1B) were used to identify CA2. In eight of the rats, brains were immunostained

for CA2 marker Purkinje Cell Protein 4 (PCP4; Sigma-Aldrich Cat# HPA005792, 1:200 dilution,

raised in rabbit) (Figure S1C) in lieu of STEP staining. Slides were then mounted using DAPI

Fluoromount-G (Southern Biotech) to visualize cell bodies. Tetrode recording sites were

identified by comparing locations across adjacent sections.

Spike sorting and unit selection for CA2 place cell analyses

Spike sorting was performed manually using graphical cluster-cutting software (MClust, A.D.

Redish, University of Minnesota, Minneapolis, Minnesota) run in MATLAB (Mathworks). Spikes

were sorted using two-dimensional representations of waveform properties (i.e., energy, peak,

and peak-to-valley difference) from four channels. A single unit was accepted for further

analysis if the associated cluster was well isolated from, and did not share spikes with, other

clusters on the same tetrode. Units were also required to have a minimum 1 ms refractory

period. Units with mean firing rates above 5 Hz were considered putative interneurons and

excluded from further analysis. In order to be considered active in the arena, a unit had to reach

a peak firing rate of at least 1 Hz. In order to be included in the sharp wave-ripple firing analysis,

a unit had to have valid clusters in both the active exploration and rest sessions. CA2 cell yields

in each condition are reported for WT rats in Table 1 and for FXS in Table 2.

Table 1. CA2 place cell yields for WT rats.

Genotype	Rat ID	Control	Odor	Visual + Odor
WT	Rat 117 ("Lauren")	7	5	0
WT	Rat 122 ("Chase")	51	32	0
WT	Rat 165 ("Gus")	34	56	26
WT	Rat 256 ("Viggo")	28	13	13
WT	Rat 391 ("Daffodil")	15	4	5
WT	Rat 418 ("Hugo")	20	35	41
	Total	155	145	85

223   Table 2: CA2 place cell yields for FXS rats.

Genotype	Rat ID	Control	Odor	Visual + Odor
FXS	Rat 120 ("Brock")	19	13	0
FXS	Rat 125 ("JR")	2	3	0
FXS	Rat 330 ("Aires")	0	1	1
FXS	Rat 395 ("Elijah")	6	1	4
FXS	Rat 445 ("Paddy")	7	7	5
FXS	Rat 442 ("Pippin")	33	35	26
	Total	72	60	40

Place cell remapping analyses

Methods used to create firing rate maps for each single unit were based on established

methods from the place cell field that were used in our lab's prior studies (Alexander et al.,

2016; Robson et al., 2025; example rate maps for the present study shown in Figure 2). First,

the arena was divided into 4 cm^2 bins. The number of spikes that occurred within each bin was

divided by the time spent in that bin to determine the firing rate. Only spikes that occurred while

the rat was traveling 5 cm/s or faster were included. The rate map was smoothed with a two-

dimensional Gaussian kernel (standard deviation = 6 cm). To determine if a place cell

remapped during presentation of a social stimulus, a Pearson correlation coefficient R was

calculated for each unit between pairs of rate maps from control and social stimuli sessions (i.e.,

A-B, B-B', B'-A', A-B', and A-A' session pairs, see Figure 1). To determine if spatial correlation

coefficients differed between WT and FXS rats, or across session pairs (e.g., A-B and A-A') or

conditions (i.e., Odor, Visual + Odor, and Control), we used a generalized linear mixed model

statistical analysis (IBM SPSS Statistics, version 29.0.2.0; see "*Experimental design and*

statistical analysis" section below). The estimated mean spatial correlations for each genotype

and condition, collapsed across session pairs, are shown in Figure 3A. Individual spatial

correlation values for each cell are shown for each session pair and each condition in Figure

3B-F.

To determine if rate remapping occurred in CA2 place cells across genotypes (i.e., WT and FXS

rats) and conditions (i.e., Odor, Visual + Odor, and Control), we calculated the rate overlap

between pairs of sessions for each place cell. The rate overlap was defined as the ratio of the

mean place field firing rate of the cell in each session, with the smaller of the two firing rates as

the numerator (as in Robson et al., 2025). This analysis allowed us to determine if CA2 place

cells changed their firing rates within a place field and therefore excluded cells that did not have

an identified place field from a given session. This allowed us to compare firing rates without the

confound of including cells that may gain or lose a field between sessions, which is defined as

global remapping, not rate remapping. To determine if rate overlap differed across genotypes

(i.e., WT and FXS rats), session pairs (i.e., A-B, A-B', B'-A', A-A', B-B'), and conditions (i.e.,

Odor, Visual + Odor, and Control), we used a generalized linear mixed model statistical analysis

(IBM SPSS Statistics, version 29.0.2.0; see "*Experimental design and statistical analysis*"

section below). The estimated mean rate overlap values for each genotype and condition,

collapsed across session pairs, are shown in Figure S2A. Individual rate overlap values for each

cell are shown for each session pair and each condition in Figure S2B-F.

Analyses of place cell properties

Spatial information (Figure 4A) was calculated as previously described (Skaggs et al., 1996)

using the following formula:

263
$$Spatial\ information = \ \sum_i P_i \frac{\lambda_i}{\lambda} log2 \frac{\lambda_i}{\lambda},$$

where i = the spatial bin index, P_i = the probability of the rat being in the ith bin, λ_i = the mean

firing rate in the ith bin, and λ = the mean firing rate of the cell.

Place fields were separately identified for each cell in each session. Potential place fields were

identified as contiguous bins of the smoothed rate map (see "*Place cell remapping analyses*"

section above) that exceeded 50% of the peak firing rate in that session. In order to be included

for further analysis, the peak firing rate of a place field had to exceed 1 Hz, and the size of the

place field had to exceed 10 cm^2.

Exploration time analysis for open field behavioral task

Exploration of the stimulus cage was quantified as previously described (Robson et al., 2025).

The arena was divided into 4 cm^2 bins, and the time spent in each bin during the first two

minutes of every session was determined for each day. These exploration maps were then

smoothed with a two-dimensional Gaussian kernel (standard deviation = 4 cm). Maps were

averaged within and then across rats within each genotype and plotted as a heat map for each

condition (Figure 5A-B). The time spent within 12 cm of the cage was then calculated (Figure

5C).

Olfactory habituation/dishabituation task

To determine if rats could detect and distinguish between social odors, we used an olfactory

habituation/dishabituation task (Yang & Crawley, 2009). First, rats were pre-acclimatized to the

test cage containing a clean dry cotton tip applicator for 15 mins to reduce novelty-induced

exploratory activity during the test. During the testing phase, rats were initially exposed to a

cotton-tip applicator with no odor for 2-3 trials ("null" trials; N1-N3 in Figure 6). This was used to

establish a baseline level of exploration for the object (i.e., the cotton-tip applicator). Following

the null trials, rats were exposed to cotton tips with an unfamiliar social odor for four consecutive

trials ("habituation" trials, H1-H4 in Figure 6). Following the habituation trials, rats were exposed

to a cotton tip with odors from a second unfamiliar social odor ("dishabituation" trial, D in Figure

6). All trials were 90 seconds in length. Social odors were prepared by swiping soiled home

cages of unfamiliar rats with cotton tip applicators. All trials were video recorded. Following

completion of the experiment, the time rats spent sniffing the cotton tip (within ~2 cm of the

cotton tip and nose pointing towards the cotton tip) during each trial was scored by two

independent scorers. The time spent sniffing was then averaged across the scorers. To account

for differences in baseline activity levels across rats, we calculated normalized difference scores

for each rat (r) and trial (t) using the following formula:

299
$$Norm.diff.score_{r,t} = \frac{t_r - N1_r}{t_r + N1_r}$$

where t_r was the sniffing time from the trial of interest for a given rat and $N1_r$ was the sniffing

time from the first null trial for that rat.

Immunohistostaining for oxytocin receptors

Rats (three WT rats and nine FXS rats, aged 25 weeks) were perfused in phosphate-buffered

saline (PBS) followed by 4% paraformaldehyde solution in PBS. Brains were stored overnight in

4% paraformaldehyde solution in PBS. On the day after perfusion, brains were cut coronally in

30 µm sections using a cryostat and mounted onto slides. Sections were kept frozen at -80

degrees Celsius until tissue from all rats had been collected. Immunostaining was then

performed simultaneously on all sections from both genotypes. First, all slides were defrosted to

room temperature and washed multiple times with water, boiling water, and Tris-buffered saline

(TBS) + 0.3% Triton X-100. Sections were then blocked for 2 hours with 5% Normal Goat

Serum (NGS). Sections were then incubated overnight with the following primary antibodies

diluted in blocking buffer: Mouse anti-RGS14 (1:1000; Antibodies Inc) and rabbit anti-oxytocin

receptor (1 µg/ml; Moses V. Chao, Robert C. Froemke, NYU School of Medicine Cat# OXTR-2

antibody, RRID:AB_2571555). The following day, slides were washed with Tris-buffered saline

with Tween-20 (TBST) and then incubated for two hours with the following secondary

antibodies: goat anti-mouse ALEXA 555, A-21422 (1:500; Thermo Fisher) and goat anti-rabbit

ALEXA 488, A-11008 (1:500, Thermo Fisher). Slides were then washed with TBST and TBS.

Lastly, slides were mounted using DAPI Fluoromount-G (Southern Biotech).

Quantification of OXTR-2 expression

Slides were imaged using a Zeiss AXIO Zoom.V16 (software Zen 3.11 Pro) microscope with an

Axiocam512 color camera with filters for imaging DAPI (blue, excitation range 350-360 nm,

emission range 460nm), Cy3 (orange-red excitation range 513-556, emission 570-613), and

EGFP (Enhanced Green Fluorescent Protein, excitation about 470, emission about 510) at a

magnification of 77x. For imaging sections that were immunostained with multiple antibodies, a

standardized exposure time was chosen for each filter: DAPI 200 ms, Cy3 350 ms, and EGFP

280 ms. The levels of oxytocin receptors in each section were estimated via the following

procedures. First, blinded experimenters took photomicrographs of WT and FXS sections using

the same microscope settings. Images were saved and subsequently analyzed by blinded

experimenters using Image J software (ImageJ 1.54g). Images taken with the blue (DAPI),

green (OXTR-2), and red (PCP4) settings were opened and calibrated to the same scale.

Regions of interest (i.e., CA3, CA2, and CA1) were defined in all 3 sets of images using the

PCP4 staining as a marker for CA2. To estimate OXTR-2 levels, background fluorescence was

removed, and estimates were obtained for mean fluorescence (average pixel intensity within

each region of interest) and total (integrated) fluorescence. To determine if OXTR-2 expression

differed between WT and FXS rat, or across hippocampal subregions, we used a generalized

linear mixed model statistical analysis (IBM SPSS Statistics, version 29.0.2.0; see

"*Experimental design and statistical analysis*" section below). The estimated corrected

fluorescence from the generalized linear mixed model for each genotype and subregion is

shown in Figure 7C.

Slice preparation for in vitro recording

Rats were anesthetized using a ketamine/xylazine cocktail (100/10 mg/kg) and then underwent

cardiac perfusions with ice-cold saline consisting of (in mM): 2.5 KCl, 1.25 NaH_2PO_4, 25

$NaHCO_3$, 0.5 $CaCl_2$, 7 $MgCl_2$, 7 dextrose, 205 sucrose, 1.3 ascorbic acid, and 3 sodium

pyruvate (bubbled constantly with 95% O_2/5% CO_2 to maintain pH at ~7.4). Following perfusion,

the brain was removed and sliced into 300 μM parasagittal sections from the middle

hippocampus using a vibrating tissue slicer (Vibratome 300, Vibratome Inc). The slices were

placed in a chamber filled with artificial cerebral spinal fluid (aCSF) consisting of (in mM): 125

NaCl, 2.5 KCl, 1.25 NaH_2PO_4, 25 $NaHCO_3$, 2 $CaCl_2$, 2 $MgCl_2$, 10 dextrose, 1.3 ascorbic acid

and 3 sodium pyruvate (bubbled constantly with 95% O_2/5% CO_2) for 30 minutes at 35°C and

then held at room temperature until time of recording.

In vitro electrophysiology

Slices were placed in a submerged, heated (33-34 C°) recording chamber and continually

perfused with aCSF (in mM): 125 NaCl, 3 KCl, 1.25 NaH_2PO_4, 25 $NaHCO_3$, 2 $CaCl_2$, 1 $MgCl_2$,

10 dextrose, and 3 sodium pyruvate (bubbled constantly with 95% O_2/5% CO_2). Ionotropic

glutamatergic and GABAergic synaptic transmission were blocked with 20 μM DNQX, 25 μM D-

AP5, and 2 μM gabazine. Pyramidal neurons within the CA2 region were visualized with a Zeiss

AxioScope under 60x magnification.

Current clamp recordings were made using a Dagan BVC-700 or Sutter IPA amplifier and

Sutterpatch acquisition software. Data were sampled at 20-50 kHz, filtered at 3 kHz, and then

digitized by a Dendrite interface or integrated A/D board in the Sutter IPA (Sutter). The internal

recording solution consisted of (in mM): 135 K-gluconate, 10 HEPES, 7 NaCl, 7 K_2-

phosphocreatine, 0.3 Na-GTP, 4 Mg-ATP (pH corrected to 7.3 with KOH). Recording electrodes

were pulled from borosilicate glass and had an open tip resistance of 4-6 MΩ. Series resistance

was compensated using the bridge balance circuit and was monitored throughout the

experiment. Recordings in which the series resistance exceeded 35 MΩ or increased by more

than 30% were excluded. The resting membrane potential was noted immediately after

establishing the whole cell recording configuration.

Drugs for in vitro electrophysiology experiments

DNQX, APV, and gabazine were purchased from Hello Bio. Thr^4-Gly^7 oxytocin (TGOT) was

purchased from Bachem. All drugs were prepared from a 1000x stock solution in water.

Experimental design and statistical analyses

Whenever possible, experimenters were blinded to genotype while collecting and analyzing

place cell data (i.e., during spike sorting). A period of limited availability of FXS rats from the

supplier resulted in unblinded collection and analysis of data from three rats (rats 418, 442, and

445). All video scoring for the olfactory habituation/dishabituation task was performed by scorers

blinded to genotype. Experimenters were blinded to genotype during OXTR-2 immunostaining

and quantification. Two experimenters conducted the in vitro electrophysiology experiments.

One was blinded to genotype, but the other was not.

We used a generalized linear mixed model approach (IBM SPSS version 29.0.2.0) to assess

differences between genotypes (i.e., WT and FXS rats) for in vivo neurophysiology, behavior,

and histology data. For all analyses, rats were set as the subjects, and genotype was included

as a fixed factor. Post-hoc tests were performed if a significant interaction or main effect was

observed and used the Bonferroni correction. For place cell analyses, multiple place cells were

nested within rats. To determine if spatial correlation coefficients (Figure 3) or rate overlap

values (Figure S2) differed across genotypes, conditions, and session pairs, condition and

session pair were included as fixed factors, and session pairs were repeated measures within

rats. A three-way interaction between genotype, condition, and session pair was included in the

model, as was a two-way interaction between genotype and condition. To assess place cell

properties (Figure 4), session was included as a repeated measure within rats. In addition to the

main effect of genotype, the model included: a three-way interaction between genotype,

condition, and session; a two-way interaction between genotype and session; and a two-way

interaction between condition and session.

To verify that our statistical tests were sufficiently powered given that a lower number of CA2

place cells were recorded in FXS rats than in WT rats (see Tables 1 and 2), we performed the

following analyses. First, we performed a simulation-based power analysis in which we

randomly down-sampled the number of CA2 place cells from WT rats to match the number

recorded in FXS rats for each condition (i.e., 72 cells in the Control condition, 60 cells in the

Odor condition, and 40 cells in the Visual + Odor condition) 100 times. We then used a

generalized linear mixed model to test for a statistically significant difference between the

Control condition and the Odor and Visual + Odor conditions for each simulation. We included

an interaction between condition and session and main effects of condition and session in the

model. Post-hoc comparisons were made using the Bonferroni correction when a significant

effect was found. We tallied the number of simulations in which the post-hoc pairwise

comparison showed a significant difference between the control and experimental conditions.

We did this separately for Control and Odor, and Control and Visual + Odor, comparisons to

obtain power estimates for each. We additionally ran permutation tests to compare the effects of

condition on spatial correlation values for each genotype and condition. The permutation tests

shuffled condition 1,000 times to obtain a null distribution for each spatial correlation value.

Monte-Carlo p-values were calculated using the formula:

$$p = \frac{N_{subset} + 1}{N_{shuffle} + 1}$$

where N_{subset} is the number of shuffles with values greater than or equal to the observed value

(two-tailed test), and $N_{shuffle}$ is the total number of shuffles.

To compare social stimuli exploration times across genotypes (Figure 5), days were included as

random factors nested within rats, as conditions were repeated on different days within rats.

Sessions were set as a repeated measure. In addition to the main effect of genotype, the model

included a three-way interaction between genotype, condition, and session and a two-way

interaction between condition and session.

To analyze olfactory stimuli investigation times in the habituation/dishabituation task (Figure 6),

trial was set as a repeated measure. In addition to the main effect of genotype, a main effect of

trial and a two-way interaction between genotype and trial was included in the model.

To compare the expression of oxytocin receptors across genotypes and hippocampal

subregions (Figure 7), multiple fluorescent estimates from brain sections were nested within

each subregion and set as repeated measures. In addition to genotype, subregion was included

as a fixed factor in the model.

Statistical tests for in vitro electrophysiology data (Figure 8) were also performed in SPSS (IBM

SPSS version 29.0.2.0). To compare the resting membrane potential of CA2 neurons across

WT and FXS rats, a univariate analysis of variance was performed with genotype as the

between-subjects factor. A general linear model was used to assess the effects of TGOT

application on CA2 neurons' resting membrane potential, with drug treatment (baseline vs.

TGOT) as a repeated measure and genotype as a between-subjects factor.

Data and code availability

446 The analysis code is available on GitHub

447 (https://github.com/mmdonahue/FMR1CA2_PUBLICSHARE). Data will be made available upon

448 reasonable request.

450 **Results**

452 *CA2 place cells in FXS rats did not remap to social stimuli*

453 Our prior studies have shown that CA2 place cells remap to social stimuli (Alexander et al.,

454 2016; Robson et al., 2025) and, specifically, that this remapping is driven by the olfactory

455 content of social stimuli (Robson et al., 2025). As FXS rats have atypical social behaviors

456 (Hamilton et al., 2014; Tian et al., 2017; Wong et al., 2020), the primary goal of this study was to

457 determine if CA2 place cells in FXS rats respond normally to the presentation of social stimuli.

458 Thus, we presented social stimuli to FXS and WT control rats while recording CA2 place cell

459 activity. We compared CA2 place cell firing patterns between WT and FXS rats across social

460 and control conditions (Figure 1).

462 We have previously shown that the presentation of a conspecific in a cage with soiled bedding

463 or the presentation of a soiled cage alone induces global remapping, but not rate remapping, in

464 a subset of CA2 place cells (Alexander et al., 2016; Robson et al., 2025). To test if similar

465 remapping occurs in CA2 place cells of FXS rats, we compared CA2 place cell firing patterns in

466 WT and FXS rats exploring an open field arena in which social stimuli (i.e., a familiar rat in a

467 soiled cage or a soiled cage alone) were presented. CA2 place cell firing patterns in WT and

468 FXS rats during presentation of social stimuli were compared to CA2 place cells recorded in a

469 control condition in which no social stimuli were presented in the stimulus cage (Figure 1).

Our recordings showed seemingly normal place cells in CA2 of FXS rats (Figure 2). To quantify

the extent to which CA2 place cells exhibited global remapping (i.e., a change in place cell firing

patterns in which place field locations change) when social stimuli were presented, we

calculated spatial correlation coefficients between rate maps from pairs of sessions (Figure 3).

We found that spatial correlation coefficients were lower for the Odor and Visual + Odor

conditions compared to the Control condition in WT rats (Figure 3; generalized linear mixed

model, no interaction between genotype, condition, and session pair ($F_{(20,2755)} = 1.2$, $p =$

0.244); significant interaction between genotype and condition ($F_{(2,2755)} = 6.9$, $p = 0.001$);

significant differences in post-hoc tests for WT rats for Control vs. Odor ($t_{(2755)} = 6.5$, $p <$

0.001) and Control vs. Visual + Odor ($t_{(2755)} = 6.8$, $p < 0.001$)). However, there were no

differences between the Control condition and either of the social conditions in FXS rats

(Figures 2B and 3; no differences in post-hoc tests for FXS rats for Control vs. Odor ($t_{(2755)} =$

1.1, $p = 0.809$) and Control vs. Visual + Odor ($t_{(2755)} = 0.7$, $p = 0.973$)). These results suggest

that CA2 place cells in FXS rats do not remap when social stimuli, namely a familiar conspecific

in a cage with social odors or social odors in the absence of a rat, are presented.

Because a lower number of place cells were recorded in FXS rats than in WT rats (see Tables 1

and 2), we performed analyses to determine whether our tests were sufficiently powered to

detect CA2 place cell remapping (Alexander et al., 2016; Robson et al., 2025) using the effect

size observed in WT rats. First, we used a simulation-based power analysis in which we the

number of CA2 place cells from WT rats was randomly down-sampled 100 times to match the

number recorded in FXS rats. We then used a generalized linear mixed model to test for a

statistically significant difference between the Control condition and the Odor and Visual + Odor

conditions for each simulation. We found that there was a significant main effect of condition

and a significant post-hoc pairwise comparison between the Control and Odor conditions 87%

of the time. There was a significant main effect of condition and a significant post-hoc pairwise

comparison between the Control and Visual + Odor conditions 92% of the time. We additionally

ran permutation tests to compare the effects of condition on spatial correlation values for each

genotype and condition. We found that spatial correlation values were lower in both the Odor

(permutation test, 1000 shuffles of condition, effect size = -0.36, $p < 0.001$) and Visual + Odor

(permutation test, 1000 shuffles of condition, effect size = -0.41, $p < 0.001$) conditions than in

the Control condition in WT rats. However, there were no significant differences between the

Control condition and the Odor or Visual + Odor conditions in FXS rats (Control vs. Odor:

permutation test, 1000 shuffles of condition, effect size = -0.09, $p = 0.27$; Control vs. Visual +

Odor, permutation test, 1000 shuffles of condition, effect size = -0.06, $p = 0.51$). These results

suggest that the lack of CA2 place cell remapping observed in FXS rats was not explained by

the lower number of place cells recorded in FXS rats.

Although CA2 place cells in FXS rats did not globally remap to a social stimulus, it is possible

that CA2 place cells respond to social stimuli by rate remapping. To test this possibility, we

calculated the rate overlap between pairs of sessions for CA2 place cells in WT and FXS rats

(Figure S2). We have previously reported that CA2 place cells did not rate remap to the

presentation of social stimuli in WT rats (Robson et al., 2025). In the present study, we also did

not observe rate remapping to social stimuli in CA2 place cells in FXS rats (generalized linear

mixed model, no interaction between genotype, condition, and session pair ($F_{(8,2718)} = 0.9$, $p =$

0.511), no interaction between genotype and condition ($F_{(2,2718)} = 0.8$, $p = 0.433$), no main

effect of genotype ($F_{(1,2718)} < 0.1$, $p = 0.901$), no main effect of condition ($F_{(2,2718)} = 1.4$, $p =$

0.252)).

CA2 place cells exhibited normal firing properties in FXS rats

The lack of remapping to social stimuli observed in CA2 place cells in FXS rats raised a

question of whether basic firing or spatial coding properties of CA2 place cells were abnormal in

FXS rats. Thus, we next assessed whether baseline firing properties of CA2 place cells differed

between WT and FXS rats. While CA1 place cell activity has been characterized in both FXS

mouse and rat models (Arbab, Battaglia, et al., 2018; Arbab, Pennartz, et al., 2018; Asiminas et

al., 2022; Donahue, Robson, & Colgin 2025; Talbot et al., 2018), basic firing and spatial coding

properties of CA2 place cells in FXS rodent models have not previously been described.

In addition to quantifying remapping across sessions in which environmental conditions change,

spatial correlation coefficients can also be calculated across sessions in an unchanging

environment to quantify place cell stability. We calculated spatial correlations between session

pairs in the Control condition and observed no significant differences between WT and FXS rats

(Figure 3; generalized linear mixed model, no difference between WT and FXS in post hoc test

for Control condition, $t(2755) < 0.1$, $p = 0.987$)). This suggests that the baseline stability of CA2

place cells in an unchanged environment is similar between WT and FXS rats.

We next examined firing properties of CA2 place cells in WT and FXS rats during individual

sessions. We found that spatial information transmitted by CA2 place cells did not differ

between WT and FXS rats across all conditions and sessions (Figure 4A; generalized linear

mixed model, no interaction between genotype, condition, and session ($F(8,2204) = 0.5$, $p =$

0.863); no interaction between genotype and session ($F(2,2204) = 0.3$, $p = 0.803$); no main

effect of genotype ($F(1,2204) = 0.2$, $p = 0.674$)). Average in-field firing rates (Figure 4B;

generalized linear mixed model, no interaction between genotype, condition, and session

($F(8,2016) = 1.1$, $p = 0.366$); no interaction between genotype and session ($F(3,2016) = 1.8$, $p =$

0.138); no main effect of genotype ($F(1,2016) = 0.2$, $p = 0.668$)) and peak firing rates (Figure

4C; generalized linear mixed model, no interaction between genotype, condition, and session

(F(8,2016) = 1.1, p = 0.340); no interaction between genotype and session (F(3,2016) = 1.9, p =

0.128); no main effect of genotype (F(1,2016) = 0.2, p = 0.673)) of CA2 place cells were also

similar between WT and FXS rats. Also, CA2 place field size (Figure 4D; generalized linear

mixed model, no interaction between genotype, condition, and session (F(8,2016) = 0.2, p =

0.277); no interaction between genotype and session (F(3,2016) = 0.3, p = 0.857); no main

effect of genotype (F(1,2016) = 0.8, p = 0.380)) and average number of place fields per place

cell per session (Figure 4E; generalized linear mixed model, no interaction between genotype,

condition, and session (F(8,2204) = 0.9, p = 0.539); no interaction between genotype and

session (F(3,2204) = 1.3, p = 0.263); no main effect of genotype (F(1,2204) = 0.7, p = 0.389))

did not differ between WT and FXS rats. Altogether, these results indicate that CA2 place cells

represent spatial locations similarly in FXS and WT rats.

FXS rats explored and perceived social stimuli normally

A possible explanation of why CA2 place cells failed to remap to social stimuli in FXS rats could

be reduced exploration of, or attention to, social stimuli in FXS rats. To explore this possibility,

we calculated the amount of time spent exploring the stimulus cage across conditions for WT

rats and FXS rats (Figure 5). We first created heat maps to visualize the amount of time rats

spent exploring locations across the open field arena and noted that both WT and FXS rats

spent a high amount of time near the stimulus cage in the Odor and Visual + Odor conditions

(Figure 5A-B). To quantify these observations, we calculated the time spent near the stimulus

cage across conditions and sessions for all rats. We found that both WT and FXS groups of rats

increased the time they spent near the stimulus cage in Session B during the Odor and Visual +

Odor conditions (Figure 5C; generalized linear mixed model, no interaction between genotype,

condition, and session (F(11,260) = 0.4, p = 0.935); significant interaction between condition

and session ($F_{(11,260)} = 5.4$, $p < 0.001$); significant differences in post-hoc tests for Session B

for Control vs. Odor ($t_{(260)} = -5.3$, $p < 0.001$) and Control vs. Visual + Odor ($t_{(260)} = -4.1$, $p <$

0.001)). This indicates that WT and FXS rats explored social stimuli to a similar extent.

We have previously shown that the olfactory content of social stimuli is necessary for CA2 place

cell remapping to social stimuli (Robson et al., 2025). Although structural and functional

changes have been identified in the olfactory system of FXS rodent models (Bodaleo et al.,

2019), these differences appear to impact olfactory sensitivity and fine odor discrimination, but

not the detection of odors or the discrimination of monomolecular odors (Kuruppath et al., 2023;

Nitenson et al., 2015). Still, it is possible that impaired olfactory perception of social stimuli

explains the lack of CA2 place cell remapping to social stimuli in FXS rats. To assess whether

olfactory perception and discrimination were impaired in FXS rats, we assessed olfactory

function in FXS rats using an olfactory habituation/dishabituation task (Figure 6). This task tests

rats' ability to recognize an odor when it is initially presented, habituate to the presence of the

same odor across repeated presentations ("habituation" trials), and distinguish between odors

after the presentation of a novel odor in the last trial ("dishabituation" trial). We found that FXS

rats could detect the presence of a social odor normally during the first habituation trial (Figure

6; generalized linear mixed model, no interaction between genotype and trial ($F_{(7,259)} = 0.6$, p

$= 0.722$); significant main effect of trial ($F_{(1,259)} = 126.9$, $p < 0.001$); significant differences in

post-hoc tests for Null 1 vs. Habituation 1 ($t_{(259)} = -4.2$, $p = 0.001$), Null 2 vs. Habituation 1

($t_{(259)} = -7.4$, $p < 0.001$), Null 3 vs. Habituation 1 ($t_{(259)} = -9.8$, $p < 0.001$)). Further, both WT

and FXS rats decreased their sniffing of the same social odor across repeated presentations

(significant difference in post-hoc test for Habituation 1 vs. Habituation 4 ($t_{(259)} = 9.2$, $p <$

0.001)), indicating normal habituation to a social odor in FXS rats. FXS rats also increased their

sniffing in the same manner as WT rats when a novel social odor was introduced during the

dishabituation trial (significant difference in post-hoc test for Habituation 4 vs. Dishabituation

($t(259)$ = -8.7, $p < 0.001$)). This indicates that FXS rats detected social odors, and distinguished

between different social odors, in the same way as WT rats.

Impaired oxytocin modulation of area CA2 in FXS rats

The neuropeptide oxytocin plays a key role in social behaviors and social memory (Froemke &

Young 2022). High expression of oxytocin receptors is a characteristic feature of area CA2

(Mitre et al. 2016), and oxytocin receptors in CA2 are essential for social functions (Raam et al.

2017, Lin et al. 2018, Tsai et al. 2022). Thus, we hypothesized that reduced responses to social

stimuli in CA2 place cells could be due to deficient oxytocin receptor expression in CA2 of FXS

rats. To test this hypothesis, we immunostained hippocampal tissue from WT and FXS rats with

OXTR-2, an antibody that is highly specific for oxytocin receptors (Mitre et al. 2016). We then

compared expression of oxytocin receptors in area CA2, and neighboring subregions CA1 and

CA3, in WT and FXS rats (Figure 7). Significant differences in the mean intensity of OXTR-2

immunostaining were observed across subregions and genotypes (Figure 7C; generalized linear

mixed model, significant interaction between genotype and subregion ($F(2,701)$ = 17.1, $p <$

0.001)). In WT rats, oxytocin receptor expression levels were higher in CA2 than in CA1 and

CA3 (generalized linear mixed model, significant main effect of subregion for WT rats ($F(2,174)$

= 93.7, $p < 0.001$) and significant differences in post-hoc tests for CA1 vs. CA2 ($p < 0.001$) and

CA3 vs CA2 ($p < 0.001$)), as expected based on earlier results from WT mice (Mitre et al. 2016).

Similarly, oxytocin receptor expression in FXS rats was higher in CA2 than in CA1 and CA3

(generalized linear mixed model, significant main effect of subregion for FXS rats ($F(2,527)$ =

459.7, $p < 0.001$) and significant differences in pairwise comparisons for CA1 vs. CA2 ($p <$

0.001) and CA3 vs CA2 ($p < 0.001$)). We additionally found that oxytocin receptor expression

levels were significantly lower in FXS rats than in WT rats across all subregions (generalized

linear mixed model, significant main effect of genotype ($F_{(1,701)}$ = 160.9, $p < 0.001$)) and in

CA2 tested individually (generalized linear mixed model, significant main effect of genotype

($F_{(1,233)}$ = 9.8, $p = 0.002$)). These results raise the possibility that reduced responses to

oxytocin in CA2 neurons contribute to diminished responses to social stimuli observed in CA2

place cells in FXS rats.

Thus, we next tested whether CA2 neurons in FXS rats exhibit reduced responses to oxytocin

receptor activation. Prior work in WT mice showed that CA2 neurons depolarize in response to

oxytocin, and such depolarizing responses were proposed to promote social remapping in CA2

(Tirko et al. 2018). Accordingly, we hypothesized that CA2 neurons would depolarize in

response to oxytocin administration in slices from WT rats but not FXS rats. To test this

hypothesis, we performed whole-cell current clamp recordings of CA2 pyramidal neurons and

tested how bath application of the specific oxytocin receptor agonist Thy^4,Gly^7-oxytocin (TGOT,

400 nM) affected the resting membrane potential. First, we found that the baseline resting

membrane potential of CA2 pyramidal neurons was more hyperpolarized in FXS slices than in

WT slices (Figure 8; n = 6 cells from 4 rats of each genotype, univariate analysis of variance,

genotype as between-subjects factor, significant effect of genotype ($F_{(1,10)}$ = 5.0, $p = 0.05$)).

Next, we found that TGOT administration affected FXS and WT neurons differently (Figure 8;

general linear model with genotype as between-subjects-factor and drug treatment as repeated

measure within subjects, significant interaction between genotype and drug treatment ($F_{(1,10)}$ =

5.4, $p = 0.04$)). Specifically, TGOT application significantly depolarized CA2 pyramidal cells in

slices from WT rats (significant difference in post-hoc pairwise comparison for WT rats for

baseline vs. TGOT (ΔV_M: 3.8 ± 0.83 mV; $t_{(5)}$ = -4.6, $p = 0.006$)), as expected (Tirko et al. 2018).

In contrast, CA2 pyramidal neurons in slices from FXS rats did not significantly depolarize in

response to TGOT application (no significant difference for FXS rats for baseline vs. TGOT

(ΔV_M: 1.2 ± 0.79 mV; t(5) = -1.472, p = 0.201)). These results suggest that deficient responses

to oxytocin in CA2 neurons may hinder activation of CA2 neurons by social stimuli and thereby

contribute to deficits in social remapping observed in CA2 place cells in FXS rats.

Discussion

A large number of studies have identified abnormalities in social behaviors in mouse and rat

models of FXS (Hamilton et al., 2014; Liu & Smith, 2009; McNaughton et al., 2008; Mineur et

al., 2006; Oddi et al., 2014; Saxena et al., 2018; Spencer et al., 2005; Tian et al., 2017; Wong et

al., 2020). In this study, we identified a neurophysiological difference in responses to social

stimuli in CA2 place cells of FXS rats. We have previously shown that a significant proportion of

CA2 place cells in WT rats changed their firing fields when a familiar rat in a soiled cage was

presented (Alexander et al., 2016; Robson et al., 2025) or when a soiled home-cage was

presented without a rat (Robson et al., 2025). In contrast, the present study revealed that CA2

place cells in FXS rats did not change their firing fields when social stimuli were presented. This

lack of social remapping in CA2 of FXS rats was not explained by reduced exploration of social

stimuli or a lack of olfactory discrimination in FXS rats. Furthermore, we found no differences in

firing rates, spatial information coding, and basic place field properties in CA2 place cells in FXS

rats, suggesting that impairments in CA2 place cell activity in FXS rats were largely specific to

responses to social stimuli.

To our knowledge, this study is the first to record CA2 place cells in a rodent model of FXS. We

found no abnormalities in baseline firing rates or spatial coding in CA2 neurons in FXS rats.

However, our experiments exclusively involved familiar environments and stimuli. Environmental

novelty has been shown to differentially affect CA1 place cells in FXS rats (Asiminas et al.,

2022). Specifically, CA1 place cells in FXS rats do not undergo normal experience-dependent

changes in spatial selectivity or firing rates (Asiminas et al., 2022). Therefore, it is possible that

CA2 place cell properties would differ between WT and FXS rats across experience in a novel

environment. Future experiments examining CA2 place cell firing patterns in FXS rats exploring

novel environments will be important to determine if CA2 place cell firing properties are

differentially affected by experience in WT and FXS rats.

Multiple studies have shown that CA2 place cells alter their firing fields during social interactions

(Alexander et al., 2016; Oliva et al., 2020; Robson et al., 2025), but the mechanism underlying

this social remapping has not yet been identified. Therefore, it is possible that impairments in

CA2 place cell remapping in FXS rats may be inherited from upstream regions or may be due to

impaired local computations within CA2. CA2 is unique as a hippocampal subregion in that it

contains an abundance of neuropeptide receptors, including oxytocin. In mice, blocking or

ablating oxytocin receptors in CA2 impairs social memory (Lin et al., 2018; Raam et al., 2017;

Tsai et al., 2022), and CA2 neurons are depolarized by activation of oxytocin receptors (Tirko et

al., 2018). It is therefore possible that the release of oxytocin in CA2 during social interactions

promotes changes in CA2 place cell firing patterns. If so, a dysregulation of oxytocin release,

reduced expression of oxytocin receptors, or impaired oxytocin signaling in CA2 of FXS rats

could contribute to the lack of remapping to social stimuli observed in FXS rats in this study. In

FXS rats in the present study, we identified reductions in oxytocin receptor expression levels

and impaired depolarizing responses to oxytocin in CA2 neurons. These oxytocinergic

impairments may contribute to deficient place cell coding of social stimuli in the following

manner. Depolarizing oxytocinergic inputs to CA2 may offset strong inhibitory inputs to CA2

neurons (Chevaleyre and Siegelbaum, 2010), thereby allowing some CA2 neurons to develop a

new place field in response to social stimuli. Considering that the overall level of activity in the

place cell network is thought to be homeostatically maintained (Kaleb et al., 2021), newly

activated place cells could trigger network competition and local inhibition within CA2, causing

some previously active cells to lose their place fields. In this way, activation of oxytocin

receptors on CA2 place cells could promote remapping to social stimuli in WT rats. In contrast,

diminished oxytocin receptor expression and weakened depolarizing responses to oxytocin

could hinder CA2 place cell remapping to social stimuli in FXS rats.

Oxytocin release in CA2 has not yet been characterized in FXS models, but reduced oxytocin

release may also potentially contribute to these deficits. Oxytocin receptors are overexpressed

in several subregions of the hippocampus, including CA2, in a separate rodent model of autism,

potentially as a compensatory mechanism for reduced oxytocin release (Bertoni et al., 2021).

Previous work has also shown that the oxytocin-mediated excitatory/inhibitory shift of GABA is

impaired in a mouse model of FXS (Tyzio et al., 2014), and that treatments that rescue this

impairment prenatally result in improvements in social behavior in adulthood (Eftekhari et al.,

2014). Related to this, an important question to be examined in future studies is whether

oxytocin administration (e.g., oxytocin uncaging; Ahmed et al., 2023) can rescue CA2 place cell

remapping in FXS rats.

CA2 receives direct inputs from other brain regions involved in social processing that may be

important for transmitting social information to CA2 and driving social responses in place cells.

The direct projection from the lateral entorhinal cortex to CA2 is necessary for social memory

(Dang et al., 2022; Lopez-Rojas et al., 2022). This direct projection arises from layer II

entorhinal cortical neurons (Dang et al., 2022). To our knowledge, layer II entorhinal cortical

neurons have not been examined in a rodent model of FXS, but neurons in other layers of the

entorhinal cortex show abnormal physiological properties in FXS mice (Deng et al., 2013; Deng

& Klyachko, 2016). Thus, an interesting hypothesis to test in future studies is the extent to which

abnormal physiology in the direct projection from the lateral entorhinal cortex to CA2 contributes

to impaired CA2 place cell remapping to social stimuli in FXS rats.

The function of CA2 place cell remapping to social stimuli remains unclear. Dorsal CA2, the

area from which the place cells in this study were recorded, projects directly to ventral CA1

(Meira et al., 2018; Okuyama et al., 2016), and this pathway is essential for social memory

(Meira et al., 2018; Tsai et al., 2022). Exogenous activation of this pathway rescues social

memory deficits in a mouse model of autism (Cope et al., 2023) and in an oxytocin receptor

knockout mouse (Tsai et al., 2022). Also, the direct projection from ventral CA1 to the prefrontal

cortex is important for social memory (Phillips et al., 2019; Sun et al., 2020). Hyperexcitability in

this pathway has been shown to contribute to social memory impairments in a mouse model of

autism (Phillips et al., 2019). While this pathway has not been specifically examined in FXS

models, FXS mice show reduced coherence between the ventral hippocampus and the

prefrontal cortex (Bhandari et al., 2024), providing a potential mechanistic link between impaired

CA2 place cell remapping in FXS rats and abnormal social behaviors.

However, the possibility remains that CA2 place cell remapping to social stimuli is

epiphenomenal. CA2 place cells also remap to the presentation of novel objects (Alexander et

al., 2016; Oliva et al., 2020), but optogenetic silencing of CA2 neurons does not impact

performance on a novel object recognition task (Oliva et al., 2020). Although inactivation of CA2

impairs social memory (Hitti & Siegelbaum, 2014; Leroy et al., 2017; Oliva et al., 2020;

Stevenson & Caldwell, 2014), it is difficult to establish if the change of firing fields in response to

social conditions is necessary for social memory or if a more general CA2 signal is sufficient. It

is possible that the impaired CA2 remapping to social stimuli in FXS rats may not functionally

contribute to abnormal social behaviors. Future studies are necessary to establish whether CA2

place cell remapping has functional consequences for social behaviors and, if so, how

impairments in CA2 place cell remapping may contribute to abnormal social behaviors in FXS

rats. Establishing these connections may help provide novel treatment targets that specifically

address social behavioral impairments in patients with FXS.

References

Ahmed, I. A., Liu, J.-J., Gieniec, K. A., Bair-Marshall, C. J., Adewakun, A. B., Hetzler, B. E., Arp,
C. J., Khatri, L., Vanwalleghem, G. C., Seidenberg, A. T., Cowin, P., Trauner, D., Chao,
M. V., Davis, F. M., Tsien, R. W., & Froemke, R. C. (2023). Optopharmacological tools for
precise spatiotemporal control of oxytocin signaling in the central nervous system and
periphery. *Research Square*, rs.3.rs-2715993. https://doi.org/10.21203/RS.3.RS-
2715993/V1

Alexander, G. M., Farris, S., Pirone, J. R., Zheng, C., Colgin, L. L., & Dudek, S. M. (2016). Social
and novel contexts modify hippocampal CA2 representations of space. *Nature
Communications*, 7(1), 1–14. https://doi.org/10.1038/ncomms10300

Arbab, T., Battaglia, F. P., Pennartz, C. M. A., & Bosman, C. A. (2018). Abnormal hippocampal
theta and gamma hypersynchrony produces network and spike timing disturbances in the
Fmr1-KO mouse model of Fragile X syndrome. *Neurobiology of Disease*, *114*(1), 65–73.
https://doi.org/10.1016/j.nbd.2018.02.011

Arbab, T., Pennartz, C. M. A., & Battaglia, F. P. (2018). Impaired hippocampal representation of
place in the Fmr1-knockout mouse model of fragile X syndrome. *Scientific Reports*, 8(1),
1–9. https://doi.org/10.1038/s41598-018-26853-z

Ashley, C. T., Wilkinson, K. D., Reines, D., & Warren, S. T. (1993). FMR1 protein: conserved RNP
family domains and selective RNA binding. *Science*, 262(5133), 563–566.
https://doi.org/10.1126/SCIENCE.7692601

Asiminas, A., Booker, S. A., Dando, O. R., Kozic, Z., Arkell, D., Inkpen, F. H., Sumera, A., Akyel,
I., Kind, P. C., & Wood, E. R. (2022). Experience-dependent changes in hippocampal
spatial activity and hippocampal circuit function are disrupted in a rat model of Fragile X
Syndrome. *Molecular Autism*, 13(1), 1–29. https://doi.org/10.1186/s13229-022-00528-z

Bakker, C. E., Verheij, C., Willemsen, R., van der Helm, R., Oerlemans, F., Vermey, M., Bygrave,
A., Hoogeveen, A. T., Oostra, B. A., Reyniers, E., De Boule, K., D'Hooge, R., Cras, P.,
van Velzen, D., Nagels, G., Martin, J. J., De Deyn, P. P., Darby, J. K., & Willems, P. J.
(1994). Fmr1 knockout mice: A model to study fragile X mental retardation. *Cell*, 78(1),
23–33. https://doi.org/10.1016/0092-8674(94)90569-X

Bertoni, A., Schaller, F., Tyzio, R., Gaillard, S., Santini, F., Xolin, M., Diabira, D., Vaidyanathan,
R., Matarazzo, V., Medina, I., Hammock, E., Zhang, J., Chini, B., Gaiarsa, J. L., &
Muscatelli, F. (2021). Oxytocin administration in neonates shapes hippocampal circuitry

and restores social behavior in a mouse model of autism. *Molecular Psychiatry*, 26(12),
7582–7595. https://doi.org/10.1038/s41380-021-01227-6

Bhandari, K., Kanodia, H., Donato, F., & Caroni, P. (2024). Selective vulnerability of the ventral
hippocampus-prelimbic cortex axis parvalbumin interneuron network underlies learning
deficits of fragile X mice. *Cell Reports*, 43(5), 114124.
https://doi.org/10.1016/j.celrep.2024.114124

Bodaleo, F., Tapia-Monsalves, C., Cea-Del Rio, C., Gonzalez-Billault, C., & Nunez-Parra, A.
(2019). Structural and functional abnormalities in the olfactory system of fragile x
syndrome models. *Frontiers in Molecular Neuroscience*, 12, 448197.
https://doi.org/10.3389/FNMOL.2019.00135/BIBTEX

Chevaleyre, V., & Siegelbaum, S. A. (2010). Strong CA2 pyramidal neuron synapses define a
powerful disynaptic cortico-hippocampal loop. Neuron, 66(4), 560-572.

Colgin, L. L., Moser, E. I., & Moser, M. B. (2008). Understanding memory through hippocampal
remapping. *Trends in Neurosciences*, 31(9), 469–477.
https://doi.org/10.1016/j.tins.2008.06.008

Cope, E. C., Wang, S. H., Waters, R. C., Gore, I. R., Vasquez, B., Laham, B. J., & Gould, E.
(2023). Activation of the CA2-ventral CA1 pathway reverses social discrimination
dysfunction in Shank3B knockout mice. *Nature Communications*, 14(1), 1–16.
https://doi.org/10.1038/s41467-023-37248-8

Dang, R., Zhou, Y., Zhang, Y., Liu, D., Wu, M., Liu, A., Jia, Z., Xie, W. (2022). Regulation of social
memory by lateral entorhinal cortical projection to dorsal hippocampal CA2. *Neuroscience*
Bulletin, 38(3), 318–322. https://doi.org/10.1007/s12264-021-00813-6

Deng, P. Y., & Klyachko, V. A. (2016). Increased persistent sodium current causes neuronal
hyperexcitability in the entorhinal cortex of Fmr1 knockout mice. *Cell Reports*, 16(12),
3157–3166. https://doi.org/10.1016/j.celrep.2016.08.046

Deng, P. Y., Rotman, Z., Blundon, J. A., Cho, Y., Cui, J., Cavalli, V., Zakharenko, S. S., &
Klyachko, V. A. (2013). FMRP regulates neurotransmitter release and synaptic information
transmission by modulating action potential duration via BK channels. *Neuron*, 77(4), 696–
711. https://doi.org/10.1016/j.neuron.2012.12.018

Donahue, M. M., Robson, E., & Colgin, L. L. (2025). Hippocampal place cell sequences are
impaired in a rat model of Fragile X Syndrome. *Journal of Neuroscience*, 45(15).

Eftekhari, S., Shahrokhi, A., Tsintsadze, V., Nardou, R., Brouchoud, C., Conesa, M., Burnashev,
N., Ferrari, D. C., & Ben-Ari, Y. (2014). Response to comment on "Oxytocin-mediated
GABA inhibition during delivery attenuates autism pathogenesis in rodent offspring."
Science, 346(6206). https://doi.org/10.1126/SCIENCE.1256009/ASSET/42325DA6-
71A9-4A0B-8740-0AB20FCDBC61/ASSETS/GRAPHIC/346_176C_F1.JPEG

Froemke, R. C., & Young, L. J. (2021). Oxytocin, neural plasticity, and social behavior. *Annual*
Review of Neuroscience, 44(1), 359-381.

Hamilton, S. M., Green, J. R., Veeraragavan, S., Yuva, L., McCoy, A., Wu, Y., Warren, J., Little,
L., Ji, D., Cui, X., Weinstein, E., & Paylor, R. (2014). Fmr1 and Nlgn3 knockout rats: Novel
tools for investigating autism spectrum disorders. *Behavioral Neuroscience*, 128(2), 103–
109. https://doi.org/10.1037/a0035988

Hitti, F. L., & Siegelbaum, S. A. (2014). The hippocampal CA2 region is essential for social
memory. *Nature*, 508(1), 88–92. https://doi.org/10.1038/nature13028

Jin, P., & Warren, S. T. (2000). Understanding the molecular basis of fragile X syndrome. *Human*
Molecular Genetics, 9(6), 901–908. https://doi.org/10.1093/HMG/9.6.901

Kaleb, K., Pedrosa, V., & Clopath, C. (2021). Network-centered homeostasis through inhibition
maintains hippocampal spatial map and cortical circuit function. Cell Reports, 36(8).

Kau, A. S. M., Tierney, E., Bukelis, I., Stump, M. H., Kates, W. R., Trescher, W. H., & Kaufmann,
W. E. (2004). Social behavior profile in young males with fragile X syndrome:
Characteristics and specificity. *American Journal of Medical Genetics*, 126A(1), 9–17.
https://doi.org/10.1002/AJMG.A.20218

Kaufmann, W. E., Cortell, R., Kau, A. S. M., Bukelis, I., Tierney, E., Gray, R. M., Cox, C., Capone,
G. T., & Stanard, P. (2004). Autism spectrum disorder in fragile X syndrome:
Communication, social interaction, and specific behaviors. *American Journal of Medical*
Genetics, 129A(3), 225–234. https://doi.org/10.1002/AJMG.A.30229

Kuruppath, P., Xue, L., Pouille, F., Jones, S. T., & Schoppa, N. E. (2023). Hyperexcitability in the
olfactory bulb and impaired fine odor discrimination in the Fmr1 KO mouse model of
Fragile   X   Syndrome.   *Journal   of   Neuroscience*,   43(48),   8243–8258.
https://doi.org/10.1523/JNEUROSCI.0584-23.2023

Leroy, F., Brann, D. H., Meira, T., & Siegelbaum, S. A. (2017). Input-timing-dependent plasticity
in the hippocampal CA2 region and its potential role in social memory. *Neuron*, 95(5),
1089-1102.e5. https://doi.org/10.1016/j.neuron.2017.07.036

Lesniak-Karpiak, K., Mazzocco, M. M. M., & Ross, J. L. (2003). Behavioral assessment of social
anxiety in females with Turner or Fragile X Syndrome. *Journal of Autism and*
Developmental   Disorders,   33(1),   55–67.
https://doi.org/10.1023/A:1022230504787/METRICS

Lin, Y. T., Hsieh, T. Y., Tsai, T. C., Chen, C. C., Huang, C. C., & Hsu, K. Sen. (2018). Conditional
deletion of hippocampal CA2/CA3a oxytocin receptors impairs the persistence of long-
term social recognition memory in mice. *Journal of Neuroscience*, 38(5), 1218–1231.
https://doi.org/10.1523/JNEUROSCI.1896-17.2017

Liu, Z. H., & Smith, C. B. (2009). Dissociation of social and nonsocial anxiety in a mouse model
of   fragile   X   syndrome.   *Neuroscience   Letters*,   454(1),   62–66.
https://doi.org/10.1016/J.NEULET.2009.02.066

Lopez-Rojas, J., de Solis, C. A., Leroy, F., Kandel, E. R., & Siegelbaum, S. A. (2022). A direct
lateral entorhinal cortex to hippocampal CA2 circuit conveys social information required
for   social   memory.   *Neuron*,   110(9),   1559-1572.e4.
https://doi.org/10.1016/j.neuron.2022.01.028

Lu, L., Igarashi, K. M., Witter, M. P., Moser, E. I., & Moser, M. B. (2015). Topography of place
maps along the CA3-to-CA2 axis of the hippocampus. *Neuron*, 87(5), 1078-1092.
https://doi.org/10.1016/j.neuron.2015.07.007

McNaughton, C. H., Moon, J., Strawderman, M. S., Maclean, K. N., Evans, J., & Strupp, B. J.
(2008). Evidence for social anxiety and impaired social cognition in a mouse model of
Fragile   X   Syndrome.   *Behavioral   Neuroscience*,   122(2),   293–300.
https://doi.org/10.1037/0735-7044.122.2.293

Meira, T., Leroy, F., Buss, E. W., Oliva, A., Park, J., & Siegelbaum, S. A. (2018). A hippocampal
circuit linking dorsal CA2 to ventral CA1 critical for social memory dynamics. *Nature*
Communications, 9(1), 1–14. https://doi.org/10.1038/s41467-018-06501-w

Merenstein, S. A., Sobesky, W. E., Taylor, A. K., Riddle, J. E., Ran, H. X. ", & Hagerman, R. J.
(1996). Molecular-clinical correlations in males with an expanded FMR1 mutation.
American Journal of Medical Genetics, 64, 388–394. https://doi.org/10.1002/(SICI)1096-
8628(19960809)64:2

Mientjes, E. J., Nieuwenhuizen, I., Kirkpatrick, L., Zu, T., Hoogeveen-Westerveld, M., Severijnen,
L., Rifé, M., Willemsen, R., Nelson, D. L., & Oostra, B. A. (2006). The generation of a
conditional Fmr1 knock out mouse model to study Fmrp function in vivo. *Neurobiology of*
Disease, 21(3), 549–555. https://doi.org/10.1016/J.NBD.2005.08.019

Mineur, Y. S., Huynh, L. X., & Crusio, W. E. (2006). Social behavior deficits in the Fmr1 mutant
mouse. *Behavioural Brain Research*, 168(1), 172–175.
https://doi.org/10.1016/J.BBR.2005.11.004

Mitre, M., Marlin, B. J., Schiavo, J. K., Morina, E., Norden, S. E., Hackett, T. A., Aoki, C. J., Chao,
M. V., Froemke, R. C. (2016). A distributed network for social cognition enriched for
oxytocin receptors. *Journal of Neuroscience*, 36(8), 2517–2535.
https://doi.org/10.1523/jneurosci.2409-15.2016

Nitenson, A. S., Stackpole, E. E., Truszkowski, T. L. S., Midroit, M., Fallon, J. R., & Bath, K. G.
(2015). Fragile X Mental Retardation Protein regulates olfactory sensitivity but not odorant
discrimination. *Chemical Senses*, 40(5), 345–350.
https://doi.org/10.1093/CHEMSE/BJV019

O'Keefe, J., & Dostrovsky, J. (1971). The hippocampus as a spatial map. Preliminary evidence
from unit activity in the freely-moving rat. *Brain Research*, 34(1), 171–175.
https://doi.org/10.1016/0006-8993(71)90358-1

Oddi, D., Subashi, E., Middei, S., Bellocchio, L., Lemaire-Mayo, V., Guzmán, M., Crusio, W. E.,
D'Amato, F. R., & Pietropaolo, S. (2014). Early social enrichment rescues adult behavioral
and brain abnormalities in a mouse model of Fragile X Syndrome.
Neuropsychopharmacology, 40(5), 1113–1122. https://doi.org/10.1038/npp.2014.291

Okuyama, T., Kitamura, T., Roy, D. S., Itohara, S., Tonegawa, S. (2016). Ventral CA1 neurons
store social memory. *Science*. https://doi.org/10.1126/science.aaf7003

Oliva, A. (2022). CA2 physiology underlying social memory. *Current Opinion in Neurobiology*, 77,
102642. https://doi.org/10.1016/j.conb.2022.102642

Oliva, A., Fernández-Ruiz, A., Leroy, F., & Siegelbaum, S. A. (2020). Hippocampal CA2 sharp-
wave ripples reactivate and promote social memory. *Nature*, 587(7833), 264–269.
https://doi.org/10.1038/s41586-020-2758-y

Pagani, J. H., Zhao, M., Cui, Z., Avram, S. K. W., Caruana, D. A., Dudek, S. M., Young, W. S.
(2015). Role of the vasopressin 1b receptor in rodent aggressive behavior and synaptic
plasticity in hippocampal area CA2. *Molecular Psychiatry*, 20(4), 490–499.
https://doi.org/10.1038/mp.2014.47

Phillips, M. L., Robinson, H. A., & Pozzo-Miller, L. (2019). Ventral hippocampal projections to the
medial prefrontal cortex regulate social memory. *eLife*, 8.
https://doi.org/10.7554/ELIFE.44182

Pieretti, M., Zhang, F., Fu, Y. H., Warren, S. T., Oostra, B. A., Caskey, C. T., & Nelson, D. L.
(1991). Absence of expression of the FMR-1 gene in fragile X syndrome. *Cell*, 66(4), 817–
822. https://doi.org/10.1016/0092-8674(91)90125-I

Raam, T., McAvoy, K. M., Besnard, A., Veenema, A. H., Sahay, A. (2017). Hippocampal oxytocin
receptors are necessary for discrimination of social stimuli. *Nature Communications*, 8(1),
2001. https://doi.org/10.1038/s41467-017-02173-0

Richter, J. D., & Zhao, X. (2021). The molecular biology of FMRP: new insights into fragile X
syndrome. *Nature Reviews Neuroscience*, 22(4), 209–222.
https://doi.org/10.1038/s41583-021-00432-0

Robson, E., Donahue, M. M., Mably, A. J., Demetrovich, P. G., Hewitt, L. T., & Colgin, L. L. (2025).
Social odors drive hippocampal CA2 place cell responses to social stimuli. *Progress in*
neurobiology, 245, 102708.

Saxena, K., Webster, J., Hallas-Potts, A., Mackenzie, R., Spooner, P. A., Thomson, D., Kind, P.,
Chatterji, S., & Morris, R. G. M. (2018). Experiential contributions to social dominance in
a rat model of fragile-X syndrome. *Proceedings of the Royal Society B: Biological*
Sciences, 285(1880). https://doi.org/10.1098/RSPB.2018.0294

Siomi, H., Siomi, M. C., Nussbaum, R. L., & Dreyfuss, G. (1993). The protein product of the fragile
X gene, FMR1, has characteristics of an RNA-binding protein. *Cell*, 74(2), 291–298.
https://doi.org/10.1016/0092-8674(93)90420-U

Skaggs, W. E., McNaughton, B. L., Wilson, M. A., & Barnes, C. A. (1996). Theta phase precession
in hippocampal neuronal populations and the compression of temporal sequences.
Hippocampus, 6(2), 149–172. https://doi.org/10.1002/(sici)1098-1063(1996)6:2<149::aid-
hipo6>3.0.co;2-K

Spencer, C. M., Alekseyenko, O., Serysheva, E., Yuva-Paylor, L. A., & Paylor, R. (2005). Altered
anxiety-related and social behaviors in the Fmr1 knockout mouse model of fragile X
syndrome. *Genes, Brain and Behavior*, 4(7), 420–430. https://doi.org/10.1111/J.1601-
183X.2005.00123.X

Stevenson, E. L., & Caldwell, H. K. (2014). Lesions to the CA2 region of the hippocampus impair
social memory in mice. *European Journal of Neuroscience*, 40(9), 3294–3301.
https://doi.org/10.1111/ejn.12689

Sun, Q., Li, X., Li, A., Zhang, J., Ding, Z., Gong, H., & Luo, Q. (2020). Ventral hippocampal-
prefrontal interaction affects social behavior via parvalbumin positive neurons in the
medial prefrontal cortex. *iScience*, 23(3), 100894.
https://doi.org/10.1016/j.isci.2020.100894

Talbot, Z. N., Sparks, F. T., Dvorak, D., Curran, B. M., Alarcon, J. M., & Fenton, A. A. (2018).
Normal CA1 place fields but discoordinated network discharge in a Fmr1-null mouse
model of Fragile X Syndrome. *Neuron*, 97(3), 684-697.e4.
https://doi.org/10.1016/j.neuron.2017.12.043

Tian, Y., Yang, C., Shang, S., Cai, Y., Deng, X., Zhang, J., Shao, F., Zhu, D., Liu, Y., Chen, G.,
Liang, J., Sun, Q., Qiu, Z., & Zhang, C. (2017). Loss of FMRP impaired hippocampal long-
term plasticity and spatial learning in rats. *Frontiers in Molecular Neuroscience*, 10, 269.
https://doi.org/10.3389/fnmol.2017.00269

Tirko, N. N., Eyring, K. W., Carcea, I., Mitre, M., Chao, M. V., Froemke, R. C., Tsien, R. W. (2018). Oxytocin transforms firing mode of CA2 hippocampal neurons. *Neuron*, 100(3), 593-608.e3. https://doi.org/10.1016/j.neuron.2018.09.008

Tsai, T.-C., Fang, Y.-S., Hung, Y.-C., Hung, L.-C., Hsu, K.-S. (2022). A dorsal CA2 to ventral CA1 circuit contributes to oxytocinergic modulation of long-term social recognition memory. *Journal of Biomedical Science*, 29(1), 50. https://doi.org/10.1186/s12929-022-00834-x

Tyzio, R., Nardou, R., Ferrari, D. C., Tsintsadze, T., Shahrokhi, A., Eftekhari, S., Khalilov, I., Tsintsadze, V., Brouchoud, C., Chazal, G., Lemonnier, E., Lozovaya, N., Burnashev, N., & Ben-Ari, Y. (2014). Oxytocin-mediated GABA inhibition during delivery attenuates autism pathogenesis in rodent offspring. *Science*, 343(6171), 675–679. https://doi.org/10.1126/SCIENCE.1247190/SUPPL_FILE/675.MP3

Verkerk, A. J., Pieretti, M., Sutcliffe, J. S., Fu, Y. H., Kuhl, D. P., Pizzuti, A., ... & Warren, S. T. (1991). Identification of a gene (FMR-1) containing a CGG repeat coincident with a breakpoint cluster region exhibiting length variation in fragile X syndrome. Cell, 65(5), 905-914.

Williams, T. A., Porter, M. A., & Langdon, R. (2014). Social approach and emotion recognition in Fragile X Syndrome. *American Journal on Intellectual and Developmental Disabilities*, 119(2), 133–150. https://doi.org/10.1352/1944-7558-119.2.133

Wong, H., Hooper, A. W. M., Niibori, Y., Lee, S. J., Hategan, L. A., Zhang, L., Karumuthil-Melethil, S., Till, S. M., Kind, P. C., Danos, O., Bruder, J. T., & Hampson, D. R. (2020). Sexually dimorphic patterns in electroencephalography power spectrum and autism-related behaviors in a rat model of fragile X syndrome. *Neurobiology of Disease*, 146, 105118. https://doi.org/10.1016/J.NBD.2020.105118

Yang, M., & Crawley, J. N. (2009). Simple behavioral assessment of mouse olfaction. *Current Protocols in Neuroscience*, 48(1), 8.24.1-8.24.12. https://doi.org/10.1002/0471142301.NS0824S48

Zheng, C., Bieri, K. W., Hsiao, Y. T., & Colgin, L. L. (2016). Spatial sequence coding differs during slow and fast gamma rhythms in the hippocampus. *Neuron*, 89(2), 398–408. https://doi.org/10.1016/j.neuron.2015.12.005

Figure Legends

Figure 1. Behavioral task. Rats explored a 1-meter x 1-meter open field arena across four 20-minute sessions per day (Sessions A, B, B', A'). A stimulus cage, which was a standard rat housing cage, was placed in a corner of the arena. In the first and final sessions (A and A') of both experimental conditions (i.e., Odor and Visual + Odor) and in all four sessions of the Control condition, a clean cage containing clean bedding was presented. In the middle two sessions (B and B') of the experimental conditions, a cage containing social stimuli was

presented. In the Odor condition, the soiled bedding of a familiar rat or rats served as the social

stimuli. In the Visual + Odor condition, a cage containing a familiar rat and the rat's soiled

bedding was presented.

Figure 2. CA2 place cells in WT and FXS rats. Color-coded firing rate maps are shown for all

place cells recorded on a single example tetrode across all conditions for WT (A) and FXS (B)

rats. Rate maps are shown scaled to the maximal firing rate (shown inset) of each cell across all

sessions. White pixels indicate places that were not visited during a session.

Figure 3. CA2 place cells changed their firing fields in response to social stimuli in WT,

but not FXS, rats. (A) The estimated means of spatial correlation coefficients from the

generalized linear mixed model are shown for each genotype across each condition for all

session pairs combined. Dots represent estimated mean values, and error bars represent the

estimated 95% confidence intervals. Spatial correlation coefficients were lower in the Odor and

Visual + Odor conditions compared to the Control condition for WT rats. Spatial correlation

coefficients did not differ across social and control conditions in FXS rats. (B-F) Spatial

correlation measures are shown for the entire sample of CA2 place cells for all pairs of sessions

across all conditions. Each marker represents a spatial correlation value for an individual place

cell. Different symbols are used for CA2 place cells recorded from different rats. Data from WT

and FXS rats are shown with blue and red symbols, respectively.

Figure 4. Basic firing and spatial coding properties of CA2 place cells did not differ

between WT and FXS rats. For all plots, individual dots represent the mean value for each rat,

and error bars represent 95% confidence intervals across all place cells within a rat. Data from

WT rats and FXS rats are shown in blue and red, respectively. (A) Spatial information did not

differ between WT and FXS rats across sessions and conditions. (B-C) CA2 place cells in FXS

rats showed normal firing rates. Peak firing rates (B) and average in-field firing rates (C) of CA2

place cells did not differ between WT and FXS rats across sessions and conditions. (D) CA2

place field sizes were similar between genotypes and across sessions and experimental

conditions. (E) The average number of place fields per place cell did not differ between

genotypes across all sessions and conditions.

Figure 5. WT and FXS rats explored social stimuli similarly. (A-B) Heat maps show mean

exploration times across locations in the arena, with the stimulus cage shown in the top right

corner. Exploration times were calculated for the first 2 minutes of each individual session,

averaged within each rat, and then averaged across rats within each genotype group for WT (A)

and FXS (B) rats. (C) Time spent exploring locations close to the cage (within 12 cm) was

calculated for each session and then averaged within a rat. Individual dots represent the mean

exploration time for each rat, and error bars represent 95% confidence intervals across all

sessions within a rat. Exploration times are shown in blue for WT rats and in red for FXS rats.

Both WT and FXS rats increased their exploration time of the stimulus cage in session B for the

Odor and Visual + Odor conditions but not the Control condition.

Figure 6. FXS rats could detect and differentiate between social odors. The normalized

times spent sniffing a cotton-tip applicator is shown for trials with no odor ("null" trials, N1-N3),

an initially unfamiliar social odor ("habituation" trials, H1-H4), and a second unfamiliar social

odor ("dishabituation" trial, D). Bars indicate mean values within genotype and trial and dots

indicate values from individual rats. Both WT (blue) and FXS (red) rats increased the time spent

sniffing between null trial 3 (N3) and habituation 1 (H1), indicating that both WT and FXS rats

successfully detected the presentation of a social odor. Both WT and FXS rats decreased the

time spent sniffing across habituation trials (H1-4), indicating that FXS rats habituated normally

to repeated presentations of the same odor. Both WT and FXS rats increased time spent

sniffing between habituation trial 4 (H4) and the dishabituation trial (D), indicating that both WT

and FXS rats distinguished between different social odors.

Figure 7. Low oxytocin receptor expression levels in CA2 of FXS rats. (A-B) Example images of CA2 immunostaining in WT (A) and FXS (B) rats for a CA2 marker (Purkinje cell protein 4 (PCP4), red, top row) and an oxytocin receptor antibody (OXTR-2, green, middle row). DAPI nuclear staining is shown in blue. Merged images for PCP4 and OXTR-2 immunostaining are shown in the bottom row. Scale bar: 500 µm. (C) Plots show fluorescence estimates (mean ± 95% confidence intervals) across hippocampal subregions for WT (blue) and FXS (red) rats. Oxytocin receptor expression across subregions was lower in FXS rats than in WT rats.

Figure 8. Oxytocin depolarized the resting membrane potential in CA2 neurons in WT rats but not FXS rats. Resting membrane potential (V_M) measurements are shown during baseline and after 15 minutes of 400 nM TGOT application. Dots indicate data from individual cells. TGOT significantly depolarized the V_M of WT CA2 neurons (blue; n = 6 cells from 4 rats) but not FXS CA2 neurons (red; n = 6 cells from 4 rats).

Figure S1. Histological identification of CA2. (A) In four rats, brains were immunostained with the CA2 marker STEP (brown) and cresyl violet (purple). The boundaries of CA2 are denoted by black lines in both images. Subregions CA1, CA2, and CA3 are labeled. Right: Two example tetrode tracks are shown. The tetrode circled in red recorded place cells from CA2, and the tetrode circled in blue recorded place cells from CA3. (B) In rats in which STEP staining did not clearly differentiate the boundaries of CA2, identification of CA2 was aided by its characteristic increase in cell body size. Boundaries of CA2 are indicated by the black arrows. A CA2 tetrode track is circled in red. (C) In eight rats, brains were immunostained with CA2-expressed protein (PCP4; red) and DAPI (blue). A CA2 tetrode track is shown.

Figure S2. CA2 place cells did not significantly alter their firing rates in response to social stimuli in WT or FXS rats. (A) The estimated means of rate overlap values from a generalized linear mixed model are shown for each genotype across each condition for all session pairs combined. Dots represent estimated mean values, and error bars represent the

1063   estimated 95% confidence intervals. Rate overlap values did not differ across genotypes or

1064   conditions. (B-F) Rate overlap values are shown for the entire sample of CA2 place cells for all

1065   pairs of sessions across all conditions. Each marker represents a rate overlap value for an

1066   individual place cell. Different symbols are used for CA2 place cells recorded from different rats.

1067   Values from WT and FXS rats are shown with blue and red symbols, respectively.

Figure 1

A

B

FXS

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A   WT

B   FXS

C

Figure 7

Figure 8

A

B

C

Figure S1

Figure S2