Biological context modulates virus–host dynamics and diversification

Rodrigo Sanchez-Martinez[1], Esther Rubio-Portillo[1], Laura Medina-Ruiz[2], Jonás Sarasa[2], María Enciso[2], Fernando Santos[1,3]*, Josefa Antón[1,3,4]*

Affiliations

[1]Department of Physiology, Genetics and Microbiology, University of Alicante, San Vicent del Raspeig, Alicante, Spain

[2]IGLS, Alicante, Spain

[3]Institute of Health and Biomedical Research of Alicante (ISABIAL), Alicante, Spain

[4]Multidisciplinary Institute of Environmental Studies Ramon Margalef, Alicante, Spain

*Corresponding authors: fernando.santos@ua.es; anton@ua.es

ABSTRACT

Virus–bacterial interactions are fundamental to microbial ecology and evolution, but they have been studied mostly under simplified laboratory conditions. To better understand how ecological complexity shapes these dynamics, we examined the extreme halophilic bacterium *Salinibacter ruber* strain M1 and the EM1 virus in the presence of additional *Sal. ruber* strains and viruses. In the short term, the presence of other strains delayed lysis and reduced EM1 virus production, indicating that community composition directly affects viral replication. In the long term, both *Sal. ruber* M1 and the EM1 virus persisted across all experimental conditions, but their evolutionary responses differed. The EM1 virus showed an increased mutation rate, reduced infectivity against the native host, and expanded host range when other viruses were present, suggesting a previously unrecognized form of virus–virus interaction, in which coexisting viruses influence each other's evolutionary paths promoting viral diversification. In contrast, *Sal. ruber* M1 exhibited higher mutation rates evolving with other strains, indicating that in our system, intraspecific competition, rather than viral pressure, drives bacterial evolution. These findings demonstrate that incorporating biological complexity reveals distinct selective pressures acting on hosts and viruses, and is therefore essential for accurately predicting virus–host evolution in natural microbial ecosystems.

INTRODUCTION

Natural ecosystems are highly complex networks where multiple biotic and abiotic factors interact dynamically (Fuhrman et al., 2015; Levin, 1998). This complexity makes it challenging to predict ecological and evolutionary phenomena since experimental studies are generally conducted under simplified laboratory conditions which do not reflect natural dynamics (Calisi & Bentley, 2009; Chen et al., 2023; Roberts et al., 2015; Zimmer & Dorea, 2023). To advance our understanding of microbial processes and predict how microbial communities respond to environmental changes or selective pressures, it is crucial to design experiments that more faithfully simulate natural conditions, incorporating the diversity of species and interactions that characterize real ecosystems (Benton et al., 2007; Blazanin & Turner, 2021; Castledine & Buckling, 2024; Jessup et al., 2004).

Viruses act as major eco-evolutionary forces in microbial ecosystems. As the most abundant biological entities on Earth, they play a central role in regulating microbial communities through

diverse infection cycles that impact nutrient cycling, genetic diversity, and ecosystem structure
(Fierer, 2017; Middelboe & Brussaard, 2017; Rohwer & Thurber, 2009). Virus–host interactions
also drive microbial evolution, triggering adaptations that influence community composition and
function. Despite their importance, many aspects of these interactions remain poorly understood,
particularly under complex, natural conditions. Although numerous studies have explored virus–
host interactions, most have relied on simplified experimental systems involving a single virus–
host pair under controlled conditions. As a result, our understanding of how these interactions
operate in more realistic settings remains limited. Investigating virus–host dynamics in
environments that better mimic natural complexity is essential to fully grasp their ecological and
evolutionary roles (Blazanin & Turner, 2021; Castledine & Buckling, 2024; Chevallereau et al.,
2022; Koskella et al., 2022). While some recent efforts have incorporated greater biological
complexity (Alseth et al., 2019, 2024; De Sordi et al., 2017; Gómez & Buckling, 2011, 2013;
Middelboe et al., 2001; Mumford & Friman, 2017), such studies remain scarce. This persistent gap
limits our ability to understand and predict how virus–host interactions shape microbial
communities in natural ecosystems.

Hypersaline aquatic environments, which constitute approximately half of continental waters,
harbor microbial communities that, at extreme salinities, are dominated by archaea (Burns et al.,
2007; Shiklomanov, 1998; Ventosa, 2006). At these salinities, *Salinibacter ruber* emerges as the
main bacterial species (Antón et al., 2002; Gomariz et al., 2015; Oren, 2011). Although *Sal. ruber*
is as halophilic as the dominant archaea at close-to-saturation salinities, it is consistently less
abundant, representing between 1% and 10% of total cells, indicating that it could be subjected to
intense selective pressure (Mora-Ruiz et al., 2018; Ventosa et al., 2015). Cultivation and
metagenomic studies reveal a high degree of intraspecific genomic diversity in *Sal. ruber*,
including the coexistence of distinct phylogroups and adaptation to local microenvironments
(Viver et al., 2015, 2023). These hypersaline habitats also harbor some of the highest recorded
viral concentrations, with virus-like particles (VLP) reaching up to 10^{10} VLP/ml and virus-to-cell
ratios ranging from 10 to 100 (Di Meglio et al., 2016; Santos et al., 2012). Given its environmental
ubiquity, cultivability, and intraspecific variation, likely shaped in part by viral predation, *Sal.*
ruber constitutes a valuable model for studying microbial ecology and virus–host interactions.

In this work, we have addressed how the ecology and evolution of a virus–host pair (i.e., *Sal. ruber* strain M1 and the EM1 virus) are affected by the presence of additional *Sal. ruber* strains and their viruses. Through experiments in which biological complexity is increased, we have demonstrated that: (i) the presence of other strains and viruses delays lysis and reduces EM1 virion production; and that (ii) the biological context influences the long-term evolution of the virus–host pair and promotes changes in viral infectivity and host range, driven by a high mutation rate under more complex contexts. These findings highlight the importance of considering biological complexity in studies of virus–host interactions. Our work not only contributes to a better understanding of these dynamics in natural systems but also suggests this complexity to be considered in practical applications such as phage therapy and microbial community engineering.

RESULTS AND DISCUSSION

Five experiments were conducted in parallel to evaluate the effects of the presence of *Sal. ruber* strains and viruses on the interactions and long-term phenotypic and genetic changes (evolution) of *Sal. ruber* strain M1 and its M1EM1 virus (referred to as EM1 hereafter) (Fig. 1a). In experiment 1, *Sal. ruber* strain M1 was cultured alone. In experiment 2, *Sal. ruber* strains M8, M31, and P18 were added to the M1 culture. These two experiments served as virus-free controls. In experiment 3, M1 was infected with its EM1 virus. Experiment 4 included M1 infected with EM1, along with the presence of strains M8, M31, and P18. Finally, experiment 5 included the same components as experiment 4, with the addition of *Sal. ruber* M31CC1 and M31CR41-2 viruses (referred to as CC1 and CR41-2, respectively), which infect both *Sal. ruber* strains M31 and M8, and M8CR-4 virus (i.e. CR4), which infects *Sal. ruber* strain M8 (Villamor et al., 2018) (Supplementary Table 1).

A short-term experiment, which included the first 9 days of incubation (corresponding to ~14 generations), evaluated several factors derived from the M1-EM1 infection process under different contexts: lysis timing, virions produced, infective viruses and community dynamics. Thereafter, all cultures were periodically transferred to fresh medium and maintained for 150 days. This constituted the "long-term experiment", aimed at studying possible evolutionary changes in the virus–host pair, while continuing to monitor key ecological and phenotypic parameters (Fig. 1b).

Increased biological complexity leads to a delay in lysis of *Sal. ruber* M1 and a reduced production of EM1 virions

In the short-term experiment, *Sal. ruber* exhibited its characteristic growth pattern in experiments 1 and 2, with a slightly higher optical density (OD) in experiment 2 (Fig. 2a; Extended Data Fig. 1a). In infection experiment 3, lysis of *Sal. ruber* M1 was observed, followed by a recovery of the population, a patter that mimic previous findings from our laboratory, in which that recovery was mainly attributed to the emergence of pseudolysogens (Sanchez-Martinez et al., 2025). In experiments 4 and 5, this OD recovery was observed earlier, which could be attributed to the growth of uninfected strains.

The production of infectious EM1 viral particles, measured as plaque-forming-units (PFU) on *Sal. ruber* M1, was significantly delayed in experiments 4 and 5 compared to experiment 3 (*t*-test p-value < 0.0001) (Fig. 2b), indicating that the presence of additional *Sal. ruber* strains delayed the lysis of strain M1 by the EM1 virus. After 96 hours, a decline in the number of infectious particles was observed, more pronounced in experiments 4 and 5 than in experiment 3 (p-value < 0.01). In all the experiments a slight decline in the number of PFUs was observed over time, with no significant differences between experiments in the number of infectious EM1 viruses produced between 6 and 9 days. Total extracellular virion concentration, measured by flow cytometry, reached 9.7×10^{10} VLP/ml within 48 hours in experiment 3, whereas in experiment 4 this concentration was 4.2×10^8 VLP/ml, a statistically significant reduction (p-value < 0.0001) (Fig. 2c), consistent with the observations of PFU counts. Experiment 5 was excluded from this figure, as the presence of multiple coexisting viruses prevents attributing VLP counts specifically to EM1 (Extended Data Fig. 1b).

High throughput microfluidics-based qPCR revealed that the concentration of the M1 strain (measured as genomes/ml) varied across experiments and time points (Fig. 3a). At 48 hours, M1 abundance dropped significantly in experiment 3 (3.9×10^6 genomes/ml), indicating active lysis of the population, while it remained at higher numbers in experiments 4 (1.2×10^9 genomes/ml) and 5 (7.6×10^8 genomes/ml) (p-value < 0.0001). By 72 hours, M1 levels decreased in all experiments, maintaining low values in experiment 3 (3.8×10^6 genomes/ml), and showing

delayed but evident reductions in experiments 4 and 5 (2.2×10^6 and 5.2×10^6 genomes/ml,

respectively) (Fig. 3a; Supplementary Fig. 1).

In the case of the bacterial community, in experiment 4 a decrease in the concentration of M8,

M31, and P18 strains was observed at 50 hours, despite the absence of their viruses (Fig. 3a). This

pattern suggests that the growth of these strains was transiently inhibited after *Sal. ruber* M1 lysis.

To test this, virus-free lysate from an M1-infected culture was added to pure cultures of M8, M31,

and P18. A clear, transient growth inhibition was observed, confirming that the M1 lysate affected

the growth of the other strains (Extended Data Fig. 2). Similar results have been reported in other

studies, where intracellular bacterial contents released upon lysis can inhibit the growth of other

bacteria (Penchuk et al., 2023; Smakman & Hall, 2022). In experiment 5, M8 showed minimal

growth. As the only strain infected by three viruses (Supplementary Table 1), M8 likely could not

withstand the high viral pressure, explaining its poor performance under these conditions.

Regarding the extracellular viruses measured by microfluidics-based qPCR, a lower concentration

of EM1 was observed at 48 hours in both experiments 4 and 5 (2.2×10^8 and 3.3×10^8 genomes/ml,

respectively) compared with experiment 3 (4.2×10^9 genomes/ml) (p-value < 0.0001) (Fig. 3b;

Supplementary Fig. 2), confirming that increased community complexity delays the infection and

lysis of *Sal. ruber* M1 by the EM1 virus. By 216 hours (day 9), experiment 4 yielded significantly

lower VLP concentration (2.4×10^{10} VLP/ml) than experiment 3 (2.8×10^{10} VLP/ml) (p-value <

0.05) (Fig. 2c). Microfluidics-based qPCR measurements also revealed that the concentration of

extracellular EM1 viruses in experiments 4 and 5 (7.6×10^9 and 3.4×10^9 genomes/ml,

respectively) was significantly lower than in experiment 3 (2.7×10^{10} genomes/ml) at the same

time point (p-value < 0.01), supporting the flow cytometry results (Fig. 3b). These findings

indicate that the presence of additional *Sal. ruber* strains reduced the overall production of EM1,

in good agreement with previous studies showing that the presence of other bacterial species can

lead to lower viral densities (Alseth et al., 2019; Mumford & Friman, 2017). In contrast, a recent

study found no differences in lysis timing or viral titers produced by *Pseudomonas aeruginosa*

PA14 when exposed to competing bacterial strains (Alseth et al., 2024).

Experiment 5 also revealed distinct patterns in the production of other extracellular viruses. CC1

virus, which infects both *Sal. ruber* M8 and M31, showed sustained high titers from 50 hours

171 onward, consistent with ongoing replication in its hosts (Fig. 3b). CR41-2 virus, which infects the

172 same strains as CC1 (M8 and M31), was only detected on days 1 and 4. Adsorption assays revealed

173 that CR41-2 adsorbs significantly less efficiently than CC1 to M31 (Supplementary Fig. 3). Thus,

174 CC1 may have outcompeted CR41-2 for access to M31, thereby limiting CR41-2 infection. This

175 observation aligns with a well-established principle of phage biology, that considers adsorption

176 efficiency as a critical determinant of successful phage infection and subsequent replication

177 (Abedon, 2023). Such virus–virus competition for shared hosts can strongly shape infection

178 dynamics, allowing more efficient viruses to suppress the adsorption of less competitive ones

179 (Bürkle et al., 2025; Turner, 2005). Finally, CR4 virus, which exclusively infects M8, exhibited

180 an early increase in viral concentration followed by a sharp decline (Fig. 3b). This decline likely

181 reflects the lack of M8 cell growth, which limited the availability of susceptible hosts and

182 consequently reduced further CR4 replication.

183 ***Sal. ruber* M1 and its EM1 virus establish a stable long-term relationship, regardless of the**

184 **biological context**

185 After the initial 9-day period, all cultures were transferred to fresh medium to monitor their long-

186 term evolution with transfers every 9-13 days over a total of 150 days (13 transfers in total,

187 corresponding to ~240 generations). At selected time points, specific parameters were assessed,

188 including changes in host range and infectivity toward the native host (Fig. 1b).

189 *Sal. ruber* M1 was maintained at high density (~10^9 genomes/ml) across all time points, regardless

190 of the presence of EM1 or other strains and viruses (Extended Data Fig. 3). The EM1 virus

191 maintained relatively stable concentrations extracellularly across all three experiments (Extended

192 Data Fig. 4; Supplementary Fig. 4). Although the EM1 titer declined by about one order of

193 magnitude between days 52 and 84, its level remained stable until the end of the study. These

194 results indicate that *Sal. ruber* M1 and its EM1 virus established a stable coexistence that persisted

195 throughout the study, regardless of the surrounding biological context. Long-term virus–host

196 coexistence has been widely reported in other systems driven by mechanisms such as arms-race

197 dynamics, spatial heterogeneity, or phase variation in gene expression (Cortés-Martín et al., 2025;

198 Kortright et al., 2022; Lourenço et al., 2020). However, a previous work from our group

199 demonstrated that EM1 can be maintained in *Sal. ruber* M1 in a pseudolysogenic state (Sanchez-

Martinez et al., 2025), which appears to underlie the stable coexistence observed here. Its establishment and maintenance do not seem to be affected by the presence of additional strains and viruses.

Additional insights emerged from the dynamics of the other community members. In experiment 5, CC1 remained at high concentrations, as well as one of their hosts, M31, which was also detected until the end of the experiment (Extended Data Fig. 3 and 4). In the case of the CR4 virus, although it was not detected extracellularly after 9 days, it persisted in the total culture until 84 days, coinciding with the extinction of its host M8 (Supplementary Fig. 4). The P18 strain, the only one not infected by any virus, disappeared in experiment 5 from all 3 replicates and, although still detectable, was outcompeted by other strains in experiments 2 and 4. This indicates that viruses were not the only force structuring the community, and that competitive interactions among bacterial strains also played a crucial role.

The biological context reduces the infectivity of the EM1 virus and alters its host range

On day 9, the infectivity of EM1, calculated as the ratio between PFU/ml (tested with the native M1 strain as the host) and extracellular EM1 genomes/ml, was maintained above 0.1% in the three experiments and remained stable at these levels for 52 days (Fig. 4a). However, by day 84, significant differences (p-value < 0.0001) emerged between experiment 3 and experiments 4 and 5. In experiment 3 infectivity remained stable but, in experiment 4, where the *Sal. ruber* M1-EM1 pair was accompanied by other strains, infectivity dropped by almost one order of magnitude. In experiment 5, the most biologically complex, this reduction in infectivity was more drastic, with a drop of nearly 4 orders of magnitude compared to experiment 3. By day 130, infectivity had decreased further in all experiments, down to 5 orders of magnitude below the initial level. These findings suggested that the EM1 virus underwent genomic changes, as further explored below, which reduced its ability to infect the native strain in all the experiments and highlighted that the biological context significantly affected the interactions between EM1 and its host.

Host range changes were also detected in the EM1 virus in experiment 5, whereas no such changes were observed in experiments 3 and 4. The evolved EM1 in experiment 5, which initially only infected the M1 strain, expanded its host range and infected the native M8 strain consistently across all three biological replicates at 84 hours (Fig. 4b). Several studies have reported host range

changes in simplified experimental evolution contexts (Bono et al., 2015; Holtzman et al., 2020;
Sant et al., 2021; Subramanian et al., 2022), but only a few have explored this phenomenon under
more complex conditions. In multispecies settings, viruses have been shown to broaden their host
range by transiently using alternative hosts as intermediates to access new hosts (De Sordi et al.,
2017), which may partly explain why this host range shift occurred in experiment 5. Moreover, it
has been observed that the presence of multiple viruses affect the host range of a given virus by
driving evolutionary changes of the host that affect its susceptibility (Avrani et al., 2011;
Chevallereau et al., 2022). These findings suggest that the observed host range expansion in EM1
was a result of selective pressures arising from increased ecological complexity. Coexistence with
other strains and viruses may have favored viral variants capable of infecting alternative hosts (i.e.,
either by exploiting new receptor variants or by overcoming host-specific resistance mechanisms
(see below)). In the case of the other viruses, none of them showed any host range changes.

The unequal decline in the EM1 virus infectivity across experiments, together with the shift in host
range observed in experiment 5, suggested that EM1 underwent different genomic changes under
the different conditions, influencing the virus–host evolutionary trajectory according to the
surrounding biotic environment. To analyze whether genomic changes could be responsible for
these differences, extracellular viromes and cellular pellets from experiments 3, 4 and 5 were
sequenced on days 9, 84 and 150. The original viral genomes were also sequenced as references.

Viral genomic diversification is enhanced by the presence of other viruses

EM1 genome remained largely stable during the early stages of the experiment, with only a single
mutation in a gene coding for a hypothetical protein detected across all treatments by day 9
(Supplementary Table 2). By day 84, mutation accumulation remained low in experiments 3 and
4 (15 and 14 mutations, respectively), mostly in non-coding regions (Extended Data Fig. 5;
Supplementary Table 2). Experiment 5 showed a markedly different pattern, with 772 mutations
distributed across the genome. A large proportion of these occurred in structural genes, particularly
those related to tail components. Two minor tail protein genes accumulated over 190 mutations,
and a long-tail fiber protein gene showed 69 changes, suggesting strong selective pressure on host
recognition machinery. This genomic diversification coincided with a sharp drop in EM1

infectivity against the native M1 host, pointing to a possible link between mutation accumulation in tail-related regions and host range evolution.

As the experiment progressed to day 150, the accumulation of mutations continued to diverge between treatments. Experiment 3 reached a total of 34 mutations, including 17 in a gene coding for a DNA methyltransferase and several nonsynonymous changes in genes for a DNA-binding protein, a minor tail protein, and a long-tail fiber protein, some of which may be linked to the observed decline in infectivity (Fig. 5). Experiment 4 accumulated 267 mutations, with a high concentration in genes related to DNA metabolism, such as a DNA primase and a single-strand DNA-binding protein. This accumulation of mutations in DNA metabolism-related genes could suggest a potential disruption in viral replication, contributing to reduced infectivity (Boddin et al., 2022; Jiang et al., 2007, 2009). In experiment 5, mutation accumulation intensified, reaching 1,331 mutations distributed across the genome. Structural and replication-related genes showed particularly high mutational loads, including 198 mutations in minor tail protein genes and over 190 combined in genes for a DNA-binding protein and a DNA primase. This translated into a mutation rate of 2.5×10^{-4} mutations per site per day in experiment 5, markedly higher than those estimated for experiments 3 and 4 (6.4×10^{-6} and 5×10^{-5}, respectively) (Supplementary Table 2).

These differences were reflected in the average nucleotide identity of sequence alignment reads (ANIr) along the EM1 genome (the mean nucleotide-level similarity between mapped reads and the reference genome). Experiments 3 and 4 showed higher ANIr values (99.70% and 99.75%, respectively) indicating a more homogeneous population, compared to experiment 5 (98.13%) on day 150 (Supplementary Table 2). The lower ANIr values observed in the presence of other viruses indicated an accelerated diversification of the viral population, with the emergence of multiple distinct genotypes or "genomovars" (defined as variants with ANI <99.5%) (Aldeguer-Riquelme et al., 2024).

The increase in genetic variability observed under complex community conditions reveals a previously overlooked role of viral coexistence in promoting diversification. Our results demonstrate that the mere presence of other viruses can substantially reshape the genetic structure of a viral population, both in terms of mutation rate and the distribution of these mutations across the genome (Supplementary Table 2; Fig. 5). This diversification was accompanied by a drastic

loss of infectivity and a shift in host range, likely driven by mutations in tail protein genes. While
previous studies have shown that viral mutation rates can vary depending on the host genotype
(Duffy, 2018; Longdon et al., 2014; Sant et al., 2021), and that exposure to multiple host genotypes
can promote tail gene diversification (Hernandez et al., 2024), our findings suggest a previously
unreported mechanism. Here, it is the presence of other viruses, more than the mere presence of
other hosts, that significantly increased the viral mutation rate. This uncovers a novel form of
virus–virus interaction, where coexisting viruses indirectly drive each other's evolutionary
trajectories. Such interactions, largely overlooked to date, could play a central role in shaping viral
evolution in natural communities characterized by high viral diversity.

Intracellular viral diversity exceeds the subset released extracellularly

Analysis of intracellular populations revealed that only part of the viral diversity generated inside
host cells was released extracellularly, indicating that the extracellular fraction corresponds to a
selected subset of intracellular genotypes (i.e., viruses maintained episomally and/or genomes
undergoing replication for virion packaging). Three genes (marked with an asterisk in Fig. 5) were
characterized by amplicon sequencing: one encoding a hypothetical protein with minimal
mutations across all experiments (used as a control), and two genes highly mutated in experiment
5, encoding a minor tail protein and a DNA-binding protein, respectively. The mutation patterns
in these genes were generally similar to those observed in the extracellular viral population,
especially in experiments 3 and 4, with few differences (Extended Data Fig. 6). Experiment 5
displayed a markedly higher number of mutations in the intracellular fraction in both the minor
tail protein and DNA-binding protein genes (63 and 30 mutations on average across replicates)
compared to experiment 3 (0 and 2 mutations) and experiment 4 (no mutations).

When comparing intracellular and extracellular fractions some differences emerged, particularly
in experiment 5. Both the minor tail protein and DNA-binding protein regions showed a high
number of mutations at nearly 100% frequency in the extracellular fraction, while the intracellular
fraction exhibited greater heterogeneity in mutation frequencies. These results support our earlier
findings that the presence of other viruses acts as a selective force, increasing the mutation rate of
EM1 and generating a larger diversity of viral genotypes. However, based on the intracellular data,

only a subset of these genotypes appears to be selected and successfully replicated for virion production.

Intraspecific interactions, rather than viral pressure, drive host genome evolution

We also analyzed genetic mutations in the host *Sal. ruber* M1. As experiments 2, 4, and 5 included additional *Sal. ruber* strains sharing large genome regions, we focused on nine genomic regions, ranging from 1.8 to 23.5 kb, specific to *Sal. ruber* M1. This strategy allowed us to unambiguously track mutations in M1 despite the presence of other closely related strains in the mixed cultures. In total, 70.1 kb were analyzed, (Supplementary Fig. 5). Mutation counts at 150 hours revealed striking differences: 9 mutations in experiment 1, 224 in experiment 2, 8 in experiment 3, 266 in experiment 4, and 14 in experiment 5, with most occurring in non-coding regions (Extended Data Fig. 7; Supplementary Table 3). These findings corresponded to mutation rates of 7,6 x 10^{-7}, 2,1 x 10^{-5}, 6,7 x 10^{-7}, 5 x 10^{-5}, 5,6 x 10^{-6} mutations per site per day, respectively (Supplementary Table 3). Our results demonstrate that *Sal. ruber* M1 experienced significantly elevated mutation rates in specific regions in those experiments where it evolved alongside multiple strains—experiments 2 and 4 (Extended Data Fig. 3).

In contrast, experiment 5, which also included other viruses, resulted in markedly lower host mutation rates. This suggests that the presence of viruses in a complex community context may constrain bacterial evolution, rather than accelerate it. This finding is striking and counterintuitive, as viruses are typically considered potent drivers of evolutionary change, and other studies showed that bacteria coevolving with multiple viruses exhibited higher mutation rates (Betts et al., 2018; Wielgoss et al., 2016). In our system, however, viral pressure may have limited evolutionary change in the host, possibly by imposing bottlenecks through lysis, constraining population dynamics, or reducing the strength of competition among bacterial strains.

The observed increase in mutation rates under competitive conditions suggests that bacterial intraspecific competition is a strong driver of genomic evolution in the host. While EM1 viral mutations primarily emerged within infected populations, *Sal. ruber* M1 genetic variability was predominantly shaped by interactions with coexisting bacterial strains. This is consistent with previous work from our laboratory demonstrating weak but significant competitive interactions between *Sal. ruber* strains in co-culture (Peña et al., 2010). Furthermore, in a recent mesocosm

experiment, we showed that strain M8 was displaced by other genotypes after being introduced at high abundance into a pond, suggesting competitive exclusion through intraspecific competition (Ramos-Barbero et al., 2024).

Several mechanisms could account for the elevated mutation rates observed in experiments 2 and 4. These include stress-induced mutagenesis, where competition triggers cellular stress responses that increase mutation frequency (Svet et al., 2023; Trejo-Hernández et al., 2014), and the selection of hypermutator phenotypes (bacterial variants with defective DNA repair systems) previously documented in *Escherichia coli* during gut colonization (Giraud et al., 2001) and *Vibrio cholerae* under antibiotic pressure (Baharoglu & Mazel, 2011). In our system, the sharp increase in mutations specifically in the absence of viral pressure suggests that competition with other bacterial strains may trigger stress responses or transient mutagenic states, rather than the stable fixation of hypermutator lineages. This is further supported by the fact that mutation rates remained low in experiment 5, despite the presence of both other strains and viruses, indicating that viral pressure may counteract or suppress the effects of competition-induced mutagenesis.

These findings support the view that microbial competition plays a central role in shaping evolutionary trajectories. Together, they underscore a critical consideration for evolutionary studies: microbial mutation rates and adaptation cannot be fully understood without accounting for the complex network of interspecies interactions that govern microbial life in natural environments.

CONCLUSIONS

The usefulness of studying virus–host interactions within microcosms that simulate natural conditions in a stepwise manner has become increasingly evident in recent years (Blazanin & Turner, 2021; Castledine & Buckling, 2024; Chevallereau et al., 2022; Koskella et al., 2022). Understanding the ecological and evolutionary dynamics of viruses and prokaryotes in their natural environments not only enriches our knowledge of the natural world but also enables us to manipulate these communities more predictably. In this study, we demonstrate that a semi-

complex biological context (encompassing *Sal. ruber* strains and their viruses) substantially
influence the ecology and evolution of the *Sal. ruber* M1-EM1 pair in controlled microcosms.

This work shows how in the short-term, the presence of additional *Sal. ruber* strains delays the
lysis of *Sal. ruber* M1 and reduces EM1 virion production. These effects were consistent across
different quantification methods, indicating that community composition has a measurable impact
on virus replication dynamics. In the long-term, *Sal. ruber* M1 and the EM1 virus established a
stable coexistence throughout the entire experiment regardless of community complexity. The
infectivity of EM1 against its original host declined faster in the presence of other viruses,
revealing that viral genomic diversification is enhanced by virus–virus interactions. This reduction
in infectivity was linked to genomic changes in EM1, leading to an expanded host range and a
sharply increased mutation rate in the presence of other viruses. Meanwhile, the bacterial host, *Sal.*
ruber M1, showed elevated mutation rates evolving alongside other strains, indicating that
intraspecific bacterial interactions—not viral pressure—drive host genome evolution.

Overall, our results highlight the critical importance of incorporating biological complexity in
studies of virus–host interactions. These findings emphasize the need to consider the multifaceted
nature of microbial communities to more accurately predict virus–host dynamics and their
ecological and evolutionary consequences in natural environments.

METHODS

Bacterial strains, viruses and experimental design

Sal. ruber strains M1, M8, M31 and P18 were aerobically grown with gentle shaking (60 rpm) at
37ºC in 25 % SW (sea water, a salt solution containing the salts present in seawater at a total
concentration of 25% weight/volume) with 0.2 % yeast extract (Rodriguez-Valera et al., 1985).
Sal ruber viruses used in this study, named M1EM-1, M31CC-1, M8CR-4 and M31CR41-2
(referred to here as EM1, CC1, CR4 and CR41-2, respectively, for convenience), were isolated
during a previous study (Villamor et al., 2018).

Five experiments were conducted in parallel (Fig. 1). In all the experiments, 25 ml of culture media
were inoculated with 7.5×10^8 cells of *Sal. ruber* M1. In experiments 2, 4, and 5, the cultures also
included 2.5×10^7 cells of each of the M8, M31, and P18 strains. In experiments 3, 4 and 5, 7.5×10^7
virus-like particles (VLP) of the strictly virulent EM1 virus (multiplicity of infection = 0.01),
which only infects the M1 strain, were added. Experiment 5 also contained 2.5×10^6 VLP of each
one of the CC1, CR4 and CR41-2 viruses that infect M8 and M31 strains (Supplementary Table
1). After inoculation, experiments were incubated for 9 days as described above, taking aliquots
once a day for culture monitoring by optical density (OD) at 600 nm and additional approaches
(see below). This was defined as the short-term experiment.

On day 9 (213 hours after inoculation), 250 µl of each experiment were transferred to 25 ml of
fresh medium and incubated as described above. After 13 days, new aliquots were collected to
quantify the bacterial and viral diversity of the cultures by microfluidics-based qPCR (see below)
and transferred and incubated in the same way. This was repeated every approximately 9-13 days
for 150 days. Days 9 to 150 were considered as the long-term experiment.

In addition, on days 9, 52, 84, 130 and 150, aliquots were taken for plaque forming units (PFU)
quantification, host range analyses of the viruses and DNA extraction and sequencing. All
experiments were conducted in triplicate.

Flow cytometry

For each day of the short-term experiment, cells/ml and VLP/ml were quantified by flow cytometry
as described in Brussaard, 2004. Briefly, 500 µl of each sample were fixed with glutaraldehyde
(0.5% final concentration) at 4ºC for 30 min, flash-frozen in liquid nitrogen, and stored at -80°C
until use. Upon thawing, samples were diluted in Tris-EDTA buffer (pH 8), stained with
SYBR™Gold (0.5X final concentration), incubated for 10 min in the dark at 80°C, and cooled for
5 min at room temperature prior to analysis.

The cytometer settings were as follows: the threshold was set in blue fluorescence (300 units),
FITC voltage = 500, SSC voltage = 300, forward scatter voltage = 500, and the flow rate was
established as low (15 µl/min). Background noise was checked on blanks, composed by TE buffer
stained with SYBR Gold. Samples were recorded with an event rate of 100-1000 events per second.

Cells and VLP counts were obtained by correcting the background measured in blanks. Cells and

VLP abundances were respectively expressed as cells/ml or VLP/ml.

Plaque assay

The number of free infective EM1 viruses at each sampling point of the short- and long-term

experiments was determined by plaque assay in experiments 3, 4 and 5. Aliquots of 150 µl were

centrifuged at 17,000 x g for 10 min and supernatants were serially diluted in sterile SW 25%. 100

µl of dilutions 10^{-4}, 10^{-6} and 10^{-8} were mixed with 500 µl of *Sal. ruber* M1 in exponential phase.

Four ml of soft agar (25% SW with 0.2% yeast extract and 0.7% agar) were added and this was

poured on plates with solid media. After the agar had solidified, the plates were incubated at 37ºC

for 10 days until plaques were visible. The number of free infective viruses (PFU/ml) was counted

and represented. On days 9, 52, 84, 130 and 150, plaque assays were also carried out to obtain

individual plaques of all viruses for experiment 5, using as hosts the native strains M8, M31 and

P18.

High throughput microfluidics-based qPCR

Individual targets (i.e. the 4 strains and the 4 viruses) were quantified in each point of the short-

term experiment, in both the total culture and the supernatant to quantify free viruses, using the

Biomark HD high throughput microfluidics-based qPCR system (Standard Biotools, South San

Francisco, USA). Primers for each target gene were designed using using Geneious software 6.1.8

(Biomatters, Auckland, New Zealand) (Supplementary Table 4). From each of the experiments,

100 µl of the culture were taken for total culture analysis and 120 µl for supernatant quantification

of free viruses. In the latter, a centrifugation step consisting of 10 min at 7,197 x g and 100 µl

supernatant transfer to a clean eppendorf tube was performed. Both were fixed with 0.5% (final

concentration) of formaldehyde for 1 hour at 4°C, diluted 10-fold in phosphate buffered saline

(PBS) 1X, and stored at 4°C.

DNA targets were then pre-amplified using the PreAmp Master Mix (Standard Biotools, South

San Francisco, USA) following manufacturer's instructions. In brief, a primer pool was prepared

by mixing the forward and reverse primers for all targets (Supplementary Table 4) and diluting the

mixture with DNA suspension buffer (10 mM Tris/HCL and 0.1 mM EDTA pH 8) to a final

concentration of 0.5 µM of each primer pair. Each sample was then pre-amplified in a total volume

of 5 uL containing 1 µL PreAmp Master Mix (Standard Biotools, South San Francisco, USA),

0.5 µL of primer pool and 3.5 µL of sample. Preamplification reactions were performed on a

Simpliamp Thermal Cycler (Applied Biosystems, Waltham, USA) using the following sample

conditions: an initial activation cycle at 95°C for 10 min followed by 14 two-step cycles

(denaturation at 95°C for 15 seconds and annealing/extension at 60°C for 4 min). The preamplified

products were subjected to Exonuclease I cleanup (New England Biolabs, Massachusetts, USA)

to remove any unincorporated primers (4 U/µL final concentration at 37°C for 30 min followed by

inactivation at 80ºC for 15 min). A 1:5 dilution using DNA suspension buffer was then performed

on the Exonuclease I treated samples and stored at -20°C.

Finally, pre-amplified DNA targets were quantified by microfluidics-based qPCR on 192x24

Dynamic Array^{TM} IFCs (Fluidigm, South San Francisco, USA) according to manufacturer's

instructions. In brief, 10X assay primer mixtures for each DNA target and sample pre-mixes for

each sample were prepared in triplicate. For the generation of the 10X assay primer mixtures, the

forward and reverse primers for a single DNA target were pooled together at a concentration of 50

uM each. 0.15 uL of this pool were then mixed with 1.35 µl of DNA suspension buffer (Tris/HCL

and 0.1 mM EDTA pH 8) and 1.5 µl of 2X Assay Loading Reagent (Standard Biotools, South San

Francisco, USA). Sample pre-mixes were made by combining 1.5 µl SsoFast^{TM} Evagreen®

Supermix 2X (BioRad, California, USA) with 0.15 µl 20X GE Sample Loading Reagent (Standard

Biotools, South San Francisco, USA) and 1.35 µl of pre-amplified DNA. The 192x24 Dynamic

Array^{TM} IFCs were then primed on a Juno controller (Standard Biotools, South San Francisco,

USA) and 3 uL of each 10X assay primer mix and sample pre-mix were transferred to the

appropriate inlets of the IFC and loaded using again the Juno controller. After loading, the IFC

was transferred to a Biomark HD instrument (Standard Biotools, South San Francisco, USA) and

qPCR reactions were performed using the following cycling conditions: 95°C for 1 min, followed

by 30 two-step cycles (95°C for 15 seconds and 60°C for 20 seconds) and a final melt curve

analysis. Data were analyzed with the Real-Time PCR Analysis Software (Standard Biotools,

South San Francisco, USA) using manually defined thresholds. All samples were run in triplicate

(including the standards and negative controls). All samples, standards, and negative controls were

run in triplicate, with negative controls included both in the pre-amplification and in the

microfluidics-based qPCR. A total of four 192x24 Dynamic Array™ IFCs were used, each with

its own standard curve consisting of five concentration points for every target, and each chip

included positive controls for the eight targets individually.

Host range through spot test and qPCR

The host range of each virus at time 0 was determined through a spot test. Briefly, 4 ml of molten

0.7% top agar of 25% SW with 0.2% yeast extract were mixed with 500 µl of bacterial cultures at

exponential phase and plated on solid medium. Once solidified, 3 µl of each virus, titered at 10^{10}

PFUs/ml and diluted 10^2, 10^4 and 10^6 times, were added to the corresponding lawn. The spotted

plates were left to dry and incubated at 37ºC for 10 days. The detection of a clearance zone was

interpreted as evidence of viral lysis while the areas exhibiting growth were attributed to resistance.

This was done by exposing the 4 native strains (*Sal. ruber* strains M1, M8, M31 and P18) to the 4

viruses (EM1, CC1, CR41-2 and CR4 viruses).

The viral fractions from experiments 3, 4 and 5, at 9, 52, 84, 130 and 150 days, were obtained by

centrifuging each replicate at 7,197 x g for 20 min and filtering the supernatant through a 0.22 µm

filter (Millipore, Burlington, USA). For experiments 3 and 4, the host range of the EM1 virus over

time was measured through a spot test against the 4 native hosts as described above.

The determination of the host range of the 4 viruses over time in experiment 5 was performed by

qPCR as follows. All the plaques obtained for each native host (see plaque assay above) for each

replicate on days 9, 52, 84, 130 and 150, were picked from the soft agar and resuspended in 1 ml

of ultrapure water. The presence of each virus in these mixtures of plaques were checked by qPCR

with specific primers (Supplementary Table 4) using SYBR Green. The experiment was carried

out with the standard run in a StepOnePlus PCR System (Life Technologies, Carlsbad, USA) in a

10 µl reaction mixture with Power SYBR Green PCR Master Mix (Applied Biosystems, Waltham,

USA). The reaction contained: 5 µl of 2X Master Mix, 0.2 µl of each 10 µM primer, 1 µl of sample

and ultrapure water to complete volume. Conditions are detailed in Supplementary Table 5. The

results were analyzed with the Applied Biosystems StepOne™ Instrument program. All samples

were run in triplicate (including the standards and negative controls).

Adsorption assay

An adsorption assay was performed in parallel with the CC1 and CR41-2 viruses and *Sal. ruber*
M31 wild type. Cultures in exponential phase were diluted with fresh medium to an OD of 0.3 and
$2x10^8$ cells were mixed with $2x10^6$ PFU (MOI=0.01) of the virus. This was considered as the initial
time. Mixtures were then incubated at 37 ºC and 60 rpm. Aliquots of 150 µl were taken along 5 h
(at 1, 2, 3, 4 and 5 h), centrifuged at 17,000 x g for 8 min, and 120 µl of the cells-free supernatant
were stored in ice. The numbers of PFU/ml were measured by plaque assay as described above.
All the experiments were conducted in triplicate.

Lysate experiment

An infection of *Sal. ruber* M1 with the EM1 virus at a MOI of 0.1 was carried out under the same
conditions as described above. After 5 days, the infection was centrifuged at 17,000 x g for 10
minutes, and the supernatant was filtered through a 0.022 µm filter (Millipore, Burlington, USA)
to remove any remaining viruses. One ml of this lysate was added to 25 ml of exponential-phase
cultures of M8, M31, and P18 strains. Controls received 1 ml of fresh medium instead. Optical
density was measured at 600 nm. Both the lysate-treated cultures and controls were conducted in
triplicate.

DNA extractions and sequencing

The three biological replicates of each experiment were pooled prior to DNA extraction, as
previous experiments in our laboratory have shown high reproducibility in the mutational
trajectories of replicate populations. Bacterial pellets from all the experiments at 150 days were
obtained by centrifugation at 17,000 x g for 10 min. The cells were washed with sterile medium,
the supernatant removed, and the cell pellets stored at -80ºC until extraction. DNA was extracted
with the DNeasy Blood & Tissue Kit (Qiagen, Hilden, Germany) following the manufacturer's
protocol, and nucleic acids eluted in 70 µL of ultrapure water.

Viruses from the same points were obtained by centrifuging at 7.197 x g for 20 min and filtering
the supernatant through a filter of 0.22 µm. Dissolved DNA was removed with DNAse I from
bovine pancreas (Sigma-Aldrich, St. Louis, USA) at 37ºC for 30 min, with an inactivation at 75ºC
for 10 min. DNA was extracted with the QIAamp MinElute Virus Kit (Qiagen, Hilden, Germany)

following the manufacturer's protocol, and nucleic acids eluted in 35 µL of ultrapure water. DNA from the viruses used for infecting at the initial time was also extracted as a reference.

All extracted cellular and viral DNAs were quantified using Qubit 2.0. Flourometer (Life Technologies, Carlsbad, USA), and sequenced on an Illumina Novaseq6000 (2 x 150 bp) at Macrogen (Macrogen, Seoul, South Korea).

Metabarcoding

Three genes from the EM1 virus were selected, and specific primers targeting ~375 bp regions were designed (Supplementary Table 6). Cell pellets were washed three times by centrifugation, and DNA was extracted and quantified as described above. Specific regions were amplified via PCR in 50 µL reactions containing: 1.5 mM MgCl$_2$, 10X buffer, 10 mM dNTPs, 5 U/µL Taq polymerase, 100 µM primers, 5 µg of the template DNA and water up to the volume. PCR conditions are detailed in Supplementary Table 7. Five ml of the amplified products were electrophoresed, stained with ethidium bromide (100 µg/mL), and visualized under UV light. The remaining product was sequenced on an Illumina Miseq (2 x 150 bp) at Fundació per al Foment de la Investigació Sanitària i Biomèdica de la Comunitat Valenciana (Fisabio, Valencia, Spain).

Mutations were identified comparing the obtained reads with the reference genomes of the viruses at initial time using Geneious software 6.1.8 (Biomatters, Auckland, New Zealand) with a minimum variant frequency of 0.01% and a coverage >50% to the mean coverage. Mutations were represented using the package 'ggplot2' v3.4.0 in R v4.2.2.

Bioinformatic analysis

Primers and adapters were removed from sequences, and reads were filtered based on quality scores using Trimmomatic v0.36.0 (Bolger et al., 2014). Trimmed reads of initial viruses were assembled using SPAdes v3.13.1 (Bankevich et al., 2012) with the trusted option using their genomes of NCBI as a reference (GenBank accession numbers MF580955, MF580958, MF580960 and MF580960).

Mutations generated in the viruses over time were identified by comparing reads from the extracellular viromes on days 9, 84 and 150 with the reference genomes of the viruses at initial

time using Geneious software 6.1.8 (Biomatters, Auckland, New Zealand) with a minimum variant

frequency of 0.01% and a coverage >50% to the mean coverage. Mutations were represented using

the package 'ggplot2' v3.4.0 in R v4.2.2. For the analysis of M1 sequences, the chromosomes of

strains M1, M8, M31 and P18 were aligned with Mauve (Darling et al., 2004) and 9 M1-specific

regions were selected where the mutations were called in the same way as for the virus.

A BLASTn was performed with the reads of each virome, using the genome of the EM1 virus as

the reference. Then, the BLASTn output was filtered by best_hit option and read coverage >70%.

The ANIr (average nucleotide identity of sequence alignment reads) along the genome of EM1

was calculated with a home-made script in windows of 50 bp and plotted. Recruitment plots were

carried out by enveomics tool (Rodriguez-R & Konstantinidis, 2016) using enve.recplot 2 (R

stadistic pluging).

REFERENCES

Abedon, S. T. (2023). Bacteriophage Adsorption: Likelihood of Virion Encounter with Bacteria and Other Factors Affecting Rates. Antibiotics, 12(4), Article 4. https://doi.org/10.3390/antibiotics12040723

Aldeguer-Riquelme, B., Conrad, R. E., Antón, J., Rossello-Mora, R., & Konstantinidis, K. T. (2024). A natural ANI gap that can define intra-species units of bacteriophages and other viruses. mBio, 15(8), e01536-24. https://doi.org/10.1128/mbio.01536-24

Alseth, E. O., Custodio, R., Sundius, S. A., Kuske, R. A., Brown, S. P., & Westra, E. R. (2024). The impact of phage and phage resistance on microbial community dynamics. PLOS Biology, 22(4), e3002346. https://doi.org/10.1371/journal.pbio.3002346

Alseth, E. O., Pursey, E., Luján, A. M., McLeod, I., Rollie, C., & Westra, E. R. (2019). Bacterial biodiversity drives the evolution of CRISPR-based phage resistance. Nature, 574(7779), 549-552. https://doi.org/10.1038/s41586-019-1662-9

Antón, J., Oren, A., Benlloch, S., Rodríguez-Valera, F., Amann, R., & Rosselló-Mora, R. (2002). *Salinibacter ruber* gen. Nov., sp. Nov., a novel, extremely halophilic member of the Bacteria from

saltern crystallizer ponds. International Journal of Systematic and Evolutionary Microbiology,

52(2), 485-491. https://doi.org/10.1099/00207713-52-2-485

Avrani, S., Wurtzel, O., Sharon, I., Sorek, R., & Lindell, D. (2011). Genomic island variability

facilitates *Prochlorococcus*–virus coexistence. Nature, 474(7353), 604-608.

https://doi.org/10.1038/nature10172

Baharoglu, Z., & Mazel, D. (2011). *Vibrio cholerae* Triggers SOS and Mutagenesis in Response

to a Wide Range of Antibiotics: A Route towards Multiresistance. Antimicrobial Agents and

Chemotherapy, 55(5), 2438-2441. https://doi.org/10.1128/aac.01549-10

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., Lesin, V. M.,

Nikolenko, S. I., Pham, S., Prjibelski, A. D., Pyshkin, A. V., Sirotkin, A. V., Vyahhi, N., Tesler,

G., Alekseyev, M. A., & Pevzner, P. A. (2012). SPAdes: A New Genome Assembly Algorithm

and Its Applications to Single-Cell Sequencing. Journal of Computational Biology, 19(5), 455-

477. https://doi.org/10.1089/cmb.2012.0021

Benton, T. G., Solan, M., Travis, J. M. J., & Sait, S. M. (2007). Microcosm experiments can inform

global ecological problems. Trends in Ecology & Evolution, 22(10), 516-521.

https://doi.org/10.1016/j.tree.2007.08.003

Betts, A., Gray, C., Zelek, M., MacLean, R. C., & King, K. C. (2018). High parasite diversity

accelerates host adaptation and diversification. Science, 360(6391), 907-911.

https://doi.org/10.1126/science.aam9974

Blazanin, M., & Turner, P. E. (2021). Community context matters for bacteria-phage ecology and

evolution. The ISME Journal, 15(11), 3119-3128. https://doi.org/10.1038/s41396-021-01012-x

Boddin, J., Ip, W.-H., Wilkens, B., von Stromberg, K., Ching, W., Koyuncu, E., Bertzbach, L. D.,

& Dobner, T. (2022). A Single Amino Acid Switch in the Adenoviral DNA Binding Protein

Abrogates Replication Center Formation and Productive Viral Infection. mBio, 13(2), e00144-22.

https://doi.org/10.1128/mbio.00144-22

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina

sequence data. Bioinformatics, 30(15), 2114-2120. https://doi.org/10.1093/bioinformatics/btu170

Bono, L. M., Gensel, C. L., Pfennig, D. W., & Burch, C. L. (2015). Evolutionary rescue and the

coexistence of generalist and specialist competitors: An experimental test. Proceedings of the

Royal Society B: Biological Sciences, 282(1821), 20151932.

https://doi.org/10.1098/rspb.2015.1932

Brussaard, C. P. D. (2004). Optimization of Procedures for Counting Viruses by Flow Cytometry.

Applied and Environmental Microbiology, 70(3), 1506-1513.

https://doi.org/10.1128/AEM.70.3.1506-1513.2004

Bürkle, M., Korf, I. H. E., Lippegaus, A., Krautwurst, S., Rohde, C., Weissfuss, C., Nouailles, G.,

Tene, X. M., Gaborieau, B., Ghigo, J.-M., Ricard, J.-D., Hocke, A. C., Papenfort, K., Debarbieux,

L., Witzenrath, M., Wienhold, S.-M., & Krishnamoorthy, G. (2025). Phage–phage competition

and biofilms affect interactions between two virulent bacteriophages and *Pseudomonas*

aeruginosa. The ISME Journal, 19(1), wraf065. https://doi.org/10.1093/ismejo/wraf065

Burns, D. G., Janssen, P. H., Itoh, T., Kamekura, M., Li, Z., Jensen, G., Rodríguez-Valera, F.,

Bolhuis, H., & Dyall-Smith, M. L. (2007). *Haloquadratum walsbyi* gen. Nov., sp. Nov., the square

haloarchaeon of Walsby, isolated from saltern crystallizers in Australia and Spain. International

Journal of Systematic and Evolutionary Microbiology, 57(2), 387-392.

https://doi.org/10.1099/ijs.0.64690-0

Calisi, R. M., & Bentley, G. E. (2009). Lab and field experiments: Are they the same animal?

Hormones and Behavior, 56(1), 1-10. https://doi.org/10.1016/j.yhbeh.2009.02.010

Castledine, M., & Buckling, A. (2024). Critically evaluating the relative importance of phage in

shaping microbial community composition. Trends in Microbiology, 32(10), 957-969.

https://doi.org/10.1016/j.tim.2024.02.014

Chen, J., Zhang, Y., Kuzyakov, Y., Wang, D., & Olesen, J. E. (2023). Challenges in upscaling

laboratory studies to ecosystems in soil microbiology research. Global Change Biology, 29(3),

569-574. https://doi.org/10.1111/gcb.16537

Chevallereau, A., Pons, B. J., van Houte, S., & Westra, E. R. (2022). Interactions between bacterial
and phage communities in natural environments. Nature Reviews Microbiology, 20(1), 49-62.
https://doi.org/10.1038/s41579-021-00602-y

Cortés-Martín, A., Buttimer ,Colin, Maier ,Jessie L., Tobin ,Ciara A., Draper ,Lorraine A., Ross
,R. Paul, Kleiner ,Manuel, Hill ,Colin, & and Shkoporov, A. N. (2025). Adaptations in gut
Bacteroidales facilitate stable co-existence with their lytic bacteriophages. Gut Microbes, 17(1),
2507775. https://doi.org/10.1080/19490976.2025.2507775

Darling, A., Mau, B., Blattner, F., & Perna, N. (2004). Mauve: Multiple Alignment of Conserved
Genomic Sequence With Rearrangements. https://genome.cshlp.org/content/14/7/1394

De Sordi, L., Khanna, V., & Debarbieux, L. (2017). The Gut Microbiota Facilitates Drifts in the
Genetic Diversity and Infectivity of Bacterial Viruses. Cell Host & Microbe, 22(6), 801-808.e3.
https://doi.org/10.1016/j.chom.2017.10.010

Di Meglio, L., Santos, F., Gomariz, M., Almansa, C., López, C., Antón, J., & Nercessian, D.
(2016). Seasonal dynamics of extremely halophilic microbial communities in three Argentinian
salterns. FEMS Microbiology Ecology, 92(12), fiw184. https://doi.org/10.1093/femsec/fiw184

Duffy, S. (2018). Why are RNA virus mutation rates so damn high? PLOS Biology, 16(8),
e3000003. https://doi.org/10.1371/journal.pbio.3000003

Fierer, N. (2017). Embracing the unknown: Disentangling the complexities of the soil microbiome.
Nature Reviews Microbiology, 15(10), 579-590. https://doi.org/10.1038/nrmicro2017.87

Fuhrman, J. A., Cram, J. A., & Needham, D. M. (2015). Marine microbial community dynamics
and their ecological interpretation. Nature Reviews Microbiology, 13(3), 133-146.
https://doi.org/10.1038/nrmicro3417

Giraud, A., Matic, I., Tenaillon, O., Clara, A., Radman, M., Fons, M., & Taddei, F. (2001). Costs
and Benefits of High Mutation Rates: Adaptive Evolution of Bacteria in the Mouse Gut. Science,
291(5513), 2606-2608. https://doi.org/10.1126/science.1056421

Gomariz, M., Martínez-García, M., Santos, F., Rodriguez, F., Capella-Gutiérrez, S., Gabaldón, T.,

Rosselló-Móra, R., Meseguer, I., & Antón, J. (2015). From community approaches to single-cell

genomics: The discovery of ubiquitous hyperhalophilic Bacteroidetes generalists. The ISME

Journal, 9(1), Article 1. https://doi.org/10.1038/ismej.2014.95

Gómez, P., & Buckling, A. (2011). Bacteria-Phage Antagonistic Coevolution in Soil. Science,

332(6025), 106-109. https://doi.org/10.1126/science.1198767

Gómez, P., & Buckling, A. (2013). Coevolution with phages does not influence the evolution of

bacterial mutation rates in soil. The ISME Journal, 7(11), 2242-2244.

https://doi.org/10.1038/ismej.2013.105

Hernandez, C. A., Delesalle, V. A., Krukonis, G. P., DeCurzio, J. M., & Koskella, B. (2024).

Genomic and phenotypic signatures of bacteriophage coevolution with the phytopathogen

Pseudomonas syringae. Molecular Ecology, 33(10), e16850. https://doi.org/10.1111/mec.16850

Holtzman, T., Globus, R., Molshanski-Mor, S., Ben-Shem, A., Yosef, I., & Qimron, U. (2020). A

continuous evolution system for contracting the host range of bacteriophage T7. Scientific Reports,

10(1), 307. https://doi.org/10.1038/s41598-019-57221-0

Jessup, C. M., Kassen, R., Forde, S. E., Kerr, B., Buckling, A., Rainey, P. B., & Bohannan, B. J.

M. (2004). Big questions, small worlds: Microbial model systems in ecology. Trends in Ecology

& Evolution, 19(4), 189-197. https://doi.org/10.1016/j.tree.2004.01.008

Jiang, C., Hwang, Y. T., Randell, J. C. W., Coen, D. M., & Hwang, C. B. C. (2007). Mutations

That Decrease DNA Binding of the Processivity Factor of the Herpes Simplex Virus DNA

Polymerase Reduce Viral Yield, Alter the Kinetics of Viral DNA Replication, and Decrease the

Fidelity of DNA Replication. Journal of Virology, 81(7), 3495-3502.

https://doi.org/10.1128/jvi.02359-06

Jiang, C., Komazin-Meredith, G., Tian, W., Coen, D. M., & Hwang, C. B. C. (2009). Mutations

That Increase DNA Binding by the Processivity Factor of Herpes Simplex Virus Affect Virus

Production and DNA Replication Fidelity. Journal of Virology, 83(15), 7573-7580.

https://doi.org/10.1128/jvi.00193-09

Kortright, K. E., Chan, B. K., Evans, B. R., & Turner, P. E. (2022). Arms race and fluctuating

selection dynamics in *Pseudomonas aeruginosa* bacteria coevolving with phage OMKO1. Journal

of Evolutionary Biology, 35(11), 1475-1487. https://doi.org/10.1111/jeb.14095

Koskella, B., Hernandez, C. A., & Wheatley, R. M. (2022). Understanding the Impacts of

Bacteriophage Viruses: From Laboratory Evolution to Natural Ecosystems. Annual Review of

Virology, 9(Volume 9, 2022), 57-78. https://doi.org/10.1146/annurev-virology-091919-075914

Levin, S. A. (1998). Ecosystems and the Biosphere as Complex Adaptive Systems. Ecosystems,

1(5), 431-436. https://doi.org/10.1007/s100219900037

Longdon, B., Brockhurst, M. A., Russell, C. A., Welch, J. J., & Jiggins, F. M. (2014). The

Evolution and Genetics of Virus Host Shifts. PLOS Pathogens, 10(11), e1004395.

https://doi.org/10.1371/journal.ppat.1004395

Lourenço, M., Chaffringeon, L., Lamy-Besnier, Q., Pédron, T., Campagne, P., Eberl, C., Bérard,

M., Stecher, B., Debarbieux, L., & De Sordi, L. (2020). The Spatial Heterogeneity of the Gut

Limits Predation and Fosters Coexistence of Bacteria and Bacteriophages. Cell Host & Microbe,

28(3), 390-401.e5. https://doi.org/10.1016/j.chom.2020.06.002

Middelboe, M., & Brussaard, C. P. D. (2017). Marine Viruses: Key Players in Marine Ecosystems.

Viruses, 9(10), Article 10. https://doi.org/10.3390/v9100302

Middelboe, M., Hagström, A., Blackburn, N., Sinn, B., Fischer, U., Borch, N. H., Pinhassi, J.,

Simu, K., & Lorenz, M. G. (2001). Effects of Bacteriophages on the Population Dynamics of Four

Strains of Pelagic Marine Bacteria. Microbial Ecology, 42(3), 395-406.

https://doi.org/10.1007/s00248-001-0012-1

Mora-Ruiz, M. del R., Cifuentes, A., Font-Verdera, F., Pérez-Fernández, C., Farias, M. E.,

González, B., Orfila, A., & Rosselló-Móra, R. (2018). Biogeographical patterns of bacterial and

archaeal communities from distant hypersaline environments. Systematic and Applied

Microbiology, 41(2), 139-150. https://doi.org/10.1016/j.syapm.2017.10.006

Mumford, R., & Friman, V.-P. (2017). Bacterial competition and quorum-sensing signalling shape

the eco-evolutionary outcomes of model in vitro phage therapy. Evolutionary Applications, 10(2),

161-169. https://doi.org/10.1111/eva.12435

Oren, A. (2011). Ecology of Halophiles. En K. Horikoshi (Ed.), Extremophiles Handbook (pp.

343-361). Springer Japan. https://doi.org/10.1007/978-4-431-53898-1_16

Penchuk, Y., Savytska, M., Kobyliak, N., Ostapchenko, D., Kolodiy, I., Onysenko, S., Tsyryuk,

O., Korotkyi, O., Grygoriev, F., & Falalyeyeva, T. (2023). Antimicrobial activity of dietary

supplements based on bacterial lysate of *Lactobacillus rhamnosus* DV. Frontiers in Cellular and

Infection Microbiology, 13. https://doi.org/10.3389/fcimb.2023.1211952

Ramos-Barbero, M. D., Aldeguer-Riquelme, B., Viver, T., Villamor, J., Carrillo-Bautista, M.,

López-Pascual, C., Konstantinidis, K. T., Martínez-García, M., Santos, F., Rossello-Mora, R., &

Antón, J. (2024). Experimental evolution at ecological scales allows linking of viral genotypes to

specific host strains. The ISME Journal, wrae208. https://doi.org/10.1093/ismejo/wrae208

Roberts, A. E. L., Kragh, K. N., Bjarnsholt, T., & Diggle, S. P. (2015). The Limitations of In Vitro

Experimentation in Understanding Biofilms and Chronic Infection. Journal of Molecular Biology,

427(23), 3646-3661. https://doi.org/10.1016/j.jmb.2015.09.002

Rodriguez-R, L. M., & Konstantinidis, K. T. (2016). The enveomics collection: A toolbox for

specialized analyses of microbial genomes and metagenomes (No. e1900v1). PeerJ Inc.

https://doi.org/10.7287/peerj.preprints.1900v1

Rodriguez-Valera, F., Ventosa, A., Juez, G., & Imhoff, J. F. (1985). Variation of Environmental

Features and Microbial Populations with Salt Concentrations in a Multi-Pond Saltern. Microbial

Ecology, 11(2), 107-115. https://doi.org/10.1007/BF02010483

Rohwer, F., & Thurber, R. V. (2009). Viruses manipulate the marine environment. Nature,

459(7244), 207-212. https://doi.org/10.1038/nature08060

Sanchez-Martinez, R., Arani, A., Krupovic, M., Weitz, J. S., Santos, F., & Antón, J. (2025).

Episomal virus maintenance enables bacterial population recovery from infection and promotes

virus–bacterial coexistence. The ISME Journal, 19(1), wraf066.
https://doi.org/10.1093/ismejo/wraf066

Sant, D. G., Woods, L. C., Barr, J. J., & McDonald, M. J. (2021). Host diversity slows
bacteriophage adaptation by selecting generalists over specialists. Nature Ecology & Evolution,
5(3), Article 3. https://doi.org/10.1038/s41559-020-01364-1

Santos, F., Yarza, P., Parro, V., Meseguer, I., Rosselló-Móra, R., & Antón, J. (2012). Culture-
Independent Approaches for Studying Viruses from Hypersaline Environments. Applied and
Environmental Microbiology, 78(6), 1635-1643. https://doi.org/10.1128/AEM.07175-11

Shiklomanov, I. A. (1998). World freshwater resources. UNESCO International Hydrological
Programme, UNESCO-IHP.

Smakman, F., & Hall, A. R. (2022). Exposure to lysed bacteria can promote or inhibit growth of
neighboring live bacteria depending on local abiotic conditions. FEMS Microbiology Ecology,
98(2), fiac011. https://doi.org/10.1093/femsec/fiac011

Subramanian, S., Dover, J. A., Parent, K. N., & Doore, S. M. (2022). Host Range Expansion of
Shigella Phage Sf6 Evolves through Point Mutations in the Tailspike. Journal of Virology, 96(16),
e00929-22. https://doi.org/10.1128/jvi.00929-22

Svet, L., Parijs, I., Isphording, S., Lories, B., Marchal, K., & Steenackers, H. P. (2023).
Competitive interactions facilitate resistance development against antimicrobials. Applied and
Environmental Microbiology, 89(10), e01155-23. https://doi.org/10.1128/aem.01155-23

Trejo-Hernández, A., Andrade-Domínguez, A., Hernández, M., & Encarnación, S. (2014).
Interspecies competition triggers virulence and mutability in *Candida albicans–Pseudomonas*
aeruginosa mixed biofilms. The ISME Journal, 8(10), 1974-1988.
https://doi.org/10.1038/ismej.2014.53

Turner, P. E. (2005). Parasitism Between Co-Infecting Bacteriophages. En R. A. Desharnais (Ed.),
Advances in Ecological Research (Vol. 37, pp. 309-332). Academic Press.
https://doi.org/10.1016/S0065-2504(04)37010-8

Ventosa, A. (2006). Unusual micro-organisms from unusual habitats: Hypersaline environments.

En H. M. Lappin-Scott, N. A. Logan, & P. C. F. Oyston (Eds.), Prokaryotic Diversity: Mechanisms

and   Significance   (pp.   223-254).   Cambridge   University   Press.

https://doi.org/10.1017/CBO9780511754913.015

Ventosa, A., de la Haba, R. R., Sánchez-Porro, C., & Papke, R. T. (2015). Microbial diversity of

hypersaline environments: A metagenomic approach. Current Opinion in Microbiology, 25, 80-

87. https://doi.org/10.1016/j.mib.2015.05.002

Villamor, J., Ramos-Barbero, M. D., González-Torres, P., Gabaldón, T., Rosselló-Móra, R.,

Meseguer, I., Martínez-García, M., Santos, F., & Antón, J. (2018). Characterization of ecologically

diverse viruses infecting co-occurring strains of cosmopolitan hyperhalophilic Bacteroidetes. The

ISME Journal, 12(2), Article 2. https://doi.org/10.1038/ismej.2017.175

Viver, T., Cifuentes, A., Díaz, S., Rodríguez-Valdecantos, G., González, B., Antón, J., & Rosselló-

Móra, R. (2015). Diversity of extremely halophilic cultivable prokaryotes in Mediterranean,

Atlantic and Pacific solar salterns: Evidence that unexplored sites constitute sources of cultivable

novelty.   Systematic   and   Applied   Microbiology,   38(4),   266-275.

https://doi.org/10.1016/j.syapm.2015.02.002

Viver, T., Conrad, R. E., Lucio, M., Harir, M., Urdiain, M., Gago, J. F., Suárez-Suárez, A., Bustos-

Caparros, E., Sanchez-Martinez, R., Mayol, E., Fassetta, F., Pang, J., Mădălin Gridan, I., Venter,

S., Santos, F., Baxter, B., Llames, M. E., Cristea, A., Banciu, H. L., … Rossello-Mora, R. (2023).

Description of two cultivated and two uncultivated new *Salinibacter* species, one named following

the rules of the bacteriological code: *Salinibacter grassmerensis* sp. nov.; and three named

following the rules of the SeqCode: *Salinibacter pepae* sp. nov., *Salinibacter abyssi* sp. nov., and

Salinibacter pampae sp. nov. Systematic and Applied Microbiology, 46(3), 126416.

https://doi.org/10.1016/j.syapm.2023.126416

Wielgoss, S., Bergmiller, T., Bischofberger, A. M., & Hall, A. R. (2016). Adaptation to Parasites

and Costs of Parasite Resistance in Mutator and Nonmutator Bacteria. Molecular Biology and

Evolution, 33(3), 770-782. https://doi.org/10.1093/molbev/msv270

Zimmer, C., & Dorea, C. C. (2023). Differences in laboratory versus field treatment performance

of point-of-use drinking water treatment methods: Research gaps and ways forward. Npj Clean

Water, 6(1), 1-7. https://doi.org/10.1038/s41545-023-00241-1

ACKNOWLEDGEMENTS

We warmly thank the Molecular Microbial Ecology group for their valuable feedback and

constructive criticism throughout this work. We also acknowledge the Genomics and Proteomics

Unit of the University of Alicante for their assistance with flow cytometry. Special thanks to

Antonio Sanchez-Amat and Maliheh Mehrshad for their insightful feedback and suggestions. We

thank Beatriz Cámara for developing the script used in the ANIr analysis and Heather Maughan

for the professional English editing and critical reading of the manuscript. This research was

supported by the projects "VIRHOST" CIPROM/2021/006 (PROMETEO 2022, Generalitat

Valenciana) to J.A., and METACIRCLE PID2021-126114NB-C41 to F.S. and J.A. (Spanish

Ministry of Science and Innovation). R.S.M. received funding for his doctoral thesis from the

Spanish Ministry of Science and Innovation PRE2019-087998. R.S.M., E.R.P., F.S. and J.A. are

members of the National Excellence Network FAGOMA (RED2022-134837-T).

AUTHOR CONTRIBUTIONS

817 F.S. and J.A. conceived the study. R.S.M., F.S., and J.A. designed the experimental approach.

818 R.S.M. performed the experiments and analyzed the sequences under the inputs and supervision

819 of F.S. and J.A. E.R.P. designed the primers and probes for each strain and virus. L.M.R., J.S. and

820 M.E. provided advice and carried out the microfluidics-based qPCR. R.S.M. drafted the original

821 manuscript. All authors contributed to manuscript final writing and approved the final paper.

DATA AVAILABILITY

824 The raw sequences used were deposited in the NCBI database with BioProject ID PRJNA1294611.

ETHICS DECLARATIONS

827   The authors declare no competing interests.

Fig. 1: **Experimental design. a)** Five experiments were conducted using *Sal. ruber* strain M1,

with experiments 2, 4, and 5 also inoculated with M8, M31 and P18 strains. Experiments 3, 4, and

5 included the EM1 virus, which specifically infects the M1 strain. In experiment 5, the CC1 and

CR41-2 viruses, which infect the M8 and M31 strains, and the CR4 virus, which infects the M8

strain, were also added. The color bars indicate the proportion of each strain (i.e. cells/ml) or virus

(VLP/ml) in the respective experiments. All experiments were performed in triplicate. **b)** In the

short-term experiment, samples were incubated for 9 days with regular monitoring. In the long-

term experiment, samples were transferred to fresh media every 9-13 days over a 150-day period

(black and red dots), with additional parameters measured only at specific points (red dots).

Extracellular viromes from experiments 3, 4, and 5 were sequenced at 9, 84, and 150 days, as well

as the original viruses (asterisks denote the points chosen for sequencing). Figure created with

Biorender.

Fig. 2: **Influence of the biological context on the virus-host dynamics in the *Sal. ruber* M1-EM1 pair. a)** Optical density (OD 600 nm) of all five short-term experiments. **b)** PFU/ml of experiments 3, 4 and 5, measured by plaque assay against the original host *Sal. ruber* M1. **c)** Extracellular VLP/ml of experiments 3 and 4 measured by flow cytometry. All the experiments were performed in triplicate. Error bars represent the standard error of the biological replicates. Asterisks indicate significant differences based on one-way ANOVA comparing experiment 3 with experiments 4 and 5. `Other strains´ refers to M8, M31 and P18 strains and `other viruses´ refers to CC1, CR4, and CR41-2 viruses.

Fig. 3: **Concentration of the different *Sal. ruber* strains and viruses in experiments 3, 4, and**

5. a) Genomes/ml of each strain in experiments 3, 4 and 5 (top to bottom) measured by microfluidic

qPCR. **b**) Free genomes/ml in the supernatant of experiments 3, 4, and 5 (top to bottom) measured

by microfluidics-based qPCR. All experiments were performed in triplicate. Error bars represent

the standard error of the biological replicates. Asterisks indicate significant differences based on

one-way ANOVA comparing experiment 3 with experiments 4 and 5.

Fig. 4: **The biological context affects the evolution of the infectivity and host range of the EM1 virus. a)** Infectivity of EM1 virus in experiments 3, 4 and 5 in the long-term experiment, defined as the ratio between PFU/ml and extracellular genomes/ml. The y-axis represents the percentage of infectious viruses. Error bars represent the standard error of the biological replicates. Asterisks indicate significant differences based on one-way ANOVA comparing experiment 3 with experiments 4 and 5. **b)** Host range of the four viruses across the three biological replicates in experiment 5. The gray area indicates viral extinction. The time points marked in black (upper part) denote those selected for genomic analysis.

Fig. 5: **Mutations in the EM1 virus genome in the three experiments at 150 days.** The three panels outlined in black represent experiments 3, 4, and 5, from top to bottom. In each panel's upper section, mutations are marked as points across the EM1 genome, with the y-axis indicating mutations frequencies. Below each mutation plot is a representation of the genome of EM1, where arrows represent the open reading frames (ORFs). The middle plot depicts the ANIr along the viral genome, with the y-axis showing the ANIr. In the lower part of each panel, a fragment recruitment plot is shown, where the EM1 genome is positioned at the top as the reference, and gray points represent the reads recruited, with the y-axis showing the percentage of identity for each mapped read. A histogram on the right side of this section indicates the distribution of reads by identity. At the right of the figure, pie charts illustrate the protein-coding genes mutated in each experiment and the number of mutations in each ORF. Proteins selected for metabarcoding are marked with asterisks at the top of the first panel.

903 **EXTENDED DATA**

905 **Extended Data Fig. 1**: **Cells/ml and VLP/ml measured with flow cytometry. a)** Cells/ml of the

906 five experiments measured using flow cytometry. **b)** VLP/ml of the experiments 3, 4 and 5

907 measured using flow cytometry. All experiments were performed in triplicate. Error bars represent

908 the standard error of the biological replicates.

Extended Data Fig. 2: *Sal. ruber* **strains M8, M31 and P18 were exposed to virus-free lysate**
of *Sal. ruber* **M1-EM1. a)** The *Sal. ruber* M1-EM1 virus-free lysate was obtained by
centrifugation and filtration of the *Sal. ruber* M1-EM1 culture supernatant through a 0.022 μm
filter. The lysate was added at 67 hours (red dotted line) to exponential phase cultures of M8, M31
and P18 strains. **b)** Growth curves of M8, M31 and P18 amended with lysate (grey) and their
corresponding unamended controls (purple). All experiments were performed in triplicate. Error
bars represent the standard error of the biological replicates. Figure created with Biorender.

920 **Extended Data Fig. 3**: **Concentration of the different *Sal. ruber* strains in the long-term**

921 **experiment.** Genomes/ml of each strain in the long-term in experiments 1, 2, 3, 4 and 5 (top to

922 bottom) measured by qPCR. All experiments were performed in triplicate. Error bars represent the

923 standard error of the biological replicates.

928 **Extended Data Fig. 4**: **Concentration of the different extracellular *Sal. ruber* viruses in the**
929 **long-term experiment.** Genomes/ml of each virus in the long-term in experiments 3, 4 and 5 (top
930 to bottom) measured by qPCR. All experiments were performed in triplicate. Error bars represent
931 the standard error of the biological replicates.

Extended Data Fig. 5: **Mutations in EM1 virus genome in the three experiments at 84 days.**
The three panels outlined in black represent experiments 3, 4, and 5, from top to bottom. In each
panel's upper section, mutations are marked as points across the EM1 genome, with the y-axis
indicating mutation frequencies. Just below each mutation plot there is a representation of the
genome of EM1, where arrows represent the open reading frames (ORFs). The middle plot depicts
the ANIr along the viral genome, with the y-axis showing the ANIr. In the lower part of each panel,
a fragment recruitment plot is shown, where the EM1 genome is positioned at the top as the
reference, and gray points represent the reads recruited, with the y-axis showing the percentage of
identity for each mapped read. A histogram on the right side of this section indicates the
distribution of reads by identity. At the right of the figure, pie charts illustrate the mutated protein-
coding genes identified in each experiment and the number of mutations in each ORF.

Extended Data Fig. 6: Mutations in the extracellular and intracellular EM1 virus genome across the three proteins selected for metabarcoding. Each column represents one of the three selected proteins. Rows correspond to extracellular and intracellular mutations from experiments 3, 4, and 5 (top to bottom). Mutations are shown as dots along the EM1 genome, with their frequencies indicated on the y-axis.

Extended Data Fig. 7: Mutations accumulated in the nine selected regions of the chromosome of *Sal. ruber* M1 at 150 days. The different panels represent experiments 1, 2, 3, 4, and 5, from top to bottom. Mutations are marked as points across the *Sal. ruber* M1 chromosome, with the y-axis indicating mutation frequencies.