A phylogenetic estimate of canine retrotransposition rates based on genome

assembly comparisons

Matthew S. Blacksmith[1,2], Anthony K. Nguyen[1,2], John V. Moran[1,3], and Jeffrey M.

Kidd[1,2*]

[1]Department of Human Genetics, University of Michigan, Ann Arbor, Michigan, United

States of America

[2]Gilbert S. Omenn Department of Computational Medicine and Bioinformatics,

University of Michigan, Ann Arbor, Michigan, United States of America

[3]Department of Internal Medicine, University of Michigan, Ann Arbor, Michigan, United

States of America

*Correspondence to J.M.K. at jmkidd@umich.edu

Abstract

Background

Due to their history of domestication and breed formation, dogs are a powerful system for studying the phenotypic impact of genetic variation. Comparison of canine genome assemblies show that retrotransposons, mobile elements which mobilize through an RNA intermediate, are a major contributor to canine genetic diversity with an eightfold and 17-fold increase of LINE-1 and SINE differences, respectively, in dogs relative to that found among humans. The frequency of dimorphic retrotransposon insertions among dogs suggests these elements have mobilized at a high rate over recent canine evolution. However, the rate at which new insertions arise has yet to be determined.

Results

We aligned a collection of genome assemblies derived from four breed dogs, a Dingo, and an American grey wolf to a Greenland grey wolf to identify dimorphic LINE-1 and SINEC insertions. Across our panel of seven canine assemblies, we identified a total of 7,428 dimorphic LINE-1s and 51,572 dimorphic SINECs. Each assembly differs from the Greenland wolf genome by an average of 3,497 LINE-1s and 25,558 SINECs. Analysis of allele sharing among samples recapitulates known relationships and reveals substantial within-breed variation. Calibrating estimates using a previously estimated single nucleotide mutation rate of 4.5×10^{-9} per base pair per generation, we estimate that new LINE-1 and SINEC and insertions have occurred at a rate of 1/184 and 1/22 births over recent canine evolution. These estimates are largely consistent across assemblies and breeds.

Conclusions

Our phylogenetic estimate of SINEC retrotransposition in canines is approximately twice as large as that estimated for *Alu* in humans, while the canine LINE-1 insertion rate is within the range of human estimates. These data suggest that although SINEC has been an outsized driver of canine genome evolution, the striking levels of canine LINE-1 and SINEC dimorphism mainly reflect high levels of long-standing genetic variation.

Keywords: Canine, Reference Assemblies, LINE-1, SINE, Mobile Elements, Mutation Rate, Genome Assemblies, Retrotransposition

Background

Dogs, *Canis lupus familiaris*, have been a human companion for over ten thousand years. Recent findings using ancient DNA show that dogs were domesticated from a now-extinct Eurasian gray wolf lineage approximately 10-40 thousand years ago (1-4). Since domestication, dogs have been subject to extensive selection by humans based on their behavior, size, and morphology (5-9). Selection became more extreme with the formation of modern breeds in the late 19[th] century (10). This active process has seen the creation of over 300 recognized breeds worldwide that show over an order of magnitude of size difference (e.g., Great Dane: 80kg, Chihuahua: 0.5kg) (11).

The natural genetic variation found in dogs, along with the unique structure of breeds, has made dogs a powerful system for mapping the molecular basis of traits and for studying the impact of selection and population bottlenecks (9, 12-14). The potential of dogs as a model system has led to the creation of a robust canine genomics research community and catalyzed the development of multiple genomic resources including numerous genome assemblies and myriad short read datasets (15, 16). One striking finding from the analysis of these resources is the high level of retrotransposon differences found among dogs. Retrotransposons are a type of mobile element that propagates throughout the genome using a "copy and paste" mechanism via an RNA intermediate (reviewed in (17)). Retrotransposons are further subdivided into two classes, those that contain long terminal repeats (LTRs) and those that do not. Non-LTR retrotransposons, which include Long INterspersed Elements (LINEs) and Short INterspersed Elements (SINEs), are highly prevalent in the canine genome (18).

Comparisons between the initial dog reference genome, derived from a Boxer

breed dog named Tasha, and low-coverage survey sequencing from a Poodle breed

dog, named Shadow, revealed over 10,000 SINEs with presence-absence differences

(known as dimorphism) between the two assemblies (18, 19). Comparison of genome

assemblies from a Boxer and a Great Dane identified over 16,000 SINECs and 1,100

LINE-1s that differed between the assemblies (20). Subsequent analyses demonstrated

that a full-length LINE-1 cloned from the Great Dane genome was active in a cell culture

based retrotransposition assay (20, 21). The cloned canine LINE-1 sequence was also

capable of driving SINEC retrotransposition in cell culture (20, 22). Recent years have

also seen an explosion in canine genome comparisons, further confirming the outsized

contribution of retrotransposon sequences to canine genetic variation (16, 20).

LINE-1, the family of LINEs currently active in mammals, including canines, is

approximately 6 kb in length and encodes two proteins (23, 24). ORF1p contains RNA

binding and nucleic acid chaperone activities, while ORF2p contains domains

responsible for nicking genomic DNA (endonuclease) and copying the element-encoded

RNA into DNA (reverse transcriptase) (25-30). LINE-1 retrotransposition begins when

LINE-1 is transcribed by RNA Pol II, using the internal promotor present in the LINE-1 5'

UTR (31, 32). LINE-1s also possess a polyadenylation signal and are dependent upon a

3' poly(A) tail to fully proceed through the retrotransposition cycle (33-35). Following

transcription, LINE-1 RNA is exported to the cytoplasm where the bicistronic RNA is

translated by an unconventional mechanism (36). In humans, ORF1p is translated to a

higher degree and is reliably present in cytosolic fractions (28, 37). By contrast, ORF2p

is produced in lower abundance and co-translationally associates with the 3' poly(A) tail

of the LINE-1 RNA that encodes it (35, 38, 39). Once bound by both ORF1p and

ORF2p, the ribonucleoprotein particle is imported back into the nucleus to undergo a

process termed target primed reverse transcription (TPRT) (21, 29, 40-42). During

TPRT, the ORF2p endonuclease activity creates a nick in genomic DNA at a new locus,

with a preference for the ORF2p canonical cleavage site of 5'-TTTTT/AA (29, 43, 44).

Subsequently, the ORF2p reverse transcriptase activity is used to generate a DNA copy

of the LINE-1 RNA which, following second strand synthesis, results in the insertion of a

copy of the LINE-1 RNA into a new genomic locus (21, 30, 45). LINE-1 insertions are

often severely 5' truncated or contain other mutations that render them inactive (46-48).

LINE-1 mediated insertions possess several hallmarks which can be used to

characterize the insertion (17) (Reviewed in (49)). The first, mentioned above, is the

presence of the canonical endonuclease cleavage site, 5'TTTTT/AA, at the site of

insertion (21, 29, 43, 44) . Second, is the presence of a 3' poly(A) sequence which is

placed by the poly(A) polymerase post-transcriptionally and is required for the binding of

ORF2p (33, 35). Third, is the presence of target site duplications (TSDs), which arise

during TPRT and flank the insertion (43, 49, 50). Finally, while not present in all LINE-1

insertions, 3' transductions are frequently identified in LINE-1 datasets (21, 33, 51).

LINE-1 has a weak termination signal and consequently, RNA polymerase II may read

through the termination signal and into downstream sequence (21, 33, 51, 52). This

process creates a transcript containing LINE-1 and downstream genomic sequences,

which can be used to infer the identify of a parent element associated with an insertion

(20, 51, 53-56).

In addition to mobilizing the transcripts that encoded them (known as *cis-*

preference (39, 41)), LINE-1 encoded proteins are able to mobilize other RNAs (57), the

primary example of which is SINEs. SINEs do not encode proteins, but instead rely

upon LINE-1 ORF2p to mobilize (22). A family of SINEs, termed SINEC, is present

throughout *Carnivora* and is abundant in the canine genome (18, 19, 58). SINEC

sequences are transcribed by RNA polymerase III and contain a 5' 'head' region derived

from tRNA-lysine, a segment of $(CT)_N$ repeats of variable lengths, and an A-rich tail (58-

60). It is hypothesized that SINEC may retrotranspose via two similar mechanisms: one

that utilizes a cellular poly(A) polymerase to generate a poly(A) tail, and one that uses

the existing poly(A) sequence as a poly(A) tail (61). LINE-1 encoded proteins can also

mobilize other cellular RNAs, leading to the production of retrocopies, also known as

processed pseudogenes or retrogenes (41, 57).

LINE-1 mediated insertions contribute to, or are associated with, numerous

canine phenotypes and diseases (7) . Perhaps the most well-known are the multiple

functional retrocopies of the *FGF4* gene that cause a chondrodysplasia-like short-

legged phenotype in breeds such as the Pembroke Welsh Corgi, Basset Hound,

Dachshund, and Nova Scotia Duck Tolling Retrievers (62, 63). LINE-1 insertions are

also implicated in progressive retinal atrophy in Swedish Vallhunds, Duchene-like

muscular dystrophy in Pembroke Welsh Corgis, and brachycephaly in several breeds

(64-66). Conditions such as polyneuropathy are caused by a SINEC insertion into

RAB3GAP1 in Alaskan Huskies (67), and variability in a SINEC insertion causes the

merle coat color pattern found in some breeds (68, 69). Importantly, a recent *de novo*

LINE-1 insertion into the *DMD* gene has been identified in a Border Collie,

demonstrating that LINE-1 continues to actively mobilize in the canine genome (70).

The frequency of dimorphic LINE-1 and SINEC insertions among dogs may

reflect both a high mobilization rate over recent canine evolution and the assortment of

standing genetic variation that arose in ancestral wolf populations (6, 7). Using LINE-1

and SINEC dimorphic elements identified by comparison of canine genome assemblies,

we produce a phylogenetic estimate of the rate of LINE-1 and SINEC retrotransposition

over recent canine evolution that is calibrated using a previously estimated rate of SNV

mutation. Our analysis confirms the high degree of LINE-1 and SINEC dimorphism in

canines, including between individuals of the same breed. We estimate that new LINE-1

and SINEC insertions arise at a rate of 1/184 and 1/22 births respectively in canines.

Methods

Identification of dimorphic loci

Structural variants that correspond to dimorphic LINE-1 or SINEC elements were

identified from seven published canine genome assemblies, including two German

Shepherd Dogs, one Boxer, one Great Dane, one Dingo, and two wolves (20, 71-76)

(Supplemental Data S1). For ease of description, we refer to each assembly based on

the shortened name given in Table 1 and Supplemental Data S1. Variants were

identified by aligning each genome to that of the Greenland wolf (G_WOLF), which was

assembled using highly accurate PacBio HiFi reads and serves as an outgroup to

domestic dogs.

Each of the six assemblies were aligned as queries using the G_WOLF

assembly as the target using minimap2 [version 2.26] (77) with options -c -x asm5 --cs.

The resulting alignment files in paf format were then sorted (sort -k6,6 -k8,8n) and

processed using the paftools.js call utility with default parameters to create a listing of

variants identified from each pairwise comparison. Single Nucleotide Variants (SNVs),

and structural variants (SVs) at least 50 bp in size, were extracted from the variant list

and filtered to include only variants on the autosomes (chr1-chr38) or X chromosome

and where the query and target sequences occur on the same chromosome.

Next, a list of exclusion regions was aggregated, including segmental

duplications identified by BISER (78) and/or fastCN (79), as well as genome assembly

gaps. Duplication coordinates from G_WOLF, GSD1, GSD2, BOX, GDN, and DNG

were obtained from Nguyen, Blacksmith, and Kidd, 2024 (80). Duplications were

identified in A_WOLF following the same procedure (80). Because Illumina reads of

A_WOLF are not directly available, we used Illumina reads from the wolf parent of

A_WOLF. Exclusion regions were defined in both the query and target. Any SV within

100 bp of an excluded region was removed from the dataset using bedtools window

with options -w 100 -v (81). SVs were then split into "insertions" and "deletions" before

being further divided into candidate LINE-1 and SINEC elements based on comparison

with RepeatMasker annotations using bedtools intersect with options -wa -f 0.7. The -f

option requires that 70% of each SV is represented in the corresponding RepeatMasker

file (82). LINEs belonging to the HAL family were excluded from analyses (83). This

criterion is designed to distinguish structural variants that correspond to dimorphic

mobile element insertions from other structural variants that contain mobile element

sequences. In our informatics pipeline, we defined insertions as variants present in an

analyzed sample but absent from G_WOLF and deletions as variants present in

G_WOLF but absent from the sample being compared. In reality, dimorphic mobile

element insertions represent two states: (i) the ancestral "empty site" allele without the

element and (ii) the derived or "filled site" allele that contains the mobile element. The

mobile element content of the filled site allele identified for each variant was determined

based on the annotation of the appropriate genome assembly. Annotations were

determined based on RepeatMasker (version 4.0.7 Database: dc20170127-

rb20170127). The outputs were then sorted (sort -k1,1 -k2,2n) and retained for later

use. The sample standard deviation for the number of SINECs present across the

dataset was calculated using numpy.std(data,ddof=1) (v1.20.3).

Identifying the hallmarks of retrotransposition in SINEC and LINE-1 loci

The hallmarks of LINE-1 retrotransposition (i.e., TSDs, 3' poly(A) sequences, and

endonuclease cleavage sites) in dimorphic mobile element containing loci were

identified by comparing the filled (SINEC or LINE-1 present) and empty (SINEC or

LINE-1 absent) sequences. For each locus, the filled and empty site sequence,

including 1,000 bp of flanking sequence in each direction, were extracted with samtools

faidx (v1.21) (84) and aligned with AGE (v0.4) (85). AGE is an alignment software

designed for the analysis of structural variants and can identify the presence of

matching sequences located at rearrangement breakpoints. Because AGE can only

accurately resolve one structural variant per alignment, if the variant identified by AGE

is more than 150 bp away from the coordinate originally identified from the minimap2

alignment, the AGE processing was repeated using a reduced flanking size of 500 bp. If

necessary, this process was additionally repeated two times using flanking sizes of 250

bp and 125 bp respectively. If consistent coordinates could not be identified, the locus

was excluded from further processing. Several other filters were also introduced,

including removal of loci which are: (**1**) less than 50 bp in length after AGE processing

or (**2**) where the number of inserted base pairs that are not called as TSD is below 30.

The presence of deletions in existing LINE-1 sequences gave rise to systemic LINE-1

false positives in this dataset. Two additional filters were used to remove these loci from

further analysis. First, for each inserted sequence identified by AGE, RepeatMasker

was used to directly assess the LINE-1 content of the inserted sequence. If less than

70% of the sequence was identified as LINE-1 sequence using this method, the locus

was discarded from further processing. Second, utilizing the original whole genome

RepeatMasker annotation, loci which have a RepeatMasker identified segment

extending beyond the inserted sequence as identified by AGE were checked for the

presence of a TSD greater than or equal to 10 bp. If less than a 10 bp TSD was

identified, these loci were discarded. For remaining SINEC and LINE-1 loci with

concordant coordinates identified by AGE and minimap2, over 99.8% of TSDs identified

by AGE show perfect identity among the upstream TSD, downstream TSD, and the

corresponding empty site. TSDs 10 bp or longer were considered "high confidence."

218   To identify 3' poly(A) tails, all sequence between TSDs, or the end of the

insertion if no TSD is present, were searched for homopolymers of "A" or "T" depending

on element orientation. Element orientation was identified by bedtools intersecting the

query sequence coordinates mentioned above with SINECs or LINE-1s identified via

RepeatMasker (81, 82). RepeatMasker segments smaller than 20 bp were not used for

orientation detection. If multiple RepeatMasker segments are detected in a single locus

and all segments are in the same orientation, the locus is fully processed; otherwise, the

locus is retained but poly(A) tails are not identified. If no RepeatMasker segments meet

this requirement the locus is removed from processing. Homopolymers separated by 5

or fewer bases from the TSD (or insertion boundary if no TSD is present) were

classified as poly(A) tails. If two or more homopolymers meet these criteria, the longer is

chosen as the poly(A) tail. Only identified segments that are 10 bp or longer are

retained as poly(A) tails.

Endonuclease cleavage sites were only identified in loci possessing a TSD of 10

bp or longer and a defined element orientation. Presuming that a locus is in the forward

orientation relative to the reference, the endonuclease cleavage site consists of the two

nucleotides upstream of the 5'TSD and the first 5 nucleotides of the TSD. All

endonuclease cleavage sites are reported on the minus strand and thus should

correlate with the known "5'-TTTTT/AA" cut site (43, 44). Sequence logoplots were

created using logomaker (v0.8) (86). Other figures associated with this section were

generated using numpy (v1.20.3) (87), pandas (v1.3.4) (88), matplotlib (v3.4.3) (89),

and scipy (v1.7.1) (90).

Visualizing loci with miropeats

Filled and empty site sequences corresponding to dimorphic elements were

extracted with samtools faidx (84). Each sequence is then investigated for repeats using

RepeatMasker with the option --species dog (82). From there, a baseline miropeats

image is generated using miropeats with options -onlyinter -s 200 (v2.02) (91). The

miropeats image is then annotated to display RepeatMasker information (v4.0.7).

Identifying SINEC and LINE-1 subfamilies

To identify which subfamily each identified variant belongs to, coordinates

corresponding to the sequence between target site duplications, or the full locus if no

TSD is present, were intersected with all SINEC loci identified in genome-wide

RepeatMasker annotation using bedtools intersect (81, 82). As in 3' poly(A)

identification, the intersected locus must be at least 20 bp in length. If multiple different

subfamilies were identified, the locus is not uniquely assigned to a subfamily.

RepeatMasker reports 21 different SINEC subfamilies. This includes SINEC_Cf, a

consensus sequence used by RepeatMasker that is not present in Repbase. Because

the tRNA-derived head region of the SINEC_Cf consensus is identical to the

SINEC2A1_Cf sequence, with the only differences being the length of the internal $(CT)_N$

repeat, the length of the encoded poly(A) tail, and the presence of an extra T in the 3'

end of the element, we combined RepeatMasker annotations into a single subfamily we

refer to as SINEC_Cf/2A1. Similarly, LINE-1 subfamilies were identified using the above

methods.

Identifying shared dimorphic SINEC and LINE-1 loci

To identify which SINEC and LINE-1 loci were shared across the dataset, we first

identified autosomal regions of the G_WOLF genome that were covered by one query

contig in each comparison, as reported by paftools.js call. The callable regions were

further filtered to remove segments corresponding to duplications and gaps in the

G_WOLF assembly as described above. We then created a bed file containing the

intervals that were not callable in at least one sample and removed any dimorphic

SINEC locus that intersected with the non-callable regions. Remaining SINEC loci were

then aggregated, sorted (with utilities sort -k 1,1 -k 2,2n -k 3,3n), and merged with

bedtools merge -d 100 -c 7,4 -o collapse, collapse. These parameters merged any loci

within 100 bp of each other in G_WOLF coordinates. Any merged loci that contained

more than one variant were removed from further processing. An UpSet plot was then

created to summarize the pattern of sharing across the sample set using the UpSetPlot

module (version 0.9.0) (92).

To estimate a phylogenetic tree from dimorphic SINECs, we encoded

presence/absence data for each sample as a sequence in fasta format with a single

nucleotide representing a single SINEC locus. We supplemented this with an additional

sequence encoding the empty site for each locus which represents the ancestral state.

A neighbor joining phylogenetic tree was then estimated from the resulting sequencing

using MEGA (v11.0.13) with the p-distance mutation model, which is simply the

proportion of sites that differ (93, 94). Uncertainty in tree typology was assed using

1,000 bootstrap replications (95). The bootstrap consensus tree was rooted using the

ancestral empty-site sequence described above. Individual SINEC loci were assigned to

specific branches on the inferred tree based on their presence/absence pattern across

samples. A second phylogenetic tree was also generated using the above methods on

dimorphic LINE-1 insertions. Final tree visualizations were generated using MEGA and

FigTree (93, 96).

Examination of within-breed variation

To determine whether variation between two GSDs is consistent with that found

within other breeds, we performed within-breed comparisons using two GSDs (GSD1

and GSD2), two Labrador retrievers (LAB1 and LAB2), and two Bernese Mountain dogs

(BMD1 and BMD2). For each pair of samples, the assembly with the smallest segment

count was designated as the target, with the other sample being the query. Segment

count was calculated as number of non-'N' segments assigned to the autosomes and

chrX (Supplemental Data S1). A minimap2 alignment with downstream variant

identification was then performed as described above. Within these samples, segmental

duplications were identified using BISER and fastCN as described in Nguyen,

Blacksmith, and Kidd (80).

Identification of dimorphic SINEC variants within GSD1

Dimorphic SINECs were identified from a dataset of structural variants identified

in GSD1 (97). Specifically, Schall et al. identified structural variants in GSD1 short reads

aligned to the GSD1 reference assembly using Manta and genotyped the variants with

GraphTyper2 (98, 99). Heterozygous autosomal deletions identified in GSD1 were

aggregated and then intersected (bedtools intersect -wa -f .9 -r) with SINECs present in

GSD1 and absent in G_WOLF (81). These parameters require 90% reciprocal overlap

between the dimorphic SINEC and the structural variant.

Extracting SNVs from inter- and intra-breed comparisons

To identify SNVs relative to G_WOLF or within a single breed, single nucleotide

substitutions (i.e., not single base pair deletions or insertions) and regions

corresponding to a single aligned segment were extracted from each minimap2

alignment as described above. For each comparison, low confidence regions from the

minimap2 target genome were filtered out. This included gaps, BISER (78) and/or

fastCN (79) segmental duplications, and tandem repeat finder (TRF) (100) identified loci

present in either the query or reference. The low complexity regions were removed from

aligned regions with bedtools subtract. Remaining segments were then intersected with

SNVs with bedtools intersect to create a final list of SNVs present on the autosomes

(81). Note that the autosomal genome segments were only filtered against low

complexity regions present in the reference (*i.e.* G_WOLF or the genome with the lower

segment count in other comparisons).

Estimating a rate of canine LINE-1 and SINEC retrotransposition in canines

To estimate a rate of SINEC retrotransposition in canines, the first requirement is

identifying the number of generations since genetic divergence between each query

genome the target (G_WOLF). We combined the filtered number of base pairs in

autosomal aligned regions with the number of autosomal SNVs and the estimate canine

mutation rate of 4.5×10^{-9} per bp per generation (range 2.6×10^{-9}/bp/gen - 7.1×10^{-9}

^{9}/bp/gen) (101) to estimate the number of generations since genome divergence

between each sample pair as was done in Nguyen, Blacksmith, and Kidd (80).

$$Genome\ Divergence\ = \frac{SNV\ count \times 0.5}{Autosomal\ genome\ length} \times \frac{1}{Wolf\ mutation\ Rate}$$

From there, we estimated the rate of LINE-1 and SINEC insertions as the ratio of

the number of generations divided by the number of autosomal dimorphic variants.

Estimates were created for dimorphic variants present only in the query, only in target,

and the average of both. As an estimate of the uncertainty in the obtained values, we

also report insertion rates implied by the range of reported SNV mutation rates. Point

estimates of the rate of retrotransposition were also recalculated using only loci which

clearly have "high confidence" TSDs of at least 10 bp.

Identifying LINE-1 3' transductions

LINE-1s were assessed for the presence of 3' transductions. First, for each

genome, the coordinates of dimorphic LINE-1s were intersected with a bedfile

containing the positions of LINE-1s in the genome annotated using bedtools intersect

(version 2.26.0), with option -wao. The longest intersecting annotation was chosen for

each locus, and the sequence of candidate 3' transductions spanning from the end of

the LINE-1 annotation to the end of annotated variant, in the orientation of the LINE-1,

were extracted. If the locus possessed a 10 bp or longer TSD, the TSD sequence was

removed from the candidate 3' transduction. Candidates shorter than 25 bp were

discarded. The position of repeats and low complexity sequence were identified in each

candidate transduction using RepeatMasker (--species dog) and sdust (82, 102). Any

candidate that did not have at least 25 bp of unmasked sequence was discarded. The

original, unmasked sequence from the remaining candidate 3' transductions was

searched against the G_WOLF reference using blat (version 35, -out=pslx -

minIdentity=95) (103). Blat alignments were removed if the match count was less than

25, if the blat target was not on an autosome or chrX, if the target coordinates are within

10 kb of the dimorphic locus, if the span of the aligned target sequence is 100 bp more

than or less than the length of the query sequence, if the reported alignment included

fewer than 25 bp of unmasked query sequence, if over 50 blat alignments were

produced for a single sequence, or if a single locus corresponded to more than five 3'

transductions in a single genome. All blat alignments were aggregated using bedtools

sort and bedtools merge (-d 100). The resulting aggregated alignments were parsed to

identify if any transduction source corresponds to multiple LINE-1s with transduced

sequence in this dataset.

We also identified if blat alignments identified loci that are adjacent to LINE-1s.

To do so, all blat alignments mentioned above were aggregated, sorted with bedtools

sort, and intersected with G_WOLF LINE-1s using bedtools window (-w 50).

Intersections were then filtered to remove LINE-1s present only in the transduced

sequence as detected by blat or LINE-1s which encompass the entire transduced

sequence. Remaining alignments were then assessed to note if 1: the transduction was

"downstream" of the LINE-1, and 2: that the orientation of the transduced sequence is in

the expected orientation. If both criteria are present, the G_WOLF LINE-1 is considered

"adjacent" to the transduction.

Finally, we investigated if any transduced sequences arise adjacent to multiple

independent dimorphic LINE-1 insertions. The locations of aligned transductions in

G_WOLF coordinates were aggregated, sorted with bedtools sort and merged with

bedtools merge (-d 100).

Rescaling human retrotransposition rates

Both phylogenetic and population modeling estimates of the rate of

retrotransposition are highly sensitive to estimates of the rate of SNV mutagenesis. This

complicates comparisons as estimates of the human SNV mutation rate have changed

over time. Early estimates such as those provided by Nachman and Crowell were

around 2.5×10^{-8} per nucleotide per generation (104). However, more recent estimates

cluster between 1.1×10^{-8} and 1.45×10^{-8} per nucleotide per generation (105-108). As a

result, we rescaled existing rates using a mutation rate of 1.3×10^{-8} mutations per

nucleotide per generation.

Additionally, estimates of effective population size (N_e) are inversely correlated to

estimates of the SNV mutation rate. Mallick et al. estimated the rate of human

heterozygosity on a population-by-population basis. Heterozygosity was mostly

constrained to the interval between .0005 and .001 (109). This estimate places the long-

term human effective population size between 10,000 and 20,000 (109, 110). For

clarity, rescaled estimates also use a human effective population size of 15,000, in the

center of the range.

Results

Pairwise comparisons reveal thousands of dimorphic SINECs

Dimorphic SINEC insertions were identified in a panel of canine genome

assemblies including two German Shepherd Dogs (GSD1, GSD2), a Boxer (BOX), a

Great Dane (GDN), a Dingo (DNG), and two grey wolves (A_WOLF, G_WOLF) (20, 71-

76) (Table 1, Supplemental Data S1). Each sample assembly was aligned as a query to

the G_WOLF reference using minimap2 (77). G_WOLF was chosen as the common

target because wolves are an outgroup relative to breed dogs and Dingos, and because

the G_WOLF genome was sequenced using PacBio HiFi sequencing. For each

pairwise genome alignment, structural variants 50 bp or larger, having at least 70%

SINEC content, and present on the autosomes (chr1-chr38) or X chromosome were

extracted for further comparisons (Fig. 1, Fig. 2). Due to differences in assembly quality

and the expected evolutionary history of the X chromosome, only variants on

autosomes are considered unless otherwise noted.

Sample Key*	Breed/Species	Sample Name	Assembly Name	Assembly Accession
G_WOLF	Greenland wolf	mCanlor1.2	mCanlor1.2	GCA_905319855.2
A_WOLF	American wolf	Clu	GCA_034620435.1_Clu-1	GCA_034620435.1
GSD1	German Shepherd Dog	Mischka	UU_Cfam_GSD_1.0/canFam4	GCA_011100685.1
GSD2	German Shepherd Dog	Nala	CanFam_GSD	GCA_008641055.3
BOX	Boxer	Tasha	Dog10K_Boxer_Tasha_1.0/canFam6	GCA_000002285.5
GDN	Great Dane	Zoey	UMICH_Zoey_3.1/canFam5	GCA_005444595.1
DNG	Dingo	Sandy	CanFam_DDS	GCA_003254725.2

*Sample Key refers to the name of the sample used in figures and legends throughout the paper.

Table 1: Sample demographics and accession information.

On average, each genome comparison identified 25,558 dimorphic SINECs

(range: 23,284-27,422) on the autosomes (Fig. 3). The number of SINECs present in

G_WOLF but absent from the aligned sample is more consistent across comparisons

(mean=12,789, std 337) than the number of SINECs present in each sample but absent

from G_WOLF (mean=12,769, std 1,592). This difference is driven in part by the deficit

of dimorphic SINECs detected in BOX (Fig. 3A). The decreased count of SINECs in

BOX is found across all autosomes (Fig. S1) and is consistent with a previous

description of biased representation of heterozygous SINECs in this assembly (73).

We next refined the coordinates of candidate dimorphic SINECs using the

alignment with gap excision (AGE) software (85). After refining the variant coordinates,

we searched for the hallmarks of retrotransposition at each locus: the presence of

TSDs, 3' poly(A) tails, and a LINE-1 endonuclease cleavage site (Fig. 4). We began by

examining variants present in an alignment between GSD1 and G_WOLF. Dimorphic

SINEC loci present in either GSD1 or G_WOLF have similar TSD and 3' poly(A)

lengths. Additionally, loci that possess a 'high confidence' (i.e., 10 bp or longer TSDs)

contain a sequence profile consistent with the known LINE-1 endonuclease consensus

cleavage site (29, 43, 44). Overall, of the variants identified in GSD1, 52.5% have high

confidence TSDs and poly(A) tails, 31.2% have only a high confidence TSD, 5.8% have

only a high confidence poly(A), 10.5% have neither hallmark, and 0.1% have

ambiguous RepeatMasker orientations (Supplemental Data S2). A comparison of TSD

and poly(A) tail lengths indicates that loci with a high confidence TSD possess a wide

range of poly(A) tail lengths while loci with short or non-existent TSDs often have short

or non-existent poly(A) tails (Fig. S2). Hallmarks are found in similar proportions across

each of the analyzed assemblies (Fig. 5). RepeatMasker analysis shows that 95.2% of

dimorphic SINECs present in GSD1 but absent in G_WOLF belong to the

SINEC_Cf/2A1 family. Restricting analysis to loci with high confidence TSDs increases

this proportion to 96.3%. A similar pattern is found for loci present in G_WOLF but

absent in GSD1 (Fig. S3).

High SINEC diversity in canines

To identify variant sharing across the dataset, we identified 2,048,395,076

autosomal bp (in G_WOLF coordinates) that passed callability criteria in each

comparison. Of the 51,572 autosomal dimorphic SINEC insertions retained for analysis,

23,865 (46.3%) were present in only one of the seven analyzed assemblies. G_WOLF,

A_WOLF, and DNG all possessed more than 5,000 singleton variants. By contrast,

each breed dog had between 1,291 and 2,293 singletons, likely a result of the loss of

variation during breed formation. BOX had the fewest singletons, and both GSD1 and

GSD2 contained fewer singletons than GDN. GSD1 and GSD2 had the most pairwise

SINEC variants in common, 1,819, that were exclusively shared, yet each also

contained 1,827 and 1,763 singletons. Close behind with 1,794 variants exclusively

shared is A_WOLF and G_WOLF. These data highlight the exceptional amount of

within-breed SINEC variation present in canines (Fig. S4).

We used dimorphic SINECs to infer a neighbor joining tree across the samples,

rooting the tree with an ancestral state sequence represented as the empty site for each

locus (Fig. 6). As expected, all breed dogs form a group with the two GSDs forming a

pair and the Boxer and Great Dane forming a pair. Outside of breed dogs is the Dingo

which is more related to breed dogs than to wolves but falls outside of the breed dog

grouping. The two wolves form a separate taxonomic pair. The tree has a starlike

phylogeny with relatively long terminal branches and comparatively shorter internal

branches. All nodes had 100% bootstrap support within this dataset except for the node

containing GDN and BOX, which had only 76% bootstrap support (Fig. S5). The

genotypes of 58.2% of the loci match the inferred tree. The top non-conforming

categories include variants present in all samples except A_WOLF (1359, 2.64%), all

samples except G_WOLF (947, 1.84%), and present in A_WOLF and DNG (657,

1.27%). A total of 1,827 and 1,763 SINECs were assigned specifically to the GSD1 and

GSD2 branches, respectively.

We performed two analyses to confirm the large amount of within-breed SINEC

variability. First, we generated within-breed alignments of genome assemblies from the

two GSDs, as well as two Labrador retrievers and two Bernese Mountain dogs, referred

to as GSD1 and GSD2, LAB1 and LAB2, and BMD1 and BMD2, respectively. In each of

these comparisons we determined the number of SINECs and SNVs that differ between

the assemblies. LABs had the most diversity, with each assembly having ~6,900

SINECs that were absent in the other. BMDs had the lowest count of dimorphic SINECs

(~4,900 per lineage). Because the amount of within-breed genetic diversity differs, we

normalized dimorphic SINEC counts by the number of SNVs found between

assemblies, revealing a relatively consistent SINEC to SNV ratio of 0.0056-0.0065. (Fig.

S6).

We note that in the GSD comparison there appears to be an increase in

dimorphic SINECs between GSD1 and GSD2 relative to the phylogenetic analysis

above. Several factors contribute to this finding. First, while there are 3,590 SINECs

present exclusively in GSD1 or GSD2 in our dataset, there is a total of 10,569 SINECs

that are present in either GSD1 or GSD2 but not both. This includes 6,979 loci that do

not fit the inferred tree topology and are not depicted in Fig. 6 showing that this method

recapitulates 94% of variants that are identified in the pairwise comparison of GSD1 and

GSD2.

Second, we compared SINEC annotations with previously identified deletions

identified from GSD1 Illumina read data aligned to the GSD1 assembly to identify loci

where the assembly contains the insertion allele at a heterozygous locus (97). This

analysis identified 4,167 heterozygous SINEC variants on the autosomes. GSD1 had a

total of 13,901 SINECs present that were absent in G_WOLF, suggesting that ~30.0%

of SINECs with differential presence between GSD1 and G_WOLF are heterozygous in

GSD1. Both measures suggest that there is extensive within-breed, and within-

individual, SINEC variation in canines.

A phylogenetic estimate of the rate of SINEC retrotransposition in canines

We estimated the rate of SINEC insertion in canines calibrated by genome

divergence implied by a published per-generation SNV mutation rate estimated in

wolves. On average, each genome differs from G_WOLF at 4,190,406 autosomal single

nucleotide positions (Fig. 7A). Assuming a mutation rate of 4.5×10^{-9} bp/gen (101), and

based on the length of the autosomal genome determined as alignable in each

comparison, we estimate that average genome divergence of each sample relative to

G_WOLF is 228,568 generations (Fig. 7B). Utilizing SINECs present in each sample

and absent in G_WOLF, as well as SINECs present in G_WOLF and absent in GSD1, a

SINEC insertion is estimated to occur in 1/18.2 (stdev = 2.0) births. This rate is

somewhat deflated due to the low estimated rate of retrotransposition in BOX (1/22.2)

births (Fig. 7C). The rate along the G_WOLF branch implied by each comparison is

relatively constant, further suggesting that the BOX genome appears to have a biased

representation of SINECs (Fig. S7). When only including SINECs with a TSD of at least

10 bp in length, the average point estimate of the rate of retrotransposition was reduced

to 1/21.9 births (Fig. 7D). We note that this estimate is highly sensitive to the assumed

SNV mutation rate. Using the lower and upper bounds of the reported SNV mutation

rate implies a stringent SINEC insertion rate of 1/13.9 to 1/38.0 births (Supplemental

Data S3).

Canine samples possess thousands of dimorphic LINE-1 insertions.

Using methods similar to those for SINEC identification, we identified dimorphic

LINE-1s among the seven analyzed assemblies. Because deletion of sequence within

existing LINE-1 elements could be erroneously identified as a dimorphic element, we

implemented two additional criteria to enrich for dimorphic LINE-1s. First, we ran

RepeatMasker directly on the inserted sequence and discarded loci that had less than

70% LINE-1 sequence. Second, we removed loci that are embedded within a LINE-1

annotation that do not have a TSD of 10 bp or longer. Using these criteria, an average

of 3,497 (range 3,070-3,757) dimorphic autosomal LINE-1 insertions were found in the

samples (Fig. S8). In contrast to the profile observed for SINECs, there is no reduction

of LINE-1s found in BOX; however, there is a reduction of dimorphic LINE-1 variants

found in the A_WOLF assembly. When looking at the variants for which the sample is

the filled site, an average of 1.90-fold excess of LINE-1 insertions on the X chromosome

relative to the similarly sized chromosome 1 was identified (Fig. S8), consistent with an

X-chromosome bias previously observed for LINE-1 (111).

Putative LINE-1 insertions were queried for the presence of the hallmarks of

LINE-1 retrotransposition. While the distributions for TSD lengths and endonuclease

cleavage site sequences were similar to those discovered in SINECs, a greater

proportion of LINE-1 loci have poly(A) tails longer than 40 bp (Fig. S9, Fig. S10). Across

the dataset, an average of 43.8% of loci possess both high confidence hallmarks in

proximity to each other. Another 27.2% have TSDs only, 10.4% have 3' poly(A)s only,

13.8% have neither hallmark, and 4.8% have multiple LINE-1 segments in different

orientations for which we did not assess the presence of hallmarks in this analysis (Fig.

8, Supplemental Data S4). In GSD1, RepeatMasker analysis shows that 81.7% of loci

belong to the most recent LINE-1 subfamily, L1_Cf, while 6.4% and 6.4% of loci are

classified as belonging to the L1_Canis1 and L1MEc subfamilies. The remaining loci

were spread across numerous subfamilies with no single subfamily comprising more

than 1% of the dataset. Any locus which had segments annotated as belonging to

multiple subfamilies by RepeatMasker was reported as unresolved (Fig. S11). When

restricting loci to resolved variants which have a high confidence TSD (Fig. 8), the

fraction of L1_Cf variants drops slightly to 79.3% while the L1_Canis1 and L1Mec

families increase to 7.6% and 7.5%, respectively.

Dimorphic LINE-1 insertions possess 3' transductions

Next, we identified 3' transductions that could be mapped to the G_WOLF

genome from dimorphic loci. For each LINE-1 containing locus, the sequence from the

end of the LINE-1 until the start of the TSD was extracted. Candidate sequences were

disqualified from further analysis if they consisted solely of repetitive or low complexity

sequence. When looking at variants present in any sample but not G_WOLF, or

G_WOLF but not GSD1 (to avoid counting G_WOLF multiple times in averages), the

average genome possessed 63 dimorphic LINE-1 insertions with eligible 3'

transductions that meet our filtering requirements. Sequences were then searched

against the G_WOLF reference genome to identify other genomic locations possessing

the same sequence (blat hits with >=95% identity). This conservative method identified

an average of 35.6 dimorphic LINE-1s per sample with qualifying transductions from the

average of 1,913 LINE-1 insertions per genome (Supplemental Data S5). This fraction

(~1.8%) is substantially lower than the ~15% of full-length LINE-1s with transductions

predicted in humans (112). There are several reasons for this discrepancy. First, our

methodology requires that at least 70% of the inserted sequence corresponds to LINE-

1. As a result, truncated LINE-1 insertions that have long 3' transductions that include

non-LINE-1 sequence may fall below the detection threshold. Consistent with this

potential bias, we note that transductions appear to be more prominent in longer LINE-

1s. For variants present in GSD1 and absent in G_WOLF 1068 (54.4%) and 494

(25.2%) are at least 500 bp and 6000 bp respectively. By contrast, 23 (85.2%) and 12

(44.4%) transduction possessing loci are at least 500 bp and 6000 bp respectively.

Second, for a *bona fide* transduction to be identified, the sequence needs to be unique

(i.e. not composed of low complexity or repetitive sequence). This parameter reduces

the number of eligible transductions by ~40%.

Focusing on variants present in GSD1 and absent in G_WOLF, a total of 27

dimorphic LINE-1s possessed 3' transductions which map to G_WOLF. Each of these

loci correspond to a single location in the G_WOLF genome. It was noted in Halo et al.

that most dimorphic LINE-1 3' transductions are "parentless" (20). Parentless loci occur

when the source of the transduced sequence is not adjacent to a LINE-1, because the

"parent" locus is dimorphic in the population. Of the 27 transduction-possessing LINE-

1s, only three were 50 bp or less downstream of a LINE-1 found in the G_WOLF

genome.

We next identified whether any transduction source corresponds to multiple

transduction events in our dataset. Ten such loci exist where one G_WOLF source

coordinate corresponds to multiple transductions across out dataset. For two pairs of

loci, there were multiple G_WOLF coordinates which could potentially exist as the

transduction source. Factoring in the ambiguous loci, a total of eight unique transduction

sources were identified which contributed to multiple independent transduction events in

the dataset. Some sources, such as G_WOLF chr21:46673641-46673765, correspond

to multiple insertions in the same genome. The chr21 locus can be found as transduced

sequence in LINE-1 insertions present on DNG chr12 and DNG chr28. Others

correspond to multiple loci in different genomes, such as G_WOLF coordinates

chr7:21713470-21713969 which correspond to transductions present at four separate

locations: chr15 in BOX and DNG, chrs 19 and 21 in DNG, and chr36 in A_WOLF.

These data suggest that "hot" LINE-1s have contributed to multiple new insertions in

recent canine evolution.

Utilizing dimorphic LINE-1 insertions to recreate canine phylogeny

LINE-1s present in the 2,048,395,076 autosomal bp that were callable in all

genomes were categorized based on presence/absence pattens across the dataset. All

samples were included in this analysis, as well as an inferred "ancestral" state which

possesses an empty site at all dimorphic LINE-1 loci. In total, 7,428 dimorphic LINE-1

loci were identified with 4,034 (54.3%) being present in only a single sample. (Fig. S12,

Fig. S13). A neighbor joining tree was inferred using dimorphic LINE-1s and rooted on

the "ancestral" state. The tree possessed the same topology as the SINEC-based tree.

Again, all nodes had 100% bootstrap support aside from the GDN-BOX node which had

80% bootstrap support. (Fig. S13). Overall, 65.6% of loci have genotype profiles that fit

the inferred tree.

Estimating the rate of LINE-1 retrotransposition in canines

Using the SNV mutation and divergence estimates discussed in the SINEC

section, we estimated the rate of LINE-1 retrotransposition in canines. We used the

same average genome divergence as used in the SINEC analysis. (Fig. 7B). We

calculated the retrotransposition rate of each sample relative to G_WOLF and of

G_WOLF relative to GSD1, to avoid counting G_WOLF multiple times. The average

rate of LINE-1 retrotransposition in our dataset was 1/130.2 live births. DNG had the

highest rate of retrotransposition with 1/114.9 live births, while there appears to be a

reduction of the rate of LINE-1 retrotransposition in A_WOLF relative to the other

samples (Fig. 9A, Fig. S14). To generate a retrotransposition rate using only high

confidence LINE-1 insertions, a restricted estimate using variants that possess a TSD of

10 bp or longer was produced. Use of the high-confidence set of LINE-1s reduces the

estimate of the rate of retrotransposition to 1/183.9 births. As in the SINEC analysis, the

upper and lower bounds of the SNV mutation rate were used to calculate upper and

lower bounds of the stringent LINE-1 retrotransposition rate as 1/116.5 and 1/318.2

births, respectively (Fig. 9B, Supplemental Data S6).

In summary, we utilize inter-reference comparisons to identify dimorphic SINEC

and LINE-1 insertions in canines. We interrogated an average of 25,558 SINECs and

3,497 LINE-1s in each sample for the hallmarks of LINE-1 retrotransposition. Most

variants possessed at least one hallmark, with SINECs being overall higher quality than

LINE-1s. Dimorphic SINECs and LINE-1s were aggregated and used to generate

neighbor joining tree which recapitulated canine phylogeny. In this analysis,

approximately half of loci are present in only a single sample (46.3% of SINECs and

54.3% of LINE-1s). LINE-1s were queried for the presence of 3' transductions, which

can be used to infer which elements have been active in recent evolutionary history.

The average sample possesses ~36 LINE-1s with transduced sequences that can be

mapped to G_WOLF coordinates. Finally, we estimate the rate of SINEC and LINE-1

retrotransposition in canines at 1/22 and 1/184 births respectively.

Discussion

In this study, we compare haploid genome assemblies from four breed dogs, a

Dingo, and two wolves to estimate the rate of SINEC and LINE-1 retrotransposition in

canines. We estimate a new high confidence LINE-1 and SINEC arising in one out of

184 and 22 live births, respectively. Several factors may bias our estimate. The first is

that each reference assembly is a haploid representation of a diploid genome. As a

result, these calculations implicitly assume that for heterozygous insertions the filled site

and empty site are represented at an equal fraction across the dataset. We note that

this assumption is violated in the BOX reference genome, which showed a bias against

filled SINEC sites that is not recapitulated for LINE-1 sites (73). The analyzed genomes

were generated using a variety of methods for assembling, polishing, gap filling, and

scaffolding. BOX was the only reference genome created using the program WTDBG2

and polished with WTDPOA-CNS, which may have influenced the differential inclusion

of SINEC variants (113). However, we note that the inferred insertion rate relative to the

G_WOLF outgroup is consistent across comparisons, except for SINEC variants in the

BOX genome as previously noted. Additionally, the level of within-breed SINEC

variation is consistent among German Shepherd Dogs, Labrador Retrievers, and

Bernese Mountain Dogs (Fig. S6), a comparison that includes genomes sequenced

using Oxford Nanopore Technologies (ONT) long reads. Although this suggests that

biased representation of heterozygous sites is unlikely to majorly skew our findings,

future studies that combine phase-resolved diploid assemblies with pangenome

analysis approaches may offer a more comprehensive view of canine genome variation.

Although the bulk of our call set contains true insertion variants, some variants

that arose through other processes, such as deletions of existing elements, may be

included. To ensure that identified variants are indeed LINE-1 mediated, insertions were

queried for the hallmarks of LINE-1 retrotransposition. These include the presence of

target-site duplications, poly(A) tails, and a LINE-1 endonuclease cleavage site. SINECs

overall had a higher fraction of high confidence TSDs than LINE-1s, 83.0% and 71.0%,

respectively. There are several reasons why the fraction of variants possessing

hallmarks is a lower bound estimate. First, when detecting poly(A) tracts, we required a

true homopolymer. Thus, even a single mutation could take a 17 bp homopolymer and

split it into two shorter homopolymers; neither of which would qualify as "high

confidence". Similarly, for TSDs, excess mutations can degrade the target site

duplication such that the left boundary, right boundary, and empty site are no longer

identical. While the AGE aligner does occasionally allow for mismatches among these

three sequences, excess differences will prevent the detection of TSDs in this dataset.

Additionally, AGE identified a fraction of loci as possessing apparent target site

deletions which can arise which can arise during TPRT (114, 115), arise from

polymorphism at the insertion site, or be assembly errors (Fig. S15). For loci present in

GSD1 and absent in G_WOLF, over 80% of loci with no TSD detected by AGE also

possess a target site deletion of at least one base.

We note that it may seem unusual that only ~51% of SINECs and ~44% of LINE-

1s possess both a TSD and a poly(A) tail. We further investigated if loci that do not

possess "high confidence" TSDs and/or poly(A) tails have evidence for the presence of

lower confidence hallmarks. To do so, we reanalyzed the variants using less stringent

parameters. Because TSDs are frequently described as being 7-20 bp on average, we

reduced the minimum TSD size to 7 bp (116). We also looked for evidence of a

degraded poly(A) tail by identifying 15 bp windows that contain at least 10 As within 30

bp of a TSD, or the end of the variant if no TSD of at least 7 bp is detected. Application

of these relaxed criteria have a dramatic effect on hallmark classification. For SINEC

variants, 7,629 (52.5%) variants present in GSD1 and absent in G_WOLF possessed

both hallmarks when applying our stringent criteria while 12,625 (~86.8%) loci had

evidence of both hallmarks when applying the relaxed criteria described above, with

only 1.3% having neither hallmark. Similar results were observed when reanalyzing

LINE-1s present in GSD1 and absent in G_WOLF, with the number of loci possessing

evidence of both hallmarks increasing from 866 (44.1%) to 1,404 (71.6%) while the

number of loci with neither hallmark was reduced from 260 (13.3%) to 49 (2.5%).

Overall, these data suggest that the high stringency filters used for our refined rate

estimate are conservative. Other factors may also contribute to the deficit of dimorphic

LINE-1s with identified poly(A) tails. For example, we did not systematically process

sequences with internal rearrangements. As a result, LINE-1s that integrated via the

twin priming mechanism are included in our data set but were not processed for poly(A)

tail identification (47).

While only an average of 1.8% of loci possessed a mappable transduction in our

dataset, this is in line with previous canine estimates. For instance, comparisons

between the GDN reference genome and Canfam3.1, which was derived from a boxer,

identified 1,121 dimorphic LINE-1s present in one genome or the other. Of them, only

18 (~1.6%) possessed identifiable 3' transductions (20).Both analyses yield a much

lower proportion of 3' transductions than identified in human genomes (55, 112). One

source of this difference may be LINE-1 3' UTR sequence divergence between species.

We noted several differences between the 3' UTRs of L1_Cf and L1HS present in

RepeatMasker library version dc20170127-rb20170127. Interestingly, the L1HS 3' UTR

possesses a single instance of the "AAUAAA" poly(A) terminator sequence located at

the start of the 3' poly(A) (117). By contrast, the L1_Cf sequence possesses two

"AAUAAA" segments located in close proximity at the 3' end of the element. The

presence of two terminator sequences may contribute to the reduced rate of L1_Cf 3'

transductions.

It is important to note that LINE-1 sequences underwent several additional filters

relative to SINECs. This is because a LINE-1 specific class of false positives was

identified in our dataset. Specifically, if an existing LINE-1 in a genome undergoes a

deletion of internal LINE-1 sequence, the assembly comparison method may incorrectly

report a false LINE-1 insertion in other assemblies which do not possess the deletion.

Thus, we removed any LINE-1 which exists within larger LINE-1 sequences and does

not possess a high confidence TSD. While the removal of these variants removed many

low-quality loci, remaining variants were not of uniform quality. To reflect this

uncertainty, we present estimates for all identified sites as well as only for those sites

with an identified TSD.

Unsurprisingly, manual review suggests that some false positives remain in our

dataset. For example, a locus found in multiple dogs but absent from G_WOLF

(G_WOLF chr6:55577098) likely formed through a mechanism other than

retrotransposition. This locus involves sequence from a L1MB4 element, a LINE-1

subfamily that was active ~150 million years ago (118). The variable sequence is

flanked by large segments of matching sequence (130 bp in length) and does not have

an inferred cut site that matches the LINE-1 endonuclease consensus sequence.

Structural variants such as this locus on chr6 may have formed through homology-

mediated processes involving LINE-1-derived sequences and may contribute to the loci

with exceptionally long inferred TSDs.

Further analysis shows that simply filtering variants based on the inferred

element subfamily is not a robust strategy. RepeatMasker classifies elements based on

sequence identity relative to a collection of subfamily consensus sequences and

attempts to create a biologically meaningful family assignment from fragmented

alignments. For LINE-1s, alignments are processed relative to consensus sequences

from different parts of the element (e.g., the 5' end, the ORF2p region, etc.). The

subfamily assignment process may misclassify elements that are fragmented, such as

occurs when a mobile element has inserted inside of an existing element (119).

Furthermore, RepeatMasker calculates sequence divergence using a set of scoring

matrices that vary based on the GC content of the analyzed region. As a result,

disparate subfamily assignments may be reported when a sequence is analyzed

individually rather than as part of its encompassing chromosomal context. Since the

RepeatMasker assignment is based on sequence identity and is not weighted by the

presence of specific changes that are diagnostic of subfamily membership,

misassignment may occur more frequently when alignments encompass segments with

few inter-subfamily differences. The subfamily assignments reported in Additional_File1

were derived by intersecting the variant coordinates with the RepeatMasker annotations

of each genome assembly and should be interpreted with caution. A more detailed

analysis of canine LINE-1 and SINEC subfamilies, informed by presence-absence

patterns found among relevant genome assemblies representing a range of

evolutionary distances, may be a fruitful line of future research. Such a detailed

subfamily description could then be used to guide the principled assignment of elements

into subfamilies, as done by the C*Alu* and LINEu algorithms (120).

In addition to the technical limitations described above, our calculation of the rate

of retrotransposition is based on several assumptions. First, the estimate assumes that

the rate of LINE-1, SINEC, and SNV mutagenesis has remained constant over the last

~750,000 years of canine evolution. Violation of this assumption could cause an over- or

underestimate of the rate of retrotransposition in canines. Additionally, our phylogenetic

estimate assumes that insertions are not subject to genetic selection. Because it is far

more likely that new insertions would be deleterious rather than advantageous, the

calculated value may underestimate the true rate of *de novo* LINE-1 or SINEC insertion.

The number of generations since divergence is an estimate of the time to

genome coalescence between each sample and G_WOLF. This estimate relies upon

the number of SNVs identified in each assembly. All assemblies are PacBio derived and

the GSD2 and DNG assemblies are supplemented with Oxford Nanopore Technologies

(ONT) long reads. Each assembly was also polished using at least one short-read

sequencing modality. Each of these differences can affect the base quality throughout

the reference assemblies and can propagate further to bias our estimated rates of

retrotransposition. It is worth noting, however, that false SNVs would lead to an

underestimation of the rate of retrotransposition by artificially increasing the number of

generations since divergence and our rate estimates are broadly consistent across

samples.

A key parameter in our estimate is the rate of SNV mutation in canines, which we

set at 4.5×10^{-9}/bp/generation. Using this estimate, the average sample is 229,004

generations diverged from G_WOLF, which corresponds to ~750,000 years of

divergence. We note that this number reflects the average time of lineage coalescence

across the genome and, due to the large effective population size found in ancestral

wolf populations, is substantially older than the estimated dog-wolf population split time

of 10,000-40,000 years ago (121). As described in our previous analysis of retrogenes

in canines, this difference is consistent with estimates of wolf demographic history (80).

After correcting for differences in the assumed mutation rate, a previous analysis

estimated that the ancestors of new-world and old-world wolves had an effective size

(Ne) of 143,000 (121). This large size corresponds to an expected time to coalescence

of two lineages in the ancestral population of 286,000 generations. After accounting for

the additional divergence following population separation, this expectation is broadly

consistent with the divergence time we inferred from genome assembly comparisons.

The largest source of error in our estimates of the LINE-1 and SINEC

retrotransposition rates is the assumed SNV mutation rate. The value we used was

derived from Illumina sequencing of a wolf pedigree with four offspring. The Koch et al.

point estimate has a large confidence interval (2.6×10^{-9}-7.1×10^{-9}) and is broadly

consistent with the rate inferred from analysis of an ancient wolf genome (2, 101).

Recently, Zhang et al. estimated a SNV mutation rate of 4.89×10^{-9} per bp per generation

based on Illumina sequencing of 404 dog trios (122). The authors noted that the dog

SNV mutation rate is correlated with paternal age and may also show differences

across breeds.

We utilized a phylogenetic approach to estimate the rate of canine LINE-1 and

SINEC retrotransposition. This method is based on the total number of insertions that

occurred in two samples and is calibrated by a known SNV mutation rate. Other

approaches to estimate the rate of retrotransposition have been employed in other

species. These include direct estimates obtained through pedigree sequencing and

population modeling approaches based on the number of insertions found in a

population (i.e., Watterson's estimator (123)). Population modelling approaches require

an estimate of the effective population size of the studied group, a value that can be

sensitive to modelling assumptions and is itself often calibrated using a SNV mutation

rate. Although conceptually simple, direct pedigree estimates are sensitive to filtering

strategies and are influenced by both false positive calls in the offspring and false

negative calls in the parents.

To place our estimates of the rate of retrotransposition in canines in a broader

context, we compared them to retrotransposition rates estimated in humans (Table 2).

We recalibrated published rate estimates to use the smaller human SNV mutation rate

implied by pedigree sequencing studies. After recalibration, the estimated rate of *Alu*

retrotransposition in humans (~1/36 – ~1/40 births) is approximately half of the SINEC

rate we estimate for canines. The human LINE-1 insertion rate estimated by

phylogenetic or population modelling approaches (~1/210 - ~1/301 births) is slightly

smaller than our estimate of the rate of LINE-1 insertions in dogs. However, the

estimate obtained from human pedigrees (~1/63) is markedly higher. The extent to

which technical factors (i.e., false positives and false negatives) and biological factors

(i.e., strong negative selection against new LINE-1 insertions) contribute to differences

in the estimated rates is unclear. The modest differences in retrotransposition rates

estimated for dogs and humans suggest that the striking levels of canine SINEC and

LINE-1 dimorphism reflect long-standing genetic variation that has been segregating in

canines throughout their history.

Author and Year	Method	Element Type	Published Rate Estimate	Rescaled Rate Estimate
Xing et al., 2009	Phylogeny	LINE-1	1/212	1/301
Feusier et al., 2019	Pedigree	LINE-1	1/63	1/63
Ewing & Kazazian, 2010	Population Modeling	LINE-1	1/140	1/210
Xing et al., 2009	Phylogeny	Alu	1/21	1/36
Feusier et al.,	Pedigree	Alu	1/40	1/40

815 **Table 2: Recalibrated estimates of LINE-1 and *Alu* retrotransposition in humans.**

816 Published rates were obtained directly from the indicated manuscripts. Rescaled rates

817 were calculated based on a human mutation rate of 1.3×10^{-8} mutations per bp per

818 generation and an N_e of 15,000.

Conclusions

By comparing seven canine genome assemblies we identified a total of 7,428 dimorphic

LINE-1s and 51,572 dimorphic SINECs. Insertion allele sharing profiles reflect known

sample relationships and reveal substantial within-breed variation. Calibrating estimates

using a previously estimated single nucleotide mutation rate, we estimate that the rate

of SINEC-1 and LINE-1 and retrotransposition is 1/22 and 1/184 births over recent

canine evolution. The rate of SINEC insertions in canines is approximately two times

larger than the rate estimated for the *Alu* element in humans, while the estimated canine

LINE-1 insertion rates are comparable, suggesting that a large pool of segregating

insertions have been maintained throughout canine evolution.

Funding

Competing Interests

J.V.M. is an inventor on patent US6150160, was a paid consultant for Gilead Sciences,

serves on the scientific advisory board of Tessera Therapeutics Inc. (where he is paid

as a consultant and has equity options), and has licensed reagents to Merck

Pharmaceutical. He also previously served on the American Society of Human Genetics

Board of Directors. The other authors do not declare competing interests.

Author Contributions

M.S.B., J.V.M. and J.M.K. designed the study and interpreted the results, M.S.B. and

A.K.N performed analyses, M.S.B. and J.M.K drafted the manuscript. M.S.B., A.K.N.,

J.V.M. and J.M.K. edited the manuscript.

Data Availability

Final outputs from locus processing including variant location, hallmark information, and

element subtype are located in additional files. Relevant code for this manuscript is

available at https://github.com/mblacksmith1996/Canine_LINE1_SINEC_Scripts.

Acknowledgements

We thank the authors of multiple canine genome assemblies for making their data

available to the community. We also thank Emily Koch and Ryan E. Mills for helpful

discussion and comments on previous versions of this study.

Figures

854
Fig. 1: Genome assembly comparison pipeline.

The flowchart depicts how single nucleotide and structural variants were identified from

canine genome assembly comparisons. Each genome assembly is aligned to the

G_WOLF assembly with minimap2 and variants identified from the resulting alignments

with paftools.js. SNVs are extracted, filtered, and combined with the fraction of the

autosomal genome aligned to estimate the number of generations since genome

divergence of the two samples. SVs are extracted, filtered, and split into insertions and

deletions and intersected with RepeatMasker annotations to identify candidate LINE-1

and SINEC loci. The SV counts and genome divergence estimates are then compared

to calculate an estimated rate of LINE-1 and SINEC insertions per generation. LINE-1

and SINEC containing loci were evaluated for the presence of the hallmarks of

retrotransposition allowing for a stricter estimate of the rate of LINE-1 and SINEC

retrotransposition. Figure created in Lucid (lucid.co).

868
Fig. 2: Visualizing a dimorphic SINEC locus.

(**A**) A miropeats image shows a dimorphic SINEC locus where the empty site is in

GSD1 (top) and the filled site is found in G_WOLF (bottom). Segments present in both

GSD1 and G_WOLF are represented as solid black rectangles connected by curved

lines. The red lines depict the identified breakpoints for a structural variant present in

G_WOLF but absent in GSD1. Repetitive sequence identified by RepeatMasker is

depicted as colored arrows, corresponding to different repeat types as indicated. The

variable sequence corresponds to a SINEC in the minus strand orientation as

highlighted by the left-facing purple arrow. (**B**) Sequences corresponding to filled and

empty site at this locus are shown. Target site duplications are shown in yellow in both

the filled and empty site. SINEC sequence is show in purple, poly(A) sequence is in

blue, and the inferred LINE-1 endonuclease cleavage site is underlined (5' TTTTT/AA).

The underlined sequence corresponds to the reverse-complement (top-strand) of the

cleaved sequenced.

Fig. 3: Inter-genome alignments reveal over 20,000 dimorphic SINECs per sample.

The bar charts depict the number of dimorphic SINECs identified in each assembly

comparison located on the autosomes (A) and on chrX (B). In each chart, bars

represent variants present in G_WOLF, while green bars represent variants present in

the sample.

Fig. 4: Dimorphic SINECs possess the hallmarks of retrotransposition.

The hallmarks of retrotransposition were investigated in SINECs present in GSD1 but

not G_WOLF (panels A-C) and G_WOLF but not GSD1 (panels D-F) according to the

GSD1-G_Wolf alignment. Histograms depicting the lengths of identified TSDs and

poly(A) tails reveal that most SINECs possess predicted structural hallmarks. A red line

depicts the cutoff of 10 bp for high confidence TSDs and 3' poly(A) tracts (panels A, B,

D, and E). Logo plots depict that the loci which possess a TSD of at least 10 bp in

length tend to possess the canonical LINE-1 EN cleavage site. The x-axis represents

the position within the motif, and the dotted vertical line represents the estimated cut

site (panels C and F). Results are shown for variants on the autosomes and chrX.

Fig. 5: Dimorphic SINECs possess the hallmarks of retrotransposition across

samples.

All SINEs were categorized as possessing TSDs (>= 10 bp), poly(A) tails (>= 10 bp),

both, neither, or having ambiguous RepeatMasker annotations. Stacked bar plots show

the distribution of categories across the dataset. The first 6 bars depict variants present

in the indicated sample but absent in G_WOLF; the final bar depicts variants present in

G_WOLF but absent in GSD1. Loci were only considered "Both" if the poly(A) was not

separated by more than 5 bp from the TSD. Results are shown for variants on the

autosomes or chrX.

Fig 6: A phylogenetic tree of canine assemblies constructed from dimorphic

SINECs.

A phylogenetic tree was estimated using a distance matrix created from dimorphic

SINEC loci. Trees were rooted on a theoretical ancestral genome for which all

dimorphic SINEs are absent. The number of dimorphic SINEC variants is depicted on

each branch.

918
919 Fig. 7: SINECs arise at a rate of ~1/20 births in canines.

Histograms visualize the number of autosomal SNVs between each sample and

G_WOLF (panel A), as well as the number of generations since divergence using a

point estimate of 4.5×10^{-9} bp/generation (panel B). The rate of retrotransposition of

SINEC was calculated based on all autosomal dimorphic SINEC loci present in each

sample and absent in G_WOLF (panel C), and a more stringent dataset requiring a TSD

of at least 10 bp each (panel D). For panels C and D, the G_WOLF bar represents

variants present in G_WOLF and absent in GSD1. Thus, the corresponding SNVs and

generations since divergence can be found in the GSD1 column. A red dotted line

shows the average rate of retrotransposition across the dataset. Error bars depict

divergence and SINEC retrotransposition rates calculated based on the reported SNP

mutation rate confidence interval of 2.6×10^{-9}/bp/generation to 7.1×10^{-9}/bp/generation.

932

Fig. 8: Forty percent of dimorphic LINE-1 loci possess a high confidence TSD and

3' poly(A) tail.

As with SINEs, LINE-1s were categorized as possessing TSDs (>= 10bp), poly(A) tails

(>= 10 bp), both, neither, or having ambiguous RepeatMasker orientations. Stacked bar

plots show the distribution of categories across the dataset. The first six bars depict

variants present in the indicated sample but absent in G_WOLF; the final bar depicts

variants present in G_WOLF but absent in GSD1. Loci were only considered "Both" if

the poly(A) was not separated by more than 5 bp from the TSD. Results are shown for

variants on the autosomes or chrX.

Fig. 9: LINE-1s arise at a rate of ~1/20 births in canines.

Similar to Fig. 7, the rate of retrotransposition of LINE-1 was calculated based on all

autosomal dimorphic LINE-1 loci present in each sample and absent in G_WOLF (panel

A), and a more stringent dataset requiring a TSD of at least 10 bp each (panel B).

G_WOLF bar represents variants present in G_WOLF and absent in GSD1. A red

dotted line shows the average rate of retrotransposition across the dataset. Error bars

depict divergence and SINEC retrotransposition rates calculated based on the reported

SNP mutation rate confidence interval of 2.6×10^{-9}/bp/generation to $7.1 \times 10^{-}$

^{9}/bp/generation.

References

1.   Pang JF, Kluetsch C, Zou XJ, Zhang AB, Luo LY, Angleby H, et al. mtDNA data indicate a single origin for dogs south of Yangtze River, less than 16,300 years ago, from numerous wolves. Mol Biol Evol. 2009;26(12):2849-64.

2.   Skoglund P, Ersmark E, Palkopoulou E, Dalen L. Ancient wolf genome reveals an early divergence of domestic dog ancestors and admixture into high-latitude breeds. Curr Biol. 2015;25(11):1515-9.

3.   Botigue LR, Song S, Scheu A, Gopalan S, Pendleton AL, Oetjens M, et al. Ancient European dog genomes reveal continuity since the Early Neolithic. Nat Commun. 2017;8:16082.

4.   Skoglund P, Götherström A, Jakobsson M. Estimation of population divergence times from non-overlapping genomic sequences: examples from dogs and wolves. Mol Biol Evol. 2011;28(4):1505-17.

5.   Schlamp F, van der Made J, Stambler R, Chesebrough L, Boyko AR, Messer PW. Evaluating the performance of selection scans to detect selective sweeps in domestic dogs. Molecular ecology. 2016;25(1):342-56.

6.   Boyko AR. The domestic dog: man's best friend in the genomic era. Genome Biol. 2011;12(2):216.

7.   Parker HG, Shearin AL, Ostrander EA. Man's best friend becomes biology's best in show: genome analyses in the domestic dog. Annu Rev Genet. 2010;44:309-36.

8.   Shearin AL, Ostrander EA. Canine morphology: hunting for genes and tracking mutations. PLoS Biol. 2010;8(3):e1000310.

9.   Ostrander EA, Wayne RK, Freedman AH, Davis BW. Demographic history, selection and functional diversity of the canine genome. Nat Rev Genet. 2017;18(12):705-20.

10.   Sutter NB, Ostrander EA. Dog star rising: the canine genetic system. Nat Rev Genet. 2004;5(12):900-10.

11.   Wayne RK, Ostrander EA. Origin, genetic diversity, and genome structure of the domestic dog. Bioessays. 1999;21(3):247-57.

12.   Marsden CD, Ortega-Del Vecchyo D, O'Brien DP, Taylor JF, Ramirez O, Vila C, et al. Bottlenecks and selective sweeps during domestication have increased deleterious genetic variation in dogs. Proceedings Of The National Academy Of Sciences Of The United States Of America. 2016;113(1):152-7.

13.   Boyko AR, Quignon P, Li L, Schoenebeck JJ, Degenhardt JD, Lohmueller KE, et al. A simple genetic architecture underlies morphological variation in dogs. PLoS Biol. 2010;8(8):e1000451.

14.   Karlsson EK, Lindblad-Toh K. Leader of the pack: gene mapping in dogs and other model organisms. Nat Rev Genet. 2008;9(9):713-25.

15.   Hoeppner MP, Lundquist A, Pirun M, Meadows JR, Zamani N, Johnson J, et al. An improved canine genome and a comprehensive catalogue of coding genes and non-coding transcripts. PLoS One. 2014;9(3):e91172.

16.   Meadows JRS, Kidd JM, Wang GD, Parker HG, Schall PZ, Bianchi M, et al. Genome sequencing of 2000 canids by the Dog10K consortium advances the understanding of demography, genome function and architecture. Genome Biol. 2023;24(1):187.

17.   Richardson SR, Doucet AJ, Kopera HC, Moldovan JB, Garcia-Perez JL, Moran JV. The
Influence of LINE-1 and SINE Retrotransposons on Mammalian Genomes. Microbiol Spectr.
2015;3(2):MDNA3-0061-2014.
18.   Lindblad-Toh K, Wade CM, Mikkelsen TS, Karlsson EK, Jaffe DB, Kamal M, et al. Genome
sequence, comparative analysis and haplotype structure of the domestic dog. Nature.
2005;438(7069):803-19.
19.   Wang W, Kirkness EF. Short interspersed elements (SINEs) are a major source of canine
genomic diversity. Genome Res. 2005;15(12):1798-808.
20.   Halo JV, Pendleton AL, Shen F, Doucet AJ, Derrien T, Hitte C, et al. Long-read assembly of
a Great Dane genome highlights the contribution of GC-rich sequence and mobile elements to
canine genomes. Proceedings Of The National Academy Of Sciences Of The United States Of
America. 2021;118(11).
21.   Moran JV, Holmes SE, Naas TP, DeBerardinis RJ, Boeke JD, Kazazian HH. High frequency
retrotransposition in cultured mammalian cells. Cell. 1996;87(5):917-27.
22.   Dewannieux M, Esnault C, Heidmann T. LINE-mediated retrotransposition of marked Alu
sequences. Nat Genet. 2003;35(1):41-8.
23.   Dombroski BA, Mathias SL, Nanthakumar E, Scott AF, Kazazian HH. Isolation of an active
human transposable element. Science. 1991;254(5039):1805-8.
24.   Scott AF, Schmeckpeper BJ, Abdelrazik M, Comey CT, O'Hara B, Rossiter JP, et al. Origin
of the human L1 elements: proposed progenitor genes deduced from a consensus DNA
sequence. Genomics. 1987;1(2):113-25.
25.   Khazina E, Weichenrieder O. Non-LTR retrotransposons encode noncanonical RRM
domains in their first open reading frame. Proc Natl Acad Sci U S A. 2009;106(3):731-6.
26.   Kolosha VO, Martin SL. High-affinity, non-sequence-specific RNA binding by the open
reading frame 1 (ORF1) protein from long interspersed nuclear element 1 (LINE-1). J Biol Chem.
2003;278(10):8112-7.
27.   Martin SL, Bushman FD. Nucleic acid chaperone activity of the ORF1 protein from the
mouse LINE-1 retrotransposon. Mol Cell Biol. 2001;21(2):467-75.
28.   Hohjoh H, Singer MF. Cytoplasmic ribonucleoprotein complexes containing human LINE-
1 protein and RNA. EMBO J. 1996;15(3):630-9.
29.   Feng Q, Moran JV, Kazazian HH, Boeke JD. Human L1 retrotransposon encodes a
conserved endonuclease required for retrotransposition. Cell. 1996;87(5):905-16.
30.   Mathias SL, Scott AF, Kazazian HH, Boeke JD, Gabriel A. Reverse transcriptase encoded
by a human transposable element. Science. 1991;254(5039):1808-10.
31.   Swergold GD. Identification, characterization, and cell specificity of a human LINE-1
promoter. Mol Cell Biol. 1990;10(12):6718-29.
32.   Athanikar JN, Badge RM, Moran JV. A YY1-binding site is required for accurate human
LINE-1 transcription initiation. Nucleic Acids Res. 2004;32(13):3846-55.
33.   Holmes SE, Dombroski BA, Krebs CM, Boehm CD, Kazazian HH. A new retrotransposable
human L1 element from the LRE2 locus on chromosome 1q produces a chimaeric insertion. Nat
Genet. 1994;7(2):143-8.
34.   Belancio VP, Whelton M, Deininger P. Requirements for polyadenylation at the 3' end of
LINE-1 elements. Gene. 2007;390(1-2):98-107.

35.   Doucet AJ, Wilusz JE, Miyoshi T, Liu Y, Moran JV. A 3' Poly(A) Tract Is Required for LINE-1
Retrotransposition. Molecular cell. 2015;60(5):728-41.
36.   Alisch RS, Garcia-Perez JL, Muotri AR, Gage FH, Moran JV. Unconventional translation of
mammalian LINE-1 retrotransposons. Genes Dev. 2006;20(2):210-24.
37.   Martin SL. Ribonucleoprotein particles with LINE-1 RNA in mouse embryonal carcinoma
cells. Mol Cell Biol. 1991;11(9):4804-7.
38.   Boeke JD. LINEs and Alus--the polyA connection. Nat Genet. 1997;16(1):6-7.
39.   Kulpa DA, Moran JV. Cis-preferential LINE-1 reverse transcriptase activity in
ribonucleoprotein particles. Nat Struct Mol Biol. 2006;13(7):655-60.
40.   Luan DD, Korman MH, Jakubczak JL, Eickbush TH. Reverse transcription of R2Bm RNA is
primed by a nick at the chromosomal target site: a mechanism for non-LTR retrotransposition.
Cell. 1993;72(4):595-605.
41.   Wei W, Gilbert N, Ooi SL, Lawler JF, Ostertag EM, Kazazian HH, et al. Human L1
retrotransposition: cis preference versus trans complementation. Mol Cell Biol.
2001;21(4):1429-39.
42.   Kubo S, Seleme MC, Soifer HS, Perez JL, Moran JV, Kazazian HH, et al. L1
retrotransposition in nondividing and primary human somatic cells. Proc Natl Acad Sci U S A.
2006;103(21):8036-41.
43.   Jurka J. Sequence patterns indicate an enzymatic involvement in integration of
mammalian retroposons. Proceedings Of The National Academy Of Sciences Of The United
States Of America. 1997;94(5):1872-7.
44.   Flasch DA, Macia Á, Sánchez L, Ljungman M, Heras SR, García-Pérez JL, et al. Genome-
wide de novo L1 Retrotransposition Connects Endonuclease Activity with Replication. Cell.
2019;177(4):837-51.e28.
45.   Cost GJ, Feng Q, Jacquier A, Boeke JD. Human L1 element target-primed reverse
transcription in vitro. EMBO J. 2002;21(21):5899-910.
46.   Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, et al. Initial sequencing
and analysis of the human genome. Nature. 2001;409(6822):860-921.
47.   Ostertag EM, Kazazian HH. Twin priming: a proposed mechanism for the creation of
inversions in L1 retrotransposition. Genome Res. 2001;11(12):2059-65.
48.   Grimaldi G, Singer MF. Members of the KpnI family of long interspersed repeated
sequences join and interrupt alpha-satellite in the monkey genome. Nucleic Acids Res.
1983;11(2):321-38.
49.   Beck CR, Garcia-Perez JL, Badge RM, Moran JV. LINE-1 elements in structural variation
and disease. Annu Rev Genomics Hum Genet. 2011;12:187-215.
50.   Van Arsdell SW, Denison RA, Bernstein LB, Weiner AM, Manser T, Gesteland RF. Direct
repeats flank three small nuclear RNA pseudogenes in the human genome. Cell. 1981;26(1 Pt
1):11-7.
51.   Moran JV. Human L1 retrotransposition: insights and peculiarities learned from a
cultured cell retrotransposition assay. Genetica. 1999;107(1-3):39-51.
52.   Moran JV, DeBerardinis RJ, Kazazian HH. Exon shuffling by L1 retrotransposition.
Science. 1999;283(5407):1530-4.
53.   Szak ST, Pickeral OK, Landsman D, Boeke JD. Identifying related L1 retrotransposons by
analyzing 3' transduced sequences. Genome Biol. 2003;4(5):R30.

54.   Beck CR, Collier P, Macfarlane C, Malig M, Kidd JM, Eichler EE, et al. LINE-1
retrotransposition activity in human genomes. Cell. 2010;141(7):1159-70.
55.   Goodier JL, Ostertag EM, Kazazian HH. Transduction of 3'-flanking sequences is common
in L1 retrotransposition. Hum Mol Genet. 2000;9(4):653-7.
56.   Macfarlane CM, Collier P, Rahbari R, Beck CR, Wagstaff JF, Igoe S, et al. Transduction-
specific ATLAS reveals a cohort of highly active L1 retrotransposons in human populations. Hum
Mutat. 2013;34(7):974-85.
57.   Esnault C, Maestre J, Heidmann T. Human LINE retrotransposons generate processed
pseudogenes. Nat Genet. 2000;24(4):363-7.
58.   Vassetzky NS, Kramerov DA. CAN--a pan-carnivore SINE family. Mamm Genome.
2002;13(1):50-7.
59.   Minnick MF, Stillwell LC, Heineman JM, Stiegler GL. A highly repetitive DNA sequence
possibly unique to canids. Gene. 1992;110(2):235-8.
60.   Bentolila S, Bach JM, Kessler JL, Bordelais I, Cruaud C, Weissenbach J, et al. Analysis of
major repetitive DNA sequences in the dog (Canis familiaris) genome. Mamm Genome.
1999;10(7):699-705.
61.   Kosushkin SA, Ustyantsev IG, Borodulina OR, Vassetzky NS, Kramerov DA. Tail Wags
Dog's SINE: Retropositional Mechanisms of Can SINE Depend on Its A-Tail Structure. Biology
(Basel). 2022;11(10).
62.   Parker HG, VonHoldt BM, Quignon P, Margulies EH, Shao S, Mosher DS, et al. An
expressed fgf4 retrogene is associated with breed-defining chondrodysplasia in domestic dogs.
Science. 2009;325(5943):995-8.
63.   Brown EA, Dickinson PJ, Mansour T, Sturges BK, Aguilar M, Young AE, et al. retrogene on
CFA12 is responsible for chondrodystrophy and intervertebral disc disease in dogs. Proc Natl
Acad Sci U S A. 2017;114(43):11476-81.
64.   Everson R, Pettitt L, Forman OP, Dower-Tylee O, McLaughlin B, Ahonen S, et al. An
intronic LINE-1 insertion in MERTK is strongly associated with retinopathy in Swedish Vallhund
dogs. PLoS One. 2017;12(8):e0183021.
65.   Marchant TW, Johnson EJ, McTeir L, Johnson CI, Gow A, Liuti T, et al. Canine
Brachycephaly Is Associated with a Retrotransposon-Mediated Missplicing of SMOC2. Curr Biol.
2017;27(11):1573-84.e6.
66.   Smith BF, Yue Y, Woods PR, Kornegay JN, Shin JH, Williams RR, et al. An intronic LINE-1
element insertion in the dystrophin gene aborts dystrophin expression and results in Duchenne-
like muscular dystrophy in the corgi breed. Lab Invest. 2011;91(2):216-31.
67.   Wiedmer M, Oevermann A, Borer-Germann SE, Gorgas D, Shelton GD, Drögemüller M,
et al. A RAB3GAP1 SINE Insertion in Alaskan Huskies with Polyneuropathy, Ocular
Abnormalities, and Neuronal Vacuolation (POANV) Resembling Human Warburg Micro
Syndrome 1 (WARBM1). G3 (Bethesda). 2015;6(2):255-62.
68.   Clark LA, Wahl JM, Rees CA, Murphy KE. Retrotransposon insertion in SILV is responsible
for merle patterning of the domestic dog. Proceedings Of The National Academy Of Sciences Of
The United States Of America. 2006;103(5):1376-81.
69.   Murphy SC, Evans JM, Tsai KL, Clark LA. Length variations within the. Mob DNA.
2018;9:26.

70.   Van Poucke M, Ledeganck L, Guo LT, Shelton GD, Bhatti SFM, Cornelis I, et al.
Exonisation of an intronic L1 element in the dystrophin gene associated with X-linked muscular
dystrophy in a Border Collie dog. Anim Genet. 2024;55(5):733-43.
71.   Wang C, Wallerman O, Arendt ML, Sundström E, Karlsson Å, Nordin J, et al. A novel
canine reference genome resolves genomic architecture and uncovers transcript complexity.
Commun Biol. 2021;4(1):185.
72.   Field MA, Rosen BD, Dudchenko O, Chan EKF, Minoche AE, Edwards RJ, et al.
Canfam_GSD: De novo chromosome-length genome assembly of the German Shepherd Dog
(Canis lupus familiaris) using a combination of long reads, optical mapping, and Hi-C.
Gigascience. 2020;9(4).
73.   Jagannathan V, Hitte C, Kidd JM, Masterson P, Murphy TD, Emery S, et al.
Dog10K_Boxer_Tasha_1.0: A Long-Read Assembly of the Dog Reference Genome. Genes
(Basel). 2021;12(6).
74.   Field MA, Yadav S, Dudchenko O, Esvaran M, Rosen BD, Skvortsova K, et al. The
Australian dingo is an early offshoot of modern breed dogs. Sci Adv. 2022;8(16):eabm5944.
75.   Sinding MS, Gopalakrishnan S, Raundrup K, Dalén L, Threlfall J, Gilbert T, et al. The
genome sequence of the grey wolf,. Wellcome Open Res. 2021;6:310.
76.   Bredemeyer KR, vonHoldt BM, Foley NM, Childers IR, Brzeski KE, Murphy WJ. The value
of hybrid genomes: Building two highly contiguous reference genome assemblies to advance
Canis genomic studies. J Hered. 2024;115(4):480-6.
77.   Li H. New strategies to improve minimap2 alignment accuracy. Bioinformatics.
2021;37(23):4572-4.
78.   Iseric H, Alkan C, Hach F, Numanagic I. Fast characterization of segmental duplication
structure in multiple genome assemblies. Algorithms Mol Biol. 2022;17(1):4.
79.   Pendleton AL, Shen F, Taravella AM, Emery S, Veeramah KR, Boyko AR, et al.
Comparison of village dog and wolf genomes highlights the role of the neural crest in dog
domestication. BMC Biol. 2018;16(1):64.
80.   Nguyen AK, Blacksmith MS, Kidd JM. Duplications and Retrogenes Are Numerous and
Widespread in Modern Canine Genomic Assemblies. Genome Biol Evol. 2024;16(7).
81.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics. 2010;26(6):841-2.
82.   Smit A, Hubley R, Green P. *RepeatMasker Open-4.0*. 2013-2015.
83.   Smit AF. Interspersed repeats and other mementos of transposable elements in
mammalian genomes. Curr Opin Genet Dev. 1999;9(6):657-63.
84.   Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years of
SAMtools and BCFtools. Gigascience. 2021;10(2).
85.   Abyzov A, Gerstein M. AGE: defining breakpoints of genomic structural variants at
single-nucleotide resolution, through optimal alignments with gap excision. Bioinformatics.
2011;27(5):595-603.
86.   Tareen A, Kinney JB. Logomaker: beautiful sequence logos in Python. Bioinformatics.
2020;36(7):2272-4.
87.   Harris CR, Millman KJ, van der Walt SJ, Gommers R, Virtanen P, Cournapeau D, et al.
Array programming with NumPy. Nature. 2020;585(7825):357-62.

88.   McKinney W. Data structures for statistical computing in Python. scipy. 2010;445(1):51-6.

89.   Hunter JD. Matplotlib: A 2D graphics environment. Computing in science & engineering. 2007;9(03):90-5.

90.   Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, et al. SciPy 1.0: fundamental algorithms for scientific computing in Python. Nat Methods. 2020;17(3):261-72.

91.   Parsons JD. Miropeats: graphical DNA sequence comparisons. Comput Appl Biosci. 1995;11(6):615-9.

92.   Nothman J. UpSetPlot. 2023.

93.   Tamura K, Stecher G, Kumar S. MEGA11: Molecular Evolutionary Genetics Analysis Version 11. Mol Biol Evol. 2021;38(7):3022-7.

94.   Nei M, Kumar S. Molecular Evolution and Phylogenetics. 1st ed: Oxford University Press; 2000.

95.   Felsenstein J. CONFIDENCE LIMITS ON PHYLOGENIES: AN APPROACH USING THE BOOTSTRAP. Evolution. 1985;39(4):783-91.

96.   Rambaut A. FigTree. 2018.

97.   Schall PZ, Kidd JM. Integrative genotyping and analysis of canine structural variation using long-read and short-read data. bioRxiv. 2025:2025.03.05.641690.

98.   Chen X, Schulz-Trieglaff O, Shaw R, Barnes B, Schlesinger F, Kallberg M, et al. Manta: rapid detection of structural variants and indels for germline and cancer sequencing applications. Bioinformatics. 2016;32(8):1220-2.

99.   Eggertsson HP, Jonsson H, Kristmundsdottir S, Hjartarson E, Kehr B, Masson G, et al. Graphtyper enables population-scale genotyping using pangenome graphs. Nat Genet. 2017;49(11):1654-60.

100.   Benson G. Tandem repeats finder: a program to analyze DNA sequences. Nucleic Acids Res. 1999;27(2):573-80.

101.   Koch EM, Schweizer RM, Schweizer TM, Stahler DR, Smith DW, Wayne RK, et al. De Novo Mutation Rate Estimation in Wolves of Known Pedigree. Mol Biol Evol. 2019;36(11):2536-47.

102.   Morgulis A, Gertz EM, Schaffer AA, Agarwala R. A fast and symmetric DUST implementation to mask low-complexity DNA sequences. J Comput Biol. 2006;13(5):1028-40.

103.   Kent WJ. BLAT--the BLAST-like alignment tool. Genome Res. 2002;12(4):656-64.

104.   Nachman MW, Crowell SL. Estimate of the mutation rate per nucleotide in humans. Genetics. 2000;156(1):297-304.

105.   Sasani TA, Pedersen BS, Gao Z, Baird L, Przeworski M, Jorde LB, et al. Large, three-generation human families reveal post-zygotic mosaicism and variability in germline mutation accumulation. Elife. 2019;8.

106.   Narasimhan VM, Rahbari R, Scally A, Wuster A, Mason D, Xue Y, et al. Estimating the human mutation rate from autozygous segments reveals population differences in human mutational processes. Nat Commun. 2017;8(1):303.

107.   Awadalla P, Gauthier J, Myers RA, Casals F, Hamdan FF, Griffing AR, et al. Direct measure of the de novo mutation rate in autism and schizophrenia cohorts. Am J Hum Genet. 2010;87(3):316-24.

108. Roach JC, Glusman G, Smit AF, Huff CD, Hubley R, Shannon PT, et al. Analysis of genetic
inheritance in a family quartet by whole-genome sequencing. Science. 2010;328(5978):636-9.
109. Mallick S, Li H, Lipson M, Mathieson I, Gymrek M, Racimo F, et al. The Simons Genome
Diversity Project: 300 genomes from 142 diverse populations. Nature. 2016;538(7624):201-6.
110. Pritchard JK. An Owner's Guide to the Human GenomeVersion: September 30, 2023.
111. Bailey JA, Carrel L, Chakravarti A, Eichler EE. Molecular evidence for a relationship
between LINE-1 elements and X chromosome inactivation: the Lyon repeat hypothesis. Proc
Natl Acad Sci U S A. 2000;97(12):6634-9.
112. Pickeral OK, Makalowski W, Boguski MS, Boeke JD. Frequent human genomic DNA
transduction driven by LINE-1 retrotransposition. Genome Res. 2000;10(4):411-5.
113. Ruan J, Li H. Fast and accurate long-read assembly with wtdbg2. Nat Methods.
2020;17(2):155-8.
114. Gilbert N, Lutz-Prigge S, Moran JV. Genomic deletions created upon LINE-1
retrotransposition. Cell. 2002;110(3):315-25.
115. Gilbert N, Lutz S, Morrish TA, Moran JV. Multiple fates of L1 retrotransposition
intermediates in cultured human cells. Mol Cell Biol. 2005;25(17):7780-95.
116. Ostertag EM, Kazazian HH. Biology of mammalian L1 retrotransposons. Annu Rev Genet.
2001;35:501-38.
117. Proudfoot NJ. Ending the message: poly(A) signals then and now. Genes Dev.
2011;25(17):1770-82.
118. Giordano J, Ge Y, Gelfand Y, Abrusan G, Benson G, Warburton PE. Evolutionary history of
mammalian transposons determined by genome-wide defragmentation. PLoS Comput Biol.
2007;3(7):e137.
119. Churakov G, Grundmann N, Kuritzin A, Brosius J, Makalowski W, Schmitz J. A novel web-
based TinT application and the chronology of the Primate Alu retroposon activity. BMC Evol
Biol. 2010;10:376.
120. Gardner EJ, Lam VK, Harris DN, Chuang NT, Scott EC, Pittard WS, et al. The Mobile
Element Locator Tool (MELT): population-scale mobile element discovery and biology. Genome
Res. 2017;27(11):1916-29.
121. Fan Z, Silva P, Gronau I, Wang S, Armero AS, Schweizer RM, et al. Worldwide patterns of
genomic variation and admixture in gray wolves. Genome Res. 2016;26(2):163-73.
122. Zhang SJ, Ma J, Riera M, Besenbacher S, Niskanen JE, Salokorpi N, et al. Determinants of
de novo mutations in extended pedigrees of 43 dog breeds. Genome Biol. 2025;26(1):305.
123. Watterson GA. On the number of segregating sites in genetical models without
recombination. Theor Popul Biol. 1975;7(2):256-76.
1244

```
>GSD1, chr1:92926167-92926198
TTTAAAAATCCCTGTGTTGGACAGCCCCAGTG

>G_WOLF, chr1:30315491-30315735/rc
AAACTGGTTTAAAAATCCCTGTGTTGGACATCCCTCGG
TGGTGCAGCGGTTTGGCGCCTGCCTGTTTGGGCCCGGG
GCGATCCTGGAGGACCCGGGATCAGGTCCACCCATCGGG
CTCGTGGTGGCAGCCGGCCTCCGCGGGCTTCTCCTTGG
GTCTCTCCTCCCCCTCTCTGCTCTCTTCCTGGTGCTATCAT
CTCACAAATAAAAATTTAAAAGGGGGGGTGGGGGGGGGG
TCCCTGTGTTGGCAGCCCCAGT
```