Abnormal shear stress induces ferroptosis in endothelial cells via KLF6 downregulation

Jingang Cui, Zhiyu Fan, Suoqi Ding[¶], Jiazhen Zhang, Huihong Shen, Syeda Armana Zaidi, Yongsheng Ding*

College of Life Sciences, University of Chinese Academy of Sciences, Beijing 101408, China
[¶] Permanent address: School of Chemistry, Beihang University, Beijing, 100191, China
* Corresponding author

Abstract

Stable laminar flow maintains vascular tone regulation, while abnormal blood flow, such as disturbed flow or extreme shear stress, causes endothelial dysfunction, but the underlying mechanism is yet to be explored. We used a microfluidic device to deform flat microchannel into tunnel-like macrochannel. The cross-sectional area of this macrochannel varies with the flow rate and the thickness of the deformable layer, creating three levels of shear stress: low (0.99 dynes/cm², LSS), medium (4.78 dynes/cm², MSS), and high (24 dynes/cm², HSS). Comparing different shear stress exposure to endothelial cells for 24 h, prominent ferroptosis features emerged under either LSS or HSS compared to MSS. These features included increased C11 BODIPY-labeled lipid peroxidation and 4-hydroxynonenal accumulation, CoQ10 depletion, reduced SLC7A11 protein expression, and diminished cell death with Ferrostatin-1 treatment. RNA-seq analysis (LSS/MSS) showed that LSS significantly downregulated transcription of cholesterol homeostasis and unfolded protein response (UPR). Compared to MSS, Western blot results showed that both LSS and HSS reduced the expression of two key enzymes (MVD and IDI1) in the mevalonate pathway, as well as the expression of two main UPR signaling regulators (PERK and BiP). Based on the binding prediction between transcription factors and gene promoters from differentially expressed genes identified through RNA-seq, we found KLF6 to be a key transcription factor. It regulates the PERK-mediated UPR and the mevalonate pathway, which are associated with SLC7A11 expression and CoQ10 synthesis, respectively. The overexpression of KLF6 restored SLC7A11 and CoQ10 levels under both LSS and HSS, significantly reducing foam cell formation, monocyte adhesion, and lipid peroxidation. Our findings reveal KLF6 as a crucial regulator of atherosclerosis induced by abnormal shear stress.

Key words: abnormal shear stress; KLF6; endothelial ferroptosis; unfolded protein response; mevalonate pathway

Introduction

Atherosclerosis is the common pathological basis of a variety of cardiovascular diseases, such as coronary heart disease, myocardial infarction, and cerebral infarction. It is characterized by the accumulation of cholesterol lipid substances in the arterial walls, forming plaques that lead to the narrowing of blood vessels. It is well recognized that the early stage of plaque formation involves endothelial cell (EC) damage, monocyte adhesion, macrophage differentiation, oxidized low-density lipoprotein (oxLDL) phagocytosis, then foam cell formation.

The primary risk factors for atherosclerosis in the circulatory system are well-known: hypertension, hyperlipidemia, and hyperglycemia. A substantial body of evidence from both animal and human specimens suggests that atherosclerosis tends to develop at the bends and branches of arteries [1]. This spatial preference is believed to be linked to changes in blood flow in these regions, marked by low shear stress or oscillatory flow [2-8]. In response to shear stress, several mechanically sensitive transcription factors associated with atherosclerosis have been identified. These include anti-atherosclerotic factors Kruppel-like Factors KLF2 [9-11] and KLF4 [9, 12-14], MEF2[15], and NRF2 [16], as well as pro-atherosclerotic factors AP-1 [17], NF-κB [18], and YAP [19, 20]. These factors play various roles in inflammation, oxidative stress, and cell proliferation. KLF2 and KLF4 have dual functional domains (transcriptional activation and repression), allowing them to regulate genes bidirectionally. They positively regulate endothelial nitric oxide synthase (eNOS) and negatively regulate vascular cell adhesion molecule-1 (VCAM-1). Unlike KLF2 and 4, KLF6 only has a transcriptional activation domain and has also been reported to be involved in the occurrence of atherosclerosis [21, 22]. Recently, azathioprine for psoriasis and apremilast for hypertension have been reported to alleviate endothelial dysfunction and inflammatory responses by restoring KLF6 expression [23, 24].

Endoplasmic reticulum (ER) stress refers to the accumulation of unfolded or misfolded proteins in the ER in response to stimuli, such as oxidative stress, metabolic disorder, and ischemic injury. The unfolded protein response (UPR) is an adaptive cellular mechanism to alleviate ER stress. The accumulation of unfolded or misfolded proteins in the ER causes the 78 kDa glucose-regulated protein (GRP78/BiP) to dissociate from three ER membrane-bound sensors: protein kinase RNA-like ER kinase (PERK), activating transcription factor 6 (ATF6), and inositol-required enzyme 1 (IRE1), thereby initiating three branches of the UPR signaling cascade. To date, various in vitro models have been used to study the impact of shear stress on the ER in endothelial cells. Initially, Fever et al. applied pro-atherosclerotic shear stress using a cone-and-plate flow device, finding that it upregulates BiP through the activation of p38 and integrin $\alpha 2\beta 1$ [26]. A decade later, Bailey et al. used a fan-shaped microfluidic chamber to apply linearly declining shear stress, demonstrating that low shear stress induces VCAM-1 expression and ER stress [27]. More recently, Kim and Woo used a cone system to apply laminar unidirectional flow (up to 12 dynes/cm²), revealing that higher shear stress inhibits ER stress-induced apoptosis via the PI3K/Akt-dependent signaling pathway [28]. Based on these in vitro models, low shear stress at atherosclerosis-susceptible sites seems to trigger ER stress and UPR, fostering inflammation and apoptosis. Conversely, high shear stress protects ECs from ER stress. Moreover, some reviews also highlight in vivo studies that show ER stress in endothelial cells at locations susceptible to atherosclerosis [8, 25]. However, there is still limited understanding of how shear stress influences the molecular mechanisms of ER stress in endothelial cells.

From a physiological point of view, the effect of shear stress should be investigated based on
fluids with physiological pressure. None of the aforementioned in vitro models of fluid shear
stress have been described with a pressure range comparable to physiological blood flow
conditions. Our group has recently developed a microfluidic chip composed of three layers of
polydimethylsiloxane (PDMS). This chip features deformed tunnel-like channels with varying
cross-sectional areas, allowing for multiple combinations of pressure and shear stress [29]. Herein,
we established low, medium, and high shear stress levels at 0.99 (LSS), 4.78 (MSS), and 24 (HSS)
dynes/cm², respectively, while keeping the on-site pressure within the physiological range of
approximately 70 mmHg. After 24 hours under three different shear stress conditions, HUVECs
exposed to both LSS and HSS exhibit significant ferroptosis features compared to those under
MSS. RNA sequencing analysis comparing LSS and MSS revealed enrichment in the inhibition of
steroid biosynthesis and the UPR pathway. Specially, the genes involved include MVD and IDI1,
enzymes in the mevalonate pathway, and BiP and PERK, which are crucial for UPR signaling.
These genes are related to ferroptosis through their effects on CoQ10 production and SLC7A11
expression in ECs. KLF6, rather than KLF2 and KLF4, is a crucial transcription factor for
ferroptosis induced by LSS and HSS. These findings offer new insights into the mechanisms of
shear stress-induced atherosclerosis.
97
Results
LSS and HSS induces ferroptosis in endothelial cells
Blood flow shear stress and pressure are two distinct yet interconnected mechanical forces
affecting the blood vessel wall. The ideal research approach involves changing one variable while
maintaining the others constant, which helps identify each specific effect on the cells. **Figure 1A**
shows the microfluidic chip, emphasizing the different shear stress conditions linked to the
cross-sectional dimensions of the fluid bodies. LSS, MSS, and HSS can be attained by modifying
the thickness of the middle layer, using the given formula and a fixed number of parameters. With
a fixed flow rate of 1.6 ml/min under varying shear stress conditions, the fluid pressure in the
tunnel-like channels fluctuates around 70 mmHg. This is based on the calculation formula we
previously reported, which primarily depends on the flow rate and the size of the downstream
guide channel [29]. **Figure 1B** shows a significantly higher number of propidium iodide
(PI)-stained dead cells under both LSS and HSS compared to MSS. This result aligns with the
change of extracellular LDH level shown in **Figure 1C**, indicating that both LSS and HSS can
lead to significant endothelial cell damage. **Figure 1D-H** highlights key characteristics of
ferroptosis, including lipid peroxidation (marked by 4-hydroxynonenal and C11 BODIPY),
decreased levels of SLC7A11 and CoQ10, and the prevention of cell death with Fer-1 treatment.

Figure 1. Abnormal shear stress induces endothelial ferroptosis. (A) Diagram of the microfluidic
device and calculation of the shear stress. Three levels of shear stress (τ) are formed in the
tunnel-like channel, depending on the varying cross-sectional area of flow: LSS = 0.99 dynes/cm²,
MSS = 4.78 dynes/cm², and HSS = 24 dynes/cm². These values were calculated using a formula
that includes the dynamic viscosity of the fluid (μ), flow rate (Q), channel width (W),
cross-sectional area (A), wetted perimeter (p_w), membrane bulge height (H), and a function of *H/W*
(*g(H/W)*). (B) Fluorescent microscopic results of PI staining in ECs (The scale bar = 250 μm); (C)
ELISA results of extracellular LDH in ECs; (D, E) Flow cytometry results of lipid peroxidation in
ECs indicated by 4-HNE and C11 BODIPY. (F, G) Protein expression of SLC7A11 and
intracellular CoQ10 level. (H) The results of Fer-1 (10 μM) treatment. Data are expressed as the
mean ± standard deviation from three or more independent experiments. Statistical significance

was determined using an unpaired t-test, with $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, and $****p <$
0.0001.
130

RNA-seq analysis shows that LSS inhibits cholesterol biosynthesis and the UPR

RNA-seq analysis was conducted to systematically explore the mechanisms of endothelial cell
damage caused by the abnormal shear stress (LSS/MSS). As shown in **Figure 2A**, LSS
significantly downregulated 298 genes and upregulated 100 genes compared to MSS (│log2│>
0.6, Qvalue < 0.05). KEGG, GO, and GSEA shown in **Figure 2B-D** shows a significant
enrichment in steroid biosynthesis and ER stress response pathways. Specially, LSS significantly
inhibited terpenoid backbone synthesis and UPR. **Figure 2E** indicates that 18 DEGs related to
lipid synthesis were significantly downregulated by LSS. Among them, HMGCS1, MVD, and
IDI1 are three genes that encode key enzymes in the mevalonate pathway. This pathway is
essential for the biosynthesis of sterol isoprenes, such as cholesterol, and non-sterol isoprenes, like
ubiquinone. These enzymes catalyze reactions that convert acetyl-CoA to
(S)-3-hydroxy-3-methylglutaryl-CoA (HMG-CoA), decarboxylate to isopentenyl diphosphate
(IPP), and isomerize between IPP and dimethylallyl diphosphate (DMAPP). These processes are
crucial for CoQ10 synthesis, an antioxidant that helps prevent ferroptosis. **Figure 2F** indicates that
11 DEGs related to the UPR were significantly downregulated by LSS. Genes like SDF2L1,
HSPA5, HSP90B1, HYOU1, and FICD are involved in protein folding, while HERPUD1 and
DERL3 encode proteins for endoplasmic reticulum-associated protein degradation (ERAD).
Additionally, PERK and CHOP play roles in UPR signaling. As LSS inhibits these UPR-related
genes that manage ER stress, the ability to resolve stress from unfolded or misfolded proteins is
likely to be disrupted. To confirm the RNA-seq analysis predictions, we evaluated the protein
expression of four genes—MVD, IDI1, PERK, and HSPA5 (BiP)—under various shear stress
conditions. As shown in **Figure 2G**, Western blot results indicate that the expression of BiP, PERK,
MVD, and IDI1 was significantly reduced by both LSS and HSS compared to MSS. Notably, HSS
has a similar effect on the expression of these proteins as LSS, despite the lack of RNA
sequencing analysis between HSS and MSS. This suggests an adaptive range of organisms to
environmental mechanical stimuli, akin to a 'Window Effect'.

Figure 2. RNA-seq reveals LSS suppresses cholesterol biosynthesis and UPR. (A) Volcano plot of DEGs (LSS vs. MSS;│log2FC│>0.6, Q<0.05); (B) KEGG pathways; (C) GO biological processes; (D) GSEA of steroid biosynthesis and UPR. NES: normalized enrichment score. FDR: false discovery rate. (E) Heatmap of DEGs in lipid synthesis and metabolism. (F) Heatmap of DEGs in UPR. (G) Protein expression of BiP, PERK, MVD, and IDI1 under different shear stress conditions. Data are expressed as the mean ± standard deviation from three or more independent experiments. Statistical significance was determined using an unpaired t-test, with *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and ****$p < 0.0001$. KEGG: Kyoto Encyclopedia of Genes and Genomes, GO: gene ontology, GSEA: gene set enrichment analysis.

KLF6 is found to regulate the PERK-mediated UPR and the mevalonate pathway.

Inhibiting PERK-mediated UPR has been shown to decrease SLC7A11 protein expression in colorectal cancer cells [30]. In our study, both LSS and HSS inhibit this pathway and the mevalonate pathway, impacting SLC7A11 expression and CoQ10 synthesis, which contributes to endothelial ferroptosis. To further understand the impact of shear stress on the UPR and mevalonate pathways, we identified 11 transcription factors with decreased expression under LSS from the downregulated DEGs, including KLF6, MEF2C, ZNF91, and ZNF185, most of which have zinc finger motifs (**Figure 3A**). Using the silicon method, the promoter regions of PERK, HSP5A, MVD, and IDI1 are predicted to contain KLF6-specific binding elements (**Figure 3B**). This is consistent with the Western blot results in **Figure 3C**, showing that both LSS and HSS significantly reduce KLF6 protein expression compared to MSS. To confirm KLF6's crucial role in regulating these target genes, we created an endothelial cell line overexpressing KLF6 for a gain-of-function experiment (**Figure 3D**). The results in **Figure 3E-F** show that the expression levels of PERK, BiP, MVD, and IDI1 were significantly higher in KLF6-overexpressing endothelial cells compared to the negative control, regardless of whether the cells were treated with LSS or HSS. Additionally, the overexpression of KLF6 largely restored the expression of these four target proteins to levels similar to those observed under MSS.

Figure 3. KLF6 regulates the UPR and mevalonate pathway. (A) Heat map of the identified TF
genes downregulated by LSS; (B) Predicted KLF6-binding motifs in promoters of PERK, HSPA5,
MVD, IDI1 (JASPAR database; matrix ID: MA0472.1); (C) Protein expression of KLF6 under
different shear stress conditions; (D) Validation of KLF6 overexpression by Western blot (OE:
overexpression, NC: negative control); (E, F) KLF6-OE largely restores the expression of PERK,
BiP, MVD, and IDI1 under LSS and HSS to levels seen under MSS. Data are expressed as the
mean ± standard deviation from three or more independent experiments. Statistical significance
was determined using an unpaired t-test, with *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and ****$p <$
0.0001.
196

Verification of KLF6's role in endothelial ferroptosis and early atherosclerotic events caused
by abnormal shear stress
As illustrated in **Figure 4A-H**, KLF6 overexpression under both LSS and HSS led to an increase
in SLC7A11 protein expression and CoQ10 levels. Conversely, it reduced the extracellular LDH
level and lipid peroxidation. These results clearly demonstrate that KLF6 overexpression can
significantly inhibit ferroptosis induced by LSS and HSS. Based on our previously developed
multi-type cell (VSMC/EC/THP-1)in vitro atherosclerosis model [29], the anti-atherosclerotic
effect of KLF6 overexpression on the early events of atherosclerosis was examined. As illustrated
in Figure 4I-L, both monocyte adhesion and intracellular lipid accumulation (Oil Red O staining)
were significantly reduced in KLF6-overexpressed endothelial cells compared to the negative
control. These preliminary findings indicate that KLF6 might play a significant role in
atherosclerosis, though further validation through animal studies is necessary.

Figure 4. KLF6 overexpression alleviates endothelial ferroptosis and early atherosclerotic events induced by LSS and HSS. (A, E) Protein expression of SLC7A11; (B, F) ELISA results of extracellular LDH; (C, G) ELISA results of intracellular CoQ10; (D, H) Flow cytometry results of lipid peroxidation indicated by C11 BODIPY; (I, K) Fluorescent microscopic results of THP-1 cells (green label) adhering to the VSMC-EC bilayer cell architecture (The scale bar = 250 μm); (J, L) Lipid accumulation visualized by Oil Red O staining (The scale bar = 25 μm). Data are expressed as the mean ± standard deviation from three or more independent experiments. Statistical significance was determined using an unpaired t-test, with $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, and $****p < 0.0001$.

Discussion

219
The endothelium responds to changes in blood flow shear stress caused by the heartbeat,
regulating vasodilation and maintaining vascular homeostasis. It is generally believed that the
cause of atherosclerosis at the bend or bifurcation of the arteries is closely related to low shear
stress or oscillatory shear stress. In vivo studies on the impact of blood flow shear stress on
endothelial cells are difficult to perform, so researchers mainly use various in vitro fluid models.
However, fluid pressure is frequently overlooked in these studies. Our model demonstrates
variations in shear stress at 0.99, 4.78, and 24 dynes/cm², while keeping fluid pressure near the
physiological range of about 70 mmHg. To the best of our knowledge, this is the first study to
show that LSS and HSS induce ferroptosis in ECs. This result contrasts with the apoptosis
observed in other fluid models, possibly due to differences in fluid pressure. Another notable
finding is that the EC damage exhibits a V-shaped trend under LSS, MSS, and HSS. This suggests
that endothelial cells can adapt to a limited range of shear stress, although the underlying
mechanisms are still not understood.
233
RNA-seq analysis revealed that LSS primarily suppresses the transcriptome of steroid synthesis
and UPR-related pathways compared to MSS. Specifically, the transcription of genes such as
PERK, BiP, MVD, and IDI1 was reduced, leading to decreased SLC7A11 protein expression and
COQ10 synthesis, thereby impairing resistance to lipid peroxidation. Mechanical signal
transduction typically involves changes in the conformation of membrane proteins, leading to the
depolymerization and remodeling of actin within the cell, activation of transcription factors in the
cytoplasm, or epigenetic regulation through histone modifications. Given the complexity of the
transduction mechanism for shear mechanical signals, we focus on identifying clues from
transcription factors that respond to shear stress. Notably, KLF2 and KLF4, which are shear
stress-sensitive transcription factors, are absent from our DEGs dataset, whereas KLF6 is included.
Most studies report that KLF6 primarily regulates cell differentiation and tissue development.
Additionally, recent research has shown that KLF6 downregulation is also linked to oscillatory
flow-induced endothelial cell inflammation and oxLDL-induced endothelial dysfunction [23, 24].
As anticipated, KLF6-overexpressing ECs can restore the BiP-PERK-SLC7A11 and
MVD-IDI1-CoQ10 axials under both low and high shear stress conditions. This indicates that
KLF6 is crucial for cell survival, consistent with recent findings that KLF6 deficiency
significantly exacerbates liver damage, cell apoptosis, and hepatic inflammation [31].
251
As shear stress rises from low to high, ferroptosis follows a V-shaped pattern, while KLF6
expression displays an inverted V-shape. This suggests that cellular ferroptosis, triggered by both
excessive and insufficient shear stress, inversely correlates with KLF6 expression. The results of
KLF6 overexpression showed that KLF6 played a significant role in reducing endothelial cell
damage and lipid peroxidation under LSS and HSS. In other words, physiologically friendly shear
stress promotes KLF6 expression and enhances the antioxidant capacity of endothelial cells.
However, the distribution of shear stress in blood vessels is uneven due to variations in vessel
diameter and shape, complicating its numerical characterization in vivo. This hinders the
assessment of how changes in shear stress impact endothelial cell function in vivo. Consequently,
most in vitro models fail to simultaneously replicate the shear stress and pressure present in actual
arterial blood flow. A key benefit of our device is its ability to maintain a stable physiological

pressure range while delivering a variety of shear stresses. This could explain the ferroptosis we
observe, which differs from that in other in vitro models due to abnormal shear stress at a
physiological pressure of 70 mmHg. The anti-atherosclerotic effect of KLF6 was also assessed
using our previously established atherosclerosis model, which involves a three-cell culture system
(SMC, EC, THP-1). This evaluation demonstrated that KLF6 overexpression reduces the adhesion
of THP-1 cells and lipid accumulation induced by LSS and HSS.

269
270   **Figure 5**. Overview of KLF6 mediated the ferroptosis induced by abnormal shear stress
271

In summary, we examined the impact of different shear stress on endothelial cells using a
microfluidic chip to simulate low, medium, and high shear stress under a physiological pressure of
70 mmHg. Abnormal shear stress levels (0.99 and 24 dynes/cm^2) significantly induce endothelial
ferroptosis by inhibiting the KLF6-UPR/Mevalonate-SLC7A11/CoQ10 dual pathway (**Fig. 5**). Our
findings reveal a novel mechanism linking shear stress to the early stages of atherosclerosis,
explaining its preference for occurring at arterial bends and branch sites.
278

Methods and materials
Antibodies and reagents
Antibodies: SLC7A11 (1:1000, ab175186, Abcam), GRP78/BIP (1:1000, 66574-1-Ig, proteintech),
PERK/EIF2AK3 (1:1000, 24390-1-AP, proteintech), KLF6 (1:1000, 14716-1-AP, proteintech),
MVD (1:1000, 15331-1-AP proteintech) IDI1 (1:1000, 11166-2-AP, proteintech), Goat
Anti-Rabbit IgG-HRP (1:10,000, ASS1006-1, Biorigin), Goat Anti-Mouse IgG-HRP (1:10,000,
ASS1007-1, Biorigin). Reagents: Ferrostatin-1 (17729, Cayman), C11-BODIPY581/591 (27086,
Cayman), Propidium iodide (P4170, Sigma-Aldrich), Laminin (L2020, Sigma-Aldrich),
Opti-MEM (31985062, Gibco), M5 HiPer Lipo2000 Transfection Reagent (MF135-01, Mei5bio),
TRIzol reagent (15596026, Invitrogen). Dulbecco's Modified Eagle Medium (DMEM) (12800017,
Gibco), Fetal bovine serum (FBS) (10099141, Gibco), LDH detection kit (C0017, Beyotime),
Protease inhibitor and phosphatase inhibitor (5872S, Cell Signaling Technology), BCA protein
detection kit (P0012S, Beyotime), 4-HNE ELISA kit (RE12203-96T, Bioroyee), Oil Red O (ORO)
(Biolabs, China), CoQ10 ELISA kit (F13041-A, Beijing huabodeyi biology science and
technology co., ltd) RNAsimple Total RNA Kit (DP419,TIANGEN BIOTECH), CFDA-SE
(APExBIO, USA).

Fabrication of microfluidic chip

The microfluidic chip was made of three PDMS layers. The bottom layer contains multiple

parallel shallow grooves (height 0.075 mm, width 6 mm, and length 35 mm). The middle layer is

an elastic thin membrane to support cell culture. The top layer is cut with multiple parallel hollow

strips for alignment with the shallow grooves of the bottom layer. The hollow strips of the top

layer constrain the bulging of the middle layer to form a tunnel-like structure (Figure 1A). The

molds used for the top and bottom layers are made of SU-8 negative photoresist embossed on a

silicon wafer. The middle layer was obtained by a spin-casted PDMS on a blank silicon wafer.

After the middle layer is cured at 65 °C for 15 min, it is contacted by the top layer with the

hollowed strips and continued to cure for 1 h to form irreversible binding. Using the plasma

oxidation, the embedded groove in the bottom layer is aligned with the hollowed strips in the

middle-top layer assembly, and then the two parts are tightly compressed to form a sealed

microchannel between the middle and the bottom layers. The assembled microfluidic chip is

further cured overnight at 65 °C for use. A series of connecting holes is pre-punched at both ends

of the guide microchannel, facilitating its connection to a circulating flow path driven by a

peristaltic pump.

Cell lines and culture

A human aortic vascular smooth muscle cell line (T/G HAVSMC) was obtained from American

Type Culture Collection (ATCC, Manassas, VA, USA). A human umbilical vein endothelial cell

line (HUVEC-T1) and a human monocytic cell line (THP-1) were purchased from the National

Infrastructure of Cell Line Resource of China (Beijing, China). T/G HA-VSMC and

HUVEC-T1were cultured in DMEM (Gibco, USA) containing 10% FBS (Gibco, USA), penicillin

(100 U/ml, Gibco, USA), and streptomycin (100 μg/ml, Gibco, USA). Monocytic THP-1 cells

were cultured in RPMI 1640 medium (Gibco, USA) supplemented with 10% FBS (Gibco, USA),

penicillin (100 U/ml, Gibco, USA), and streptomycin (100 μg/ml, Gibco, USA). The cell culture

was maintained at 37 °C in 5% CO_2 in a humidified incubator.

Cell death assay

Cell death were evaluated by PI staining and LDH release assays. After 24 h shear stress treated,

the cells were washed with PBS three times and incubated with 5 μg/mL PI in the dark at 37 °C

for 30 min. The images were collected with a fluorescence microscope (DMI6000, Leica). The

LDH release was detected by a LDH detection kit (C0017, Beyotime) according to the

manufacturer's protocol. The data were collected by a multi-function microplate reader (Synergy

H1 Hybrid Reader, BioTeK).

Lipid peroxidation measurement

The cell lipid peroxidation was measured by C11-BODIPY 581/591 probe. After 24 h shear stress

treated, the cells were washed with PBS three times and incubated with 10 μM C11-BODIPY

581/591 in the dark at 37 °C for 30 min, then the cells were analyzed by flow cytometry.

ELISA for 4-HNE and CoQ10

The content of 4-HNE in the cells was detected by 4-HNE ELISA kit (RE12203-96T, Bioroyee)

according to the manufacturer's protocol. The content of CoQ10 in the cells was detected by
CoQ10 ELISA kit (F13041-A, Beijing huabodeyi biology science and technology co., ltd)
according to the manufacturer's protocol.

Western blotting
Cells were treated with shear stress for 24 h, total proteins in the cells were isolated by RIPA lysis
buffer (P0013B, Beyotime) with the complete protease inhibitor and phosphatase inhibitor (5872S,
Cell Signaling Technology) and protein concentration was determined using a BCA protein
detection kit (P0012S, Beyotime). After separation by SDS-PAGE, the proteins were transferred to
a PVDF membrane (ISEQ00010, Merck Millipore). The membrane was blocked with 5% non-fat
milk in TBST (20 mM Tris, 137 mM NaCl, 0.1% Tween- 20, pH7.4) at room temperature for 1 h,
incubated with the primary antibodies at 4 °C for overnight, and then the membrane was incubated
with anti-mouse or anti-rabbit IgG with HRP linked secondary antibody, followed by
chemiluminescence detection (WBKLS0100, Merck Millipore) with a gel imaging analysis
system (ChampChemi 910 Plus, Sagecreation).

RNA-seq data analysis
Total RNA was extracted from the shear stress treated endothelial cells using RNAsimple Total
RNA Kit (DP419,TIANGEN BIOTECH) according to the manufacturer's protocol. RNA
quantification and purity were determined using Nanodrop One (Thermo Fisher Scientific).
Qualified RNA samples were sent to BGI-Shenzhen (Shenzhen, China) for strand-specific library
construction using the NEBNext Ultra II RNA Library Prep Kit (NEB #E7770), followed by
sequencing on an Illumina NovaSeq 6000 platform with 150-bp paired-end reads. Raw data
processing and downstream analyses were conducted on BGI's Dr. TOM platform
(https://biosys.bgi.com).

Lentiviral transduction of HUVEC-T1
For the KLF6 overexpression experiment, two lentiviral particles were obtained from GENEWIZ:
one containing the human KLF6 gene (plvx-puro-KLF6) and the other an empty plvx-puro vector.
These were used for KLF6 overexpression and as a negative control, respectively. In brief,
HEK293T cells were transfected with either plvx-puro vector or plvx-puro-KLF6 constructs with
psPAX.2 and pMD2.G second-generation lentiviral packaging system. After 48 h incubation, the
lentivirus particles in the medium were collected and filtered to infect the HUVEC-T1 cells. After
48 h infection, 4 μg/mL puromycin was added into the cell culture medium to obtain the stable cell
lines with successful transduction.

THP-1 cell adhesion assay
THP-1 cells were incubated with 10 μM CFDA-SE (APExBIO, USA) for 15 minutes at 37 °C.
The stained THP-1 cell solution, at a density of 2.0×10^5 cells/ml, was then added to the
circulating medium. After the cells were treated with shear stress for 24 h, the number of attached
THP-1 cells was counted using a fluorescent microscope (Leica, DMI 6000B).

Oil Red O staining
After the cells were treated with shear stress for 24 h, they were rinsed with PBS three times, and

then fixed in 4% paraformaldehyde for 30 min, and then washed again with PBS three times. After the cells were stained with Oil Red O solution (ORO) (Biolabs, China) for 1 h at room temperature, the stained cells were incubated with 60% isopropyl alcohol for 2 min and a hematoxylin solution (Beijing Biodragon Immunotechnologies, China) for 3 min, followed by rinsing the cells with the tap water. The sample was covered with glycerin and then mounted with a slide. ORO-stained cells were photographed under the bright-field condition by a microscope (Leica Microsystems). ImageJ software was used for quantification after setting the appropriate threshold.

Transcription factor binding site prediction

Putative KLF6-binding sites in promoter regions of PERK (EIF2AK3), HSPA5 (BiP), MVD, and IDI1 were predicted using the JASPAR database (https://jaspar.elixir.no/; version 2022). Genomic sequences (2000 bp upstream of transcription start sites) were analyzed with the KLF6 position weight matrix (MA0472.1). Binding motifs containing GC-box elements (consensus: 5′-CACCC-3′) with a relative profile score threshold >85% were identified.

Statistical analysis

The data are reported as the means ± SD. All data used for the statistical analysis were from three independent experiments. Statistics were performed using an unpaired two-tailed Student's t-test. P values < 0.05 were considered statistically significant. All statistical analyses were performed with GraphPad Prism 5.

Acknowledgment

This work was supported by the National Natural Science Foundation of China (NNSFC) (31571481, 31771585, and 32070799).

Declaration of conflict of interest

The authors have no conflict of interest to disclose.

Author Contributions

All authors, including JC (Jingang Cui), FZY (Zhiyu Fan), SD (Suoqi Ding), J.Z. (Jiazhen Zhang), HS (Huihong Shen), SAZ (Syeda Armana Zaidi), and YD (Yongsheng Ding), were involved in the acquisition, analysis, and interpretation of data according to their respective roles. YD conceived and designed the study, drafted the manuscript, and approved the final version for submission.

Data availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

References:

1.   Martorell, J., et al., *Extent of flow recirculation governs expression of atherosclerotic and thrombotic biomarkers in arterial bifurcations.* Cardiovasc Res, 2014. **103**(1): p. 37-46.

2.   Wu, J., et al., *Disturbed flow impairs MerTK-mediated efferocytosis in aortic endothelial cells during atherosclerosis.* Theranostics, 2024. **14**(6): p. 2427-2441.

3.   Chen, C., et al., *Activation of GPR30 with G1 inhibits oscillatory shear stress-induced adhesion of THP-1 monocytes to HAECs by increasing KLF2.* Aging (Albany NY), 2021. **13**(8): p. 11942-11953.

4.   Rashad, S., et al., *Epigenetic response of endothelial cells to different wall shear stress magnitudes: A report of new mechano-miRNAs.* J Cell Physiol, 2020. **235**(11): p. 7827-7839.

5.   Bibli, S.I., et al., *Shear stress regulates cystathionine gamma lyase expression to preserve endothelial redox balance and reduce membrane lipid peroxidation.* Redox Biol, 2020. **28**: p. 101379.

6.   Sun, Z., et al., *Activation of GPR81 by lactate inhibits oscillatory shear stress-induced endothelial inflammation by activating the expression of KLF2.* IUBMB Life, 2019. **71**(12): p. 2010-2019.

7.   Qiao, C., et al., *Laminar Flow Attenuates Macrophage Migration Inhibitory Factor Expression in Endothelial Cells.* Sci Rep, 2018. **8**(1): p. 2360.

8.   Davies, P.F., et al., *The atherosusceptible endothelium: endothelial phenotypes in complex haemodynamic shear stress regions in vivo.* Cardiovasc Res, 2013. **99**(2): p. 315-27.

9.   Tamargo, I.A., et al., *HEG1 Protects Against Atherosclerosis by Regulating Stable Flow-Induced KLF2/4 Expression in Endothelial Cells.* Circulation, 2024. **149**(15): p. 1183-1201.

10.   Lingrel, J.B., et al., *Myeloid-specific Kruppel-like factor 2 inactivation increases macrophage and neutrophil adhesion and promotes atherosclerosis.* Circ Res, 2012. **110**(10): p. 1294-302.

11.   Atkins, G.B., et al., *Hemizygous deficiency of Kruppel-like factor 2 augments experimental atherosclerosis.* Circ Res, 2008. **103**(7): p. 690-3.

12.   Lu, Q., et al., *Shear-Sensitive lncRNA AF131217.1 Inhibits Inflammation in HUVECs via Regulation of KLF4.* Hypertension, 2019. **73**(5): p. e25-e34.

13.   He, M., et al., *Atheroprotective Flow Upregulates ITPR3 (Inositol 1,4,5-Trisphosphate Receptor 3) in Vascular Endothelium via KLF4 (Kruppel-Like Factor 4)-Mediated Histone Modifications.* Arterioscler Thromb Vasc Biol, 2019. **39**(5): p. 902-914.

14.   Sharma, N., et al., *Myeloid Kruppel-like factor 4 deficiency augments atherogenesis in ApoE-/- mice--brief report.* Arterioscler Thromb Vasc Biol, 2012. **32**(12): p. 2836-8.

15.   Lu, Y.W., et al., *MEF2 (Myocyte Enhancer Factor 2) Is Essential for Endothelial Homeostasis and the Atheroprotective Gene Expression Program.* Arterioscler Thromb Vasc Biol, 2021. **41**(3): p. 1105-1123.

16.   Kim, M., et al., *Laminar flow activation of ERK5 protein in vascular endothelium leads to atheroprotective effect via NF-E2-related factor 2 (Nrf2) activation.* J Biol Chem, 2012. **287**(48): p. 40722-31.

17.   Lee, J.Y., et al., *Fluid shear stress regulates the expression of Lectin-like oxidized low density lipoprotein receptor-1 via KLF2-AP-1 pathway depending on its intensity and pattern in endothelial cells.* Atherosclerosis, 2018. **270**: p. 76-88.

18.   Fang, F., et al., *Cathepsin K contributed to disturbed flow-induced atherosclerosis is dependent on integrin-actin cytoskeleton-NF-kappaB pathway.* Genes Dis, 2023. **10**(2): p. 583-595.

19.   Walther, B.K., et al., *Mechanotransduction-on-chip: vessel-chip model of endothelial YAP mechanobiology reveals matrix stiffness impedes shear response.* Lab Chip, 2021. **21**(9): p. 1738-1751.

20.   Jiang, M.C., et al., *Thioridazine protects against disturbed flow-induced atherosclerosis by inhibiting RhoA/YAP-mediated endothelial inflammation.* Acta Pharmacol Sin, 2023. **44**(10): p. 1977-1988.

21.   Atkins, G.B. and M.K. Jain, *Role of Kruppel-like transcription factors in endothelial biology.* Circ Res, 2007. **100**(12): p. 1686-95.

22.   Syafruddin, S.E., et al., *Two Sides of the Same Coin: The Roles of KLF6 in Physiology and Pathophysiology.* Biomolecules, 2020. **10**(10).

23.   Wei, G., et al., *The protective effects of azilsartan against oscillatory shear stress-induced endothelial dysfunction and inflammation are mediated by KLF6.* J Biochem Mol Toxicol, 2021. **35**(6): p. 1-8.

24.   Wang, H., et al., *Apremilast ameliorates ox-LDL-induced endothelial dysfunction mediated by KLF6.* Aging (Albany NY), 2020. **12**(19): p. 19012-19021.

25.   Tabas, I., *The role of endoplasmic reticulum stress in the progression of atherosclerosis.* Circ Res, 2010. **107**(7): p. 839-50.

26.   Feaver, R.E., et al., *GRP78 upregulation by atheroprone shear stress via p38-, alpha2beta1-dependent mechanism in endothelial cells.* Arterioscler Thromb Vasc Biol, 2008. **28**(8): p. 1534-41.

27.   Bailey, K.A., et al., *Atherosusceptible Shear Stress Activates Endoplasmic Reticulum Stress to Promote Endothelial Inflammation.* Sci Rep, 2017. **7**(1): p. 8196.

28.   Kim, S. and C.H. Woo, *Laminar Flow Inhibits ER Stress-Induced Endothelial Apoptosis through PI3K/Akt-Dependent Signaling Pathway.* Mol Cells, 2018. **41**(11): p. 964-970.

29.   Cui, J., et al., *Microfluidic-to-macrofluidic: A simple in vitro model of atherosclerosis induced by fluidic stimulation.* Biomicrofluidics, 2023. **17**(4): p. 044106.

30.   Saini, K.K., et al., *Loss of PERK function promotes ferroptosis by downregulating SLC7A11 (System Xc(-)) in colorectal cancer.* Redox Biol, 2023. **65**: p. 102833.

31.   Li, J., et al., *KLF6 alleviates hepatic ischemia-reperfusion injury by inhibiting autophagy.* Cell Death Dis, 2023. **14**(7): p. 393.

498