Adeno-associated virus (AAV)-TBX18 does not generate biological pacemaker activity, unlike AAV-Hcn2

Jianan Wang[1], Mathilde R. Rivaud[1,2], Mischa Klerk[1], Arie R. Boender[1,3], Ruud N. Visser[1], Rinske Sparrius[1], Hee Young Lee[4], Karel van Duijvenboden[1], Yuting Yang[3], Emiel J.M. Kramer[3], Kyung Ho Park[4], Larry C. Park[4], Silke Schrödel[5], Christian Thirion[5], Eric Ehrke-Schulz[6], Anja Ehrhardt[6], Osne F. Kirzner[3,7], Klaus Neef[3,8], Hanno L. Tan[3,8,9], Arie O. Verkerk[1,9], Vincent M. Christoffels[1], Gerard J.J. Boink[1,3,10]

[1]Department of Medical Biology, Amsterdam Cardiovascular Sciences, Amsterdam University Medical Centers, University of Amsterdam, Meibergdreef 15, 1105 AZ, Amsterdam, the Netherlands

[2]Laboratory of Experimental Cardiology, Department of Cardiology, Leiden University Medical Center (LUMC), Albinusdreef 2, 2300 RC, Leiden, the Netherlands

[3]PacingCure BV, Roetersstraat 35, 1018 WB, Amsterdam, the Netherlands

[4]Naason Science Inc, 123 Saengmyung-ro, Cheongju, Korea

[5]Revvity Gene Delivery GmbH, Am Haag 6, 82166 Graefelfing, Germany

[6]Virology and Microbiology, Center for Biomedical Education and Research, Department Human Medicine, Faculty of Health, University Witten/Herdecke, Stockumer Stre 10, 58455 Witten, Germany

[7]Department of Anaesthesiology, Amsterdam University Medical Centers, De Boelelaan 1117, 1081 HV, Amsterdam, the Netherlands

[8]Netherlands Heart Institute, Moreelsepark 1, 3511 EP, Utrecht, the Netherlands

[9]Department of Clinical and Experimental Cardiology, Amsterdam Cardiovascular Sciences, Amsterdam University Medical Centers, University of Amsterdam, Meibergdreef 15, 1105 AZ, Amsterdam, the Netherlands

[10]Department of Cardiology, Amsterdam Cardiovascular Sciences, Amsterdam University Medical Centers, University of Amsterdam, Meibergdreef 9, 1105 AZ, Amsterdam, the Netherlands

Address for correspondence: Gerard J.J. Boink, Amsterdam University Medical Centers, Meibergdreef 9, 1105AZ, Amsterdam, the Netherlands. E-mail: g.j.boink@amsterdamumc.nl

Conflict-of-interest statement

J.W., A.R.B., Y.Y., E.J.M.K., O.F.K., K.N., and H.L.T. are employees of PacingCure BV. O.F.K., H.L.T,. and G.J.J.B. report ownership interest in PacingCure BV. S.S. and C.T. were employees of Revvity Gene Delivery GmbH, a wholly-owned subsidiary of PerkinElmer Inc. C.T. receives shares from the company. H.Y.L, K.H.P., and L.C.P. are employees of Naason Science Inc. L.C.P reports ownership interest in Naason Science Inc. J.W., R.V., V.M.C., and G.J.J.B. filed patent applications concerning application of uORF technologies. Other authors declare no conflicts of interest.

Abstract

Gene therapy-based biological pacemakers have been proposed as an alternative to their hardware-based counterparts. In this context, short-term ectopic expression of the T-box transcription factor 18 (TBX18) in the ventricle reportedly generated potent short-term pacemaker function in various animal models. Here, we investigated the effect of adeno-associated virus (AAV)-mediated long-term expression of TBX18, and compared the outcome to that of the pacemaker ion channel Hcn2. Our findings revealed that CMV-driven ectopic TBX18 expression in mouse hearts led to severe cardiac fibrosis. At lower, non-fibrogenic levels, TBX18 maintained its transcriptional function but failed to induce pacemaker phenotypes. TBX18-expressing cells showed suppressed expression of key working myocardial genes, but the pacemaker gene program was not induced. Electrophysiological studies showed abnormal automaticity in TBX18-expressing cells, combined with prolonged repolarization and various current changes. However, no hyperpolarization-activated funny current was detected. In a complete AV-block rat model, AAV-mediated Hcn2 expression induced robust ectopic pacemaker activity in the presence of isoproterenol, whereas TBX18 expression neither generated such activities, nor augmented Hcn2-mediated pacing. In conclusion, at functional non-fibrogenic levels, TBX18 is neither sufficient nor necessary to induce pacemaker activity. In contrast, Hcn2 generates reliable pacing, making it a more viable candidate for biological pacemaker development.

Key words

TBX18, heart, biological pacemaker, cardiomyocytes, adeno-associated viral vector

1 Introduction

Symptomatic bradycardia can pose a significant risk to life if left untreated. Electrical

pacemakers have been used to successfully manage this condition, effectively

restoring and maintaining normal cardiac rhythm. However, shortcomings inherent to

these devices, such as limited battery life, complications associated with surgical

procedures and inadequate autonomic modulation, underscore the necessity for more

advanced therapeutic strategies (1, 2).

Gene therapy-based biological pacemakers are a promising alternative to electronic

pacemakers (1, 3, 4). By modulating gene expression to mimic function of sinoatrial

nodal (SAN) pacemaker cells, biological pacemakers aim to induce spontaneous

diastolic depolarizations, capable of driving the heart at physiologically relevant rate.

One approach is to manipulate ion channels (e.g. Hcn2 and Kir2.1) in order to increase

inward current and/or decrease outward current during diastole (5-9). Transduced

cardiomyocytes would then spontaneously depolarize and function as pacemaker cells.

A potentially more comprehensive approach is based on transcription factor

overexpression to reprogram cardiomyocytes into pacemaker cells that are functionally

and phenotypically similar to SAN pacemaker cells (10, 11). These reprogramed

cardiomyocytes are expected to emulate SAN cells not only in their

electrophysiological properties, but also in their morphology and gene expression

profiles.

T-box transcription factor (TBX) 3 and 18 are required for the development of the SAN

and as such represent obvious targets for the biological pacemaker development (12,

13). Ectopic atrial TBX3 expression was shown to impose pacemaker phenotype using

a transgenic expression system activated during early embryogenesis (14). However,

tamoxifen-induced TBX3 expression in adult cardiomyocytes failed to do so but solely

suppressed working myocardial phenotype (10). On the other hand, adenoviral (AdV)

vector-mediated TBX18 overexpression was reported to converted ventricular

chamber cardiomyocytes into pacemaker cells and generated biological pacing in both

small and large animal models of heart block (11, 15, 16). Moreover, TBX18

overexpression in subsidiary atrial pacemaker tissue was shown to increase the

spontaneous beating rates (17). However, the AdV vector used in these studies is

highly immunogenic and, as a result, provides only transient transgene expression,

impeding clinical translation (18, 19). In line with this immunological issue, pacing

function waned during the follow-up period of several weeks. Alternatively, this

transient pacemaker phenotype could also be due to insufficient reprogramming. Direct

injection of TBX18 mRNA was recently explored as an alternative, but this approach

was complicated by the need transforming growth factor B (TGFB) inhibition (as a co-

factor to suppress fibrosis), and remained suboptimal in terms of pacemaker durability

(20).

Adeno-associated viral (AAV) vectors are non-pathogenic, have low immunogenicity

and can sustain long-term expression lasting for years (4, 21). Systemic delivery of

AAV vector has been used in the field of cardiac gene therapy for over a decade,

providing robust transgene expression in the whole heart (22-25). Recently, local

delivery of AAV vectors has been demonstrated by several groups to achieve efficient

transgene expression in large animal models (26, 27). AAV vectors have been

approved by the US regulatory authorities for human application, further illustrating the

strong potential for clinical translation (28). We hypothesized that ectopic AAV-

mediated TBX18 expression would convert chamber cardiomyocytes into pacemaker

cells and thereafter maintain a stable pacemaker phenotype. In order to test this

hypothesis, it was critical to reduce TBX18 expression to non-cytotoxic levels. This

allowed us to conduct further mechanistic studies on TBX18-expressing

cardiomyocytes that were not possible in previous studies due to the transient

expression systems employed (11, 15, 20). These mechanistic studies indicated that

TBX18 expression *in vivo* did not induce a pacemaker phenotype, but induced

abnormal automaticity. In the absence of local tissue inflammation (that has

confounded previous studies utilizing AdV vectors), this induced phenotype was

unable to generate any meaningful pacemaker activity in a complete AV-block rat

model, while AAV-Hcn2 generated reliable pacing in the presence of isoproterenol.

Results

Adenoviral vectors support ectopic pacing in rats with complete AV-block

Previous studies showed that AdV-mediated TBX18 overexpression in ventricular cardiomyocytes (CM) induced the formation of ectopic pacemaker activity (11, 15, 16). To be able to recapitulate these findings, we used an established surgical AV-block rat model (29, 30). Electrical AV node ablation was performed and the successful generation of complete AV-block was confirmed by disassociation between P waves and QRS complexes on the surface cardiac electrocardiogram (ECG) (Figure 1A). We injected AdV vector expressing FLAG-tagged human TBX18 (AdV-TBX18) or GFP (AdV-GFP) as a control intramyocardially to the apex of the heart (Figure 1B). Saline injection was used as sham control. One week later, ECGs were measured before and after isoproterenol administration. Successful transgene delivery was confirmed by immunohistochemistry (Figure 1C). ECG analyses (Figure 1D) revealed that both AdV-TBX18 and AdV-GFP animals had significantly more profound ectopic pacing from the injection site and higher heart rate following isoproterenol administration when compared to the saline group (Figure 1, E and F). However, no significant difference was observed between AdV-TBX18 and AdV-GFP animals regarding the frequency of the ectopic pacing or the heart rate (Figure 1, E and F). Picrosirius red staining showed extensive cardiac fibrosis at the injection site of both AdV-TBX18 and AdV-GFP animals (Figure 1, G and H). These results suggested that the ectopic pacing observed was likely based on local tissue inflammation generated by the AdV vector used, rather than by TBX18 overexpression.

TBX18 ectopic expression generates fibrotic scars in the mouse heart

To circumvent the use of AdV vector and at the same time achieve long-term TBX18 expression, we injected adult mice with AAV vectors containing CMV promoter and FLAG-tagged human TBX18 (AAV-CMV-TBX18). AAV vectors containing CMV promoter and FLAG-tagged GFP were used as control (AAV-CMV-GFP). To our surprise, a profound fibrotic scar was observed at the injection site in 9 out of 10 TBX18-injected mice, 4 weeks post injection, while such scar was absent from all 9 GFP-injected mice (Figure 2, A and B). Interestingly, the single TBX18 heart in which fibrosis was not detected was found to express TBX18 at the lowest level among the 10 TBX18 hearts (Supplemental Figure 1), suggesting that the fibrosis is likely related to the high levels of TBX18 overexpression, typically generated with conventional CMV-driven vectors. To investigate the progression of fibrosis, we performed histological studies on hearts harvested 4 days, 7 days and 28 days post injection of AAV-CMV-TBX18. Cardiac fibrosis appeared as early as 7 days post injection and progressed into substantial myocardial scars at 28 days post injection (Figure 2C). Immunofluorescence staining revealed that TBX18 expression accompanied the fibrotic tissues (Figure 2D). These results indicate that high levels of TBX18 overexpression are cytotoxic and lead to severe cardiac fibrosis.

Optimized AAV vector expressing TBX18 circumvents fibrosis while maintaining transcriptional functionality

We hypothesize that the fibrosis observed was due to the supra-physiological levels of TBX18 expression. To reduce expression levels of TBX18, we added an upstream open reading frame (uORF) in the 5' UTR of the transgene (Figure 3A), which was previously reported to reduce the translational efficiency.(31, 32) To test this approach in the context of an AAV expression cassette, we started with exploratory GFP

(transfection) and TBX18 (transduction) experiments. As expected, introduction of this

uORF did not change the transfection efficiency (fraction of GFP-positive cells), while

effectively reducing GFP expression levels (Figure 3, B-D). Lowering the plasmid dose

significantly reduced GFP expression level but also lowered the fraction of transduced

cells (Figure 3, B-D). In a similar experiment, introduction of this uORF significantly

reduced TBX18 expression level by 42% (Figure 3, E and F), confirming the effect of

the uORF in our TBX18 overexpression vector. In neonatal rat ventricular myocytes

(NRVM), introduction of the uORF and switching from the ubiquitously active CMV

promoter to the cardiomyocyte-specific cTnT promoter reduced the TBX18 expression

level to 25% and 39% to that of that from the original AAV-CMV-TBX18 vector,

respectively (Figure 3, G and H). The combination of both further reduced the TBX18

expression level to 1% of the original vector and allows cardiomyocyte-specific

expression (Figure 3G and 3H). We then sought to evaluate the function of this

optimized AAV vector containing both cTnT promoter and uORF (AAV-cTnT-uORF-

TBX18), which drives TBX18 at low expression levels and specifically in

cardiomyocytes. NRVMs were transduced with both AAV-CMV-TBX18 and AAV-cTnT-

uORF-TBX18 at a multiplicity of infection of 50,000. AAV-CMV-GFP was used as

negative control while previously reported AdV vector bicistronically expressing TBX18

and GFP served as positive control (11). Quantitative real time-polymerase chain

reaction (RT-qPCR) showed a significant upregulation of pacemaker gene *Hcn4* and

down regulation of chamber genes *Gja1*, *Scn5a*, *Kcnj2*, and *Pln* in all three TBX18

vectors compared to the negative control. No significant differences were observed

between cells transduced with AAV-CMV-TBX18 and AAV-cTnT-uORF-TBX18 (Figure

3I and 3J), indicating that the canonical transcription factor functionality of TBX18 was

maintained in the optimized TBX18 vector.

We subsequently studied fibrosis and myocardial scarring in mice injected with AAV-cTnT-uORF-TBX18, and compared outcomes to mice injected with the conventional AAV-CMV-TBX18 and two vectors expressing TBX18 incorporating either CMV-uORF or cTnT (without uORF; Figure 4A). Hearts injected with AAV-CMV-GFP were used as control. Picrosirius red staining illustrated that hearts injected with AAV-CMV-TBX18 formed an extensive fibrotic scar at the injection site with no viable myocardium left 4 weeks post injection (Figure 4B). On the contrary, hearts injected with AAV-cTnT-uORF-TBX18 had virtually no fibrosis, which was comparable to control (Figure 4, B and C). The other two groups showed intermediary outcomes. To study the long-term effect, we also performed analyses on hearts harvested 8 weeks post injection. Injections with AAV-CMV-TBX18 were not performed for this long-term study for animal welfare reasons. Fibrosis in hearts injected with AAV-cTnT-uORF-TBX18 did not differ from that in control hearts (Figure 4, D and E). These data indicated long-term tolerance of the AAV-cTnT-uORF-TBX18 vector. As AAV-cTnT-uORF-TBX18 vector is functionally equivalent to the other three vectors and tolerant for long-term expression, we continued our analyses with this vector.

TBX18 suppresses the working myocardium phenotype

We focused the *in vivo* functional analysis on AAV-cTnT-uORF-TBX18, using AAV-CMV-GFP as a control (Figure 5A). Immunohistochemistry revealed expression of TBX18 or GFP at the injection sites, indicating robust transgene delivery (Figure 5, B and C). At 4 weeks post injection, Connexin43 (Cx43, encoded by *Gja1*), a direct downstream target of TBX18 (33) was efficiently suppressed in the TBX18-positive region when compared to the TBX18-negative region (Figure 5C). In control animals injected with AAV-GFP, Cx43 was homogenously detected across the GFP-positive and GFP-negative regions (Figure 5C). However, expression of the pacemaker ion

channel Hcn4 remained undetectable at 4 or 8 weeks post injection, despite robust

TBX18 expression (Figure 5B). As the positive control, Hcn4 was clearly detectable at

the AV nodal region (Figure 5B).

To identify and dissect the transduced tissues, we modified AAV-cTnT-uORF-TBX18

by adding a self-cleaving p2A-GFP as a fluorescent marker (AAV-cTnT-uORF-TBX18-

GFP) (Figure 5D). The same vector expressing nanoluciferase (AAV-cTnT-uORF-

Nluc-GFP) was used as control. Two weeks post injection, the GFP-positive region

was isolated by micro-dissection and used for quantitative RT-PCR to evaluate

changes in gene expression. Expression of exogenous TBX18 was confirmed,

whereas endogenous expression of *Tbx18* remained unaffected at low level (Figure

5E). A significant downregulation of the direct target *Gja1* indicates the transcriptional

functionality of TBX18 delivered (Figure 5E). Expression of cardiac fibrosis marker

Postn (34, 35) was not affected, confirming the non-cytotoxic expression level of

TBX18 (Figure 5E). We observed significant down-regulation of the cardiac Na$^+$

channel encoding gene *Scn5a* and K$^+$ channel encoding gene *Kcnj5* (Figure 5F), which

are important for cardiac conduction and modulation of resting membrane potential,

respectively. K$^+$ channel encoding genes *Kcnj2* and *Kcnd2* were non-significantly

downregulated (p = 0.11 for *Kcnj2* and p = 0.06 for *Kcnd2*) (Figure 5F). On the other

hand, *Hcn4* was not induced by TBX18 overexpression (Figure 5F), in line with our

immunohistochemistry results. Ventricular stress marker *Nppa* was significantly

upregulated (Figure 5F). RNA sequencing analysis confirmed these findings, and

further strengthened the notion that working myocardial genes were suppressed but

pacemaker genes were not induced (Supplemental Figure 2).

TBX18 expression generates abnormal automaticity in mouse cardiomyocytes

To assess the electrophysiological consequences of TBX18 expression in cardiomyocytes, single left ventricular cardiomyocytes were isolated from mouse hearts 2 weeks post injection and were subjected to patch-clamp analysis. Spontaneous beating was clearly visible under the microscope in the isolated TBX18-expressing cardiomyocytes. Therefore, we proceeded to measure both spontaneous and overdrive-stimulated (4 Hz) action potentials (AP). Control cardiomyocytes did not generate spontaneous APs in 7 out of the 8 cardiomyocytes measured, while 8 out of the 8 TBX18-expressing cardiomyocytes exhibited spontaneous activity (Figure 6, A and B). The spontaneous APs of the TBX18-expressing cardiomyocytes had oscillations around plateau level and the time of such oscillations varied between approximately 2 to 10 seconds. We then performed 4 Hz overdrive stimulation to measure the AP parameters in detail. Typical recordings are shown in Figure 6C and average data are summarized in Figure 6D. All AP parameters measured differed between TBX18-overexpressing cardiomyocytes and control. The resting membrane potential of TBX18-expressing cardiomyocytes was depolarized by 5.8 mV, and the maximal AP upstroke velocity was significantly reduced, resulting in a significantly lower AP amplitude (Figure 6D). The AP durations at 20, 50 and 90% of repolarization were all significantly longer (Figure 6D).

We subsequently applied a generic voltage-clamp protocol to test net membrane currents in control and TBX18-expressing cardiomyocytes. Example current tracings and average voltage-current relationships indicate a difference in inward rectifier K_+ current (I_{K1}) and outward potassium currents (I_K) between TBX18-expressing cells and control (Figure 6, E-H). Compared to control, TBX18-expressing cardiomyocytes

presented a 23% reduction in I_{K1} (at -100 mV) (Figure 6G), coincident with the down-regulated *Kcnj2* expression found in our RT-qPCR analysis. TBX18-expressing cardiomyocytes also presented a 39% reduction in I_K (at 50 mV) (Figure 6H), coincident with the down-regulated *Kcnd2* expression found in our RT-qPCR analysis. In the voltage-clamp experiment, time-dependent, inward currents that activated upon the hyperpolarizing voltage steps were not observed, indicating absence of the funny current (I_f) in both control and TBX18-overexpressing cardiomyocytes (Figure 6I). We next proceeded with Na$^+$ current (I_{Na}) measurement which revealed a 71.3% reduction in TBX18-overexpressing cardiomyocytes (Figure 6, J and K), consistent with the reduced maximal AP upstroke velocity and *Scn5a* expression. Whole-cell capacitance was also recorded as a measure of cell size, which was not different between TBX18-expressing and control cardiomyocytes (Figure 6L).

Taken together, our patch-clamp data revealed that ectopic TBX18 expression in mouse cardiomyocytes generates a distinctive electrophysiological phenotype that is incapable of sustaining a regular rhythm. The observed spontaneous depolarizations in these TBX18-expressing cardiomyocytes were the result of potassium channel dysregulation, rather than from the canonical mechanisms seen in SAN pacemaker cells.

AAV-mediated Hcn2 expression generates ectopic ventricular pacemaker activity while TBX18 expression does not

Previous studies have suggested that interference with the inward rectifier current, via overexpression of a dominant negative variant of Kir2.1 can possibly liberate endogenous pacemaker activity (36), or support Hcn2-based pacemaker activity (8, 37). We therefore proceeded with testing AAV-cTnT-uORF-TBX18 in the complete AV-

block rat model (Figure 7A). In doing so, we compared outcomes to overexpression of the pacemaker channel Hcn2 and also tested the combination of Hcn2 and TBX18 overexpression. Successful transgene delivery was confirmed via immunofluorescence staining (Figure 7B). At baseline, all animals showed similar frequency of ectopic pacing and average heart rate. Neither TBX18-injected animals nor saline-injected animals responded to the isoproterenol administration (Figure 7C-7E). However, intriguingly, Hcn2-injected animals exhibited robust ectopic pacing in response to isoproterenol (Figure 7, C and D), as illustrated by the increase in average heart rate (Figure 7E). Co-injection of AAV-cTnT-Hcn2 and AAV-cTnT-uORF-TBX18 did not further augment Hcn2-based pacemaker function (Figure 7, C-E). No significant cardiac fibrosis was observed at the injection site of both saline and AAV-injected animals (Figure 7, F and G). Taken together, our data shows that AAV-mediated Hcn2 expression generates a viable pacemaker phenotype in response to isoproterenol, while TBX18 expression does not.

Analysis the transcriptional changes induced by *Hcn2* and *TBX18* expression in NRVMs

To investigate the impact of *Hcn2* or *TBX18* expression on cardiomyocytes, we transduced NRVMs with the *Hcn2* vector and all four *TBX18* vectors used in this study, followed by RNA-sequencing. AAV vector with the cTnT promoter and no coding sequence was used as control. Principal component analysis revealed a clear segregation of the samples based on *Hcn2* and on *TBX18* expression. All samples of the same group clustered together, and all four TBX18 groups clustered closely based on overall expression profiles (Figure 8A). Differentially expressed genes across the groups clustered into three distinct expression profiles (Figure 8B). The Hcn2 and TBX18 groups shared similarity in gene expression profiles corresponding to gene

cluster 1 and 2, where cluster 1 marks the upregulated genes and cluster 2 marks the down regulated genes compared to control. Notably, expression of the transcription factors necessary for SAN development (e.g. *Isl1*, *Shox2*, *Tbx3*) were not induced in any of the TBX18 groups (Figure 8, B and D), in line with our findings (Figure 3, Figure 5 and Figure 6) and other studies using NRVMs and mice (38, 39).

Genes in cluster 3 were specifically suppressed in the TBX18 groups, and were related to various signaling pathways (e.g. *Vegfa*, *Bmp3*, *Pln*), ion channels (e.g. *Scn5a*, *Kcnj5*, *Cacna2d2*) and cell-cell communications (*Gja1*, *Gja5*, *Cadh17*) (Figure 8, B-E). Cluster 1 genes were induced in both TBX18 and Hcn2 groups compared to control, suggesting that the elevated expression of those genes may reflect a broader consequence of transgene overexpression, such as disruption of the normal maturation process and/or pathological remodeling of the ventricular myocytes. Notably, genes in this cluster, such as *Hcn4*, *Myl4* and *Nppa*, are typically expressed only in fetal, but not adult ventricular cardiomyocytes (40-42). We observed a 3-fold decrease in *Hcn4* expression over 5 days of culture in untreated NRVMs (Supplemental Figure 3), suggesting that the observed increase in *Hcn4* in TBX18 and Hcn2 groups compared to control may be caused by attenuated downregulation that normally occurs in untreated NRVMs during their maturation. Reactivation of these fetal genes in adult ventricular myocytes has been associated with cardiac disease and pathological remodeling (43-45).

We then compared the transcriptional profiles of NRVMs transduced with Hcn2 and TBX18 from the current study, as well as NRMV transduced with TBX18 from Foster et al. (39), to those of postnatal day 4 (P4) mouse SAN pacemaker cells (46) (Figure 8, F-H). Our analysis revealed that neither Hcn2 nor TBX18 induced a transcriptional signature resembling that of native SAN pacemaker cells. For example, the expression

of the TBX18-induced level of *Hcn4* and *Isl1* remains 48-fold and 3600-fold lower than

those in native newborn SAN pacemaker cells, respectively, whereas common

cardiomyocyte genes were expressed at comparable levels. Comparison between

these datasets to embryonic day 16.5 (E16.5) mouse SAN pacemaker cells (47) also

showed similar results (Supplemental Figure 4). These results together suggest that

transducing ventricular myocytes with Hcn2 or TBX18 does not result in the generation

of *bona fide* SAN-like pacemaker cells. Furthermore, these data indicate the Hcn2-

based biological pacing is likely attributable to the functional modification of working

cardiomyocytes through overexpression of Hcn2 itself, as anticipated.

Discussion

In this study, we ectopically expressed TBX18 in adult murine ventricular myocardium using AAV-mediated delivery with the goal to generate long-term pacemaker function (11, 15, 16). However, we observed that conventional overexpression of TBX18 leads to severe cardiac fibrosis, which could be prevented by modification of the expression cassette to reduce protein expression levels of TBX18 while maintaining its transcriptional function. We found that TBX18 expression at non-cytotoxic levels suppressed working myocardial genes, in accordance with previous findings (11, 15, 38), but did not induce a genuine pacemaker phenotype. As a result, AAV-mediated TBX18 expression did not induce a pacemaker phenotype in AV-block rats, in contrast to Hcn2 expression, which generated robust ventricular pacing in response to isoproterenol.

We observed ectopic pacing activities from both AdV-GFP-injected and AdV-TBX18-injected animals using a complete AV-block rat model, suggesting that these activities are induced by the AdV vector itself instead of the transgene delivered. Previous studies showed similar results where AdV-TBX18 generated automaticity in rat complete AV-block models however not significantly different from AdV-GFP-induced automaticity (29, 30). Moreover, AdV infection rapidly provokes inflammation and infiltration of host immune cells in hearts (48). Recombinant AdV vectors, albeit being replication-defective, still activate both innate and adaptive immunity, leading to an inflammatory response (18, 19, 49). Indeed, we and others found severe fibrosis at the injection sites of AdV-GFP and AdV-TBX18 (30). These immune/inflammatory responses make it difficult to specifically study the effect of candidate biological pacemaker genes, and might also contribute to the inconsistent outcomes in different studies utilizing AdV-TBX18 (11, 20, 30).

In our study, AAV vectors were used to deliver TBX18 because of their low immunogenicity, allowing us to study the effect of TBX18 overexpression without the confounding effects of a severe immune response. Moreover, the use of the AAV vector system also would enable long-term expression of TBX18, aiming to facilitate a chronic treatment effect. In our experiments, ectopic expression of TBX18 using a conventional CMV-driven AAV vector was highly cytotoxic and therefore fibrogenic in mouse hearts. The absence of fibrosis in control where GFP was expressed from the same vector ruled out other potential non-transgene related causes such as mechanical trauma from injection or immune responses to viral vectors. Indeed, others have also found that TBX18 induced significantly more fibrosis than GFP when expressed from directly injected mRNA or from AdV vectors (30, 50). Inhibition of TGFB signaling using small molecules has been reported to mitigate such fibrosis and maintains the TBX18-induced automaticity *in vitro* and *in vivo* (30). However, TGFB signaling is also heavily involved in epithelial to mesenchymal transition (51), which is the proposed mechanism of TBX18-mediated conversion from cardiomyocytes to pacemaker cells (39). Furthermore, Tgfb signaling plays a crucial role in many physiological processes, such as tissue development, homeostasis and injury repair (52). These pleiotropic effects of TGFB inhibition are therefore generally undesirable from a translational perspective.

In order to attenuate fibrosis, we lowered expression levels of TBX18. We did not lower the viral titer to reduce expression, because AAV transduction follows a Poisson distribution (53), hence lower transduction, will lead to a mosaic pattern of transduced and un-transduced cells, making it more difficult to reach the critical mass needed for reliable pacemaker performance. Instead, a uORF was inserted in the 5'UTR of the TBX18 expression cassette. The theoretical advantage of this approach is that it allows

to reduce protein levels without the need to reduce transduction efficiency or

transcriptional activity of the expression cassette. Both applying the uORF and

switching to a cardiac-specific promoter that is less strong than CMV (54, 55)

significantly reduced TBX18-induced fibrosis in mouse hearts, and the combination

completely eliminated it. This combination reduced the expression level of TBX18 to

approximately 1% of the original level but still maintained its transcriptional function,

as the suppression of its direct downstream target *Gja1* (33) and other target genes

remained robust both *in vitro* as well as *in vivo*. Our RNA sequencing results showed

similar transcriptional profiles across all four TBX18 groups further supporting the

functional equivalence of TBX18 delivered by different vectors.

Our *in vitro* and *in vivo* molecular experiments showed that ectopic expression of

TBX18 suppressed the expression of chamber myocardial genes *Gja1*, *Scn5a, Kcnj5*

and *Kcnj2*, in agreement with previous studies using a transgenic mouse line and AdV-

mediated overexpression (11, 15, 38). This finding is further supported by our

electrophysiological experiments where I_{K1}, I_K and I_{Na} significantly decreased in

TBX18-expressing cells. In terms of expression of pacemaker genes, neither the

TBX18-transduced NRVMs from our study nor the ones from others (39, 56) show

resemblance to the transcription profiles of the *bona fide* SAN pacemaker cells. For

instance, genes for essential SAN pacemaker transcription factors ISL1, SHOX2 and

TBX3, each essential for SAN pacemaker cell differentiation and maintenance, are not

induced by TBX18 (57). Contradictory responses of *Hcn4* to ectopic TBX18 expression

in NRVMs have been reported with two studies showing induction (11, 39) and another

study showing reduction (56). In our study we observed higher expression of *Hcn4* in

NRVMs transduced with TBX18 compared to control. However, a similar increase was

also observed in NRVMs transduced with Hcn2, indicating the induction is not specific

to TBX18 but may be a more general response to gene overexpression. The apparent

induction of *Hcn4* expression levels may reflect the attenuated maturation of

transduced NRVMs and failure to downregulate *Hcn4.* In fact, the induction of *Hcn4* by

TBX18 has only been reported using neonatal cardiomyocytes (11, 20, 39) except one

study using juvenile pig heart where improper reference tissue was used as control

(15). In our adult mouse model we did not observed the induction of *Hcn4* expression

at either mRNA or protein levels, nor did we find I_f in the TBX18-expressing

cardiomyocytes. These results are consistent with those reported by Greulich *et al.,* in

which misexpression of *Tbx18* in fetal mouse hearts suppressed the expression of

working cardiomyocyte genes, but did not induce the expression of pacemaker genes

(38). These results are also in line with the previous reports that TBX18 is primarily a

transcriptional repressor rather than an activator (58, 59). Furthermore, the decrease

in cell size was also considered as the result of the transition from mature

cardiomyocytes to pacemaker cells (11), which we did not observe in the TBX18-

expressing cardiomyocytes isolated from mice. Taken together, our results indicate

that ectopic expression of TBX18 in cardiomyocytes is impacting on the expression of

particular genes involved in ion handling that are expressed preferentially in working

cardiomyocytes, thereby altering AP characteristics, but TBX18 is not converting

NRVMs or adult rodent (i.e. mouse and rat) ventricular cardiomyocytes into pacemaker

cells.

Interestingly, abnormal automaticity was observed in isolated TBX18-expressing

cardiomyocytes, despite the lack of I_f. These cells exhibited spontaneous

depolarization, resulting in APs exhibiting early afterdepolarization (EAD)-like

oscillation near the plateau. Since no I_f was detected in these cells, we interpret this

spontaneous depolarization likely to stem from the decrease of I_{K1}, which destabilizes

the resting membrane potential. Indeed, suppression of I_{K1} by itself is enough to generate automaticity in cardiomyocytes, when achieved by overexpression of a dominant negative mutant of *Kcnj2* (36, 60). Of note, in the previous study with AdV-mediated TBX18 overexpression, the amplitude of I_{k1} decrease is much higher than that of I_f increase, suggesting the major cause of the reported Ad-TBX18-induced pacing activity might come from I_{K1} suppression (11). The EAD-like oscillations are likely the result of a decrease in I_K, resulting in long APs with repolarization disorders.

Previous studies have shown TBX18-mediated biological pacing in rats, guinea pigs and pigs when TBX18 was injected at the apex (20), AV junction (15) or His-bundle (16), where the cardiac conduction system is substantially present. Cells within the cardiac conduction system express *Hcn4*, have potential automaticity and display a spindle-shaped morphology distinct from ventricular myocytes (61). Expressing TBX18 in these cells may lead to an increased beating rate due to the intrinsic pacemaker properties of Purkinje cells, and will be responsive to beta-adrenergic stimulation due to the presence of Hcn4. This could potentially explain the discrepancies observed across different studies.

In the presence of beta-adrenergic stimulation, we observed robust biological pacemaker activity in animals injected with AAV-cTnT-Hcn2. To the best of our knowledge, this is the first comprehensive *in vivo* demonstration of an AAV-based biological pacemaker. Functional biological pacemaker engineering via overexpression of ion channels and constitutively active mutants has been extensively studied using AdV vectors (6-9, 36). However, the induction of long-term function with such interventions critically relies on sustained transgene overexpression, which now seems to be feasible with AAV-mediated gene transfer.

In conclusion, ectopic TBX18 expression in cardiomyocytes leads to cytotoxicity and fibrosis, while TBX18 expression at non-cytotoxic levels suppresses several working myocardial genes. Nevertheless, it does not induce a pacemaker gene program, nor does it generate or enhance a biological pacemaker. On the other hand, biological pacemaker outcomes of AAV-cTnT-Hcn2 were robust, provided that isoproterenol was present. This outcome represents a highly encouraging finding, supporting further exploration and testing of Hcn2-based dual transgene strategies (6-8).

Methods

Sex as a biological variable

Our study exclusively examined female mice and rats. It is unknown whether the findings are relevant for male mice and rats.

Vector preparation

AAV production. The transfer plasmids used for AAV production in this study were all derived from pAAV-CMV-GFP (Penn Vector Core, University of Pennsylvania) which contains AAV serotype 2 inverted terminal repeats (ITRs). Coding sequences of human *TBX18* with C-terminal FLAG tag and murine *Hcn2* with N-terminal FLAG tag were synthesized and cloned into AAV transfer plasmids. Primers containing FLAG sequence were sued to amplify GFP, which was then cloned into to the backbone. Oligos with sequence GGCCGGCCGCCATGGCCAGTTTCTACCGGT and GGCCACCGGTAGAAACTGGCCATGGCGGCC were annealed and inserted in front of transgene coding sequence to introduce an upstream open reading frame. AAV6 vectors were produced by double transfection of HEK293T. Low passage HEK293T cells were plated in thirty 145 mm dishes at the density of 1.5×10^7 cells per dish in DMEM-GlutaMax (ThermoFisher Scientific) containing 10% fetal bovine serum (FBS) (Sigma) and 1% penicillin-streptomycin (P/S) (ThermoFisher Scientific). For AAV6 vectors, cells were transfected next day with 33 µg pDP6 and 16 µg AAV transfer plasmids per dish using linear polyethylenimine (PEI) (Polysciences Inc). Medium was replaced during transfection to DMEM containing 1% P/S. Three days after transfection, cells were collected by centrifugation and the medium was concentrated by tangential flow filtration using the ÄKTA flux s system (GE Healthcare) to a final volume of 20 mL. Cells and medium were then combined, frozen and thawed twice followed by DNAseI,

Rnase A, and Benzonase treatment. AAV vectors were purified by iodixanol density-

gradient ultracentrifugation overnight. The AAV-containing fraction was then collected

and concentrated to 1 mL by buffer exchange to PBS containing 0.001% Pluronic F68

using Amicon Ultra-15 100 kDa centrifugal filter units (Millipore). Concentrated AAV

vectors were aliquoted and stored at -80°C until use.

AdV production. AdV-TBX18-GFP vector used in *in vitro* experiment was purchased

from abmgood (Richmond, Canada). AdV vectors used in animal experiments were

produced in Revvity Gene Delivery GmbH (Graefelfing, Germany). The coding

sequence of GFP and N-terminal FLAG tagged human TBX18 was cloned into a pO6

shuttle plasmid and transferred via recombination in a BAC vector containing the

genome of a replication deficient Ad5-based vector deleted in E1/E3 genes as

previously described by Ruszics et al (62). Recombinant viral DNA was released from

the purified BAC-DNA by restriction digest with Pac I and transfected into HEK293 cells.

Viral particles were released via freeze/thaw treatment and amplified on HEK293. Final

release of viral particles was performed via Na-Deoxycholate treatment followed by

CsCl ultracentrifugation and buffer exchange on PD10 columns. Infectious units (IU)

were determined via immunhistochemical detection of the adenoviral hexon protein in

HEK293 cells infected with serial dilutions using anti hexon antibody (Santa Cruz).

NRVM isolation and transduction

NRVMs were isolated from 1- to 2-day-old rats as described previously (63). NRVMs

were cultured in Tung medium, which is M199 containing (in mmol/L) NaCl 137 , KCl

5.4, $CaCl_2$ 1.3, $MgSO_4$ 0.8, $NaHCO_3$ 4.2, KH_2PO_4 0.5,Na_2HPO_4 0.3, and supplemented

with 20 units/100 mL penicillin, 20 µg/100 mL streptomycin, 2 µg/100 mL vitamin B12,

and 2% or 10% fetal bovine serum (FBS). Tung medium containing 10% FBS (10%

Tung) was only sued on the first day. Cells were cultured in monolayers or in low

density single cell preparations on fibronectin-coated plates at 37°C in 5% CO_2. AAV

vector transduction was performed at multiplicity of infection (MOI) = 10,000 in 2%

Tung medium. AdV vector transduction was performed at MOI = 1 in culture media.

Medium was changed to 2% Tung medium the day after transduction, and afterwards

was refreshed every 2 days till analyses.

Western blot

Cells were washed twice with cold Dulbecco's PBS and incubated with RIPA buffer

supplemented with cOMPLETE protease inhibitor (Roche) on ice for 2 min. Cell lysates

were gather using cell scraper, collected and transferred to a microcentrifuge tube. Cell

debris was removed by centrifugation and the supernatant was used for further

analysis. Protein concentration was determined using Pierce BCA Protein Assay Kit

(Thermo Scientific) according to manufacturer's manual. Two and twenty ug of the cell

lysates were loaded onto each well in the HEK and NRVM experiments, respectively.

Cell lysates were separated by 4–20% precast polyacrylamide gel (Bio-Rad) and

transferred to PVDF membranes (Bio-Rad). After blocking with 5% skim milk in Tris-

buffered saline with Tween 20 (5% skim milk/TBST) for one hour at room temperature,

membranes were incubated with Rabbit-anti-FLAG (Sigma, F7425, 1:500) and Mouse-

anti-B-actin antibodies (Sigma, A5441, 1:5000) in 5% skim milk/TBST overnight at 4°C.

Next day, membranes were washed twice with TBST and incubated with Donkey-anti-

Rabbit (Sigma, NA9340, 1:5000) and Sheep-anti-Mouse (Sigma, NA9310, 1:5000)

antibodies for 2 hours at room temperature. Membranes were washed twice with TBST

and once with water, followed by chemiluminescence detection using Amsersham ECL

detection reagents (Cytiva).

In vivo gene delivery in mice

Six- to eight-week old female FVB mice were used for experiments. Mice were injected subcutaneously with buprenorphine (0.075 mg/kg) and carprofen (0.05 mg/kg) for analgesia, at least 30 min prior to surgery. Anesthesia was induced with 4% isoflurane in 1 L/min O_2. Mice were shaved, intubated and placed on a heating mat to maintain body temperature. Subsequently, an analgesic mixture consisting of lidocaine (2 mg/kg) and bupivacaine (3 mg/kg) was applied subcutaneously at the site of the incision. Anesthesia was maintained using ventilation with 2% isoflurane in 1 L/min O_2. Left thoracotomy was performed to expose the apex of the heart at the fourth intercostal space. To inject the vector into the apex, a 10 µL Hamilton syringe fitted with a 31-G needle (13 mm, point style 4) was inserted from the anterior LV towards the apex. Five µL of the viral vector solution was slowly administered at each injection site to administer a total volume of 20 µL viral vectors containing 1×10^{11} viral genomes. Mice were randomly selected to receive control or experimental vectors. The thoracotomy and skin were closed with a C-1 12 mm cutting needle with a 6-0 silicone coated braided silk wire (Sofsilk, Covidien). Post-surgery analgesia consisted of 4 days of *ad libitum* carprofen (Rimadyl Cattle, 0.06 mg/mL) in drinking water and wet food.

In vivo gene delivery in rats

Six-month old female Sprague Dawley rats were used for experiments. Anesthesia was induced with 5% isoflurane in 2 L/min O_2. Rats were shaved, intubated and placed on a heating mat to maintain body temperature. Meloxicam (5 mg/kg) and Buprenorphine SR (1 mg/kg) were delivered subcutaneously for analgesia. Anesthesia was maintained using ventilation with 2% isoflurane in 2 L/min O_2. Partial right thoracotomy was performed to create an ambulatory complete AV-block model, as previously reported.(64) Briefly, the right atrial appendage was ligated and retracted

to expose the AV node region with its characteristic fat pad. Monopolar electrosurgical current was delivered subepicardially to the AV nodal region via a sharp needle to ablate the region. Monopolar electrosurgical current was delivered subepicardially to the AV nodal region via a sharp needle to ablate the region at five watts for 60 sec. If AV block was not generated, the current delivery was repeated under the same condition. ECG was taken one week after to confirm stable complete AV-block. Upon validating stable complete AV-block, rats were randomly selected to receive control or experimental vectors. A partial left thoracotomy was performed to expose the apex of the heart at the fourth intercostal space and inject 1×10^9 viral genomes AdV vectors or 2.5×10^{11} viral genomes AAV vectors, suspended in 50 µL. Control groups were injected with equivalent volume of saline. For rats injected with AAV vectors and its saline control, 3 mg/kg rapamycin was intraperitoneally injected every other day to control the immune response and improve the transduction efficiency. ECGs were recorded for 60 minutes one week after vector injection. Isoproterenol was injected intraperitoneally at a dose of 1 mg/kg, 15 min after the start of recording.

RNA isolation, cDNA synthesis, Reverse-transcription PCR and RNA sequencing

Cells were collected in lysis buffer RP1 (Macherey-Nagel). Tissues were stored immediately in RNAlater, and at a later timepoint homogenized using tissue homogenizer. RNA was isolated using Nucleospin RNA kit (Macherey-Nagel) according to the manufacturer's protocol. For RT-PCR, complementary DNA library was transcribed from 500-1000 ng total RNA with oligo-dT primers (125 µmol/L) and the Superscript II system (Invitrogen). RT-qPCR was performed using the LightCycler 480 Real-Time PCR system (Roche). For comparison across different treatment groups, relative expression of the transgene mRNA was normalized to the geomean

of reference genes *Hprt1* and *Rpl4*. Fold change was calculated over the mean of the

control group. For *Hcn4* expression over different time points, relative expression of

the transgene mRNA was normalized to the reference gene *Hprt1* and fold change was

calculated over day 0. For RNA sequencing, cDNA was prepared with KAPA mRNA

Hyperprep Kit (Roche, KK8580). Total RNA (1 µg) was purified and sheared into small

fragments followed by cDNA synthesis and ligation of the adapter. The quality and size

distribution of the cDNA library templates were validated with Agilent DNA 1000 on

2100 Bioanalyzer (Agilent Technologies). cDNA samples were pooled per lane of

paired-end 150 bp sequencing on a NovaSeq X Plus instrument (Ilumina).

Differential expression analysis

Reads　　　were　　　trimmed　　　using　　　Trim　　　Galore!

(https://https://github.com/FelixKrueger/TrimGalore.com/fenderglass/Flye)　　　and

mapped to rn6 build of the rat transcriptome or mm10 build of the mouse transcriptome

using STAR (65). Differential expression analysis was performed using DESeq2 (66).

Unsupervised hierarchical clustering was performed on genes differentially expressed

(between X, Y and Y) using the R package pheatmap, version 1.0.12. (http://cran.r-

project.org/web/packages/pheatmap/index.html). Gene Ontology (GO) terms and

Kegg pathways were found using DAVID (67). For the comparison to Goodyer et al.,

the fold change of insignificantly differentially expressed genes were set to 1 where

adjusted p-value < 0.01 was considered significant.

Histological staining

Immunofluorescence staining. Hearts were harvested and fixed in 4% PFA

overnight, embedded in paraffin and sectioned at 7 µm thickness. Sections were

deparaffinized and dehydrated by a series of ascending ethanol concentrations. For

antigen retrieval, sections were boiled in unmasking solution (H3300, Vector). Sections were blocked in 4% bovine serum albumin (BSA) and incubated in 2% BSA with nuclear staining and primary antibodies. The primary antibodies used were: chicken anti-GFP (1:500, Aves Labs, GFP-1020), rabbit anti-FLAG (1:200, Millipore, F7425), Goat anti-TBX18 (1:50, Santa Cruz, SC-17869) and DAPI (1 μg/mL, Sigma, D9542). Fluorescence images were acquired using a Leica DM6000 fluorescence microscope or a Leica TCS SP8 confocal microscope (Leica Microsystems).

Picrosirius red staining. Hearts were harvested and fixed in 4% PFA overnight, embedded in paraffin and sectioned at 7 μm thickness. Sections were deparaffinized and dehydrated by a series of ascending ethanol concentrations and stained with picrosirius red. Images were acquired using a Leica DM5000 fluorescence microscope (Leica Microsystems). Image analysis was performed on 4 non-contiguous sections from the anterior of the heart ImageJ. Total area of the left ventricle and the area of fibrosis were measured from which the percentage of fibrosis was calculated.

Patch clamp experiments

Cell preparation and data acquisition. Left ventricular cardiomyocytes of mice were isolated by an enzymatic dissociation procedure (68). Cells were stored at room temperature for at least 45 min in a modified Tyrode's solution containing (in mmol/L): NaCl 140, KCl 5.4, $CaCl_2$ 1.8, $MgCl_2$ 1.0, glucose 5.5, HEPES 5.0; pH 7.4 (NaOH) before use. Cells were put into a recording chamber on the stage of an inverted microscope (Nikon Diaphot, Melville, New York, NY, USA), and single GFP-positive quiescent mouse cardiomyocytes with smooth surfaces were selected for electrophysiological measurements. Action potentials (APs) and net membrane currents were recorded with the amphotericin-perforated patch-clamp technique, while the sodium current (I_{Na}) was measured with the ruptured patch clamp, using an

Axopatch 200B amplifier (Molecular Devices, Sunnyvale, CA, USA). Pipettes (resistance 2–3 MΩ) were pulled from borosilicate glass capillaries (TW100F-3; World Precision Instruments) and filled with solutions as indicated below. Voltage control, data acquisition, and analysis were realized with custom software (69). Signals were low-pass-filtered with a cutoff of 5 kHz and digitized at 40 kHz (APs), 5 kHz (net currents), and 20 kHz (I_{Na}). Potentials were corrected for the calculated liquid junction potential.(70) Cell membrane capacitance (C_m) was estimated by dividing the time constant of the decay of the capacitive transient in response to 5 mV hyperpolarizing voltage clamp steps from –40 mV by the series resistance (69). For net currents and I_{Na} measurements, C_m and series resistance were compensated for by at least 60 and 80%, respectively.

AP measurements. APs were recorded at 36 ± 0.2°C using the modified Tyrode's solution as bath solution. Pipette solution contained (in mmol/L): K-gluconate 125, KCl 20, NaCl 5.0, amphotericin-B 0.44, HEPES 10, and pH 7.2 (KOH). The single cells from dtTBX18-GFP positive cells were spontaneously beating. We measured spontaneously APs as well as APs APs elicited at an overdrive stimulation of 4 Hz by 3-ms, ≈20–40% suprathreshold current pulses, through the patch pipette. Averages AP parameters of stimulated cells were taken from 10 consecutive APs and we analyzed the maximal diastolic potential (MDP), maximum AP upstroke velocity (V_{max}), AP amplitude (APA), AP plateau potential (APplat), and APD at 20, 50, and 90% repolarization (APD_{20}, APD_{50}, and APD_{90}, respectively). Parameters from 10 consecutive APs were averaged.

Net membrane current measurements. APs measurements were followed by voltage clamp experiments to determine which changes in currents underlay the AP differences between WT and TBX18 overexpressing cells. To make a direct

comparison between MDP and background K^+ currents within one cell, and to ensure

that AP shapes and the remaining viable cardiomyocytes in the recording chamber

stay undistorted for biophysical analysis, these measurements were performed without

specific channel blockers or modified solutions. Membrane currents were examined by

500 ms voltage-clamp steps every 2 s to membrane potentials ranging from -100 to

+50 mV from a holding potential of -40 mV. The inward rectifier K^+ current (I_{K1}) and

delayed rectifier K^+ current (I_K) were defined as the quasi steady-state current at the

end of the voltage-clamp steps at potentials negative or positive to -30 mV, respectively.

The transient outward K^+ currents ($I_{to's}$) was defined as the difference between the peak

inward current and the current amplitude at the end of the 500 ms depolarizing voltage

clamp step. All currents were normalized for C_m. Presence of the hyperpolarization-

activated, funny current (I_f) was tested with 2000 ms voltage-clamp steps every 5 s to

membrane potentials ranging from -120 to -40 mV from a holding potential of -40 mV.

Differences between the current at the begin and the current at the end of the

hypolarizing voltage clamp step was defined as I_f.

I_{Na} measurements. *I_{Na} was* characterized at room temperature using a bath solution

containing (in mmol/L): NaCl 7, CsCl 133, $CaCl_2$ 1.8, $MgCl_2$ 1.2, glucose 11.0, HEPES

5.0, and pH 7.4 (CsOH). Nifedipine (5 µmol/L) was added to block the L-type calcium

current. Pipettes were filled with solution containing (in mmol/L): NaCl 3, CsCl 133,

$MgCl_2$ 2.0, Na_2ATP 2.0, TEA-Cl 2.0, EGTA 10.0, HEPES 5.0, and pH 7.2 (CsOH). I_{Na}

amplitudes and voltage dependency of I_{Na} activation were measured by 50 ms

depolarizing voltage clamp steps from a holding potential of -120 mV. A two-step

protocol consisting of series of 500 ms pulses between -140 and 40 mV from a holding

potential of -120 mV, followed by a second 50 ms step to -20 mV, was used to establish

the voltage dependency of I_{Na} inactivation. Cycle lengths of the protocols were 5 s. I_{Na}

was defined as the difference between peak current and steady-state current and I_{Na}

density was calculated by dividing current amplitudes by C_m. Voltage dependence of

activation and inactivation were determined by fitting a Boltzmann function

$(y=[1+\exp\{(V-V1/2)/k\}]^{-1})$ to the individual plots, where $V_{1/2}$ is the voltage of half-

maximal (in)activation and k the slope factor (in mV).

Statistical analysis

GraphPad Prism version 10 software (GraphPad Software Inc.) was used for statistical

analysis. Numerical data are presented as mean ± standard error of the mean (SEM)

of at least three biological replicates. Two independent groups were compared by

Mann-Whitney test and multiple independent groups were compared by analysis of

variance (ANOVA) followed by the *post-hoc* Holm-Šídák test or Fisher's LSD test for

multiple comparisons. Categorical data were presented as absolute values and

percentages. Categorical data were compared by Fisher's exact test. All tested were

performed two-sided and statistical significance was considered for $p < 0.05$.

Study approval

Amsterdam UMC (the Netherlands): All mice were treated in accordance with the EC

Directive 2010/63/EU of the European Parliament and of the Council on the protection

of animals used for scientific purposes and using protocols approved and monitored

by the Animal Welfare Body of Amsterdam UMC.

Naason Science Inc. (Republic of Korea): Rat study was carried out according to Korea

Food and Drug Administration guidelines for the care and use of laboratory animals

and approved by the Institutional Animal Care and Use Committee.

Data availability

The datasets generated or analyzed during the studies reported herein are available

in the Supporting Data Values file provided in a single Excel sheet, and repeats and

supporting data sets can be obtained from the corresponding author on reasonable

request.

Author contribution

JW designed and conducted experiments, analyzed data, drafted the manuscript, and oversaw the project. MRR, MK, ARB, RNV, RS, HYL and KvD contributed to the conduct of experiments. KHP and LCP oversaw the conduct of rat experiments. SS, CT, EE and AE contributed to the vector production. OFK, KN and HLT designed experiments and revised the manuscript. AOV designed, conducted experiments, and analyzed data. VM and GJJB designed experiments, drafted and revised the manuscript, and oversaw the project.

Acknowledgement

This work is supported by funding from The European Research Council starting grant 714866 and associated proof-of-concept grant 899422 (to G.J.J.B.), Health Holland LentiPace II (to G.J.J.B. and H.L.T.), Horizon 2020 Eurostars (E114245 and E115484; to G.J.J.B. and V.M.C.); Dutch Research Council Open Technology Program 18485 (to H.L.T. and G.J.J.B.); The European Innovation Council (EIC; TRANSITION - Project 101099608 – TRACTION; to V.M.C. and G.J.J.B.); and The Dutch Research Council OCENW.GROOT.2019.029 (to V.M.C.). We would like to thank Otto J. Mulleners for his help in RNA sequencing analysis.

Reference

1. Rosen MR, Robinson RB, Brink PR, and Cohen IS. The road to biological pacing. *Nat Rev Cardiol.* 2011;8(11):656-66.
2. Amin MS, Matchar DB, Wood MA, and Ellenbogen KA. Management of recalled pacemakers and implantable cardioverter-defibrillators: a decision analysis model. *JAMA.* 2006;296(4):412-20.
3. Rosen MR. Gene therapy and biological pacing. *N Engl J Med.* 2014;371(12):1158-9.
4. Boink GJ, and Robinson RB. Gene therapy for restoring heart rhythm. *J Cardiovasc Pharmacol Ther.* 2014;19(5):426-38.
5. Plotnikov AN, Bucchi A, Shlapakova I, Danilo P, Jr., Brink PR, Robinson RB, et al. HCN212-channel biological pacemakers manifesting ventricular tachyarrhythmias are responsive to treatment with I(f) blockade. *Heart Rhythm.* 2008;5(2):282-8.
6. Boink GJ, Nearing BD, Shlapakova IN, Duan L, Kryukova Y, Bobkov Y, et al. Ca(2+)-stimulated adenylyl cyclase AC1 generates efficient biological pacing as single gene therapy and in combination with HCN2. *Circulation.* 2012;126(5):528-36.
7. Boink GJ, Duan L, Nearing BD, Shlapakova IN, Sosunov EA, Anyukhovsky EP, et al. HCN2/SkM1 gene transfer into canine left bundle branch induces stable, autonomically responsive biological pacing at physiological heart rates. *J Am Coll Cardiol.* 2013;61(11):1192-201.
8. Cingolani E, Yee K, Shehata M, Chugh SS, Marban E, and Cho HC. Biological pacemaker created by percutaneous gene delivery via venous catheters in a porcine model of complete heart block. *Heart Rhythm.* 2012;9(8):1310-8.
9. Qu J, Plotnikov AN, Danilo P, Jr., Shlapakova I, Cohen IS, Robinson RB, et al. Expression and function of a biological pacemaker in canine heart. *Circulation.* 2003;107(8):1106-9.
10. Bakker ML, Boink GJ, Boukens BJ, Verkerk AO, van den Boogaard M, den Haan AD, et al. T-box transcription factor TBX3 reprogrammes mature cardiac myocytes into pacemaker-like cells. *Cardiovasc Res.* 2012;94(3):439-49.
11. Kapoor N, Liang W, Marban E, and Cho HC. Direct conversion of quiescent cardiomyocytes to pacemaker cells by expression of Tbx18. *Nat Biotechnol.* 2013;31(1):54-62.
12. Wiese C, Grieskamp T, Airik R, Mommersteeg MT, Gardiwal A, de Gier-de Vries C, et al. Formation of the sinus node head and differentiation of sinus node myocardium are independently regulated by Tbx18 and Tbx3. *Circ Res.* 2009;104(3):388-97.
13. Mommersteeg MT, Dominguez JN, Wiese C, Norden J, de Gier-de Vries C, Burch JB, et al. The sinus venosus progenitors separate and diversify from the first and second heart fields early in development. *Cardiovasc Res.* 2010;87(1):92-101.
14. Hoogaars WM, Engel A, Brons JF, Verkerk AO, de Lange FJ, Wong LY, et al. Tbx3 controls the sinoatrial node gene program and imposes pacemaker function on the atria. *Genes Dev.* 2007;21(9):1098-112.
15. Hu YF, Dawkins JF, Cho HC, Marban E, and Cingolani E. Biological pacemaker created by minimally invasive somatic reprogramming in pigs with complete heart block. *Sci Transl Med.* 2014;6(245):245ra94.
16. Dawkins JF, Hu YF, Valle J, Sanchez L, Zheng Y, Marban E, et al. Antegrade Conduction Rescues Right Ventricular Pacing-Induced Cardiomyopathy in Complete Heart Block. *J Am Coll Cardiol.* 2019;73(13):1673-87.
17. Choudhury M, Black N, Alghamdi A, D'Souza A, Wang R, Yanni J, et al. TBX18 overexpression enhances pacemaker function in a rat subsidiary atrial pacemaker model of sick sinus syndrome. *J Physiol.* 2018;596(24):6141-55.

18. Hendrickx R, Stichling N, Koelen J, Kuryk L, Lipiec A, and Greber UF. Innate immunity to adenovirus. *Hum Gene Ther.* 2014;25(4):265-84.

19. Shirley JL, de Jong YP, Terhorst C, and Herzog RW. Immune Responses to Viral Gene Therapy Vectors. *Mol Ther.* 2020;28(3):709-22.

20. Wolfson DW, Kim NK, Lee KH, Beyersdorf JP, Langberg JJ, Fernandez N, et al. Transient pacing in pigs with complete heart block via myocardial injection of mRNA coding for the T-box transcription factor 18. *Nat Biomed Eng.* 2024.

21. Li C, and Samulski RJ. Engineering adeno-associated virus vectors for gene therapy. *Nat Rev Genet.* 2020;21(4):255-72.

22. Kupatt C, Hinkel R, Pfosser A, El-Aouni C, Wuchrer A, Fritz A, et al. Cotransfection of vascular endothelial growth factor-A and platelet-derived growth factor-B via recombinant adeno-associated virus resolves chronic ischemic malperfusion role of vessel maturation. *J Am Coll Cardiol.* 2010;56(5):414-22.

23. Beeri R, Chaput M, Guerrero JL, Kawase Y, Yosefy C, Abedat S, et al. Gene delivery of sarcoplasmic reticulum calcium ATPase inhibits ventricular remodeling in ischemic mitral regurgitation. *Circ Heart Fail.* 2010;3(5):627-34.

24. Raake PW, Hinkel R, Muller S, Delker S, Kreuzpointner R, Kupatt C, et al. Cardio-specific long-term gene expression in a porcine model after selective pressure-regulated retroinfusion of adeno-associated viral (AAV) vectors. *Gene Ther.* 2008;15(1):12-7.

25. Weber C, Neacsu I, Krautz B, Schlegel P, Sauer S, Raake P, et al. Therapeutic safety of high myocardial expression levels of the molecular inotrope S100A1 in a preclinical heart failure model. *Gene Ther.* 2014;21(2):131-8.

26. Liu S, Li K, Wagner Florencio L, Tang L, Heallen TR, Leach JP, et al. Gene therapy knockdown of Hippo signaling induces cardiomyocyte renewal in pigs after myocardial infarction. *Sci Transl Med.* 2021;13(600).

27. Gabisonia K, Prosdocimo G, Aquaro GD, Carlucci L, Zentilin L, Secco I, et al. MicroRNA therapy stimulates uncontrolled cardiac repair after myocardial infarction in pigs. *Nature.* 2019;569(7756):418-22.

28. Philippidis A. BioMarin's ROCTAVIAN Wins Food and Drug Administration Approval As First Gene Therapy for Severe Hemophilia A. *Hum Gene Ther.* 2023;34(15-16):665-8.

29. Kim NK, Wolfson D, Fernandez N, Shin M, and Cho HC. A rat model of complete atrioventricular block recapitulates clinical indices of bradycardia and provides a platform to test disease-modifying therapies. *Sci Rep.* 2019;9(1):6930.

30. Fan J, Kim NK, Fernandez N, Kim TY, Li J, Wolfson D, et al. Inhibition of Tgfβ signaling enables durable ventricular pacing by TBX18 gene transfer. *BioRxiv.* 2022.

31. Ferreira JP, Overton KW, and Wang CL. Tuning gene expression with synthetic upstream open reading frames. *Proc Natl Acad Sci U S A.* 2013;110(28):11284-9.

32. Wang XQ, and Rothnagel JA. 5'-untranslated regions with multiple upstream AUG codons can support low-level translation via leaky scanning and reinitiation. *Nucleic Acids Res.* 2004;32(4):1382-91.

33. Kapoor N, Galang G, Marban E, and Cho HC. Transcriptional suppression of connexin43 by TBX18 undermines cell-cell electrical coupling in postnatal cardiomyocytes. *J Biol Chem.* 2011;286(16):14073-9.

34. Zhao S, Wu H, Xia W, Chen X, Zhu S, Zhang S, et al. Periostin expression is upregulated and associated with myocardial fibrosis in human failing hearts. *J Cardiol.* 2014;63(5):373-8.

35. Minicucci MF, Santos PP, Rafacho BP, Goncalves AF, Ardisson LP, Batista DF, et al. Periostin as a modulator of chronic cardiac remodeling after myocardial infarction. *Clinics (Sao Paulo).* 2013;68(10):1344-9.

36. Miake J, Marban E, and Nuss HB. Biological pacemaker created by gene transfer. *Nature.* 2002;419(6903):132-3.

37. Li Y, Wang K, Li Q, Hancox JC, and Zhang H. Reciprocal interaction between IK1 and If in biological pacemakers: A simulation study. *PLoS Comput Biol.* 2021;17(3):e1008177.

38. Greulich F, Trowe MO, Leffler A, Stoetzer C, Farin HF, and Kispert A. Misexpression of Tbx18 in cardiac chambers of fetal mice interferes with chamber-specific developmental programs but does not induce a pacemaker-like gene signature. *J Mol Cell Cardiol.* 2016;97:140-9.

39. Foster DB, Gu JM, Kim EH, Wolfson DW, O'Meally R, Cole RN, et al. Tbx18 Orchestrates Cytostructural Transdifferentiation of Cardiomyocytes to Pacemaker Cells by Recruiting the Epithelial-Mesenchymal Transition Program. *J Proteome Res.* 2022;21(10):2277-92.

40. Liang X, Wang G, Lin L, Lowe J, Zhang Q, Bu L, et al. HCN4 dynamically marks the first heart field and conduction system precursors. *Circ Res.* 2013;113(4):399-407.

41. Strohman RC, Micou-Eastwood J, Glass CA, and Matsuda R. Human fetal muscle and cultured myotubes derived from it contain a fetal-specific myosin light chain. *Science.* 1983;221(4614):955-7.

42. Horsthuis T, Houweling AC, Habets PE, de Lange FJ, el Azzouzi H, Clout DE, et al. Distinct regulation of developmental and heart disease-induced atrial natriuretic factor expression by two separate distal sequences. *Circ Res.* 2008;102(7):849-59.

43. Cerbai E, Pino R, Sartiani L, and Mugelli A. Influence of postnatal-development on I(f) occurrence and properties in neonatal rat ventricular myocytes. *Cardiovasc Res.* 1999;42(2):416-23.

44. Giovou AE, Gladka MM, and Christoffels VM. The Impact of Natriuretic Peptides on Heart Development, Homeostasis, and Disease. *Cells.* 2024;13(11).

45. Gudbjartsson DF, Holm H, Sulem P, Masson G, Oddsson A, Magnusson OT, et al. A frameshift deletion in the sarcomere gene MYL4 causes early-onset familial atrial fibrillation. *Eur Heart J.* 2017;38(1):27-34.

46. Vedantham V, Galang G, Evangelista M, Deo RC, and Srivastava D. RNA sequencing of mouse sinoatrial node reveals an upstream regulatory role for Islet-1 in cardiac pacemaker cells. *Circ Res.* 2015;116(5):797-803.

47. Goodyer WR, Beyersdorf BM, Paik DT, Tian L, Li G, Buikema JW, et al. Transcriptomic Profiling of the Developing Cardiac Conduction System at Single-Cell Resolution. *Circ Res.* 2019;125(4):379-97.

48. Padget RL, Zeitz MJ, Blair GA, Wu X, North MD, Tanenbaum MT, et al. Acute Adenoviral Infection Elicits an Arrhythmogenic Substrate Prior to Myocarditis. *Circ Res.* 2024;134(7):892-912.

49. Kreppel F, and Hagedorn C. Capsid and Genome Modification Strategies to Reduce the Immunogenicity of Adenoviral Vectors. *Int J Mol Sci.* 2021;22(5).

50. Amin AS, Asghari-Roodsari A, and Tan HL. Cardiac sodium channelopathies. *Pflugers Arch.* 2010;460(2):223-37.

51. Xu J, Lamouille S, and Derynck R. TGF-beta-induced epithelial to mesenchymal transition. *Cell Res.* 2009;19(2):156-72.

52. Chen PY, Qin L, and Simons M. TGFbeta signaling pathways in human health and disease. *Front Mol Biosci.* 2023;10:1113061.

53. Prasad KM, Xu Y, Yang Z, Acton ST, and French BA. Robust cardiomyocyte-specific gene expression following systemic injection of AAV: in vivo gene delivery follows a Poisson distribution. *Gene Ther.* 2011;18(1):43-52.

54. Nguyen HX, Wu T, Needs D, Zhang H, Perelli RM, DeLuca S, et al. Engineered bacterial voltage-gated sodium channel platform for cardiac gene therapy. *Nat Commun.* 2022;13(1):620.

55. Konkalmatt PR, Beyers RJ, O'Connor DM, Xu Y, Seaman ME, and French BA. Cardiac-selective expression of extracellular superoxide dismutase after systemic injection of adeno-associated virus 9 protects the heart against post-myocardial infarction left ventricular remodeling. *Circ Cardiovasc Imaging.* 2013;6(3):478-86.

56. Chou PC, Liu CM, Weng CH, Yang KC, Cheng ML, Lin YC, et al. Fibroblasts Drive Metabolic Reprogramming in Pacemaker Cardiomyocytes. *Circ Res.* 2022;131(1):6-20.

57. van Eif VWW, Devalla HD, Boink GJJ, and Christoffels VM. Transcriptional regulation of the cardiac conduction system. *Nat Rev Cardiol.* 2018;15(10):617-30.

58. Rivera-Reyes R, Kleppa MJ, and Kispert A. Proteomic analysis identifies transcriptional cofactors and homeobox transcription factors as TBX18 binding proteins. *PLoS One.* 2018;13(8):e0200964.

59. Farin HF, Bussen M, Schmidt MK, Singh MK, Schuster-Gossler K, and Kispert A. Transcriptional repression by the T-box proteins Tbx18 and Tbx15 depends on Groucho corepressors. *J Biol Chem.* 2007;282(35):25748-59.

60. Sekar RB, Kizana E, Cho HC, Molitoris JM, Hesketh GG, Eaton BP, et al. IK1 heterogeneity affects genesis and stability of spiral waves in cardiac myocyte monolayers. *Circ Res.* 2009;104(3):355-64.

61. Ideker RE, Kong W, and Pogwizd S. Purkinje fibers and arrhythmias. *Pacing Clin Electrophysiol.* 2009;32(3):283-5.

62. Ruzsics Z, Lemnitzer F, and Thirion C. Engineering adenovirus genome by bacterial artificial chromosome (BAC) technology. *Methods Mol Biol.* 2014;1089:143-58.

63. Rohr S, Scholly DM, and Kleber AG. Patterned growth of neonatal rat heart cells in culture. Morphological and electrophysiological characterization. *Circ Res.* 1991;68(1):114-30.

64. Kim D, Lim K, Kim ST, Yoon SH, Kim K, Ryu SM, et al. Genome-wide target specificities of CRISPR RNA-guided programmable deaminases. *Nat Biotechnol.* 2017;35(5):475-80.

65. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics.* 2013;29(1):15-21.

66. Love MI, Huber W, and Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 2014;15(12):550.

67. Huang da W, Sherman BT, and Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc.* 2009;4(1):44-57.

68. Man JCK, Bosada FM, Scholman KT, Offerhaus JA, Walsh R, van Duijvenboden K, et al. Variant intronic enhancer controls *SCN10A-short* expression and heart conduction. *Circulation.* 2021;144(3):229-42.

69. Verkerk AO, and Wilders R. Injection of I_{K1} through dynamic clamp can make all the difference in patch-clamp studies on hiPSC-derived cardiomyocytes. *Front Physiol.* 2023;14:1326160.

70. Barry PH, and Lynch JW. Liquid junction potentials and small cell effects in patch-clamp analysis. *J Membr Biol.* 1991;121(2):101-17.

Figures

Figure 1 Injection of adenoviral vectors (AdV) leads to ectopic pacing in a complete atrioventricular block (AV-block) rat model.

(**A**) An example ECG tracing showing complete AV-block in rats after AV node ablation. P wave and QRS complex are indicated on top. (**B**) Experimental design. (**C**) Immunofluorescence staining images of GFP and TBX18 in rat hearts injected with AdV-GFP and AdV-TBX18, respectively. Scale bar = 50 μm. (**D**) Example ECG tracings of rats injected with saline, AdV-GFP and AdV-TBX18, before and after the administration of isoproterenol. (**E**) Percentage of ectopic pacing and (**F**) average heart rate in rats injected with saline (n = 4), AdV-GFP (n = 6) and AdV-TBX18 (n = 6). (**G**) Picrosirius red staining images and (**H**) fibrosis quantification of rat hearts injected with

saline (n = 4), AdV-GFP (n = 6) and AdV-TBX18 (n = 6). Scale bar = 300 μm. Data are
shown as mean ± SEM. Data were compared using two-way ANOVA (E & F) or one
way ANOVA (H) with *post-hoc* Fisher's LSD test . $*p < 0.05$; $**p < 0.01$; $***p < 0.001$;
ns, not significant.

Figure 2 AAV-mediated TBX18 overexpression induces cardiac fibrosis in
mouse hearts.

(**A**) Macroscopy images of mouse hearts injected with AAV vectors overexpressing
GFP or TBX18. Black star indicates the fibrotic scar at the injection site. (**B**) Incidence
of hearts with fibrotic scar. Number of animals with fibrotic scar and total number of
animals studied are presented within the associated bar charts. P value was calculated
using Fisher's exact test. (**C**) Picrosirius red staining images of mouse hearts injected
with AAV-CMV-TBX18 at 4, 7 and 28 days post injection (dpi). Scale bar = 1 mm. (**D**)
Immunofluorescence staining images of mouse hearts injected with AAV-CMV-TBX18
at 7 and 28 dpi. Scale bar = 100 μm.

Figure 3 Dose-titrated expression cassette achieved low yet functional TBX18 expression.

(**A**) Schematic representation of expression cassette with and without upstream open reading frame (uORF). Translation start sites are made bold. (**B**) Direct fluorescence images of HEK cells transfected with GFP and uORF-GFP plasmids. Yellow arrows indicate the GFP-positive cells. (**C**) Percentage of GFP-positive cells and (**D**) mean fluorescence intensity of HEK cells transfected with GFP or uORF-GFP plasmids (n = 3). (**E**) Example western blot and (**F**) quantification of TBX18 protein expression in HEK cells transfected with TBX18 or uORF-TBX18 plasmids (n = 3). (**G**) AAV vectors used for transduction and western blot of TBX18 protein expression in transduced cardiomyocytes. (**H**) Quantification of TBX18 protein expression in cardiomyocytes transduced with various TBX18 vectors (n = 3). (**I**) Expression of pacemaker genes and (**J**) working myocardial genes in cardiomyocytes transduced with various TBX18 vectors (n = 6) and one GFP vector (n = 5). Data are presented as mean ± SEM. Data were compared using Mann-Whitney test (H), one-way ANOVA with *post-hoc* Holm-Šídák test (C, D & F) or two-way ANOVA with *post-hoc* Holm-Šídák test (I & J). *p < 0.05; **p < 0.01; ns, not significant.

Figure 4 Long-term TBX18 expression delivered by cTnT-uORF-TBX18 vector does not induce cardiac fibrosis.

(**A**) AAV vectors used for intramyocardial injection. (**B**) Picrosirius red staining images of GFP- or TBX18- injected mouse hearts 4 weeks post injection (wpi). Black bar at the top indicates the anticipated transgene expression level. Scale bar = 1 mm (upper) or 200 μm (lower). (**C**) Quantification of the cardiac fibrosis in the left ventricle of the injected hearts (n = 3). (**D**) Picrosirius red staining images of GFP- or TBX18-injected mouse hearts 8 wpi expressing GFP or TBX18. Black bar at the top indicates the anticipated transgene expression level. Scale bar = 1 mm (upper) or 200 μm (lower). (**E**) Quantification of cardiac fibrosis in the left ventricle of the injected hearts (n = 3). Data are presented as mean ± SEM. Data were compared using one-way ANOVA with *post-hoc* Fisher's LSD test. **p < 0.01; ns, not significant.

Figure 5 TBX18 expression at non-cytotoxic level suppresses working myocardial genes but does not induce pacemaker gene *Hcn4*.

(**A**) Experimental design for histological analyses. (**B**) Immunofluorescence staining images of Hcn4 in TBX18-injected animals 4 and 8 weeks post injection (wpi). Scale bar = 50 μm. (**C**) Immunofluorescence staining image of Cx43 at injection sites of GFP- or TBX18- injected animals 4 wpi. Scale bar = 50 μm. (**D**) Experimental design for molecular analyses. (**E**) Expression level of human *TBX18*, *Tbx18*, *Gja1* and *Postn* determined by quantitative PCR (n = 4 for Nluc and n = 5 for TBX18). (**F**) Expression level of *Scn5a*, *Kcnj2*, *Kcnj5*, *Kcnd2*, *Hcn4* and *Nppa* determined by quantitative PCR (n = 4 for Nluc and n = 5 for TBX18). Data are shown as mean ± SEM. Data were compared using Mann-Whitney test. *p < 0.05.

Figure 6 Long-term TBX18 expression at non-cytotoxic level leads to ionic remodeling of mouse cardiomyocytes.

(**A**) Example action potentials (APs) recorded from TBX18-expressing and control cardiomyocytes (spontaneous / not stimulated). (**B**) Incidence of spontaneous activity. Number of cells with spontaneous activity and total number of cells studied are presented within the associated bar charts. (**C**) Example APs recorded from TBX18-expressing and control cardiomyocyte paced at 4 Hz. Inset: AP upstroke velocities. (**D**) Average AP parameters. RMP: resting membrane potential; APA: AP amplitude; $APD_{20/50/90}$: AP durations at 20%/50%/90% repolarization; V_{max}: maximal AP upstroke velocity (n = 8). (**E**) Example current traces at -100 and +50 mV of TBX18-expressing and control cardiomyocytes from a generic voltage clamp. (**F**) Current-voltage (I-V) relationships of the quasi steady-state current. (**G**) Current densities of the inwardly rectifying K$^+$ current (I_{K1}) measured at -100 mV. (**H**) Current densities of the delayed rectifier outward K$^+$ current (I_K) measured at 50 mV. (**I**) I-V relationships of the hyperpolarization-activated funny current (I_f). (**J**) I-V relationships of Na$^+$ current (I_{Na}). (**K**) Peak I_{Na} density measured at -40 mV. (**L**) Cell capacitance reflecting cell size. (F-L) n = 4 for Nluc and n = 5 for TBX18. Data are shown as mean ± SEM. Data were

1   compared using Fisher's exact test (B) or Mann–Whitney test (D, G, H, K, L). *p < 0.05;

2   **p < 0.01; ns, not significant.

Figure 7 AAV-mediated expression of Hcn2 generates robust ectopic pacing in the presence of isoproterenol while AAV-mediated expression of TBX18 does not.

(**A**) Experimental design. (**B**) Expression of TBX18 and Hcn2 in rat hearts injected with AAV vector expressing TBX18 and Hcn2, respectively. Scale bar = 50 μm. (**C**) Example ECG tracings of rats injected with saline or various AAV vectors before and after the administration of isoproterenol. (**D**) Percentage of ectopic pacing and (**E**) average heart rate in rats injected with saline or various AAV vectors . (**F**) Picrosirius red staining images and (**G**) fibrosis quantification of rat hearts injected with saline, AAV vector

expressing TBX18, AAV vector expressing Hcn2, or both. (D-G) n = 5 for saline, n = 4 for TBX18, n = 6 for Hcn2 and TBX18 + Hcn2. Scale bar = 300 μm. Data are shown as mean ± SEM. Data were compared using two-way ANOVA (D & E) or one-way ANOVA (G) with *post-hoc* Fisher's LSD test. *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$; ns, not significant.

Figure 8. RNA sequencing analysis reveals the transcriptional effects of Hcn2 and TBX18 expression in NRVMs.

(**A**) Principal component analysis of the cardiomyocytes transduced with control, Hcn2 and various TBX18 vectors. (**B**) Unsupervised functional annotation heatmap of NRVMs transduced with control, Hcn2 and various TBX18 vectors. (**C-E**) Volcano plots showing transcripts differentially expressed between NRVMs transduced with (C) control and Hcn2, (D) control and TBX18, and (E) Hcn2 and TBX18. (**F-H**) Scatter plots showing gene expression in NRVMs transduced with TBX18 or Hcn2 in comparison to postnatal day 4 (P4) mouse sinoatrial node (SAN) pacemaker cells. (F) Hcn2-

1   transduced NRVMs from the current study (G) TBX18-transduced NRVMs from the
2   current study. (H) TBX18- transduced NRVMs from Foster et al. (39).