Quantitative mapping of methionine sensitivity to oxidation in the copper-bound PcuC chaperone

Lionel Tarrago[a,b,1,]*, Lise Molinelli[a,c], Maya Belghazi[d], Mathilde Tribout[b,2], David Lemaire[b], Pierre

Legrand[e], Sandrine Grosse[b], David Pignol[b], Monique Sabaty[b], Thierry Tron[c], Pascal Arnoux[b,]*

From:

[a] Aix Marseille Univ, INRAE, BBF, Marseille, France

[b] Aix Marseille Univ, CEA, CNRS, BIAM, Saint Paul-Lez-Durance, France

[c] Aix Marseille Univ, CNRS, Centrale Marseille, ISM2, Marseille, France

[d] Aix Marseille Univ, CNRS, BIP, IMM, Marseille, France

[e] Synchrotron SOLEIL, L'Orme des Merisiers, F-91192, Gif-sur-Yvette, France

[1] Permanent address: Aix Marseille Univ, INRAE, BBF, Marseille, France

[2] Present address: Aix Marseille Univ, CNRS, LCB, IMM, Marseille, France

*Correspondence: Lionel Tarrago (lionel.tarrago@inrae.fr) or Pascal Arnoux (pascal.arnoux@cea.fr)

Keywords: cofactor biosynthesis, copper chaperone, metalloprotein, methionine oxidation, oxaziridine

Abstract

Copper is typically coordinated by histidine, cysteine, or methionine in proteins, and these residues are particularly sensitive to oxidation. However, it remains unclear whether copper-coordinating residues are more prone to oxidation than non-coordinating ones, and how their susceptibility changes between the apo and copper-bound states. The copper chaperone PcuC, important for cytochrome c oxidase assembly in bacteria, contains a canonical binding site composed of two histidines and two methionines ($H51x_nM63x_{22}H86xM88$), as well as a disordered C-terminal extension enriched in methionine and histidine. To quantify methionine oxidation sensitivity in both apo- and Cu-bound PcuC, we used a methionine-specific oxaziridine probe combined with mass spectrometry and compared labeling patterns to those generated by ^{18}O-labeled hydrogen peroxide. We show that methionine residues display distinct oxidation sensitivities in the apoprotein, and that the oxaziridine reacts similarly to $H_2^{18}O_2$. Importantly, this probe enables quantification of methionine oxidation independently of hydroxyl radicals generated by copper-driven Fenton chemistry, which lacks residue specificity. In the copper-bound form, Cu binding strongly alters methionine reactivity, with a marked increase in oxidation of the coordinating Met63 and Met88. Structural analysis revealed that two copper ions occupy the canonical site, while the C-terminal extension does not contribute to coordination. Comparison of structural features and oxidation sensitivity showed that methionine susceptibility correlates with solvent exposure in the folded domain, but with local positive charge in the disordered region. These findings demonstrate that copper coordination modulates methionine oxidation, and that oxaziridine-based probes provide a powerful tool for mapping oxidation sensitivity in (metallo)proteins.

1. Introduction

Metals are essential in life, especially within proteins where they serve as critical cofactors for various biochemical reactions. Copper plays essential roles in biology as redox center in proteins involved in electron transfer, redox catalysis, and metal homeostasis [1]. Copper is commonly coordinated by histidine (His), cysteine (Cys), and methionine (Met) residues, with His and Cys predominantly found in redox-active sites while Met is mostly found in copper transporters and chaperone [2]. Copper can also bind proteins implicated in neurodegenerative diseases, such as α-synuclein and the prion protein [3,4]. The diversity of copper sites and ligands reflects the metal's versatility, but also the need for precise control to ensure correct metalation and prevent oxidative damage. Indeed, the copper-coordinating residues Cys, His, and Met are particularly sensitive to oxidation, leading to radical or oxygenated forms. Oxidants like hydrogen peroxide (H_2O_2) or hypochlorite have distinct reactivity towards amino acids, and the reaction of H_2O_2 with iron or copper in Fenton(-like) reaction generates strongly oxidant hydroxyl radicals [5,6]. While hypochlorite and hydroxyl radicals have broad reactivity and affects most amino acids and the protein backbone, H_2O_2 reacts specifically with Cys and Met [7]. In particular, Met can be oxidized into sulfoxide (MetO) and sulfone ($MetO_2$) [7,8]. Protein oxidation may have diverse consequences, from damage to regulated changes in function or activity. However, little is known about the sensitivity to oxidation of Cys, His, or Met within copper-binding proteins. It is not known whether these residues that are present in copper coordination sites, either in the apo- or copper-bound state, present a susceptibility to oxidation different from non-coordinating counterparts. Understanding the sensitivity of copper proteins to oxidation is crucial, as it can lead to their inactivation, as observed for superoxide dismutase and lytic polysaccharide monooxygenases [9–11]. It can also affect structural properties of α-synuclein and the prion protein, with potential pathologic outcomes [12,13]. In peptides mimicking copper-binding sites of the eukaryotic Ctr1 copper transporter, few evidence indicates that Cys is highly sensitive to oxidation contrary to Met or His [14,15]. Oxidation sensitivity of copper-binding Met in α-synuclein has also been observed, though quantitative data are lacking. In bacteria, such mechanisms remain poorly understood, despite their particular relevance for periplasmic proteins, which

are exposed to copper-induced hydroxyl radical production and rely on detoxification and export systems

[5,16,17]. Consistently, it has been shown that the expression of the methionine sulfoxide reductase (Msr)

P/Q system, that repairs periplasmic protein by reducing oxidized Met [18,19], is induced by the presence

of an excess of copper in the periplasm through the activation of the CusSR two-component system [20].

Bacteria membranes and mitochondria contain the respiratory chain with cytochrome c oxidase (Cox)

which possesses two Cu atoms in the Cu_A center of its periplasmic and catalytic subunit CoxB, necessary

for the reduction of oxygen [21]. Bacterial CoxB requires two chaperones for its assembly: PrrC,

homologous to eukaryotic Sco proteins, and PcuC, a cupredoxin-like protein. This protein is often referred

to as PCu_AC but we use PcuC in this article to follow uniform nomenclature of bacterial proteins. The

sequence of events allowing Cu insertion in CoxB has been evaluated in several bacterial species [22–25].

In the model proposed for *Cereibacter sphaeroides*, PcuC acquires copper from the periplasm and

transfers it to PrrC, which then assembles the Cu_A center with one Cu(I) and one Cu(II) [25]. Cu ions are

sequentially transferred from one coordinating site to another [26,27]. Particularly, CoxB coordinates the

two Cu with two Cys, two His, one Met and one Glu in a $Hx_{34}CxExCx_3Hx_2M$ motif [23,28]. PrrC uses two

Cys and one His in the Cx_3Cx_nH motif to coordinate a Cu(II) ion [23]. It has been shown that copper-

coordinating Cys of both CoxB and PrrC can form disulfide bonds that need to be reduced by the

dedicated thiol-reductase TplA before copper insertion [29]. PcuC is widely conserved among bacteria

and most homologs possess a Cu(I)-binding motif composed of two Met and two His ($Hx_nMx_{22}HxM$) and

have an additional C-terminal extension rich in Met and His, proposed to bind a second copper ion

[22,23,30]. Other types lack this extension or bind copper in a His-brace [23,31]. The *C. sphaeroides*

PcuC is a representative model for studying copper's role in protein oxidation.

In this work, we evaluated the sensitivity of PcuC's Met to oxidation, either in the apo or in the

copper-loaded protein by using two molecules reacting with Met: i) [18]O-labeled hydrogen peroxide, and ii)

an oxaziridine probe (**Figure 1A**). We quantified the level of oxidation of each Met by trypsin proteolysis

and liquid chromatography coupled with tandem mass spectrometry (LC-MS/MS). H_2O_2 is a natural

oxidant potentially encountered by bacteria. In the absence of Cys, as it is the case for PcuC, it allows the

specific oxidation of Met [7,19]. The use of [18]O-labeling avoid the quantification of artifactual Met

oxidation potentially arising in proteomic analysis [32–34]. However, H_2O_2 can react with the metal in the

copper-loaded protein, generating highly reactive hydroxyl radicals that may oxidize peptide bonds and

residues other than Met. To specifically quantify methionine sensitivity to oxidation in the copper-loaded

protein independently of Fenton reaction, we used an oxaziridine probe designed for selective reactivity

with Met [35,36]. This probe operates via nitrogen atom transfer, where the electrophilic nitrogen of the

strained N–O bond reacts with the thioether sulfur, forming a stable sulfimide adduct (**Figure 1B**). We

first assessed Met sensitivity in the apo form using $H_2^{18}O_2$ and then with the oxaziridine probe, allowing

comparison of both approaches. Then, we applied the probe to the copper-loaded protein to evaluate the

impact of Cu coordination on Met oxidation. Finally, we determined the three-dimensional structures of

both apo and copper-bound PcuC and integrated these structural insights with oxidation data to identify

determinants of Met susceptibility to oxidation.

2. Results and Discussion

2.1. Methionine residues of the PcuC apoprotein have various sensitivity to oxidation

The copper chaperone PcuC contains eight Met residues: one isolated in the first half of the primary sequence (Met25), two pairs clustered together (Met61 and Met63, and Met88 and Met90), and three near the C-terminus (Met128, Met132, and Met135) (**Figure 2A**). Using the purified recombinant apo-PcuC we first confirmed that it contains oxidizable Met and thus is relevant to quantify susceptibility to Met oxidation. It has up to seven MetO formed simultaneously by oxidation with H_2O_2 and most of which are reducible by the methionine sulfoxide reductase MsrP (**Supplementary Results 2.1, Figure S1**). To quantify the sensitivity of the Met in the apo-PcuC individually, we incubated the protein with 1,000 molar equivalents of $H_2^{18}O_2$ for 2 to 60 minutes and oxidized residue were quantified (**Figure 2B, S2B**). An SDS-PAGE analysis of the samples revealed a delay in protein migration, as observed for non-isotopic H_2O_2, confirming that oxidation occurred (**Figure S2A**). The identification and quantification of ^{18}O-Met revealed that the very most abundant form was ^{18}O-sulfoxide, with only traces of ^{18}O-containing sulfone detected (**Figure S2B**). The search for oxidized form of His (oxoHis) revealed only extremely low levels, confirming that Met were the target of oxidation by $H_2^{18}O_2$ (**Figure S2C**). The analysis of the kinetics of oxidation revealed striking differences in the reactivity of the Met residues (**Figure 2B, E**). Indeed, Met25 remained reduced throughout the treatment while Met135 was rapidly and extensively oxidized, up to ~93% after 60 minutes. Met63, which is part of the copper-coordinating site, was the second most oxidized residue, reaching ~45% by the end of the kinetics. The other Met showed intermediate levels of oxidation, all remaining below 50%. Interestingly, the oxidation kinetics of Met132 displayed a unique pattern compared to the other reactive Met residues. While for the other residues oxidation level early reached a plateau or increased linearly (Met61, Met128), Met132 oxidation level remained low at early timepoints but eventually increased to ~42%, like the level observed for Met63 (**Figure 2B**). This behavior suggests that Met132 may be a secondary target for oxidation, becoming oxidized only after the more sensitive residues, Met63, Met90, and Met135. Based on final oxidation levels, Met reactivity in

137 apo-PcuC ranked from least to most oxidized: Met25, Met128, Met88, Met61, Met90, Met132, Met63,

138 Met135.

Figure 2. Utilization of $H_2^{18}O_2$ and of an oxaziridine probe to quantify oxidizable Met in apo- and Cu-loaded PcuC. (**A**) Amino acid sequence of PcuC in single letter. The eight Met are shown in *red* with numbering above, amino acids on *green* background correspond to the periplasmic peptide signal and *black* triangles indicate Cu-coordinating residues. Met are numbered considering Asp as first residue after removal of the peptide signal. (**B**) Time-resolved percentage of oxidation of the eight Met of the apo-PcuC after incubation with $H_2^{18}O_2$. Apo-PcuC was incubated with 1,000 molar equivalents (equiv.) of $H_2^{18}O_2$ for 2, 5, 15, 30, 45 and 60 min and the reaction stopped with catalase. The presented values correspond to the sum of all oxidation forms involving ^{18}O (^{18}O-sulfoxide, ^{18}O-sulfone and $^{16}O/^{18}O$-sulfone). (**C**) Percentage of labeling of the apo-PcuC after incubation with the oxaziridine probe. Apo-PcuC was incubated with 0, 0.1, 0.5, 1, 5 or 10 equiv. of the oxaziridine probe and the reaction stopped by desalting. (**D**) Percentage of labeling of the Cu-loaded PcuC after incubation with the oxaziridine probe. Cu-loaded PcuC was incubated with the oxaziridine probe similarly to the apoprotein. (**E**) Heatmap showing the percentage of oxidation or labeling of the eight Met residues of PcuC, using a *blue*-to-*red* color scale, with *blue* and *red* representing the lowest and highest values of modification, respectively. Underlined Met63 and Met88 belong to the Cu-binding site.

2.2. The oxaziridine probe reactivity toward Met is similar to that of hydrogen peroxide

With the aim of quantifying Met sensitivity in the copper-loaded PcuC, we used an oxaziridine probe

[35,36]. First, we validated its ability to bind the apo-PcuC and its potential usability to quantify the

sensitivity of PcuC's Met to oxidation (**Supplementary Results 2.2, Figure S3**). Our first approach was

to determine if it can report similar results to those obtained using $H_2^{18}O_2$ with the apo-PcuC. To do so, we

designed a slightly different strategy as chemical probes are usually used at several concentrations to

identify reactive residues in activity-based protein profiling or related approaches [35,37]. Thus, we

incubated the apo-PcuC with 0 to 10 molar equivalents of probe and quantified the labeling of each Met

after trypsin proteolysis and LC-MS/MS (**Figure 2C, E**). The overall pattern closely matched the one

obtained with $H_2^{18}O_2$: Met25 was unlabeled, while Met135 was the most reactive, reaching ∼83% with 10

equivalents of probe. Both Met88 and Met128 were found labeled around 5%. Similarly, Met132 labeling

percentage (∼44%) was nearly identical to the values obtained with $H_2^{18}O_2$. Met88 and Met128 showed

low labeling (~5%), and Met132 reached ~44%, consistent with $H_2^{18}O_2$ results. Differences were observed

for Met63 and Met90, that were less labeled (~20% and ~11%, respectively), whereas Met61 showed

higher labeling (~51%). At the highest probe concentration, the reactivity ranking of Met residues in apo-

PcuC was: Met25, Met88, Met128, Met90, Met63, Met132, Met61, Met135. Consistently with the results

obtained with $H_2^{18}O_2$, this shows that copper-coordinating Met63 and Met88 do not present a sensitivity to

oxidation particularly different from those of the other residues. These results indicated that the use of the

oxaziridine probe is relevant to quantify Met sensitivity to oxidation in copper-loaded PcuC. To our

knowledge, although it was previously used to identify oxidizable Met [35,38], oxaziridine probes'

reactivity was never compared to physiological oxidants. This showed that the oxaziridine probe

effectively mimics H_2O_2-induced oxidation and is very likely relevant to identify redox-active Met

residues in proteins from various biological contexts.

2.3. Cu binding strongly alters PcuC's Met sensitivity to oxidation

Then, we wanted to evaluate whether the presence of copper in PcuC affects the Met sensitivity to oxidation using the same approach. First, we incubated the protein with increasing concentration of $CuSO_4$ and observed that two coppers could be bound to the protein using non-denaturing ESI-MS (**Figure S4A**), as shown for the *B. diazoefficiens* homolog [23]. As anticipated, the incubation of the copper-loaded PcuC with 1,000 equivalents of $H_2^{18}O_2$ resulted in a quick and complete protein degradation (**Figure S4B**). This was very likely due to Fenton-like reaction occurring between the peroxide and the copper, either bound to the protein or free in the solution because of remaining trace despite the desalting. Using the probe on the copper-loaded PcuC revealed a very different profile than for the apo protein (**Figure 2D, E**). Indeed, we first noticed that all Met were labeled, including Met25. By comparison with the apo-PcuC, we observed strong increases for Met25, Met63, Met88 and Met90 labeling. The most labeled became Met61, followed by Met132, while the level of labeling of Met135 strongly decreased. Interestingly, the reactivity of the copper-ligand Met63 and Met88 increased by 2.1-fold and 3.5-fold, respectively (**Figure 2D, E**). This indicates that copper-binding increases Met sensitivity to oxidation. At the highest probe concentration, the reactivity ranking in Cu-loaded PcuC was: Met128, Met25, Met88, Met135, Met63, Met90, Met132, and Met61. This change in reactivity could be attributed to alterations in the electronic density on sulfur atoms, as copper coordination withdraws electron density, increasing the electrophilicity of sulfur. Additionally, copper binding might stabilize the Met lateral chain in a conformation that favors oxidation. The apparent decrease in the sensitivity of Met135 is most likely due to the increase in sensitivity of the other Met residues, particularly those involved in copper coordination, as it is unlikely that the structural properties have been modified for this Met residue (the C-terminal part remained unfolded, *vide infra*).

2.4. PcuC coordinates two Cu ions at the canonical Cu binding site

To evaluate whether structural determinants can explain the observed differences of Met reactivity, we solved the tridimensional structures of the apo and copper-bound PcuC by X-ray crystallography (**Figure 3**). Apo structure of PcuC was solved at 1.05 Å resolution in a $P4_32_12$ space group obtained from a crystal

grown in sodium citrate at pH 8.5 (**Table S1**). The structure is typical for homologs of this protein family,

i.e. a seven stranded β-barrel with one side of this barrel containing three additional strands, which include

the conserved Met and His residues involved in copper binding. Two parts of the protein could not be

resolved: a first from Ala55 to Met61, and a second corresponding to the C-terminal part, starting from

Arg124 (and therefore containing Met128, Met132 and Met135) (**Figure 3A**). These parts are very likely

flexible in the protein, and the C-terminal part was predicted to be disordered (**Figure S5**). To obtain

experimental data on copper binding to PcuC, we added ten copper equivalents to the protein and re-

screened for crystallization. We obtained a first crystal that clearly showed two sites of high electron

density, very likely corresponding to metals, but was affected by translational non-crystallographic

symmetry (**Supplementary Results 2.4**). Then, we obtained a second crystal that diffracted to 2.75 Å

resolution and the elucidated structure showed the presence of two copper ions (**Table S1, Figure 3B**).

One copper (Cu1) is bound to Met88 Sδ (2.6 Å) and His86 Nδ (2.1 Å), similarly to observed in the PcuC

homologs from *Streptomyces lividans* [39] and *Thermus thermophilus* [22]. However, in these homologs

the copper is coordinated by two other ligands (corresponding to His51 and Met63), whereas in the

structure of *C. sphaeroides* PcuC a water molecule linked the copper to Nδ of His51 (2.7 Å; Cu1-HOH =

3 Å), and Met63 adopted such a conformation that its side chain was not in the coordination sphere of Cu1

(the distance Cu1 to Met63 Sδ is 5.5Å). Finally, what we interpreted as a citrate molecule, very likely

coming from the crystallization media, participated in the coordination sphere of this copper with a

bidentate interaction from one carboxylate moiety (Cu1-OA1 = 2.4 Å and Cu1-OA2 = 2.9Å). Overall,

although the position of this copper binding site is conserved between the three homologs structurally

characterized, the coordination sphere is different in *C. sphaeroides* PcuC. The additional copper ion

(Cu2) observed in the structure of *C. sphaeroides* PcuC was bound by Nε of His53 (1.8 Å) and three

oxygen atoms from the three carboxylate moieties of citric acid (2.7 Å, 3.1Å and 3.1 Å; **Figure 3B**).

Altogether, these results show that PcuC could bind two coppers without the involvement of the unfolded

C-terminal part. This finding was unexpected given the results obtained for the *B. diazoefficiens* PcuC

homolog [23]. While it is possible that the two chaperones function differently, several elements support a

unified model that reconciles both sets of results. In this model, a first copper ion is stably coordinated in

the canonical site (high-affinity), while a second copper ion oscillates between various positions along the

C-terminal extension and the coordination site observed in our structure (low-affinity) (**Supplementary**

Discussion).

Figure 3. Structural characterization of apo-PcuC and Cu-loaded PcuC. Overall structure of the apo

form of PcuC (PDB 9REZ) **(A)** and the Cu-loaded form of PcuC (PDB 9RF0) **(B)**. **(C)** Superimposition of

both the apo-and copper-bound form of PcuC with the side chains of the discussed residues shown in

stick. **(D)** Details of the Cu-binding site of *S. lividans* PcuC (PDB 3ZJA).

2.5. Met oxidation sensitivity correlates with solvent exposure in the folded region but with positive charge in the unfolded C-terminal extension

To understand the factors influencing Met oxidation in PcuC, we analyzed the solvent exposure and local charges surrounding each Met in both apo- and Cu-bound structures. As the unfolded C-terminal extension containing Met128, Met132, and Met135 was unresolved, we used an AlphaFold model to complete the analysis (**Supplementary Results 2.5, Figure S6A**). In the folded region (encompassing Met25 to Met90), solvent exposure values were consistent between the resolved and modeled structures. Met25 showed minimal exposure, while Met61 and Met63 were among the most exposed. Met128, Met132, and Met135, in the unfolded C-terminal extension, displayed high solvent exposure in the model (**Figure 4A**). We assumed that the C-terminal extension remains unstructured in both the apo- and Cu-PcuC, and the exposure values for these Met are similar to the model. To determine if oxidation sensitivity correlates with solvent exposure, we plotted the values of oxidation determined with ^{18}O-labeled hydrogen peroxide and oxaziridine as function of the Met exposure values for both the apo- and the copper-loaded PcuC structures (**Figure 4C, D**). In apo-PcuC, oxidation levels correlated positively with solvent exposure for Met residues in the folded region (R^2 = 0.48 for $H_2^{18}O_2$ and R^2 = 0.90 for oxaziridine). A similar positive correlation was observed using the solvent exposure values from the AlphaFold model (**Figure S6B**). Surprisingly, the C-terminal Met residues showed a strong negative correlation between oxidation and solvent exposure (R^2 = 0.99). Similar trends were observed in Cu-loaded PcuC, though with lower correlation values (**Figure 4C**). Strikingly, the most exposed Met128 was one of the least oxidized whether with $H_2^{18}O_2$ or oxaziridine, and with or without copper. To explain this discrepancy in the disordered region, we considered the electrostatic environment. Surface charge maps revealed that, while most of the protein surface is neutral or negatively charged, the C-terminal region includes a localized positive charge due to a lysine (**Figure S7A, B, C**). We quantified the partial charge within 5.5 Å of each Met sulfur (**Figure 4D**). In the folded domain, no correlation was found between oxidation and local charge (**Figure S7D, E**). However, for the C-terminal Met residues, we observed a strong positive correlation between oxidation sensitivity and positive local charge (R^2 = 0.92–0.97 in apo-PcuC) (**Figure**

4E). These results indicated that the oxidation sensitivity of the Met belonging to the unfolded C-terminal part of PcuC was positively correlated with the positive partial charge surrounding the sulfur atom of the Met. Together, these results reveal that distinct factors govern metal oxidation in different structural contexts of PcuC: solvent exposure is crucial in the folded part, while local positive charge controls reactivity in the unfolded part. These results are consistent with the fact that solvent exposure plays very likely a major role in Met sensitivity to oxidation, as shown previously [40–42], but that other factors should be considered [43,44]. This suggests that Met sensitivity to oxidation might be protein and Met dependent and should be determined for each protein of interest.

Figure 4. Correlation of Met oxidation sensitivity with solvent exposure and local charge. (**A**) Heatmap showing the solvent exposure (Å²) of the Met, using a *blue*-to-*red* color scale, with *blue* and *red* representing the lowest and highest exposure, respectively. *n.d., not determined.* Underlined Met63 and Met88 belong to the Cu-binding site. (**B**) Plot of oxidation level of the Met as function of solvent exposure for the apo-PcuC after 60 min oxidation with $H_2^{18}O_2$ with independent linear regressions for Met of the folded part and the unfolded C-terminal part. (**C**) Plot of oxidation level of the Met as function of solvent exposure for the apo- and Cu-loaded PcuC after exposure to 10 equivalents of oxaziridine with independent linear regressions for Met of the folded part and the unfolded C-terminal part. (**D**) Heatmap showing the partial charge of the environment surrounding the sulfur of the Met, using a *blue*-to-*red* color scale, with *blue* and *red* representing the more negative and positive charge, respectively. *n.r., not resolved.* Underlined Met63 and Met88 belong to the canonical Cu-binding site. (**E**) Plot of oxidation of the Met from the C-terminal unfolded part determined with $H_2^{18}O_2$ and oxaziridine as function of the partial charge of the environment surrounding the sulfur of the Met determined on the AlphaFold model for the apo- and Cu-loaded PcuC with linear regressions.

3. Conclusion

In this study, we showed that the recently developed oxaziridine probe reacts with Met residues similarly to $H_2{}^{18}O_2$, using the copper chaperone PcuC from *C. sphaeroides* as a model. This suggests the probe is a reliable tool for quantitatively mapping Met oxidation sensitivity in any protein, including in metalloproteins. Copper-coordinating Met residues are not significantly more or less sensitive to oxidation than non-coordinating ones. We also provide new insights into the function of the conserved, disordered C-terminal extension present in most PcuC homologs. This region could play a dual role: facilitating copper incorporation into the coordination site, as suggested for copper transporters and oxidases [16,45], and protecting against oxidative stress via Met135, a highly oxidation-sensitive residue. Met135 may act as a local scavenger of reactive species and be regenerated by the periplasmic reductase MsrP, ensuring efficient copper delivery and CoxB assembly, ultimately supporting proper bacterial respiration.

4. Materials and methods

4.1. Protein and chemicals

PcuC was produced and purified as described [46]. [18]O-labeled hydrogen peroxide and dibenzocyclooctyne-Cyanine5 fluorophore (DBCO-Cy5) were purchased from Sigma-Aldrich. Initial concentration of commercial $H_2^{18}O_2$ was estimated to be ~450 mM using FOX method [47]. The oxaziridine probe ((4-azidopiperidin-1-yl)(3-phenyl-1,2-oxaziridin-2-yl)methanone) was synthetized according to [36] and validated by NMR (**Figure S8**). Protein concentrations were determined using the Bradford method (Bio-Rad Protein Assay Dye Reagent Concentrate). SDS-PAGE were performed using Bolt[TM] pre-cast gel system (ThermoFisher) using 12% polyacrylamide gels and 2-(N-Morpholino)ethanesulfonic acid (MES) running buffer. Visible and fluorescent imaging were performed using Amersham ImageQuant™ 800 (Cytiva).

4.2. PcuC oxidation, Cu-loading and labeling with the oxaziridine probe

Apo-PcuC (70 µM) was oxidized with 10, 100, 500 or 1,000 molar equivalents (equiv.) of H_2O_2 in 50 mM 3-(N-Morpholino)propanesulfonic acid (MOPS) buffer at pH 8.0 for 1 h and the reaction was stopped by adding excess dithiothreitol (DTT) (100 mM) and subsequent desalting using Sephadex G-10 (PD MidiTrap G-10, Cytiva). For oxidation with $H_2^{18}O_2$, PcuC (100 µM), apo or Cu-loaded, was incubated for 2, 5, 15, 30, 45 and 60 min with 1,000 equiv. of $H_2^{18}O_2$ in 50 mM MOPS, pH 8.0, then the reaction was stopped with the addition of 0.071 U.µl^{-1} of catalase followed by desalting. PcuC Cu-loading was performed by incubating the protein (200 µM) with 2 to 10 equiv. of $CuSO_4$ in MOPS 50 mM followed by desalting. For Cu-loading before oxaziridine labeling, 5 equiv. of $CuSO_4$ were used. Oxaziridine labeling was made by incubating PcuC (14 or 100 µM) with 0, 0.1, 0.5, 1, 5 or 10 equiv. of the probe for 20 or 30 min in 50 mM MOPS, pH 8.0 followed by desalting. For in-gel fluorescence analysis, oxaziridine-labeled PcuC was incubated with 10 equiv. of DBCO-Cy5 overnight in 50 mM MOPS, pH 8.0 and excess reagent was removed by ultrafiltration (5 kDa cutoff, Vivaspin 500, Sartorius).

4.3. Mass spectrometry analyses

ESI-MS analyses were performed as described [19]. For proteolysis and LC-MS/MS analyses, 20 µL of PcuC solution (7 µM) in 50 mM MOPS buffer (pH 8.0) with 20% acetonitrile (ACN) were digested by adding 3 µL of trypsin/Lys C (Promega) at a concentration of 12.5 ng.µL^{-1} in 0.1% formic acid (FA) for 3 h at 37°C. For TMT labeling, the manufacturer's protocol (ThermoFisher) was followed. Peptides from each experimental condition (various concentrations of oxaziridine or different reaction times with $H_2^{18}O_2$) were incubated with one of the six 6-plex TMT reagents, consisting of reporter ions 126, 127, 128, 129, 130, and 131 (0.8 mg in 41 µL of anhydrous ACN) for 1 h at room temperature. The reactions were quenched by adding 2 µL of 5% ammonium hydroxide for 15 minutes. For each experiment (oxaziridine or $H_2^{18}O_2$), the six conditions were pooled and diluted in 300 µL of 0.1% trifluoroacetic acid (TFA). The labeled peptides were filtered using a C18 column (ThermoFisher) to remove excess TMT reagents, eluted with 2×300 µL of 50% ACN/0.1% TFA, and dried using a SpeedVac. The dried samples were reconstituted in 2% ACN/0.1% FA to achieve a theoretical protein concentration of 1 µg.µL^{-1}. The samples were injected into an LC-MS/MS system using a high-performance nano-liquid chromatography (UHPLC RSLC, ThermoFisher) equipped with online desalting (PEPMAP C18, 5 µm, 300 Å trap column, 300 µm × 5 mm), coupled to a Q-Exactive Plus hybrid mass spectrometer (ThermoFisher). The analytical column used was an Acclaim PepMap RSLC capillary column (nanoViper C18, 2 µm, 100 Å, 75 µm × 15 cm). The gradient for peptide elution was as follows: solvent A: 0.1% FA, solvent B: ACN with 0.1% FA. The gradient program included the following steps: (1) 2.5–25% B over 57 min, (2) 25–50% B over 6 min, (3) 50–90% B in 1 min, and (4) 90% B for 10 min. A data-dependent acquisition (DDA) method was employed with a Top10 approach, where the 10 most abundant peptides were selected for fragmentation in each survey scan. MS settings were: precursor mass range 375 to 1400 m/z, resolution 70,000, AGC target 3×10^6, maximum IT 100 ms. MS/MS parameters: fragment mass range 100 to 2000 m/z, resolution 70,000, AGC target 2×10^5, and normalized collision energy of 27 (33 for TMT-labeled samples). Peak list files in .mgf format (Mascot Generic Format) were generated using Proteome Discoverer 2.4 (ThermoFisher). Protein identification was carried out using MASCOT 2.3 software, querying a database

containing PcuC and common contaminants (246 sequences, including trypsin, keratins, etc.). Search

parameters were set as follows: trypsin as the digestion enzyme, up to 4 missed cleavages, precursor mass

tolerance of 10 ppm, product ion mass accuracy of 0.02 Da, with fixed modifications of TMT6plex (N-

term) and TMT6plex (K), and variable modifications including deamidation (Gln, Asn), ^{16}O-oxidation

(Met, monoisotopic mass = 15.994915 Da), ^{18}O-oxidation (Met, monoisotopic mass = 17.99916), ^{16}O-

dioxidation (Met, monoisotopic mass = 31.989829 Da), ^{18}O-dioxidation (Met, monoisotopic mass =

35.998321 Da), ^{16}O/^{18}O-dioxidation (Met, monoisotopic mass = 33.994075 Da) or oxaziridine

modification (Met, monoisotopic mass = 167.080710 Da). Quantification was performed by exporting

reporter ion intensity data for all peptides to Excel using Mascot software.

4.4. PcuC crystallization and structure analysis

Crystals of PcuC were obtained by the sitting drop method using 96 wells plates and a pipetting robot

(TECAN Evoware). The apo protein was concentrated to 26 mg.mL^{-1} and crystals were obtained by

mixing 1 µL of protein with 1 µL of 1.2 M sodium citrate and 0.1 M Tris pH8.5. Crystals quickly

appeared and, after supplementing the mother liquor with 15% glycerol, a few crystals were flash frozen

directly in liquid nitrogen. Crystals were collected at the SOLEIL synchrotron (France) and the structure

of apo PcuC was then solved at 1.05 Å resolution in a $P4_32_12$ space group. To obtain experimental data on

copper binding to PcuC, ten copper equivalents were added to the protein and crystallization of the protein

was rescreened. Crystals were obtained in a condition containing 25% PEG 3350 and 20 mM CaCl$_2$, 20

mM CdCl$_2$ and 20 mM CoCl$_2$. This crystal form diffracted to 2.1 Å resolution, however it is affected by a

translational NCS (tNCS) that, together with the three molecules in the asymmetric unit, appeared to be

difficult to refine ($R = 30.2\%$; $R_{\text{free}} = 36.5\%$). This structure is therefore not presented, although it clearly

showed three sites of high electron density, all of them relatively close to each other. Data collection at

different wavelengths (copper and cobalt edges) also indicated that one site was occupied by copper,

whereas the two others were cadmium ions. Additional screens were used to find a different crystal form

that would not grow in a complex mixture of metal and that would eventually not be affected by tNCS. A

crystal form was obtained in a condition containing 20% PEG 6000, 1 M LiCl$_2$ and 0.1 M citric acid at pH

4. It diffracted to 2.75Å resolution and was refined to a R and R_{free} of 24.2% and 30.4%, respectively.

Prediction of intrinsically disordered region was made using IUpred2A [48]. Solvent exposure of Met

was determined using the "*get_area*" function with default parameters and partial charges were

determined using APBS Electrostatics plugin of PyMol 3.1.0a0 (open-source build).

Supplementary information

Supplementary Results, Supplementary Discussion, Figure S1-S8, Table S1, Supplementary References.

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE [49,50] partner repository with the dataset identifier PXD062219. Structure data have been deposited in Protein Data Bank (PDB) with accession code 9REZ for the apo form and 9RF0 for the copper-bound form.

Acknowledgments

This research was funded by the Commissariat à l'Energie Atomique et aux Energies Alternatives (CEA) and by the project METOXIC [ANR 16-CE11-0012]. This work received support from the French government under the France 2030 investment plan, as part of the Initiative d'Excellence d'Aix-Marseille Université —IM2B (AMX-19-IET-006) for L.M. and L.T. – A*MIDEX (AMX-21-PEP-028) for L.T. The authors would like to thank the NovoNordisk foundation (OxyMiST project, grant number NNF20OC0059697).

CRediT author statement

Lionel Tarrago: Conceptualization, Methodology, Validation, Formal analysis, Investigation, Resources, Data Curation, Writing - Original Draft, Writing - Review & Editing, Visualization, Supervision, Funding acquisition, Project administration. **Lise Molinelli:** Conceptualization, Methodology, Validation, Formal analysis, Investigation, Data Curation, Writing - Review & Editing. **Maya Belghazi:** Conceptualization, Methodology, Validation, Formal analysis, Investigation, Data Curation, Writing - Review & Editing. **Mathilde Tribout:** Formal analysis, Investigation, Data Curation. **David Lemaire:** Methodology, Validation, Formal analysis, Investigation, Data Curation. **Pierre Legrand:** Methodology, Validation, Formal analysis, Investigation, Data Curation. **Sandrine Grosse:** Formal analysis, Investigation, Data

Curation. **David Pignol**: Conceptualization, Methodology, Supervision, Funding acquisition, Writing -

Review & Editing. **Monique Sabaty:** Conceptualization, Methodology, Supervision, Funding acquisition,

Writing - Review & Editing. **Thierry Tron:** Conceptualization, Methodology, Supervision, Funding

acquisition, Writing - Review & Editing. **Pascal Arnoux:** Conceptualization, Methodology, Validation,

Formal analysis, Investigation, Resources, Data Curation, Writing - Original Draft, Writing - Review &

Editing, Visualization, Supervision, Funding acquisition, Project administration.

Declaration of generative AI in scientific writing

During the preparation of this work the authors used ChatGPT to improve the clarity and readability of the

text. After using this tool, the author reviewed and edited the content as needed and take full responsibility

for the content of the publication

References

[1] F.W. Outten, B.S. Twining, Metal Homeostasis, in: Wiley Encyclopedia of Chemical Biology, John Wiley & Sons, Ltd, 2008: pp. 1–10. https://doi.org/10.1002/9780470048672.wecb324.

[2] J.T. Rubino, K.J. Franz, Coordination chemistry of copper proteins: How nature handles a toxic cargo for essential function, Journal of Inorganic Biochemistry 107 (2012) 129–143. https://doi.org/10.1016/j.jinorgbio.2011.11.024.

[3] R.M. Rasia, C.W. Bertoncini, D. Marsh, W. Hoyer, D. Cherny, M. Zweckstetter, C. Griesinger, T.M. Jovin, C.O. Fernández, Structural characterization of copper(II) binding to α-synuclein: Insights into the bioinorganic chemistry of Parkinson's disease, Proc Natl Acad Sci U S A 102 (2005) 4294–4299. https://doi.org/10.1073/pnas.0407881102.

[4] J.H. Viles, F.E. Cohen, S.B. Prusiner, D.B. Goodin, P.E. Wright, H.J. Dyson, Copper binding to the prion protein: Structural implications of four identical cooperative binding sites, Proceedings of the National Academy of Sciences 96 (1999) 2042–2047. https://doi.org/10.1073/pnas.96.5.2042.

[5] J. Arguello, D. Raimunda, T. Padilla-Benavides, Mechanisms of copper homeostasis in bacteria, Frontiers in Cellular and Infection Microbiology 3 (2013). https://www.frontiersin.org/article/10.3389/fcimb.2013.00073 (accessed May 9, 2022).

[6] J.A. Imlay, Where in the world do bacteria experience oxidative stress?, Environ. Microbiol. 21 (2019) 521–530. https://doi.org/10.1111/1462-2920.14445.

[7] C.L. Hawkins, M.J. Davies, Detection, identification, and quantification of oxidative protein modifications, J. Biol. Chem. 294 (2019) 19683–19708. https://doi.org/10.1074/jbc.REV119.006217.

[8] M.J. Davies, The oxidative environment and protein damage, Biochim. Biophys. Acta 1703 (2005) 93–109. https://doi.org/10.1016/j.bbapap.2004.08.007.

[9] M.K. Tiwari, P.M. Hägglund, I.M. Møller, M.J. Davies, M.J. Bjerrum, Copper ion / H2O2 oxidation of Cu/Zn-Superoxide dismutase: Implications for enzymatic activity and antioxidant action, Redox Biol 26 (2019) 101262. https://doi.org/10.1016/j.redox.2019.101262.

[10] B. Bissaro, Å.K. Røhr, G. Müller, P. Chylenski, M. Skaugen, Z. Forsberg, S.J. Horn, G. Vaaje-Kolstad, V.G.H. Eijsink, Oxidative cleavage of polysaccharides by monocopper enzymes depends on H2O2, Nat. Chem. Biol. 13 (2017) 1123–1128. https://doi.org/10.1038/nchembio.2470.

[11] D. Kracher, M. Andlar, P.G. Furtmüller, R. Ludwig, Active-site copper reduction promotes substrate binding of fungal lytic polysaccharide monooxygenase and reduces stability, Journal of Biological Chemistry 293 (2018) 1676–1687. https://doi.org/10.1074/jbc.RA117.000109.

[12] E. Ponzini, A. De Palma, L. Cerboni, A. Natalello, R. Rossi, R. Moons, A. Konijnenberg, J. Narkiewicz, G. Legname, F. Sobott, P. Mauri, C. Santambrogio, R. Grandori, Methionine oxidation in α-synuclein inhibits its propensity for ordered secondary structure, J Biol Chem 294 (2019) 5657–5665. https://doi.org/10.1074/jbc.RA118.001907.

[13] N.D. Younan, R.C. Nadal, P. Davies, D.R. Brown, J.H. Viles, Methionine oxidation perturbs the structural core of the prion protein and suggests a generic misfolding pathway, J Biol Chem 287 (2012) 28263–28275. https://doi.org/10.1074/jbc.M112.354779.

[14] J.T. Rubino, P. Riggs-Gelasco, K.J. Franz, Methionine motifs of copper transport proteins provide general and flexible thioether-only binding sites for Cu(I) and Ag(I), J. Biol. Inorg. Chem. 15 (2010) 1033–1049. https://doi.org/10.1007/s00775-010-0663-9.

[15] J.T. Rubino, M.P. Chenkin, M. Keller, P. Riggs-Gelasco, K.J. Franz, A comparison of methionine, histidine and cysteine in copper(i)-binding peptides reveals differences relevant to copper uptake by organisms in diverse environments, Metallomics 3 (2011) 61–73. https://doi.org/10.1039/c0mt00044b.

463   [16]   U. Contaldo, D. Savant-Aira, A. Vergnes, J. Becam, F. Biaso, M. Ilbert, L. Aussel, B. Ezraty, E. Lojou, I.
464   Mazurenko, Methionine-rich domains emerge as facilitators of copper recruitment in detoxification
465   systems, Proceedings of the National Academy of Sciences 121 (2024) e2402862121.
466   https://doi.org/10.1073/pnas.2402862121.

467   [17]   L. Macomber, C. Rensing, J.A. Imlay, Intracellular copper does not catalyze the formation of
468   oxidative DNA damage in Escherichia coli, J Bacteriol 189 (2007) 1616–1626.
469   https://doi.org/10.1128/JB.01357-06.

470   [18]   A. Gennaris, B. Ezraty, C. Henry, R. Agrebi, A. Vergnes, E. Oheix, J. Bos, P. Leverrier, L. Espinosa, J.
471   Szewczyk, D. Vertommen, O. Iranzo, J.-F. Collet, F. Barras, Repairing oxidized proteins in the
472   bacterial envelope using respiratory chain electrons, Nature 528 (2015) 409–412.
473   https://doi.org/10.1038/nature15764.

474   [19]   L. Tarrago, S. Grosse, M.I. Siponen, D. Lemaire, B. Alonso, G. Miotello, J. Armengaud, P. Arnoux, D.
475   Pignol, M. Sabaty, Rhodobactersphaeroides methionine sulfoxide reductase P reduces R- and S-
476   diastereomers of methionine sulfoxide from a broad-spectrum of protein substrates, Biochem. J.
477   475 (2018) 3779–3795. https://doi.org/10.1042/BCJ20180706.

478   [20]   A. Vergnes, C. Henry, G. Grassini, L. Loiseau, S. El Hajj, Y. Denis, A. Galinier, D. Vertommen, L.
479   Aussel, B. Ezraty, Periplasmic oxidized-protein repair during copper stress in E. coli: A focus on the
480   metallochaperone CusF, PLoS Genet 18 (2022) e1010180.
481   https://doi.org/10.1371/journal.pgen.1010180.

482   [21]   F.L. Sousa, R.J. Alves, M.A. Ribeiro, J.B. Pereira-Leal, M. Teixeira, M.M. Pereira, The superfamily of
483   heme-copper oxygen reductases: types and evolutionary considerations, Biochim Biophys Acta
484   1817 (2012) 629–637. https://doi.org/10.1016/j.bbabio.2011.09.020.

485   [22]   L.A. Abriata, L. Banci, I. Bertini, S. Ciofi-Baffoni, P. Gkazonis, G.A. Spyroulias, A.J. Vila, S. Wang,
486   Mechanism of Cu(A) assembly, Nat. Chem. Biol. 4 (2008) 599–601.
487   https://doi.org/10.1038/nchembio.110.

488   [23]   F. Canonica, D. Klose, R. Ledermann, M.M. Sauer, H.K. Abicht, N. Quade, A.D. Gossert, S. Chesnov,
489   H.-M. Fischer, G. Jeschke, H. Hennecke, R. Glockshuber, Structural basis and mechanism for
490   metallochaperone-assisted assembly of the CuA center in cytochrome oxidase, Sci Adv 5 (2019)
491   eaaw8478. https://doi.org/10.1126/sciadv.aaw8478.

492   [24]   L. Hederstedt, Diversity of Cytochrome c Oxidase Assembly Proteins in Bacteria, Microorganisms 10
493   (2022) 926. https://doi.org/10.3390/microorganisms10050926.

494   [25]   A.K. Thompson, J. Gray, A. Liu, J.P. Hosler, The roles of Rhodobacter sphaeroides copper
495   chaperones PCu(A)C and Sco (PrrC) in the assembly of the copper centers of the aa(3)-type and the
496   cbb(3)-type cytochrome c oxidases, Biochim. Biophys. Acta 1817 (2012) 955–964.
497   https://doi.org/10.1016/j.bbabio.2012.01.003.

498   [26]   A.K. Boal, A.C. Rosenzweig, Structural Biology of Copper Trafficking, Chem. Rev. 109 (2009) 4760–
499   4779. https://doi.org/10.1021/cr900104z.

500   [27]   R.A. Festa, D.J. Thiele, Copper: An essential metal in biology, Current Biology 21 (2011) R877–R883.
501   https://doi.org/10.1016/j.cub.2011.09.040.

502   [28]   S. Iwata, C. Ostermeier, B. Ludwig, H. Michel, Structure at 2.8 A resolution of cytochrome c oxidase
503   from Paracoccus denitrificans, Nature 376 (1995) 660–669. https://doi.org/10.1038/376660a0.

504   [29]   H.K. Abicht, M.A. Schärer, N. Quade, R. Ledermann, E. Mohorko, G. Capitani, H. Hennecke, R.
505   Glockshuber, How periplasmic thioredoxin TlpA reduces bacterial copper chaperone Sco1 and
506   cytochrome oxidase subunit II (CoxB) prior to metallation, J Biol Chem 289 (2014) 32431–32444.
507   https://doi.org/10.1074/jbc.M114.607127.

508   [30]   L. Banci, I. Bertini, S. Ciofi-Baffoni, E. Katsari, N. Katsaros, K. Kubicek, S. Mangani, A copper(I)
509   protein possibly involved in the assembly of CuA center of bacterial cytochrome c oxidase, Proc.
510   Natl. Acad. Sci. U.S.A. 102 (2005) 3994–3999. https://doi.org/10.1073/pnas.0406150102.

[31] O.S. Fisher, M.R. Sendzik, M.O. Ross, T.J. Lawton, B.M. Hoffman, A.C. Rosenzweig, PCuAC domains
from methane-oxidizing bacteria use a histidine brace to bind copper, J. Biol. Chem. 294 (2019)
16351–16363. https://doi.org/10.1074/jbc.RA119.010093.

[32] H. Liu, G. Ponniah, A. Neill, R. Patel, B. Andrien, Accurate Determination of Protein Methionine
Oxidation by Stable Isotope Labeling and LC-MS Analysis, Anal. Chem. 85 (2013) 11705–11709.
https://doi.org/10.1021/ac403072w.

[33] J.Q. Bettinger, K.A. Welle, J.R. Hryhorenko, S. Ghaemmaghami, Quantitative Analysis of in Vivo
Methionine Oxidation of the Human Proteome, J Proteome Res 19 (2020) 624–633.
https://doi.org/10.1021/acs.jproteome.9b00505.

[34] L. Molinelli, E. Drula, J.-C. Gaillard, D. Navarro, J. Armengaud, J.-G. Berrin, T. Tron, L. Tarrago,
Methionine oxidation of carbohydrate-active enzymes during white-rot wood decay, Appl Environ
Microbiol 90 (2024) e0193123. https://doi.org/10.1128/aem.01931-23.

[35] S. Lin, X. Yang, S. Jia, A.M. Weeks, M. Hornsby, P.S. Lee, R.V. Nichiporuk, A.T. Iavarone, J.A. Wells,
F.D. Toste, C.J. Chang, Redox-based reagents for chemoselective methionine bioconjugation,
Science 355 (2017) 597–602. https://doi.org/10.1126/science.aal3316.

[36] S.K. Elledge, H.L. Tran, A.H. Christian, V. Steri, B. Hann, F.D. Toste, C.J. Chang, J.A. Wells, Systematic
identification of engineered methionines and oxaziridines for efficient, stable, and site-specific
antibody bioconjugation, Proc. Natl. Acad. Sci. U.S.A. (2020).
https://doi.org/10.1073/pnas.1920561117.

[37] E. Weerapana, C. Wang, G.M. Simon, F. Richter, S. Khare, M.B.D. Dillon, D.A. Bachovchin, K.
Mowen, D. Baker, B.F. Cravatt, Quantitative reactivity profiling predicts functional cysteines in
proteomes, Nature 468 (2010) 790–795. https://doi.org/10.1038/nature09472.

[38] D. He, H. Feng, B. Sundberg, J. Yang, J. Powers, A.H. Christian, J.E. Wilkinson, C. Monnin, D.
Avizonis, C.J. Thomas, R.A. Friedman, M.D. Kluger, M.A. Hollingsworth, P.M. Grandgenett, K.A.
Klute, F.D. Toste, C.J. Chang, I.I.C. Chio, Methionine oxidation activates pyruvate kinase M2 to
promote pancreatic cancer metastasis, Mol Cell 82 (2022) 3045-3060.e11.
https://doi.org/10.1016/j.molcel.2022.06.005.

[39] K.L.I.M. Blundell, M.A. Hough, E. Vijgenboom, J.A.R. Worrall, Structural and mechanistic insights
into an extracytoplasmic copper trafficking pathway in Streptomyces lividans, Biochemical Journal
459 (2014) 525–538. https://doi.org/10.1042/BJ20140017.

[40] E.J. Walker, J.Q. Bettinger, K.A. Welle, J.R. Hryhorenko, A.M. Molina Vargas, M.R. O'Connell, S.
Ghaemmaghami, Protein folding stabilities are a major determinant of oxidation rates for buried
methionine residues, Journal of Biological Chemistry (2022) 101872.
https://doi.org/10.1016/j.jbc.2022.101872.

[41] I. Dalle-Donne, R. Rossi, D. Giustarini, N. Gagliano, P. Di Simplicio, R. Colombo, A. Milzani,
Methionine oxidation as a major cause of the functional impairment of oxidized actin, Free Radical
Biology and Medicine 32 (2002) 927–937. https://doi.org/10.1016/S0891-5849(02)00799-2.

[42] H. Pan, K. Chen, L. Chu, F. Kinderman, I. Apostol, G. Huang, Methionine oxidation in human IgG2 Fc
decreases binding affinities to protein A and FcRn, Protein Science 18 (2009) 424–433.
https://doi.org/10.1002/pro.45.

[43] E. Vandermarliere, B. Ghesquière, V. Jonckheere, K. Gevaert, L. Martens, Unraveling the
specificities of the different human methionine sulfoxide reductases, Proteomics 14 (2014) 1990–
1998. https://doi.org/10.1002/pmic.201300357.

[44] J.Q. Bettinger, M. Simon, A. Korotkov, K.A. Welle, J.R. Hryhorenko, A. Seluanov, V. Gorbunova, S.
Ghaemmaghami, Accurate Proteomewide Measurement of Methionine Oxidation in Aging Mouse
Brains, J Proteome Res 21 (2022) 1495–1509. https://doi.org/10.1021/acs.jproteome.2c00127.

[45] S. Kar, S. Sen, S. Maji, D. Saraf, Ruturaj, R. Paul, S. Dutt, B. Mondal, E. Rodriguez-Boulan, R.
Schreiner, D. Sengupta, A. Gupta, Copper(II) import and reduction are dependent on His-Met

clusters in the extracellular amino terminus of human copper transporter-1, Journal of Biological
Chemistry 298 (2022). https://doi.org/10.1016/j.jbc.2022.101631.
[46] L. Tarrago, S. Grosse, D. Lemaire, L. Faure, M. Tribout, M.I. Siponen, M. Kojadinovic-Sirinelli, D.
Pignol, P. Arnoux, M. Sabaty, Reduction of Protein Bound Methionine Sulfoxide by a Periplasmic
Dimethyl Sulfoxide Reductase, Antioxidants 9 (2020) 616. https://doi.org/10.3390/antiox9070616.
[47] S.P. Wolff, [18] Ferrous ion oxidation in presence of ferric ion indicator xylenol orange for
measurement of hydroperoxides, in: Methods in Enzymology, Academic Press, 1994: pp. 182–189.
https://doi.org/10.1016/S0076-6879(94)33021-2.
[48] B. Mészáros, G. Erdős, Z. Dosztányi, IUPred2A: context-dependent prediction of protein disorder as
a function of redox state and protein binding, Nucleic Acids Research 46 (2018) W329–W337.
https://doi.org/10.1093/nar/gky384.
[49] Y. Perez-Riverol, C. Bandla, D.J. Kundu, S. Kamatchinathan, J. Bai, S. Hewapathirana, N.S. John, A.
Prakash, M. Walzer, S. Wang, J.A. Vizcaíno, The PRIDE database at 20 years: 2025 update, Nucleic
Acids Res 53 (2025) D543–D553. https://doi.org/10.1093/nar/gkae1011.
[50] E.W. Deutsch, N. Bandeira, Y. Perez-Riverol, V. Sharma, J.J. Carver, L. Mendoza, D.J. Kundu, S.
Wang, C. Bandla, S. Kamatchinathan, S. Hewapathirana, B.S. Pullman, J. Wertz, Z. Sun, S. Kawano, S.
Okuda, Y. Watanabe, B. MacLean, M.J. MacCoss, Y. Zhu, Y. Ishihama, J.A. Vizcaíno, The
ProteomeXchange consortium at 10 years: 2023 update, Nucleic Acids Res 51 (2023) D1539–
D1548. https://doi.org/10.1093/nar/gkac1040.
578