barbieQ: An R software package for analysing barcode count data from lineage tracing experiments

Liyang Fei[1, 2], Jovana Maksimovic[1, 2, +,] Alicia Oshlack[1, 2, *, +]

[1] Peter MacCallum Cancer Centre, Melbourne, Victoria 3000, Australia

[2] Sir Peter MacCallum Department of Oncology, The University of Melbourne, Victoria 3010, Australia

[*] To whom correspondence should be addressed.

[+] These authors jointly supervised this work.

Abstract

Motivation: A "clone" encompasses a progenitor cell and its progeny cells. Tracking clonal composition as cells differentiate or evolve is useful in many fields. Various single-cell lineage tracing (clonal tracking) technologies use unique DNA barcodes that are passed from progenitor cells to their offspring. The barcode count for each sample indicates cell number in clones. However, analysis of barcode count data is often bespoke and relies on visualisations and heuristics. A generalized workflow for preprocessing and robust statistical analysis of barcode count data across protocols is needed.

Results: We introduce *barbieQ*, a Bioconductor R package for analysing barcode count data across groups of samples. It provides data-driven quality control and filtering, extensive visualisations, and two statistical tests: 1) Differential barcode proportion (differences in proportions between sample groups), and 2) Differential barcode occurrence (differences in presence/absence odds between groups). Both tests handle complex experimental designs using regression models and rigorously account for sample-to-sample variability. We validated both tests on semi-simulated, real data and a case study, demonstrating that they hold their size, are sufficiently powered to detect true differences, and outperform existing approaches.

Contact: Alicia.oshlack@petermac.org

Keywords

lineage tracing, clonal tracking, bioinformatics, data analysis, single-cell, proportion analysis, DE analysis

Introduction

Cellular DNA barcoding is a powerful cell clonal tracking lineage tracing) technique also referred to as lineage tracing. It incorporates unique DNA barcodes into a cell's genome, thereby genetically labelling each cell in a population.[1] The DNA barcodes are inherited by daughter cells, allowing the origin of each progeny cell to be identified. Collectively, an initial cell and all its progeny are referred to as a "clone" or "lineage". Tracking clonal behaviour reveals underlying clonal heterogeneity, leading to widespread use of DNA barcoding in studies of haematopoietic stem cell differentiation, tumour cell evolution, and embryonic development.[2–7] While each barcode serves as a unique clonal identifier, the total number of cells in each clone can be determined by sequencing and counting the barcode frequencies, thereby estimating relative clonal abundance. Based on the observed clonal abundance patterns, many studies then investigate the features of the initial cells, such as gene expression, gene activity and epigenomic patterns, that relate to clonal outcomes.[8–11]

Cellular genetic barcoding involves several steps, including barcode sequence design, barcode incorporation into cells, and barcode sequencing, each supported by several published protocols.[12–24] These protocols have inspired a series of bespoke workflows, protocol-specific bioinformatics tools, and generalised software for barcode sequence recovery.[25–32] Typically, next-generation-sequencing (NGS) combined with barcode-specific PCR amplification is used to efficiently capture barcode-specific DNA from bulk cell samples. The *CellBarcodeSim* software can simulate barcode data, including the technical effects introduced throughout the barcoding processes.[32,33]

While barcode extraction and alignment approaches vary, these pipelines eventually output a matrix of
barcode read counts across samples. A barcode matrix typically contains thousands of barcodes across tens
to hundreds of samples. The overall read count for a single barcode represents the number of cells derived
from the originally barcoded cell. However, in studies such as *in vivo* haematopoietic stem cell tracking, not
all progeny cells can be collected, so barcode counts reflect the relative abundance of each lineage in a
sample. This contrasts with studies such as *in vitro* tumour heterogeneity tracking, where all progeny cells
can be collected and sequenced, allowing for barcode counts to be interpreted as absolute abundance.
However, even in the second scenario, researchers may not sequence all of progeny cells collected or know
the exact total cell count. Consequently, it is more common and practical to interpret barcode count as relative
or proportional abundance.

Several bioinformatics tools that offer a range of visualization approaches and heuristic interpretation
strategies have been developed to analyse barcode count matrixes. *barcodetrackR* on Bioconductor, and
CellDestiny and *bartools* on GitHub[28–31,34–37] are widely used and share some common features. All three tools
typically use heatmaps to visualise the abundance of each barcode per sample. *BarcodetrackR* and *bartools*
display the log normalised barcode proportion or counts per million (CPM) for selected barcodes (e.g. top 10
most abundant per sample), whilst *CellDestiny* displays the arcsine transformed proportion across all
barcodes. Additionally, *CellDestiny* provides line plots and bubble plots, whilst *bartools* provides stacked bar
plots and bubble plots for displaying barcode abundance across multiple samples.

Furthermore, all three tools also summarise sample characteristics using the per-sample Shannon diversity
index and number of features (detected unique barcodes per sample). For assessing sample differences,
barcodetrackR uses MDS plots and *bartools* uses PCA plots. Both assess sample similarities by displaying
sample pair-wise correlations using the Pearson correlation coefficient. Notably, *CellDestiny* characterises the
distribution of barcode abundance within each sample by visualising the cumulative barcode proportion per
sample.

However, the three tools differ in how they characterize barcode changes between sample groups.
barcodetrackR calculates the log fold change (LFC) of barcode proportions between two groups of samples
and categorizes barcodes using a heuristic threshold. *CellDestiny* compares relative barcode abundance
across groups by first applying column-wise and then row-wise normalisation to the barcode count matrix,
then classifying the barcodes as either group-specific or universal based on whether each group passes a
heuristic threshold. Neither of these tools provides a statistical test. *bartools* tests the odds of barcode
presence/absence between two groups of progeny cells using a hypergeometric test, typically applied in the
context of two clusters in single cell barcoding data.

Table 1. Feature comparison of existing tools for barcode count analysis

		barcodetrackR	CellDestiny	bartools
barcode abundance per sample	Visualisation	Heat map	Heat map Bubble plot Line plot	Heat map Bubble plot Stacked bar plot
	Normalization	Log proportion	Arcsine proportion	Log CPM
	Filtering	Top 10 most abundant	Rely on QC	Top 10 most abundant
Sample characteristics	Characteristics	Shannon Index Num. of features	Shannon Index Num. of features Distribution of barcode abundance per sample	Shannon Index Num. of features
	Dissimilarity	MDS plot		PCA plot
	Similarity	Sample pair Pearson correlation		Sample pair Pearson correlation
Barcode characteristics	Abundance	Compare LFC on proportion to a threshold	Compare row-normalised proportion to a threshold	
	Presence/absence			Hypergeometric test on odds of presence/absence

Lineage tracing experiments often involve complex experimental designs, including multiple time points, genotypes, treatments, and both biological and technical replicates. For example, in a xenograft-based haematopoiesis study, the clonal output was determined by a combination of different cell types, tissues and time points from which cells were collected.[28,38] Despite this complexity, existing tools lack statistical frameworks capable of appropriately testing for differences between groups of samples. Specific limitations include,

1) The LFC approach, the relative proportion approach or the hypergeometric test cannot directly handle multiple conditions within a single factor.
2) None of the approaches can robustly address within-group variability. Technical replicates can be aggregated by *CellDestiny* and *bartools*, however, they cannot rigorously address the differences within replicates and between samples.
3) Both LFC and relative proportion approaches rely on heuristic thresholds to characterise differences in barcode abundance between samples, which can result in subjective and unreproducible interpretations.

As such, there are a lack of robust and rigorously validated statistical testing models specifically designed for population based (bulk) barcode count data that can adequately address barcode variability between conditions. A study associated with the DEBRA approach has explored adapting the *DESeq*, *DESeq2*, and *edgeR* models for barcode count data, however, it lacks thorough validation and comprehensive assessment of the performance of these models.[39–41]

This paper presents the *barbieQ* R Bioconductor package, which provides barcode normalization, data-driven filtering, rich visualisations and statistical testing accommodating the multifactorial designs that are common in the lineage tracing field to enable reproducible analysis.[42] The statistical tests in *barbieQ* build upon a linear regression framework to account for variation in barcodes arising from various sources and can robustly detect significant changes in both barcode abundance and barcode presence/absence between groups of samples. We have used four publicly available datasets to demonstrate and validate *barbieQ*. We show that *barbieQ* controls false discovery rate using both real and semi-simulated datasets and demonstrate its ability to detect real barcode differences in public data.

Results

Package overview

The *barbieQ* package workflow consists of data preprocessing and statistical testing, supported by corresponding visualisations. Briefly, preprocessing loads in the barcode counts and produces a "*barbieQ*" object, preparing data for efficient and accurate interpretation. Statistical testing of the preprocessed "*barbieQ*" object identifies barcodes with significant changes between sample groups in either occurrence or proportion. These results can be visualised using dedicated functions. (Figure 1)

More specifically, preprocessing starts by using "createBarbieQ" to read in a *matrix* of the barcode counts, optionally including a *data.frame* with sample metadata, which initializes a "*barbieQ*" object that is updated throughout the workflow. The "*barbieQ*" object is built upon the *SummarizedExperiment* structure in *R*, consisting of assays storing raw and processed count matrices, barcode-oriented metadata, and sample-oriented metadata. Based on the imported barcode count matrix, the "createBarbieQ" function calculates the barcode counts per million (CPM), proportion and ranking per sample, as well as the presence/absence (occurrence), and saves the results for each metric as a separate matrix in assays. The definitions of those metrics are listed in Table 2. The function also calculates the diversity index, number of unique barcodes (n_barcode) and number of total barcode counts (n_count) for each sample and saves those sample features as part of the sample data (column data), along with any imported sample metadata.

Next, "clusterCorrelatingBarcodes" and "collapseBarcodes" can be used to detect and collapse any
barcodes co-occurring in the same cell. "plotBarcodePairCorrelation" can be used to inspect the
correlation between every pair of barcodes. Filtering barcodes with, "tagTopBarcodes" and
"subsetBarcodes" tags barcodes as top/bottom contributors and filters out low-count/low-contribution
barcodes. Visulations using "plotBarcodeProportion", "plotBarcodePerato" and
"plotBarcodeSankey" can be used for inspecting each barcode's contribution before and after filtering.
Each step stores the results in the "*barbieQ*" object (Table 2), and recalculates assays such as CPM,
proportion, rank, and occurrence when barcodes are collapsed or filtered.

The preprocessed "*barbieQ*" object, generated by collapsing barcodes within the same cell and removing low-
abundance barcodes, provides a clean set of barcodes for statistical testing and allows accurate estimation
of barcode proportions.

133
<div align="center">Table 2. barbieQ object structure and components</div>

Type	Component	Definition	Initiated by Function
Assay	Count	Barcode count per sample	createBarbieQ
	CPM	Barcode count / total barcode count per sample * 1M	createBarbieQ
	Proportion	Barcode count / total barcode count per sample	createBarbieQ
	Occurrence	Presence/absence: barcode CPM >=1 or CPM <1	createBarbieQ
	Rank	Rank of barcode count per sample	createBarbieQ
	isTop	Top/bottom: barcode being main contributor or low count per sample	tagTopBarcodes
Sample data (*columnData*)	SampleMetadata	Sample experimental conditions (factors)	createBarbieQ
	N_barcode	Number of detected barcodes (total occurrence) per sample	createBarbieQ
	N_count	Number of total barcode counts (library size) per sample	createBarbieQ
	Diversity index	Sample Shannon diversity index	createBarbieQ
Barcode data (*rowData*)	Cluster	Barcode cluster of high correlation	clusterCorrelatingBarcodes
	isTop	Barcode determined as top/bottom considering all samples	tagTopBarcodes
	Significance	Significance of barcode abundance changes between sample groups	testBarcodeSignif

For statistical testing, "testBarcodeSignif" is applied to a "*barbieQ*" object and requires a user-defined design
matrix indicating the sample conditions and other experimental factors, as well as user-defined contrasts that
specify which sample groups to compare. Users can choose to test either differential occurrence or differential
proportions of barcodes between the specified sample groups. The results are stored in barcode data in the
"*barbieQ*" object (Table 2) and can be visualised by applying "plotBarcodeMA", "plotBarcodePValue",
"plotSignifBarcodeProportion", or "plotSignifBarcodeHeatmap" to the "*barbieQ*" object.

Additional visualizations, such as "plotSamplePairCorrelation", "plotBarcodeHeatmap" and barcode
PCA, which use barcode CPM, are available before or after statistical testing, to inspect sample heterogeneity
and barcode variability.

Figure 1. Workflow of *barbieQ* package. *barbieQ* consists of three modules: preprocessing, statistical testing, and visualisation. 1) Preprocessing starts by creating a "*barbieQ*" object, and updates it by collapsing highly correlated barcodes, and filtering out lowly abundant barcodes, ensuring an accurate and clean set of barcodes for subsequent interpretation and visualization. Dedicated functions for each step are listed. 2) Statistical testing applies to the preprocessed "*barbieQ*" object with choices for differential occurrence or differential proportion test, and stores the results in the object, which can be visualised. 3) Dedicated visualisations are available at any stage.

Barcode Preprocessing with barbieQ

We demonstrate the effectiveness of the implemented preprocessing functions in barbieQ by applying to the example dataset of 30 selected samples from the monkey HSPC data.

Collapsing highly correlated barcodes

The "clusterCorrelatingBarcodes" function is used to identify clusters of highly correlated barcodes, which are likely barcodes that have been integrated into the same initial cell and, therefore, co-exist in the progeny cells. This phenomenon has been previously observed and, if not properly accounted for, leads to overcounting of cells and distortion of proportions violating the assumption of one-to-one mapping of barcodes to cells.[31] To address the issue, this function calculates the correlation of the proportions across samples of every pair of barcodes and classifies highly correlated pairs based on thresholds for correlation coefficient and mean proportion. The function provides several options for correlation methods and proportion transformation methods; by default, it uses Pearson correlation coefficient (r) of asin-sqrt transformed proportion. Highly correlated pairs are grouped into clusters containing two or more barcodes. Barcodes within a cluster are considered to co-exist in one clone and should be merged. "collapseBarcodes" selects the representative barcode as the one with the highest mean proportion to maintain a consistent barcode-clone association.

We applied this approach to the monkey HSPC data. For efficiency, low-count barcodes were filtered out beforehand using the methods described in the next section. Barcode pairs with a correlation coefficient greater than 0.95 and an average mean proportion exceeding 0.001 were classified as paired barcodes belonging to a cluster (Figure 2A). In total, 9 barcodes formed 4 clusters (Figure 2B), each containing 2~3 barcodes, and the mean proportion of the barcodes within a cluster varied (Figure 2C). These barcode

172 clusters were subsequently collapsed for downstream analysis. These findings are consistent with a
173 published study, and unpublished data we analysed, where paired barcodes were validated through single-
174 cell RNA sequencing of the initial cells, confirming the co-existence and co-expression of the DNA barcodes
175 pairs (Supplementary Figure 1A).[31] Preprocessing results for three additional datasets (AML data, HSPC
176 xenograft data, and mixture data) can be found in Supplementary Figure 1.

Barcode filtering

177
The "tagTopBarcodes" function uses a data-driven approach to classify barcodes as "top" and "bottom"
contributors, guiding filtering to retain "top" barcodes that contribute substantially to the total counts while
removing "bottom" barcodes with minimal contributions. Barcode proportions are calculated for each sample
and ranked from highest to lowest. A cumulative proportion is then calculated at each rank, starting from the
most abundant barcode. Barcodes ranked ahead of the cumulative proportion cutoff at an empirical threshold
(default = 0.99) are tagged as "top" in a sample, indicating they are the most abundant. Across the dataset,
barcodes tagged as "top" in at least n samples (n being the minimum sample group size) are classified as
"top" and retained across the whole dataset. This approach ensures that consistently low-abundance
barcodes are filtered out while preserving those present in at least one sample group.

We applied this approach to the monkey HSPC dataset, using the default proportion cutoff of 0.99 and a
minimum group of n = 6; this was chosen because the 30 samples feature 5 groups of cell types with a
minimum group size of 6. This resulted in the classification of 1140 barcodes as "top" barcodes out of a total
of 16598. The mean proportion of each barcode was visualised using the *barbieQ* function
"plotBarcodeProportion" (Figure 2D). These "top" barcodes represent 6.9% of all barcodes accounting for
91.3% of total barcode counts in the dataset as shown in Figure 2E using the "plotBarcodeSankey"
visualisation. By filtering out consistently low abundance "bottom" barcodes, we retained a focused, clean,
and interpretable set of "top" barcodes (Figure 2F).

Figure 2. Preprocessing results on monkey HSPC data. (**A**) Mean proportion (asin-sqrt transformed) and Pearson correlation coefficients for each pair of barcodes out of the selected not-low-count barcodes using the "plotBarcodePairCorrelation" function; (**B**) Cluster of highly correlated barcodes visualised using the "clusterCorrelatingBarcodes" function; (**C**) Number of barcodes in each cluster (**top**) and Log mean CPM of each barcode within its cluster (**bottom**); (**D**) Proportion of each barcode's mean CPM out of all barcodes (outside circle), color-coded by "top" or "bottom" contributors as tagged by the filtering strategy (inside circle), using the "plotBarcodePareto" function; (**E**) Fraction of "top" or "bottom" contributor barcodes and their cumulative mean proportion, using the "plotBarcodeSankey" function; (**F**) Heatmap of barcodes on proportion before (**left**) and after (**right**) filtering as visualised using the "plotBarcodeHeatmap" function; heatmap colour scale mapped to asin-sqrt transformed proportion but labelled by raw proportion.

Performance of *barbieQ* statistical testing

To assess differences in barcode counts between sample groups that represent clonal outcomes, *barbieQ* can analyse barcode changes in two ways: differential proportion and differential occurrence. Differential proportion identifies barcodes with significantly different proportions between the samples of two conditions, whilst differential occurrence identifies barcodes that are more frequently present in samples of one condition compared to another.

Both statistical tests are based on linear regression models that assess each barcode individually, while accounting for barcode variability between samples groups. A key feature of regression models is their ability

to handle complex experimental designs using a design matrix that accounts for multiple factors and
estimates the variance associated with each factor. This enables the tests to determine whether the factor of
interest significantly changes with respect to barcode occurrence or proportion, while accounting for the
effects of other confounding factors. The differential proportion test leverages the "*limma*" linear regression
framework.[43,47] Inspired by Phipson et al. 2022 it models barcode proportions after applying an arcsine square
root (asin-sqrt) transformation to stabilize the variance across the proportion range to help satisfy the
assumptions of the linear regression.[48] The logit transformation is an alternative transformation available in
barbieQ. Differential occurrence uses a logistic regression model combined with Firth's penalised likelihood to
reduce bias in the maximum likelihood estimates of regression coefficients.[44,49] This correction addresses
inflated standard error that can arise in extreme cases of perfect separation, which are more likely to occur
in small sample size cases (e.g., n = 3), a scenario not uncommon in clonal tracking studies. The model tests
barcode presence or absence in each sample as a binary value, as determined by CPM ≥ threshold (default
to 1) for presence, and CPM < threshold for absence.

The function "testBarcodeSignif" allows users to specify the statistical test and define the factor of interest
through a design matrix with the desired contrast. The results of the test are saved to the *rowData* of the
barbieQ object, aligned with the corresponding barcodes. They contain key statistics such as group-wise
means and differences in proportion or odds ratios in occurrence, along with *P*-values adjusted for multiple
testing using the Benjamin-Hochberg method by default.[50] The test results can be visualised using the
functions listed in Figure 1.

Type I error control under null scenario using real data
To evaluate the Type I error (false positives) of the statistical tests in *barbieQ*, we applied them to a variety of
semi-simulated "null" datasets. From several datasets (mixture data, HSPC xenograft data, and AML data),
we selected sets of samples with homogeneous conditions, where no biologically meaningful differences
were expected. For each sample set, we generated 100 simulated null datasets by randomly assigning
samples into two groups without replacement. We then applied the "testBarcodeSignif" function to perform
differential occurrence and differential proportion tests, assessing the number of barcodes with significant
changes (*P*-value < 0.05) between the two groups, which represent false positives. We also varied the
number of samples per group to determine the effect of group size on Type I error rate (n = 3, 4, ..., up to half
the total number of the null samples).

Figure 3A shows the design of the Mixture dataset. Cells from Pool1 and Pool2 were counted and mixed in
equal ratios to produce a mixed pool. Twelve baseline samples (each from the mixture of Pool 1 and Pool 2
in equal proportions) with varying total cell numbers (n = 20, 40, 80, 160, 330, and 660k, two replicates each)
were used for Type 1 error evaluation as relative barcode abundance should be consistent regardless of the
technical effects related to total cell number.

Barcode proportions were generated by adding 1 to the counts, addressing undersampling artifacts arising
from low library sizes in some samples. Our filtering retained 1553 barcodes from the total 3998
(Supplementary Figure1). To investigate how transformation influences the differential proportion test, we
applied the test after the arcsine square root (asin-sqrt), logit and no transformation. We found that the
differential proportion test, across all transformations controlled the median Type I error rate below 0.05
across 100 simulations (Figure 3C). The differential occurrence test also held its size (Figure 3C). We further
binned barcodes into 10 quantiles based on their proportion (Figure 3D), with the asin-sqrt transformation
showing the most consistent performance across bins while logit transformation resulted in a higher Type 1
error for lower proportions and no transformation had the reverse trend (Figure 3E).

We also assessed the differential proportion test when the number of barcodes in an experiment varied,
noting that barcode proportions generally decrease as the number of barcodes increase. To do this
generated datasets with differing total numbers of barcodes (N = 12, 32, 64, 101, 169, 269, 394, 617, 1060,
1553) by applying various filtering thresholds (0.1, 0.2, 0.3, .4, 0.5, 0.6, 0.7, 0.8, 0.9, 0.95) in the
tagTopBarcode function to the Mixture data, and extracting the top barcodes (Figure 3G). We re-calculated

the barcode proportion within each dataset and repeated the Type I error rate assessment as previously
described. The three transformations showed similar performance and the Type I error rate became more
stable towards 0.05 as barcode size increased (Figure 3H). In addition, we assessed the Type I error rate of
the differential proportion test as a function of the number of barcodes by using the raw barcode proportion
for various sets of barcodes from the Mixture data. The raw proportion is generally smaller than the updated
proportion (Figure 3G). Of the three transformations, asin-sqrt showed the most consistent Type I error rate
control across barcode number (Figure 3H).

Finally, we evaluated Type I error rate in two further datasets (Supplementary Figure 2). For the HSPC
xenograft data, samples collected from the same donor (C21, C22 and C23) were treated as null samples,
despite the differences in time points, tissues, and cell types. For the AML data, samples treated by DMSO
were taken as null samples, despite being collected at different times. All tests maintained Type I error rates
under 0.05. A similar trend in terms of the effect of transformations was also observed in these data.

Power to detect true positives in real data

Using a second subset of samples in the preprocessed Mixture data, we evaluated the power of the
differential proportion test to detect true positives. Twenty-four perturbed samples were generated by
introducing additional Pool1 cells into the baseline level: sampling a certain number of cells from the mixed
pool (n= 20, 40, 80 or 160 thousand) and adding a certain ratio of cells from Pool1 (α = 0.18, 0.27, or 0.35),
with 2 replicates in each case. Specifically, (n × α) cells from Pool1 were added to the n cells from the mixed
pool to generate a perturbed sample. The 12 baseline samples were considered as perturbation level α = 0.
(Figure 3A) This design results in a known difference in barcode proportions (for all barcodes) between
samples with different α.

We gauged the power of the differential proportion test based on the fraction of barcodes with *P*-values <
0.05, obtained from testing three contrasts using the "testBarcodeSignif" function: group$_{\alpha=0.18}$, group$_{\alpha=0.27}$,
and group$_{\alpha=0.35}$ v.s. group$_{\alpha=0}$, respectively, each with asin-sqrt, logit and no transformation (see methods). To
assess the power at different proportion ranges, we binned barcodes into 10 quantiles based on their
proportion as previously described. Across all transformations, the differential proportion test tended to detect
more significance in quantile bins of higher proportion or under stronger contrast, but logit generally showed
lower power (Figure 3F) except at the smallest proportions.

As before, we also assessed the power of the differential proportion test under scenarios of different numbers
of barcodes. We generated datasets by selecting various numbers of top barcodes from the Mixture data as
previously described, and likewise, re-calculated barcode proportion per sample and repeated the power
assessment for each dataset. The three transformations showed similar power, with all demonstrating greater
power with lower number of barcodes and greater proportions (Figure 3I). To investigate the impact of the
way proportion is calculated, we repeated this process by using the raw barcode proportion when taking
subsets of barcodes at each barcode size and found asin-sqrt showed greater power than logit across
barcode sizes (Figure 3I). All tests showed greater power under stronger contrast.

The coefficients of the contrasts obtained from the "testBarcodeSignif" function are the estimated
differences between sample groups for each barcode which can be predicted from the design (see methods).
Both asin-sqrt and no transformation accurately estimated the barcode proportion change, while logit tended
to over-estimate the change (Supplementary Figure 3A). Residuals from asin-sqrt-transformed proportion
were the most Normally distributed (Supplementary Figure 3B), in line with the assumption of the linear
regression model implemented in the "testBarcodeSignif" function.

Figure 3. Performance of *barbieQ* statistical testing. (**A**) **Schematic of mixture experiment** showing the generation of baseline
samples and perturbed samples in the Mixture dataset; (**B**) **Barcode proportion data used in evaluations** displaying proportion per
sample; Samples are annotated by total number of cells and perturbation levels; (**C**) **Type** I **error rates** of (**left**) differential proportion
test with three transformation methods and (**right**) differential occurrence test, taking the faction of barcodes with raw *P*-value < 0.05

obtained from each simulation (100 simulations per group size); (**D**) **Barcode proportion quantiles:** boxplot of barcode mean
proportion across baseline samples in each group, binned into 10 quantiles of the mean proportion; (**E**) **Type** I **error rate as a**
function of barcode quantiles: taking the fraction of barcodes with P-value < 0.05 within each quantile bin from Figure 3**D** using
differential proportion test with three transformations in baseline samples; (**F**) As in **E** but showing power as determined by the fraction
of p-values<0.05 in perturbed samples; (**G**) **Density of barcode proportion** in datasets generated by isolating different sets of
barcodes, (**top**) with proportion updated per sample or (**bottom**) keeping raw proportion; datasets color-coded by number of total
barcodes in datasets; (**H**) **Type** I **error rate as a function of total barcode number:** taking the faction of barcodes with P-value <
0.05 obtained from testing each set of barcodes in Figure 3**F** using differential proportion test with three transformations from baseline
samples; (**I**) As in **H** but showing power as determined by the fraction of p-values <0.05 in perturbed samples.

Case study

Reanalysis of monkey HSPC data

Two important features of the *barbieQ* package are its data-driven filtering strategy and robust statistical tests.
To demonstrate how *barbieQ* advances upon previous approaches in the clonal tracking and lineage tracing
fields, we applied the *barbieQ* analytical workflow to a real-world dataset and compared the results with those
reported in the original publication. The example data is a subset of 30 samples from the monkey HSPC
dataset, collected across 6 time points and sorted into 5 different cell types (Figure 4A). In total, 16,603
unique barcodes were observed in this dataset, with barcode proportions ranging from a maximum of 0.14,
a mean of approximately 6×10^{-5} and a median of 0 (Figure 4B). These barcode proportions are variable
between cell types but relatively stable between time points within the same cell type (Figure 4B).

We preprocessed the raw barcode count data following the steps outlined in Figure 2, including aggregating
highly correlated barcodes and filtering low-count barcodes. Compared to the "topN" filtering approaches
used in the original study, *barbieQ* retained a larger number of barcodes (Figure 4C). *barbieQ* retained 8% of
barcodes, accounting for 93% of the total counts while *top10-per-sample* retained 0.5% of barcodes
contributing 43% of total counts, and *top20-per-sample* retained 1% of barcodes contributing 51% of total
counts (Figure 4D). Figure 4E shows a heatmap of barcode logCPM, annotating barcodes retained by both
barbieQ and *top10-per-sample* strategies or exclusively by one. Notably, barcodes exclusively retained by
barbieQ still show informative logCPM across samples. The 4 barcodes retained by *top10-per-sample* only
showed a contribution in only few samples. An additional advantage of *barbieQ* is avoiding arbitrarily selecting
a *topN* threshold for every new dataset, making the filtering process less subjective.

Next, we applied the statistical testing methods implemented in *barbieQ* by reproducing the comparisons from
the original publication using the exact same sets of barcodes. The first analysis included 81 barcodes that
were selected by the *top10-per-sample* filtering strategy out of the 30 samples (Figure 4A, B). We used
barbieQ to test barcode differences between the NK$^{CD56-CD16+}$ samples and other cell types (T, B, Gr and
NK$^{CD56+CD16-}$), treating each individual time point as a factor. We used a design matrix that sets the 5 cell types
as 5 factors and 6 time points as 6 additional factors with the contrast as NK$^{CD56-CD16+}$ - (T + B + Gr +
NK$^{CD56+CD16-}$) / 4. We then applied the differential proportion test, using the "testBarcodeSignif" function,
with the asin-sqrt transformation. We identified 23 barcodes with significantly increased proportion in NK$^{CD56-CD16+}$
samples, including 6 novel barcodes and 17 overlapping with those selected by the original publication,
which used a heuristic cutoff for barcode proportion ratios between NK$^{CD56-CD16+}$ and other cell types at each
time point respectively (Figure 4F). We inspected the log fold change (LFC) in barcode proportions (NK$^{CD56-CD16+}$
/ other cell types) at different time points and observed that the 6 novel barcodes showed robust LFC
although with lower effect size (Figure 4F).

We also applied the same test to a larger set of barcodes retained using the *barbieQ* filtering strategy, including
1,028 barcodes that contribute 91.3% to the 30 samples. We identified three times more barcodes showing
significantly increased proportion in NK$^{CD56-CD16+\ cells}$ and collectively contributing more to the NK$^{CD56-CD16+}$
samples (Figure 4G, H). We inspected the LFC of the significant barcodes obtained based on each filtering
strategy. (Figure 4J) Using the *barbieQ* filtering strategy allowed us to keep more barcodes with robust LFC
and relatively high abundance.

The second analysis used the subset of 8 NK$^{CD56-CD16+}$ samples from the monkey HSPC dataset, collected at
various time points. We used the 39 barcodes retained using the *top10-per-sample* filtering strategy to match
the set of barcodes in the original publication. We then used the "testBarcodeSignif" function to test
barcodes that change over time within NK$^{CD56-CD16+}$ samples. Specifically, each sample was given a numeric
factor for the number of months post transplantation. We identified 17 barcodes with significantly increased
and 8 with decreased proportions over time (Figure 4I). The *barbieQ* testing results robustly categorized
barcode kinetics, compared to the barcodes selected by visual inspection of heatmap dendrograms as used
in the original publication.

Figure 4 Applying *barbieQ* to the monkey HSPC dataset. (**A**) Experimental design for a subset of the monkey HSPC dataset, where 30 samples contained 5 cell types sharing 6 time points; (**B**) Barcode proportion per sample in the subset Monkey HSPC dataset, with each barcode represented by a unique colour; Samples are annotated by cell type and time to collection in months; (**C**) UpSet plot showing the overlap in retained barcodes using *barbieQ* filtering, *top10-per-sample* and *top20-per-sample* strategies; (**D**) Faction of retained barcodes and fraction of their total proportion to the whole dataset. Filtering strategies from left to right: *barbieQ*, *top10-per-sample*, and *top20-per-sample*; (**E**) Heatmap of retained barcodes by either *barbieQ* or *top10-per-sample.* Right annotations indicate the overlap between the two approaches; (**F**) Box plot of the LFC for each barcode. A dot refers to the LFC in proportion between to $NK^{CD56-CD16+}$ and other cell types at a given time points (6 time points in total). Barcode were ordered by *barbieQ*-estimated coefficient for the contrast, from low to high. Barcodes were categorized into identified as $NK^{CD56-CD16+}$-associated by *barbieQ*, heuristic LFC, both and neither method. (**G**) UpSet plot showing the overlap in identified $NK^{CD56-CD16+}$-associated barcodes using *barbieQ* "testBarcodeSignif" based on *barbieQ* filtering and *top10-per-sample* strategies. (**H**) Total barcode proportion per sample for identified $NK^{CD56-CD16+}$-associated barcodes based on *top10-per-sample* (left) and *barbieQ* filtering strategy (right), respectively. (**I**) Box plot of the LFC for each barcode. A dot refers to the LFC in proportion between to $NK^{CD56-CD16+}$ and other cell types at a given time point (6 time points in total). Barcodes were ordered by mean proportion across cell types and time points, from low to high. Barcodes were categorized into identified as $NK^{CD56-CD16+}$-associated by *barbieQ*, based on the set of *barbieQ* filtering and top10-per-sample, and both strategies. (**J**) Heatmap of 39 barcodes retained from 8 $NK^{CD56-CD16+}$ samples using *barbieQ* filtering. Samples were order over time. Right annotations indicate barcodes being identified with significantly increasing, decreasing kinetics or not-significant (n.s.) over time. Barcodes were grouped into 5 patterns using the left dendrogram.

Discussion

Lineage tracing experiments provide powerful insights into cellular development and evolution, but the complexity of the data presents significant analytical challenges. Current approaches, typically based on pairwise comparisons, may fail to account for the inherent complexity of the experimental design, potentially leaving important insights unexplored. Here, we presented *barbieQ*, a new R Bioconductor package tailored to the analysis of lineage tracing barcode count data that provides an integrated platform for quality control, filtering, visualisation and statistical testing. Built upon the *SummarizedExperiment* infrastructure, it ensures compatibility with existing workflows. *barbieQ*'s core function testBarcodeSignif provides statistical tests for identifying differential barcode proportions and differential barcode occurrence. These tests enable the robust identification of barcode differences between sample groups, whilst comprehensively accounting for variability resulting from multiple technical and biological conditions. *barbieQ*'s data-driven approach to preprocessing using the functions clusterCorrelatingBarcodes and tagTopBarcodes, guarantees a clean, reproducible and informative set of barcodes for testing. Additionally, the built-in visualisation functions can be used to inspect barcode features throughout an analysis, ensuring transparency and interpretability at every stage of the workflow.

We demonstrated *barbieQ*'s preprocessing capabilities by applying the clusterCorrelatingBarcodes function to four barcode datasets: monkey HSPC, AML data, HSPC xenograft data, and mixture data. In the first three, we found highly correlated barcode clusters likely to be the multiple barcodes that were incorporated into the same initial cells during preparation. We previously applied this approach in the analysis of an unpublished dataset and identified clusters of highly correlated barcodes, which were also detected in the same initial cells by single cell sequencing. Although this approach was validated as useful for confirming barcode uniqueness, its reliance on heuristic thresholds may limit generalisation across different datasets. We also applied the tagTopBarcodes function to the four datasets to identify the main contributing barcodes across samples. We showed that our data-driven method effectively filters the data whilst retaining the informative barcodes. In the case study, we compared the *barbieQ* filtering strategy to the commonly used *top-10* (or *20*)-per-sample approach in the analysis of Monkey HSPC data, where our approach retained a greater number of potentially essential barcodes.

Using the Mixture data, we evaluated the *barbieQ*'s statistical tests by examining Type I error rate by simulating sample groups under null conditions and the power by detecting known barcode differences using the function testBarcodeSignif. We showed that both statistical tests, differential proportion and differential occurrence, held their sizes. The differential proportion test combines *limma* with commonly-used transformation methods for proportion data; the asin-sqrt and logit transformations, in order to accommodate the assumptions of normally distributed data and mean variance relationship in the linear regression model.[48]

Based on our assessment, we recommend using asin-sqrt in barcode data analysis for its superior
performance. Specifically, we found asin-sqrt transformation had: 1) consistent Type I error rate across
proportion levels, across datasets with different numbers of barcodes and with or without updating proportion
within selected barcodes, 2) greater power to detect true positives, and 3) improved residual Normality.

We also applied the *barbieQ* workflow to a case study analysis of the monkey HSPC data, in which samples
differed across time points and cell types. We repeated analyses from the original paper using the *barbieQ*
differential proportion test and identified a novel set of barcodes with significantly higher proportion in mature
NK cells compared with other cell types. We also robustly identified a set of barcodes showing significantly
increasing proportion over time in the mature NK cell cohort, some of which were only described using a
heatmap in the orginal paper. This demonstrates that *barbieQ* can recapitulate existing results derived using
heuristics and visualisation, whilst robustly and reproducibly identifying additional barcodes of interest.
Furthermore, given that *barbieQ*'s differential proportion test is built upon the versatile *limma* framework, its
usage can be extended beyond simple group comparisons, such as fitting "splines" to identify more complex
barcode kinetics.

In conclusion, *barbieQ* is a powerful R Bioconductor package for robustly and reproducibly detecting
biologically meaningful changes in lineage tracing barcode count data. The *barbieQ* package is available from
Bioconductor.[42]

Methods

Software development and implementation

The *barbieQ* package was developed using R 4.5.0 and Bioconductor version 3.21
(https://www.bioconductor.org/packages/release/bioc/html/barbieQ.html). To ensure high performance and
interoperability within the R Bioconductor ecosystem we leveraged established libraries such as:

- *Dplyr (https://cran.r-project.org/web/packages/dplyr/index.html)* version 1.1.4 for data manipulation, *tydyr*
(https://cran.r-project.org/web/packages/tidyr/index.html) version 1.3.1 for numerical calculations,
SummarizedExperiment
(https://bioconductor.org/packages/release/bioc/html/SummarizedExperiment.html) version 1.37.0
and *S4Vectors* (https://bioconductor.org/packages/release/bioc/html/S4Vectors.html) version 0.45.4
for data structure.

- *ggplot2* (https://cran.r-project.org/web/packages/ggplot2/index.html) version 3.5.1 and
ComplexHeatmap (https://www.bioconductor.org/packages/release/bioc/html/ComplexHeatmap.html) version
2.23.0 for drawing plots.

- and *stats (https://cran.r-project.org/doc/manuals/r-patched/packages/stats/refman/stats.html)* version 4.5.0,
limma (https://bioconductor.org/packages/release/bioc/html/limma.html) version 3.63.11 and *logistif*
(https://cran.r-project.org/web/packages/logistf/index.html) version 1.26.0 for statistical modelling.[43,44]

The package was structured based on the *SummarizedExperiment* S4 class, allowing data manipulation and
processing directly within the data structure. Processed results are saved to back to the same structure,
extended with additional features to seamlessly interoperate with upstream barcode alignment tools that also
use *SummarizedExperiment* or *DGEList* from *edgeR (https://bioconductor.org/packages/release/bioc/html/edgeR.html)*
for data handling.[45]

Data description

Several published datasets, briefly described below, were used to demonstrate preprocessing and evaluate the statistical tests implemented in *barbieQ*. Raw barcode count data were obtained following sequence-level QC and barcode library alignment, which had been completed by the original authors.

Monkey HSPC Data

A subset of publicly available data from a study on monkey hematopoietic stem and progenitor cell (HSPC) clonal expansion *in vivo* using barcoding technique.[46] The monkey HSPC data were analysed using the *barcodetrackR* package and made available via the compatible barcodetrackR Data repository on GitHub (https://github.com/dunbarlabNIH/barcodetrackRData/tree/main/datasets/WuC_etal).[28] Herein, we used data from monkey "ZG66" including 16,603 barcodes across all samples, where we further selected 30 samples to be used here.

Briefly, unique barcodes were initially integrated into the DNA of HSPCs and subsequently passed to progeny cells. At a series of time points, progeny cells were collected from blood and sorted into various cell types, including T cells, B cells, granulocytes (Gr), and natural killer (NK) cells for $NK^{CD56+CD16-}$ and $NK^{CD56-CD16+}$ subtypes. Barcode counts across different cell types were used to interpret the patterns of HSPC differentiation. The original study focused on identifying barcodes (clones) with higher abundance in $NK^{CD56-CD16+}$ samples compared to other cell type samples.

AML Data

Publicly available data from a study of acute myeloid leukaemia (AML) clones, investigating the heterogeneity in their response to various therapeutic drugs *in vitro*, which was originally analysed and introduced with *bartools*.[31] Briefly, AML cells were barcoded at the DNA level using the SPLINTR system, recognized as individual clones, and their population was expanded under exposure to different drugs (Arac, IBET), at various doses and DMSO as a negative control.[22] Samples were collected from each treatment condition at a series of time points. This barcode count matrix contains 1,811 barcodes, across 41 samples, under the described conditions.

HSPC Xenograft Data

Public data from a study of engraftment tracking of human umbilical cord blood HSPC xenotransplanted into mice.[38] The data were analysed in the *barcodetrackR* publication and made available via the compatible barcodetrackR Data repository on GitHub (https://github.com/dunbarlabNIH/barcodetrackRData/tree/main/datasets/BelderbosME_etal).[28] In this study, human cord blood HSPC cells from 20 individual donors were isolated and barcoded at the DNA level respectively. Sets of starting clones from different donors (n = 20) were transplanted into eith one or two recipient mice. For each mouse, progeny cells were collected from peripheral blood at multiple time points, as well as from various tissues. From each collection, cells were sorted into different cell types forming individual samples, with unsorted cells also retained. Herein, we used data of recipient mice with sufficiently engrafted HSPC clones, containing 10,149 barcodes across 8 donors, with 199 samples, under the described conditions.

Mixture Data

This dataset was generated to simulate both true and null changes in barcode abundance by mixing cells from two barcoded cell line samples.[39] Cells from each cell line were divided into two pools (Pool1 and Pool2), each incorporating distinct clonal tracking barcodes into their DNA, ensuring no overlap between pools. Cells in each pool were counted and mixed in an equal ratio to produce a mixed pool. Twelve baseline samples were sampled from the mixed pool at various cell numbers. Twenty-four perturbed samples were generated by sampling a certain number of cells from the mixed pool and adding a certain ratio of cells from Pool1, with 2 replicates in each case. This dataset contains 3998 barcodes, across 36 samples as described, as well as two reference samples representing the unmixed Pool1 and Pool2 barcode counts.

Data analysis

All data was analysed using the functions provided in barbieQ. Complete documentation of these
can be found in the analysis on GitHub (https://github.com/Oshlack/barbieQ-paper-analysis).

True positive calculations for Mixture data

The mixture pool contains a mix of cells with barcodes from Pool1 and Pool2, for which the barcodes didn't
overlap. Let x_n be the proportion of barcode, n = (1, ..., 1553).

507
$$X_{Pool1} = \begin{bmatrix} x_1 \\ \vdots \\ x_m \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \qquad X_{Pool2} = \begin{bmatrix} 0 \\ \vdots \\ 0 \\ x_{m+1} \\ \vdots \\ x_n \end{bmatrix}$$

508 A baseline sample theoretically consisted of an equal mixture of total barcode numbers from Pool1 and
509 Pool2, leading to barcode proportions being half of the reference samples.

510
$$X_0 = \frac{1}{2}(X_{Pool1} + X_{Pool2})$$

511 A sample with a perturbation level of α theoretically leads to a known change in proportion for the barcodes
512 from both Pool1 and Pool2.

513
$$X_\alpha = \frac{\frac{1}{2} + \alpha}{(1 + \alpha)} X_{Pool1} + \frac{\frac{1}{2}}{(1 + \alpha)} X_{Pool2}, \qquad \alpha = 0.18, 0.27, 0.35$$

The designed difference in barcode proportions between perturbed samples and baseline samples can
therefore be calculated for each transformation.

$\Delta = f(X_\alpha) - f(X_0), \qquad where\ f_{asin-sqrt}(x) = \sin^{-1}(\sqrt{x}), f_{logit}(x) = \log\frac{x}{1-x}, f_{no\ trans}(x) = x, \qquad x \in (0,1)$

Therefore, for this data we modelled the 36 samples using a design matrix that set factors for each
perturbation level (α = 0, 0.18, 0.27, and 0.35), with additional factors accounting for the cell number in the
equal mixture (n= 20, 40, 80, 160, 330, and 660 thousand). We then applied the "testBarcodeSignif"
function to test barcode differential proportion under three contrasts: group$_{\alpha=0.18}$, group$_{\alpha=0.27}$, and group$_{\alpha=0.35}$
v.s. group$_{\alpha=0}$, respectively, each with asin-sqrt, logit and no transformation. After preprocessing, a total of
1553 barcodes were tested using their raw proportion, and all barcodes were designed to change under each
contrast.

Acknowledgements

We would like to thank our collaborators, Mark Dawson, Elanor Wainright, Ali Motazedian, Enid Lam, Dane
Vassiliadis, Henrietta Holze, who brought in valuable data, interesting biology questions and inspiring
discussion. We would like to thank our colleagues and friends, Caitlin Page who gave advice on the package
development, and Lizhong Chen who helped improve code efficiency. And we acknowledge the clonal
tracking community, the discussion with whom benefited in a thorough understanding of the background and
potential application for this research.

Funding

This work has been supported by National Health and Medical Research Council [GNT1196256,
GNT2026098].

Conflict of Interest: none declared.

References

(1) Bramlett, C.; Jiang, D.; Nogalska, A.; Eerdeng, J.; Contreras, J.; Lu, R. Clonal Tracking Using Embedded Viral Barcoding and High-Throughput Sequencing. *Nat. Protoc.* **2020**, *15* (4), 1436–1458. https://doi.org/10.1038/s41596-019-0290-z.

(2) Emert, B. L.; Cote, C. J.; Torre, E. A.; Dardani, I. P.; Jiang, C. L.; Jain, N.; Shaffer, S. M.; Raj, A. Variability within Rare Cell States Enables Multiple Paths toward Drug Resistance. *Nat. Biotechnol.* **2021**, *39* (7), 865–876. https://doi.org/10.1038/s41587-021-00837-3.

(3) Gutierrez, C.; Al'Khafaji, A. M.; Brenner, E.; Johnson, K. E.; Gohil, S. H.; Lin, Z.; Knisbacher, B. A.; Durrett, R. E.; Li, S.; Parvin, S.; Biran, A.; Zhang, W.; Rassenti, L.; Kipps, T. J.; Livak, K. J.; Neuberg, D.; Letai, A.; Getz, G.; Wu, C. J.; Brock, A. Multifunctional Barcoding with ClonMapper Enables High-Resolution Study of Clonal Dynamics during Tumor Evolution and Treatment. *Nat. Cancer* **2021**, *2* (7), 758–772. https://doi.org/10.1038/s43018-021-00222-8.

(4) Naik, S. H.; Perié, L.; Swart, E.; Gerlach, C.; van Rooij, N.; de Boer, R. J.; Schumacher, T. N. Diverse and Heritable Lineage Imprinting of Early Haematopoietic Progenitors. *Nature* **2013**, *496* (7444), 229–232. https://doi.org/10.1038/nature12013.

(5) Lin, D. S.; Zhang, S.; Schreuder, J.; Tran, J.; Sargeant, T.; Metcalf, D.; Ng, A. P.; Weber, T. S.; Naik, S. H. A Multi-Track Landscape of Haematopoiesis Informed by Cellular Barcoding and Agent-Based Modelling. bioRxiv March 30, 2024, p 2024.03.28.587126. https://doi.org/10.1101/2024.03.28.587126.

(6) Chan, M. M.; Smith, Z. D.; Grosswendt, S.; Kretzmer, H.; Norman, T. M.; Adamson, B.; Jost, M.; Quinn, J. J.; Yang, D.; Jones, M. G.; Khodaverdian, A.; Yosef, N.; Meissner, A.; Weissman, J. S. Molecular Recording of Mammalian Embryogenesis. *Nature* **2019**, *570* (7759), 77–82. https://doi.org/10.1038/s41586-019-1184-5.

(7) Bandler, R. C.; Vitali, I.; Delgado, R. N.; Ho, M. C.; Dvoretskova, E.; Ibarra Molinas, J. S.; Frazel, P. W.; Mohammadkhani, M.; Machold, R.; Maedler, S.; Liddelow, S. A.; Nowakowski, T. J.; Fishell, G.; Mayer, C. Single-Cell Delineation of Lineage and Genetic Identity in the Mouse Brain. *Nature* **2022**, *601* (7893), 404–409. https://doi.org/10.1038/s41586-021-04237-0.

(8) Biddy, B. A.; Kong, W.; Kamimoto, K.; Guo, C.; Waye, S. E.; Sun, T.; Morris, S. A. Single-Cell Mapping of Lineage and Identity in Direct Reprogramming. *Nature* **2018**, *564* (7735), 219–224. https://doi.org/10.1038/s41586-018-0744-4.

(9) Lin, D. S.; Tian, L.; Tomei, S.; Amann-Zalcenstein, D.; Baldwin, T. M.; Weber, T. S.; Schreuder, J.; Stonehouse, O. J.; Rautela, J.; Huntington, N. D.; Taoudi, S.; Ritchie, M. E.; Hodgkin, P. D.; Ng, A. P.; Nutt, S. L.; Naik, S. H. Single-Cell Analyses Reveal the Clonal and Molecular Aetiology of Flt3L-Induced Emergency Dendritic Cell Development. *Nat. Cell Biol.* **2021**, *23* (3), 219–231. https://doi.org/10.1038/s41556-021-00636-7.

(10) Li, L.; Bowling, S.; McGeary, S. E.; Yu, Q.; Lemke, B.; Alcedo, K.; Jia, Y.; Liu, X.; Ferreira, M.; Klein, A. M.; Wang, S.-W.; Camargo, F. D. A Mouse Model with High Clonal Barcode Diversity for Joint Lineage, Transcriptomic, and Epigenomic Profiling in Single Cells. *Cell* **2023**, *186* (23), 5183-5199.e22. https://doi.org/10.1016/j.cell.2023.09.019.

(11) Laisné, M.; Lupien, M.; Vallot, C. Epigenomic Heterogeneity as a Source of Tumour Evolution. *Nat. Rev. Cancer* **2025**, *25* (1), 7–26. https://doi.org/10.1038/s41568-024-00757-9.

(12) VanHorn, S.; Morris, S. A. Next-Generation Lineage Tracing and Fate Mapping to Interrogate Development. *Dev. Cell* **2021**, *56* (1), 7–21. https://doi.org/10.1016/j.devcel.2020.10.021.

(13) Chen, C.; Liao, Y.; Peng, G. Connecting Past and Present: Single-Cell Lineage Tracing. *Protein Cell* **2022**, *13* (11), 790–807. https://doi.org/10.1007/s13238-022-00913-7.

(14) Deng, L.-H.; Li, M.-Z.; Huang, X.-J.; Zhao, X.-Y. Single-Cell Lineage Tracing Techniques in Hematology: Unraveling the Cellular Narrative. *J. Transl. Med.* **2025**, *23* (1), 270. https://doi.org/10.1186/s12967-025-06318-4.

(15) Sun, J.; Ramos, A.; Chapman, B.; Johnnidis, J. B.; Le, L.; Ho, Y.-J.; Klein, A.; Hofmann, O.; Camargo, F. D. Clonal Dynamics of Native Haematopoiesis. *Nature* **2014**, *514* (7522), 322–327. https://doi.org/10.1038/nature13824.

(16) Wagner, D. E.; Weinreb, C.; Collins, Z. M.; Briggs, J. A.; Megason, S. G.; Klein, A. M. Single-Cell Mapping of Gene Expression Landscapes and Lineage in the Zebrafish Embryo. *Science* **2018**, *360* (6392), 981–987. https://doi.org/10.1126/science.aar4362.

(17)   Spanjaard, B.; Hu, B.; Mitic, N.; Olivares-Chauvet, P.; Janjuha, S.; Ninov, N.; Junker, J. P. Simultaneous Lineage Tracing and Cell-Type Identification Using CRISPR–Cas9-Induced Genetic Scars. *Nat. Biotechnol.* **2018**, *36* (5), 469–473. https://doi.org/10.1038/nbt.4124.

(18)   Kong, W.; Biddy, B. A.; Kamimoto, K.; Amrute, J. M.; Butka, E. G.; Morris, S. A. CellTagging: Combinatorial Indexing to Simultaneously Map Lineage and Identity at Single-Cell Resolution. *Nat. Protoc.* **2020**, *15* (3), 750–772. https://doi.org/10.1038/s41596-019-0247-2.

(19)   Weinreb, C.; Rodriguez-Fraticelli, A.; Camargo, F. D.; Klein, A. M. Lineage Tracing on Transcriptional Landscapes Links State to Fate during Differentiation. *Science* **2020**, *367* (6479), eaaw3381. https://doi.org/10.1126/science.aaw3381.

(20)   Bowling, S.; Sritharan, D.; Osorio, F. G.; Nguyen, M.; Cheung, P.; Rodriguez-Fraticelli, A.; Patel, S.; Yuan, W.-C.; Fujiwara, Y.; Li, B. E.; Orkin, S. H.; Hormoz, S.; Camargo, F. D. An Engineered CRISPR-Cas9 Mouse Line for Simultaneous Readout of Lineage Histories and Gene Expression Profiles in Single Cells. *Cell* **2020**, *181* (6), 1410-1422.e27. https://doi.org/10.1016/j.cell.2020.04.048.

(21)   Oren, Y.; Tsabar, M.; Cuoco, M. S.; Amir-Zilberstein, L.; Cabanos, H. F.; Hütter, J.-C.; Hu, B.; Thakore, P. I.; Tabaka, M.; Fulco, C. P.; Colgan, W.; Cuevas, B. M.; Hurvitz, S. A.; Slamon, D. J.; Deik, A.; Pierce, K. A.; Clish, C.; Hata, A. N.; Zaganjor, E.; Lahav, G.; Politi, K.; Brugge, J. S.; Regev, A. Cycling Cancer Persister Cells Arise from Lineages with Distinct Programs. *Nature* **2021**, *596* (7873), 576–582. https://doi.org/10.1038/s41586-021-03796-6.

(22)   Fennell, K. A.; Vassiliadis, D.; Lam, E. Y. N.; Martelotto, L. G.; Balic, J. J.; Hollizeck, S.; Weber, T. S.; Semple, T.; Wang, Q.; Miles, D. C.; MacPherson, L.; Chan, Y.-C.; Guirguis, A. A.; Kats, L. M.; Wong, E. S.; Dawson, S.-J.; Naik, S. H.; Dawson, M. A. Non-Genetic Determinants of Malignant Clonal Fitness at Single-Cell Resolution. *Nature* **2022**, *601* (7891), 125–131. https://doi.org/10.1038/s41586-021-04206-7.

(23)   Goyal, Y.; Busch, G. T.; Pillai, M.; Li, J.; Boe, R. H.; Grody, E. I.; Chelvanambi, M.; Dardani, I. P.; Emert, B.; Bodkin, N.; Braun, J.; Fingerman, D.; Kaur, A.; Jain, N.; Ravindran, P. T.; Mellis, I. A.; Kiani, K.; Alicea, G. M.; Fane, M. E.; Ahmed, S. S.; Li, H.; Chen, Y.; Chai, C.; Kaster, J.; Witt, R. G.; Lazcano, R.; Ingram, D. R.; Johnson, S. B.; Wani, K.; Dunagin, M. C.; Lazar, A. J.; Weeraratna, A. T.; Wargo, J. A.; Herlyn, M.; Raj, A. Diverse Clonal Fates Emerge upon Drug Treatment of Homogeneous Cancer Cells. *Nature* **2023**, *620* (7974), 651–659. https://doi.org/10.1038/s41586-023-06342-8.

(24)   Weber, T. S.; Biben, C.; Miles, D. C.; Glaser, S. P.; Tomei, S.; Lin, C.-Y.; Kueh, A.; Pal, M.; Zhang, S.; Tam, P. P. L.; Taoudi, S.; Naik, S. H. LoxCode in Vivo Barcoding Reveals Epiblast Clonal Fate Bias to Fetal Organs. *Cell* **2025**, *0* (0). https://doi.org/10.1016/j.cell.2025.04.026.

(25)   Lin, D. S.; Kan, A.; Gao, J.; Crampin, E. J.; Hodgkin, P. D.; Naik, S. H. DiSNE Movie Visualization and Assessment of Clonal Kinetics Reveal Multiple Trajectories of Dendritic Cell Development. *Cell Rep.* **2018**, *22* (10), 2557–2566. https://doi.org/10.1016/j.celrep.2018.02.046.

(26)   Rodriguez-Fraticelli, A. E.; Wolock, S. L.; Weinreb, C. S.; Panero, R.; Patel, S. H.; Jankovic, M.; Sun, J.; Calogero, R. A.; Klein, A. M.; Camargo, F. D. Clonal Analysis of Lineage Fate in Native Haematopoiesis. *Nature* **2018**, *553* (7687), 212–216. https://doi.org/10.1038/nature25168.

(27)   Thielecke, L.; Cornils, K.; Glauche, I. genBaRcode: A Comprehensive R-Package for Genetic Barcode Analysis. *Bioinformatics* **2020**, *36* (7), 2189–2194. https://doi.org/10.1093/bioinformatics/btz872.

(28)   Espinoza, D. A.; Mortlock, R. D.; Koelle, S. J.; Wu, C.; Dunbar, C. E. Interrogation of Clonal Tracking Data Using barcodetrackR. *Nat. Comput. Sci.* **2021**, *1* (4), 280–289. https://doi.org/10.1038/s43588-021-00057-4.

(29)   Hadj Abed, L.; Tak, T.; Cosgrove, J.; Perié, L. CellDestiny: A RShiny Application for the Visualization and Analysis of Single-Cell Lineage Tracing Data. *Front. Med.* **2022**, *9*, 919345. https://doi.org/10.3389/fmed.2022.919345.

(30)   Putri, G. H.; Pires, N.; Davidson, N. M.; Blyth, C.; Al'Khafaji, A. M.; Goel, S.; Phipson, B. Extraction and Quantification of Lineage-Tracing Barcodes with NextClone and CloneDetective. November 20, 2023. https://doi.org/10.1101/2023.11.19.567755.

(31)   Holze, H.; Talarmain, L.; Fennell, K. A.; Lam, E. Y.; Dawson, M. A.; Vassiliadis, D. Analysis of Synthetic Cellular Barcodes in the Genome and Transcriptome with BARtab and Bartools. *Cell Rep. Methods* **2024**, 100763. https://doi.org/10.1016/j.crmeth.2024.100763.

(32)   Sun, W.; Perkins, M.; Huyghe, M.; Faraldo, M. M.; Fre, S.; Perié, L.; Lyne, A.-M. Extracting, Filtering and Simulating Cellular Barcodes Using CellBarcode Tools. *Nat. Comput. Sci.* **2024**, *4* (2), 128–143. https://doi.org/10.1038/s43588-024-00595-7.

(33)   *TeamPerie/CellBarcodeSim at v1*. GitHub. https://github.com/TeamPerie/CellBarcodeSim (accessed 2025-09-11).

(34)   *barcodetrackR*. Bioconductor. http://bioconductor.org/packages/barcodetrackR/ (accessed 2025-09-11).

(35)   TeamPerie/CellDestiny, 2025. https://github.com/TeamPerie/CellDestiny (accessed 2025-09-11).

(36)   Phipsonlab/CloneDetective, 2023. https://github.com/phipsonlab/CloneDetective (accessed 2025-11-07).

(37)   Vassiliadis, D. DaneVass/Bartools, 2025. https://github.com/DaneVass/bartools (accessed 2025-09-11).

(38)   Belderbos, M. E.; Jacobs, S.; Koster, T. K.; Ausema, A.; Weersing, E.; Zwart, E.; Haan, G. de; Bystrykh, L. V. Donor-to-Donor Heterogeneity in the Clonal Dynamics of Transplanted Human Cord Blood Stem Cells in Murine Xenografts. *Biol. Blood Marrow Transplant.* **2020**, *26* (1), 16–25. https://doi.org/10.1016/j.bbmt.2019.08.026.

(39)   Akimov, Y.; Bulanova, D.; Timonen, S.; Wennerberg, K.; Aittokallio, T. Improved Detection of Differentially Represented DNA Barcodes for High-throughput Clonal Phenomics. *Mol. Syst. Biol.* **2020**, *16* (3), e9195. https://doi.org/10.15252/msb.20199195.

(40)   Love, M. I.; Huber, W.; Anders, S. Moderated Estimation of Fold Change and Dispersion for RNA-Seq Data with DESeq2. *Genome Biol.* **2014**, *15* (12), 550. https://doi.org/10.1186/s13059-014-0550-8.

(41)   Robinson, M. D.; McCarthy, D. J.; Smyth, G. K. edgeR: A Bioconductor Package for Differential Expression Analysis of Digital Gene Expression Data. *Bioinformatics* **2010**, *26* (1), 139–140. https://doi.org/10.1093/bioinformatics/btp616.

(42)   *barbieQ*. Bioconductor. http://bioconductor.org/packages/barbieQ/ (accessed 2025-09-11).

(43)   Ritchie, M. E.; Phipson, B.; Wu, D.; Hu, Y.; Law, C. W.; Shi, W.; Smyth, G. K. Limma Powers Differential Expression Analyses for RNA-Sequencing and Microarray Studies. *Nucleic Acids Res.* **2015**, *43* (7), e47. https://doi.org/10.1093/nar/gkv007.

(44)   Heinze, G.; Schemper, M. A Solution to the Problem of Separation in Logistic Regression. *Stat. Med.* **2002**, *21* (16), 2409–2419. https://doi.org/10.1002/sim.1047.