Novel biomaterial-based synovial fluid analysis reveals protective microRNA signatures

in a mouse model of acute synovitis-driven osteoarthritis

Authors

Kei Takahata1, Kohei Arakawa2, Takashi Yasuda3, Saaya Enomoto4, Takuya Katashima5,

Sakai Takamasa5, Takanori Kokubun1, 6*

Affiliations

1 Graduate School of Health, Medicine, and Welfare, Saitama Prefectural University,

Saitama, Japan

2 Laboratory of Animal Regeneration Systemology, Department of Life Sciences, School of

Agriculture, Meiji University, Kanagawa, Japan

3 Faculty of Advanced Life Sciences, Hokkaido University, Hokkaido, Japan

4 Saitama Prefecture General Rehabilitation Center, Saitama, Japan

5 Graduate School of Engineering, The University of Tokyo

6 Department of Physical Therapy, School of Health and Social Services, Saitama Prefectural

University, Koshigaya, Saitama, Japan

Correspondence

[1,6]Takanori Kokubun, Ph.D.

820 San-nomiya, Koshigaya, Saitama, 343-8540. Japan

Tel: +81 48-973-417617

kokubun-takanori@spu.ac.jp

Abstract

Objective: Synovial fluid micro-RNAs have been investigated to clarify the Osteoarthritis (OA) development, however, analyzing them in the murine models had not been established. The purpose of this study was to develop a novel biomaterial-based method to collect murine synovial fluid and elucidate onset mechanism of OA with acute synovitis and chronic mechanical stress.

Method: Twelve-week-old C57BL/6J males (n = 72) were divided to the ACL-Transection (ACL-T), ACL-rupture (ACL-R), and Intact groups. We performed joint instability test and histological analysis for cartilage degeneration and synovitis at 2, 6, and 10 weeks. Real time PCR was conducted for articular cartilage at 2 weeks and synovium at 2, 6, and 10 weeks. Tetra-slime was injected into the knee joint and solidified slime including synovial fluid was analyzed in digital PCR at 2 weeks.

Results: No difference was observed in joint instability between both ACL injury models. Unlike the Control and ACL-R groups, acute synovitis and secondary cartilage degeneration were induced in the ACL-T group at 2 weeks. The ACL-T group also exhibited an increase of Mmp-3, Tnf-α, Ifn-γ, and inos in synovium, and miR145-5p and miR149-5p in synovial fluid compared to the ACL-R group. However, no histological and biological differences were observed at 6 and 10 weeks.

Conclusion: We successfully analyzing synovial fluid miRNAs in mice model using

Tetra-Slime. Acute synovitis caused secondary cartilage degeneration through MMP-3, TNF-

α, and M1 macrophage activation in synovium during the early stage, whereas miR145-5p

and miR149-5p in synovial fluid were upregulated to prevent the progression. However,

mechanical stress had a much greater impact on cartilage degeneration in the long run rather

than synovitis.

Key Words:

Introduction

Knee Osteoarthritis (OA) is one of the most common musculoskeletal diseases, characterized by whole−joint pathology, including synovitis and subchondral bone sclerosis, resulting from cartilage degeneration[1]. These structural changes affect each other via synovial fluid and progress irreversibly, as articular cartilage is avascular and has a poor healing capacity. To achieve the prevention of cartilage degeneration, elucidating the intra-articular changes that occur during early OA progression before cartilage degeneration is required; however, it is assumed that early detection is difficult in humans due to the variability of symptoms and disease progression in the early stages of cartilage degeneration. [2, 3]

In the knee joint space, synovial fluid fills and mediates bidirectional interaction between the joint tissues. Synovial fluid is an ultrafiltrate of blood plasma and is primarily composed of hyaluronan and lubricin, which play a significant role as a joint space lubricant and a nutrient source for intra-articular tissues. [4] Previous studies have reported that Interleukin (IL)-6, IL-8, and Tumor Necrosis Factor (TNF)-α in synovial fluid are correlated with OA severity [5]. Additionally, Matrix Metalloproteinase (MMP)-3 could be a biomarker for detecting early-stage OA. [6] Furthermore, circulating microRNAs (miRNAs) in the synovial fluid have been recently highlighted as a novel biomarker. [7] miRNAs are short, non-coding RNAs that are typically 18-25 nucleotides in length, which are used to investigate various diseases as they are stable compared to other RNAs degraded by ribonucleases.[8] It has

been reported that miR-23a-3p, miR-24-3p, miR-27b-3p, miR-29c-3p, miR-34a-5p, and miR-

186-5p in the synovial fluid were expressed in the late-stage OA compared to the early-stage.

[9] In addition, synovial fluid miR-210 increased significantly in the early and late-stage OA

than in healthy individuals. [10] Therefore, synovial fluid has been identified as a potential

target to detect the intra-articular environment in the OA progression. Many clinical studies

have been investigating the synovial fluid biomarker to find OA earlier in human [11], which

has yet to be established. Thus, preclinical experiments using animal models are crucial for

understanding the mechanism of OA onset. However, synovial fluid analysis, especially for

miRNAs in rodent models, is limited because the amount of synovial fluid is tiny. In a few

studies, joint lavage was performed as follows: 25-50 µL of saline was initially injected into

the knee joint, and then the synovial fluid was drawn out.[12-14] However, it is difficult to

collect a sufficient amount of synovial fluid and present relevant data due to the tiny knee

joints in mice.[15]

Recently, many kinds of biomaterials were developed and applied for drug-

deliveries, tissue engineering, and medical diagnostics. [16-18] Tetra-slime is composed of

reversible boronic ester bonds between phenylboronic acid and diol groups, and its hardness

can be changed by adjusting the precursor concentration and pH. Moreover, it can also be

dissolved by sorbitol as sugars attach to phenylboronic acid and cleave ester bonds. [19] Based

on these material properties, we hypothesized that harvesting and dissolving the solidified

Tetra-Slime, which was injected into the knee joint, would allow for the efficient collection of

synovial fluid. The purpose of this study was to develop a novel method to analyze synovial

fluid miRNAs in murine knee joints using biomaterials, Tetra-slime,[20] instead of joint lavage

methods to reveal the onset mechanism of knee OA based on the synovial fluid dynamics.

Using this method in surgical Anterior Cruciate Ligament Transection (ACL-T) and Non-

Invasive ACL rupture (ACL-R) mice models, we aimed to elucidate the characteristics of

each rodent model in terms of inflammation and serve fundamental insights to understand

human pathology in detail.

MATERIALS AND METHODS

Animals and Experimental Design

This study was approved by the Animal Research Committee of Saitama Prefectural

University (approval number: 2020-6). The animals were handled according to relevant

legislation and institutional guidelines for humane animal treatment. Seventy-two 12-week-

old C57BL/6 male mice were randomized into the Invasive ACL-T group (n = 36) and Non-

Invasive ACL-R group (n = 36). The contralateral knee joint in the ACL-R group was used as

the control group. Six mice in each group were sacrificed at 2, 6, and 10 weeks, and the target

tissues were collected for each analysis (Fig. 1). All mice were housed in plastic cages under a

12-hour light/dark cycle. The mice were allowed unrestricted movement within the cage and

had free access to food and water.

Fig. 1. Experimental design. We made the Invasive ACL-T, Non-Invasive ACL-R, and
Control groups. Contralateral knee in the ACL-R model was used as Control group. At 2, 6,
and 10 weeks, anterior drawer test and histological analysis for synovitis and cartilage
denegation in the medial and lateral compartments were performed. Articular cartilage at 2
weeks and synovium at 2, 6, and 10 weeks were harvested, and mRNAs were analyzed in real
time PCR. In addition, synovial fluid was collected at 2 weeks and miRNAs level were
assessed.

Creating ACL-T and ACL-R models

All procedures were performed on the left knee joint of each mouse under a

combination of anesthetics (medetomidine, 0.375 mg/kg; midazolam, 2.0 mg/kg; and

butorphanol, 2.5 mg/kg). The medial joint capsule was cut to expose the intra-articular joint.

ACL was transected with scissors to avoid causing injuries to other tissues. After checking the

anterior tibial translation manually, the joint capsule and skin incision were closed with 5-0

nylon threads. The ACL-R model was created based on our previous study.[21] Briefly, the knee

joint was fixed at 90 degrees using surgical tape on a stand, then the force along the long axis

of the femur was applied by the thumb's tip slowly and carefully. Applied force was stopped

quickly after hearing a distinct popping sound, which indicates ACL rupture; thereafter,

anterior tibial translation was confirmed manually as well.

Ex vivo radiographic analysis for knee joint instability

After dissection at each time point, knee joints were collected (each group, n = 6) and

an anterior drawer test was performed to evaluate joint instability (Fig. 2A). Based on our

previous study [21, 22], the proximal tibia was pulled forward with a constant force spring (0.05

kgf; Sanko Spring Co., Ltd., Fukuoka, JPN), and X-ray images were taken at this position. In

the acquired image, the tibial tuberosity and the anterior end of the femoral condyle were set

as landmarks, then the anterior tibial displacement was measured by the linear distance from

the perpendicular line from the anterior end of the femoral condyle to the tibial tuberosity

using Image J (National Institutes of Health, Bethesda, MD, USA).

Histological analysis for cartilage degeneration and synovitis

Knee joints were fixed using 4% paraformaldehyde for 1 day, decalcified in 10%

ethylenediaminetetraacetic acid for 2 weeks, dehydrated in 70% and 100% ethanol and

xylene, and embedded in paraffin blocks after performing radiographic analysis. The samples

were cut in the sagittal plane (thickness, 7 μm) using a microtome (ROM-360; Yamato Kohki

Industrial Co., Ltd., Saitama, JPN). Then, safranin-O/fast green staining was performed to

assess the articular cartilage degeneration with the Osteoarthritis Research Society

International (OARSI) histopathological grading system [23] and synovitis score [24] in the medial

and lateral compartments. These scores were assessed by two independent observers blinded

to all other sample information. We evaluated the contact area not covered by the meniscus on

the tibial plateau and the synovium located inside the infrapatellar pad. The mean of the

observer's scores was used as a representative value.

Real-time PCR for articular cartilage and synovium

Articular cartilage on the medial and lateral tibial plateau at 2 weeks and synovium at

2, 6, and 10 weeks were collected and stored in a deep freezer until analysis (each group, n =

6). Based on the manufacturer's instructions, total RNA was extracted from these tissues

using the ISOGEN (NIPPON GENE CO., LTD., Tokyo, JPN). cDNA synthesis was

performed from the total RNA by Thermal Cycler Palm-Cycler (Corbett Research, SYD,

AUS) with High-Capacity cDNA Reverse Transcription kit (Applied Biosystems, CA, USA).

Subsequently, real-time PCR was conducted to assess the expression of following genes using

the StepOne-Plus system (Applied Biosystems, Waltham, MA, USA); *Mmp-*

3(Mm00440295), Tissue inhibitor of metalloproteinase-1 (*Timp-1:* Mm01341361),

Transforming growth factor-β1 (*Tgf-β1:* Mm01178820), *Tnf-α* (Mm00443258), *Il-1β*

(Mm00434228), *Il-6* (Mm00446190), *Il-10* (Mm01288386), Interferon-γ (*Ifn-*

γ :Mm01168134), inducible nitric oxide synthase（*inos* :Mm00440502_m1), and Toll like

receptor-4 (*Tlr4* :Mm00445273_m1). We used Glyceraldehyde-3-phosphate dehydrogenase

(*Gapdh* :Mm99999915) as a housekeeping gene, and relative expression of genes was

detected by the 2-△△Ct method [25].

Digital PCR for synovial fluid miRNA using Tetra-Slime

207   In this study, we selected the Tetra-slime to collect the murine synovial fluid. 4-arm

PEG BFA and 4-arm PEG Gluconic (XIAMEN SINOPEG BIOTECH CO., LTD., Fujian,

CHN) were resolved with 50 mM phosphate-buffered saline (pH 3.0) to adjust 0.2 g/ml

concentration respectively. We injected both solutions by 25 μL into the knee joint for live

mice, then kept the needle for a while to avoid leaking the solution (Fig. 7A). After 3 minutes,

mice were sacrificed and the quadriceps and patellar were removed to expose the intra-

articular joints, then solidified Tetra-Slime was collected. We added 100 μL of sorbitol

solution with a 0.5 g/ml concentration to the collected Tetra-Slime and stirred slowly to

dissolve and separate from the surrounding intra-articular tissue, then obtained the pure

solution with synovial fluid biomolecules released. Based on the manufacturer's instructions,

miRNA was extracted from the synovial fluid solution using the miRNeasy Micro Kit

(QIAGEN, LTD. Tokyo, JPN). cDNA synthesis was performed by QIAamplifier 96

(QIAGEN, LTD. Tokyo, JPN) with miRCURY LNA RT Kit (QIAGEN, LTD. Tokyo, JPN).

Subsequently, digital PCR (dPCR) was conducted using the QIAcuity One (QIAGEN, LTD.

Tokyo, JPN) and target genes were *hsa-miR-145-5p* (YP00204483) and *hsa-miR-149-5p*

(YP00204321). We normalized the target gene with the housekeeping gene, 5S rRNA

(YP00203906), for each sample, and then calculated the relative expression level by

standardizing with the Control group. UniSp6 (YP00203954) was also used as a spike-in

control.

Statistical analysis

Statistical analysis was performed using RStudio version 1.4.1106. For the results of

anterior tibial displacement by the anterior drawer test, relative expression of genes in

synovium and cartilage by real-time PCR, and miRNA data measured using dPCR, one-way

analysis was performed initially, then the Tukey-Kramer test was used as a post-hoc test. We

utilized the value of ΔCt for the real-time PCR data and the relative expression value for the

dPCR data. Whereas the Kruskal-Wallis test was used to compare the OARSI and synovitis

scores, and the Steel-Dwass method was used for the subsequent multiple comparisons.

Parametric data are expressed as the means ± 95% confidence intervals (CI); in contrast, the

non-parametric data are expressed as the medians ± interquartile ranges. Statistical

significance was set at $p < 0.05$. CIs were shown in the Supplementary data.

Results

ACL-T and ACL-R groups showed similar joint instability

The anterior displacement in the ACL-T and ACL-R groups was significantly higher

than that in the Control groups at 2 weeks (ACL-T vs. Control, $p < 0.001$, ACL-R vs. Control,

$p < 0.001$), 6 weeks (ACL-T vs. Control, $p < 0.001$, ACL-R vs. Control, $p < 0.001$), and 10

weeks (ACL-T vs. Control, $p < 0.001$, ACL-R vs. Control, $p < 0.001$). However, there was no

significant difference between the ACL-T and ACL-R groups at all time points. These results

suggested that the effect of mechanical stress on OA progression was identical in both ACL-

deficient models.

Fig. 2. Evaluation of the joint instability *ex vivo* using soft X-ray and original anterior drawer

device. At 2, 6, and 10 weeks, the amount of anterior tibial displacement was significantly

increased in the ACL-T and ACL-R groups compared to that in the Control group. However,

there was no significant difference between both ACL injury models. Data are presented as

the mean ± 95% CI.

ACL-T group induced acute synovitis in the early stage

For the medial compartment, severe synovitis with increased cellularity, enlargement

of the synovial lining cell layer, and inflammatory infiltration was observed in the ACL-T

group (Fig. 3A). Moderate synovitis with cell multiplication and cell layer enlargement was

confirmed in the ACL-R group. Compared with the Control group, synovitis score was

significantly higher in the ACL-T group (p = 0.009) and ACL-R group (p = 0.012) (Fig 3B).

Interestingly, ACL-T group increased score significantly compared with the ACL-R group (p

= 0.032). At 6 weeks, moderate synovitis characterized by some enlargement and increased

cellularity developed in both ACL-deficient groups, and the synovitis score in the ACL-T

group (p = 0.01) and ACL-R group (p = 0.01) was significantly higher than in the Control

group. At 10 weeks, similar moderate to severe synovitis was observed in both ACL-deficient

groups, and the synovitis score in the ACL-T group (p = 0.009) and ACL-R group (p = 0.01)

was significantly elevated compared to the Control group.

For the lateral compartment, moderate and mild synovitis occurred in both ACL-

deficient groups. Synovitis score was significantly higher in the ACL-T group (p = 0.009) and

ACL-R group (p = 0.01) compared with the Control group (Fig. 3C-D). At 6 and 10 weeks,

moderate synovitis was observed in the ACL-T and ACL-R groups, and both synovitis scores

increased significantly compared with the Control group (6 weeks: ACL-T vs. Control, p =

0.01; ACL-R vs. Control, p = 0.013, 10 weeks: ACL-T vs. Control, p = 0.016; ACL-R vs.

Control, p = 0.01).

Fig. 3. Histological analysis of synovitis using safranin-O/fast green staining and synovitis score. (A-B) In the medial compartment, the synovitis score in the ACL-T and ACL-R groups was significantly higher than that in the Control group at 2 weeks. Moreover, the score in the ACL-T group was also elevated significantly compared to the ACL-R group. At 6 and 10 weeks, both ACL injury groups significantly increased the synovitis score compared with the Control group. (C-D) In the lateral compartment, the ACL-T and ACL-R groups exhibited a significant increase of synovitis score compared to the Control group at all timepoints. Data are presented as the median ± interquartile range. Black scale bar, 100 μm.

Secondary cartilage degeneration was induced by acute synovitis in the ACL-T group

To examine the tissue interaction between early synovitis and cartilage degeneration, histological analysis was performed. For the medial compartment, some fibrillation of the superficial cartilage was observed in the ACL-T and ACL-R groups at 2 weeks (Fig. 4A). Additionally, small clefts and loss of cartilage were confirmed only in the ACL-T group. The

OARSI score of the ACL-T group was significantly higher than that of the Control group ($p =$

0.009) (Fig. 4B). At 6 weeks, cartilage fibrillation and mild to moderate vertical cracks into

the calcified layer developed in both ACL-deficient groups. Compared with the Control

group, the OARSI score was significantly elevated in the ACL-T group ($p = 0.022$) and ACL-

R group ($p = 0.014$). At 10 weeks, moderate to severe cartilage degeneration with loss of the

superficial layer was observed in both ACL-deficient groups. Compared with the Control

group, the OARSI score significantly increased in the ACL-T group ($p = 0.01$) and ACL-R

group ($p = 0.01$).

298   For the lateral compartment, cartilage degeneration with loss of safranin staining,

small fibrillation, and vertical clefts was confirmed in the only ACL-T group at 2 weeks (Fig.

4C). The OARSI score in the ACL-T group was significantly higher compared with that in

the Control group ($p = 0.018$) and ACL-R group ($p = 0.02$) (Fig. 4D). At 6 weeks, small

fibrillation and clefts in the superficial cartilage layer occurred in both ACL-deficient groups;

however, there was no significant difference in the OARSI score between all groups. At 10

weeks, mild cartilage degeneration with small fibrillation and vertical clefts in the superficial

cartilage layer was observed in both ACL-deficient groups. Compared with the Control

group, the OARSI score was significantly higher in the ACL-T group ($p = 0.012$) and ACL-R

group ($p = 0.044$). In summary of the histological analysis, the ACL-T group induced acute

synovitis and cartilage degeneration in the first 2 weeks compared with the ACL-R group,

although no difference in the joint instability. Cartilage degeneration in both ACL-deficient

models progressed in the late stage, as well as synovitis.

Fig. 4. Histological analysis of cartilage degeneration using safranin-O/fast green staining and
OARSI score. (A-B) In the medial compartment, the OARSI score in the ACL-T group was
significantly higher than that in the Control group at 2 weeks. At 6 and 10 weeks, both ACL
injury groups significantly increased the OARSI score compared with the Control group. (C-
D) In the lateral compartment, the OARSI score in the ACL-T group increased significantly
compared the Control and ACL-R groups at 2 weeks. At 10 weeks, both ACL injury groups
exhibited a significant increase of OARSI score compared to the Control group. Data are
presented as the median ± interquartile range. Black scale bar, 100 μm.

ACL-R group induced catabolic factors and inflammation cytokine expression in the

articular cartilage

To explore the molecular mechanism between acute synovitis and cartilage

degeneration, we performed real-time PCR for the articular cartilage at 2 weeks because

cartilage degeneration occurred in the whole area moderately after 6 weeks. In the medial

tibial cartilage compared with the Control group, ACL-T group significantly increased the

expression of *Mmp-3* ($p < 0.001$), *Timp-1* ($p = 0.009$), and *Tnf-α* ($p = 0.008$) (Fig. 5A-C).

ACL-R group also significantly exhibited an increase of *Mmp-3* ($p < 0.001$), *Timp-1* ($p <$

0.001), and *Tnf-α* ($p < 0.001$) compared with the Control group. Moreover, *Tnf-α* in the ACL-

R group was significantly higher than in the ACL-T group ($p = 0.005$) (Fig. 5C).

In the lateral tibial cartilage, ACL-T group significantly exhibited an increase of

Mmp-3 ($p < 0.001$) and *Timp-1* ($p = 0.036$) compared with the Control group (Fig. 5E-F).

Mmp-3, *Timp-1*, and *Tnf-α* in the ACL-R group were significantly upregulated compared with

the Control group (*Mmp-3*; $p < 0.001$, *Timp-1*; $p < 0.001$, *Tnf-α*; $p = 0.008$) and ACL-T group

(*Mmp-3*; $p = 0.03$, *Timp-1*; $p = 0.036$, *Tnf-α*; $p < 0.001$) (Fig. 5E-G). However, no significant

differences were observed in the *Ifn-γ* between all groups for the medial and lateral tibial

cartilage (Fig. 5D, H). *Tgf-β1*, *Il-1β*, *Il-6*, and *Il-10* were exhibited in the Supplementary

Figure 1.

Medial Tibial Cartilage

Lateral Tibial Cartilage

Fig. 5. Real time PCR analysis for medial and lateral tibial articular cartilage at 2 weeks. (A-D) In the medial cartilage, *Mmp-3*, *Timp-1*, and *Tnf-α* expression in the ACL-T and ACL-R groups were significantly increased compared to that in the Control group. In addition, the ACL-R group exhibited a significant increase of *TNF-α* compared to the ACL-T group. (E-H) In the lateral cartilage, the expression of *Mmp-3*, *Timp-1*, and *Tnf-α* in both ACL injury groups were significantly upregulated compared to that in the Control group. *Mmp-3* and *Timp-1* were also significantly increased in the ACL-R group compared to the ACL-T group. Data are presented as the mean ± 95% CI.

ACL-T group increased the expression of catabolic factors, inflammatory cytokines, and M1 macrophage markers in the synovium

At 2 weeks, *Mmp-3* expression was significantly higher in the ACL-T group (p < 0.001) and ACL-R group (p < 0.001) than the Control group (Fig. 6A). Furthermore, the ACL-T group also exhibited an increase of *Mmp-3* expression significantly compared with the ACL-R group (p = 0.048). *Timp-1* was elevated significantly in the ACL-T (p = 0.002)

and ACL-R group (p < 0.001) compared with the Control group (Fig. 6B). The ACL-T group

significantly increased *Tnf-α*, *Ifn-γ*, and *inos* compared with the Control (*Tnf-α*; p = 0.001, *Ifn-*

γ; p = 0.021, *inos*; p = 0.017) and ACL-R group (*Tnf-α*; p = 0.004, *Ifn-γ*; p = 0.012, *inos*; p =

0.009) (Fig. 6C-E). However, no significant difference was observed in the *Tlr4* expression

among groups (Fig. 6F). The results for *Tgf-β1*, *Il-1β*, *Il-6,* and *Il-10* at 2 weeks, all results at

6 weeks, and 10 weeks were described in the Supplementary Figures 2-4.

363   In summary, the RT-qPCR results showed the increased expression of the *Mmp-3*,

364   *Timp-1* of *Mmp-3* inhibitor [26], and *Tnf-α* in cartilage of the ACL-R group, although it induced

365   less synovitis and cartilage degeneration. Unlike the results of cartilage, synovium in the

366   ACL-T group showed upregulation of *Ifn-γ,* M1 macrophage polarization factor, and *iNos*,

367   M1 macrophage marker[27-30] , in addition to *Mmp-3* and *Tnf-α*. However, the *Mmp-3* and *Tnf-α*

368   expressions were not increased at 6 and 10 weeks.

Synovium

A *MMP-3 / GAPDH*

Relative mRNA level

p < 0.001
p < 0.001   p = 0.048

Control   ACL-T   ACL-R

B *TIMP-1 / GAPDH*

Relative mRNA level

p < 0.001
p = 0.002

Control   ACL-T   ACL-R

C *TNF-α / GAPDH*

Relative mRNA level

p = 0.001   p = 0.004

Control   ACL-T   ACL-R

D *IFNn-γ / GAPDH*

Relative mRNA level

p = 0.021   p = 0.012

Control   ACL-T   ACL-R

E *iNOS / GAPDH*

Relative mRNA level

p = 0.017   p = 0.009

Control   ACL-T   ACL-R

F *TLR4 / GAPDH*

Relative mRNA level

Control   ACL-T   ACL-R

Fig. 6. (A) Real time PCR analysis for synovium at 2 weeks. (A) *Mmp-3* expression in the

ACL-T and ACL-   that in the Control group.

Mo

Tin

(C-

the

bet

Th

T g

G

3 minutes later

❑ Injection of Tetra-Slime   ❑ Collecting Tetra-Slime from joint cavity

Tetra-Slime

4-arm PEG-MA   8-arm PEG-Adtronic

synovial fluid biomelecules

381 exp

382 inc

383 low

H *145-5p / 5S rRNA*

Relative miRNA level

Control   ACL-T   ACL-R

I *149-5p / 5S rRNA*

Relative miRNA level

Control   ACL-T   ACL-R

expression of miR145-5p[31] and miR149-5p[32, 33], which are downregulated by TNF-α, was

investigated. dPCR results showed the increased expression of *miR145-5p* and *miR149-5p* in

the ACL-T group compared with other groups, although this difference was not statistically

significant (Fig

Fig. 7. (A) Conceptual diagram of a novel method to collect synovial fluid biomolecules in
the murine knee joints using Tetra-Slime. After dissolving 4-arm PEG-BFA and 4-arm PEG-
Gluconic in 50 mM phosphate-buffered saline (pH 3.0) and adjusting the concentration to 0.2
g/ml, 50 μl was injected into the knee joint. 3 minutes later, solidified Tetra-Slime including
synovial fluid was collected and it was resolved by 100 μL of sorbitol solution with a 0.5 g/ml
concentration to separate from the surrounding intra-articular tissue. (B-C) The ACL-T group
exhibited an increase of miR145-5p and miR149-5p expression in synovial fluid compared to
the Control and ACL-R groups at 2 weeks. Data are presented as the mean ± 95% CI.

Discussion

Synovial fluid analysis contributes to understanding molecular changes within knee

joint capsule and detecting OA earlier. However, analysis methods for mice had not been

established. In this study, we developed a novel method to collect and analyze synovial fluid

miRNAs in murine knee joints using Tetra-slime, which can adjust its hardness. Using this

method, we aimed to investigate knee OA pathology in each Invasive and Non-Invasive ACL

injury mouse models in terms of tissue-interaction via synovial fluid. Our data clarified that

acute synovitis in the ACL-T model induced secondary cartilage degeneration in the early

stage compared with the ACL-R model; however, the progression of these histological

changes was identical between both models in the late stage. Real-time PCR revealed that

Mmp-3, *Tnf-α*, and M1 macrophage activation in synovium might be involved in the

secondary cartilage degeneration. We successfully established a novel method for analyzing

murine synovial fluid, and dPCR revealed that circulating *miR145-5p* and *miR149-5p* in

synovial fluid may suppress OA progression caused by acute synovitis.

Mechanical stress plays a significant role in maintaining intra-articular homeostasis;

however, excessive stress induces synovitis and cartilage degeneration. During the early stage

of OA progression, synovitis is generally initiated and driven by damage-associated

molecular patterns (DAMPs) released from the extracellular matrix to the synovial fluid.[34-37]

DAMPs activate TLRs in the synovium, leading to cartilage degeneration through the

inflammatory mediators and MMPs.[38, 39] In this study, we demonstrated that the ACL-T

group significantly induced synovitis at 2 weeks compared with the ACL-R group. However,

there was no difference in joint instability between groups. Furthermore, real-time PCR

revealed no differences in *Tlr4* expression, the receptor for DAMPs [39] in the synovium,

among all groups. Considering that synovitis is generally induced by DAMPs through

cartilage degeneration, our result indicates that acute synovitis at 2 weeks in the ACL-T group

is attributed to the surgical intervention, not secondary pathogenesis. Previous studies

reported that cartilage degeneration following ACL injury initiates in the medial compartment

due to joint instability. [40-42] However, the OARSI score in the lateral compartment of the

ACL-T group was significantly higher than that of the ACL-R group at 2 weeks, suggesting

that acute synovitis might cause secondary cartilage degeneration in the early stage.

We revealed the effect of acute synovitis on cartilage degeneration during the OA

progression, then performed molecular biological analysis to examine the tissue interaction

between synovium and cartilage. MMP-3 in the synovium of OA patients was positively

related to the severity [43], and it degrades cartilage ECM and activates other serine proteases

besides MMP-1, MMP-9, and MMP-13 [44]. Previous studies have shown that the depletion of

synovial macrophages suppresses MMP-3 expression both in vivo [45]and in vitro[46]. Our data

showed that *Mmp-3*, *Tnf-α*, *Ifn-γ*, and *inos* were upregulated in the synovium of the ACL-T

group at 2 weeks, which supported a previous study. This suggests that M1 macrophages

likely affected the tissue interaction between acute synovitis and secondary cartilage

degeneration. Unlike the synovial molecular changes, the expression of *Mmp-3* and *Tnf-α* in

the cartilage increased significantly in the ACL-R group compared with the ACL-T group at

the same time point. These results suggest that articular cartilage may initiate intra-articular

degeneration more preferentially than synovium in the OA induced by joint instability and

less inflammation, which supports our previous study. [21] The expression of *Timp-1*, which is

an MMP-3 inhibitor [26], was also elevated significantly; therefore, it is estimated that cartilage

degeneration occurred slightly in the ACL-R model at 2 weeks.

446   Synovial fluid is the most suitable biofluid for exploring the onset mechanism of OA

because it directly reflects the changes in various intra-articular tissues. The importance of

miRNA epigenetic regulation as a cellular communication is becoming increasingly

recognized, and many studies have recently demonstrated its effectiveness in the synovial

fluid of OA patients [47, 48]. However, no study investigated the changes in synovial fluid

miRNA in the mice OA model due to the difficulty of collection and analysis. Here, we

established a novel method to collect synovial fluid effectively from the murine knee joint

using Tetra-Slime. dPCR showed a non-specific reaction of Tetra-Slime to 5s rRNA

expression (Supplementary result), meaning that it enabled the analysis of synovial fluid

components accurately. Previous studies reported that TNF-α downregulated both miR145-5p

[31] and miR149-5p [32] in chondrocytes, leading to cartilage degeneration. However, our results

clarified that circulating *miR145-5p* and *miR149-5p* expression in synovial fluid increased

moderately at 2 weeks in the ACL-T group despite inducing moderate OA pathology. This

suggests that miR145-5p and miR149-5p in synovial fluid might be produced from

chondrocytes to protect against secondary cartilage degeneration by acute synovitis.

461   In contrast to the histological OA pathologies observed at 2 weeks, there were no

462   significant differences in the synovitis and OARSI score at 6 and 10 weeks between the ACL-

463   T and ACL-R groups. This suggests that mechanical stress with joint instability has a much

464   greater impact on cartilage degeneration in the long term, rather than acute synovitis, in the

465   progression of OA.

466   This study has two limitations: first, we could not exclude the intra-patellar fat pad

(IFP) when synovium was collected from the knee joint. Therefore, the result of real-time

PCR for synovium might reflect the biological changes of IFP to synovitis and mechanical

stress. The second is that miRNA-derived cells were not clarified. Synovial fluid components

reflect biological changes of various articular tissues, not only synovium and cartilage;

therefore, we need to perform additional in vitro experiments to investigate the tissue

interaction between synovium and cartilage in detail.

473   In conclusion, we successfully established a novel method for collecting synovial

fluid in the murine knee joint using Tetra-slime. During the OA progression, acute synovitis

led to secondary cartilage degeneration in the early stage, which might be attributed to the

increase of MMP-3, TNF-α, and M1 macrophage activation in the synovium. Among this

tissue interaction, synovial fluid might play a role in preventing cartilage degeneration by

increasing miR145-5p and miR149-5p. Whereas in the middle and late stages, mechanical

stress was a more critical factor in the progression of cartilage degeneration rather than

synovitis (Fig. 8).

Fig. 8. Schematic mechanism of OA development with acute synovitis and chronic
mechanical stress. Acute synovitis caused secondary cartilage degeneration in the early stage
and MMP-3, TNF-α, and M1 macrophage activation in the synovium were involved in this
tissue interaction. Furthermore, miR145-5p and miR149-5p were secreted into the synovial
fluid to protect secondary cartilage deterioration during the same time. Unlike the early stage,
continuous mechanical stress had a much greater impact on cartilage degeneration rather than
synovitis after the middle stage.

Acknowledgments

493   The author(s) received no financial support for the research, authorship, and/or

494 publication of this article. The study design and summary scheme were created with

495 BioRender.com.

Author contributions

All authors approved the final submitted manuscript.

Study design: KT.

Data collection: KT

Histological analysis: KT, KA, and SE.

Biological analysis: KT, SY, TS, and TK.

Manuscript composition: KT, TK.

Role of funding source

506   This study was supported by the Grant-in-Aid for JSPS Research Fellows (22J23384)

507 to KT and Grant-in-Aid for JSPS Challenging Research (Exploratory)(21K19724) to TK.

Competing interest statement

510   All authors have no conflicts of interest related to the manuscript.

References

1. Han S. Osteoarthritis year in review 2022: biology. Osteoarthritis Cartilage 2022;30:1575–82.

2. Karsdal MA, Christiansen C, Ladel C, Henriksen K, Kraus VB, Bay–Jensen AC. Osteoarthritis—a case for personalized health care? Osteoarthritis Cartilage 2014;22:7–16.

3. Kuyinu EL, Narayanan G, Nair LS, Laurencin CT. Animal models of osteoarthritis: classification, update, and measurement of outcomes. J Orthop Surg Res 2016;11:19.

4. Peters H, Rockel JS, Little CB, Kapoor M. Synovial fluid as a complex molecular pool contributing to knee osteoarthritis. Nat Rev Rheumatol 2025;21:447–64.

5. Plsikova Matejova J, Spakova T, Harvanova D, Lacko M, Filip V, Sepitka R, et al. A Preliminary Study of Combined Detection of COMP, TIMP–1, and MMP–3 in Synovial Fluid: Potential Indicators of Osteoarthritis Progression. Cartilage 2021;13:1421S–30S.

6. Rajandran SN, Ma CA, Tan JR, Liu J, Wong SBS, Leung YY. Exploring the Association of Innate Immunity Biomarkers With MRI Features in Both Early and Late Stages Osteoarthritis. Front Med (Lausanne) 2020;7:554669.

7. Tavallaee G, Rockel JS, Lively S, Kapoor M. MicroRNAs in Synovial Pathology Associated With Osteoarthritis. Front Med (Lausanne) 2020;7:376.

8. Cortez MA, Bueso–Ramos C, Ferdin J, Lopez–Berestein G, Sood AK, Calin GA. MicroRNAs in body fluids—the mix of hormones and biomarkers. Nat Rev Clin Oncol 2011;8:467–77.

9. Li YH, Tavallaee G, Tokar T, Nakamura A, Sundararajan K, Weston A, et al. Identification of synovial fluid microRNA signature in knee osteoarthritis: differentiating early– and late–stage knee osteoarthritis. Osteoarthritis Cartilage 2016;24:1577–86.

10. Xie W, Su W, Xia H, Wang Z, Su C, Su B. Synovial Fluid MicroRNA–210 as a Potential Biomarker for Early Prediction of Osteoarthritis. Biomed Res Int 2019;2019:7165406.

11. Boffa A, Merli G, Andriolo L, Lattermann C, Salzmann GM, Filardo G. Synovial Fluid Biomarkers in Knee Osteoarthritis: A Systematic Review and Quantitative Evaluation Using BIPEDs Criteria. Cartilage 2021;13:82S–103S.

12. Sousa–Valente J, Calvo L, Vacca V, Simeoli R, Arevalo JC, Malcangio M. Role of TrkA signalling and mast cells in the initiation of osteoarthritis pain in the monoiodoacetate model. Osteoarthritis Cartilage 2018;26:84–94.

13. Keeble J, Blades M, Pitzalis C, Castro da Rocha FA, Brain SD. The role of substance P in microvascular responses in murine joint inflammation. Br J Pharmacol 2005;144:1059–66.

14. Keeble J, Russell F, Curtis B, Starr A, Pinter E, Brain SD. Involvement of transient receptor potential vanilloid 1 in the vascular and hyperalgesic components of joint inflammation. Arthritis Rheum 2005;52:3248–56.

15.   Seifer DR, Furman BD, Guilak F, Olson SA, Brooks SC, 3rd, Kraus VB. Novel synovial fluid recovery
method allows for quantification of a marker of arthritis in mice. Osteoarthritis Cartilage
2008;16:1532-8.
16.   Hong S, Yu T, Wang Z, Lee CH. Biomaterials for reliable wearable health monitoring: Applications
in skin and eye integration. Biomaterials 2025;314:122862.
17.   Wang H. Biomaterials in Medical Applications. Polymers (Basel) 2023;15.
18.   Trucillo P. Biomaterials for Drug Delivery and Human Applications. Materials (Basel) 2024;17.
19.   Fang G, Bian Z, Liu D, Wu G, Wang H, Wu Z, et al. Water-soluble diboronic acid-based fluorescent
sensors recognizing d-sorbitol. New Journal of Chemistry 2019;43:13802-9.
20.   Katashima T, Kudo R, Naito M, Nagatoishi S, Miyata K, Chung UI, et al. Experimental Comparison of
Bond Lifetime and Viscoelastic Relaxation in Transient Networks with Well-Controlled
Structures. ACS Macro Lett 2022;11:753-9.
21.   Takahata K, Arakawa K, Enomoto S, Usami Y, Nogi K, Saitou R, et al. Joint instability causes
catabolic enzyme production in chondrocytes prior to synovial cells in novel non-invasive ACL
ruptured mouse model. Osteoarthritis Cartilage 2022.
22.   Arakawa K, Takahata K, Enomoto S, Oka Y, Ozone K, Nakagaki S, et al. The difference in joint
instability affects the onset of cartilage degeneration or subchondral bone changes. Osteoarthritis
Cartilage 2022;30:451-60.
23.   Glasson SS, Chambers MG, Van Den Berg WB, Little CB. The OARSI histopathology initiative -
recommendations for histological assessments of osteoarthritis in the mouse. Osteoarthritis
Cartilage 2010;18 Suppl 3:S17-23.
24.   Krenn V, Morawietz L, Burmester GR, Kinne RW, Mueller-Ladner U, Muller B, et al. Synovitis
score: discrimination between chronic low-grade and high-grade synovitis. Histopathology
2006;49:358-64.
25.   Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative
PCR and the $2^{-\Delta\Delta C_T}$ Method. Methods 2001;25:402-8.
26.   Takahashi K, Goomer RS, Harwood F, Kubo T, Hirasawa Y, Amiel D. The effects of hyaluronan on
matrix metalloproteinase-3 (MMP-3), interleukin-1a (IL-1a), and tissue inhibitor of
metalloproteinase-1 (TIMP-1) gene expression during the development of osteoarthritis.
Osteoarthritis Cartilage 1999;2:182-90.
27.   Zhang H, Cai D, Bai X. Macrophages regulate the progression of osteoarthritis. Osteoarthritis
Cartilage 2020;28:555-61.
28.   Wu CL, Harasymowicz NS, Klimak MA, Collins KH, Guilak F. The role of macrophages in
osteoarthritis and cartilage repair. Osteoarthritis Cartilage 2020;28:544-54.
29.   Fernandes TL, Gomoll AH, Lattermann C, Hernandez AJ, Bueno DF, Amano MT. Macrophage: A
Potential Target on Cartilage Regeneration. Front Immunol 2020;11:111.
30.   Zhao K, Ruan J, Nie L, Ye X, Li J. Effects of synovial macrophages in osteoarthritis. Front Immunol
2023;14:1164137.

31. Hu G, Zhao X, Wang C, Geng Y, Zhao J, Xu J, *et al.* MicroRNA-145 attenuates TNF-alpha-driven
cartilage matrix degradation in osteoarthritis via direct suppression of MKK4. Cell Death Dis
2017;8:e3140.

32. Santini P, Politi L, Vedova PD, Scandurra R, Scotto dAbusco A. The inflammatory circuitry of miR-
149 as a pathological mechanism in osteoarthritis. Rheumatol Int 2014;34:711-6.

33. Chiu YS, Bamodu OA, Fong IH, Lee WH, Lin CC, Lu CH, *et al.* The JAK inhibitor Tofacitinib inhibits
structural damage in osteoarthritis by modulating JAK1/TNF-alpha/IL-6 signaling through Mir-
149-5p. Bone 2021;151:116024.

34. de Lange-Brokaar BJ, Ioan-Facsinay A, van Osch GJ, Zuurmond AM, Schoones J, Toes RE, *et al.*
Synovial inflammation, immune cells and their cytokines in osteoarthritis: a review.
Osteoarthritis Cartilage 2012;20:1484-99.

35. Nefla M, Holzinger D, Berenbaum F, Jacques C. The danger from within: alarmins in arthritis. Nat
Rev Rheumatol 2016;12:669-83.

36. Lopes EBP, Filiberti A, Husain SA, Humphrey MB. Immune Contributions to Osteoarthritis. Curr
Osteoporos Rep 2017;15:593-600.

37. Lambert C, Zappia J, Sanchez C, Florin A, Dubuc JE, Henrotin Y. The Damage-Associated Molecular
Patterns (DAMPs) as Potential Targets to Treat Osteoarthritis: Perspectives From a Review of the
Literature. Front Med (Lausanne) 2020;7:607186.

38. Liu-Bryan R. Synovium and the innate inflammatory network in osteoarthritis progression. Curr
Rheumatol Rep 2013;15:323.

39. Bartels YL, van Lent P, van der Kraan PM, Blom AB, Bonger KM, van den Bosch MHJ. Inhibition of
TLR4 signalling to dampen joint inflammation in osteoarthritis. Rheumatology (Oxford)
2024;63:608-18.

40. Pauly HM, Larson BE, Coatney GA, Button KD, DeCamp CE, Fajardo RS, *et al.* Assessment of cortical
and trabecular bone changes in two models of post-traumatic osteoarthritis. J Orthop Res
2015;33:1835-45.

41. Maerz T, Newton MD, Kurdziel MD, Altman P, Anderson K, Matthew HW, *et al.* Articular cartilage
degeneration following anterior cruciate ligament injury: a comparison of surgical transection and
noninvasive rupture as preclinical models of post-traumatic osteoarthritis. Osteoarthritis
Cartilage 2016;24:1918-27.

42. Gilbert SJ, Bonnet CS, Stadnik P, Duance VC, Mason DJ, Blain EJ. Inflammatory and degenerative
phases resulting from anterior cruciate rupture in a non-invasive murine model of post-traumatic
osteoarthritis. J Orthop Res 2018;36:2118-27.

43. Chen JJ, Huang JF, Du WX, Tong PJ. Expression and significance of MMP3 in synovium of knee joint
at different stage in osteoarthritis patients. Asian Pac J Trop Med 2014;7:297-300.

44. Murphy G, Nagase H. Progress in matrix metalloproteinase research. Mol Aspects Med
2008;29:290-308.

45.   Blom AB, van Lent PL, Libregts S, Holthuysen AE, van der Kraan PM, van Rooijen N, *et al.* Crucial
622   role of macrophages in matrix metalloproteinase-mediated cartilage destruction during
623   experimental osteoarthritis: involvement of matrix metalloproteinase 3. Arthritis Rheum
624   2007;56:147–57.

625   46.   Bondeson J, Wainwright SD, Lauder S, Amos N, Hughes CE. The role of synovial macrophages and
626   macrophage-produced cytokines in driving aggrecanases, matrix metalloproteinases, and other
627   destructive and inflammatory responses in osteoarthritis. Arthritis Res Ther 2006;8:R187.

628   47.   Munjal A, Bapat S, Hubbard D, Hunter M, Kolhe R, Fulzele S. Advances in Molecular biomarker for
629   early diagnosis of Osteoarthritis. Biomol Concepts 2019;10:111–9.

630   48.   Palma AD, Cheleschi S, Pascarelli NA, Tenti S, Galeazzim M, Fioravanti A. Do microRNAs have a key
631   epigenetic role in osteoarthritis and in mechanotransduction? Clin Exp Rheumatol 2016;35 518–
632   26.