Communication networks of wild zebra finches (*Taeniopygia castanotis*)

Hagedoorn, K.[1]*, Tschirren, N.[1]*, ter Avest, E. [1], Tyson, C. [1], Snijders, L. [1], Griffith, S. C. [2, 3], Loning, H. [1#],

Naguib, M. [1#+]

[1] Behavioural Ecology Group, Wageningen University & Research, Wageningen, The Netherlands

[2] School of Natural Sciences, Macquarie University, Sydney, New South Wales, Australia

[3] School of Biological, Earth and Environmental Sciences, University of New South Wales, Sydney,

New South Wales, Australia

* shared first authors

[#] shared last authors

[+] Corresponding author

[+] Marc Naguib, marc.naguib@wur.nl

Behavioural Ecology Group, Wageningen University, De Elst 1, 6708 WD Wageningen, The

Netherlands

Abstract

Communication networks are widespread across species, permitting information flow and facilitating social connections across space and time. In birds, communication networks are well studied in territorial species with long-range songs connecting individuals across space, where unintended listeners extract information from others' signalling interactions. Yet acoustic signals also play important social roles at short range, forming communication networks that connect individuals within larger social units. Wild zebra finches (*Taeniopygia castanotis*) provide a unique model system to examine such communication networks in a non-territorial species with multiple males singing at shared locations in close temporal proximity. Zebra finches breed in loose colonies in the Australian arid zone, typically stay near their partner, move around in pairs or small groups, and gather at social hotspots, thus forming dynamic, potentially multi-level, societies. Here, we used the individually distinctive male song, recorded over 1,835 song bouts at two breeding sub-colonies and three social hotspots to quantify singing activity. We identified 163 males based on spectrographic similarities, verified by a deep learning model (BirdNET), and constructed communication networks based on temporal singing proximity at shared locations, social hotspots and the breeding colony. We reveal higher singing activity at social hotspots, with no dawn song and singing peaking at different times of day between breeding sites and social hotspots. Communication networks, with distinct males singing in close temporal proximity, were apparent in both contexts, with larger networks at hotspots. These networks were not strongly nested, yet many individuals met repeatedly at either the hotspot or breeding sites and some even maintained associations across breeding colonies and hotspots, providing building blocks for a multi-level society. These networks may facilitate synchronized foraging and breeding as adaptations to a harsh and unpredictable environment. Additionally, our approach offers a novel road map for widening the understanding of communication networks.

Keywords: Communication networks, birdsong, multi-level society, social network, zebra finch, BirdNET, fission-fusion societies, vocal communication, environmental unpredictability

Introduction

Animals vary in communication and sociality, ranging from simple to complex signalling interactions and from solitary living to highly complex social groupings and networks (Ward & Webster, 2016). Sociality and communication are tightly linked, yet while social networks typically refer to spatial proximity of individuals (Croft et al., 2008), communication networks have been mainly considered for long-range signalling interactions where individuals do not associate closely (McGregor, 2005). In many cases, these signals peak at specific times of day, such as in the dawn song of territorial songbirds (Staicer et al., 1996; Kunc et al., 2005), and the singing and singing interactions can influence which individuals perceive the signal and how they respond (McGregor & Dabelsteen, 1996; Naguib et al., 1999; Todt & Naguib, 2000; McGregor, 2005; Foote et al., 2010; Logue, 2021). Such long-range signals may enable information transfer across large distances and can thus be crucial to determine movements, social decisions, such as mate choice, breeding decisions, responses to predators and other aspects of social synchronisation (Catchpole & Slater, 2008; Snijders & Naguib, 2017).

Studies on these long-distance networks have shown that information is not only gathered from a singer and its song but also by eavesdropping on the dynamics of vocal interactions among others, information that is not available when attending to singers separately, leading to the initial definition of communication networks (McGregor & Dabelsteen, 1996). Such relative differences can be important for decision making, such as reproductive decisions (Otter et al., 1999; Schmidt et al., 2006) and priorities in territory defence (Peake et al., 2001; Mennill & Ratcliffe, 2004; Naguib et al., 2004; Peake et al., 2005; Fitzsimmons et al., 2008; Amy et al., 2010; Foote et al., 2010). Eavesdropping on vocal interactions among others has been at the core of communication network approaches to birdsong (Peake, 2005; Naguib et al., 2011). Yet, biologically, a communication network, defined as

patterns of potential information flow, would be much larger, also including information extracted

from individual songs and from direct interactions. Such wider approaches can be useful to better

understand the link between communication networks and spatial networks, as well as to obtain a

better understanding of information flow across large and complex social organizations in animal

societies (Snijders & Naguib, 2017)

Because information from signals affects social decision making, including the initiation, maintenance

and dissolution of social bonds, they can be expected to play a key role in the social organisation of

non-territorial animal societies. For instance, in dynamic and more complex animal organisations such

as multi-level fission-fusion societies, in which not only individuals but also social units at higher levels

of organisation fission and fuse (Papageorgiou et al., 2019; Papageorgiou & Farine, 2021), signals

carrying individually specific information might be key facilitators allowing individuals to maintain one-

to-one or pair-to-pair connections within larger social units. However, the quantitative use of

individually-distinct signals by animals in dynamic, potentially multi-level, societies has received little

attention.

The zebra finch (*Taeniopygia castanotis*) in its natural habitat, the Australian arid zone, offers a highly

suitable study system to address these knowledge gaps. Zebra finches provide a unique model

organism for many lab-based studies (Griffith & Buchanan, 2010), and their dynamic social organisation

offers scope to broaden the traditional concept of communication networks in order to better

understand the role of signals in social organisation. Zebra finches live throughout the Australian arid

zone and move around together with their mate (Tyson et al., 2024) and are encountered mostly as a

pair and not in a group (McCowan et al., 2015). These pairs gather at social hotspots also during non-

breeding times (Loning et al., 2023a) and additionally during breeding will associate with others in their

loose breeding colonies (Brandl et al., 2021). When resources are highly clumped, birds can form

temporarily large groups, such as near water (Zann, 1996; McCowan et al., 2015). These fission-fusion

dynamics suggest that they might live in multi-level societies (e.g. pairs within loose groups, loose

groups within larger aggregations). Since male zebra finches each sing a single unique song motif with

minor variation (Sossinka & Böhner, 1980; Sturdy et al., 1999) and both sexes have individually distinct

calls (Elie & Theunissen, 2018), their dynamic social organisation is most likely facilitated by these

distinctive vocal signals, allowing receivers to potentially identify others in different social settings and

in different locations. Zebra finch vocalisations are very short-distance signals, with an average range

of nine meters for their song, and the loudest calls in their repertoire have an average range of only 14

meters (Loning et al., 2021). These individually distinct close-range vocalisations therefore likely

function to establish and maintain social connections through information exchange at gathering

points, such as at the breeding area (Brandl et al., 2019c), at foraging sites (Brandl et al., 2019a; Brandl

et al., 2021) or at the social hotspots (Loning et al., 2023a).

We took advantage of the wild zebra finch social system, in which birds vocalise and encounter

conspecifics at different locations, to integrate communication networks and spatial networks (Snijders

& Naguib, 2017; Loning et al., 2024) and so examine the network structure of a highly dynamic wild

bird society. We quantified male singing activity throughout the day in a wild population of zebra

finches in the Australian Outback, recording singing both at breeding sites and social hotspots (Loning

et al., 2023a) using automated audio recorders. Using the individually distinctive male song (Sossinka

& Böhner, 1980; Sturdy et al., 1999), we used singing as a proxy for presence at nests and hotspots

and determined if different males would repeatedly sing at the same time at these locations. From

these songs, we created communication networks and expected larger networks at the hotspots, with

more individuals heard singing and associating there than in the breeding colonies. We also expected

a hierarchical social structure with individuals at the hotspots interacting with conspecifics they

associated with in the breeding colony, as well as others. Although only males sing, leading here to

male-focussed networks, pairs stay almost constantly together (Tyson et al., 2024), suggesting that

male song networks reflect a network among pairs.

Methods

The study was conducted from September to October 2024 in the Gap Hills area of the Fowlers Gap

Arid Zone Research Station, New South Wales, Australia. This area consists of open shrubland with low

bushes such as bluebush (*Maireana* sp.) and saltbush (*Rhagodia* sp.) as well as higher vegetation along

natural creeks, mostly shrub-like trees such as prickly wattle (*Acacia victoriae*) and dead finish (*Acacia*

tetragonophylla). There is an artificial water reservoir in the area (Fig. 1), and zebra finches at this site

use nestboxes to breed, which are placed on poles near trees for shelter (Griffith et al., 2008).

Nestboxes are placed in the area in six sub-colonies of about 30 boxes each.

Audio data collection

We recorded vocal activity at nestboxes and nearby social hotspots using time-programmable audio

recorders (AudioMoth; Hill et al., 2019). AudioMoths were set to record from 90 minutes before

sunrise to 90 minutes after sunset (settings: 32 kHz sample rate; medium gain; energy saver mode on).

We monitored breeding activity by checking all nestboxes in the area every three days and attached

AudioMoths to nestbox poles of active nests. We identified social hotspot locations by walking

repeated transects, registering recurring observations of zebra finches and checking for the presence

of faecal droppings. We combined this information with hotspot locations obtained in the same way

in previous years, as they are generally temporally stable (Loning et al., 2023b). We then observed

these potential hotspots three times for one hour between 0800 and 1230 hours on different days to

confirm that they were actively used by zebra finches. We then placed AudioMoths at three identified

social hotspots (all three were also in use in the previous year). Two social hotspots were located at

the edges of a breeding sub-colony (colony D), and one was located between another sub-colony

(colony E) and the central water reservoir (Fig. 1).

Figure 1. Map showing the field site at Gap Hills with positions of analysed nestboxes and social

hotspots. Tree icons denote social hotspots, house icons denote nestboxes selected for audio analysis,

with six located in breeding colony D and three in breeding colony E. The central dam, creek lines and

highway are visible. Satellite image retrieved from Google Maps.

Audio analysis

For each of nine nestboxes (Fig. 1), we selected one day of audio recordings for analysis, assuming that

one day is sufficient to record the locally breeding birds. We selected the day with the most favourable

weather condition (i.e. no rain, little wind) during the egg-laying phase as males sing most during this

phase of breeding (Loning et al., 2023b). The selected nests all produced fledglings but were active at

different times during the study period. For the three hotspots, we selected audio from seven different

days, two days for hotspots 1 and 2 and three days for hotspot 3 (Fig. 1). Where possible, we analysed

hotspots on the same day as a nearby nestbox (at most five days difference), to minimise

environmental variation such as wind affecting movement and sociality. In total, we analysed 248

hours of audio recordings, consisting of 140 hours of recordings at nine nestboxes (one day per nestbox) and 108 hours at three social hotspots (two to three days per hotspot).

Within these audio recordings, we first assigned zebra finch songs to individual males by listening to the song recordings and visually inspecting their sound spectrograms (window type: Hamming, window size: 1024) in Audacity® (version 3.7.1; Audacity Team, 2024), and later verified this using deep learning logarithms (see below). Zebra finch song has a distinct structure, with each male repeatedly singing an individually distinct motif (series of syllables), which is readily recognizable on spectrograms (Fig. 2) although some intraindividual variation exists (Sossinka & Böhner, 1980; Sturdy et al., 1999). A gap of more than five seconds between song motifs separated song bouts. We assigned a recording quality score to each song bout on a scale of one to 10. Only songs of high quality, typically with a score of six or higher, were assigned to an individual bird, while songs of lower quality were not included here. We established a database containing clear spectrograms of individual birds and compared newly encountered songs to this database. If song element frequencies, shapes and sequence within motifs, as well as the motif organisation of a new song, matched that of an already encountered individual, we assigned it to the same individual. Ambiguous cases were checked by a second observer. If there was no consensus on the classification, that song was not assigned to a specific individual. Out of the total 2,552 song bouts present in our recordings (248 hours), we assigned 1,835 song bouts (71% of song bouts) to 163 individuals (mean ± SD: 11.3 ± 15.9 song bouts per individual, range 1-113).

Based on the occurrence of individual songs, we determined the number of different locations visited by each individual. We quantified song activity over the day by summing the number of seconds containing song within each half-hour block, allowing us to compare singing activity at nestboxes and social hotspots.

Figure 2. Example spectrograms of song bouts from three different individuals. The y-axis shows the

frequency in kHz and the x-axis time in seconds. Darker shades correspond to a higher sound energy.

Different elements of the motifs are annotated. IN denotes introductory notes, E1, E2 and E3 denote

different song element types. The individuals (one per row) differ in element sequence, frequencies,

and shapes. Intraindividual differences such as variation in motif duration and content are visible, e.g.

individual 2 does not have element E2 in the second motif. Spectrograms were made using Audacity.

BirdNET validation of audio analysis

To validate whether or not the songs we had manually assigned to an individual based on visual inspections of sound spectrograms were indeed more similar to each other than to songs that we assigned to other individuals while accounting for potential bias and unavoidable variation in recording conditions, we used BirdNET (model version 2.4; (Kahl et al., 2021) as an impartial observer. BirdNET is a deep-learning model trained to recognise the vocalisations of over 6,000 bird species to the species level (including zebra finches), but it has not been trained to distinguish between zebra finch individuals. BirdNET was used as a pre-trained model without additional training or fine-tuning on our dataset, as this would have required using our individual labels, which we intended to validate independently. BirdNET's deep neural network processes each sound fragment and extracts a set of 1,024 numerical features. These so-called embeddings are a compact representation that captures the patterns relevant for distinguishing bird species (McGinn et al., 2023). While originally designed to support species classification, the embeddings can also be treated as a 1,024-dimensional feature vector that characterises the acoustic properties of individual bird sound recordings.

We extracted BirdNET embeddings representing zebra finch songs for subsequent intra- and inter-individual comparison. Since BirdNET internally uses three-second segments for classification, we trimmed song bouts to the first three seconds and also kept the last 3 seconds for bouts over 6 seconds long. Song bouts shorter than three seconds were left unaltered (BirdNET pads segments shorter than three seconds). This resulted in 2,617 song segments (belonging to 163 individuals) for which we extracted the corresponding embeddings using BirdNET-Analyzer (GUI version 2.1.1; settings: overlap 0 seconds, batch size 1, no speed modification, bandpass frequency 0-15 kHz). We then calculated cosine distances (i.e. 1 - cosine similarity, which measures the angle between vectors irrespective of their magnitude) between the embeddings for all combinations of segments using SciPy's *spatial.distance.pdist* (with metric="cosine") function in Python (version 3.13.5). This resulted in a combined 3.4 million intra- and interindividual comparisons. We further divided comparisons into

three categories based on where the two compared segments originated: comparisons made between

segments from (1) the same one-hour audio recording, (2) the same date, and (3) different dates.

These categories reflect a potential increase in recording condition differences and a decrease in

certainty of songs actually having been sung by the same individual.

Video recordings at social hotspots and analysis

To quantify zebra finch visit durations for constructing communication networks, we filmed the three

social hotspot trees using tripod-mounted action cameras (GoPro Hero5 Black, resolution of 1280x720,

25 FPS, narrow field of view) for a total of six days (one to three days per social hotspot) from a height

of about 150 cm at a distance that allowed the whole tree to be in frame in order to approximate visit

durations by scoring arrivals and departures. To estimate a conservative (i.e. minimum) average time

zebra finches spend at a social hotspot, we analysed 64 hours of video footage of the three social

hotspots. Each departure from and arrival at the hotspot was scored using BORIS (Behavioural

Observation Research Interactive Software; version 9.2.1). For each arrival, we matched the next

available departure to calculate the minimum duration each bird spent on the tree by keeping a queue

of arrival times, matching each arrival to the earliest unmatched departure. To compute visit durations,

we excluded visits lacking matching pairs attributable to missed arrivals or departures on video and

birds staying for extended periods. To limit the influence of extreme outliers and focus on typical visit

durations, bird visits lasting longer than one hour were excluded from the main analysis. This cutoff

was based on preliminary data exploration, which showed that durations exceeding one hour were

rare and probably represented atypical or faulty events. Applying this filter improved the robustness

of summary statistics while retaining the majority of observations. In total, we recorded 686 bird visits

with a mean duration of 1,263 seconds (± 21 minutes; median 1,364s ± 722s; IQR 634 – 1,750s).

Communication network characteristics

To determine the connections between individuals, we considered individuals that sang within 1,263 seconds of each other (the mean visit duration at social hotspots, see above) as being present at the same time (associating) and as being socially interacting. Using these associations, we determined social network edges, representing connections between two individuals. We constructed networks (1) based on all associations and (2) based only on repeated associations. For individuals that were heard both at a nestbox and a social hotspot (N = 24), we calculated the number of associations and the number of unique associations (i.e. the number of unique individuals associated with) they had at nestboxes and social hotspots. If the bird was heard at multiple nestboxes or multiple social hotspots, we used the total number of associations or the number of unique associations and calculated how long each individual sang (summed duration of all the song bouts) at each type of location.

Statistical analysis

All statistical analyses were performed using R (version 4.5.1; R Core Team, 2025). Linear mixed-effect models (LMMs) and generalised linear mixed-effect models (GLMMs) were constructed using the *lmer* and *glmer* functions from the 'lme4' package, respectively (Bates et al., 2015). Model assumptions were checked using the *simulateResiduals* and *testDispersion* functions (for Poisson GLMMs) from the 'DHARMa' package (Hartig, 2024). We tested for statistical significance of variables with two categories using estimated marginal means (EMM) with the *emmeans* function from the 'emmeans' package (Lenth, 2025). We also used this function to obtain the reported mean ± SE and contrasts for all mixed model data. To test interaction effects and variables with more than two categories, we compared an interaction model to a model without the interaction term with a likelihood-ratio test, using the function *anova* from the 'stats' package.

To test for differences in singing at the nestboxes and social hotspots, we compared the total duration of song and the number of individuals heard per day between social hotspots and nestboxes using a

LMM and a GLMM with a Poisson distribution, respectively. For both models, we accounted for the

different number of observation days for the different locations (N = 1 for all nestboxes, N = 2 for

hotspot 1 and 2, N = 3 for hotspot 3) by adding the date and the location as non-nested random

intercepts.

For each comparison between BirdNET embeddings, we calculated the cosine distance. We then used

a LMM to test the effect of comparison type (intra- or interindividual) as well as song origin (from the

same one-hour audio recording, the same date, or different dates) and their interaction on cosine

distance. We included the assigned individuals of both compared songs (the individual for song 1 and

for song 2 was the same in the case of an intraindividual comparison) as crossed random intercepts.

To quantify the communication/social networks, we obtained network metrics and graphs using the

igraph package (Csárdi & Nepusz, 2006). For the nestboxes and social hotspots, we calculated and

compared the number of nodes (order) and edges (size) of the repeated association network (defined

as dyads with two or more interactions between them) per day using Poisson GLMMs with the same

random effect structure (see above). For these order and size models, we set the maximum number

of iterations to 100,000 and changed the optimizer to 'bobyqa' to solve convergence issues. We then

compared the total number of associations per individual and the number of unique associations

between breeding sites and social hotspots using Poisson GLMMs with individual ID as a random

intercept. For these models, we included song duration in minutes as a covariate.

Ethical note

We collected all audio and video data using non-invasive methods. Audio and video equipment was

placed while minimising disturbance. Approval was granted by the Macquarie University Animal

Ethics Committee, and the study was conducted under a New South Wales DPIE Scientific Licence

SL100378, Ethics ARA 2018/02.

Results

Singing activity at breeding sites and social hotspots

In 246 hours of recordings, comprising nine nestboxes (breeding sites) at two breeding sub-colonies

and three hotspots, a total of 163 males were identified individually by their song. Most individuals

were heard at only one location (N = 126), with fewer individuals heard at more than one location (two

locations: N = 23, three locations: N = 9, four locations: N = 4, five locations: N = 1). In total, 24

individuals (14.7%) were heard at both a social hotspot and the breeding sites. Zebra finch males did

not sing at dawn, i.e. the hour before sunrise, at either location. Birds started singing around the same

time of the day at both locations, breeding sites and social hotspots (Fig. 3a). Yet, at the breeding sites,

most song occurred earlier in the day between sunrise and noon. At the social hotspots, most song

occurred during the second half of the day. Furthermore, song continued at the social hotspots for

longer, with a higher prevalence of song after sunset than at the breeding sites. The average duration

of song in minutes per day at social hotspots (mean ± SE: 23.9 ± 4.4, N = 7) was significantly higher than

at breeding sites (6.8 ± 3.5, N = 9; EMM: t(6.4) = 3.2, P = 0.02; Fig. 3b). Furthermore, significantly more

individuals were heard during the day at social hotspots (21.2 ± 4.9, N = 7) than at the breeding sites

(8.7 ± 1.6, N = 9; EMM: z = 3.3, P = 0.001; Fig. 3c).

Figure 3. (a) The average distribution of song over the day for breeding sites (nestboxes) and social hotspots. The time is scaled to an average day of 12.5 hours. Sunrise, solar noon, and sunset are denoted by the dashed vertical lines. The y-axis shows the proportion of total daily song that occurred per half hour. The average duration of song in minutes per day (b) and the number of individuals heard per day (c) was higher at social hotspots than at breeding sites. * denotes $P < 0.05$, *** denotes $P = 0.001$.

BirdNet validation of song classification

Songs that we assigned by visual inspection of sound spectrograms as produced by the same individual were more similar using BirdNet analyses, as cosine distance for intra-individual song comparisons (0.304 ± 0.004) was significantly lower than for inter-individual comparisons (0.362 ± 0.004; EMM: z = 156.3, P < 0.001; Fig. 4). There was also a significant interaction between song comparison type (same individual and different individual) and sound origin (same bout, same day, different day) (likelihood-ratio test: $\chi^2(2)$ = 578.5, P < 0.0001). The difference between intra-individual and inter-individual song comparisons was largest for compared songs from the same one-hour recording (0.070 ± 0.001), smaller for the same date (0.053 ± 0.001), and smallest for different dates (0.050 ± 0.001; all P < 0.0001; Fig. 4). However, the BirdNET embeddings probably did not exclusively reflect zebra finch song, but also reflected recording conditions, as sound origin differences also affected the cosine distance (Fig. 4). Overall, songs originating from the same one-hour recording had lower cosine distances (0.305 ± 0.004) than those recorded at the same date (0.340 ± 0.004), and these in turn were lower than those comparisons with songs recorded on different dates (0.354 ± 0.004; likelihood-ratio test: $\chi^2(2)$ = 19,346, P < 0.0001). These differences likely cannot be attributed exclusively to potential errors in individual assignment as not only the intra-individual, but also the inter-individual comparisons originating from the same one-hour recording had lower cosine distances (intra: 0.270 ± 0.004 ; inter: 0.340 ± 0.004) than those made between recordings of the same date (intra: 0.313 ± 0.004 ; inter: 0.366 ± 0.004) and those of different dates (intra: 0.329 ± 0.004 ; inter: 0.380 ± 0.004; EMM: all pairwise z > 16.7 and P < 0.0001; Fig. 4).

Figure 4. Comparisons of extracted BirdNET embeddings as validation of individual identification. We calculated cosine distances between embeddings of 2,617 songs, resulting in 3.4 million comparisons. The cosine distance or dissimilarity ranges from zero (vectors have the same direction) to two (vectors have the opposite direction). Across song origins, cosine distance was lower for intraindividual song comparisons (i.e. between songs of the same individual as assigned based on spectrographic similarities) compared to interindividual comparisons. This difference was biggest when recording conditions were similar (i.e. with both songs originating from the same one-hour audio recording), whereas it was smallest when conditions were potentially most different (songs recorded on different days). Each song comparison type-origin combination differed from the other five (EMM: all pairwise $z > 16.7$ and $P < 0.0001$).

Communication network characteristics

Zebra finches showed non-uniform network structures at both breeding sites and social hotspots. 151 out of 163 identified individuals had an association. However, most of the 473 observed dyads consisted of only one association (86% of dyads), while only a small fraction of dyads associated three or more times (3.2%; Fig. 5a). When we filtered the network to represent repeated associations (defined as dyads with two or more interactions between them), 59 individuals and 66 dyads remained (Fig. 5b). Of these 66 dyads, five dyads showed associations at both location types, indicating that most individuals, within the analysed time frame, have social ties either in a breeding colony or at social hotspots, but not both.

Figure 5. The social network for (a) all associations and (b) all repeated associations, i.e. dyads that were identified together at least two times. The pie shapes represent individuals (nodes), with coloured slices representing the relative amount of time they spent singing in different locations. The size of the node scales by the total amount of time that node was heard singing. The lines represent the song-based associations (edges). The colours of the edges correspond to the location where that singing took place.

When taking all associations into account, the network on a given day was larger at social hotspots

than at the breeding sites. At social hotspots, networks had more nodes (58.6 ± 24.8; EMM: z = 3.3, P

< 0.001) and more edges (48.1 ± 20.0; EMM: z = 3.3, P = 0.001) than at the breeding sites (13.1 ± 4.0

and 11.4 ± 3.5, respectively). Results were qualitatively similar for the repeated association network

(containing only those dyads with two or more associations) on a given day. Also, at social hotspots,

networks had more nodes (9.2 ± 3.5; EMM: z = 2.7, P = 0.01) and more edges (9.5 ± 4.2; EMM: z = 2.1,

P = 0.04) than at breeding sites (3.6 ± 1.3 and 3.9 ± 1.5, respectively).

Network at the individual level

In total, 24 individuals were recorded singing at both a social hotspot and a breeding site. Birds that

sang longer had more associations, and this relation was stronger at social hotspots than at breeding

sites (likelihood-ratio test: $\chi^2(1)$ = 4.0, P = 0.05; Fig. 6a). Furthermore, these individuals had more

associations at social hotspots than at breeding sites (social hotspot = 8.6 ± 1.2, breeding site = 3.9 ±

0.6; EMM: *z = 6.6, P < 0.0001*; Fig. 6b). Individuals also had more unique associations (i.e. associated

with more individuals) at social hotspots than at breeding sites (social hotspot = 7.0 ± 1.0, breeding

site = 3.4 ± 0.5; EMM: *z = 5.9, P < 0.0001*; Fig. 6c).

Figure 6. Total number of associations and number of unique associations at the individual level. Only

individuals that were heard at both a breeding site and a social hotspot are shown. (a) Birds that sang

more had more associations, and this relation was stronger at social hotspots than at breeding sites.

(b) Individuals had more associations at the social hotspot than at the breeding site. (c) Individuals met

with more unique conspecifics at the social hotspot than at the breeding site. Circle size corresponds

to the number of individuals. *** denotes P = 0.001.

Discussion

Here we reveal distinct locations at which wild zebra finches sing in close spatial and temporal

proximity, namely their breeding colony and at social hotspots. Song at these locations was never

produced at dawn, in contrast to many territorial species, but peaked at different times of day, at the

breeding colony in the morning and at the social hotspots in the afternoon. At both locations, multiple

individually identified birds were singing in close temporal proximity in a communication network. The

networks at the breeding colony and social hotspots were distinct, with networks at the social hotspots

being larger than those in the breeding colony. Most individuals were heard at one location only, and

most dyads of males were heard together once, yet we also identified several males that sang together

at different times. Most dyads that associated in the breeding colony were not heard together at a

social hotspot and vice versa, yet a few dyads (five) were identified at multiple locations. These findings

suggest that within this relatively short time frame studied, breeding sites and social hotspots served

as distinct sites for social interactions, with a few individuals and dyads forming a bridge between

them. These findings also suggest that the males of neighbouring breeding pairs do not move together

as a unit to these hotspots, which can be hundreds of meters away from their breeding site, but that

the decision to visit social hotspots is likely more a decision taken by an individual or a pair.

The higher singing activity at the social hotspots was predicted, as social hotspots are defined as

gathering places (Loning et al., 2023a) and zebra finch males sing all year round at various locations

(Loning et al., 2023b). Singing in different social contexts is interesting as it might be triggered

mechanistically through different internal and external processes (Waas et al., 2000; Wang et al.,

2012), while also having different functions and implications. Singing at the breeding colony may

primarily reinforce pair bonds and indicate local breeding activity, thus providing reproductive cues to

neighbours (Loning et al., 2023b), which are known to coordinate their breeding with one another

(Brandl et al., 2019c). Singing at social hotspots, or in similar contexts in other species where individuals

substantially vocalize in groups forming similar types of communication networks (Cure et al., 2009;

Major & Jones 2011; Calcari et al., 2021; Devogel et al., 2024) might have a wider social trigger and

provides information to larger groups, potentially transferring information about resources, predation

risk and reproductive states and allowing individuals to create social bonds, and perhaps coordinate

breeding at a broader local scale.

The absence of dawn song at either location highlights an important distinction to the daily singing

activity of well-studied territorial songbirds in the Northern hemisphere, where dawn song is strongly

associated with territory defence and mate choice (Catchpole & Slater, 2008). These functions appear

only marginally relevant in species like zebra finches, as they are not territorial and most of their singing

occurs after pair formation. Consequently, song in such species is considered to have its primary

function within established pairs and in social groups (Loning et al., 2024). Also, the very clear

individual distinctiveness of the single individual zebra finch song motif appears almost unique among

songbirds and is likely to facilitate establishing and maintaining social bonds in their dynamic social

organisation. Moreover, the difference in peak singing times at breeding colonies and social hotspots

suggest behavioural routines which may allow individuals to predict when to meet whom, or how many

others, at which location. This pattern might have been specific to the breeding period, as overall birds

may visit hotspots more frequently when they are not breeding, and thus do not have to trade off

activities for self-maintenance and socialising with reproduction, and are not tied to regular visits to a

breeding site. Taken together, the communal singing pattern of zebra finches leading to

communication networks, along with the individual distinctiveness of the song and other differences

in singing compared to territorial species, widens our view on the function of birdsong, and provides

the potential to study how information is shared across communication networks and wider social

organisations.

Next to a higher singing activity, social hotspots also included larger song networks, with more distinct

individuals and singing associations than in the breeding sub-colonies, further supporting the idea that

social hotspots serve as meeting places (Loning et al., 2023a). The larger hotspot network could

facilitate enhanced social information exchange across familiar and unfamiliar individuals, which can

be increasingly valuable in the relatively harsh and unpredictable environment the zebra finch inhabits

(Rafacz & Templeton, 2003; Rubenstein & Lovette, 2007). One benefit could be increased information

on the breeding status of other individuals, which could inform breeding synchronization (Mariette &

Griffith, 2012b; Brandl et al., 2021) as males also sing more in groups during breeding periods than in

periods without breeding (Loning et al., 2023b). While individuals can obtain breeding information by prospecting others' (Mariette & Griffith, 2012a; Brandl et al., 2019b), social hotspots could provide access to information from a wider area and in a more centralised, readily available location. In zebra finches, reproduction is opportunistic and not only regulated by photoperiod, but instead strongly dependent on rainfall and subsequent food availability (Zann, 1994; Zann et al., 1995). Maintaining social ties with (previously successful) breeding neighbours across contexts and over time could facilitate synchrony across multiple breeding attempts (Brandl et al., 2019a; Brandl et al., 2021). Moreover, in captive zebra finches, colony sounds were found to increase breeding synchronisation (Waas et al., 2000), and the vocalizations at hotspots could have a similar effect. While the existence of social hotspots has been shown earlier (Loning et al., 2023a), our finding that the same individuals repeatedly can be identified to meet, expands on earlier findings, showing that the hotspots are not just anonymous gathering sites, but reveal a social structure with repeated encounters. The combination of encountering known and unknown individuals would allow for different levels of information exchange that can be relevant for various movement, foraging and breeding decisions in such an unpredictable environment (Benskin et al., 2002; Aplin et al., 2012; Mariette & Griffith, 2013).

Listening to other birds at social hotspots could also be valuable to gain other types of information. Zebra finch song output reflects, for instance, not only breeding stage (Loning et al., 2023b), but also early developmental background (Holveck et al., 2008; Honarmand et al., 2015), condition (Ritschard & Brumm, 2012) and testosterone levels (Cynx et al., 2005) and may therefore signal condition, foraging success, and fitness. Any such more subtly coded information might be more readily accessible from an already familiar song by known individuals. Indeed, our finding that at least some individuals can be identified to meet repeatedly, suggests that the hotspots are not just anonymous gathering sites but locations to encounter specific familiar individuals, facilitating such more subtle information exchange. Such stable preferential associations have been found in numerous bird species and could have multiple additional benefits (Griffiths et al., 2004; Kurvers et al., 2013; Rose & Croft, 2017; Beck et al., 2020). For example, associating with familiar individuals could reduce the frequency

or intensity of aggressive interactions, thereby conserving energy that could be allocated to vigilance

or predator avoidance (Griffiths et al., 2004). The combination of encountering familiar and less

familiar individuals would also allow for different levels of information exchange that can be relevant

for various movement and foraging decisions in such an unpredictable environment (Benskin et al.,

2002; Aplin et al., 2012; Mariette & Griffith, 2013). Individuals with more social connections potentially

have greater benefits since they have access to more information from others (Wey et al., 2008),

leading to higher efficiency in problem-solving and foraging (Morand-Ferron & Quinn, 2011; Aplin et

al., 2015; Snijders et al., 2018; Snijders et al., 2021). Having a larger network could also aid social

learning, as individuals learn more effectively from familiar conspecifics (Swaney et al., 2001; Cadieu

& Cadieu, 2004). Finally, gathering at hotspots with many conspecifics may reduce the risk of predation

through dilution or increased vigilance (Beauchamp, 2008; Lehtonen & Jaatinen, 2016). Social network

structure indeed can be shaped by predation risk, which is known to strengthen associations in other

species (Kelley et al., 2011; Firth et al., 2017; Heathcote et al., 2017). In our study population, predation

pressure likely varies between years (Naguib et al. 2024), which likely influences temporal variation in

network structures. Further, breeding state may influence social network structure, as many species,

including zebra finches, associate mainly in pairs during breeding periods, but form larger groups

during non-breeding periods (Griesser et al., 2009; Wang & Lu, 2014; Farine & Whitehead, 2015;

Kurvers et al., 2020). This suggests that the loose structure we observed may reflect the breeding

season, and networks could be more cohesive outside this period.

The network observed in this study primarily consisted of single encounters between individuals. Only

14% of dyads encountered each other multiple times, with only 3% interacting three or more times,

indicating that birds were not preferentially associating with specific others within the short time frame

of our analysis. The low rate of repeated dyadic interactions in our data, therefore, points to a loose,

more ephemeral social structure. Yet, most likely the dyads recorded as interacting only once actually

interacted more frequently, either on different days or at other locations. This is supported by the

observation that the relationship between total singing time and the number of detected interactions

did not reach an asymptote, suggesting that additional data collection most likely will reveal a substantially larger and more connected network. In addition, the use of song as a proxy for presence has limitations, as zebra finches do not sing continuously, yet both males and females additionally communicate through frequent calling. Individuals were found to sing for just over half of the time that they were actually present at a hotspot (Loning et al., 2023a), meaning that some associations likely have been missed. Combining acoustic analysis with observations, including the individually specific calls, or a different methodology using an automated radio tracking system or PIT-tags (Brandl et al., 2021; Tyson et al., 2024) could provide a more complete picture of the networks. Such tracking systems would also capture female associations, which are overlooked when relying on song alone, although given their close spatial association (Mariette & Griffith, 2012a; McCowan et al., 2015; Tyson et al., 2024), pair members are expected to form highly similar networks (Brandl et al., 2019a).

The two relatively separate networks at the breeding colony and the social hotspots shown in this study, combined with previous findings of stable pair bonds (Griffith et al., 2010; Tyson et al., 2024), suggest that their social system may share some features of a multi-level social structure, with pairs as the core unit, distinct social associations at the nest and social trees as distinct higher level social units and potentially fusing of these social units later in the season. However, other defining characteristics of a multi-level society were less apparent. Individual membership within grouping levels was flexible, with few repeated associations, resulting in a relatively sparse network. Furthermore, the breeding site network was not clearly nested within the social hotspot network, as we initially predicted, and association strengths did not differ substantially between the breeding colony and social hotspot levels. This does not fit the traditional definition of a multi-level society as a hierarchical social structure (Grueter et al., 2020). Further work is needed to determine whether zebra finch societies show the temporal stability and hierarchical nesting characteristics of a multi-level society or if wild zebra finch social organisation is more consistent with a generally more fluid fission-fusion social structure (Aureli et al., 2008). It is possible that a hierarchical network becomes more evident with longer-term or finer-scale data. Meanwhile, our findings give new insight into the social structure of a widely used model

species. Given this widespread use of zebra finches as a model species in the lab, a clearer

understanding of zebra finch natural social dynamics may also improve the interpretation of lab

findings and inform best practices in captive housing (Griffith et al., 2017).

Taken together, our findings show that wild zebra finches sing near their breeding site and at social

hotspots at different times of day, with a distinct lack of dawn song. This highlights that the context

and potential function of their singing differ from those of classically studied territorial songbirds, as

previously noted for other song characteristics (Loning et al., 2024). Our establishment of using

BirdNET embeddings for the validation of individual classification demonstrates a promising approach

to scale up the study of communication networks, opening the door for more advanced future studies.

Male song revealed different levels of social organisation, with individuals connecting with more and

different conspecifics at social hotspots than at the breeding colony, and a few individuals maintained

associations across both sites, linking sub-units and suggesting elements of multi-level societies where

information can flow through vocal interactions. However, the overall low strength of the networks

suggests that birds mostly move around as pairs, meeting others opportunistically at social hotspots

or the breeding area. This reflects a social organization that is at the interface between a multi-level

society of stable hierarchical subgroups and a more dynamic fission-fusion society with individualistic

network structures. Such a dynamic system might be adaptive in the harsh and unpredictable

environment of the Australian Outback, as birds have stable social subunits (the pair) connected to

different familiar individuals at different locations, providing relevant information on resource

acquisition, predation risk or breeding state. Encountering a variety of other, socially less stable

connected individuals will allow both males and females to draw information from a wider network,

and may be important to gather population-wide information, for instance, on dispersal and

immigration patterns, as well as potential novel food sources, such as when unfamiliar individuals are

in better condition.

Acknowledgements & Artificial Intelligence statement

We are grateful to Marie-Anne Bellon and Tis Voortman for their assistance during fieldwork and Vicki and Garry Dowling from Fowlers Gap Arid Zone Research Station for their support. We thank Joan Heikamp and Louis Diekmann for video analysis and Aneesh Chauhan for advice on the deep-learning aspect of the data validation.

References

Amy, M., Sprau, P., de Goede, P., & Naguib, M. (2010). Effects of personality on territory defence in communication networks: a playback experiment with radio-tagged great tits. *Proceedings of the Royal Society B, 277*, 3685-3692. doi: https://doi.org/10.1098/rspb.2010.0598

Aplin, L. M., Farine, D. R., Morand-Ferron, J., Cockburn, A., Thornton, A., & Sheldon, B. C. (2015). Experimentally induced innovations lead to persistent culture via conformity in wild birds. *Nature, 518*(7540), 538-541. doi: 10.1038/nature13998

Aplin, L. M., Farine, D. R., Morand-Ferron, J., & Sheldon, B. C. (2012). Social networks predict patch discovery in a wild population of songbirds. *Proceedings of the Royal Society B: Biological Sciences, 279*(1745), 4199-4199. doi: 10.1098/RSPB.2012.1591

Aureli, F., Schaffner, C. M., Boesch, C., Bearder, S. K., Call, J., Chapman, C. A., . . . van Schaik, C. P. (2008). Fission-fusion dynamics: new research frameworks. *Current Anthropology, 49*, 627-654.

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software, 67*, 1-48. doi: 10.18637/jss.v067.i01

Beauchamp, G. (2008). What is the magnitude of the group-size effect on vigilance? *Behavioral Ecology, 19*(6), 1361-1368. doi: 10.1093/BEHECO/ARN096

Beck, K. B., Farine, D. R., & Kempenaers, B. (2020). Winter associations predict social and extra-pair mating patterns in a wild songbird. *Proceedings of the Royal Society B, 287*(1921). doi: 10.1098/RSPB.2019.2606

Benskin, C. M. H., Mann, N. I., Lachlan, R. F., & Slater, P. J. B. (2002). Social learning directs feeding preferences in the zebra finch, *Taeniopygia guttata*. *Animal Behaviour, 64*(5), 823-828. doi: 10.1006/ANBE.2002.2005

Brandl, H. B., Farine, D. R., Funghi, C., Schuett, W., & Griffith, S. C. (2019a). Early-life social environment predicts social network position in wild zebra finches. *Proceedings of the Royal Society, London, Series B,, 286*, 20182579.

Brandl, H. B., Griffith, S. C., Farine, D. R., & Schuett, W. (2021). Wild zebra finches that nest synchronously have long-term stable social ties. *Journal of Animal Ecology, 90*, 76–86. doi: https://doi.org/10.1111/1365-2656.13082

Brandl, H. B., Griffith, S. C., Laaksonen, T., & Schuett, W. (2019b). Begging calls provide social cues for prospecting conspecifics in the wild zebra finch (*Taeniopygia guttata*). *The Auk, 136*(2). doi: 10.1093/auk/ukz007

Brandl, H. B., Griffith, S. C., & Schuett, W. (2019c). Wild zebra finches choose neighbours for synchronized breeding. *Animal Behaviour 151*, 21-28. doi: https://doi.org/10.1016/j.anbehav.2019.03.002

Cadieu, N., & Cadieu, J. C. (2004). The influence of free interactions and partner familiarity on social transmission in the young canary. *Animal Behaviour, 67*(6), 1051-1057. doi: https://doi.org/10.1016/j.anbehav.2003.10.004

Calcari, C., Pilenga, C., Baciadonna, L., Gamba, M., & Favaro, L. (2021). Long-term stability of vocal individuality cues in a territorial and monogamous seabird. *Animal Cognition, 24*, 1165–1169. doi: https://doi.org/10.1007/s10071-021-01518-z

Catchpole, C. K., & Slater, P. J. B. (2008). *Bird Song: biological themes and variation*. New York: Cambridge University Press.

Croft, D. P., James, R., & Krause, J. (2008). *Exploring animal social networks*. Princeton:

Princeton University Press.

Csárdi, G., & Nepusz, T. (2006). The igraph software package for complex network research.

Complex Systems(1695).

Cure, C., Aubin, T., & Mathevon, N. (2009). Acoustic convergence and divergence in two

sympatric burrowing nocturnal seabirds. *Biological Journal of the Linnean Society, 96*,

115–134. doi: https://doi.org/10.1111/j.1095-8312.2008.01104.x

Cynx, J., Bean, N. J., & Rossman, I. (2005). Testosterone implants alter the frequency range of

zebra finch songs. *Hormones and Behavior, 47*, 446-451.

Devogel, M., Le Fur, R., Grissot, A., Kidawa, D., Araya-Salas, M., & Wojczulanis-Jakubas, K.

(2024). Vocal interactions of breeding partners predict duration of incubation bout in an

Artic seabird. *Journal of Ornithology, 165*, 987-995.

Elie, J. E., & Theunissen, F. E. (2018). Zebra finches identify individuals using vocal signatures

unique to each call type. *Nature Communications, 9*(1), 4026. doi: 10.1038/s41467-018-

06394-9

Farine, D. R., & Whitehead, H. (2015). Constructing, conducting and interpreting animal social

network analysis. *Journal of Animal Ecology, 84*(5), 1144-1163. doi:

https://doi.org/10.1111/1365-2656.12418

Firth, J. A., Voelkl, B., Crates, R. A., Aplin, L. M., Biro, D., Croft, D. P., & Sheldon, B. C. (2017).

Wild birds respond to flockmate loss by increasing their social network associations to

others. *Proceedings of the Royal Society B: Biological Sciences, 284*(1854). doi:

10.1098/RSPB.2017.0299

Fitzsimmons, L. P., Foote, J. R., Ratcliffe, L. M., & Mennill, D. J. (2008). Eavesdropping and

communication networks revealed through playback and an acoustic location system.

Behavioral Ecology, 19(4), 824-829. doi: https://doi.org/10.1093/beheco/arn036

Foote, J. R., Fitzsimmons, L. P., Mennill, D. J., & Ratcliffe, L. M. (2010). Black-capped chickadee

622   dawn choruses are interactive communication networks. *Behaviour, 147*(10), 1219-

623   1248. doi: 10.1163/000579510x513761

624   Griesser, M., Barnaby, J., Schneider, N. A., Figenschau, N., Wright, J., Griffith, S. C., . . . Russell,

625   A. F. (2009). Influence of winter ranging behaviour on the social organization of a

626   cooperatively breeding bird species, the apostlebird. *Ethology, 115*(9), 888-896. doi:

627   10.1111/J.1439-0310.2009.01678.X

628   Griffith, S. C., & Buchanan, K. L. (2010). The Zebra Finch: the ultimate Australian supermodel.

629   *Emu, 110*(3), V-xii. doi: 10.1071/MUv110n3_ED

630   Griffith, S. C., Crino, O. L., Adkins-Regan, E., Alonso-Alvarez, C., Andrew, S. C., Bailey, I. E., . . .

631   Williams, T. D. (2017). Variation in reproductive success across captive populations:

632   methodological differences, potential biases and opportunities. *Ethology, 123*, 1-29.

633   Griffith, S. C., Holleley, C. E., Mariette, M. M., Pryke, S. R., & Svedin, N. (2010). Low level of

634   extrapair parentage in wild zebra finches. *Animal Behaviour, 79*(2), 261-264. doi:

635   10.1016/J.ANBEHAV.2009.11.031

636   Griffith, S. C., Pryke, S. R., & Mariette, M. (2008). Use of nest-boxes by the Zebra Finch

637   (*Taeniopygia guttata*): implications for reproductive success and research. *Emu - Austral*

638   *Ornithology, 108*(4), 311-319. doi: 10.1071/MU08033

639   Griffiths, S. W., Brockmark, S., Höjesjö, J., & Johnsson, J. I. (2004). Coping with divided attention:

640   the advantage of familiarity. *Proceedings of the Royal Society of London. Series B:*

641   *Biological Sciences, 271*(1540), 695-699. doi: 10.1098/RSPB.2003.2648

642   Grueter, C. C., Qi, X., Zinner, D., Bergman, T., Li, M., Xiang, Z., . . . Swedell, L. (2020). Multilevel

643   organisation of animal sociality *Trends in Ecology and Evolution* (Vol. 35, pp. 834-847):

644   Elsevier Ltd.

645   Hartig, F. (2024). DHARMa: Residual diagnostics for hierarchical (multi-level / mixed) regression

646   models. *R package version 0.4.7, https://github.com/florianhartig/dharma*. .

Heathcote, R. J., Darden, S. K., Franks, D. W., Ramnarine, I. W., & Croft, D. P. (2017). Fear of

predation drives stable and differentiated social relationships in guppies. *Scientific*

Reports, 7, 41679. doi: https://doi.org/10.1038/srep41679

Holveck, M. J., de Castro, A. C. V., Lachlan, R. F., ten Cate, C., & Riebel, K. (2008). Accuracy of

song syntax learning and singing consistency signal early condition in zebra finches.

Behavioral Ecology, 19(6), 1267-1281. doi: 10.1093/beheco/arn078

Honarmand, M., Riebel, K., & Naguib, M. (2015). Nutrition and peer group composition in early

adolescence: impacts on male song and female preference in zebra finches. *Animal*

Behaviour, 107, 147-158.

Kahl, S., Wood, C. M., Eibl, M., & Klinck, H. (2021). BirdNET: A deep learning solution for avian

diversity monitoring. *Ecological Informatics, 61*, 101236.

Kelley, J. L., Morrell, L. J., Inskip, C., Krause, J., & Croft, D. P. (2011). Predation risk shapes social

networks in fission-fusion populations. *Plos One, 6*(8). doi:

https://doi.org/10.1371/journal.pone.0024280

Kunc, H. P., Amrhein, V., & Naguib, M. (2005). Seasonal variation of dawn song and its relation to

mating success in the nightingale (*Luscinia megarhynchos*). *Animal Behaviour, 70*, 1265-

1271.

Kurvers, R. H., Prox, L., Farine, D. R., Jongeling, C., & Snijders, L. (2020). Season-specific

carryover of early life associations in a monogamous bird species. *Animal Behaviour,*

164, 25-37.

Kurvers, R. H. J. M., Adamczyk, V. M. A. P., Kraus, R. H. S., Hoffman, J. I., van Wieren, S. E., van

der Jeugd, H. P., . . . Jonker, R. M. (2013). Contrasting context dependence of familiarity

and kinship in animal social networks. *Animal Behaviour, 86*(5), 993-1001. doi:

10.1016/J.ANBEHAV.2013.09.001

Lehtonen, J., & Jaatinen, K. (2016). Safety in numbers: the dilution effect and other drivers of

group life in the face of danger. *Behavioral Ecology and Sociobiology, 70*(4), 449-458.

doi: 10.1007/S00265-016-2075-5/TABLES/1

Lenth, R. (2025). _emmeans: Estimated marginal means, aka least-squares means. *R package*

version 1.11.2, https://CRAN.R-project.org/package=emmeans. doi:

doi:10.32614/CRAN.package.emmeans

Logue, D. M. (2021). Countersinging in birds. *Advances in the Study of Behavior, 53*, 1-61. doi:

https://doi.org/10.1016/bs.asb.2021.03.001

Loning, H., Fragueira, R., Naguib, M., & Griffith, S. C. (2023a). Hanging out in the outback: the

use of social hotspots by wild zebra finches. *Journal of Avian Biology*, e03140. doi:

https://doi.org/10.1111/jav.03140

Loning, H., Griffith, S. C., & Naguib, M. (2021). Zebra finch song is a very short-range signal in the

wild: evidence from an integrated approach. *Behavioral Ecology, 33*(1), 37-46. doi:

10.1093/BEHECO/ARAB107

Loning, H., Griffith, S. C., & Naguib, M. (2024). The ecology of zebra finch song and its

implications for vocal communication in multi-level societies. *Philosophical*

Transactions B, 379(1905). doi: 10.1098/RSTB.2023.0191

Loning, H., Verkade, L., Griffith, S. C., & Naguib, M. (2023b). The social role of song in wild zebra

finches. *Current Biology, 33*, 372-380. doi: https://doi.org/10.1016/j.cub.2022.11.047

Major, H. L., & Jones , I. L. (2011). An experimental study of the use of social information by

prospecting nocturnal burrow-nesting seabirds. *The Condor: Ornithological*

Applications, 113, 572–580. doi: https://doi.org/10.1525/cond.2011.100088

Mariette, M. M., & Griffith, S. C. (2012a). Conspecific attraction and nest site selection in a

nomadic species, the zebra finch. *Oikos, 121*(6), 823-834. doi: 10.1111/j.1600-

0706.2011.20014.x

Mariette, M. M., & Griffith, S. C. (2012b). Nest visit synchrony is high and correlates with

reproductive success in the wild Zebra finch *Taeniopygia guttata*. *Journal of Avian*

Biology, 43(2), 131-140. doi: 10.1111/j.1600-048X.2012.05555.x

Mariette, M. M., & Griffith, S. C. (2013). Does coloniality improve foraging efficiency and nestling

provisioning? A field experiment in the wild Zebra Finch. *Ecology, 94*(2), 325-335.

McCowan, L. S. C., Mariette, M. M., & Griffith, S. C. (2015). The size and composition of social

groups in the wild zebra finch. *Emu, 115*(3), 191-198.

McGinn, K., Kahl, S., Peery, M. Z., Klinck, H., & Wood, C. M. (2023). Feature embeddings from

the BirdNET algorithm provide insights into avian ecology. *Ecological Informatics, 74*,

101995.

McGregor, P. K. (2005). *Communication networks*. Cambridge: Cambridge University Press.

McGregor, P. K., & Dabelsteen, T. (1996). Communication networks. In D. E. Kroodsma & E. H.

Miller (Eds.), *Ecology and Evolution of Acoustic Communication in Birds* (pp. 409-425).

Cornell: University Press.

Mennill, D. J., & Ratcliffe, L. M. (2004). Do male black-capped chickadees eavesdrop on song

contests? A multi-speaker playback experiment. *Behaviour, 141*, 125-139. doi:

https://doi.org/10.1163/156853904772746637

Morand-Ferron, J., & Quinn, J. L. (2011). Larger groups of passerines are more efficient problem

solvers in the wild. *Proceedings of the National Academy of Sciences of the United*

States of America, 108(38), 15898-15903. doi:

10.1073/PNAS.1111560108/SUPPL_FILE/SM01.MOV

Naguib, M., Amrhein, V., & Kunc, H. P. (2004). Effects of territorial intrusions on eavesdropping

neighbors: communication networks in nightingales. *Behavioral Ecology, 6*, 1011-1015.

Naguib, M., Fichtel, C., & Todt, D. (1999). Nightingales respond more strongly to vocal leaders in

simulated dyadic interactions. *Proceedings of the Royal Society B, 265*, 537-542. doi:

https://doi.org/10.1098/rspb.1999.0669

Naguib, M., Kunc, H. P., Sprau, P., Roth, T., & Amrhein, V. (2011). Communication networks and spatial ecology in nightingales. *Advances in the Study of Behavior, 43*, 239-271.

Otter, K., McGregor, P. K., Terry, A. M. R., Burford, F. R. L., Peake, T. M., & Dabelsteen, T. (1999). Do female great tits (*Parus major*) assess males by eavesdropping? A field study using interactive song playback. *Proceedings of the Royal Society of London B, 266*(1426), 1305-1309. doi: https://doi.org/https://doi.org/10.1098/rspb.1999.0779

Papageorgiou, D., Christensen, C., Gall, G. E. C., Klarevas-Irby, J. A., Nyaguthii, B., Couzin, I. D., & Farine, D. R. (2019). The multilevel society of a small-brained bird. *Current Biology, 29*(21), R1120-R1121.

Papageorgiou, D., & Farine, D. R. (2021). Multilevel societies in birds. *Trends in Ecology & Evolution, 36*, 15-17. doi: https://doi.org/10.1016/j.tree.2020.10.008.

Peake, T. M. (2005). Eavesdropping in communication networks. In P. K. McGregor (Ed.), *Communication networks* (pp. 13 - 37). Cambridge: Cambridge University Press.

Peake, T. M., Matessi, G., McGregor, P., L., & Dabelsteen, T. (2005). Song type matching, song type switching and eavesdropping in male great tits. *Animal Behaviour, 69*, 1063-1068.

Peake, T. M., Terry, A. M. R., McGregor, P. K., & Dabelsteen, T. (2001). Male great tits eavesdrop on simulated male-to-male vocal interactions. *Proceedings of the Royal Society London B, 268*(1472), 1183-1187. doi: https://doi.org/10.1098/rspb.2001.1648

Rafacz, M., & Templeton, J. J. (2003). Environmental unpredictability and the value of social information for foraging starlings. *Ethology, 109*(12), 951-960. doi: 10.1046/J.0179-1613.2003.00935.X

Ritschard, M., & Brumm, H. (2012). Zebra finch song reflects current food availability. *Evolutionary Ecology, 26*(4), 801-812. doi: 10.1007/S10682-011-9541-3/FIGURES/3

Rose, P. E., & Croft, D. P. (2017). Social bonds in a flock bird: Species differences and seasonality in social structure in captive flamingo flocks over a 12-month period. *Applied Animal Behaviour Science, 193*, 87-97. doi: 10.1016/J.APPLANIM.2017.03.006

Rubenstein, D. R., & Lovette, I. J. (2007). Temporal environmental variability drives the evolution

of cooperative breeding in birds. *Current Biology, 17*(16), 1414-1419. doi:

10.1016/J.CUB.2007.07.032

Schmidt, R., Kunc, H. P., Amrhein, V., & Naguib, M. (2006). Responses to interactive playback

predict future mating status in nightingales. *Animal Behaviour, 72*, 1355-1362. doi:

https://doi.org/10.1016/j.anbehav.2006.03.022

Snijders, L., Krause, S., Tump, A. N., Breuker, M., Ortiz, C., Rizzi, S., . . . Kurvers, R. H. J. M.

(2021). Causal evidence for the adaptive benefits of social foraging in the wild.

Communications Biology, 4, 1-8.

Snijders, L., Kurvers, R. H., Krause, S., Ramnarine, I. W., & Krause, J. (2018). Individual- and

population-level drivers of consistent foraging success across environments. *Nature*

Ecology and Evolution, 2, 1610–1618. doi: https://doi.org/10.1038/s41559-018-0658-4

Snijders, L., & Naguib, M. (2017). Communication in animal social networks: a missing link? .

Advances in the Study of Behavior, 49, 297-359. doi:

https://doi.org/10.1016/bs.asb.2017.02.004

Sossinka, R., & Böhner, J. (1980). Song types in the zebra finch *Poephila guttata castanotis*.

Zeitschrift für Tierpsychology, 53, 123-132.

Staicer, C. A., Spector, D. A., & Horn, A. G. (1996). The dawn chorus and other diel patterns in

acoustic signaling. In D. E. Kroodsma & E. H. Miller (Eds.), *Ecology and evolution of*

acoustic communication in birds. London: Cornell University Press.

Sturdy, C. B., Phillmore, L. S., & Weisman, R. G. (1999). Note types,, harmonic structure, and

note order in the songs of zebra finches (*Taeniopygia guttata*). *Journal of Comparative*

Psychology, 113, 194-203.

Swaney, W., Kendal, J., Capon, H., Brown, C., & Laland, K. N. (2001). Familiarity facilitates social

learning of foraging behaviour in the guppy. *Animal Behaviour, 62*(3), 591-598. doi:

10.1006/ANBE.2001.1788

Todt, D., & Naguib, M. (2000). Vocal interactions in birds: The use of song as a model in

communication. *Advances in the Study of Behavior, 29*, 247-296. doi: 10.1016/s0065-

3454(08)60107-2

Tyson, C., Loning, H., Griffith, S. C., & Naguib, M. (2024). Constant companions: wild zebra finch

pairs display extreme spatial cohesion. *Biology Letters, 20*(11). doi:

10.1098/RSBL.2024.0519

Waas, J. R., Caulfield, M., Colgan, P. W., & Boag, P. T. (2000). Colony sound facilitates sexual

and agonistic activities in royal penguins. *Animal Behaviour, 60*, 77-84.

Wang, C., & Lu, X. (2014). Dispersal in kin coalition throughout the non-breeding season to

facilitate fine-scale genetic structure in the breeding season: evidence from a small

passerine. *Ethology, 120*(10), 1003-1012. doi: 10.1111/ETH.12273

Wang, G., Harpole, C. E., Trivedi, A. K., & Cassone, V. M. (2012). Circadian regulation of bird

song, call, and locomotor behavior by pineal melatonin in the zebra finch. *Journal of*

Biological Rhythms, 27(2), 145-155. doi:

10.1177/0748730411435965/SUPPL_FILE/DS_10.1177_0748730411435965.PDF

Ward, A., & Webster, M. (2016). *Sociality: The Behaviour of Group-Living Animals*. Cham:

Springer International Publishing.

Wey, T., Blumstein, D. T., Shen, W., & Jordán, F. (2008). Social network analysis of animal

behaviour: a promising tool for the study of sociality. *Animal Behaviour, 75*(2), 333-344.

doi: 10.1016/J.ANBEHAV.2007.06.020

Zann, R. A. (1994). Reproduction in a zebra finch colony in South—Eastern Australia - the

significance of monogamy, precocial breeding and multiple broods in a highly mobile

species. *Emu, 94*, 285-299.

Zann, R. A. (1996). *The zebra finch: A synthesis of field and laboratory studies*: Oxford University

Press.

Zann, R. A., Morton, S. R., Jones, K. R., & Burley, N. T. (1995). The timing of breeding by zebra

finches in relation to rainfall in Central Australia. *Emu, 95*(3), 208-222. doi:

10.1071/MU9950208