Assessing the digit organisation of focal hand dystonia using 7T functional MRI

[1]Michael Asghar, [1,2,3]Rosa Sanchez-Panchuelo, [1]Paul Glover, [1,4,5]Daisie Pakenham, [1]George C. O'Neill, [6]Denis Schluppeck, [7]Miles Humberstone, [1,2]Susan Francis

Affiliations:

[1]Sir Peter Mansfield Imaging Centre, School of Physics and Astronomy, University of Nottingham, United Kingdom.

[2]NIHR Nottingham Biomedical Research Centre, Nottingham University Hospitals NHS Trust and the University of Nottingham, Nottingham, United Kingdom

[3]University Hospitals Birmingham NHS Foundation Trust, United Kingdom.

[4]Clinical Neurophysiology, Birmingham Women's and Children's NHS Foundation Trust, Birmingham, United Kingdom.

[5]Institute of Health and Neurodevelopment (IHN), Aston University, Birmingham, United Kingdom.

[6]School of Psychology, University of Nottingham, Nottingham, United Kingdom.

[7]Nottingham University Hospitals NHS Trust, Nottingham, United Kingdom.

Pages: 31

Figures: 8

Supplementary Figures: 4

Tables: 1

Word count: 8436

ABSTRACT

Introduction: Focal hand dystonia (FHD) is typically treated with Botulinum Toxin (BTX/BoNT-A/Botox/ Onabotulinum Toxin Type A) injected into digit and wrist muscles to provide temporary symptomatic relief from muscle spasm. We use 7T functional MRI (fMRI) to study 1) differences in somatosensory and motor cortical digit maps between FHD patients and age and sex-matched healthy volunteers (HV), 2) changes in cortical digit maps of FHD patients after BTX treatment.

Methods: Four FHD patients underwent behavioural measures and 7T fMRI at two timepoints: ~30 days after BTX treatment at peak efficacy (Post-BTX), immediately prior to their next injection ~100 days after treatment when therapeutic effects had declined (Pre-BTX). HVs were scanned at a single time point. Behavioural measures included temporal discrimination (TDT), amplitude thresholding (AMP) and spatial acuity grating orientation tasks. A somatosensory travelling wave (TW) digit-mapping fMRI task was performed on dominant/affected and non-dominant/un-affected hands, and a motor TW digit-mapping task on the dominant/affected hand. Fourier analysis derived digit maps which were compared to a probabilistic digit atlas, and population receptive field (pRF) mapping performed. Structural and resting state fMRI data were acquired.

Results: Spatial acuity was lowered in the dominant/affected hand in FHD compared to HVs, while TDT and AMP measures showed no significant differences. Somatosensory and motor TW cortical digit maps in FHD and HV showed clear digit ordering. Somatosensory and motor cortical digit maps were more disordered in the dominant/affected hand than the non-dominant hand for FHD Post and Pre-BTX compared to HVs (DICE coefficient/Figure of Merit). Compared to the atlas, HV maps were in-line for both hands, while in FHD central tendency (CT) was lower in the dominant/affected hand than non-dominant hand. pRF sizes were larger in the FHD Pre-BTX group compared to HV and Post-BTX for the dominant hand.

Conclusion: Behaviourally, FHD patients had lower spatial acuity than HVs. FHD digit maps had typical D1-D5 ordering, but DICE coefficients identified a disordered dominant hand in FHD patients compared to HVs at both timepoints. The FHD Pre-BTX group had larger pRF sizes than Post-BTX and HVs, suggesting BTX treatment coincides with more localised touch perception.

58 **Keywords**: Focal Hand Dystonia, botulinum toxin, functional MRI, somatosensory

59 function, spatial acuity, digit mapping

1. INTRODUCTION

Focal Hand Dystonia is a sensorimotor disorder presenting with involuntary movements, tremors and weakness in the hand (1). Early brain imaging studies of focal hand dystonia (FHD) have suggested disordered somatosensory function demonstrated by reduced distances between the cortical representation of digits and increased overlapping activity compared to healthy controls (2,3). However, because the cortical digit maps of somatosensation and motor function representing the hand are at the millimetre scale, these findings of disorder are now being questioned as higher spatial resolution fMRI become available, motivating this work.

Functional MRI (fMRI) studies probing the cortical digit representation use block (4,5), phase-encoding (6) and event-related (7) designs to deliver sensory stimulation to the fingers, mostly in healthy individuals. In the past decade, there has been a push to investigate digit maps at higher magnetic field strength, due to the increased BOLD contrast and spatial resolution (8,9). Somatotopic digit mapping in human primary somatosensory cortex using a travelling-wave (TW) phase-encoding fMRI paradigm has been shown to be highly reproducible in healthy participants at 7T (10) but show inter-participant variability. To address this, we developed a probabilistic somatosensory atlas (11) to define digit dominance and inter-participant variability in healthy participants. This atlas provides a standardised template for comparison with patient groups, such as those people with Focal Hand Dystonia (FHD).

Prior studies in dystonia have principally used motor tasks. A study in dystonic musicians (12) showed greater activity in ipsilateral premotor area and reduced activity in the cerebellum during a right-handed tapping task, suggesting irregular neural activity and compensatory mechanisms compared to healthy controls. In contrast, a study of a cued movement task in FHD patients (13) showed a decrease in ipsilateral activity and an increase in contralateral activity associated with the affected hand. A further study (14) showed that motor performance to a button press task was similar between controls and FHD patients, while the motor preparation section of the task showed reduced activation in premotor and postcentral gyrus for FHD patients, implying impaired early stages of motor control. A 3T fMRI study using passive extension and active finger presses of individual fingers in musician's dystonia (n=9) (15) discerned individual digits

and clear somatotopy but showed no difference in the spatial geometry (expansion or spatial shift of digit representation) for musicians with and without dystonia. This suggestion of the intact cortical digit maps in musician's dystonia contests the idea that task-specific dystonia alters cortical maps and lends weight to an alternative hypothesis that task-specific dystonia is due to a higher-order disruption of skill encoding. A recent high spatial resolution (0.75x0.75x0.89mm) laminar 7T MRI Vascular Space Occupancy (VASO) study showed in a small group of individuals with FHD (n=4) a breakdown of digit somatotopy and increased activation in superficial cortico-cortical input layers during finger tapping compared to unaffected hemispheres, suggested to be due to reduced inhibition (16).

Together, these studies of FHD suggest widespread differences in sensorimotor areas and subcortical structures. However, these prior studies have generally been performed at a field strength of 3T or less and have used spatial analyses (e.g. centre of gravity of each digit based on vertices and searchlight analysis) in volumetric space. There is therefore a need for high spatial resolution high field fMRI studies using surface-based methods to provide more accurate and more sensitive results (17) and better digit delineation across gyral and sulcal boundaries. More recently, fMRI studies have measured how receptive fields of neurons in the somatosensory cortex are distributed, for example by using a 2D grid of stimulators to map within- and between-digit characteristics. Computational models have been used to investigate more fine-grained characteristics of the hand, such as population receptive field (pRF) size, in healthy participants (18–21).

Resting state fMRI (rs-fMRI) has also shown alterations in functional connectivity (FC) in dystonia. In Writer's Cramp (WC) (22) reduced functional connectivity compared with controls was shown when placing seeds in left lateral premotor cortex, left thalamus, left/right pallidum to study the projection to the symptomatic left primary sensorimotor cortex. There were also reductions in FC from primary motor cortex premotor, frontal and somatosensory areas. Reductions in FC were also shown in cervical dystonia, in regions related to the M1-SMA motor network (23). Another study showed abnormal recruitment of parietal and premotor cortices, as well as distinct patterns of sensorimotor integrations between musician's and non-musician's dystonia (24). Unique patterns for dystonic tremor, based on a reduction in FC were found in

higher-level cortical, basal ganglia, and cerebellar regions (25). Further, a study of
cervical dystonia and blepharospasm (26) suggested that focal dystonia is a disorder of
specific networks and not just a result of basal ganglia alterations.

Behaviourally, there is evidence of raised temporal thresholds in patients as
determined with temporal discrimination tasks and spatial discrimination thresholds
using a grating orientation task (GOT) (27,28).

A symptomatic treatment of FHD involves injections of Botulinum Toxin (BTX), into
the muscles controlling digit and /or wrist movement. This can induce relief from
involuntary muscle spasm associated with FHD. A prior fMRI study in patients with
cervical dystonia affecting the neck showed an increase in activation in sensorimotor
cortex, 4 weeks after BoNT injection (29).

Here we used high spatial resolution 7T fMRI and surface-based analyses to study
cortical somatotopy digit maps and the pRF digit size and perform rs-fMRI in a group of
FHD patients and healthy controls (HV). Patients were scanned at two timepoints in the
BTX treatment cycle of, ~30 days after localised BTX injection at peak efficacy (Post-BTX),
and immediately prior to their next injection ~100 days after treatment when therapeutic
effects have declined (Pre-BTX). At each timepoint, behavioural tasks of temporal
discrimination, amplitude thresholding and spatial acuity were collected.

We test the following hypotheses:

1. Behavioural thresholds (temporal and spatial) are raised in FHD patients compared
to healthy controls.

2. In FHD, digit maps derived from TW mapping (11) are blurred/have poorer delineation
compared to healthy controls,

3. pRF sizes are increased in FHD patients compared to healthy controls.

4. Functional connectivity (FC) to the somatosensory cortex is reduced in FHD
compared to healthy controls.

For these measures we predict FHD responses at 4 weeks following BTX treatment are
more like those of healthy controls compared those at 3 months following BTX treatment.

2. MATERIALS AND METHODS

The study was conducted with the approval of the University of Nottingham Faculty of Medicine and Health Sciences Ethics Committee for healthy volunteers and of the Health Research Authority Research Ethics Committee (REC 17/EM/0368) for FHD patients. FHD patients were recruited from the Botulinum Toxin Clinic at Queens Medical Centre. Eligibility criteria included patients 18 and 75 years who were clinically assessed to have FHD and presented with unilateral symptoms on their dominant hand. Exclusion criteria included participants with a history of neurological illness other than FHD and MRI contraindications. All participants gave written informed consent and completed a MR safety screening form and a handedness form at each visit.

Six patients (2 visits) with Focal Hand Dystonia (FHD) (age mean ± stdev: 54 ± 11 yrs, 4 male) were recruited to be scanned at 7T at both 4 weeks (20-27days) (Post-BTX) and 3 months (105-119 days) after (Pre-BTX) Onabotulinumtoxin type-A (Botox, BTX) treatment (30), along with age-matched healthy controls (HVs) (age mean ± stdev: 42 ± 7 yrs. At each visit, participants completed 1hour of behavioural data collection followed by 7T functional and structural MRI scanning lasting 1hour. In all FHD participants, the affected hand was also the dominant hand.

2.1 Behavioural data acquisition

Behavioural data collection included a temporal discrimination task ("TDT") and an amplitude threshold task ("AMP") coded in MATLAB (The Mathworks, Natick, Massachusetts), and spatial acuity grating orientation task ("GOT"). These tasks were performed using identical piezoelectric stimulators to those used in the fMRI study, while the GOT used Perspex domes.

TDT: The temporal discrimination task was performed on the tips of digits D2 and D3 of both the left and right hands using piezo stimulators, as well as a Brain Gauge device (31). On each trial, two brief stimuli were presented sequentially to D2 and D3, and participants had to indicate the digit they felt the stimulus appear first verbally; the experimenter noted down their responses. To estimate each participant's temporal discrimination threshold, we used an adaptive 1-up, 2-down staircase procedure

implemented via the *upDownStaircase()* function from MGL (32). This staircase method

converges on the stimulus level at which participants respond correctly 70.7% of the

time (33). The staircase ended after 8 reversals (changes in staircase direction).

AMP: To determine, participants completed a two-interval forced choice task to estimate

on the left and right index finger (D2) at 30 Hz (match fMRI paradigm) and 200 Hz. On each

trial, participants were presented with two brief tactile stimuli in succession: One

stimulus was at baseline, near-threshold amplitude, the other was a test stimulus at

higher amplitude, adjusted trial-by-trial via a 1-up, 2-down staircase procedure. The

order of the baseline and test stimulus was randomized across trials. Participants were

asked to indicate which of the two stimuli felt stronger. The task consisted of 60 trials,

and the threshold estimate was taken as the mean amplitude across the final reversals.

This procedure adaptively estimated the smallest amplitude difference the participant

could reliably detect above baseline.

GOT: This grating orientation task was used to assess spatial acuity (34) of the tip of D2,

due to time constraints only one hand was assessed: for FHD patients the affected hand

was studied, and for controls the dominant hand. Participants were presented with

machine-cut Perspex domes, each with a raised grating surface of a specific spatial

frequency (defined by the groove-to-groove gap width in mm). On each trial, the

experimenter applied a dome to the fingertip in one of two orientations: 0°: aligned with

the long axis of the finger ("forwards"), 90°: perpendicular to the long axis ("sideways").

The participant, unable to see the dome, was asked to report whether the grating was

oriented along (0°) or across (90°) the digit, by saying "forwards" or "sideways". Dome

orientation was randomised on each trial using MATLAB code. Each grating width was

tested in 20 trials (10 trials per orientation), and orientations were pseudorandomised

within each spatial frequency. Accuracy was recorded manually. Different stimulus sets

were used for patient and controls to ensure a suitable range for estimating threshold

performance in both groups (Controls: 0.75, 1.0, 1.2, 1.5, 1.75, 2.0, 2.5, 3.0 mm; 2)

Patients: 0.75, 1.0, 1.5, 1.75, 2.0, 2.5, 3.0, 4.0 mm), since patients were unable to reliably

discriminate orientation even at larger widths.

In addition, 10 healthy participants (age mean $\pm$ stdev: 36 $\pm$ 11 yrs) were recruited

to assess the repeatability of the behavioural tasks. The participants underwent the

same procedure as those in the main study (TDT using piezo, AMP and GOT). The tasks

were each performed twice, separated by a two-week interval. This allowed both the

within- and between-participant coefficient of variation (CV) of measures to be

computed, and the comparison of the CV to (Mikkelsen et al. 2020) (35) who assessed

reproducibility of vibrotactile detection and discrimination.

2.2 MRI data acquisition

Data were collected on a 7 T Philips Achieva scanner using a 32-channel receive coil

(Nova Medical). Travelling wave (TW) fMRI data was collected to assess somatotopy and

motortopy. Somatosensory/motortopy fMRI scans were collected using an axial 48 slice

GE-EPI BOLD acquisition with TE/TR = 25/2000 ms, SENSE 1.5, 1.5 mm isotropic

resolution collected using multiband (MB) factor 3.

Somatosensory stimuli were delivered to the fingertips (D1/D2/D3/D4/D5) using

MR-compatible piezo-electric stimulators driven at 30 Hz vibrotactile frequency (Dancer

Design, UK; http://www.dancerdesign.co.uk/). Stimulation was on for 4s per digit, with

20s per cycle, for 8 cycles. This was performed in the forward and reverse direction on

each hand. (10). Somatotopic maps were generated for both the left and right hands in

each scan session (Post-BTX and Pre-BTX for FHD patients).

Motortopy was performed in the dominant hand. During the motortopy task,

participants were visually cued by 5 blinking (1Hz) circles to perform in-air tapping with

the same timings as the somatosensory TW task. During the motortopy task, an

accelerometer glove, designed in-house to monitor 3-axes of displacement, was

attached to the participant's dominant hand and used to track the precise timings of the

in-air tapping for each of the five digits.

Resting state fMRI data was collected comprising 72 axial slices centred on the

fMRI acquisition stack with a TE/TR=25/1500 ms, SENSE 1.5, 1.5mm isotropic resolution,

using MB factor 4.

Structural data comprised a whole head T_1-weighted MPRAGE (1 mm isotropic

resolution) and PSIR (0.7 mm isotropic resolution), and high-resolution (0.5x0.5x1.5

mm^3) T$_2$*-weighted FLASH image, these were all collected with the same slice
prescription as the fMRI data.

2.3 MRI analysis

Following quality control (tSNR, movement <1mm), only four participants were used
going forward. fMRI data were distortion-corrected (FSL TOPUP), motion-corrected and
aligned to the FLASH image. Data were high-pass filtered with a 0.01 Hz cutoff (100 s) to
remove low-frequency trends. fMRI data were then analysed using custom software
within mrTools (36) implemented in MATLAB. We performed several analyses to study
the spatial representation of the digits and compare FHD patients to healthy controls, as
described in Sections 2.3.1-2.3.5.

For these analyses, tissue segmentation and cortical reconstruction of the T$_1$-
weighted volumes was performed using Freesurfer (http://surfer.nmr.mgh.harvard.edu/)
(37). Reconstructed cortical surfaces were flattened in a patch around the hand
representation in post-central gyrus (radius, 55mm to 75 mm, depending on anatomy),
using the mrFlatMesh algorithm (Vista software,
https://github.com/vistalab/vistasoft/tree/master/mrAnatomy/mrFlatMesh). Freesurfer
was used to estimate, for each participant, the location of primary and secondary
sensorimotor Brodmann areas (BAs 1, 2, 3a, 3b) based on the histological analysis (38).
The maximum-probability map for these was imported into participant-space to define
ROIs for each Brodmann area. Data were registered to the *fsaverage* template
(Freesurfer).

We also confirmed our digit representations from phase analysis of the TW data
against a General Linear Model (GLM) analysis to confirm correspondence between
analysis methods **(see Supplementary Methods).**

2.3.1 Somatotopy and motortopy phase analysis of TW data

Somatotopy and motortopy data were analyzed using standard, Fourier based
methods (in mrTools), producing coherence and phase maps. To account for shifts in the
fMRI response due to the haemodynamic lag, the forward and reverse scans were
averaged, as described in (7). Coherence, phase, and amplitude of the best-fitting
sinusoid were then computed as described in (39). The phase map was displayed on the

flattened cortical patch with a threshold of c>0.3 (coherence); this corresponds to an uncorrected p value of p<0.0004. First the digit correspondence between the somatosensory task data and the probabilistic Atlas were compared using Dice coefficients in fsaverage space for healthy volunteers and FHD patients for each visit. Then digit correspondence of the somatosensory and motor data was assessed using Dice coefficients between the HVs and FHD patients for each visit (Post-BTX and Pre-BTX), and between FHD Post-BTX and Pre-BTX. For each group, we also assessed Dice coefficients related to the extent of correspondence of motor and somatosensory maps in the Dominant hand. For the comparison between HV and FHD patients the Figure of Merit (FoM) was calculated from the Dice scores to summarise the response of each digit; this metric penalises degeneracy and poor representation (40).

$$\text{FoM}_j = r(m_j, X_i) - \frac{1}{n_{\text{digit}} - 1} \sum_k r(m_j, X_{k \neq i})$$

which defines the Dice (r) of the participant's digit ROI (m) to its paired ROI (X) subtracting the mean of the Dice from the other digit ROIs. A FoM score of 1 represents perfect overlap with the target digit, whilst a lower score suggests the digit overlaps with the non-target digit or the representation is degenerate (i.e. overlaps with many digits).

2.3.2 Comparison of data to the somatosensory probabilistic atlas

ROIs of each digit from the TW phase analysis were normalised to standard-space (*fsaverage*) and compared to the somatosensory probabilistic atlas (11). Surface projection and normalisation followed the methods described in O'Neill *et al*., i.e., using Freesurfer's *mri_vol2surf* to project coherence maps, phase maps, and digit ROIs into the participant's surface space, followed by a surface-to-surface transformation to MNI-305 space with *mri_surf2surf*. Since surface-based registration has previously been shown to outperform volume comparisons (11), only surface-based transformations were performed in this work. Full probabilistic maps (FPMs) were defined for each group (HC, and FHD for both Post-BTX and Pre-BTX), in which probabilities at each vertex were defined as the number of participants for which a particular digit was assigned to this voxel/vertex, divided by the total number of participants.

For the somatosensory data, central tendency (CT) was measured between the
FHD patients' digits and the atlas digits. CT defines how centrally located a digit is
relative to a probabilistic atlas digit (1=perfect overlap, >1=resides centrally, <1=resides
peripherally) (41). CT quantifies how much a single participant's digit ROI overlaps with
high probability areas of the atlas. For example, a small focal ROI that is within the atlas
ROI would give a high central tendency. A larger ROI that encompasses the atlas ROI
would also give a large central tendency if the two centres were aligned.

2.3.3 Population receptive field (pRF) mapping
This analysis followed the methods described in (21) for somatosensory and motortopy
tasks, which was adapted from a pRF method developed for visual cortex (42). Briefly, a
1-dimensional Gaussian model was combined with the spatio-temporal description of
the stimulus, producing a predicted "cortical" response. To account for haemodynamic
effects, this was then convolved with a HRF and then fit to the measured timeseries using
non-linear least squares optimisation. This method produces cortical maps of "between
digit" (equivalent to the TW Fourier analysis, albeit parameterized), pRF "size" (defined
from the standard deviation of the 1D Gaussian best fit for each voxel) and adjusted r^2
(the goodness of fit of the pRF model to the fMRI timeseries, pRF outputs were threshold
by adjusted $r^2 > 0$ (since adjusted r^2 can be negative). ROIs were calculated by
intersecting the binned digit phase value (D1-D5) with the FreeSurfer Brodmann areas
(3a,3b,1,2), creating 20 ROIs for each participant. pRF size was extracted from each ROI
and normalised (converted to z-scores) to account for inter-participant variability in
absolute pRF size.

2.3.4 Resting state connectivity
Data were analysed using the CONN toolbox in MATLAB (RRID:SCR_009550) (43)
(Whitfield-Gabrieli and Nieto-Castanon 2012). First, the default preprocessing pipeline
was used which included realignment, outlier detection, co-registration, normalisation
to the MNI template, and 5 mm spatial smoothing. Denoising was performed using
aCompCor (44), scrubbing, and temporal band pass filtering [0.008 0.09Hz]. Seed-based
functional connectivity (FC) measures to cortical brain regions were then assessed by
placing seed ROIs in both the left (L) and right (R) post-central gyrus taken from the

Harvard-Oxford atlas ("atlas.PostCG r" and "atlas.PostCG l"). FC was represented by
Fisher-transformed bivariate correlation coefficients from a weighted GLM, defined
separately for each pair of seed and target areas, modelling the association between
their BOLD signal timeseries. Cluster-level inferences were based on parametric
statistics from Gaussian Random Field Theory; results were threshold using a cluster-
forming $p<0.001$ voxel-level threshold, and familywise corrected p-FDR < 0.05 cluster-
size threshold. Group independent component analysis (groupICA) was performed. FC
was compared for 1) HC vs FHD (for both Post-BTX and Pre-BTX), and 2) FHD between
visits (Post-BTX vs Pre-BTX).

Data and code availability statement: Code to generate results and figures, and the
accompanying derived data is freely available on the Mendeley data repository:
https://data.mendeley.com/datasets/tvt7mg5bf3/1.

3. RESULTS

3.1 Behavioural measures: Spatial acuity using the "GOT" task was significantly reduced
in FHD for both Post-BTX and Pre-BTX groups compared with healthy controls (Mean HV
(std): 1.87 (0.19), Mean Post-BTX (std): 2.96 (0.93), Mean Pre-BTX (std): 3 (0.88); one-way
ANOVA $p<0.05$), **Figure 1A**.

In the healthy control repeatability group, within-participant CV for behavioural
measures were: "GOT": 6.2 ± 1.7 %; "TDT": 40 ± 12 % (R) 46 ± 11 % (L) (Mikkelson: 46%);
"AMP": 21 ± 6 % (R) 28 ± 8 % (L) (Mikkelson: 31%), as shown in **Figure 1B** and
Supplementary Table 1 which provides between-participant CVs. These CVs are in-line
with those reported by Mikkelson (35).

There were no significant differences in the "TDT" tasks or "AMP" task between the FHD
group and healthy controls, or for FHD patients between Post-BTX and Pre-BTX visits
(**Figure 1C&D**).

3.2 Comparing Travelling Wave digit maps between HV and FHD patients

Qualitatively, FHD travelling wave somatotopy and motor phase analysis digit maps

showed similar digit-specific activity patterns to HVs, with FHD patterns similar between

Post-BTX and Pre-BTX visits, **Figure 2**.

Quantitatively, Dice coefficient matrices of somatosensory phase analysis digit

correspondence assessed in *fsaverage* space between groups are shown in **Figure 3.**

When comparing matrices between HV and FHD groups for the somatosensory task

(**Figure 3Ai**) and motor task (**Figure 3Aii**), lower Dice coefficients were seen in the

dominant affected hand compared to the non-dominant hand for HV versus Post-BTX

and HV versus Pre-BTX (P<0.02). This is shown in **Figure 3Aiv**) by the lower FoM of the

Dice scores in the Dominant affected hand. However, no significant difference in the

Dice matrices was detected between Post-BTX and Pre-BTX conditions. **Figure 3Aiii** is a

graphic that explains how the DICE matrix affects the FoM score.

In response to the somatosensory task the dominant hand had larger digit ROIs than non-

dominant digit ROIs for FHD and HV groups (P<0.03), with D5 being significantly smaller

than all other digit (D1-D4) ROIs (P=0.004) (see **Supplementary Figure 1**).

Somatosensory and motortopy task dominant hand digit maps showed strong spatial

correspondence, with clear diagonal high Dice coefficients in each HV, FHD Post-BTX,

FHD Pre-BTX group, **Figure 4**. For all participants, there was good agreement in digit

correspondence of the TW GLM "winner-take-all" maps with the standard phase analysis

maps shown by the high diagonal Dice matrix coefficients (**Supplementary Figure 2**).

3.3 Comparing digit maps to the probabilistic atlas

The Dice of the TW phase analysis digit maps in response to the somatosensory task with

the Atlas digit ROIs showed a trend for a stronger diagonal in the HV compared to FHD

group, particularly for the Dominant (affected) hand, **Figure 5A** (mean Dice diagonal: HV

L/R: 0.634 ±0.009/0.62±0.02; Post-BTX L/R: 0.61 ±0.02/0.60±0.01; Pre-BTX L/R: 0.58

±0.02/0.58±0.01). This is also reflected in the central tendency (CT) matrices (**Figure 5B**)

which also considers how centrally located a digit is. A within group paired t-test between

Dominant and Non-dominant hand CT scores, showed that for both FHD visits (Post-BTX

and Pre-BTX) the Dominant hand had significantly smaller CT values than the non-Dominant hand (p<0.05, Bonferroni corrected).

3.4 Population Receptive Field (pRF) mapping

Mean (absolute) pRF size (across hemispheres and digits) across groups was HV (1.7±1.8), HV Atlas (1.4±1.2), FHD Post-BTX (2.0±1.9), FHD Pre-BTX (1.7±1.9), these results are in-line with our previous study in HVs (1.9±1.6 2D Gaussian model, mean±stdev) (21).

Comparing group mean pRF sizes for the somatosensory task, the FHD Pre-BTX group had larger pRF sizes in the Dominant (affected) hand than the FHD Post-BTX, HV, and Atlas. There were no differences in pRF size between groups in the Non-dominant hand for the somatosensory task, or in the motor pRF data for the Dominant hand (**Figure 6**). There were no differences in pRF sizes from the motortopy data in the Dominant hand (the only hand on which the motor task was performed). There was no overall significant difference in pRF size between hands across groups. Motor pRF size was larger than somatosensory pRF size for both HV and FHD groups (**Supplementary Figure 3).**

Table 1 and **Figure 7** show the pRF size ordering across digits in each group for non-dominant and dominant hands for the somatosensory task and dominant hand for the motor task. While HV and FHD Post-BTX showed greater pRF sizes for D2,3,4,5 than D1 this was not seen for FHD Pre-BTX in the dominant hand. In the non-dominant hand, all groups had similar pRF size trends, e.g. D2,3,4 > D1. **Table 1** also shows pRF results separated by Brodmann areas, with BA2 larger than BA3a and 3b in all groups in the dominant hand, however, in the non-dominant hand, only Post-BTX and Atlas groups show BA2 > BA3b.

3.5 Resting state fMRI

There was a significant (p<0.001, uncorrected) difference in seed-based connectivity between the seed in left postcentral gyrus (contralateral to dominant hand) and projection to right postcentral gyrus, with lower connectivity in FHD, **Figure 8**. There was no difference in FC in FHD between Post-BTX and Pre-BTX visits.

4 DISCUSSION

This work has assessed behavioural and functional measures of digit localisation and population receptive field (pRF) size in Focal Hand Dystonia (FHD) patients compared to healthy volunteers, and the effect of treatment with Botulinum Toxin (BTX) using high spatial resolution fMRI.

4.1 Behavioural temporal and spatial thresholds in FHD

The spatial acuity threshold measured using the grating orientation task (GOT) was larger in FHD patients compared to healthy controls, supporting the hypothesis of reduced spatial acuity in FHD (28). In contrast, temporal-based temporal discrimination task (TDT) measures did not show any significant differences between groups, this may be due to large variability in the 'TDT' measure which was consistent with previous literature (35).(Mikkelsen et al., 2020)

4.2 Dominant (affected) and Non-Dominant (unaffected) hand digit maps in FHD

Digit maps on the flattened cortical surface showed clear digit-specific activity patterns for both the travelling wave (TW) phase-analysis and general linear model (GLM) winner-takes-all maps in healthy volunteers (HV) and FHD patients. These findings contest early work showing widespread digit disruption in FHD (2,45) but supports more recent work that have shown intact digit representation within primary sensorimotor cortex of individuals with musician's dystonia (15). There were however significantly lower Dice coefficients in the dominant affected hand of the FHD group compared to the HV group, with the HV group better matching the probabilistic atlas; this points to a degree of disorganisation or increased variability in the digit representations in the affected hand likely due to abnormal sensorimotor integration (28). It should be noted that early functional magnetic resonance imaging (fMRI) studies (2) were performed at coarser spatial resolution and assessed digit maps in un-flattened space based on Euclidean distances between digits, so differences may have arisen due to folding structure or limitations of spatial resolution or altered amplitude of responses in dystonia patients previously.

It was also shown that the dominant hand had larger digit regions of interest (ROIs) than
non-dominant hand for both FHD and HV groups. Digit 5 (D5) was significantly smaller
than all other ROIs, agreeing with literature on this digit having the least representation
in terms of size, for an overview of cortical magnification of digits, see (46).
For all groups there was large overlap in the digit maps in postcentral gyrus for the
somatosensory and motor tasks as evidenced by Dice coefficients. This overlap reflects
the anatomical and functional relationship of primary S1 and M1, which form an
integrated sensorimotor representation of the digits (47).
470
4.3 Comparison of central tendency scores of digit maps to the probabilistic atlas
The central tendency matrices showed good comparison to the atlas for HV and FHD
patients. However, in FHD patients the dominant hand had a significantly lower central
tendency than the non-dominant hand. This suggests that in FHD, the dominant hand
has larger less centrally matched digit representations whilst the non-dominant hand
has smaller more centrally represented digits.
It is noted that the differences between groups may be due to atypical HVs, however,
when comparing groups to the Atlas, the HVs perform equal to or better than FHD
patients (**Figure 5B).**
480
4.3 Population receptive field sizes in FHD
For the dominant hand, pRF sizes were significantly larger for the FHD Pre-BTX group
compared to the HV group and the FHD Post-BTX group. Smaller pRF sizes indicate better
tactile acuity, while larger pRFs indicate reduced sensitivity to spatial frequency. This
suggests that BTX treatment reduces pRF size to a similar level to HVs, but this increases
again ~3 months later. This finding is consistent with lower Dice scores in the dominant
hand, identifying subtle disruption in the dominant affected hand at the BTX visit.
pRF analysis showed that in all groups and hands, the thumb had the smallest
pRF size (D2, D3, D4, D5 > D1). This ordering is consistent with electrophysiological and
primate literature, whereby D1 has the highest receptor density, smallest receptive fields
compared to the other digit tips and the largest cortical magnification (48,49).
Previous fMRI pRF studies have also shown D5 to be larger than the other digits.
Interestingly, we showed in some cases that the pRF size of D3 and D4 was greater than

D5, where other studies have found D5 to have the largest pRF density size. This may be
due to fewer good fits of the pRF fitting algorithm at D5 (46) as less volume is dedicated
to D5 in the somatosensory hand map (50).
When comparing pRF size between the dominant and non-dominant hands there was no
significant difference (**Supplementary Figure 3)**.
Our finding of no significant difference between dominant and non-dominant hands in
digit representations and pRF size in our healthy participants agrees with previous work
which showed no significant differences in volume, activation strength, and Euclidean
distance between hands (51), or tactile acuity (52).
Motor pRF sizes, measured across both post-central and pre-central gyrus in
response to the motor task, were larger than somatosensory pRF size. This is consistent
with previous literature which showed larger pRF sizes in solely precentral gyrus (53), and
electrophysiologically where complex motor movements have been reported (54).
507
4.5 Resting state functional connectivity in FHD:
Resting state data identified lower functional connectivity (FC) in FHD as compared to
the HV group, but no differences between FHD Post-BTX and Pre-BTX groups. This is
consistent with previous literature, which shows lower FC in sensorimotor regions as
compared to controls at rest (22,55) in more participants (n=16, 15, respectively), and
the lower sample size here can explain the fewer differences between groups seen here.
514
515   *4.6 Limitations*
516   This study had a relatively small sample size for FHD patients but suggests alterations in
517   the dominant (affected) hand. Future studies should build on this work with a larger study
518   in FHD patients. In future it would also be of interest to compare motor task responses
519   in both the Dominant and non-Dominant hand.
520
In conclusion, spatial acuity was raised in FHD patients compared to healthy controls.
The FHD group had some alterations in their Dominant affected hand with both lower
Dice scores and larger pRF sizes between Post-BTX and Pre-BTX. These results provide
some evidence to support a neuronal change in the cortical representation of the

affected hand in focal hand dystonia and that BTX treatment in FHD patients can alter
pRF size in the dominant affected hand.

Data availability statement

The raw data supporting the conclusions of this article will be made available by the
authors on request.

Ethics statement

The study was conducted with the approval of the University of Nottingham Faculty of
Medicine and Health Sciences Ethics Committee for healthy volunteers and of the Health
Research Authority Research Ethics Committee (REC 17/EM/0368) for FHD patients. All
participants gave written informed consent. The studies were conducted in accordance
with the local legislation and institutional requirements. The participants provided their
written informed consent to participate in this study.

Author contributions

MA: study design, data collection, data analysis, manuscript draft. RS: study design, data
collection, data analysis. PG: data collection, manuscript editor. DP: data collection,
manuscript editor. GON: data collection, manuscript editor. DS: data analysis,
manuscript editor. MH: data collection, manuscript editor. SF: study design, data
collection, data analysis, manuscript draft. All authors contributed to the article and
approved the submitted version.

Funding

This work was supported by the Medical Research Council [grant number
MR/M022722/1].

Non-standard abbreviations

AMP – Amplitude Thresholding Task
BOLD – Blood Oxygenation Level Dependent contrast
BTX – Botulinum toxin

CT – Central Tendency

FC – Functional Connectivity

FHD – Focal Hand Dystonia

fMRI – Functional magnetic resonance imaging

FoM – Figure of Merit

GOT – Grating Orientation Task

Post-BTX – Focal Hand Dystonia patients ~30 days after Botulinum toxin treatment

Pre-BTX – Focal Hand Dystonia patients ~100 days after Botulinum toxin treatment

pRF – Population Receptive Field

rs-fMRI – Resting State fMRI

TDT – Temporal Discrimination Task

Acknowledgements

We thank the focal hand dystonia participants and healthy volunteers who took part in

the study.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial

or financial relationships that could be construed as a potential conflict of interest.

Generative AI statement

The author(s) declare that no Generative AI was used in the creation of this manuscript.

Contribution to the Field Statement – not for in draft to upload on website on submission

Focal hand dystonia is a neurological condition that affects the digits and can be treated with botulinum toxin (BoNT-A) injections to relieve muscle spasm. This treatment is symptomatic, and it is unclear how it affects brain areas responsible for digit movement and sensation.

We used high-resolution functional magnetic resonance imaging to assess whether digit representations differ between healthy people and individuals with focal hand dystonia. Patients were scanned at two timepoints: (1) shortly after BoNT-A treatment, when its effects were strongest, and (2) three months later, when the effects had worn off.

Patients' digit maps showed the typical digit ordering overall and compared well to a normative probabilistic atlas. However, we detected subtle differences in FHD patients: (1) The dominant affected hand maps in individuals with FHD showed more spatial disorder in digit to healthy controls. (1) Sensory receptive sizes in the dominant affected hand were smaller shortly after BoNT-A treatment, then enlarged again after three months, suggesting the treatment may temporarily alter brain organization.

These findings add to our understanding of how digits are represented in hand dystonia, and how they respond to treatment.

604 **Tables**

606 **Table 1:** Summary of significant differences in pRF size by digit and Brodmann area.

	pRF by digits			pRF by Brodmann area		
	Somato Non-Dom	**Somato Dom**	**Motor**	**Somato Non-Dom**	**Somato Dom**	**Motor**
HV	D2,3,4,5>D1 D4>D3	D2,3,4,5>D1	D5>D1,2,3,4	No difference	BA2>BA3a&3b	BA6>BA3a,3b&1
FHD Post-BTX	D2,3,4,5>D1 D4>D2 D3,4 >D5	D2,3,4,5>D1	D3,4>D1	BA2>BA3b	BA2>BA3a	BA2>BA3a BA4a>BA3a
FHD Pre-BTX	D2,3,4>D1. D4>D2. D3, 4 > D5	No difference	D2,3,4,5>D1	No difference	BA2>BA3a&3b	No difference
Atlas	D2,3,4>D1 D2,3,4>D5	D2,3,4>D1 D3, 4>D5	Not acquired	BA2>BA3a&3b	BA2>BA3a&3b	Not acquired

Figures

Figure 1: Behavioural results in healthy volunteer and focal hand dystonia Post-BTX and Pre-BTX visits for A) Spatial acuity grating orientation task ("GOT") applied to the tip of the dominant (affected) hand. Participants who only performed the behavioural tasks and did not proceed to do the fMRI session are colour coded. B) Within-participant coefficient of variation (CV) computed from two visits in 10 healthy controls. CVs are comparable to Mikkelsen et al. 2020 (35); C) Temporal discrimination task ("TDT") using piezoelectric stimulators and Brain Gauge (BG), and D) Amplitude discrimination ("AMP") at 31 and 200 Hz.

Figure 2: A) Somatosensory and motor travelling wave (TW) tasks performed during fMRI. An accelerometer glove was worn during the motor task to quality control participant movements. B) Example TW phase analysis digit maps to the Somatosensory and Motor task for an FHD participant at Post-BTX and Pre-BTX visits. Brodmann areas projected from the FreeSurfer segmentation also shown. C) Non-dominant somatosensory task TW map from Post-BTX visit projected to 'fsaverage' space. D) Example accelerometer trace from the forward run, post-BTX, dominant hand of the participant shown in B&C) for the motor task.

Figure 3: Ai) Dice matrices of digit patterns between study groups (HV versus FHD Post-BTX visit, HV versus FHD Pre-BTX visit and FHD BTX versus NoBTX) for (i) somatosensory task for the Non-dominant (left, non-affected) and Dominant (right, affected) hand and (ii) motor task for the Dominant hand. Aiii) Graphic to outline how key patterns and how the Figure of Merit (FoM) relates to the DICE matrices. Aiv) Figure of Merit (FoM) computed for each DICE matrix in (Ai) and (Aii). Note the FoM tends to be lower for the dominant hand. Significant difference (p<0.05, corrected for multiple comparisons) in FoM shown by brackets above plot.

Figure 4: DICE matrices of digit patterns between motor and somatosensory maps for the Dominant (right, affected) hand for each study group. Strong correspondence is seen for the HVs and FHD patients at both visits.

Figure 5: A) Dice matrices of digit patterns for the somatosensory task between HV, FHD groups compared with the probabilistic atlas. B) Matrices of central tendency (CT) scores for HV and FHD Post-BTX and Pre-BTX visits. The CT score represents how centrally located the digits are to the probabilistic atlas (1=perfect overlap, >1=resides centrally, <1=resides peripherally). Note no significant difference in CT matrix between Non-dominant and dominant hand for HV vs Atlas, while there was a significant difference (P<0.001) in CT matrix between Non-dominant and dominant hand for FHD vs Atlas for both Post-BTX and Pre-BTX visits.

Figure 6: Population receptive field size across all digits, split into the study groups (HV, FHD Post-BTX, and FHD Pre-BTX) as well as the Atlas HVs. A) non-dominant (left) hand and B) dominant (right) hand for the somatosensory task; C) dominant (right) hand for the motor task. * indicates a significant difference ($p < 0.05$), and the height of the bracket indicates which group had the larger pRF size.

Figure 7: Box plots of population receptive field size, with z-scores shown for each digit (x-axis) and coloured by Brodmann area. Each row is a different group (HV, FHD Post-BTX, FHD Pre-BTX, Atlas HVs). Red text on each plot indicates significant differences between digits. Left column is non-dominant (left) hand and middle column is dominant (right) hand for the somatosensory task; and right column is dominant (right) hand for the motor task.

Figure 8: A) Resting state seed-based functional connectivity analysis. Seeds placed in post-central gyrus, and the projection maps for each group are shown. B) CONN cortical ROIs based on Harvard Oxford parcellation are shown. The green crosshair marks the postcentral gyrus. C) Significant difference between groups for a seed in the left (contralateral to dominant hand) postcentral gyrus projecting to right postcentral gyrus. This indicates reduced FC in FHD.

Supplementary Figure 1

ROIs are shown (RD1 and LD1) for each group. Voxel counts are shown for each group (HV, BTX, NoBTX), for left) non-dominant and right) dominant hands. Each data point corresponds to a single participant's digit ROI in fsaverage space.The dominant hand was larger than the non-dominant hand across groups. D5 was significantly smaller in all groups.

Ai) Winner-take-all digit maps

Aii) DICE between GLM and phase analysis digit maps

738 **Supplementary Figure 2**

739 Figure A) GLM analysis of travelling wave fMRI data for digit localisation. i) "winner-take-

740 all" maps shown for an example participant from each group. ii) DICE coefficient

741 matrices showing excellent correspondence between the travelling wave GLM and

742 phase analysis ('PH') for digit localisation.

759 **Supplementary Figure 3**

760 pRF size separated by hand dominance, for each group. Motor pRFs across groups were

761 significantly larger than somatosensory pRFs.

Supplementary Figure 4

Travelling wave and population receptive field flat maps for a single example participant from each group 1) post-Botox; 2) pre-Botox; 3) healthy volunteers; 4) atlas healthy volunteer.

References

1. The Dystonia Society. Focal Hand Dystonia [Internet]. 2014 [cited 2008 Jul 20]. Available from: http://www.dystonia.org.uk/index.php/professional-research/types-of-dystonia/focal-hand-dystonia

2. Butterworth S, Francis S, Kelly E, McGlone F, Bowtell R, Sawle G V. Abnormal cortical sensory activation in dystonia: An fMRI study. Movement Disorders. 2003;18(6):673–82.

3. Nelson AJ, Blake DT, Chen R. Digit-specific aberrations in the primary somatosensory cortex in writer's cramp. Ann Neurol. 2009;66(2):146–54.

4. Martuzzi R, van der Zwaag W, Farthouat J, Gruetter R, Blanke O. Human finger somatotopy in areas 3b, 1, and 2: A 7T fMRI study using a natural stimulus. Hum Brain Mapp [Internet]. 2014;35(1):213–26. Available from: http://www.ncbi.nlm.nih.gov/pubmed/22965769

5. Schweizer R, Voit D, Frahm J. Finger representations in human primary somatosensory cortex as revealed by high-resolution functional MRI of tactile stimulation. Neuroimage. 2008;42(1):28–35.

6. Sanchez-Panchuelo RM, Francis S, Bowtell R, Schluppeck D. Mapping Human Somatosensory Cortex in Individual Subjects With 7T Functional MRI. J Neurophysiol [Internet]. 2010;103(5):2544–56. Available from: http://jn.physiology.org/cgi/doi/10.1152/jn.01017.2009

7. Besle J, Sanchez-Panchuelo RM, Bowtell R, Francis S, Schluppeck D. Single-subject fMRI mapping at 7 T of the representation of fingertips in S1: a comparison of event-related and phase-encoding designs. J Neurophysiol [Internet]. 2013;109(9):2293–305. Available from: http://jn.physiology.org/cgi/doi/10.1152/jn.00499.2012

8. Schluppeck D, Sanchez-Panchuelo RM, Francis ST. Exploring structure and function of sensory cortex with 7T MRI. Neuroimage [Internet]. 2016;(October 2016):1–8. Available from: http://dx.doi.org/10.1016/j.neuroimage.2017.01.081

9. van der Zwaag W, Schafer A, Marques JP, Turner R, Trampel R. Recent applications of UHF-MRI in the study of human brain function and structure: a review. NMR Biomed. 2016;29(9):1274–88.

10. Sanchez-Panchuelo RM, Besle J, Mougin O, Gowland P, Bowtell R, Schluppeck D, et al. Regional structural differences across functionally parcellated

Brodmann areas of human primary somatosensory cortex. Neuroimage [Internet]. 2014;93:221–30. Available from: http://linkinghub.elsevier.com/retrieve/pii/S1053811913002942

11. O'Neill GC, Sengupta A, Asghar M, Barratt EL, Besle J, Schluppeck D, et al. A probabilistic atlas of finger dominance in the primary somatosensory cortex. Neuroimage [Internet]. 2020;217(August 2019):116880. Available from: https://doi.org/10.1016/j.neuroimage.2020.116880

12. Kadota H, Nakajima Y, Miyazaki M, Sekiguchi H, Kohno Y, Amako M, et al. An fMRI study of musicians with focal dystonia during tapping tasks. J Neurol. 2010 Jul 9;257(7):1092–8.

13. Kimberley TJ, Pickett KA. Differential activation in the primary motor cortex during individual digit movement in focal hand dystonia vs. healthy. Restor Neurol Neurosci. 2012 Jun 1;30(3):247–54.

14. Jankowski J, Paus S, Scheef L, Bewersdorff M, Schild HH, Klockgether T, et al. Abnormal Movement Preparation in Task-Specific Focal Hand Dystonia. PLoS One. 2013 Oct 22;8(10):e78234.

15. Sadnicka A, Wiestler T, Butler K, Altenmüller E, Edwards MJ, Ejaz N, et al. Intact finger representation within primary sensorimotor cortex of musician's dystonia. Brain. 2023 Apr 1;146(4):1511–22.

16. Huber L, Kassavetis P, Gulban OF, Hallett M, Horovitz SG. Laminar VASO fMRI in focal hand dystonia patients. Dystonia. 2023 Feb 1;2.

17. Tucholka A, Fritsch V, Poline JB, Thirion B. An empirical comparison of surface-based and volume-based group studies in neuroimaging. Neuroimage. 2012 Nov;63(3):1443–53.

18. Schellekens W, Petridou N, Ramsey NF. Detailed somatotopy in primary motor and somatosensory cortex revealed by Gaussian population receptive fields. Neuroimage [Internet]. 2018;179(May):337–47. Available from: https://doi.org/10.1016/j.neuroimage.2018.06.062

19. Puckett AM, Bollmann S, Junday K, Barth M, Cunnington R. Bayesian population receptive field modeling in human somatosensory cortex. Neuroimage [Internet]. 2020;208(March 2019):116465. Available from: https://doi.org/10.1016/j.neuroimage.2019.116465

20. Wang L, Zhang Z, Okada T, Li C, Chen D, Funahashi S, et al. Population Receptive Field Characteristics in the between- And Within-Digit Dimensions of

the Undominant Hand in the Primary Somatosensory Cortex. Cerebral Cortex.
2021;31(10):4427–38.

21.   Asghar M, Sanchez-Panchuelo R, Schluppeck D, Francis S. Two-Dimensional
Population Receptive Field Mapping of Human Primary Somatosensory Cortex.
Brain Topogr. 2023 Nov 1;36(6):816–34.

22.   Dresel C, Li Y, Wilzeck V, Castrop F, Zimmer C, Haslinger B. Multiple changes
of functional connectivity between sensorimotor areas in focal hand dystonia. J
Neurol Neurosurg Psychiatry [Internet]. 2014;85(11):1245–52. Available from:
http://jnnp.bmj.com/cgi/doi/10.1136/jnnp-2013-307127

23.   Pan P, Wei S, Ou Y, Jiang W, Li W, Lei Y, et al. Reduced Global-Brain
Functional Connectivity and Its Relationship With Symptomatic Severity in
Cervical Dystonia. Front Neurol. 2020 Jan 10;10.

24.   Bianchi S, Fuertinger S, Huddleston H, Frucht SJ, Simonyan K. Functional and
structural neural bases of task specificity in isolated focal dystonia. Movement
Disorders. 2019 Apr 6;34(4):555–63.

25.   DeSimone JC, Archer DB, Vaillancourt DE, Wagle Shukla A. Network-level
connectivity is a critical feature distinguishing dystonic tremor and essential
tremor. Brain. 2019 Jun 1;142(6):1644–59.

26.   Giannì C, Pasqua G, Ferrazzano G, Tommasin S, De Bartolo MI, Petsas N, et
al. Focal Dystonia. Neurology. 2022 Apr 5;98(14).

27.   Molloy FM, Carr TD, Zeuner KE, Dambrosia JM, Hallett M. Abnormalities of
spatial discrimination in focal and generalized dystonia. Brain.
2003;126(10):2175–82.

28.   Sanger TD, Tarsy D, Pascual-Leone A. Abnormalities of spatial and temporal
sensory discrimination in writer's cramp. Movement Disorders. 2001;16(1):94–
9.

29.   Nevrlý M, Hluštík P, Hok P, Otruba P, Tüdös Z, Kaňovský P. Changes in
sensorimotor network activation after botulinum toxin type A injections in
patients with cervical dystonia: a functional MRI study. Exp Brain Res. 2018 Oct
1;236(10):2627–37.

30.   Cohen LG, Hallett M, Geller BD, Hochberg F. Treatment of focal dystonias of
the hand with botulinum toxin injections. J Neurol Neurosurg Psychiatry
[Internet].   1989;52(3):355–63.   Available   from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1032411&tool=pmc
entrez&rendertype=abstract

31. Tommerdahl M. corticalmetrics.com. 2017 [cited 2017 Jul 17]. Brain Gauge.
Available from: https://www.corticalmetrics.com/braingaugeplus

32. Gardner JL, Merriam EP, Larsson J, Schluppeck D. MGL: Visual psychophysics
stimuli and experimental design package. 2018 Jun 28 [cited 2025 Sep
5];(Version 2.0). Available from: https://zenodo.org/records/1299497

33. Levitt H. Transformed Up-Down Methods in Psychoacoustics. J Acoust Soc Am
[Internet]. 1971;49(2B):467–77. Available from:
http://asa.scitation.org/doi/10.1121/1.1912375

34. Craig JC. Grating orientation as a measure of tactile spatial acuity. Somatosens
Mot Res. 1999;16(3):197–206.

35. Mikkelsen M, He J, Tommerdahl M, Edden RAE, Mostofsky SH, Puts NAJ.
Reproducibility of flutter-range vibrotactile detection and discrimination
thresholds. Sci Rep. 2020 Dec 1;10(1).

36. Gardner JL, Merriam EP, Schluppeck D, Besle J, Heeger DJ. mrTools: Analysis
and visualization package for functional magnetic resonance imaging data. 2018
Jun 28 [cited 2019 Sep 3]; Available from: https://zenodo.org/record/1299483

37. Fischl B, Sereno MI, Dale AM. Cortical Surface-Based Analysis. Neuroimage
[Internet]. 1999;9(2):195–207. Available from:
http://linkinghub.elsevier.com/retrieve/pii/S1053811998903962

38. Fischl B. FreeSurfer. Neuroimage. 2012;62(2):774–81.

39. Besle J, Sanchez-Panchuelo RM, Bowtell R, Francis S, Schluppeck D. Event-
related fMRI at 7T reveals overlapping cortical representations for adjacent
fingertips in S1 of individual subjects. Hum Brain Mapp. 2014;35(5):2027–43.

40. O'Neill GC, Tewarie PK, Colclough GL, Gascoyne LE, Hunt BAE, Morris PG, et
al. Measurement of dynamic task related functional networks using MEG.
Neuroimage. 2017;146:667–78.

41. Eickhoff SB, Paus T, Caspers S, Grosbras MH, Evans AC, Zilles K, et al.
Assignment of functional activations to probabilistic cytoarchitectonic areas
revisited. Neuroimage [Internet]. 2007;36(3):511–21. Available from:
http://dx.doi.org/10.1016/j.neuroimage.2007.03.060

42. Dumoulin SO, Wandell BA. Population receptive field estimates in human visual
cortex. Neuroimage. 2008;39(2):647–60.

43. Whitfield-Gabrieli S, Nieto-Castanon A. Conn: A Functional Connectivity
Toolbox for Correlated and Anticorrelated Brain Networks. Brain Connect. 2012
Jun 1;2(3):125–41.

44. Behzadi Y, Restom K, Liau J, Liu TT. A component based noise correction
method (CompCor) for BOLD and perfusion based fMRI. Neuroimage. 2007 Aug
1;37(1):90–101.

45. Elbert T, Candia V, Altenmüller E, Rau H, Sterr A, Rockstroh B, et al. Alteration
of digital representations in somatosensory cortex in focal hand dystonia.
Neuroreport. 1998;9(16):3571–5.

46. Janko D, Thoenes K, Park D, Willoughby WR, Horton M, Bolding M.
Somatotopic Mapping of the Fingers in the Somatosensory Cortex Using
Functional Magnetic Resonance Imaging: A Review of Literature. Vol. 16,
Frontiers in Neuroanatomy. Frontiers Media S.A.; 2022.

47. Ejaz N, Hamada M, Diedrichsen J. Hand use predicts the structure of
representations in sensorimotor cortex. Nat Neurosci [Internet].
2015;18(7):1034–40.   Available   from:
http://www.nature.com/doifinder/10.1038/nn.4038

48. Merzenich MM, Kaas JH, Wall J, Nelson RJ, Sur M, Felleman D. Topographic
reorganization of somatosensory cortical areas 3b and 1 in adult monkeys
following restricted deafferentation. Neuroscience. 1983;8(1):33–55.

49. Sutherling WW, Levesque MF, Baumgartner C. Cortical sensory representation
of the human hand. Neurology. 1992 May;42(5):1020–1020.

50. Overduin SA, Servos P. Distributed digit somatotopy in primary somatosensory
cortex. Neuroimage [Internet]. 2004;23(2):462–72. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/15488396

51. Schweisfurth MA, Frahm J, Farina D, Schweizer R. Comparison of fMRI digit
representations of the dominant and non-dominant hand in the human primary
somatosensory cortex. Front Hum Neurosci. 2018 Dec 10;12.

52. Sathian K, Zangaladze A. Tactile spatial acuity at the human fingertip and lip:
Bilateral symmetry and interdigit variability. Neurology. 1996 May 1;46(5):1464–
1464.

53. Schellekens W, Petridou N, Ramsey NF. Detailed somatotopy in primary motor
and somatosensory cortex revealed by Gaussian population receptive fields.

Neuroimage [Internet]. 2018;179(May):337–47. Available from:
https://doi.org/10.1016/j.neuroimage.2018.06.062

54.   Brown AR, Teskey GC. Motor Cortex is Functionally Organized as a Set of
Spatially Distinct Representations for Complex Movements. Journal of
Neuroscience. 2014 Oct 8;34(41):13574–85.

55.   Mohammadi B, Kollewe K, Samii A, Beckmann CF, Dengler R, Münte TF.
Changes in resting-state brain networks in writer's cramp. Hum Brain Mapp.
2012 Apr;33(4):840–8.