Auxiliary data, quality assurance and quality control for wearable light loggers and optical radiation dosimeters

Johannes Zauner [1, 2] [0000-0003-2171-4566] *#

Oliver Stefani[3] [0000-0003-0199-6500] #

Gianfranco Bocanegra[4] [0009-0004-5280-4657]

Carolina Guidolin[2] [0009-0007-4959-2667]

Björn Schrader[3] [0000-0002-8387-922X]

Ljiljana Udovicic[5] [0000-0002-6035-8309]

Manuel Spitschan[1, 2, 6, 7,] [0000-0002-8572-9268] *

TUM School of Medicine and Health, Department Health and Sports Sciences, Chronobiology & Health, Technical University of Munich, Munich, Germany

Max Planck Institute for Biological Cybernetics, Max Planck Research Group Translational Sensory & Circadian Neuroscience, Tübingen, Germany

Lucerne University of Applied Sciences and Arts, Lucerne, Switzerland

The Hague University of Applied Sciences

Federal Institute for Occupational Safety and Health (BAuA), Dortmund, Germany

TUM Institute for Advanced Study (TUM-IAS), Technical University of Munich, Garching, Germany

TUMCREATE Ltd., Singapore, Singapore

* Corresponding author: johannes.zauner@tum.de , manuel.spitschan@tum.de

#JZ and OS are shared first-authors

Authors other than JZ, OS, and MS are in alphabetical order

Abstract

Wearable light loggers and optical radiation dosimeters are increasingly used in chronobiology and circadian health research, yet their data often lack contextual information (e.g., sleep, activity, environmental conditions) and may be compromised by non-wear periods, compliance issues, or technical faults. To address these limitations, we conducted expert interviews (n=21) and a survey (n=16) to distil and iteratively develop auxiliary data and quality-control strategies that enhance the accuracy and interpretability of wearable light measurements. From this process, we established a six-domain auxiliary data framework encompassing wear/non-wear logging, sleep monitoring, light-source context, participant behaviour, user experience, and environmental light levels. Survey responses showed strong consensus on the value of auxiliary information (mean importance 4.0/5), with sleep and wear-time tracking rated as the most essential additions. To support practical adoption, we provide implementation tools, including extensions to the open-source R package LightLogR, enabling streamlined integration of wearable and auxiliary data as well as systematic quality assurance and control. Experts agreed that combining contextual records with rigorous QA/QC procedures substantially improves the reliability of field-collected light-exposure data. These recommendations and tools aim to help researchers in chronobiology, wearable sensing, and health sciences maximise data quality and enhance interpretation in real-world light-exposure studies.

Introduction

Light affects humans beyond the visual aspect through autonomous pathways originating in the eye [1,2]. These so-called non-visual or non-image-forming (NIF) effects of ocular light play a significant role in regulating circadian rhythms, such as the 24-hour sleep-wake cycle, and directly or indirectly affect neurological, endocrine, metabolic, and immune systems [3]. Ideally, ocular light exposure should follow the natural light-dark cycle and its spectral and spatial properties. However, people in modern societies are often exposed to too little light during the day due to predominantly indoor activities and insufficient outdoor time. They are also exposed to too much light at night, for example, during night shift work. [4-7] To best support physiology, sleep, and wakefulness in healthy adults, recommendations for daytime, evening, and nighttime indoor light exposure have been published [8]. However, little is known about human light exposure under naturalistic conditions, and a growing but fragmented landscape of researchers is taking on the task to fill this gap [9].

In order to assess the negative impact of modern living conditions on well-being, performance, and health, studies are required that measure personal light exposure and link these measurements to health-related outcomes. By now, studies have shown relationships of personal light exposure patterns with personal well-being and pathological outcomes, such as obesity and type II diabetes, major depression, anxiety, seasonal depressive disorder, and other pathologies related to metabolic and mental health [10-19].

The effects of light on human health and well-being have historically been studied in laboratory settings, examining mechanisms and dose- or phase-response relationships [20], and from studies over the past two decades, many principled relationships and mechanisms have been established in connecting ocular light exposure to health and health-related outcomes [8,20]. While these highly controlled studies remain the gold standard for establishing causal and mechanistic relationships, the relevance and magnitude of real-life effects are best studied through personal light exposure data in field studies, measured through wearable devices [21,22]. For example, laboratory studies revealing mechanistic effects of evening display light on human physiology (e.g., melanopic irradiance dependent effects on sleep latency, melatonin suppression, and alertness)[23,24] could be modulated by prior light history, arguing for careful assessment and reporting of pre-

exposure conditions [25]. By contrast, real-world outcomes are shaped by
heterogeneous daylight exposure, content/arousal, etc, which can attenuate or
outweigh isolated light-mechanism effects. Recent field-oriented reviews therefore
down-weight pure "brightness/arousal" accounts and emphasize usage patterns
when explaining sleep impacts [26].

When combined with physiological measurements, wearable data contribute to
understanding how laboratory findings can be translated and applied to everyday
environments [4,11,15].

Emerging evidence suggests that an increasing number of research groups are
integrating wearable light loggers into their protocols, ranging from small but
comprehensively recorded samples [4-6,10,16,27-31] to hundreds or even tens of
thousands of participants in large cohort studies [11,13,14,32-35]. Wearable designs have
also found their way into laboratory studies. As prior light history may influence the
outcome of laboratory experiments investigating non-visual light effects, sleep, or
chronobiology in general, it is recommended to monitor and report light exposure
using wearable light loggers even before the actual experiment [21,36]. The studies
employ wearable devices to measure or estimate the ocular exposure to light and
other spectral ranges of optical radiation. The devices are most often attached to a
participant's wrist, at chest level, or, most optimally, positioned at eye level [21,37,38].
The output of these devices is commonly a time series of photopic illuminance, but
depending on the device, other quantities are stored as well [39]. These can include
alpha-opic quantities like melanopic EDI, activity, temperature, direct sensor outputs,
or even (reconstructed) spectral power distributions [40,41].

Furthermore, light recordings from wearables on their own – even if they are of high
fidelity from a measurement perspective – often lack essential information beyond
light about the environment and the participant, e.g., non-wear time, physical activity,
or climate-based information [40]. This means that additional measures are regularly
required to contextualise and categorise the light and optical radiation data, which
can be used either as covariates in the analysis or as quality indicators in data
preprocessing, such as to detect non-wear [42]. Beyond collecting new data, essential
steps must be taken to ensure data quality throughout a study. Data from wearables
can easily be compromised, for example, by clothing covering the device or if a
device's internal clock is not properly synchronised with local time. These and other

confounding aspects can be mitigated through various strategies used during the

study design, data collection, and data analysis, which significantly affect the utility

and actual accuracy of the data [31,39,42]. Strategies include factors such as how

participants are briefed and motivated, the device placement, and the choice of

analysis software.

In this study, we identified factors that can influence data quality from wearable

devices used to measure light exposure and optical radiation. We also gathered and

developed strategies to improve the utility of the collected data. The study is part of

the MeLiDos project (Metrology for wearable light loggers and optical radiation

dosimeters), dedicated to providing tools, standards, best practice guidelines, and

FAIR (findable, accessible, interoperable, and reusable) data for the collection and

analysis of wearables in research studies [9]. In a series of interviews performed by a

MeLiDos subtask group with researchers familiar with the use of wearable light

loggers, the experts expressed doubts about wearer compliance with the devices and

the reliability of the recorded data [43].

This article focuses on the most important measures that can help to augment

wearable data, beyond obvious technical aspects (e.g., measurement accuracy). We

prioritized these because technical aspects are thoroughly treated in prior reviews [41],

while these complementary factors emerged as under-addressed yet high-impact

areas for improving data quality. The measures are divided into auxiliary data

(extending the data beyond the wearable itself), quality assurance measures

(increasing the likelihood of good compliance), and quality control (reducing the

number of faulty data points). The auxiliary measures were surveyed and refined with

researchers from across the field. The mitigating strategies were derived from

discussions of common issues experienced by consortium members of the MeLiDos

project and expert researcher interviews [43]. This publication summarises the

outcomes and provides guidance for auxiliary data and mitigation strategies to

maximise the utility when using wearable devices in a research study.

Results

Auxiliary data

Definition: Auxiliary data in the context of this study is defined as time-dependent
(i.e. time-stamped) data relevant to the analysis of light logging data but not
automatically collected through wearable light logger devices, as visualised in
Figure 1.

149   --- Figure 1 to be inserted here ---

150 Two simple examples are sleep-wake logs and non-wear diaries. **Figure 2** shows a
151 visualised example of how auxiliary data extends the utility of personal light exposure
152 data from a wearable.

153   --- Figure 2 to be inserted here ---

Scope and outcome format: All instruments in this section consist of questionnaire
items and/or a set of instructions for participants. Applying these in a study result in a
list of state changes connected to timestamps and participants. A simple example is
shown in Table 1. The tracked states are:

- wear / non-wear

- sleep / wake

- light environment

- activity

Some states may be tracked already through the wearable light logger or other
devices participants use. These would not be considered auxiliary data as per the
definition above. In that case, the researcher must decide whether the redundant
tracking through the auxiliary data strategy adds enough benefit during analysis for a
particular state.

167   --- Table 1 to be inserted here ---

The tracked state changes can be converted to timespans of states, which relate to
measurements of light and augment their usage during analysis. The MeLiDos
funded software package LightLogR [39] features functions specifically to importing,
handling, and merging auxiliary data to data from wearable light loggers in an easy
and robust fashion. This includes an optional setting for upper thresholds to the
length a state can sensibly persist, like 24 hours for a single wake period. The
package was recently updated to version 0.9.2 ("Sunrise") with functionality
specifically targeted at auxiliary data and states.

Wear log

Objective: Track wear and non-wear times throughout the day. To gain significant

confidence, data from multiple measures should be combined.

Suggested measures:

- A digital logbook with several options, as laid out in the procedure

- A black, light-sealed bag or container to put the wearable light logger into
during non-wear time while awake (note: auxiliary data would be times of zero-
lux measurements during wake times) unless light measurement outside wear
time is needed for context information

- An event trigger/button on the wearable light loggers

Possible procedure: Throughout the day, participants are instructed to report their

wear and non-wear time in a digital logbook, e.g. through a smartphone. Specifically,

they have five choices of wear log entry:

- 1 = "Taking the light glasses off"

- 2 = "Putting the light glasses on"

- 3 = "Taking the light glasses off before sleep and placing them on a nightstand
or flat surface"

- 4 = "Leaving <study location > and its surroundings (e.g. 60 km radius)"

- 5 = "Re-entering < study location > and its surroundings (e.g. 60 km radius)".

In case of 1, they should put their device into the black bag or container provided by

the investigator. If an event button on the light logger exists and is not utilised in

another way, for options 1 to 3, participants may also press the button to signal an

event occurring, and in the case of 1, they are asked to confirm whether they place

the light glasses in a black bag provided to them and if they are in movement.

Options 4 and 5 are introduced to control for potential differences in personal light

exposure due to environmental availability rather than behaviour. They might be

dropped if they are too specific for a study. For all five wear log entry choices,

participants must state whether they are logging a present or a past event. That way,

a lapse in procedure can be corrected by the participant, and the investigator can

decide whether and under which circumstances to accept past events.

To increase compliance, a researcher might consider allowing certain non-wear

periods without requiring logging (e.g., one minute).

Utility: Tracking non-wear time allows for the invalidation of light exposure data time

periods that do not capture personal light exposure.

Further information: The data repository [44] contains one possible implementation

for the wear log as it was used for a data collection effort in Tübingen, Germany.

Potentially, a sixth option could be added: "Putting the light glasses on upon waking

up".

Morning sleep log

Objective: Collect sleep and wake times for each day, as well as some additional
sleep-related information

Suggested measures:

-   A digital logbook with several options, as laid out in the procedure

Possible procedure: Every morning after waking up, participants fill in the core
Consensus Sleep Diary [45] consisting of 9 items to assess their sleep timing, sleep
duration during the night and their subjective sleep quality. This last item is scored on
a five-point scale (1 = "Very poor" to 5 = "Very good").

A researcher might decide to only include sleep/wake timing to minimise the burden
on participants. This already allows the utility as described below.

Utility: Collecting sleep and wake times allows for metric calculation based on sleep-
wake episodes instead of the 24-hour day, which can be more meaningful for certain
metrics. It further allows for an assessment of how close a participant's light
exposure is to the Brown, et al. [8] recommendations based on daytime, pre-sleep, and
sleep times.

Further information: The data repository [44] contains one possible implementation of
the digital logbook, which was used for a data collection effort in Tübingen, Germany.

The morning sleep log will have to be adjusted (and renamed) for shift workers,
where participants can be queried after each sleep period. Similarly, for certain
research questions, capturing naptimes might be important. In this case, the log
could be adjusted.

Some wearable devices can be worn during sleep, e.g. by using wristbands. These
devices may track sleep-wake times automatically through actigraphy. However, they
increase the error relating to the measure of interest, i.e., corneal exposure.

Light exposure diary

Objective: Collect types of light environments that a participant occupied throughout

the day.

Suggested measures:

- An analogue form as laid out in the procedure

- A digital logbook to confirm compliance as laid out in the procedure

- Alternatively, a digital log might be used throughout the day for an ecological
momentary assessment. However, this can affect the lighting environment in a
significant way during night times (i.e, in a dark environment, there would
always be partly "smartphone" or "laptop" light)

Possible procedure: Every evening, participants must fill in a modified version of the

Harvard Light Exposure Assessment (H-LEA [46]). This is referred to as "mH-LEA" and

is done on paper using a form provided by the experimenter during the in-person visit

at the start of the study. Participants are asked to report, for each hour of the day, the

main light source they are exposed to. The main light source is described as "the

main light source in their environment". They can choose between 8 light categories:

- L = "Electric light source indoors (e.g.: lamps such as LEDs)"

- S = "Electric light source outdoors (e.g.: street lights)"

- I = "Daylight indoors (through windows)"

- O = "Daylight outdoors (including being in the shade)"

- E = "Emissive displays (e.g.: smartphone, laptop etc.)"

- D = "Darkness (outdoors and/or indoors)"

- W = "Light entering from outside during sleep (e.g., daylight, streetlights)").

If they believe they are exposed to a significant combination of lights within the same

hour, they can choose from the following combinations:

- "L+I"

- "L+E"

- "I+E"

- "S+O"

- "D+W"

To ensure that participants complete this task, they send a picture of the completed

form every night and upload it to a shared folder (separate for each participant)

where the experimenter can check compliance. Furthermore, they are asked to rate

their confidence in their answers ("How sure are you about the light exposure and

activity categories you chose?"), where they can answer using a 5-point scale

ranging from 1 = "Not confident at all" to 5 = "Completely confident".

Alternatively, a digital form could be used to capture the mH-LEA directly. However,

as mentioned above, this can affect the lighting environment in a significant way

during night times (i.e, in a dark environment, there would always be partly

"smartphone" or "laptop" light)

Utility: Allows for connecting subjective assessments of environmental light

conditions to objective light exposure.

Further information: *Supplementary information 1 (C)* shows one possible

implementation for the digital logbook and the analogue form, as it was used for a

data collection effort in Tübingen, Germany.

The diary might be split up into two sections, each covering 12 hours instead of 24.

This can reduce recollection errors by the participants.

Behaviour/Exercise diary

Objective: Collect types of exercise throughout the day.

Suggested measures:

- An analogue form as laid out in the procedure

- A digital logbook to confirm compliance as laid out in the procedure

- Alternatively, a digital log might be used throughout the day for an ecological
momentary assessment.

Possible procedure: Every evening, participants are required to complete an
experience diary form. This is done on paper using a form provided by the
experimenter during the in-person visit at the start of the study. Participants are
asked to report, for each hour of the day, the activity they performed in that hour.
With regards to their activity, they could choose between 8 categories

- 1 = "Sleeping in bed"

- 2 = "Awake at home"

- 3 = "On the road with public transport/car"

- 4 = "On the road with bike/on foot"

- 5 = "Working in the office/from home"

- 6 = "Working outdoors (including lunch break outdoors)

- 7 = "Free time outdoors (e.g. garden/park etc.)

- 8 = "Other: please specify (e.g. sport)"

To ensure that participants complete this task, they send a picture of the completed
form every night and upload it to a shared folder (separate for each participant)
where the experimenter can check compliance. Furthermore, they are asked to rate
their confidence in their answers ("How sure are you about the light exposure and
activity categories you chose?"), where they can answer using a 5-point scale
ranging from 1 = "Not confident at all" to 5 = "Completely confident".

Alternatively, a digital form could be used to directly capture the exercise diary.

Utility: Allows the connection of different types of activity to objective light exposure.

Further information: The data repository [44] contains one possible implementation
for the digital logbook and the analogue form, as it was used for a data collection
effort in Tübingen, Germany.

Experience log

Objective: Collect positive and negative experiences associated with the wearable

Suggested measures:

- A digital logbook to confirm compliance as laid out in the procedure

Possible procedure: At any time, participants can note down positive and negative

experiences associated with using the wearable in a digital logbook. The questions

cover a wide range from what kind of experience, where it occurred and who was

involved, to whether the experience resulted in any behavioural change.

Utility: It gives insights into a possible behavioural impact of the wearable on the

participant, which might affect "regular, everyday" behaviour.

Further information: The data repository [44] contains one possible implementation

for the digital logbook as it was used for a data collection effort in Tübingen,

Germany.

Environmental light levels

Objective: Collect objective comparative light levels that reflect the natural lighting

conditions in the study location of a participant.

Suggested measures:

- A calibrated light logger of the same type and model as used by the
participant, mounted in a fixed, unobstructed position, such as a high-up
rooftop. This is the measure as laid out in the procedure.
- Estimates of ground-level illuminance based on, e.g., satellite data or weather
and time data. Unobstructed outdoor light levels can be derived from data
provided by meteorological institutes, which typically measure global
irradiance and allow for relatively straightforward conversion into illuminance
values.

Possible procedure: The rooftop-mounted light logger collects light data during the

same period as a participant. If the sensor is not rated for outdoor use, the sensor

should have a clear cover, e.g., a plastic or glass dome (see **Figure 3**). We

recommend a horizontal mount because the sensor can only generally reflect the

environmental exposure potential.

354   --- Figure 3 to be inserted here ---

Utility: Gives insights into the relative light exposure a participant got compared to

what would be available under natural conditions in this area. While it cannot

precisely re-cap the environmental light levels at a participant's exact position, it can

do so generally.

Further information: *LightLogR* [39] has functions to connect and compare reference

light levels to personal light exposure data from participants, regardless of whether

they were collected with the same measurement epoch or equal time points.

For additional information, weather and ambient temperature data can provide vital

context to certain behaviours affecting personal light exposure.

Auxiliary data survey

Sixteen researchers participated in a survey collecting feedback on auxiliary data

across 28 questions. They had the option to skip questions. Thus, the number of

respondents (N) is provided for each question.

Q1, agreement with the definition of auxiliary data (N=15): All participants agreed

with the definition of auxiliary data as described, with one participant adding that the

utility of the experience questionnaire is not clear with regards to how it relates to the

main light logger data.

It is of note that the definition of auxiliary data was adjusted slightly to cover data that

are not recorded **automatically** by a device. Otherwise, event triggers or zero-lux

readings enforced through the black bag would not be covered.

Q2, previous experience with auxiliary data (N=16): Thirteen participants collected

auxiliary data before, and three had not. Of those that have previous experience, two

collected data paper-based, six collected digitally, and five used a hybrid method of

both. Participants could also provide additional information about their collection

methods, and REDCap was mentioned three times, Qualtrics and MyCAP once

each.

Q3, most effective collection method (N=16): All researchers agree that data

should be collected digitally (n=8) or in a hybrid form (n=8). Votes for the hybrid form

additionally mentioned that qualitative information or open-ended questions may be

better collected via paper or interview, that participants should be given a choice

about the modality, and that retrospective questionnaires about time points in the

past day can profit from the paper form by providing a simple visualisation while filling

the form in.

Q4, the importance of auxiliary data (N=16): Overall, the importance of auxiliary

data is rated 4.0 on a 1 (isn't required or helpful) to 5 (essential) scale. No participant

indicated the lowest two scores (1 or 2), three gave it a middle rating (3), 10

participants selected the second highest option (4), and three participants believed

light exposure measurements would be worthless without auxiliary data (5).

Q5, relevance of auxiliary data (N=15): Ranking the six categories of light exposure

reveal that sleep/wake is the most relevant category for participants (average rank

5.0), followed by wear/non-wear (4.85), the light environment (4.13), a tie between

behaviour/exercise and environmental light levels (3.07), and finally experience with
the light logger (1.54).

Q6, other domains of auxiliary data (N=10): participants note other useful data
would be on

- Clothing and accessories (e.g., hats), especially glasses/sunglasses

- Reasons for not wearing a device (author note: this has since been included in
the wear log)

- A more detailed light environment (e.g., curtains)

- For intervention studies, feedback on expectations and acceptability of the
intervention, symptoms, etc.

- Distinguishing work day and free day (author note: this could be extracted
from the exercise diary)

- Weather conditions and temperature (author note: this aspect was added to
the environmental light levels section)

- One researcher noted that the sleep diary might not work for shift workers
(author note: this caveat was added to the morning sleep log section)

Wear/Non-wear (Q7 N=16; Q8 N=16; Q9 N=5): The domain is rated as important
(3.6 on a scale of 1, not required or helpful, to 5, essential). Researchers believe the
wear log captures that domain of auxiliary data sufficiently well (4.1 on a 1 to 5 scale,
where 1 means it does not cover the domain at all, 2: not well, 3: partly, 4:
sufficiently, 5: completely).

- Some researchers mentioned doubts as the log is quite detailed, thus
increasing participant burden and potentially reducing compliance.

- One researcher suggested another option for "putting the light glasses on
upon waking up" (author note: this was added as a potential addition to the
wear log).

- Three researchers suggest that some way of automated non-wear detection
would be much preferred. (author note: this has been added as an option in
the wear log section)

Sleep/Wake (Q10 N=16; Q11 N=16; Q12 N=2): The domain is rated as important
(4.0 on a scale of 1, not required or helpful, to 5, essential). Researchers believe the

morning sleep log captures that domain of auxiliary data sufficiently well (4.0 on a 1
to 5 scale, where 1 means it does not cover the domain at all, 2: not well, 3: partly, 4:
sufficiently, 5: completely).

- Some researchers note that automated detection through actigraphy can be
preferable.

- One researcher suggests extending it to include nap times (author note: this
option has been added to the morning sleep log section).

Light exposure diary (Q13 N=16; Q14 N=16; Q15 N=4): The importance of the
domain is rated as average (3.4 on a scale of 1, not required or helpful, to 5,
essential). Researchers believe the light exposure diary captures that domain of
auxiliary data sufficiently well (3.9 on a 1 to 5 scale, where 1 means it does not cover
the domain at all, 2: not well, 3: partly, 4: sufficiently, 5: completely).

- Three researchers noted the high participant burden for this log,

- One researcher believed participants might not be able to assess different
light contributions correctly.

- One researcher believes the log should be split in two, each covering 12 hours
instead of 24 hours (author note: this option has been added to the light
exposure diary section). Further, the utility might be highly dependent on the
research question.

Behaviour / Exercise log (Q16 N=16; Q17 N=16; Q18 N=3): The importance of the
domain is rated as average (3.2 on a scale of 1, not required or helpful, to 5,
essential). Researchers believe the exercise log captures that domain of auxiliary
data sufficiently well (4.1 on a 1 to 5 scale, where 1 means it does not cover the
domain at all, 2: not well, 3: partly, 4: sufficiently, 5: completely).

- As with the previous question, there is some doubt about the utility of these
data compared to the added participant burden.

Experience log (Q19 N=16; Q20 N=16; Q21 N=3): The importance of the domain is
rated as average (3.0 on a scale of 1, not required or helpful, to 5, essential).
Researchers believe the experience log captures that domain of auxiliary data
sufficiently well (3.8 on a 1 to 5 scale, where 1 means it does not cover the domain at
all, 2: not well, 3: partly, 4: sufficiently, 5: completely).

- While researchers overall agree that it is a useful questionnaire, some note
that it might be too complex and could be phrased more generally and with
open-ended comments.

- One researcher notes that standards such as SUS and WEAR questionnaires
are only partly suitable for usability testing of light loggers.

- Another researcher believes this log is only helpful in limited validation studies,
and the nature of the log would make it hard to condense the data.

Environmental light (Q22 N=15; Q23 N=15; Q24 N=2): The domain is rated as
important (3.7 on a scale of 1, not required or helpful, to 5, essential). Researchers
believe the experience log captures that domain of auxiliary data sufficiently well (3.9
on a 1 to 5 scale, where 1 means it does not cover the domain at all, 2: not well, 3:
partly, 4: sufficiently, 5: completely).

- Two researchers note that global illuminance levels do not precisely reflect
environmental light levels,

- One researcher suggests using satellite data as an easier proxy.

- One researcher voices concerns about feasibility, while another believes it is a
nice and easy addition.

- Another suggestion is to add information on the mounting (author note:
satellite data and mounting information were added to the environmental light
section).

Q25 Impact on own research (N=15): Two-thirds of the researchers (n=10) believe
that the auxiliary data approach "[…] will definitely have some impact on [their]
research design". One-third (n=5) believe "it might have some impact […]". None
believe "it will have no impact […]".

Q26 general feedback (N=5): Researchers note that the utility of the different
domains depends heavily on the context, study, population, and other factors, and
that other methods to add or get around these types of auxiliary data could be
sought.

Q27, Q28 (N=15 each): Thirteen researchers would like to be mentioned as
contributing to the auxiliary data questionnaire in reports and publications. Those
who have not joined the group of authors have been added to the acknowledgements

section. Two would like to remain anonymous. Fourteen researchers would be willing
to be contacted again regarding auxiliary data, and one would not.

Non-technical strategies for improving light data integrity

As outlined before, the quality of light exposure data collected by wearable light
dosimeters can be significantly compromised by improper compliance and misuse,
thus reducing accuracy to reflect the corneal exposure of the wearer. This highlights
the importance of providing clear instructions, involving participants in training
sessions and incorporating feedback mechanisms to improve compliance with study
protocols. Addressing these factors is critical to obtaining accurate and reliable light
exposure data, which can ultimately affect study results and outcomes. The following
aspects have been collected from internal rounds of discussions and in-depth
interviews with researchers who have used wearable devices in their studies. Table 2
provides a succinct overview of the topics, challenges, and possible strategies.

--- Table 2 to be inserted here ---

Non-compliance and misuse

• Inconsistent use
If participants do not wear the dosimeters consistently - whether due to
forgetfulness, discomfort or lack of understanding - the data collected will be
incomplete. This can lead to gaps in exposure data, making it difficult to draw
accurate conclusions about light exposure patterns. Participants may forget to
wear the dosimeter during critical exposure periods, resulting in data that does
not accurately reflect their actual light exposure.

• Incorrect placement
Improper placement of dosimeters can lead to inaccurate estimations of ocular
light exposure. For instance, when worn on the wrist instead of near the eye
(such as on spectacle frames), the device may primarily capture ambient
rather than corneal illuminance, resulting in misleading data. Obstruction by
clothing or hair can further impair the sensor's ability to detect light accurately.
This can lead to underestimation of exposure levels, particularly for light that
would naturally reach the eyes.

• Equipment misuse
Participants and even experimenters may not understand how to properly
handle or calibrate the device. For example, if the dosimeter is switched off or

not activated correctly, it won't record any data, leading to data loss.

Misleading feedback about the recording state could also lead to data loss. If

experimenters do not follow the manufacturer's instructions on how to use the

device (e.g., how to adjust settings or interpret readings), they may misuse it,

resulting in erroneous data collection.

•   Lack of training and education

Without sufficient training on the importance of proper use and compliance,

participants may not realise how their actions affect data quality. This could

lead to careless behaviour, such as not following instructions on how to wear

the dosimeter or ignoring reminders to wear it consistently. If participants do

not receive real-time feedback on whether they are using the dosimeter

correctly, they may continue to make the same mistakes without realizing it.

For example, a dosimeter that doesn't provide acute feedback on incorrect

positioning may lead to prolonged periods of incorrect use.

•   Environmental and social factors

Participants may feel self-conscious about wearing a conspicuous dosimeter

in public, leading to avoidance of use in social situations. This may bias the

data as the dosimeter may not be worn during important exposure periods.

Participants may remove the dosimeter during certain activities (such as

exercise or work), not realising that it's important to collect data during these

times. This results in incomplete datasets that do not represent true exposure

levels.

544 Strategies to improve the quality of data collection

•   Ease of wear

The placement of the dosimeter may be uncomfortable or inconvenient,

especially when worn on glasses, which may cause participants to avoid using

it or to wear it incorrectly. Devices that are small, lightweight and comfortable

significantly increase compliance by minimising disruption to the wearer.

Wearers are more likely to follow instructions if they forget they are wearing

the device. Making devices waterproof or able to be worn in different positions

without compromising accuracy further increases wearer comfort and

compliance. Devices with long battery life reduce the burden on participants

by limiting the frequency of charging, which can lead to data loss. Avoiding

555   frequent charging or providing devices with rechargeable options that last for

556   weeks ensures continuous data collection.

557   •   Minimise behavioural disturbance

558   Devices should avoid being visually intrusive or obvious, particularly around

559   the eyes, which may cause wearers to behave unnaturally due to concerns

560   about how they will be perceived by others. Visible or bulky devices could alter

561   behaviour, thereby compromising the validity of the study. Discreet, near-eye

562   level devices are preferred, but should not resemble cameras to avoid being

563   mistaken for recording devices.

564   •   Intrinsic and extrinsic motivation

565   In some instances, participants may not fully grasp the importance of the study

566   or lack intrinsic motivation to follow the instructions. If they do not understand

567   how crucial accurate data is – and how they can contribute toward that end –

568   they may not make the effort to use the dosimeter correctly. Therefore,

569   educating participants about the scientific importance of the study can foster a

570   sense of purpose, leading to higher intrinsic motivation and increased

571   compliance. Offering financial rewards, personal feedback reports, or other

572   incentives can provide external motivation. Wearers are more likely to follow

573   instructions if they perceive a tangible benefit, such as receiving data on their

574   own light exposure.

575   •   Device misuse

576   Participants may not have a clear understanding of the correct procedures for

577   wearing and using the dosimeter, resulting in incorrect data collection;

578   therefore, participants should receive thorough training on how to use the

579   equipment, with regular check-ins or home visits by supervisors to ensure

580   compliance. Reminders and instructions should be simple, emphasising the

581   importance of the research and the correct use of the equipment to reinforce

582   participant commitment. To prevent participants from changing their behaviour

583   based on light exposure feedback, any real-time data, such as battery life or

584   exposure levels, should only be visible to researchers. Providing this feedback

585   only to the researcher avoids biasing the results while still ensuring the

586   functionality of the device.

588 Strategies to improve the quality of data processing: Exclusion and calibration
589 (quality control)

• Transparency of calibration
A consistent concern voiced by researchers in the interviews is the lack of
transparency in how instruments are calibrated by manufacturers.
Standardised calibration procedures, including inter-device comparisons and
position-specific validations (e.g. wrist vs. chest), are essential to ensure
reliable measurements. Improving the transparency of the calibration process
will increase researchers' confidence and improve data quality, especially in
low-light or extreme conditions.

• Exclusion criteria
Researchers often exclude data taken under certain conditions, such as below
20 lux, due to concerns about accuracy in low light or overly bright
environments (e.g. saturated levels). Establishing clear exclusion criteria
based on well-understood device limitations ensures that only reliable data is
used in the analysis.

• Data cleaning and interpretation
Depending on the device, participant compliance, and other factors, it is
commonly necessary to invalidate some of the measurements recorded by a
device. Reasons include aspects such as:
o out of range measurements (beyond sensor saturation and at the noise
level)
o non-wear
o device occlusion
o recordings outside relevant time frames (trimming)
Often, these invalid data are randomly distributed, but due to non-wear at
times of day at which specific activities occur (e.g., showering, contact sports,
swimming, …), light exposure data may be missing systematically.
The following sanity checks are recommended:
o Visualisation of raw time series for the available sensors
o Visualisation of histograms to identify distributions of data and outliers
Combining device data with participant-reported activities (e.g. diaries, apps –
see the section on auxiliary data above) is a common strategy to contextualise
light exposure, but such methods are considered unreliable by some due to

participant forgetfulness. Automated data collection or digital wearable time

recording devices coupled with motion sensors are recommended for greater

accuracy.

There is a need for standardisation of data cleaning methods (e.g.

identification of non-wear times, processing of light data by body position).

Standard protocols will ensure that data from different studies are comparable,

thus improving the overall quality of research results.

•   Data compression and connectivity

Large datasets can lead to interruptions in downloads, creating a risk of data

loss. Software developers should prioritise efficient data compression

techniques without compromising quality. Reliable connectivity (Bluetooth,

cable or infrared) should be emphasised to avoid transmission problems

during data extraction.

635 Discussion

The utility of personal light exposure data heavily depends on key decisions and

strategies before, during, and after data collection. In this study, we present essential

aspects, both to maximise the quality of light exposure data themselves, as well as

for auxiliary data to the time series.

The auxiliary data strategy in this publication covers six domains: wear and non-wear

of the devices, sleep, light sources, behaviour, experience, and environmental light

levels. Sixteen experts in the field were surveyed towards this strategy. All

researchers agree on the definition of auxiliary data, and most of them have previous

experience collecting it. Furthermore, all surveyed participants agree that auxiliary

data are important, with a third believing them "essential". Of the different modalities

of auxiliary data, sleep and wear data are considered to be the most important, but

none are considered irrelevant. Importantly, the presented solutions were (on

average) considered as sufficient to cover each domain, and all researchers believe

the strategy will or might have some impact on their future research. This suggests a

robust basis to collect auxiliary data in these domains.

Besides these encouraging results, many answers caution that some domains or the

whole approach would heavily increase participant burden, a concern we share. The

benefits of additional data have to be weighed against potentially reduced participant
compliance, errors, or, depending on the study goal, questionable utility of a given
domain. Thus, it has to be emphasised that we do not recommend every future study
to include the auxiliary data strategy in full and as laid out here. Rather, we want to
make researchers aware of domains of auxiliary data that have been of importance in
the past and suggest a possible implementation to capture these data. During study
design, each domain and measure has to be weighed against the additional burden
for participants, the study population's willingness and ability to comply, and
alternative ways to collect or add the data. For example, the experience log can be a
good tool to assess different wearing positions in a pilot study, but might be dropped
from a larger-scale collection effort. Ideally, as devices and analysis tools become
more sophisticated, the need for some of these domains will decrease, e.g., when
wear times are assessed and recorded by the devices themselves. Furthermore,
contextual or environmental data may become easier to collect, as interfaces to
public services such as meteorological institutes become more interoperable and
accessible.

Collection of non-wear data was rated as important by researchers participating in
our survey. While self-reported methods (e.g., wear logs) were generally considered
effective, some researchers emphasised the need for automated detection. A recent
study by Guidolin et al. (2025)[42] evaluated a low-illuminance cluster–based algorithm
for detecting non-wear periods and found good agreement with self-reported wear
logs (ground truth), provided that participants stored the device in a light-blocking bag
during non-wear time. Importantly, their analysis showed that accounting for non-
wear periods affected only half of the light exposure metrics investigated (7 of 14),
and even among those, most differences were small. Significant effects were
primarily observed when comparing uncorrected (raw) data to corrected datasets,
rather than between correction methods. For instance, average differences between
raw and wear log–corrected data for key duration metrics (TAT250, TAT1000) were
minimal—approximately 2 and 1 minutes in total, respectively. These results suggest
that the influence of non-wear time on derived metrics is generally modest but can
become relevant in studies focusing on specific light exposure periods (e.g.,
evenings). In the absence of a standardised non-wear detection framework,
researchers should therefore align their data collection and preprocessing approach
with the specific aims and sensitivity requirements of their study. To improve the

quality and usability of light exposure data, future studies should adopt participant-

centred strategies, including optimised device placement, clear instructions, real-time

feedback, a discreet and weather-resistant design, motivational incentives and robust

data quality control, to balance measurement accuracy with compliance and comfort.

There are four resources we want to highlight in that regard:

-    The open-source software package LightLogR by Zauner, et al. [39] provides

standardised routines to ingest data from many wearable devices, detect and

handle gaps and irregularities, merge data from different sources (such as

wear or sleep/wake), visualise, and perform metric calculations. Using these

standard analysis pipelines not only reduces time spent preparing the data,

but also reduces errors and differences when implementing routines on a by-

workgroup basis. Compared to the version at the time of the referenced paper,

a recent update (0.9.2 Sunrise) [47] has added many functions related to states,

which are at the core of auxiliary data.

-    A recent survey by Zauner, et al. [48] with over 150 participants from around the

world explored wearer preferences regarding eight different wearing positions

for wearable devices. Surveyed contexts include work, home, social, and

sport, which have slight but significant effects on user preferences. This

survey can be one resource to guide decisions on device type and placement

based on study duration and contexts.

-    A research guide by the Research data alliance (RDA) working group on

optical radiation exposure and visual experience data (in active development,

https://rda-wg-visualexperiencedata.github.io/ResearcherGuide/) is a rich

resource for many technical and non-technical data quality aspects.

-    A device specification tool, developed as another part of the MeLiDos project

contains many aspects relating to data quality and wearability (https://tscnlab-

wearable-devices-specification.share.connect.posit.cloud) [49]. The tool

combines these aspects into a specification sheet that facilitates device

selection and procurement.

-    The project *A day in daylight* [50], as part of the *Daylight Awareness Week 2025*

combines light exposure data from researchers around the globe (47

participants), collected on the day of the solar equinox with a rich event log

(over 1700 log entries). These log entries relate to or directly encapsulate

many auxiliary data concepts. While this project is not a traditional research
study, and the sample is biased, the conditional light exposure tables still
highlight the relevance and effect of auxiliary data. Part of that information is
contained in **Figure 2**, but many more contexts are available in the accessible
project dashboard (https://tscnlab.github.io/2025_ADayInDaylight/).

A measure not discussed in this study, but still noteworthy, is that random acts of
non-compliance or misuse become less relevant for light exposure metrics as the
data collection duration increases. In a bootstrap analysis of a month-long collection
period per person, Biller, et al. [6] found that even 6 hours of missing data per day did
not significantly affect light exposure metrics across the whole period, with shorter
collection periods obviously more sensitive to missingness than longer ones. While
missing data is not equal to measurement changes due to non-compliance or
misuse, the trend towards less effect over longer periods could still hold.

There are noteworthy limitations to our study. As field studies with wearable light
loggers and optical radiation dosimeters are still an emerging field of research, the
suggestions and measures provided above must be considered optional rather than
required. They reflect best practices available at the time of publication and need to
be considered in the context of a given study protocol. Depending on a study's goal
or scope, only some suggested measures might be useful or sensible. Participant
burden is also an aspect to be considered carefully.

Further, the measures are not infallible, and, in many cases, their benefit cannot be
quantified. This limitation is mostly grounded in how the measures were collected: as
experience-based solutions to problems researchers encountered in their work. As
such, they must be considered a byproduct of research rather than the actual topic,
which would be required for a quantitative assessment of their benefit.

Methods

Data collection

Data, i.e. strategies and questionnaires, were collected in multiple rounds of

discussion among the participating MeLiDos consortium members in 2024.

In addition, in-depth interviews were conducted with 21 researchers experienced in

the use of light loggers (invited: 101). While the full results of these interviews will be

reported separately [43], this paper extracts and expands upon the key findings

specifically related to strategies for augmenting data quality.

Furthermore, a survey was conducted in the third quarter of 2024 for the auxiliary

data. The survey was hosted on SurveyMonkey and was sent out to 106

researchers. Invited researchers were experts known to MeLiDos consortium

members, the CIE joint technical committee JTC20, and corresponding authors from

papers cited in a recent review about metrics used in wearable light logger studies by

Hartmeyer and Andersen [51]. Sixteen (n=16) researchers participated in the survey,

providing additional insights and suggestions, which were consequently implemented

into the results. The survey was accompanied by a preliminary version of the

auxiliary data presented in this study. The 28 survey questions ranged from general

questions about the nature and utility of auxiliary data to questions specifically

targeted at individual aspects of auxiliary data in the preliminary document, such as

sleep or non-wear. The survey results can be found in *Supplementary*

information S1.

In general, both the survey and interview participants (i.e., experts and researchers)

can be considered a form of convenience sample. However, given the emerging yet

relatively small size of the research field, this sample represents a substantial

proportion of the relevant expert community, considering that the invited group

stemmed from a comprehensive review of the field with a size of only about 100

researchers.

Data analysis

774 Collected data consisted of long-form interviews, questionnaires, and (mostly) free-

775 form survey replies. Data were condensed through manual, topical analysis, and

776 expert feedback was manually implemented.

Declaration statements

Data availability

All relevant data are part of this publication, either contained in the manuscript or its supplements.

Code availability

The auxiliary data questionnaires (PDF) and REDCap questionnaires for the *Morning sleep diary*, the *Wear log*, the *Light Exposure diary*, and the *Experience log* are available from the GitHub repository (https://github.com/tscnlab/ZaunerEtAl_npjBiolTimingSleep_2025), archived on Zenodo [44].

Acknowledgments

We would like to thank the following researchers (in alphabetical order) and three anonymous researchers that participated in the auxiliary data survey and contributed to the auxiliary data strategy as it is presented here: Tianren Chen, Débora B. Constantino, Altug Didikoglu, Fatemeh Fazlali, Gena Glickman, Steffen Hartmeyer, Rafael Lazar, Renske Lok, Elise McGlashan, Karin Smolders, and Jamie Zeitzer

The authors were financially supported by MeLiDos. The project (22NRM05 MeLiDos) has received funding from the European Partnership on Metrology, co-financed by the European Union's Horizon Europe Research and Innovation Programme and by the Participating States.
On the use of generative AI and AI-assisted technologies in the writing process: The authors used ChatGPT during the preparation of this work. After using this tool, the authors reviewed and edited the content as needed and take full responsibility for the content of the publication.

The following use of AI by contributor roles is based on the CRediT taxonomy. Funding acquisition, project administration, resources, and supervision were deemed irrelevant in this context and thus removed:

Conceptualization: no

Data curation: no

Investigation: no

Formal analysis: no

Methodology: no

Software: no

Validation: no

Visualization: no

Writing – original draft: abstract refinement

Writing – review & editing: improve readability and language

Authors' contributions

Conceptualisation: J.Z., O.S., C.G.

Data Curation: J.Z., O.S., C.G.

Formal Analysis: -

Funding Acquisition: M.S., B.S.

Investigation: J.Z., O.S., G.B.

Methodology: J.Z., O.S., C.G.

Project Administration: M.S., B.S., J.Z., O.S

Resources: -

Software: -

Supervision: M.S.

Validation: -

Visualisation: J.Z.

Writing – Original Draft Preparation: J.Z., O.S.

Writing – Review & Editing: All authors

Competing interests

JZ, OS, and MS are guest editors in the npj special issue "Light exposure in the real

world".

JZ declares the following potential conflict of interest in the past five years (2021-

2025). Funding: Received research funding from Reality Labs Research.

MS declares the following potential conflicts of interest in the past five years (2021–2025). Academic roles: Member of the Board of Directors, Society of Light, Rhythms, and Circadian Health (SLRCH); Chair of Joint Technical Committee 20 (JTC20) of the International Commission on Illumination (CIE); Member of the Daylight Academy; Chair of Research Data Alliance Working Group Optical Radiation and Visual Experience Data. Remunerated roles: Speaker of the Steering Committee of the Daylight Academy; Ad-hoc reviewer for the Health and Digital Executive Agency of the European Commission; Ad-hoc reviewer for the Swedish Research Council; Associate Editor for LEUKOS, journal of the Illuminating Engineering Society; Examiner, University of Manchester; Examiner, Flinders University; Examiner, University of Southern Norway. Funding: Received research funding and support from the Max Planck Society, Max Planck Foundation, Max Planck Innovation, Technical University of Munich, Wellcome Trust, National Research Foundation Singapore, European Partnership on Metrology, VELUX Foundation, Bayerisch-Tschechische Hochschulagentur (BTHA), BayFrance (Bayerisch-Französisches Hochschulzentrum), BayFOR (Bayerische Forschungsallianz), and Reality Labs Research. Honoraria for talks: Received honoraria from the ISGlobal, Research Foundation of the City University of New York and the Stadt Ebersberg, Museum Wald und Umwelt. Travel reimbursements: Daimler und Benz Stiftung. Patents: Named on European Patent Application EP23159999.4A ("System and method for corneal-plane physiologically-relevant light logging with an application to personalized light interventions related to health and well-being"). With the exception of the funding source supporting this work, M.S. declares no influence of the disclosed roles or relationships on the work presented herein.

References

Berson, D. M., Dunn, F. A. & Takao, M. Phototransduction by retinal ganglion cells that set the circadian clock. Science **295**, 1070-1073 (2002). https://doi.org/10.1126/science.1067262

Hattar, S., Liao, H. W., Takao, M., Berson, D. M. & Yau, K. W. Melanopsin-containing retinal ganglion cells: architecture, projections, and intrinsic photosensitivity. Science **295**, 1065-1070 (2002). https://doi.org/10.1126/science.1069609

Schmidt, T. M., Chen, S. K. & Hattar, S. Intrinsically photosensitive retinal ganglion cells: many subtypes, diverse functions. Trends Neurosci **34**, 572-580 (2011). https://doi.org/10.1016/j.tins.2011.07.001

Didikoglu, A., Mohammadian, N., Johnson, S., van Tongeren, M., Wright, P., Casson, A. J., Brown, T. M. & Lucas, R. J. Associations between light exposure and sleep timing and sleepiness while awake in a sample of UK adults in everyday life. Proc Natl Acad Sci U S A **120**, e2301608120 (2023). https://doi.org/10.1073/pnas.2301608120

Price, L. L. A., Khazova, M. & Udovicic, L. Assessment of the Light Exposures of Shift-working Nurses in London and Dortmund in Relation to Recommendations for Sleep and Circadian Health. Ann Work Expo Health **66**, 447–458 (2022). https://doi.org/10.1093/annweh/wxab092

Biller, A. M., Zauner, J., Cajochen, C., Gerle, M. A., Kalavally, V., Mohamed, A., Rottländer, L., Seah, M.-Y., Stefani, O. & Spitschan, M. Physiologically-relevant light exposure and light behaviour in Switzerland and Malaysia. bioRxiv, 2025.01.07.631760 (2025). https://doi.org/10.1101/2025.01.07.631760

Agbeshie, G. K., Duah Junior, I. O., Andoh, A. K. A., Ampong, J., Mensah, N. A. O., Ampoma-Mensah, A. Y., Zauner, J., Spitschan, M. & Akuffo, K. O. Physiologically relevant real-world light exposure and its behavioural and environmental determinants in Kumasi, Ghana. Open Res Eur **5**, 300 (2025). https://doi.org/10.12688/openreseurope.21304.1

Brown, T. M., Brainard, G. C., Cajochen, C., Czeisler, C. A., Hanifin, J. P., Lockley, S. W., Lucas, R. J., Munch, M., O'Hagan, J. B., Peirson, S. N., Price, L. L. A., Roenneberg, T., Schlangen, L. J. M., Skene, D. J., Spitschan, M. *et al.* Recommendations for daytime, evening, and nighttime indoor light exposure to best support physiology, sleep, and wakefulness in healthy adults. PLoS Biol **20**, e3001571 (2022). https://doi.org/10.1371/journal.pbio.3001571

Spitschan, M., Zauner, J., Nilsson Tengelin, M., Bouroussis, C. A., Caspar, P. & Eloi, F. Illuminating the future of wearable light metrology: Overview of the MeLiDos Project. Measurement **235**, 114909 (2024). https://doi.org/10.1016/j.measurement.2024.114909

Nowozin, C., Wahnschaffe, A., de Zeeuw, J., Papakonstantinou, A., Hadel, S., Rodenbeck, A., Bes, F. & Kunz, D. Living in Biological Darkness II: Impact of Winter Habitual Daytime Light on Night-Time Sleep. Eur J Neurosci **61**, e16647 (2025). https://doi.org/10.1111/ejn.16647

Burns, A. C., Windred, D. P., Rutter, M. K., Olivier, P., Vetter, C., Saxena, R., Lane, J. M., Phillips, A. J. K. & Cain, S. W. Day and night light exposure are

associated with psychiatric disorders: an objective light study in >85,000
people. Nat Ment Health **1**, 853–862 (2023). https://doi.org/10.1038/s44220-
023-00135-8

12 Gubin, D., Danilenko, K., Stefani, O., Kolomeichuk, S., Markov, A., Petrov, I.,
Voronin, K., Mezhakova, M., Borisenkov, M., Shigabaeva, A., Yuzhakova, N.,
Lobkina, S., Petrova, J., Malyugina, O., Weinert, D. *et al.* Light Environment of
Arctic Solstices is Coupled With Melatonin Phase-Amplitude Changes and
Decline of Metabolic Health. J Pineal Res **77**, e70023 (2025).
https://doi.org/10.1111/jpi.70023

13 Windred, D. P., Burns, A. C., Rutter, M. K., Ching Yeung, C. H., Lane, J. M.,
Xiao, Q., Saxena, R., Cain, S. W. & Phillips, A. J. K. Personal light exposure
patterns and incidence of type 2 diabetes: analysis of 13 million hours of light
sensor data and 670,000 person-years of prospective observation. Lancet
Reg Health Eur **42**, 100943 (2024).
https://doi.org/10.1016/j.lanepe.2024.100943

14 Windred, D. P., Burns, A. C., Lane, J. M., Olivier, P., Rutter, M. K., Saxena,
R., Phillips, A. J. K. & Cain, S. W. Brighter nights and darker days predict
higher mortality risk: A prospective analysis of personal light exposure in
>88,000 individuals. Proc Natl Acad Sci U S A **121**, e2405924121 (2024).
https://doi.org/10.1073/pnas.2405924121

15 Burns, A. C., Saxena, R., Vetter, C., Phillips, A. J. K., Lane, J. M. & Cain, S.
Time spent in outdoor light is associated with mood, sleep, and circadian
rhythm-related outcomes: a cross-sectional and longitudinal study in over
400,000 UK Biobank participants. J Affect Disord **295**, 347–352 (2021).
https://doi.org/10.1016/j.jad.2021.08.056

16 Gubin, D., Danilenko, K., Stefani, O., Kolomeichuk, S., Markov, A., Petrov, I.,
Voronin, K., Mezhakova, M., Borisenkov, M., Shigabaeva, A., Yuzhakova, N.,
Lobkina, S., Weinert, D. & Cornelissen, G. Blue Light and Temperature
Actigraphy Measures Predicting Metabolic Health Are Linked to Melatonin
Receptor Polymorphism. Biology (Basel) **13**, 22 (2023).
https://doi.org/10.3390/biology13010022

17 Leger, D., Bayon, V., Elbaz, M., Philip, P. & Choudat, D. Underexposure to
light at work and its association to insomnia and sleepiness: a cross-sectional
study of 13,296 workers of one transportation company. J Psychosom Res **70**,
29-36 (2011). https://doi.org/10.1016/j.jpsychores.2010.09.006

18 Gubin, D., Julia, B., Oliver, S., Sergey, K., Liina, D., Alexander, M., Aislu, S.,
Germaine, C. & Weinert, D. Light exposure predicts COVID-19 negative status
in young adults. Biol Rhythm Res **55**, 535-546 (2024).
https://doi.org/10.1080/09291016.2024.2427608

19 Gubin, D., Kolomeichuk, S., Danilenko, K., Stefani, O., Markov, A., Petrov, I.,
Voronin, K., Mezhakova, M., Borisenkov, M., Shigabaeva, A., Boldyreva, J.,
Petrova, J., Weinert, D. & Cornelissen, G. Light Exposure, Physical Activity,
and Indigeneity Modulate Seasonal Variation in NR1D1 (REV-ERBalpha)
Expression. Biology (Basel) **14**, 231 (2025).
https://doi.org/10.3390/biology14030231

20 Brown, T. M. Melanopic illuminance defines the magnitude of human circadian
light responses under a wide range of conditions. J Pineal Res **69**, e12655
(2020). https://doi.org/10.1111/jpi.12655

21 Spitschan, M., Smolders, K., Vandendriessche, B., Bent, B., Bakker, J. P.,
Rodriguez-Chavez, I. R. & Vetter, C. Verification, analytical validation and

clinical validation (V3) of wearable dosimeters and light loggers. Digit Health **8**,
20552076221144858 (2022). https://doi.org/10.1177/20552076221144858

22   Spitschan, M. & Joyce, D. S. Human-Centric Lighting Research and Policy in
the Melanopsin Age. Policy Insights Behav Brain Sci **10**, 237-246 (2023).
https://doi.org/10.1177/23727322231196896

23   Schollhorn, I., Stefani, O., Lucas, R. J., Spitschan, M., Slawik, H. C. &
Cajochen, C. Melanopic irradiance defines the impact of evening display light
on sleep latency, melatonin and alertness. Commun Biol **6**, 228 (2023).
https://doi.org/10.1038/s42003-023-04598-4

24   Cajochen, C., Frey, S., Anders, D., Spati, J., Bues, M., Pross, A., Mager, R.,
Wirz-Justice, A. & Stefani, O. Evening exposure to a light-emitting diodes
(LED)-backlit computer screen affects circadian physiology and cognitive
performance. J Appl Physiol **110**, 1432-1438 (2011).
https://doi.org/10.1152/japplphysiol.00165.2011

25   Schöllhorn, I., Stefani, O., Blume, C. & Cajochen, C. Seasonal Variation in the
Responsiveness of the Melanopsin System to Evening Light: Why We Should
Report Season When Collecting Data in Human Sleep and Circadian Studies.
Clocks Sleep **5**, 651-666 (2023). https://doi.org/10.3390/clockssleep5040044

26   Bauducco, S., Pillion, M., Bartel, K., Reynolds, C., Kahn, M. & Gradisar, M. A
bidirectional model of sleep and technology use: A theoretical review of How
much, for whom, and which mechanisms. Sleep Med Rev **76**, 101933 (2024).
https://doi.org/10.1016/j.smrv.2024.101933

27   Guidolin, C., Aerts, S., Agbeshie, G. K., Akuffo, K. O., Aydin, S. N., Baeza-
Moyano, D., Bolte, J., Broszio, K., Cantarero-Garcia, G., Didikoglu, A.,
Gonzalez-Lezcano, R. A., Joosten-Ma, H., Melero-Tur, S., Tengelin, M. N.,
Perez Gutierrez, M. C. *et al.* Protocol for a prospective, multicentre, cross-
sectional cohort study to assess personal light exposure. BMC Public Health
24, 3285 (2024). https://doi.org/10.1186/s12889-024-20206-4

28   Estevan, I., Tassino, B., Vetter, C. & Silva, A. Bidirectional association
between light exposure and sleep in adolescents. J Sleep Res **31**, e13501
(2022). https://doi.org/10.1111/jsr.13501

29   Hand, A. J., Stone, J. E., Shen, L., Vetter, C., Cain, S. W., Bei, B. & Phillips, A.
J. K. Measuring light regularity: sleep regularity is associated with regularity of
light exposure in adolescents. Sleep **46**, zsad001 (2023).
https://doi.org/10.1093/sleep/zsad001

30   da Costa Lopes, L., Ribeiro da Silva Vallim, J., Tufik, S., Louzada, F. &
D'Almeida, V. Associations between real-life light exposure patterns and sleep
behaviour in adolescents. J Sleep Res **33**, e14315 (2024).
https://doi.org/10.1111/jsr.14315

31   Stefani, O., Marek, R., Schwarz, J., Plate, S., Zauner, J. & Schrader, B.
Wearable Light Loggers in Field Conditions: Corneal Light Characteristics,
User Compliance, and Acceptance. Clocks Sleep **6**, 619-634 (2024).

32   Lok, R., Ancoli-Israel, S., Ensrud, K. E., Redline, S., Stone, K. L. & Zeitzer, J.
M. Timing of outdoor light exposure is associated with sleep-wake
consolidation in community-dwelling older men. Front Sleep **2**, 1268379
(2023). https://doi.org/10.3389/frsle.2023.1268379

33   Li, Q., Xu, Y. X., Lu, X. Z., Shen, Y. T., Wan, Y. H., Su, P. Y., Tao, F. B.,
Chen, X. & Sun, Y. Impact of bedroom light exposure on glucose metabolic
markers and the role of circadian-dependent meal timing: A population-based
cross-sectional study. Ecotoxicol Environ Saf **290**, 117589 (2024).
https://doi.org/10.1016/j.ecoenv.2024.117589

Wallace, D. A. Light exposure differs by sex in the US, with females receiving less bright light. npj Biol Timing Sleep **1**, 16 (2024). https://doi.org/10.1038/s44323-024-00016-y

Wallace, D. A., Redline, S., Sofer, T. & Kossowsky, J. Environmental Bright Light Exposure, Depression Symptoms, and Sleep Regularity. JAMA Netw Open **7**, e2422810 (2024). https://doi.org/10.1001/jamanetworkopen.2024.22810

Roguski, A., Needham, N., MacGillivray, T., Martinovic, J., Dhillon, B., Riha, R. L., Armstrong, L., Campbell, I. H., Ferguson, A., Hilgen, G., Lako, M., Ritter, P., Santhi, N., von Schantz, M., Spitschan, M. *et al.* Investigating light sensitivity in bipolar disorder (HELIOS-BD). Wellcome Open Res **9**, 64 (2024). https://doi.org/10.12688/wellcomeopenres.20557.1

Hartmeyer, S. L., Webler, F. S. & Andersen, M. Towards a framework for light-dosimetry studies: Methodological considerations. Light Res Technol **55**, 377–399 (2022). https://doi.org/10.1177/14771535221103258

Stampfli, J. R., Schrader, B., di Battista, C., Häfliger, R., Schälli, O., Wichmann, G., Zumbühl, C., Blattner, P., Cajochen, C., Lazar, R. & Spitschan, M. The Light-Dosimeter: A new device to help advance research on the non-visual responses to light. Light Res Technol **55**, 474–486 (2023). https://doi.org/10.1177/14771535221147140

Zauner, J., Hartmeyer, S. & Spitschan, M. LightLogR: Reproducible analysis of personal light exposure data. J Open Source Softw **10**, 7601 (2025). https://doi.org/10.21105/joss.07601

van Duijnhoven, J., Hartmeyer, S. L., Didikoglu, A., Stefani, O., Houser, K. W., Kalavally, V. & Spitschan, M. Measuring light exposure in daily life: A review of wearable light loggers. Build Environ **274**, 112771 (2025). https://doi.org/10.1016/j.buildenv.2025.112771

Danilenko, K. V., Stefani, O., Voronin, K. A., Mezhakova, M. S., Petrov, I. M., Borisenkov, M. F., Markov, A. A. & Gubin, D. G. Wearable Light-and-Motion Dataloggers for Sleep/Wake Research: A Review. Appl Sci **12**, 11794 (2022). https://doi.org/10.3390/app122211794

Guidolin, C., Zauner, J., Hartmeyer, S. L. & Spitschan, M. Collecting, detecting and handling non-wear intervals in longitudinal light exposure data. bioRxiv, 2024.12.23.627604 (2024). https://doi.org/10.1101/2024.12.23.627604

Bocanegra, G., Stefani, O., Zauner, J., Spitschan, M. & Joosten-Ma, H. Wearable Light Loggers and Dosimeters for Health Impact Measurement: A User Experience Study. SSRN **5047905**, (2024). https://doi.org/10.2139/ssrn.5047905

Zauner, J., Stefani, O., Bocanegra, G., Guidolin, C., Schrader, B., Udovicic, L. & Spitschan, M. Auxiliary data questionnaires. Zenodo **17535086**, (2025). https://doi.org/10.5281/zenodo.17535086

Carney, C. E., Buysse, D. J., Ancoli-Israel, S., Edinger, J. D., Krystal, A. D., Lichstein, K. L. & Morin, C. M. The consensus sleep diary: standardizing prospective sleep self-monitoring. Sleep **35**, 287-302 (2012). https://doi.org/10.5665/sleep.1642

Bajaj, A., Rosner, B., Lockley, S. W. & Schernhammer, E. S. Validation of a light questionnaire with real-life photopic illuminance measurements: the Harvard Light Exposure Assessment questionnaire. Cancer Epidemiol Biomarkers Prev **20**, 1341-1349 (2011). https://doi.org/10.1158/1055-9965.EPI-11-0204

47 Zauner, J., Tsukimori, E. & Spitschan, M. LightLogR Newsletter, Edition #01.
Zenodo **16308351**, (2025). https://doi.org/10.5281/zenodo.16308351
48 Zauner, J., Biller, A. M. & Spitschan, M. Impact of light logger and dosimeter
placement on wearability and appeal in real-life settings. Open Res Eur **5**, 293
(2025). https://doi.org/10.12688/openreseurope.20985.1
49 Zauner, J., Stefani, O., Biller, A. M., Guidolin, C. & Spitschan, M. Web-based
specification tool for wearable light loggers and optical radiation dosimeters
(Version   1.0.1)   [Software].   Zenodo   **1.04ga-fd22**,   (2025).
https://doi.org/10.17617/1.04ga-fd22
50 Murukesu, R. R., Zauner, J. & Spitschan, M. A day in daylight. Zenodo
17467253, (2025). https://doi.org/10.5281/zenodo.17467253
51 Hartmeyer, S. L. & Andersen, M. Towards a framework for light-dosimetry
studies: Quantification metrics. Light Res Technol **56**, 337-365 (2023).
https://doi.org/10.1177/14771535231170500

1072

1073

1074 **Figures**

1075

Figure 1. Schematic of auxiliary data in the context of other study data. Data collected in any
field study using wearable devices sit on a spectrum of time-independent data (i.e., are valid
independent of the time these data were recorded, like participant height or eye colour), and
time-dependent data (i.e., which are valid/relevant only for part of the time, like an
illuminance measurement). Some of these data can be considered metadata of the study,
while others are part of the main data collection effort, e.g., as dependent variables or
confounder controls. A host of secondary data that should be considered are auxiliary data,
which sit on the same end of the time-dependency spectrum as data from wearables and
give context to the wearable data. These data can help to filter and validate wearable data
during preprocessing and allow for additional covariates during analysis.

1086

Summary of melanopic EDI across log entries
Question/Characteristic: What is your general context?

	Distribution[1]							Histogram[2]	Timeline[3]	Mean[1]	Episode duration[4]	Total duration
	0%	2.5%	17%	Median	83%	97.5%	100%					
Outdoors	0	0	14	1,075	8,711	37,422	91,415			5,156	36m 13s (273 eps.)	6d 20h (7%)
Mixed	0	0	16	221	3,777	20,829	76,834			2,459	56m 59s (184 eps.)	1w 6h (8%)
Indoors	0	0	3	54	314	1,638	67,476			319	2h 25m (444 eps.)	6w 2d (48%)
Bed	0	0	0	0	0	32	5,222			5	7h 18m (84 eps.)	3w 4d (27%)
Non-wear	0	0	0	0	22	513	32,405			228	2h 12m (104 eps.)	1w 2d (10%)

[1] Melanopic equivalent daylight illuminance (lx)

[2] Scaled histograms show 66% of data in dark blue, 95% of data in dark & light blue, and the most extreme 5% of data in grey. The median is shown as a red dot.

[3] Timeline plots show the median of the condition (yellow dot) against the rest of the data (grey dot). Colored bands show the interquartile range. Size of the dots indicates the relative number of instances, i.e., large dots indicate many occurances at a given time (relative to other times).

[4] Average duration of an episode in this category. eps. = Episode(s)

Figure 2. Example of the effect of auxiliary data to contextualise personal light exposure. The figure-table shows distributional characteristics of the melanopic EDI time series, a histogram, timeline plot, the mean, the mean duration of the states and number of episodes, as well as total duration for five conditions (Outdoors, Mixed indoor-outdoor, Indoors, Bedtime, and non-wear time. The effect of the auxiliary data is especially visible in the timeline plots, where the light exposure during specific states, collected through methods in the auxiliary data framework, vary strongly from the rest of the data (yellow vs. grey).

1097

1098   **Figure 3**. Rooftop set-up for environmental light logging. A: ActLumus light logger (Condor

1099   Instruments) horizontally placed in water-resistant set-up. B: Position of the set-up on the

1100   rooftop during measurements.

1101 **Tables**

1102

1103   Collected state changes:

Datetime	State_Change	Participant ID
2023-08-28 23:40:00	sleep	Ao12b
2023-08-29 09:37:00	wake	Ao12b
2023-08-29 23:40:00	sleep	Ao12b
2023-08-30 09:21:00	wake	Ao12b
2023-08-30 23:15:00	sleep	Ao12b
2023-08-31 09:47:00	wake	Ao12b

1104   Statechanges transformed to states allows for quality checks (e.g., is the length of a state sensible,
1105   does it fall within a sensible time window of the day), and manipulation to derived states (e.g., Brown
1106   recommendations split the wake times into a "daytime" phase, and an "evening" phase, 3 hours before
1107   sleep):

Interval	State	ID
2023-08-28 23:40:00 - 2023-08-29 09:37:00	sleep	Ao12b
2023-08-29 09:37:00 - 2023-08-29 23:40:00	wake	Ao12b
2023-08-29 23:40:00 - 2023-08-30 09:21:00	sleep	Ao12b
2023-08-30 09:21:00 - 2023-08-30 23:15:00	wake	Ao12b
2023-08-30 23:15:00 - 2023-08-31 09:47:00	sleep	Ao12b
2023-08-31 09:47:00 - NA	wake	Ao12b

1108   A set of functions in the open-source package LightLogR (*add_states* and *interval2data*) facilitate the
1109   robust combination of state data with light exposure time series, even across complex hierarchical
1110   datasets.

Datetime	Melanopic_EDI / lx	Sleep_State	ID

...	...	...	...
2023-08-29 23:39:40	55.5	wake	Ao12b
2023-08-29 23:39:50	79.1	wake	Ao12b
2023-08-29 23:40:00	71.32	sleep	Ao12b
2023-08-29 23:40:10	23.00	sleep	Ao12b
2023-08-29 23:40:20	1.22	sleep	Ao12b
2023-08-29 23:40:30	8.91	sleep	Ao12b
...	...	...	...

1111

1112 **Table 1. Example table for collecting state changes in sleep/wake for one participant**

1113 **and considerations on how these relate to actual measurements of light exposure**

1114

1115

Category	Challenges	Strategies
Methodology (Device placement and usage)	• Wrist placement: Less accurate due to clothing coverage and remoteness to the eye. • Lanyard-worn and clip-worn devices are susceptible to occlusion by jackets. • Glasses placement: More accurate but inconvenient and noticeable.	• Offer less intrusive placement options for balance between accuracy and comfort.
Instructions and guidance	• Unclear instructions. • Lack of reminders. • No feedback on incorrect positioning, battery status, or device inactivity.	• Provide clear instructions & simplify instructions with visual aids. • Use automated reminders & provide real-time feedback.
Social/Environmental factors	• Social discomfort in public settings. • Environmental conditions (e.g., weather, physical activities) affect usage.	• Use devices that are discreet. • Offer guidance for handling the device in different conditions.
Motivational feedback (Incentives and rewards)	• No incentive system. • Lack of feedback on performance.	• Introduce rewards (e.g., gift cards, gamification). • Provide regular feedback on progress and impact.
Data processing (Quality control techniques)	• Need for clear criteria to exclude non-compliant data. • Handling outlier data. • Identifying periods when the device was worn improperly.	• Define exclusion criteria. • Employ auxiliary data strategies • Use advanced algorithms to detect improper wear. Apply robust outlier analysis.

1116

1117   **Table 2.** Summary of quality assurance and quality control strategies

1118   **Supplementary information**

1119   **Supplementary information S1.** PDF document containing the (anonymous) survey results
1120   for the auxiliary data strategy. Email addresses in Q27 and Q28 have been removed for
1121   privacy reasons. Names in Q27 have been preserved, as participants opted to be named as
1122   contributors.

1123

1124

Wearable devices	Device metadata (model, fidelity, etc.)	Dataset metadata (instructions, location, etc.)	Device reports (measurement overviews, battery health, etc.)	Event markers (button presses, etc.)	Device measurements Time series of, e.g., light, optical radiation, distance, activity

Wearable data

Time-independent data → Time-series data

Other sources	Study metadata (groups, contributors, etc.)	Participant metadata (chronotype, age, gender, etc.)	Outcome measures (Melatonin level, health parameters, etc.)	Log / Diary data, regular questionnaires (sleep times, non-wear, exercise, etc.)	Time series from other sources (environmental light level, climate data, etc.)

Auxiliary data

Summary of melanopic EDI across log entries

Question/Characteristic: What is your general context?

	Distribution[1]							Histogram[2]	Timeline[3]	Mean[1]	Episode duration[4]	Total duration
	0%	2.5%	17%	**Median**	83%	97.5%	100%					
Outdoors	0	0	14	**1,075**	8,711	37,422	91,415			5,156	36m 13s (273 eps.)	6d 20h (7%)
Mixed	0	0	16	**221**	3,777	20,829	76,834			2,459	56m 59s (184 eps.)	1w 6h (8%)
Indoors	0	0	3	**54**	314	1,638	67,476			319	2h 25m (444 eps.)	6w 2d (48%)
Bed	0	0	0	**0**	0	32	5,222			5	7h 18m (84 eps.)	3w 4d (27%)
Non-wear	0	0	0	**0**	22	513	32,405			228	2h 12m (104 eps.)	1w 2d (10%)

[1] Melanopic equivalent daylight illuminance (lx)

[2] Scaled histograms show 66% of data in dark blue, 95% of data in dark & light blue, and the most extreme 5% of data in grey. The median is shown as a red dot.

[3] Timeline plots show the median of the condition (yellow dot) against the rest of the data (grey dot). Colored bands show the interquartile range. Size of the dots indicates the relative number of instances, i.e., large dots indicate many occurances at a given time (relative to other times).

[4] Average duration of an episode in this category. eps. = Episode(s)