Expanding vaginal microbiome pangenomes via a custom MIDAS database reveals *Lactobacillus crispatus* accessory genes associated with cervical dysplasia

Claire A. Dubin[1,2], Chunyu Zhao[3,4], Katherine S. Pollard[3,4,5], Tomiko Oskotsky[1,2,6], Jonathan L. Golob[7]*, Marina Sirota[1,2,8]*

1. March of Dimes Prematurity Research Center at the University of California San Francisco, San Francisco, CA, USA
2. Bakar Computational Health Sciences Institute, University of California San Francisco, San Francisco, CA, USA
3. Institute of Data Science & Biotechnology, Gladstone Institutes, San Francisco, California, USA
4. Data Science, Chan Zuckerberg Biohub Network, San Francisco, California, USA
5. Department of Epidemiology & Biostatistics, University of California San Francisco, San Francisco, California, USA
6. Division of Clinical Informatics and Digital Transformation, Department of Medicine, University of California, San Francisco, San Francisco, CA, USA
7. Division of Allergy and Infectious Diseases, University of Washington School of Medicine, Seattle, WA, USA
8. Department of Pediatrics, University of California San Francisco, San Francisco, CA, USA
*These authors contributed equally to this work.

ABSTRACT

The vaginal microbiome plays a central role in reproductive health. Vaginal microbiome dysbiosis increases risk of many adverse reproductive health outcomes, but most studies have focused on associations at the species level. The potential contribution of intraspecies microbial variation, especially gene content differences across bacterial strains, remains underexplored in reproductive health contexts. The Metagenomic Intra-Species Diversity Analysis (MIDAS) framework enables such analyses but depends on comprehensive reference databases. We constructed a MIDAS-compatible pangenome database from over 18,000 genomes in the Vaginal Microbiome Genome Collection (VMGC). Compared to the Genome Taxonomy Database (GTDB)-derived reference, the VMGC database expanded pangenomes of prevalent vaginal species, better capturing vaginal-specific intraspecies diversity. Applying this database to a cervical dysplasia cohort, we identified thirteen *Lactobacillus crispatus* accessory genes significantly associated with increased risk of cervical dysplasia, including a HicAB toxin-antitoxin system, three transcriptional regulators, and three phage-derived genes. These findings highlight the utility of body site-specific reference resources for uncovering intraspecies microbial variation relevant to reproductive health.

INTRODUCTION

The microbiome is increasingly recognized as an essential factor in health and disease throughout the human body.[1] While the gut is the most well-studied system in microbiome research, microbial communities in other body sites also play an important role in host physiology and disease susceptibility.[2,3] Bidirectional interactions between the vaginal microbiome and host play a key role in reproductive health.[4] Disruption to this balance, often called vaginal dysbiosis, is linked to adverse reproductive health outcomes, including preterm birth[5–7] and endometriosis.[8,9] Vaginal dysbiosis involves a shift away from *Lactobacillus*-dominated communities toward more diverse and potentially pathogenic taxa.[4] Pathogenic bacteria in the vaginal microbiome compromise the epithelial barrier and weaken immune responses, heightening susceptibility to infection by other pathogenic bacteria, yeast, and viruses.[4,10] For example, persistent human papillomavirus (HPV) infection is the primary precursor to cervical dysplasia,[11,12] the precancerous growth of abnormal cells on the cervix. Cervical dysplasia is a significant public health concern due to its potential to progress to cervical cancer, the fourth most common cancer in women.[12,13] Further clarifying the role of the vaginal microbiome in preventing or promoting cervical dysplasia is essential for elucidating its underlying etiology, identifying biomarkers, and developing therapeutics.

While the majority of vaginal microbiome studies have focused on composition at the species-level, intraspecies genomic variation also plays a role in the functional potential and ecological dynamics of the vaginal microbiome. *Lactobacillus crispatus*, a widely prevalent species often regarded as a marker of a healthy vaginal microbiome, has a large accessory genome compared to other common vaginal *Lactobacillus* species,[14] and frequent gain and loss of genes across *L. crispatus* strains results in genetic and phenotypic variation among members of the species. *L. crispatus* vaginal isolates vary in their ability to utilize glycogen sources, produce hydrogen peroxide, and survive acidic stress.[15] Genes encoding mucin-binding proteins and bacteriocins, which enhance adhesion to host tissues and inhibit pathogen growth, respectively, are variably present across *L. crispatus* strains.[15–17] Vaginal *L. crispatus* strains differ in their ability to suppress the growth of bacterial and fungal pathogens, and *L. crispatus* and other common vaginal microbiome species exhibit

different co-occurrence patterns at the strain level.[18] These observations suggest that *L. crispatus* may not universally inhibit progression to a dysbiotic vaginal microbiome state.[19] Recognizing and characterizing this intraspecies variation is essential for identifying reliable biomarkers of disease and developing probiotics to support reproductive health.

The Metagenomic Intra-Species Diversity Analysis (MIDAS) pipeline is a versatile tool for measuring intraspecific variation at the nucleotide and gene level from shotgun metagenomic sequencing data.[20,21] MIDAS can be used to quantify bacterial gene content in metagenomic samples by mapping metagenomic reads to a preconstructed pangenome reference database. Intraspecies gene presence/absence data generated by MIDAS can be used in metagenome-wide association studies (MWAS) to identify bacterial accessory genes associated with clinical outcomes. For instance, recent work using the MWAS-based model microSLAM identified a fructoselysine utilization operon in *Faecalibacterium prausnitzii* that is negatively associated with inflammatory bowel disease.[22] Given the demonstrated utility of MIDAS and microSLAM in identifying accessory genes associated with disease states and the mounting evidence for intraspecies genotypic and phenotypic variation in the vaginal microbiome, we sought to optimize this framework to identify associations between accessory genes in common vaginal microbiome species and reproductive health.

In this work, we address the limited representation of vaginal-derived genomes in existing MIDAS pangenome databases by developing a pipeline for constructing MIDAS-compatible custom pangenome reference databases and applying it to construct a pangenome database from a collection of over 18,000 vaginal genomes. We demonstrate that this database expands the pangenomes of six prevalent vaginal microbiome species to better reflect the genetic diversity of the vaginal microbiome compared to existing MIDAS pangenome databases. We then apply the newly-constructed database to quantify gene content in vaginal microbiome samples and identify thirteen *L. crispatus* accessory genes associated with increased risk of cervical dysplasia, including a toxin-antitoxin system and phage-derived genes of unknown function.

RESULTS

Construction of a MIDAS-compatible pangenome reference database from vaginal genomes

Reference-based gene content analyses are dependent on the quality of the reference database; genes that are absent from the reference cannot be measured, and an excess of irrelevant genes can introduce unnecessary noise. Two prebuilt MIDAS-compatible databases were previously constructed from genomes in the Unified Human Gastrointestinal Genome collection (UHGG)[23] and the Genome Taxonomy Database r202 (GTDB).[24] UHGG exclusively contains genomes from the human gut microbiome, and although GTDB contains a broader set of genomes, genomes from gastrointestinal samples make up approximately 35% of human-derived GTDB genomes with specified origin. In contrast, genomes from other human body sites such as the female reproductive tract, oral cavity, and skin make up only 1%, 2%, and 4%, respectively. Recent collections of genome assemblies derived from metagenomic sequencing data have increased the number of available genomes for prevalent species in these less-studied body sites.[25–27] Expanding pangenome databases to include these metagenome-assembled genomes may better reflect the microbial genetic diversity within these body sites. We set out to develop a MIDAS-compatible reference set of genomes specifically for use with vaginal microbiome specimens to refine and expand the landscape of genes available for analysis of gene content and associations with reproductive health.

The Vaginal Microbiome Genome Collection (VMGC) assembled prokaryotic genomes from publicly available metagenomic and single-isolate sequencing data from vaginal samples. To construct a vaginal microbiome-specific pangenome database from this data, the 19,542 VMGC prokaryotic genomes and their GTDB 214.1 taxonomic assignments were downloaded from Huang et al (2024).[25] There is a median of 19 input genomes per species (IQR 7.0-93.5), and genome quality ranges from medium-quality (52%), to high-quality (43%), and near-complete (5%) as calculated in Huang et al. (2024).[25] We developed a Nextflow pipeline to streamline construction of a MIDAS-compatible pangenome database from a user-specified set of genomes and their

species assignments as described in Zhao et al. (2023)[21] and Smith et al. (2025)[20] (Figure 1A). Using this

pipeline, we created a pangenome reference database for each prokaryotic species with at least five assigned

genomes, totaling 18,970 genomes across 179 species, 81 genera, and 42 families. Predicted genes from all

genomes assigned to a species were clustered by average nucleotide identity (ANI) thresholds ranging from

75-99%. Higher ANI thresholds can distinguish subtle differences among highly similar genes, while lower

thresholds can group more distant homologs. To balance detection of gene variants and grouping of gene

families, we focused our pangenome analysis on 90% ANI gene clusters. The number of 90% ANI gene

clusters per species ranges from 713 to 18,533 (IQR 2358.0-4409.5) and is positively correlated with the

number of input genomes (Figure 1B, Spearman $\rho = 0.57$, $p = 5e\text{-}17$). This trend is consistent across all tested

ANI thresholds, with pangenome size decreasing as gene clusters are merged at lower ANI thresholds (Figure

S1, Table S1).

Comparison to GTDB-only database

We next compared the ability of the VMGC-derived MIDAS pangenome reference database to reflect the

genetic diversity of organisms in the vaginal microbiome as compared to the GTDB-derived reference.

Pangenome expansion widens the set of prokaryotic genes that can be tested for differential prevalence

across strains and associated with clinical data. We compared pangenome sizes (number of 90% ANI gene

clusters) of 129 species with one-to-one mappings across GTDB versions r202 and r214.1, which were used to

determine taxonomic classification in the VMGC and GTDB databases, respectively. Of these species, 66%

(85/129) have a larger pangenome in the database constructed from VMGC genomes, and 34% (44/129) have

a larger GTDB pangenome (Figure 2A). Differences in pangenome size are positively correlated with

differences in the number of genomes per species in each database (Figure S2, Spearman $\rho = 0.92$, $p = 8e\text{-}$

55). The five species with the greatest difference in pangenome size between GTDB and VMGC, all with larger

GTDB pangenomes, are gut-associated species (*Phocaeicola vulgatus, Enterococcus faecalis, Klebsiella*

pneumoniae) and/or widely sampled pathogenic species (*Acinetobacter baumannii, Escherichia coli*). Species

with a larger VMGC pangenome include all four hallmark *Lactobacillus* species used to classify vaginal

community state types (*L. crispatus, L. gasserii, L. iners,* and *L. jensenii*),[28] as well as species associated with

vaginal microbiome dysbiosis, including six *Bifidobacterium vaginale* species (collectively referred to as

Gardnerella vaginalis in NCBI taxonomy) and *Fannyhessea vaginae* (formerly named *Atopobium vaginae*).

We used MIDAS to quantify the relative abundance of 129 species with one-to-one mappings across GTDB

versions among 352 vaginal samples from Norenhag et al. (2024)[29] (Figure S3). Of the 86 species with

relative abundance greater than 1% in at least one vaginal sample, 76% had larger pangenomes in the

database built from VMGC genomes. VMGC has a larger pangenome and more input genomes for all six

species with relative abundance greater than 5% in at least 5% of samples (Figure 2A-B). We compared

pangenome completeness between databases using rarefaction analysis on these six species. Tail slopes

from collector's curves (calculated using the four largest subsampling points per database) were lower in

VMGC than GTDB across all species (Wilcoxon $p = 0.0312$), indicating that VMGC-derived pangenomes are

approaching saturation while GTDB pangenomes continue to accumulate novel gene content as more

genomes are added (Figure 2C-D).

VMGC-based pangenome database expands accessory gene repertoire in prevalent

vaginal bacteria

Pangenome expansion is most meaningful when it uncovers biologically relevant genes, rather than merely

increasing the number of genes associated with a species. To further characterize differences between

pangenome reference databases constructed from VMGC and GTDB genomes, we analyzed the overlap and

distinctiveness of the gene clusters across databases. We rebuilt pangenome databases with the set of all

genomes in VMGC and GTDB for the six prevalent species identified above. The resulting 90% ANI gene

clusters were classified as follows: VMGC-only (containing sequences from genomes found exclusively in

VMGC), GTDB-only (containing sequences from genomes found exclusively in GTDB), or shared (containing

sequences from genomes found in VMGC and GTDB; i.e., at minimum, a single sequence from a genome

found in both GTDB and VMGC, or two sequences: one from a genome found only in VMGC and one from a

genome found only in GTDB). The number of VMGC-only genes was larger than the number of GTDB-only

genes for all six species (Figure 3A).

To see how commonly the database-specific genes are detected in vaginal samples, we used MIDAS and the

combined reference database to measure gene presence and absence in vaginal samples from Norenhag et

al. (2024).[29] While *L. crispatus* has a relatively large number of GTDB-only genes, less than 3% (74/2686) of

these genes are detected among the 130 vaginal samples with *L. crispatus* present (Figure 3B). Across all six

species, shared genes are more frequently detected than VMGC-only and GTDB-only genes, and more

VMGC-only genes are detected than GTDB-only genes (Figure 3B-C). Among genes detected in at least one

sample, VMGC-only genes are found at a significantly higher frequency than GTDB-only genes in *L. crispatus*

and *B. vaginale* (two-sided Mann Whitney U $p = 7 \times 10^{-6}$ and 0.002 respectively). Higher-frequency accessory

genes are particularly valuable in association analyses, as they provide greater statistical power to detect

associations with clinical traits.

Accessory genes often encode functions that are relevant to fitness in a specific environment. We used a

permutation analysis to identify differentially prevalent PFAM domains in VMGC-only and GTDB-only gene

clusters. *L. crispatus* GTDB-only centroids were enriched for eleven PFAM domains, including

glycosyltransferase domains, cell surface-related domains, and a bacteriophage abortive infection AbiH

domain (Table 1). In contrast, *L. crispatus* VMGC-only centroids were enriched for three domains associated

with integrated phage sequences: BRO family N-terminal domain, AAA domain, and Terminase_1 (Table 1). *L.*

crispatus VMGC-only genomes had a higher number of predicted viral regions compared to GTDB-only

genomes (Mann Whitney U $p = 0.001$, Figure S4). More broadly, *L. crispatus* genomes from human hosts had

a higher number viral regions than those from non-human hosts ($p = 1 \times 10^{-5}$), and genomes from the human

female reproductive tract had more predicted viral regions than genomes from the human gastrointestinal tract

($p = 0.012$) and the combined set of genomes from all other human body sites ($p = 0.014$) (Figure S4). This

finding aligns with the known prevalence of viruses and bacteriophage in the vaginal microbiome,[30,31] and

demonstrates that the site-specific database is better equipped to capture virus-bacteria interactions at the

genomic level.

Thirteen *Lactobacillus crispatus* accessory genes are associated with increased cervical dysplasia risk

We applied the VMGC pangenome database to shotgun metagenomic data from a Swedish cervical dysplasia cohort[29] to evaluate its effectiveness in detecting intraspecies associations with clinical phenotypes. We focused our intraspecies analysis on *L. crispatus*, the most abundant species across samples. Of 131 samples with *L. crispatus*, 82 are from healthy controls and 49 are from individuals with cervical dysplasia. We focused our association analysis on 75% ANI gene clusters, representing broader functional gene groups, to increase statistical power by reducing multiple testing. We identified 832 75% ANI gene clusters detected in 15% to 75% of these samples. We used the MWAS-based tool microSLAM to model the associations between the presence/absence vectors of these genes and disease state, also including *L. crispatus* relative abundance, a propensity score calculated from five clinical metadata variables, and population structure as covariates (see Methods). We identified thirteen accessory genes associated with increased risk of cervical dysplasia (locFDR < 0.2; OR = 2.48-6.19; Figure 4A). These genes encode six hypothetical proteins, a protein with a domain of unknown function, three transcriptional regulators, an ion transporter, and both components of the HicAB type II toxin-antitoxin system. All significant genes are more frequently detected in cervical dysplasia samples than in samples from healthy controls (Figure 4B). The ion transporter is a VMGC-only gene. Two groups of genes have similar co-occurrence patterns across samples (Jaccard similarity >= 0.8) (Figure 4C).

Genomic context of cervical dysplasia-associated genes

To further elucidate potential functions of the cervical dysplasia-associated genes, we analyzed their genomic context in high-quality *L. crispatus* isolates. We identified all *L. crispatus* genome assemblies available on NCBI constructed from single-isolate (non-metagenomic) sequencing and a 'complete genome' assembly level. We used BLASTn to identify predicted coding sequences from these 22 strains with homology to the thirteen significant gene sequences. The significant genes are variably detected across the *L. crispatus* isolate genomes (Figure 5A), consistent with our variable detection of them in genomes. Seven strains had two or fewer cervical-dysplasia associated genes detected, suggesting that these may be ideal probiotic candidates for treatment of cervical dysplasia. One of these strains, M247, was shown to increase HPV clearance in a

preliminary trial (N=35).[32] In a larger case-control study (N=160), the group treated with oral M247 had significant regression of cervical lesions, though their overall HPV clearance rates did not significantly differ from controls.[33] This demonstrates the potential for our framework to identify potential probiotic strains based on accessory gene content.

The sets of genes with similar co-occurrence patterns across samples were co-located in syntenic regions of the isolate genomes (Figure 5B and 5C). We used Operon-mapper[34] to predict co-regulated genes in these regions. A group of five cervical dysplasia-associated genes encoding two hypothetical proteins, an XRE family transcriptional regulator, a MarR family transcriptional regulator, and a protein with a domain of unknown function are genomically adjacent, but only the latter two genes are predicted to be co-regulated (Figure 5B). The group of three co-occurring genes encoding hypothetical proteins are located within an approximately 40 kb region with phage-related annotations and are predicted to be part of a single operon (Figure 5C, S5). GeNomad also predicted this region to be related to the *Caudoviricetes* phage class (Figure S5). The HicAB toxin-antitoxin system components are predicted to be part of an operon that also includes two hypothetical proteins (Figure 5D). The LacI transcriptional regulator is part of a potential carbohydrate utilization operon (Figure 5E). The remaining hypothetical protein is predicted to be co-regulated with a GNAT family N-acetyltransferase (Figure 5F), and the sodium transporter is not predicted to be co-regulated with any other genes (Figure 5G). The cervical dysplasia-associated genes span diverse functional categories and genomic contexts; the prevalence of proteins with unknown functions in and near the set of cervical dysplasia-associated genes highlights the need for functional studies to investigate the mechanistic basis of these associations.

DISCUSSION

In this study, we constructed a vaginal microbiome-specific pangenome database using the Vaginal Microbiome Genome Collection and demonstrated its utility in identifying associations between bacterial accessory genes and clinical phenotypes. This is important, because previously published pangenome

databases are largely derived from gastrointestinal and environmental organisms, limiting their ability to reflect

the genetic diversity of bacteria from other body sites such as the human vaginal microbiome. For hallmark

Lactobacillus species, as well as dysbiosis-associated organisms like *Fannyhessea vaginae* and

Bifidobacterium vaginale, use of VMGC genomes increased pangenome size and completeness, with percent

increase in size ranging from 21% to 332%. Pangenome expansion uncovered accessory genes that were

frequently detected in vaginal samples and enriched for environment-specific functions. For *L. crispatus*,

vaginal genomes were enriched for regions of viral origin and PFAM domains associated with integrated

phage, consistent with prior observations that viral elements are widely prevalent in the vaginal niche.[31,35] One

of these cervical dysplasia-associated genes (*nhaC* ion transporter) is found in the VMGC pangenome

database but not in the GTDB database, highlighting the importance of using a body site specific database in

MWAS analyses.

We identified thirteen *L. crispatus* accessory genes associated with increased risk of cervical dysplasia,

despite the species' overall protectiveness against cervical dysplasia and other adverse reproductive health

outcomes.[36] These include transcriptional regulators, toxin-antitoxin system components, and proteins of

unknown function, several of which were embedded within phage-associated regions or putative operons. The

HicAB toxin-antitoxin system promotes biofilm formation and inhibits translation to slow bacterial growth in

response to stress, promoting persistence during antibiotic treatment and acid stress in *Burkholderia*

pseudomallei and *Acetobacter pasteurianus.*[37–39] Other type II antitoxin systems are involved in abortive

infection of phage DNA; while a similar role has been proposed for the HicAB system via comparative

genomics,[40] experimental validation is needed to confirm this role. Given the well-documented relationship

between HPV infection and cervical dysplasia,[11] the significance of three genes in a phage-derived region is

particularly notable and warrants further study of virus-bacteria interactions in this context. Experimental work

is needed to understand any mechanistic role of these genes; they may be directly involved in the etiology of

cervical dysplasia, or they may make *L. crispatus* more permissive to the growth of pathogenic species,

increasing the likelihood of transitioning to a dysbiotic state that favors viral infection and persistence.

The heterogeneity within *L. crispatus* populations underscores the need to move beyond species-level

assessments when studying host-microbiome relationships. In particular, distinguishing beneficial from neutral

or potentially pathogenic sublineages may be critical for understanding why *L. crispatus* dominance is not

universally protective and for guiding design of probiotic interventions. Norenhag, et al. 2024 identified amino

acid biosynthesis and sugar degradation pathways enriched in healthy controls and nucleotide and

peptidoglycan biosynthesis pathways enriched in cases of cervical dysplasia.[29] *L. crispatus* strongly

contributed to the pathways enriched in healthy controls, but within-species variation of these pathways was

not reported. This is the first study to our knowledge to analyze bacterial intraspecies variation in relation to

cervical dysplasia and points to the analytic gain from a richer, more tailored, reference database.

Unlike existing scripts for constructing a MIDAS-compatible pangenome database, our Nextflow pipeline is

automated, highly parallelized, and compatible with multiple executors. This workflow may be particularly

useful for constructing pangenome databases from genome collections recently published for the oral and skin

microbiomes,[26,27] as well as keeping reference sets updated and up-to-date with new data. VIRGO2, an

expansive catalog of non-redundant vaginal microbiome genes pooled across species, is also an excellent

resource for measuring genetic diversity in the vaginal microbiome.[41] Our reference database contains

pangenomes constructed for each species individually, and adjustable ANI clustering thresholds to suit the

desired level of granularity. Our approach may be particularly advantageous for studies focused on specific

species or for seamless integration with the MIDAS pipeline.

This study has several limitations. We did not stratify cases by severity of dysplasia, as the sample sizes within

each subgroup would be too small for statistical analysis. Access to healthcare influences the availability of

cervical dysplasia screening and screening practices vary across countries,[42] so individuals included in this

study may not fully represent the broader population of people with and without cervical dysplasia. Our

methodology for accurate measurement of gene content relies on sufficient sequencing depth across several

samples, so it may be challenging to apply this framework to less abundant or less prevalent species. The top

dysplasia-associated *L. crispatus* gene is a hypothetical protein, underscoring the need for approaches to

improve the functional annotation of vaginal pangenomes. Finally, the significant genes identified here are

associations rather than causal determinants of cervical dysplasia; experimental validation is necessary to

confirm their roles and characterize the genes with unknown function.

Together, our work establishes both a resource and a methodological framework for expanding MIDAS

pangenome databases to better represent body site-specific microbiomes. The VMGC-based database

improves the resolution of metagenomic analyses in the vaginal microbiome, reveals novel accessory genes of

potential clinical importance, and demonstrates that intraspecies diversity within key taxa like *L. crispatus* may

shape reproductive health outcomes. Extending similar approaches to other body sites and diseases will help

uncover overlooked functional variation within the human microbiome and elucidate its role in health and

disease.

METHODS

Categorization of GTDB genome sources

We downloaded metadata for the 258,405 GTDB genomes used in the GTDB-based pangenome database via

the NCBI datasets command line tool version 18.0.2.[43] We identified 98,680 genomes with the words "human"

or "homo sapiens" in the Assembly BioSample Host, Assembly BioSample Source type, Assembly BioSample

Isolation source, Assembly BioSample Models, and/or Assembly BioProject Lineage Title. We manually

categorized the 187 most common isolation sources into the following categories: skin, gastrointestinal,

respiratory, female reproductive tract, oral, blood, urine, other, unknown, and non-human. These manual

annotations categorized the source of 91% (89,483/98,680) of human-associated genomes. We used the

gemma-2-9b-it[44] large language model to categorize the 3,022 remaining isolate sources corresponding to 9%

of genomes. We removed genomes whose sources remained unknown (n = 33,573) or were recategorized as

non-human (n = 532) before calculating the proportions of genomes isolated from each body site. For the six

species depicted in Figure 2B, we manually verified and re-annotated genomes whose sources were

categorized as unknown and non-human through a literature search.

Genome assembly download

Prokaryotic vaginal genomes and their taxonomic assignments to GTDB r214.1 were downloaded from Huang

et al. (2024).[25] Accession numbers and GTDB r202 species assignments for 258,405 genomes in the

preconstructed pangenome database generated from GTDB r202 genomes were downloaded via MIDAS

version 3.0.1.[20] We used gtdb-taxdump version v0.5.0[45] to map taxonomic assignments across versions r202

and r214.1 of GTDB. Species not present in both database versions and species with many-to-one mappings

across versions were excluded from the comparison of pangenome databases. We verified that genomes from

the remaining species were assigned to the same species in GTDB versions r202 and r214.1, and we

downloaded GTDB genome assemblies via the NCBI datasets command line tool version 18.0.2.[43]

Nextflow pipeline for constructing custom pangenome databases

Nextflow is an HPC and workflow orchestration system widely used in biomedical research notable for

versatility and scalability. We created a Nextflow pipeline to streamline construction of a pangenome database

from a user-specified set of genomes and their species assignments. The steps for constructing a pangenome

reference database are described in Zhao et al. (2023)[21] and Smith et al. (2025)[20] and Figure 1A. Briefly, each

genome is annotated via Prokka,[46] and predicted protein-coding sequences are clustered at 99% ANI using

VSEARCH.[47] The centroids of each cluster are reclustered at 95, 90, 85, 80, and 75% ANI, and refined using

CD-HIT.[48] Centroid sequences of 99% ANI clusters are functionally annotated using eggNOG.[49] For each

genome, regions associated with antimicrobial resistance, viral elements, and plasmid/mobile element-

associated regions are identified using ResFinder,[50] GeNomad,[51] and MobileElementFinder,[52] respectively.

We verified that our Nextflow pipeline outputs identical results to the original database build scripts published in

Smith et al. (2025)[20] for two sample species. The database build from VMGC genomes took 199 CPU days (16

real-time days) on a high performance computing cluster, distributed across 506 nodes equipped with Intel and

AMD processors and 12-256 CPUs and 48-1512 GiB of RAM per node. Most processes used 4 or fewer CPUs

and less than 16GB of RAM; the Nextflow pipeline dynamically increased CPU and RAM requests for more

computationally-intensive processes. While this work was completed on a Sun Grid Engine-managed cluster,

Nextflow enables easy transfer to SLURM, AWS, GCP, and other HPC systems.

 Cervical dysplasia samples

We downloaded shotgun metagenomic sequencing reads from European Nucleotide Archive projects

PRJEB72778 and PRJEB72779 and corresponding metadata from Norenhag et al. (2024).[29] We used

Geneshot[53] to run a quality control workflow on the sequencing data, using Trim Galore version 0.6.6[54] to

remove adapter sequences, bwa version 0.7.18[55] to identify sequences mapping to the human genome version

hg38, and samtools version 1.20[56] to filter out the non-microbial reads. We removed samples with fewer than

250,000 reads remaining after quality control, leaving 352 samples for analysis (175 samples from healthy

controls and 177 samples from individuals with cervical dysplasia).

We quantified species-level composition in each sample using the MIDAS run_species command and merged

results across samples with the merge_species command. We considered a species to be present in a sample

if the median coverage across single-copy marker genes was at least 0.75 and relative abundance was at least

1%. 138 and 137 species were detected in at least one sample when using the VMGC and GTDB-based

pangenome database as a reference, respectively.

Rarefaction analysis

We selected six species with a relative abundance greater than 5% in at least 5% of samples for rarefaction

analysis. For each species and database, rarefaction was performed by selecting a subset of genomes,

rebuilding the pangenome database with only those genomes as input, and calculating the number of 90%

average nucleotide identity gene clusters in the resulting pangenome database. The number of genomes

selected began with n-1 genomes, where n is the total number of genomes available for that species, and

decreased by 10% until 2 genomes remained. Ten replicates at each number of genomes were performed.

Gene cluster uniqueness

Genomes from the six species selected above were assigned categories: VMGC-unique (only present in the

VMGC database), GTDB-unique (only present in the GTDB database), or common (present in both the VMGC

and GTDB databases). For each species, we rebuilt the pangenome database with the combined set of

VMGC-unique, GTDB-unique, and common genomes as input. 90% ANI gene clusters were categorized as

"VMGC-only" if they only contained sequences derived from VMGC-unique genomes, "GTDB-only" if they only

contained sequences derived from GTDB-unique genomes, and shared if they contained sequences from a

common genome or sequences from both VMGC-unique and GTDB-unique genomes.

PFAM enrichment

We used eggNOG-generated PFAM domain annotations to test for functional enrichment in VMGC- and

GTDB-only gene clusters for each species. $N_{clusters}$ represents the total number of VMGC-only or GTDB-only

90% ANI gene clusters for a species, and N_{PFAM} represents the number of database-unique gene clusters with

a given PFAM domain. For each PFAM with N_{PFAM} greater than or equal to ten, we took 10,000 samples of

$N_{clusters}$ gene clusters from the set of all gene clusters for the species. We used the proportion of samples with

more than N_{PFAM} gene clusters with the given PFAM annotation as an empirical *p* value. *p* values for each

species were adjusted for multiple hypothesis testing via the Benjamini Hochberg procedure.[57]

Gene presence/absence

The MIDAS run_genes command was used to measure gene content in each vaginal microbiome sample for

all species detected in the sample with median coverage of marker genes greater and 2, reads mapping to at

least 50% of marker genes, and mean depth of at least 5 across all detected genes for the species. Gene

clusters with copy number of 0.35 or greater were considered present in a sample. Results of the gene-level

analysis were merged across samples via the merge_genes command, with genes clustered at 90% ANI for

pangenome database comparisons and 75% ANI for the cervical dysplasia association analysis. The six-

species combined database was used as a reference for the database comparisons and the pangenome

database constructed from VMGC genomes was used as a reference in the cervical dysplasia association

analysis.

Propensity score calculation

For each of 21 metadata variables reported in Norenhag et al. (2024), we constructed a logistic regression

model: outcome ~ variable + 1, with outcome = 0 representing healthy controls and outcome = 1 representing

cervical dysplasia. Five variables with *p* < 0.1 in the univariate models were used in the propensity score:

LowRisk_HPV, PotentialHR_HPV, Gyn_problem_24hrs, Menses_yesno, and Gyn_BloodyDischarge. Each of

these variables is binary. LowRisk_HPV and PotentialHR_HPV did not have missing values for any samples.

The variable Menses_yesno was missing data for 7% of samples; we imputed missing values for this column

based on subject age. Subjects with AgeAtSampling < 52 were imputed with True, and subjects with

AgeAtSampling >= 52 were imputed with False. 52 was selected as the threshold because the average age of

menopause is 51-52.[58] Gyn_problem_24hrs and Gyn_BloodyDischarge had low frequency of nonzero values

(16% and 2% of samples, respectively), and for these columns we imputed missing values (10% of samples for

both variables) with the mode (0). We fit a logistic regression model with the matrix of selected variables as

features and outcome as the response variable. The predicated probability of each sample belonging to the

disease group based on this model was used as a propensity score.

Cervical dysplasia association analysis

We used the microSLAM R package version 1.0.0.9[22] to model associations between *L. crispatus* gene content

and outcome (*y*). We calculated fixed effects of covariates (α) using a baseline generalized linear model

(GLM) y ~ propensity score + *L. crispatus* relative abundance. We constructed a gene presence/absence

matrix from 75% ANI gene clusters detected in 15-75% of samples with mean depth of at least 5 across

detected *L. crispatus* genes (n = 702 genes and 131 samples). Next we calculated a gene relatedness matrix

(GRM), a pairwise similarity matrix based on gene presence/absence data used as a proxy for population

structure across samples. To calculate the random effects between population structure and outcome (*b*), we

combined the baseline GLM and the GRM in a random-effects generalized linear model. Permutation of the tau

variable from this model indicated there was not a significant correlation between population structure and

outcome (*p* = 0.34). Lastly, we fit a mixed effects model for each gene presence absence vector *g:*

$$logit(y_i) = X_i\alpha + \beta g_i + b_i + \varepsilon.$$

where *i* represents a sample, *X* is the covariate matrix, and *β* represents the effect size of the gene. All models

and statistical tests except for the baseline GLM were implemented via microSLAM. We used the locfdr R

package version 1.1.8[59] to calculate false discovery rates from the *z* scores derived from the mixed effects

model. Significant gene clusters were annotated via BLASTx[60] of the centroid sequences.

Analysis of *L. crispatus* isolate genome assemblies

We identified 22 *L. crispatus* genome assemblies available on NCBI with sequencing data from a single

isolate, assembly level of "complete assembly," and available coding sequence annotations. Accession

numbers for these isolates are provided in Table S2. We used BLASTn version 2.16.0[60] with minimum query

and subject coverage of 75% and minimum identity of 75% to identify predicted coding sequences in each

strain with homology to the centroids of 75% ANI gene clusters for *L. crispatus.* The gene with the highest bit

score was selected for each coding sequence. If a gene did not have any matches from a strain, it was

considered absent in that strain. We used the Operon-mapper web tool[34] to predict the location of operons and

geNomad[51] to predict viral regions in each strain. PyGenomeViz version 1.6.1[61] was used for visualizations in

Figures 5B-G and Supplementary Figure 4. The PyGenomeViz integration of MMseqs2 reciprocal best hit

search identified sequences with identity of at least 50% across at least 80% of the query and target

sequences.

Data Availability

10 The VMGC pangenome database will be hosted on AWS and made publicly available upon publication.

Code Availability

12 The Nextflow pipeline for constructing custom MIDAS-compatible pangenome databases and code for data

13 analysis in this paper will be posted on Github and made publicly available upon publication.

FIGURES

Figure 1. Construction of a custom pangenome database. A. Schematic for Nextflow pipeline for constructing custom pangenome database from genomes of a given species. B. Number of genomes vs. number of 90% ANI gene clusters for each species with at least 5 genomes in VMGC.

Figure 2. Comparison of pangenome databases constructed from VMGC and GTDB genomes. A. Comparison of the number of 90% ANI gene clusters in the pangenome among shared species in databases constructed from VMGC and GTDB genomes. Species with relative abundance greater than 5% in at least 5% of samples are labeled in bold. B. Comparison of the number of genomes and their origins for six prevalent vaginal microbiome species. *Bifidobacterium vaginale* is a member of the clade referred to as *Gardnerella vaginalis* in NCBI taxonomy, and *Fannyhessea vaginae* was formerly known as *Atopobium vaginae*. C-D. Rarefaction of pangenome size for six selected species using subsampled genomes from VMGC (C) and GTDB (D).

Figure 3. Detection of shared and unique gene clusters in pangenomes constructed from VMGC and GTDB genomes. A. Venn diagrams of 90% ANI gene cluster overlap for six prevalent vaginal microbiome species. B. Proportion of gene clusters detected in samples from Norenhag et al. (2024), stratified by cluster classification. C. Frequency of detection of gene clusters detected in at least one sample, stratified by cluster classification.

Figure 4. *L. crispatus* accessory genes associated with increased cervical dysplasia risk. A. Effect size and significance of 832 accessory genes used as input into association analysis. B. Prevalence of significant genes across samples; significant genes are detected in a larger proportion of cervical dysplasia samples compared to healthy controls. C. Presence and absence of significant genes across samples, clustered by similarity of gene presence/absence vectors. Genes in red brackets have pairwise Jaccard similarity greater than or equal to 0.8 for all genes in the group.

Figure 5. Detection and genomic context of cervical dysplasia-associated genes in NCBI *L. crispatus* isolates. A. Presence/absence of cervical dysplasia-associated genes among 22 *L. crispatus* genome assemblies from single-isolate sequencing. **B-G.** 5kb genomic context windows around significant genes among NCBI isolates with the given gene(s) detected. Gray connections across isolates indicate genes are reciprocal best hits as identified by MMseqs2, with sequence identity of at least 75% across at least 75% of both sequences. Brown segments indicate operons predicted by Operon-mapper.

TABLES

Table 1: PFAM enrichment

Species	Enrichment	PFAM	BH-adjusted p
Lactobacillus crispatus	Enriched in GTDB-only centroids	Bacteriophage abortive infection AbiH	0.0080
		YSIRK type signal peptide	0.0080
		LPXTG cell wall anchor motif	0.0080
		Methylase_S	0.0080
		Glycosyl transferase family 2	0.0080
		Glycosyl transferases group 1	0.0080
		N-6 DNA Methylase	0.0140
		Polysaccharide biosynthesis C-terminal domain	0.0224
		Glycosyltransferase Family 4	0.0224
		Hexapep	0.0420
		SLAP (surface layer-associated protein) domain	0.0440
	Enriched in VMGC-only centroids	BRO family, N-terminal domain	0.0080
		AAA domain	0.0356
		Terminase_1	0.0430
Fannyhessea vaginae	Enriched in VMGC-only centroids	CHAP domain	0.0190
		Methylase_S	0.0190

REFERENCES

1. Rackaityte E, Lynch SV. The human microbiome in the 21st century. *Nat Commun*. 2020;11(1):5256. doi:10.1038/s41467-020-18983-8

2. Reynoso-García J, Miranda-Santiago AE, Meléndez-Vázquez NM, et al. A complete guide to human microbiomes: Body niches, transmission, development, dysbiosis, and restoration. *Front Syst Biol*. 2022;2. doi:10.3389/fsysb.2022.951403

3. Asangba AE, Mugisha L, Rukundo J, et al. Large Comparative Analyses of Primate Body Site Microbiomes Indicate that the Oral Microbiome Is Unique among All Body Sites and Conserved among Nonhuman Primates. *Microbiol Spectr*. 2022;10(3):e01643-21. doi:10.1128/spectrum.01643-21

4. Zhu B, Tao Z, Edupuganti L, Serrano MG, Buck GA. Roles of the Microbiota of the Female Reproductive Tract in Gynecological and Reproductive Health. *Microbiol Mol Biol Rev*. 2022;86(4):e00181-21. doi:10.1128/mmbr.00181-21

5. Liao J, Shenhav L, Urban JA, et al. Microdiversity of the vaginal microbiome is associated with preterm birth. *Nat Commun*. 2023;14(1):4997. doi:10.1038/s41467-023-40719-7

6. Fettweis JM, Serrano MG, Brooks JP, et al. The vaginal microbiome and preterm birth. *Nat Med*. 2019;25(6):1012-1021. doi:10.1038/s41591-019-0450-2

7. Golob JL, Oskotsky TT, Tang AS, et al. Microbiome preterm birth DREAM challenge: Crowdsourcing machine learning approaches to advance preterm birth research. *Cell Rep Med*. 2023;5(1):101350. doi:10.1016/j.xcrm.2023.101350

8. Qing X, Xie M, Liu P, et al. Correlation between dysbiosis of vaginal microecology and endometriosis: A systematic review and meta-analysis. *PLOS ONE*. 2024;19(7):e0306780. doi:10.1371/journal.pone.0306780

9. MacSharry J, Kovács Z, Xie Y, et al. Endometriosis specific vaginal microbiota links to urine and serum N-glycome. *Sci Rep*. 2024;14(1):25372. doi:10.1038/s41598-024-76125-2

10. Dabee S, Passmore JAS, Heffron R, Jaspan HB. The Complex Link between the Female Genital Microbiota, Genital Infections, and Inflammation. *Infect Immun*. 2021;89(5). doi:10.1128/iai.00487-20

11. Sharifian K, Shoja Z, Jalilvand S. The interplay between human papillomavirus and vaginal microbiota in cervical cancer development. *Virol J*. 2023;20(1):73. doi:10.1186/s12985-023-02037-8

12. Perkins RB, Wentzensen N, Guido RS, Schiffman M. Cervical Cancer Screening: A Review. *JAMA*. 2023;330(6):547-558. doi:10.1001/jama.2023.13174

13. Sung H, Ferlay J, Siegel RL, et al. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin*. 2021;71(3):209-249. doi:10.3322/caac.21660

14. Bhattacharya A, Das S, Bhattacharjee MJ, Mukherjee AK, Khan MR. Comparative pangenomic analysis of predominant human vaginal lactobacilli strains towards population-specific adaptation: understanding the role in sustaining a balanced and healthy vaginal microenvironment. *BMC Genomics*. 2023;24(1):565. doi:10.1186/s12864-023-09665-y

15. Mancabelli L, Mancino W, Lugli GA, et al. Comparative Genome Analyses of Lactobacillus crispatus Isolates from Different Ecological Niches Reveal an Adaptation of This Species to the Human Vaginal Environment. *Appl Environ Microbiol*. 2021;87(8):e02899-20. doi:10.1128/AEM.02899-20

16. Santarelli G, Sanguinetti M, Delogu G, De Maio F. Comparative genomic analysis of bacteriocin genes in Lactobacillus crispatus strains. *Sci Rep*. 2025;15(1):20798. doi:10.1038/s41598-025-08321-7

17. Nori SRC, Walsh CJ, McAuliffe FM, et al. Strain-level variation among vaginal Lactobacillus crispatus and Lactobacillus iners as identified by comparative metagenomics. *Npj Biofilms Microbiomes*. 2025;11(1):49. doi:10.1038/s41522-025-00682-1

18. Pace RM, Chu DM, Prince AL, Ma J, Seferovic MD, Aagaard KM. Complex species and strain ecology of the vaginal microbiome from pregnancy to postpartum and association with preterm birth. *Med N Y N*. 2021;2(9):1027-1049. doi:10.1016/j.medj.2021.06.001

19. Argentini C, Fontana F, Alessandri G, et al. Evaluation of Modulatory Activities of Lactobacillus crispatus Strains in the Context of the Vaginal Microbiota. *Microbiol Spectr*. 2022;10(2):e02733-21. doi:10.1128/spectrum.02733-21

20. Smith BJ, Zhao C, Dubinkina V, et al. Accurate estimation of intraspecific microbial gene content variation in metagenomic data with MIDAS v3 and StrainPGC. *Genome Res*. 2025;35(5):1247-1260. doi:10.1101/gr.279543.124

21. Zhao C, Dimitrov B, Goldman M, Nayfach S, Pollard KS. MIDAS2: Metagenomic Intra-species Diversity Analysis System. *Bioinformatics*. 2023;39(1):btac713. doi:10.1093/bioinformatics/btac713

22. Goldman M, Zhao C, Pollard KS. Improved detection of microbiome-disease associations via population structure-aware generalized linear mixed effects models (microSLAM). *PLOS Comput Biol*. 2025;21(5):e1012277. doi:10.1371/journal.pcbi.1012277

23. Almeida A, Nayfach S, Boland M, et al. A unified catalog of 204,938 reference genomes from the human gut microbiome. *Nat Biotechnol*. 2021;39(1):105-114. doi:10.1038/s41587-020-0603-3

24. Parks DH, Chuvochina M, Rinke C, Mussig AJ, Chaumeil PA, Hugenholtz P. GTDB: an ongoing census of bacterial and archaeal diversity through a phylogenetically consistent, rank normalized and complete genome-based taxonomy. *Nucleic Acids Res*. 2022;50(D1):D785-D794. doi:10.1093/nar/gkab776

25. Huang L, Guo R, Li S, et al. A multi-kingdom collection of 33,804 reference genomes for the human vaginal microbiome. *Nat Microbiol*. 2024;9(8):2185-2200. doi:10.1038/s41564-024-01751-5

26. Cha JH, Kim N, Ma J, et al. A high-quality genomic catalog of the human oral microbiome broadens its phylogeny and clinical insights. *bioRxiv*. Preprint posted online April 21, 2025:2025.03.10.642329. doi:10.1101/2025.03.10.642329

27. Liu Y, Li Z, Zhang C, et al. A human skin microbiome reference catalog and the skin microbial landscape of plateau adults. *iMetaOmics*. 2025;2(2):e70000. doi:10.1002/imo2.70000

28. Ravel J, Gajer P, Abdo Z, et al. Vaginal microbiome of reproductive-age women. *Proc Natl Acad Sci U S A*. 2011;108(Suppl 1):4680-4687. doi:10.1073/pnas.1002611107

29. Norenhag J, Edfeldt G, Stålberg K, et al. Compositional and functional differences of the vaginal microbiota of women with and without cervical dysplasia. *Sci Rep*. 2024;14(1):11183. doi:10.1038/s41598-024-61942-2

30. Jakobsen RR, Haahr T, Humaidan P, et al. Characterization of the Vaginal DNA Virome in Health and Dysbiosis. *Viruses*. 2020;12(10):1143. doi:10.3390/v12101143

31. Gupta P, Singh MP, Goyal K. Diversity of Vaginal Microbiome in Pregnancy: Deciphering the Obscurity. *Front Public Health*. 2020;8. doi:10.3389/fpubh.2020.00326

32. DI Pierro F, Criscuolo AA, Dei Giudici A, et al. Oral administration of Lactobacillus crispatus M247 to papillomavirus-infected women: results of a preliminary, uncontrolled, open trial. *Minerva Obstet Gynecol.* 2021;73(5):621-631. doi:10.23736/S2724-606X.21.04752-7

33. Dellino M, Cascardi E, Laganà AS, et al. Lactobacillus crispatus M247 oral administration: Is it really an effective strategy in the management of papillomavirus-infected women? *Infect Agent Cancer.* 2022;17:53. doi:10.1186/s13027-022-00465-9

34. Taboada B, Estrada K, Ciria R, Merino E. Operon-mapper: a web server for precise operon identification in bacterial and archaeal genomes. *Bioinformatics.* 2018;34(23):4118-4120. doi:10.1093/bioinformatics/bty496

35. Jakobsen RR, Haahr T, Humaidan P, et al. Characterization of the Vaginal DNA Virome in Health and Dysbiosis. *Viruses.* 2020;12(10):1143. doi:10.3390/v12101143

36. Norenhag J, Du J, Olovsson M, Verstraelen H, Engstrand L, Brusselaers N. The vaginal microbiota, human papillomavirus and cervical dysplasia: a systematic review and network meta-analysis. *BJOG Int J Obstet Gynaecol.* 2020;127(2):171-180. doi:10.1111/1471-0528.15854

37. Xia K, Han C, Xu J, Liang X. Toxin-antitoxin HicAB regulates the formation of persister cells responsible for the acid stress resistance in Acetobacter pasteurianus. *Appl Microbiol Biotechnol.* 2021;105(2):725-739. doi:10.1007/s00253-020-11078-w

38. Butt A, Higman VA, Williams C, et al. The HicA toxin from Burkholderia pseudomallei has a role in persister cell formation. *Biochem J.* 2014;459(2):333-344. doi:10.1042/BJ20140073

39. Jørgensen MG, Pandey DP, Jaskolska M, Gerdes K. HicA of Escherichia coli Defines a Novel Family of Translation-Independent mRNA Interferases in Bacteria and Archaea. *J Bacteriol.* 2009;191(4):1191-1199. doi:10.1128/JB.01013-08

40. Gerdes K. Diverse genetic contexts of HicA toxin domains propose a role in anti-phage defense. *mBio.* 2024;15(2):e0329323. doi:10.1128/mbio.03293-23

41. France MT, Chaudry I, Rutt L, et al. VIRGO2: Unveiling the Functional and Ecological Complexity of the Vaginal Microbiome with an Enhanced Non-Redundant Gene Catalog. *bioRxiv.* Published online March 4, 2025:2025.03.04.641479. doi:10.1101/2025.03.04.641479

42. Bruni L, Serrano B, Roura E, et al. Cervical cancer screening programmes and age-specific coverage estimates for 202 countries and territories worldwide: a review and synthetic analysis. *Lancet Glob Health.* 2022;10(8):e1115-e1127. doi:10.1016/S2214-109X(22)00241-8

43. O'Leary NA, Cox E, Holmes JB, et al. Exploring and retrieving sequence and metadata for species across the tree of life with NCBI Datasets. *Sci Data.* 2024;11(1):732. doi:10.1038/s41597-024-03571-y

44. Team G, Riviere M, Pathak S, et al. Gemma 2: Improving Open Language Models at a Practical Size. *arXiv.* Preprint posted online October 2, 2024. doi:10.48550/arXiv.2408.00118

45. Shen W, Ren H. TaxonKit: A practical and efficient NCBI taxonomy toolkit. *J Genet Genomics.* 2021;48(9):844-850. doi:10.1016/j.jgg.2021.03.006

46. Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinforma Oxf Engl.* 2014;30(14):2068-2069. doi:10.1093/bioinformatics/btu153

47. Rognes T, Flouri T, Nichols B, Quince C, Mahé F. VSEARCH: a versatile open source tool for metagenomics. *PeerJ.* 2016;4:e2584. doi:10.7717/peerj.2584

48. Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics*. 2012;28(23):3150-3152. doi:10.1093/bioinformatics/bts565

49. Huerta-Cepas J, Szklarczyk D, Heller D, et al. eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Res*. 2019;47(D1):D309-D314. doi:10.1093/nar/gky1085

50. Florensa AF, Kaas RS, Clausen PTLC, Aytan-Aktug D, Aarestrup FM. ResFinder – an open online resource for identification of antimicrobial resistance genes in next-generation sequencing data and prediction of phenotypes from genotypes. *Microb Genomics*. 2022;8(1):000748. doi:10.1099/mgen.0.000748

51. Camargo AP, Roux S, Schulz F, et al. Identification of mobile genetic elements with geNomad. *Nat Biotechnol*. 2024;42(8):1303-1312. doi:10.1038/s41587-023-01953-y

52. Johansson MHK, Bortolaia V, Tansirichaiya S, Aarestrup FM, Roberts AP, Petersen TN. Detection of mobile genetic elements associated with antibiotic resistance in Salmonella enterica using a newly developed web tool: MobileElementFinder. *J Antimicrob Chemother*. 2021;76(1):101-109. doi:10.1093/jac/dkaa390

53. Minot SS, Barry KC, Kasman C, Golob JL, Willis AD. geneshot: gene-level metagenomics identifies genome islands associated with immunotherapy response. *Genome Biol*. 2021;22(1):135. doi:10.1186/s13059-021-02355-6

54. Krueger F, James F, Ewels P, et al. FelixKrueger/TrimGalore: v0.6.10 - add default decompression path. Published online February 2, 2023. doi:10.5281/zenodo.7598955

55. Li H, Durbin R. Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics*. 2009;25(14):1754-1760. doi:10.1093/bioinformatics/btp324

56. Twelve years of SAMtools and BCFtools | GigaScience | Oxford Academic. Accessed August 27, 2025. https://academic.oup.com/gigascience/article/10/2/giab008/6137722?login=false

57. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J R Stat Soc Ser B Methodol*. 1995;57(1):289-300.

58. Gold EB, Bromberger J, Crawford S, et al. Factors Associated with Age at Natural Menopause in a Multiethnic Sample of Midlife Women. *Am J Epidemiol*. 2001;153(9):865-874. doi:10.1093/aje/153.9.865

59. Efron B, Turnbull B, Narasimhan B, Strimmer K. locfdr: Computes Local False Discovery Rates. Published online July 15, 2015. Accessed August 27, 2025. https://cran.r-project.org/web/packages/locfdr/index.html

60. Camacho C, Coulouris G, Avagyan V, et al. BLAST+: architecture and applications. *BMC Bioinformatics*. 2009;10:421. doi:10.1186/1471-2105-10-421

61. Shimoyama Y. pyGenomeViz: A genome visualization python package for comparative genomics. Published online May 2024. Accessed August 27, 2025. https://github.com/moshi4/pyGenomeViz