Trehalose Transport Dynamics Underpin a Metabolic Trade-Off between

Exogenous Uptake and Endogenous Synthesis in Lepidopteran Insects

Bhagyashri Y. Chaudhari[1, 2*] , Vikram J. Nichit[1, 2*], Vitthal T. Barvkar[3], Rakesh

Shamsunder Joshi[1, 2*]

[1]Biochemical Sciences Division, CSIR-National Chemical Laboratory, Dr. Homi

Bhabha Road, Pune, 411008, Maharashtra, India

[2]Academy of Scientific and Innovative Research (AcSIR), Ghaziabad 201002, Uttar

Pradesh, India

[3]Department of Botany, Savitribai Phule Pune University, Pune, 411007, Maharashtra,

India

***Corresponding author**

Dr. Rakesh Shamsunder Joshi

Biochemical Sciences Division, CSIR – National Chemical Laboratory

Dr. Homi Bhabha Road, Pune – 411 008, INDIA

Email: rs.joshi@ncl.res.in

Tel (office): 020-2590-2247

Abstract

Trehalose is the major insect hemolymph sugar and plays a diverse role. Its level is regulated endogenously by the dynamics of biosynthesis and distribution by sugar transporters (STs). The metabolic trade-off between trehalose synthesis and uptake remains poorly understood, despite its critical role in homeostasis. Here, we examined the role of a gut-specific trehalose transporter, *HaST46*, in regulating this metabolic trade-off in *Helicoverpa armigera*, a Lepidopteran pest model. Integrated transcriptomics analysis and functional analyses revealed that *HaST46* acts as a diet-responsive transporter, localised to the posterior midgut, with trehalose preference. Its expression is modulated in response to dietary trehalose availability, enhancing the efficient exogenous trehalose uptake while attenuating its endogenous synthesis and conserving energy. Functional perturbation through overexpression and silencing revealed a feedback-regulated mechanism in which *HaST46* expression showed strong correlation with trehalose metabolising enzymes and other HaSTs isoforms to maintain systemic trehalose homeostasis. Overall, our findings reveal a metabolic trade-off between exogenous trehalose uptake and endogenous synthesis mediated by gut-specific sugar transporters.

Keywords: Trehalose, Sugar uptake, Metabolism, Transporters, *Helicoverpa armigera*, Gut

1. Introduction

Insects exhibit a remarkable metabolic adaptation that contributes to their ecological success. Carbohydrates serve as a primary energy source for fuelling flight, development, reproduction, diapause, and stress tolerance (Thorat et al. 2012; Feofilova et al. 2014). Under stress conditions, insects must precisely regulate their energy acquisition, storage, and mobilisation (Zhang et al. 2019). Trehalose is the predominant hemolymph sugar in insects, fulfilling both structural and metabolic functions (Becker et al. 1996; Matsushita and Nishimura 2020; Nishimura 2020; Murphy and Wyatt 1965; Shukla et al. 2015). Its transport and metabolism are essential for insect growth and adaptability (Tellis, Kotkar, et al. 2023). Trehalose is primarily synthesised in the fat body through a conserved pathway involving *trehalose-6-phosphate synthase* (*TPS*) and *trehalose-6-phosphate phosphatase* (*TPP*) (Murphy and Wyatt 1965; Becker et al. 1996). Following synthesis, trehalose is transported into the hemolymph and then distributed to the target tissues as per the energy demand through trehalose transporters (TRETs) (Tellis, Chaudhari, et al. 2023).

Insects can ingest a wide range of foods to obtain complementary nutrients during nutritional scarcity (Koyama et al. 2020; Raubenheimer and Jones 2006). They further optimise energy utilization through metabolic balancing, enabling adaptation to diverse dietary conditions (Smith et al. 2018; Chown and Gaston 1999). Trehalose synthesis requires a continuous supply of ATP and carbon skeletons from glycolysis, glycogenolysis, or gluconeogenesis (Miyamoto and Amrein 2017). Depending on resource availability, resource insects may shift their metabolism to suppress endogenous synthesis and prioritise exogenous uptake (Izadi 2025). Hemolymph trehalose levels range from 5 to 50mM, which is maintained by continuous exchange

via TRETs for effective utilisation (Thompson 2003; Al Hinai et al. 2025; Wyatt 1967;

Kikawada et al. 2007). TRETs are key in regulating metabolic trade-offs under

controlled and stressed conditions (Leyria et al. 2020). The midgut is the primary site

of sugar and nutrient absorption, detoxification, and enzymatic activity (Huang et al.

2015). To meet the metabolic demand, insects rely on sugar transporters such as

TRETs, sodium-glucose co-transporters (SGLTs), and facilitative glucose transporters

(GLUTs) to manage dietary sugar acquisition and use (Chaudhari et al. 2025).

Phytophagous insects such as *H. armigera* can acquire trehalose directly from

their diet (Shukla et al. 2015; Tellis, Kotkar, et al. 2023). In *H. armigera*, trehalose-

specific sugar transporters (*HaSTs*) showed diversity and spatio-temporal expression

patterns (Tellis, Chaudhari, et al. 2023). In dipteran and hymenopteran insects,

glucose absorption is primarily mediated by SGLTs. and GLUTs (Wang et al. 2016).

In *Drosophila melanogaster,* the diet-responsive GLUT homolog CG4797 and SGLT

homolog dSGLT1 have been shown to influence development, fertility, and survival

(Chaudhari et al. 2025). The activity of these transporters is closely linked to the insulin

and TOR signalling pathways, facilitating metabolic plasticity by coupling nutrient

availability with growth regulation (Riera et al. 2016). This critical trait allows insects to

cope with environmental stressors, shift between trophic modes, and optimise energy

expenditure during metamorphosis or overwintering (Danks 2007). The interplay

between trehalose metabolism and other physiological systems, including

neuroendocrine signalling, symbiotic relationships, and hormonal regulation, facilitates

nutrient sensing, energy homeostasis, and ecological adaptability (Tellis, Kotkar, et al.

2023).

Despite advances in understanding trehalose metabolism within insects, the

integrated dynamics between endogenous trehalose synthesis and exogenous uptake

remain poorly understood in many model and non-model organisms. Lepidopteran

pests such as *H. armigera* serve as valuable models for metabolic studies due to their

high dietary adaptability, rapid development, and stress resilience (Riaz et al. 2021).

Elucidating how these insects regulate trehalose metabolism in response to diet offers

insights into the metabolic homeostasis and its impact on physiological processes

(Dawkar et al. 2016;,Guan et al. 2024;, Tian et al. 2025;, Zhou et al. 2023). Here, we

shed light on the functional role of the trehalose transporter, *HaST46,* in mediating the

metabolic trade-off between trehalose's exogenous uptake and endogenous

synthesis. Through differential gene expression analysis, qRT-PCR, LC-MS-based

metabolomics, enzyme activity assays, and genetic manipulation, we investigated how

dietary trehalose influences *HaSTs* dynamics and activity, mainly in the gut. This study

advances the knowledge of sugar transport, metabolic homeostasis and nutritional

physiology.

2. Results

2.1. *HaST46* showed posterior midgut-specific expression at the transcript and

protein levels

We analyzed the gut-specific expression and proteomic presence of selected putative

trehalose transporters, *HaSTs*, using publicly available *H. armigera* transcriptomic and

proteomic datasets (Ioannidis et al. 2022). Our previous study elucidated five putative

trehalose transporters with selective abundance in specific tissues, among which

HaST46 and *HaST69* were found to be present in gut-specific transcriptomic data

(Figure 1a). While only *HaST46* showed expression in proteomic data, indicating its

proteomic level abundance in the mid-gut **(Figure 1b)**. In the transcriptomics dataset,

HaST46 and *HaST69* showed high expression levels across larval stages (L2, L3, L4)

and in different gut regions, including foregut (FG), anterior midgut (AMG), middle

midgut (MMG), posterior midgut (PMG), and hindgut (HG) in fifth instar (L5) larvae.

HaST46 expression peaked in the gut during the L4 larval stage and in the PMG of L5

larvae **(Figure 1a)**. Given that feeding activity peaks during the fourth and fifth larval

instars, elevated expression of the sugar transporter likely supports increased

metabolic demand. Collectively, these data highlight that *HaST46* is a putative key

facilitator of sugar transport across the PMG during the energetically demanding

developmental stages and the period of active feeding. Transcriptomics comparisons

were conducted between the gut compartments of L5 larvae fed an artificial diet (AD)

versus those fed a plant diet (PD)-cotton to assess the diet-dependent expression

dynamics. Interestingly, *HaST46* expression was elevated exclusively in the PMG of

L5 larvae fed a plant diet (PD) compared to AD, whereas *HaST69* exhibited the

opposite trend **(Figure 1c-d)**. The presence of HaST46 in the PMG, which is a primary

site for nutrient absorption during active feeding, suggests its putative role in sugar

absorption from feed. The selective higher expression of *HaST46* in larvae consuming

a plant diet supports its involvement in sugar assimilation from plant (complex) food

materials. This further suggests that the dietary composition possibly compensates for

the metabolic demand.

2.2. HaST46 showed a preference toward trehalose transport

Functional assays with 10mM trehalose exposure to Sf9 cells overexpressing HaST46

revealed a significant increase in intracellular sugar accumulation in *HaST46*-

expressing cells compared with the vector control **(Figure 1e and f)**. LC-MS based

quantification revealed that *HaST46*-expressing cells accumulated ~1.5-fold more

trehalose relative to controls, while glucose uptake exhibited a moderate increase

(~1.2-fold), suggesting a trehalose preference of HaST46. Furthermore, the structural

alignment of HaST46 with the human GLUT1 transporter (PDB: 4PYP) showed a high

degree of conservation in transmembrane domain architecture **(Figure 1g)**. Molecular

docking studies further revealed that CYS416 and ASN312 of HaST46 formed

conserved hydrogen bonds with both glucose and trehalose. At the same time,

TRP420 and ALA193 contributed additional interactions that stabilized trehalose

binding **(Figure 1g)**. These findings support the hypothesis that HaST46 functions as

a trehalose-preferred transporter, likely contributing to sugar uptake and homeostasis

in *H. armigera.*

2.3. Exogenous trehalose feeding upregulates *HaST46* expression and trehalose

catabolism genes in *H. armigera*

Upon exogenous trehalose feeding, we observed optimal growth and fitness in the

case of 50mM trehalose containing diet. In comparison to this, feeding with 10mM

trehalose doesn't showed any noteworthy physiological impact, and 100 mM trehalose

lead to sugar-associated toxicity **(Supplementary File S5, Figures S1 and S2)**.

Feeding of 50mM trehalose induced significant changes in the expression of trehalose

transporters (TRETs) and metabolic genes as revealed by qRT-PCR analysis.

HaST46 expression was upregulated upon trehalose feeding **(Figure 2a)**. In response

to this, *HaTPS/TPP* were downregulated, while *HaTreh-1* showed upregulation upon

feeding a diet with 50mM trehalose **(Figure 2b-c)**. This alteration in the *HaTPS/TPP*

expression level suggests a putative reduction in the energy-intensive endogenous

trehalose biosynthesis and increased hydrolysis of exogenous trehalose for energy

needs. Notably, *HaST69* was downregulated on a 50mM TD, indicating that it is

potentially not involved in trehalose absorption from the gut **(Supplementary File S5,**

Figure S3). Residual TPP activity was slightly reduced, while trehalase activity was

high in larvae fed the 50mM TD ($p < 0.01$) **(Figure 2d-e)**. Metabolomic quantification

showed that the insect fed on the trehalose-containing diet showed relatively more

accumulation compared to the control **(Figure 2f)**. Furthermore, increased

accumulation of ATP and low ADP/AMP indicates little demand for ATP hydrolysis in

trehalose fed insects **(Figure 2f)**.This evidence suggests a preference for hydrolytic

metabolism under elevated dietary trehalose, which is likely to convert absorbed

trehalose into glucose for systemic energy needs.

2.4. *HaST46* showed gut-specific upregulation in response to trehalose feeding

Tissue-specific gene expression upon 50mM trehalose feeding was further analyzed

to elucidate the dynamics of trehalose transport. The relative expression of *HaST46*

and *HaTPP* was reduced in hemolymph **(Figure 3a-b)**, while the expression of

HaTreh1 and *HaTreh2* was nearly unchanged **(Figure 3c-d)**. The gut showed a

significant ~3-fold increase in *HaST46* expression under a 50mM TD **(Figure 3e)**,

indicating an increase in the need to transport and use excess trehalose in the gut

tissue. As the midgut is the primary site for transporter-mediated dietary trehalose

uptake, *HaST46* overexpression was found to be relevant. *HaTPS/TPP* expression

was found to be unchanged, while *HaTreh1* and *HaTreh2* expression were

significantly high **(Figure 3f-h)**. *HaST46* expression in the fat body (key tissue for

trehalose synthesis) was found to be unchanged **(Figure 3i)**. Interestingly, the

expression of *HaTPS/TPP* decreased significantly, suggesting a reduced endogenous

trehalose biosynthesis **(Figure 3j)**. *HaTreh1* and *HaTreh2* expression were relatively

high in the fat body of trehalose-fed insects **(Figure 3k-l)**. These observation suggests

that dietary trehalose downregulates internal trehalose synthesis through feedback

inhibition.

2.5. Metabolomic Profiling Reveals Trehalose Trade-off

LC-MS analysis was performed for hemolymph of insects fed on artificial diet (AD) and

a trehalose-rich diet (TD) to quantify systemic metabolic changes. TD-fed larvae

exhibited relatively elevated levels of trehalose, particularly in the hemolymph

(Supplementary File_S5, Figure S5). In addition, downstream glycolytic

intermediates such as glucose-6-phosphate and pyruvate were elevated, supporting

the notion that absorbed trehalose is rapidly catabolized into usable energy. These

results confirm that trehalose supplementation alters gene expression and shifts the

metabolic landscape toward catabolic flux, optimizing energy production in the

presence of abundant dietary sugar in *H. armigera.*

2.6. *HaST46* knockdown induces compensatory upregulation of the trehalose

anabolism TPS*/TPP* gene

RNA interference (RNAi) was employed to silence gut-prevalent *HaST46* expression.

We achieved ~80% reduction in the *HaST46* transcripts levels compared with the

empty vector **(Figure 4a)**. Notably, insects with *HaST46* silencing that were fed a 50

mM TD exhibited a compensatory increase in *HaST46* expression, suggesting a

feedback mechanism and highlighting its critical role in trehalose absorption from the

gut **(Figure 4b)**. Furthermore, trehalase activity was elevated in TD-fed insects

compared with AD-fed insects **(Figure 4c-d)**. LC-MS analysis revealed a decrease in

trehalose levels in the insect body upon silencing *HaST46*, indicating putative

involvement in the dietary trehalose absorption and resulting in an overall decrease in

endogenous trehalose content **(Figure. 4e)**. However, upon trehalose feeding to *dsHaST46* insects, the trehalose level was almost the same as in the control **(Figure 4f)**. These findings highlight the potential role of *HaST46 in* trehalose transport. The impact of *HaST46* silencing on nutritional efficiency was assessed by measuring nutritional indices in both the control and silenced groups. Nutritional indices were almost the same in both groups, such as the control and *HaST46* silenced insects **(Supplementary File_S5, Figure S8-9)**.

The expression of other trehalose transporter genes (*HaST09*, *HaST29*, *HaST64*, and *HaST69*) was examined in *dsHaST46*-fed larvae to explore potential compensatory mechanisms. Silencing *HaST46* resulted in significant downregulation of these genes, suggesting overall impact due to reduced trehalose content upon silencing (Supplementary File_S5, Figure 12a). *HaST09*, *HaST29*, *HaST64*, and *HaST69* exhibited upregulation in exogenous trehalose, although the functional significance of these changes requires further investigation (Supplementary File_S5, Figure 12b). These results underscore the pivotal role of *HaST46* in modulating trehalose transport and metabolism.

2.7. *HaST46* overexpression showed gut specificity and may enhance dietary trehalose

Transient overexpression of *HaST46* was confirmed with qRT-PCR analysis, showing ~1000-fold higher expression of *HaST46* in treated larvae compared with the control **(Figure 5a)**. *HaST46* overexpression in larvae did not significantly alter trehalase activity, suggesting a maintained balance between trehalose synthesis and absorption to preserve homeostasis **(Figure 5b)**. Further, we observed the upregulation of *HaST46* in all three tissues of the *HaST46-OE* insects upon trehalose feeding

compared with the control group. Interestingly, there was a significant upregulation of

HaST46 in the gut, indicating the potential preferred localization of overexpressed

HaST46 in the gut tissue **(Figure 5c-e)**. However, there was a significant increase in

the *HaST46* transcript level in a group of insects fed on a trehalose-rich diet in the gut,

followed by hemolymph and fat **(Figure 5c-e)**. Trehalose levels were slightly high in

TD-fed *HaST46*_OE insects **(Figure 5f-g)**. At the same time, trehalase activity was

also found to be high in TD-fed *HaST46*_OE insects **(Figure 5h)**. These results

suggest that *HaST46* facilitates the uptake of dietary trehalose and reinforces its role

in sugar homeostasis and energy balance in Lepidopterans.

3. Discussion

Trehalose is key in energy homeostasis, functioning as a circulating sugar and

substrate for rapid energy mobilization. The identification and functional

characterization of *HaST46* as a trehalose-specific transporter localized to the

posterior midgut, principal site for nutrient uptake, highlights its putative function in

dietary sugar absorption **(Figure 1a and b)** (Terra et al. 2023). Furthermore, the

nutritionally responsive expression patterns of *HaST46* suggest that its expression is

coupled to the diet composition and larval metabolic demand. The higher expression

of *HaST46* in complex or plant diet reflects an adaptive shift to facilitate the uptake of

structurally and compositionally diverse carbohydrates in natural food sources **(Figure**

1c and d). This adaptive plasticity aligns with earlier reports of the gut-specific

modulation of nutrient transporters in response to dietary sugars in lepidopteran larvae

(Azzouz et al. 2005), underscoring the ecological and physiological relevance of

transporter regulation in herbivorous pests.

The substrate preference of HaST46 for trehalose over glucose represents a

functional specialization **(Figure 1e and f)**. Despite structural conservation with

GLUT1, HaST46 exhibits trehalose preference over glucose, suggesting an

evolutionary divergence toward the hemolymph sugar composition of holometabolous

insects **(Figure 1g)**. This specificity likely enhances transport efficiency and minimizes

competitive inhibition, optimizing trehalose handling in sugar-rich dietary contexts.

At the metabolic level, the changes in trehalose biosynthesis and degradation

pathways in response to transporter activity reflects a coordinated homeostatic

system. The downregulation of endogenous trehalose synthesis in the presence of

abundant dietary trehalose suggests a resource-efficient switch that curtails ATP-

intensive biosynthesis in favour of direct absorption **(Figure 2a to e)**. Increased level

of ATP compared to low level of ADP/AMP suggest low demand of ATP hydrolysis in

exogenous supply of trehalose **(Figure 2f)**. This trade-off supports the notion that

insects prioritize the least metabolically costly route to meet carbohydrate

requirements, a trait advantageous for survival in fluctuating nutritional environments.

Furthermore, systemic metabolic shifts, evidenced by increased glycolytic

intermediates in trehalose-fed larvae, indicate that transporter-mediated uptake

directly fuels energy metabolism, facilitating rapid developmental transitions. This

catabolic reprogramming toward glucose-generating pathways mirrors the metabolic

flux observed during feeding stages in other insects and provides a mechanistic

explanation for the physiological benefits of dietary trehalose.

RNAi-mediated silencing reduces *HaST46* transcript abundance and triggers

compensatory responses at both transcriptomic and enzymatic levels **(Figure 4a and**

b). Trehalose biosynthetic genes upregulation on *HaST46* knockdown implies

compensation for decreased absorption by enhancing endogenous production.

However, the simultaneous downregulation of other transporter isoforms suggests a

response to reduced overall trehalose levels. Notably, despite transporter knockdown,

nutritional indices remained relatively unaffected, indicating that *H. armigera* maintains

systemic metabolic resilience, likely through alternative sugar transporters or

compensatory physiological mechanisms. Conversely, *HaST46* overexpression did

not disrupt homeostasis, suggesting that while the transporter enhances trehalose

uptake, the system possesses regulatory checkpoints that prevent metabolic overload

(Figure 5a and b).

Together, these findings highlight role of *HaST46* in trehalose acquisition and

regulation. Its dynamic expression, functional specificity, and involvement in feedback-

controlled metabolic shifts position it as a key mediator of dietary sugar assimilation.

Given that trehalose metabolism is unique to insects and absent in vertebrates,

HaST46 offers a promising molecular target for the selective disruption of energy

metabolism in lepidopteran pests.

4. Materials and Methods

4.1 Differential gene expression and proteomic signature analysis

Publicly available *H. armigera* gut transcriptomic and proteomic datasets (Ioannidis et

al. 2022) were used for expression profiling of *HaSTs*. Out of 5 tissue specific *HaSTs*,

only *HaST46* and *HaST6*9 were mapped to gut transcriptomics and proteomics. For

transcriptomics, raw fragments per kilobase of transcript per million mapped reads

(FPKM) values were extracted, $\log_2(x+1)$ transformed, and row-wise Z-score

normalized before hierarchical clustering using Euclidean distance and complete

linkage in the pheatmap package (R v4.x). For proteomics, the absence or presence

of HaSTs in the proteomics set was denoted in binary form. Heatmaps were generated

using TBtools 1.09 (Chen et al. 2020) to visualise stage- and tissue-specific

expression patterns. For bar plots, mean FPKM values ± SEM were calculated across

biological replicates.

4.2 *HaST46* overexpression in Sf9 cells

For overexpression, *HaST46* was cloned into the pIBV5 vector and sequence-

confirmed. Sf9 cells   (4×10^6) were seeded in a T25 cell culture flask (Corning, NY,

USA) with 1 mL serum-free SF900 II SFM media (Thermo Scientific, Waltham, MA,

USA) and allowed to adhere by incubating for 30 minutes at 27 °C. For transfection,

1.5 µg of either the EGFP_pIBV5 control plasmid or the *HaST46-pIBV5* plasmid was

diluted in 100 µL SF-900 II SFM (without antibiotics). Separately, 8µL Cellfectin II

reagent (Gibco, Waltham, MA, USA) was diluted in 100 µL Sf-900 II SFM and

incubated for 30 minutes. The DNA and lipid solutions were combined (total 200 µL)

and incubated for an additional 30 min to allow complex formation. The transfection

mix was added dropwise while gently rocking the flask for an even distribution. Flasks

were incubated for 5-7 hours at 27°C. After the incubation, the media was removed,

and fresh Sf-900 II SFM containing blasticidin (Gibco, Waltham, MA, USA) antibiotic

was added and incubated for 48 hours. Overexpression was confirmed by qRT-PCR.

For the sugar uptake assay, 5mM and 10mM concentrations of glucose and trehalose

were added to the media, and the cells were incubated for varying time points (0, 5,

and 10 minutes). Following incubation, the media and cells were collected for

metabolite extraction and analysis. Sugar uptake was determined from changes in

extracellular and intracellular sugar concentrations by metabolite profiling.

4.3 Structural modelling and docking

Three dimensional structure of HaST46 was predicted using Alphafold 3 server (https://alphafoldserver.com/). Structural aligment of best model was performed using PyMol (The PyMOL Molecular Graphics System, Version 3.0 Schrödinger, LLC.) using Human Glut1 (PDB ID: 4PYP) structure as reference. Binding site residues of Glut1 were used to predict the binding pocket of HaST46 and molecular docking of glucose and trehalose was performed using AutoDock Tool as described earlier (Barbole et al. 2024).

4.4 Quantitative Real-Time PCR (qRT-PCR) Analysis

Total RNA was extracted from 80–100 mg of insect cells/tissues using the TRIzol reagent (Invitrogen, Waltham, MA, USA). To eliminate DNA contamination, RNA was treated with RQ1 RNase-free DNase. First-strand cDNA was synthesised from 5 µg RNA using oligo-dT primers and the High-Capacity cDNA Reverse Transcription Kit (Applied BioSystem, Foster, CA, USA), following the manufacturer's protocol. Gene expression was quantified using a 7500 Fast Real-Time PCR System (Applied Biosystems, Foster, CA, USA) with Takara TB Green Premix Ex Taq II. Primer list provided in supplementary information (S3). Expression dynamics, including transcript abundance (2-ΔCt) and fold change (2-ΔΔCt), were visualised using GraphPad software. The expression of five *HaSTs*, *trehalose-6-phosphate synthase* (*TPS*), *trehalose-6-phosphate phosphatase* (*TPP*), and trehalase was assessed in trehalose-fed tissues, with the gut as the reference for fold change in fifth instar larvae. *HaSTs* expression was further evaluated in haemolymph, fat body, and silencing assays, using control expression as the reference.

4.5 Metabolite Analysis Using LC-Orbitrap-MS

To extract metabolites, 80-100mg of finely crushed sample powder was mixed with 500 µL of 80% MS-grade methanol to ensure efficient metabolite extraction (J.T. Baker, Center Valley, PA, USA). Methods were carried out according to the procedures described in the supplementary method 4.

4.6 Insect rearing

Larval populations of *H. armigera* were obtained from the ICAR-National Bureau of Agricultural Insect Resources (NBAIR), Bengaluru, India, and maintained on a chickpea-based artificial diet (AD) for multiple generations. Insects were reared at 25 ± 1°C, 70 ± 10% relative humidity, and a 16:8 h light: dark photoperiod. Adult mating boxes, containing equal numbers of males and females, were provided with 10% (w/v) sucrose solution and 1% (w/v) vitamin E, with a muslin cloth for egg collection (Dawkar et al. 2016).

4.7 Exogenous trehalose feeding and tissue collection for the bioassay

For exogenous trehalose feeding in the *H. armigera* bioassay, freshly moulted second instar larvae were fed AD supplemented with 10mM, 50mM, or 100mM trehalose for 6 days to assess the impact on nutritional indices efficiency of conversion of ingested food (ECI), efficiency of conversion of digested food (ECD), and approximate digestibility (AD) and expression of sugar transporter genes (*HaSTs*). Three biological replicates (n = 10 larvae each) were used, and the samples were flash-frozen in liquid nitrogen and stored at -80°C. For tissue collection, fifth instar larvae were starved for 1-2 hours followed by washing with phosphate-buffered water and dissected under RNase-free conditions to collect haemolymph, gut tissues (foregut, midgut, hindgut), and fat body. The samples were flash-frozen and stored at -80°C.

4.8 In vitro Targeted Putative Sugar Transporter Gene *HaST46* Silencing via dsRNA Feeding

Silencing of *the HaST46* gene was performed using microbial-based dsRNA production. To generate dsRNA targeting *HaST46*, a 400-bp fragment was PCR amplified using Phusion DNA Polymerase (Thermo Fisher Scientific, Waltham, MA, USA) and a Proflex PCR machine (Thermo Fisher Scientific, Waltham, MA, USA) with primers (**Supplementary File S3**). Detailed methods are provided in **Supplementary method 2**.

4.9 Transient overexpression of *HaST46*

For overexpression, *HaST46* was cloned into the pIB/V5 vector (Thermo Fisher Scientific, Waltham, MA, USA) and sequence-verified. Larvae were starved for 3 h before injection. Further process were carried out according to the procedures described in the supplementary method 3. A mixture of 1500 ng of each plasmid and lipofectamine (Invitrogen, Thermo Fisher Scientific Inc., Waltham, MA, USA) (1:1 v/v), incubated for 30 minutes, was injected into the haemocoel between the 5th and 7th abdominal segments. The EGFP pIB/V5 plasmid served as the control (Barbole et al. 2024).

4.10 Enzymatic activity

The α,α-trehalase activity was determined by measuring the amount of glucose released from the hydrolysis of α,α-trehalose using the dinitrosalicylic acid (DNSA) reagent (Sigma-Aldrich, Merck KGaA, St. Louis, MO, USA). By monitoring the release of inorganic phosphate (Pi) from the crude extract using the malachite-green reagent (Sigma-Aldrich, St. Louis, MO, USA) to assess the enzymatic activity of trehalose 6-

phosphate phosphatase (TPP). The assay was conducted following the modified

protocol of (Klutts et al. 2003). Methods were performed as described in

Supplementary method 4b.

4.11 Statistical analysis

All experiments contain 3 or more biological replicates. Data were expressed as mean

± SE using GraphPad Prism v8.0 (GraphPad Software, San Diego, CA, USA). The

unpaired Student's t-test. Asterisks indicate significant changes compared to control

(*P-value < 0.05; **P-value < 0.01; ***P-value < 0.001, ****P-value < 0.0001).

Declaration of competing interests

The authors have no competing interests of a financial or personal nature.

Acknowledgments

Fellowship support from CSIR-UGC-NET to BYC and VJN is acknowledged. ANRF

(GAP336726) partially supports this work.

Contributions

BYC designed and conducted the experiments, performed data curation, formal

analysis, validation, and visualisation, and wrote and edited the original draft. VJN

performed only transfection in the Sf9 cell line, whereas cloning, qRT, LC-MS sample

preparation, and data analysis were performed by BYC. VTB carried out LC-MS

experiments, and RSJ performed the analysis. RSJ conceptualisation, experiment

design, supervision, provision of resources, funding acquisition, investigation,

visualisation, docking, and methodology. BYC, VTB, and RSJ contributed to writing

and revising the manuscript and approved the final version.

Data availability statement

Data are available in the figures of the article and its supplementary materials.

Conflict of interest

The authors declare that they have no conflicts of interest.

References:

Feofilova EP, Usov AI, Mysyakina IS, Kochkina GA. 2014. Trehalose: Chemical

structure, biological functions, and practical application. Microbiology. 83:184–194

Thorat LJ, Gaikwad SM, Nath BB. 2012. Trehalose as an indicator of desiccation

stress in Drosophila melanogaster larvae: a potential marker of anhydrobiosis.

Biochem Biophys Res Commun. 419(4):638–642

Zhang D-W et al. 2019. Insect behavior and physiological adaptation mechanisms

under starvation stress. Front Physiol. 10:163

Becker A, Schlöder P, Steele JE, Wegener G. 1996. The regulation of trehalose

metabolism in insects. Experientia. 52:433–439

Matsushita R, Nishimura T. 2020. Trehalose metabolism confers developmental

robustness and stability in Drosophila by regulating glucose homeostasis. Commun

Biol. 3(1):170

Nishimura T. 2020. Feedforward regulation of glucose metabolism by steroid

hormones drives a developmental transition in Drosophila. Curr Biol. 30(18):3624–

3632

Murphy TA, Wyatt GR. 1965. The enzymes of glycogen and trehalose synthesis in

silk moth fat body. J Biol Chem. 240(4):1500–1508

Shukla E, Thorat LJ, Nath BB, Gaikwad SM. 2015. Insect trehalase: physiological

significance and potential applications. Glycobiology. 25(4):357–367

Tellis MB, Kotkar HM, Joshi RS. 2023. Regulation of trehalose metabolism in

insects: from genes to the metabolite window. Glycobiology. 33(4):262–273

Tellis MB, Chaudhari BY, et al. 2023. Trehalose transporter-like gene diversity and
dynamics enhances stress response and recovery in Helicoverpa armigera. Gene.
862:147259

Koyama T, Texada MJ, Halberg KA, Rewitz K. 2020. Metabolism and growth
adaptation to environmental conditions in Drosophila. Cell Mol Life Sci. 77(22):4523–
4551

Raubenheimer D, Jones SA. 2006. Nutritional imbalance in an extreme generalist
omnivore: tolerance and recovery through complementary food selection. Anim
Behav. 71(6):1253–1262

Smith RL, Soeters MR, Wüst RCI, Houtkooper RH. 2018. Metabolic flexibility as an
adaptation to energy resources and requirements in health and disease. Endocr
Rev. 39(4):489–517

Chown SL, Gaston KJ. 1999. Exploring links between physiology and ecology at
macro-scales: the role of respiratory metabolism in insects. Biol Rev. 74(1):87–120

Miyamoto T, Amrein H. 2017. Gluconeogenesis: An ancient biochemical pathway
with a new twist. Fly (Austin). 11(3):218–223

Izadi H. 2025. Endocrine and enzymatic shifts during insect diapause: a review of
regulatory mechanisms. Front Physiol. 16:1544198

Thompson SN. 2003. Trehalose—the insect 'blood' sugar. Adv In Insect Phys.
31(Supplement C):205–285

Al Hinai MS, Rehman A, Siddique KHM, Farooq M. 2025. The Role of Trehalose in

Improving Drought Tolerance in Wheat. J Agron Crop Sci. 211(3):e70053

Kikawada T et al. 2007. Trehalose transporter 1, a facilitated and high-capacity

trehalose transporter, allows exogenous trehalose uptake into cells. Proc Natl Acad

Sci U S A. 104(28):11585–11590. https://doi.org/10.1073/pnas.0702538104

Wyatt GR. 1967. The Biochemistry of Sugars and Polysaccharides in Insects. Adv In

Insect Phys. 4(C):287–360. https://doi.org/10.1016/S0065-2806(08)60210-6

Leyria J, Orchard I, Lange AB. 2020. What happens after a blood meal? A

transcriptome analysis of the main tissues involved in egg production in Rhodnius

prolixus, an insect vector of Chagas disease. PLoS Negl Trop Dis. 14(10):e0008516

Huang J-H, Jing X, Douglas AE. 2015. The multi-tasking gut epithelium of insects.

Insect Biochem Mol Biol. 67:15–20

Chaudhari BY, Pradhan AG, Joshi RS. 2025. Metabolic gatekeepers: Dynamic roles

of sugar transporters in insect metabolism and physiology. Insect Mol Biol. 34(1):1–

18

Wang X, Li E, Chen L. 2016. A review of carbohydrate nutrition and metabolism in

crustaceans. N Am J Aquac. 78(2):178–187

Riera CE, Merkwirth C, De Magalhaes Filho CD, Dillin A. 2016. Signaling networks

determining life span. Annu Rev Biochem. 85(1):35–64

Danks H V. 2007. The elements of seasonal adaptations in insects. Can Entomol.

139(1):1–44

Riaz S et al. 2021. A review on biological interactions and management of the cotton

bollworm, Helicoverpa armigera (Lepidoptera: Noctuidae). J Appl Entomol.

145(6):467–498

Dawkar V V. et al. 2016. The expression of proteins involved in digestion and

detoxification are regulated in Helicoverpa armigera to cope up with chlorpyrifos

insecticide. Insect Sci. 23(1):68–77. https://doi.org/10.1111/1744-7917.12177

Guan L et al. 2024. The Role of TcCYP6K1 and TcCYP9F2 influences trehalose

metabolism under high-CO_2 stress in Tribolium castaneum (Coleoptera). Insects.

15(7):502

Tian Z et al. 2025. Trehalose metabolism mediates trade-offs between reproduction

and survival in beet webworm, Loxostege sticticalis, under heat stress. Pest Manag

Sci. 81(2):903–911

Zhou Y-F et al. 2023. Characterization of the TcCYPE2 gene and its role in

regulating trehalose metabolism in response to high CO_2 stress. Agronomy.

13(9):2263

Ioannidis P et al. 2022. A spatiotemporal atlas of the lepidopteran pest Helicoverpa

armigera midgut provides insights into nutrient processing and pH regulation. BMC

Genomics. 23(1):75

Barbole RS et al. 2024. Chitinase inhibition induces transcriptional dysregulation

altering ecdysteroid-mediated control of Spodoptera frugiperda development.

Iscience. 27(3)

Terra WR, Ferreira C, Silva CP. 2023. Overview of insect midgut function. In:

Molecular physiology and evolution of insect digestive systems. Springer; p 13–26

Azzouz H et al. 2005. Effects of plant protease inhibitors, oryzacystatin I and

soybean Bowman–Birk inhibitor, on the aphid Macrosiphum euphorbiae (Homoptera,

Aphididae) and its parasitoid Aphelinus abdominalis (Hymenoptera, Aphelinidae). J

Insect Physiol. 51(1):75–86

Chen C et al. 2020. TBtools: an integrative toolkit developed for interactive analyses

of big biological data. Mol Plant. 13(8):1194–1202

Klutts S et al. 2003. Purification, cloning, expression, and properties of mycobacterial

trehalose-phosphate phosphatase. J Biol Chem. 278(4):2093–2100

Figure Legends

Figure 1. Spatial expression and functional characterisation of *HaST46* (a) Pro-

tein abundance across larval stages (L2, L3, L4) and gut regions (FG: foregut; AMG:

anterior midgut; MMG: median midgut; PMG: posterior midgut; HG: hindgut) in L5 lar-

vae reared on an artificial diet. (b) DGE patterns comparing larvae fed on an artificial

diet and a plant-based diet (cotton) across different gut compartments.

(c, d) Expression levels of *HaST46* and *HaST69* in the posterior midgut under the two

dietary regimes. Data represent mean ± SEM of biological replicates. Statistical signif-

icance was determined using an unpaired *t*-test ($P < 0.05$). (e) Verification of *HaST46*

overexpression by qRT-PCR. (f) Quantification of D-glucose and trehalose in cell su-

pernatants and lysates using LC–MS/MS after supplementation of the culture medium

with glucose (5 mM, 10 mM) or trehalose (5mM, 10mM) at 0, 5, and 10 min intervals.

All data are presented as mean ± SEM. Statistical analysis was performed using an

unpaired *t*-test ($P < 0.05$). (g) Structural conservation of HaST46 with human GLUT1

transporter. Molecular interaction of HaST46 with glucose and trehalose.

Figure 2. Gene expression and enzymatic activity upon exogenous trehalose

feeding in *H. armigera*. (a–c) qRT-PCR analysis of *HaST46* and trehalose metabo-

lism-related genes following dietary trehalose supplementation. Transcript abundance

was normalised to reference genes and expressed as fold change ($2^{\wedge-\Delta\Delta Ct}$).

(d, e) In vivo activity of trehalase and trehalose-6-phosphate phosphatase (TPP) in

larvae fed a 50mM trehalose-enriched diet or an artificial diet.

(f) Quantification of trehalose, ATP, ADP and, AMP in whole larvae using LC–MS un-

der the same dietary conditions. Data are presented as mean ± SEM, with statistical

significance assessed by an unpaired *t*-test ($P < 0.05$).

Figure 3. Tissue-specific expression of *HaST46* and trehalose metabolism genes following trehalose feeding. (a–c) qRT-PCR analysis showing expression patterns in haemolymph, fat body, and gut tissues. Transcript levels were normalised to reference genes and expressed as fold change ($2^{-\Delta\Delta Ct}$) relative to control. Data represent mean ± SEM, with statistical differences determined by an unpaired *t*-test ($P < 0.05$).

Figure 4. Silencing of *HaST46* in *H. armigera*. (a, b) qRT-PCR analysis of *HaST46* expression in whole-body samples from control insects (empty L4440 vector) and dsRNA-treated insects, either fed on an artificial diet or an artificial diet supplemented with 50mM trehalose. (c, d) In vivo trehalase and TPP activities under the same conditions. (e, f) Trehalose levels quantified in whole insects using LC–MS/MS. All values are mean ± SEM, with statistical analysis performed using an unpaired *t*-test ($P < 0.05$).

Figure 5. Overexpression of *HaST46* in fifth instar larvae of *H. armigera*. (a) qRT-PCR expression of *HaST46* in whole-body (b) In vivo trehalase activity in whole-body samples following *HaST46* overexpression. (c-e) Expression analysis of *HaST46* in haemolymph, fat body, and gut tissues under four feeding conditions: artificial diet, trehalose-rich diet (50mM), *HaST46* overexpression with artificial diet, and *HaST46* overexpression with trehalose-rich diet (50mM). Transcript levels were normalised to reference genes and expressed as fold change ($2^{-\Delta\Delta Ct}$). (f-g) LC–MS/MS-based quantification of trehalose in haemolymph after *HaST46* overexpression. (h) In vivo trehalase activity in in haemolymph, fat body, and gut tissues samples following *HaST46* overexpression. Data are shown as mean ± SEM, with significance assessed using an unpaired *t*-test ($P < 0.05$).

Supplementary information

Supplementary information S1: H. armigera STs gene sequences

Supplementary information S2: Name of putative HaSTs and its corresponding

gene id from NCBI gene database and scaffold id from transcriptome data

Supplementary information S3: HaST primers used for real-time PCR analysis and

silencing experiment

Supplementary information S4: General methods for real-time PCR, TPP and tre-

halase enzymatic acitivites.

Supplementary information S5: *H. armigera* growth and phenotype upon silencing

and upregulation. Also real-time expression analysis of HaSTs related.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5