A Mammalian Genomic Signature Shaped by Single Nucleotide Variants Regulating

Transcriptome Integrity and Diversity

Jian Yang[1,*], Samuel Ogunsola[2,*], Jason Wong[3,*], Aydan Wang[2], Roby Joehanes[4,5]

Daniel Levy[4,5], Shalini Sharma[3,†], Chunyu Liu[1,4 †], Jiuyong Xie[2, †]

[1]Department of Biostatistics, Boston University School of Public Health, Boston, MA, 02118,

USA

[2]Department of Physiology & Pathophysiology, Max Rady College of Medicine, Rady Faculty

of Health Sciences, University of Manitoba, Winnipeg, MB, R3E 0J9, Canada

[3]Department of Basic Medical Sciences, College of Medicine-Phoenix, University of Arizona,

Phoenix, AZ, USA.

[4] Framingham Heart Study, Framingham, MA 01702, USA

[5]Population Sciences Branch, Division of Intramural Research, National Heart, Lung, and

Blood Institute, National Institutes of Health, Bethesda, MD 20892, USA

Keywords: Non-coding regions, REPAG, GWAS, cryptic splicing, 2nd transesterification step

The authors declare no potential conflicts of interest.

†: Correspondence to: shalinijs@arizona.edu, liuc@bu.edu, and xiej@umanitoba.ca, phone: +1

204 975-7774

Abstract

Many functional features of mammalian genomic sequences remain poorly understood. Here, we identify a widely evolved genomic signature, G-tract-AG motifs consisting of guanine tracts closely upstream of AG dinucleotides, which is significantly associated with single-nucleotide variants (SNVs) identified in genome-wide association studies (GWAS), particularly within non-coding regions (NCRs). Approximately 9,000 such G-tracts within human genes are disrupted by variants of the *cis*-splicing quantitative trait loci (sQTLs) in the Genotype-Tissue Expression (GTEx) project. Functionally, G-tracts repress splicing at the adjacent 3′ AG, primarily by stalling the second transesterification step. Disruption of the G-tracts by SNVs relieves this repression, enabling splicing and generating novel transcript isoforms. These G-tract-disrupting SNVs are in *cis* across the majority of protein-coding genes and are among thousands of rare variants causing genetic diseases. Our findings provide mechanistic insights into the maintenance of transcriptome integrity while still allowing diversity by a mammalian-evolved genomic signature, particularly for the NCR SNVs in association with diverse traits and a new framework for their functional annotation.

Introduction

Many sequence features of mammalian genomes, predominantly composed of non-coding regions (NCRs), remain largely unknown ([1](#)). While cryptically spliced products from the NCR pose challenges to transcript integrity ([2](#)), novel mRNA isoforms driven by genetic variations could contribute to transcript diversity in evolution ([3,4](#)). It remains challenging, however, to identify the functional sequence motifs and single nucleotide variants (SNVs) in the NCR, particularly those reported in genome-wide association studies (GWAS) ([5](#)).

Several sequence motifs have been identified to prevent cryptic splicing of endogenous transcripts, such as pyrimidine-, UG- or CA-rich motifs ([2,6-10](#)). Lengthening uridine tracts also allow the emergence of new exons in primate evolution ([4](#)). Distinct from these motifs, G-tracts are made of $(G)_{\geq 3}$ purines. They are abundant in mammalian introns often as splicing enhancers ([11-13](#)), but not typically anticipated to be within the consensus sequence of the 3′ splice site (SS), $(Y)_{20}NY\underline{AG}$ in humans (Y: C or T; N: A, C, G or T) ([14](#)). When G-tracts do occur inside this region, they repress splicing as <u>r</u>egulatory <u>e</u>lements between the <u>p</u>olypyrimidine tract (<u>P</u>y) and 3′ <u>AG</u> dinucleotides (REPAG) ([15-19](#)). REPAGs have been found within annotated 3′SS in eukaryotes from yeast to human and are enriched the most in mammals ([18](#)), likely contributing to the extensive use of alternative splicing ([17,20,21](#)).

Despite the abundance of G-tracts in mammalian introns ([11](#)), only about 2,000 REPAG-harbouring 3′SS have been identified in the human genome ([17,18](#)). Interestingly, in addition to annotated 3′SS, cryptic 3′SS in cancer tissues with splicing factor mutations also show REPAG enrichment ([18](#)). This observation suggests that additional G-tract-AG signatures may lie deeper within introns or other non-coding regions, where they could serve additional roles.

Here, we searched the human genome and identified millions of these signatures, which are largely in the NCRs conserved in mammals. We show their potent ability to repress or de-repress splicing depending on the SNVs and highlight a distinct mechanism by which genetic

variants in these regions could control splicing in their association with mammalian traits or

diseases. These key sequences may also assist in the functional annotation of SNVs in NCRs

(22-25).

Results

1. Human G-tract-AG signatures: genome-wide presence, enrichment in GWAS-mapped

regions and conservation among mammalian species

To identify genome-wide G-tract-AG signatures, we searched both strands of the human

genome based on the criteria applied to the annotated 3´SS (18) (Fig. 1A). Initially, we

identified and evaluated 111 millions of AG dinucleotides within splice site-like sequences

(3´SSL) that had splicing signals greater than 0.1, the lower boundary of most constitutive

exons (S_Fig. 1), according to their MaxENT scores (26). Although numerous such 3´SSLs

possess potential splicing signals, currently annotated 3´SS is only around 0.3 million (14).

This discrepancy suggests that the majority of these 3´SSLs remain either unused or actively

repressed in cells. Consistent with this repression hypothesis, roughly 6 million of these 3′SSLs

(about 5.4%) harbor the G-tract-AG signature - an enrichment of 2,630 times over that

observed in annotated 3′SS (18). This substantial enrichment underscores the potential

functional importance of G-tract-AG signatures within these 3′SSL genomic regions.

To identify G-tract-AG signatures that are close to or disrupted by single nucleotide

variants (SNVs), we first filtered their 3′SSL sequences (Fig. 1A, 23nt: 18nt + AG + 3nt, by

the MaxENT algorithm(26)) for the presence of any common SNVs from the TOPMed (Trans-

Omics for Precision Medicine, minor allele frequency MAF $\geq$ 0.05) database. This analysis

identified approximately 1.8 million unique SNVs, including 41,353 that overlap with the

GWAS hits. These SNVs were significantly enriched in the GWAS loci (32.1%) compared to

random SNVs in the human genome (30%, Fig. 1B, enrichment index, see also Methods). In

contrast, there was no significant enrichment of SNVs in the CACA- or TGCATG-3′SSL in

GWAS in parallel searches. Moreover, more than 95% of such GWAS SNVs were found in

the NCR, mainly introns and intergenic regions (Fig. 1C). Further, within the genomic start-

end boundaries of 3,202 GWAS genes, we identified 4,447 SNVs that directly disrupt the G-

tracts [$(G)_{3-11}$] in the sense strand (*in cis*, Fig. 1A, S_Table Ia and S_Fig. 2). Together, these

data support a significant association of the G-tract-AG signatures with the GWAS SNVs.

Manual examination of a group of G-tract-AG signatures indicated that most of them

were in mammals only (97%, n = 37), based on the MultiZ-alignment sequences of 100

vertebrates from the UCSC Genome Browser ([27]) (S_Fig. 3). However, they exhibited higher

divergence than those in the annotated 3′SS ([15],[17]), for instance, about 25% of them found in

primates only. In the two representative examples of different levels of conservation (Fig. 1D),

the G-tract-AG signature in *MAX* is conserved in 80.6% of the species (n = 62), and the

signature in *QSOX1* is conserved in 7.3% only (n = 41). The G-tract disruption occurred in

other species as well, by A, C or T in guinea pig and primates, among other changes within the

3′SSLs (see also S_Figs. 4 & 5). By MaxENT scores, the G-tract disruption increased the

splicing signal to or close to that of the human SNV-T variants in a few species (*MAX* in guinea

pig and *QSOX1* in chimpanzee, gorilla and orangutan).

To systematically assess the conservation/divergence of the genomic signatures and G-

tract-disrupting SNVs located between human gene start and end boundaries, we obtained their

phyloP scores from the alignment of 470 mammalian genomes ([27]). The mean phyloP scores

for the G tract, G-tract-disrupting SNV, AG and G-tract-AG are -0.34, -0.61, -0.14 and -0.24,

respectively (n > 107K motifs, SEM = 0.004 each, S_Fig. 2B). The mean scores are

significantly higher for the GWAS than the non-GWAS group (Fig. 1E), consistent with the

reported association between higher phyloP scores (constraint) and higher enrichment of

GWAS SNVs (28). Based on the phyloP scores and presence or absence of corresponding

signature motifs in the MultiZ alignment(27), we estimate that about 63% of the signatures are

limited to primates (G-tract-AG phyloP score ≤ 0), while the remaining signatures (phyloP

score > 0) extend beyond primates to other mammals, with only a small fraction $(< 0.1\%)$ also

found in fish. Therefore, the G-tract-AG signatures or the G-tract-disruptive SNVs are mainly

found in mammals with varying degrees of divergence or conservation.

These data, together with the REPAG repression of splicing (15,18), imply a much wider

role for the G-tract-AG signature in the NCR of mammalian genomes, beyond its control of

alternative splicing of annotated 3′SS (15-17).

2. G-tract repression and SNV-caused de-repression of splicing from the 3′ AG

To determine whether the G-tract-disruptive SNVs are globally associated with splicing,

we examined156,380 such SNVs *in cis* within gene start-end boundaries in the human genome,

and cross-referenced them with the *cis*-sQTLs (splicing quantitative trait loci, false discovery

rate, FDR<0.05, n = 486,725 unique IDs) from the Genotype-Tissue Expression Project (GTEx)

v10 (29). Of these, approximately 9,000 SNVs were significant sQTLs (5.62%, S_Table_Ib),

representing a 1.58-fold enrichment compared to randomly selected SNVs (n=1,000 sets of

156,380 SNVs, matched by their MAF and transcription start sites (TSS), $p = 2.2E\text{-}16$, one

sample t-test). Of the 4447 GWAS SNVs, the sQTL proportion is even higher (14%). Thus,

these G-tract-disruptive SNVs are significantly enriched for sQTLs, supporting their potential

role in splicing and impact on the transcriptome.

To validate the potential effects of the G-tracts and SNVs on splicing, we chose the above

two examples for their evolutionary conservation or divergence, increased MaxENT scores

upon disruption by SNVs and deep intron locations. We examined the effects in mini-gene

splicing reporter assays (Fig. 2 and S_Fig. 5), which we have employed to consistently

demonstrate the G-tract repression of splicing in ten REPAG reporters (15,17,19) (S_Table II).

The *MAX* SNV rs762810-T is in the upstream 3′SS of exon 4 (E4), in a mammal-specific but

much less conserved region than the E3 region (Fig. 2A). The E4 inclusion is at about 20% of

the E3 by their RNA-Seq reads in blood cells, but invisible in the other 53 GTEx tissues

including kidney and skin. Moreover, the intron 3 excision ratio is significantly higher in

samples with the rs762810-T/T than those with the G/G genotype (S_Fig. 6) (29). The *MAX*

G-tract repression and SNV de-repression effects were indeed validated in the splicing

reporters (Fig. 2B): in human kidney HEK293T cells, the $(G)_4$-carrying reporter skipped the

E4. In contrast, the $(G)_3$T-carrying reporter with a transversion of the last G to the rs762810-T

resulted in 66% E4 usage (in molar ratio), and the $(A)_4$ reporter with G to A transitions resulted

in 98% E4 usage. Thus, it is the $(G)_4$ that represses but the rs762810-T that de-represses splicing

from the 3′ AG. The *QSOX1* rs3767199-T is in the first intron, 398nt downstream of exon 1

(S_Fig. 5C). Like the rs762810-T, this variant also de-repressed the G-tract repression of

splicing from its 3′ AG splice site, albeit to a smaller extent in the reporter assay.

To visualize the SNV genotype effect on splicing in humans, we examined the 3′SS

intron/exon ratios of RNA-Seq reads, as described previously (30), in genotype- and RNA-Seq

data-matched whole blood cell samples from participants in the Framingham Heart Study (FHS)

(50). We found intron/exon junction-reads ratios of 1.0, 0.7, and 0.2 for individuals with G/G,

G/T, or T/T genotypes of rs762810, respectively (Fig. 2C), indicating no splicing, partial

splicing, or near-complete splicing of the 3′SS of the blood cell-specific exon. We further

conducted association analyses with genotype or additive models between 3' AG usage

(measured as 1 - ratio of intron/exon junction-reads) and 3,302 SNVs located within G-tracts

up to 13nt upstream of AG acceptor sites (S_Table_I), using RNA-seq data from 100 FHS

participants. By the two models (S_Table_I and S_Files_1 & 2), we identified 241 pairs of

SNV and G-tract-AG sites where the mean 3′ AG usage was increased in the non-G variant

(H)-harboring genotype over the G/G homozygotes (Fig. 2D, $p < 2.0E\text{-}18$). The 3' AG usage

was not correlated with the 3'SS MaxENT score (Pearson coefficient $r = -0.05$, $p = 0.47$),

likely due to the inclusion of G-tract-AG as training decoy sites (~20%) for the MaxENT

algorithm([26]). However, the mean value of the 3' AG usage was correlated inversely with the

G-tract lengths (3-8nt, $r = -0.94$, $p < 0.02$), where the additive effect by the H variants is evident

in the $(G)_{3\text{-}6}$ groups (Fig. 2D). In contrast, there was no such a correlation between the 3' AG

usage with the distance of the $(G)_3$-tracts to the 3′ AG ($r = -0.03$, $p = 0.92$). Therefore, the

human genotype and allele effects strongly support that a large proportion of the G-tracts

repress and the G-tract-disrupting SNVs de-repress splicing for 3' AG usage in a G-tract length-

dependent way.

Taken together, these data support that the G-tract-AG signatures have dual functions in

the NCR: (1) to repress splicing from the 3′ AG with the G-tract as a functional REPAG to

protect transcript integrity and (2) to permit splicing for the emergence of novel exon (e.g., E4

in *MAX*) or transcript isoforms in mammals upon disruption of the G-tract by a SNV for

transcript diversity.

3. Splicing repression at the two transesterification steps by the REPAG

For the mechanism of splicing repression by the G-tracts, we have shown previously that

a REPAG element with its bound hnRNP (heterogeneous nuclear ribonucleoprotein) H1

represses splicing and reduces Py-binding by an early spliceosome factor U2AF65 ([15],[17]). The

hnRNP H1 binding is disrupted by mutating the G-tract ([15]), and by the *MAX* SNV-T variant

in structure modelling (S_Fig. 7). However, it is still unclear which of the two

transesterification steps of splicing is repressed by such G-tracts. Our *in vitro* assay of splicing

intermediates indicated that both steps were repressed but primarily at the 2^{nd} step.

In the *in vitro* splicing assay, we analyzed the kinetics of lariat formation in a pre-mRNA

containing either the REPAG or its mutant as in our previous report ([7](#)) (Fig. 3A). Compared

to the AdML vector, the presence of REPAG caused a significant delay by about 6 minutes in

generating the lariat intermediate (lanes 1-3 vs. 4-6, and Fig. 3B). The delay was reversed upon

mutating the REPAG sequence (lanes 7-9). Thus, REPAG delays the generation of the first

step product, the lariat intermediate, which is consistent with the reduction of U2AF65 binding

to the Py ([15](#)).

Interestingly, the second step of splicing was also significantly delayed by REPAG but

far more than would be expected from a simple delay of the first-step (Fig. 3B). The half-life

of lariat intermediates in conversion to the lariat was approximately 10, 21, and 45min for the

AdML, REPAG mutant, and REPAG reporters, respectively (Mean, ±1.2, ±4.4, ±6.5, SEM, n

= 6, 3 and 4, Fig. 3C), indicating a significant delay by the REPAG. Additionally, a less

pronounced second-step repression was also observed for the REPAG mutant, likely due to the

longer spacer between the Py and 3′ AG as reported ([31](#)).

Together, the results from the splicing intermediate assays demonstrate that the REPAG

unusually represses both transesterification steps with a stronger effect on the 2^{nd} step of

splicing.

4. Widespread impact of *in cis* disruption of the G-tracts of the genomic signatures by

SNVs

The impact of the genomic G-tract-AG signature is likely widespread on mammalian

genes. In the GWAS, the 3,202 genes that contain G-tract-disrupting SNVs *in cis* are strongly

enriched for a variety of health-related traits including heart disease and cancer by DAVID

functional clustering analysis (Fig. 4A, S_Table Ia). About 77% of them are protein-coding

genes. In the human genome, the 156,380 G-tract-disrupting SNVs are *in cis* within 23,479

unique genes, including about 71% of the 20,078 protein-coding genes in the hg38 (Fig. 4B,

S_Table Ib). These figures together with the sQTL enrichment and functional data on splicing

(Figs. 2-3 and S_Fig. 5) imply widespread impact by the G-tract-AG signatures and their SNVs

on the transcriptomes and proteomes. For example, the *MAX* E4 usage contributes to two

transcripts, resulting in truncated protein isoforms lacking the nuclear localization signal (NLS)

or dimerization domain (S_Fig. 8) (27,32-34). Several similarly truncated MAX isoforms

showed cell growth-promoting instead of the tumor suppressor effect by the full length protein

(33,35-37). The E4-promoting T-variant is associated with higher blood cell counts in GWAS

(38), enriched among individuals with melanomas and associated with E4 ectopic inclusion in

melanoma tumor cells (S_Fig. 9), implying its association with cell growth as well. The usage

of the *QSOX1* cryptic splice site caused by the rs3767199-T variant would lead to intron

retention and nonsense-mediated mRNA decay reducing full-length protein levels (S_Fig. 5A).

Related to this anticipation, reduced protein level in *Qsox1* knockout mice caused lower pulse

pressure, an effect also associated with the rs3767199-T variant in GWAS (39,40). Moreover,

the sQTL of predominantly primate G-tract-AG signatures are highly enriched in cadherin

family genes ($p < 1E-10$), which are important for cell-cell communications, while those also

conserved in other mammals are enriched in genes encoding proteins with signal transduction

domains (C2 or SH3, $p < 1E-4$). These distinct enrichment patterns suggest that the signatures

may have contributed to the evolution of lineage-specific functional properties in primates and

other mammals.

Beyond the genome-wide common SNVs, we also identified 2,145 rare genetic variants

(MAF<0.01) that disrupt the G-tracts of the signatures in the human gene mutation database

(HGMD) and ClinVar (41,42) (Fig. 4B, S_Fig. 10 and S_Tables III). For instance, the

c.392G>T variant of the glucose-6-phosphate dehydrogenase (*G6PD*) gene causes a neutral

change of amino acids (G131V) but deficient enzyme activity in hemizygotes (43-45). The

discrepancy could be well explained by RNA level changes instead, by aberrant splicing due

to T variant-disruption of the G-tract resulting in 44aa deletion at the N-terminal region (Fig.

4C). The variant enzyme will lose the Y112 for NADP$^+$ binding and catalytic activity of the

dehydrogenase (46).

Conclusion

Taken together, our data define an evolved mammalian genomic signature (G-tract-AG)

enriched for GWAS SNVs and GTEx sQTLs, and demonstrate that G-tract-disruptive SNVs

act through a distinct molecular mechanism to control splicing. The G-tract, as a two-step tight

splicing repressor, not only regulates alternative splicing of annotated 3′SS as previously

reported(15), but also represses the latent AG splice sites from splicing in the introns of the

genome. The 3′ AG becomes a splice site upon disruption of the G-tract by SNVs, leading to

the generation of novel transcript isoforms. Therefore, it is likely that the widespread G-tract-

AG bipartite motifs regulate mammalian transcriptome integrity and diversity depending on

the SNV.

Our finding provides not only a functional genomic signature to explain how

transcriptome integrity and diversity are controlled in mammalian species by the signatures

and their SNVs, but also a valuable tool for annotating similar sequence signatures and SNVs

in the genomes. Identifying more of such signatures will help cover a much larger proportion

of GWAS SNVs such as those in the CACA-3′SSL (Fig. 1B) or even coding regions (Fig. 1C,

and 4D), or in other species (Fig. 1D). It will be particularly helpful for the identification of

functional SNVs underestimated by existing prediction algorithms such as MaxENT or

SpliceAI (23). For examples, the *MAX* and *QSOX1* SNVs were scored either below the

confidence level 0.5 or not even scored by SpliceAI, as were 99% of the GWAS SNVs (Fig.

2A, S_Figs. 11, and S_Table IV). Further genome-wide functional annotation of the signatures

will provide a global view of the balanced control between the maintenance of transcriptome

integrity and the evolution of transcriptome diversity, as well as genetic differences among

individuals or mammalian species. Moreover, detailed mechanistic studies on the motifs,

splicing factors, and local structures will also provide further insights into the signature's

unusual effect on the 2^{nd} step of splicing, as found in other cases (47,48). Together, these studies

on the signatures could help bridge the mammalian genome with transcriptome integrity and

diversity but also advance personalized medicine through understanding and correcting cryptic

splicing caused by SNVs (49).

Materials and Methods

***In vitro* splicing assay**: To the 3′ splice site of the adenovirus splicing reporter (50), we inserted

the REPAG gggtgggggagtg upstream of the PRMT5 exon 3 and its mutant aaataaaaaagtg (15).

The *in vitro* splicing was carried out in HeLa nuclear extracts as reported (50). The optimal

time range for the time course measurements of lariat products was determined by pre-tests to

avoid RNA degradation after prolonged incubation *in vitro*.

Human genome search, G-tract-AG harbouring 3´SSL and phyloP score analysis: The

GRCh38/hg38 reference genome was searched on both strands by identifying $(G)_{\geq 3}$ (i.e., at

least three consecutive guanines) within 1 to 13nt upstream of AG dinucleotides (18). The 3´SS

23nt (18nt upstream plus 3nt downstream of AG) region was filtered for MaxENT 3´SS

strength measurement (26), and for the presence of TOPMed common SNVs (MAF $\geq$ 0.05)

(51). Briefly, to identify potential 3´SS AG sites, we developed a pipeline that first searched

for genomic coordinates of AG dinucleotides on both strands across the human genome, then

output the corresponding 23nt sequences covering the AG (5´- 18nt + AG + 3nt -3´, e.g. 5´-

ttccaaacgaacttttgtaggga-3´). The likelihood of each sequence functioning as a valid 3´SS was

quantified using MaxEntScan (26). To map SNVs within the 23nt 3′SSL sequences, we used

the TOPMed SNV annotation to retrieve allele and MAF information and integrated these with

SNVs reported in the GWAS Catalog(5). The TOPMed SNV annotation has been developed

from deep sequencing on over 200,000 human genomes, with an average coverage of

approximately 30-fold (51). In this study, we selected SNVs with MAF $\geq$ 0.05 (approximately

3.5% of all TOPMed variants) from the TOPMed SNV annotation to identify variants within

the 23nt region. To obtain MaxENT scores for the 23nt sequences, we replaced the reference

alleles with the alternative alleles and recalculated splice site strength using MaxEntScan.

To identify G-tract-AG-harbouring 3´SSL, we extracted 23nt sequences (Fig. 1A)

containing $(G)_{\geq 3}$ between the 6th and 18th nucleotides using our in-house pipeline. The

consensus sequence motifs from these 23nt regions were then obtained using the online MEME

motif analysis tool ([52]).

The phyloP scores of genomic motifs, SNVs and 3´SSL were obtained from the Hiller

lab's analysis of 470-mammal alignment through the UCSC Genome webserver

(https://hgdownload.soe.ucsc.edu/goldenPath/hg38/phyloP470way/) ([27]). The >107K scores

of each group were averaged by their nucleotide lengths. The cutoff scores used to distinguish

signatures present predominantly in primates (≤ 0) from those also in other mammals (> 0) was

manually verified using the UCSC Genome Browser's MultiZ alignment of 100 vertebrates.

Enrichment analysis of G-tract-AG-harbouring 3′SSL among GWAS loci, GTEx

sQTLs or health traits/diseases. Some SNVs in the GWAS Catalog ([5],[53]) lack allele

information or allele frequencies. Therefore, we annotated the GWAS SNVs using TopMed

SNV data. We selected SNVs with MAF ≥ 0.05 and p-value $\leq$ 5E-08 from the GWAS Catalog,

resulting in 128,991 distinct GWAS SNVs. These SNVs were categorized into 20 groups

according to five MAF bins (0.01, 0.05, 0.1, 0.2, 0.3, and 0.5) and four distance bins (0kb, 1kb,

10kb, 100kb, and 1Mb) from the transcription start site (TSS) of the closest genes, creating 20

matching groups ([54]). Based on the same criteria, we generated 20 corresponding groups from

all TOPMed SNVs. Specifically, TOPMed SNVs were categorized into five MAF bins (0.01,

0.05, 0.1, 0.2, 0.3, and 0.5) and four distance bins (0 kb, 1 kb, 10 kb, 100 kb, and 1 Mb) from

the transcription start site (TSS) of the closest genes, yielding 20 matching groups parallel to

those defined for the GWAS SNVs. From each group, we randomly selected the same number

of SNPs as in the corresponding GWAS groups, yielding 128,991 SNPs in total. To minimize

bias, we further generated 1000 random sets of 128,991 SNVs using the same procedure.

We calculated the proportion of GWAS SNVs located within G-tract-AG-harbouring

3′SSL (n = 29,284). We also calculated the proportion of the 128,991 randomly selected SNVs

within these 3′SSL across 1000 sets. Fold-enrichment was calculated as the ratio of the

proportion of GWAS SNVs within 3′SSL to the average proportion of matched SNVs within

3′SSLs across the 1000 sets. GTEx v10 sQTLs of 54 tissues were filtered to obtain 486,725

unique sQTLs, which was used to identify sQTLs located within the G-tract-AG-harbouring

3′SSL. One thousand random sets of SNVs matched by number, MAF and distance to TSS

were generated using the same method described above. Enrichment ratio was calculated

similarly as well.

Disease enrichment/cluster analysis was carried out using the DAVID functional

clustering analysis at the NIH webserver (https://davidbioinfomatics.nih.gov/) with the

Genetic Association Database (GAD) (55,56), which is a curated archive of human genetic

association studies of complex diseases and disorders. GAD annotations include genes linked

to reported disease associations, which allows enrichment analysis of gene–disease

relationships. Genes from our list were tested against the GAD annotation set. Enrichment was

assessed using DAVID's modified Fisher's exact test (EASE score). Three sets of control

genes, 3202 per set, were randomly picked from 66,832 stable IDs of human genes for the same

analyses.

Study participants: This study used WGS and RNA-seq data from selected samples of 1,544

Third Generation cohort participants (mean age 47±9 years, 52% women, 100% American

White) (57), at the second examination cycle (2008–2011). Protocols for genetic material

collection during regular health examinations were approved by the Institutional Review Board

at Boston Medical Center. All participants provided written informed consent for genetic

studies, and the research was conducted in accordance with relevant guidelines and regulations.

RNA-Seq in the Framingham Heart Study: Peripheral whole blood samples were collected

at the second examination cycle (2008-2011). RNA was isolated from peripheral whole blood

using PAXgene™ tubes (PreAnalytiX, Hombrechtikon, Switzerland) to collect 2.5 mL samples

from FHS participants (57). Samples were incubated at room temperature for 4 hours for RNA

stabilization and then stored at −80 °C. Prior to extraction, tubes were thawed at room

temperature for 16 hours. Total RNA was extracted with a PAXgene Blood RNA Kit at the

FHS Genetics Laboratory. Following centrifugation and washing, white blood cell pellets were

lysed in guanidinium buffer. RNA quality was assessed by absorbance at 260 and 280 nm using

a NanoDrop ND-1000, and integrity was evaluated with the Agilent Bioanalyzer 2100

microfluidic.

RNA sequencing was conducted at the University of Washington Northwest Genomics

Center, an NHLBI TOPMed reference laboratory, following standard protocols. Data

processing took place at the same facility. Base calls were generated on the NovaSeq 6000, and

unaligned BAM files were converted to FASTQ format using SamTools bam2fq. Sequence

quality was assessed with the FASTX-toolkit, and alignment to GRCh38 with GENCODE

release 30 was performed using STAR. FastQC provided initial quality control metrics, and

STAR created a genome index to generate BAM files at the gene and transcript levels(58). The

TOPMed RNA-seq pipeline utilized RNA-SeQC for standard quality control metrics from

aligned reads(59).

Whole genome sequencing: DNA from peripheral whole blood from 7,215 FHS participants

underwent whole genome sequencing (WGS) at the Broad Institute, a TOPMed contracting

laboratory. Details on DNA extraction, processing, and variant identification followed standard

procedures (51). Consistent sequencing and data processing criteria were applied, aligning

DNA sequences to human genome build GRCh38. The resulting BAM files were sent to

TOPMed's Informatics Research Center (IRC) for re-alignment and standardized BAM file

generation. This study utilized WGS data from Freeze 10a, which included approximately 208

million SNPs across TOPMed cohorts.

Visualization of RNA-Seq reads corresponding to rs762810 (G>T) genotypes: We

identified the same FHS participants with both RNA-Seq and WGS data. We then randomly

selected three participants with GG, GT, and TT genotypes of rs762810 (G > T, MAF = 0.25

for T). We applied Integrative Genomics Viewer (IGV) ([60](#)) to visualize the RNA-Seq reads

count in response to the genotypes of rs762810, GG, GT, and TT.

Association analysis between SNVs and 3′ AG usage: We quantified 3′ AG usage of the

3′SSLs following previously described methods for intron/exon ratios of RNA-Seq reads

within 25nt flanking the intron-exon junctions ([61](#),[62](#)), in 100 randomly selected FHS

participants. The 25nt downstream of the 3' AG were considered as exon sequences here.

Briefly, RNA-seq reads were first aligned to the reference genome hg38 and reads mapped

within 25nt flanking the intron-exon junctions overlapping the plus-strand G-tract-AG

signatures were extracted using the *featureCounts* in Subread package([63](#)). The 3' AG usage

was quantified as: 1 - the ratio of (intron reads)/(exon reads), within 25nt flanking the junction.

Additional quality control was applied to the intron-exon junction ratios. Specifically, we

excluded AG sites with missing ratio values caused by zero-read counts on both sides of the

AG site. We also removed sites with negative ratio values below -0.1. We focused on SNVs

located within G-tracts up to 13nt upstream of AG sites (S_Table_I). Two association models

were applied to examine the relationship between SNV genotypes and 3' AG usage in the 100

FHS participants. In the genotype model, genotypes were treated categorically (0/0, 0/1 or 1/1),

using the homozygous reference (0/0) as the baseline when all three genotypes were present.

If "0/0" was absent due to limited sample size, the genotype with the largest number of samples

was used as the reference. SNVs with only one observed genotype were excluded. In the

additive model, association was tested using alternative allele dosage (0, 1, or 2) as predictor

for the ratios in the same 100 sample. Covariates included sex, blood cell counts, five genotype-

based principal components (PCs) to account for population stratification, and 15

transcriptome-based PCs to adjust for unknown confounders affecting gene expression(64).

Given the limited sample size, statistical significance was assessed with a liberal threshold of

$p < 0.05$. Violin plots were generated to show the associations between SNP genotypes and the

ratios (S_files 1 and 2). To make sure the analysis method correctly reflects the splicing

changes, we pre-tested it with the *MAX* E4 upstream 3'SS and obtained significant ratio changes

consistent with the RNA-Seq and splicing reporter results in figure 2A-C.

Structure modelling of protein-RNA complexes: We used AlphaFold 3.0(65) to model the

hnRNP H1 protein-RNA structure and changes by the SNV. We first confirmed the prediction

accuracy by aligning the top-ranked AlphaFold models with the corresponding crystal or NMR

structures of a group of RNA-binding proteins in complex with RNA oligos (≤ 30nt) from the

protein data bank PDB(66). The root mean square deviation (RMSD) was 0.559 angstrom (±

0.049, mean ± SEM, n = 21 pairs of predicted and PDB structures) after alignment using

PyMOL (Molecular Graphics System, Version 3.0 Schrödinger, LLC). This deviation is

significantly less than the threshold of 3.0 (p < 0.0001, one sample two-tailed t-test) for

structural differences (67). We then modelled the protein-RNA complex structures between

hnRNP H1 qRRM1 and qRRM2 and the REPAG-containing RNA oligos 5´-UGGGGCU-3´

and variant 5´-UGGGUCU-3´of the 3´SS of *MAX* intron 3.

Mini-gene splicing reporter assay in HEK293T cells: G-tract, SNV or mutant reporters of

MAX intron 3 or *QSOX1* intron 1 were made using the Gibson mutagenesis system(68), and

verified by Sanger sequencing. Briefly, plasmids were transfected into HEK293T cells for

overnight before RNA extraction for RT-PCR assay of spliced variants, as previously

reported(69).

Plasmid Construction: Minigene splicing reporters were constructed using the Gibson

Assembly Master Mix (New England Biolabs(68)) following the manufacturer's protocol.

Linear PCR fragments were generated using primers with overlapping sequences

corresponding to the *ApaI/BglII* linearized pDUP175 vector. The fragments were designed with

the NEBuilder Assembly Tool.

For the pDUP175-MAX-E4-WT plasmid, a genomic DNA fragment containing exon 4

(81bp) and partial flanking introns (27 bp upstream and 22 bp downstream) was PCR amplified

from human genomic DNA. The primers used were MAXE4-WT-DUP175 (5′-

CTTGGGTTTCTGATAGGGCCCTGACCTCTCCCCTTCTTGGGGCTCAGGTG-3′) and

MAX_3′SS_DUP175_REV (5′-

TCTCTGTCTCCACATGCCCAACCTGGAGTCTCTGGTAC-3′).

For the pDUP175-STREX-QSOX1-3′SS-WT plasmid, the 3′ splice site of the STREX exon

from the pDUPST1 vector was replaced by a 27-nt sequence from intron 1 of QSOX1 using

PCR with primers QSOX1_DupST1_3SS'_WT (5′-

CTTGGGTTTCTGATAGGGCCCTGACACCTTACCGTTTCCTGGGCCAGCCAAGATG

TCCATCTAC-3′) and DUP175_DupST1_3′SS_REV (5′-

TCTCTGTCTCCACATGCCCAAGACATGGAACTGTGTGTG-3′). Mutant plasmids were

created by amplifying the PCR fragments from the wild-type plasmid using complementary

mutant-specific primers and cloned using the Gibson Assembly Master Mix. All plasmid

constructs were verified through DNA sequencing.

Cell Culture and Transfection: Human embryonic kidney (HEK293T) cells were cultured at

37°C with 5% CO_2. The cells were grown in Dulbecco's Modified Eagle Medium (DMEM)

supplemented with 10% fetal bovine serum (FBS) and 1% penicillin-streptomycin-glutamine

(PSG). For transfections, 1μg of plasmid DNA was used per well in a 6-well plate, and

transfection was performed using Lipofectamine 3000 (Invitrogen), according to the

manufacturer's protocol.

RT-PCR: Reverse transcription was carried out based on a previously described procedure (7).

Briefly, for reverse transcription, 1μg of total RNA (for MAX) or cytoplasmic RNA (for

QSOX1) was added to a 10μl RT reaction and incubated at 45°C for 50min, followed by a 5-

minute incubation at 90°C. For PCR amplification, 1μl of the RT product was used in a 24μl

reaction. The PCR cycling was set for 30 cycles using primers DUP8a (5′-

GACACCATGCATGGTGCACC-3′) and DUP10 (5′-CAAAGGACTCAAAGAACCTCTG-

3′). PCR products were resolved on agarose gels, visualized under UV light, and photographed

using a digital camera. Band intensities were quantified using ImageJ software (70). Exon

inclusion efficiency was calculated as the molar percentage of the included products by their

intensity relative to the total intensity of both included and skipped products after normalization

to the production lengths.

Data Availability Statement: The whole genome sequencing (WGS) and RNA sequencing

(RNA-seq) data from the Framingham Heart Study (FHS) are available at the dbGaP database

under accession codes, phs000007.v32.p13 (FHS)

[https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=phs000007.v34.p15].

These data are available under restricted access to ensure confidentiality and protect participant

privacy. Access can be obtained by submitting an ancillary study proposal and obtaining IRB

approval. Timelines for the approval process range from 3–6 weeks. Researchers must first

obtain an eRA Commons account, which is typically set up within 3-7 business days if the

applicant's institution is already registered with NIH (institutional registration may add an

additional 2-4 weeks).

The WGS and RNA-seq data from Genotype-Tissue Expression (GTEx) Project can be

obtained through the dbGaP database under accession code phs000424.v9.p2. Researchers

must submit a Data Access Request (DAR) through dbGaP, including institutional and IRB

approvals. Review and approval of data access requests typically requires 2–6 weeks. Summary

statistics are publicly available through the GTEx Portal (https://gtexportal.org).

Acknowledgements: We thank the donors for the Framingham Heart Study for sharing their

genotype and RNA-Seq information, and Dr. Ren-Jang Lin for helpful comments. The work is

supported by a discovery grant (RGPIN-2022-05023) by the Natural Sciences and Engineering

Council of Canada (NSERC) to J.X., by NIH (GM127464 and P30CA023074) and Arizona

Biomedical   Research   Centre   (ABRC: RFGA2022-010-30)   to   S.S.,

and by NIH R01AA028263 and R01HL15569 to C.L. The Framingham Heart Study (FHS)

was   supported   by   NIH   contracts   N01-HC-25195,   HHSN268201500001I,   and

75N92019D00031. RNA-Seq was supported in part by the Division of Intramural Research

(D. Levy, Principal Investigator) and an NIH Director's Challenge Award (D. Levy, Principal

Investigator).

479
480   **Author contributions**: JX conceptualized the project with LC and SS. JY, SO and JW carried

481   out the bioinformatics analyses, cell culture or *in vitro* splicing reporter assays, respectively.

482   AW contributed to the REPAG MEME and AlphaFold3 structural analyses, RJ and DL

483   contributed to the FHS RNA-Seq data. JX wrote the manuscript, with critical revisions by LC

484   and SS. All authors have read and approved the manuscript.

References

1. Shabalina, S.A. and Spiridonov, N.A. (2004) The mammalian transcriptome and the function of non-coding DNA sequences. *Genome Biol*, **5**, 105.

2. Ling, J.P., Pletnikova, O., Troncoso, J.C. and Wong, P.C. (2015) TDP-43 repression of nonconserved cryptic exons is compromised in ALS-FTD. *Science*, **349**, 650-655.

3. Gilbert, W. (1978) Why genes in pieces? *Nature*, **271**, 501.

4. Attig, J., Ruiz de Los Mozos, I., Haberman, N., Wang, Z., Emmett, W., Zarnack, K., Konig, J. and Ule, J. (2016) Splicing repression allows the gradual emergence of new Alu-exons in primate evolution. *Elife*, **5**.

5. Sollis, E., Mosaku, A., Abid, A., Buniello, A., Cerezo, M., Gil, L., Groza, T., Gunes, O., Hall, P., Hayhurst, J. *et al.* (2023) The NHGRI-EBI GWAS Catalog: knowledgebase and deposition resource. *Nucleic Acids Res*, **51**, D977-D985.

6. McClory, S.P., Lynch, K.W. and Ling, J.P. (2018) HnRNP L represses cryptic exons. *RNA*, **24**, 761-768.

7. Lei, L., Cao, W., Liu, L., Das, U., Wu, Y., Liu, G., Sohail, M., Chen, Y. and Xie, J. (2018) Multilevel Differential Control of Hormone Gene Expression Programs by hnRNP L and LL in Pituitary Cells. *Mol Cell Biol*, **38**.

8. Das, U., Nguyen, H. and Xie, J. (2019) Transcriptome protection by the expanded family of hnRNPs. *RNA Biol*, **16**, 155-159.

9. Zarnack, K., Konig, J., Tajnik, M., Martincorena, I., Eustermann, S., Stevant, I., Reyes, A., Anders, S., Luscombe, N.M. and Ule, J. (2013) Direct competition between hnRNP C and U2AF65 protects the transcriptome from the exonization of Alu elements. *Cell*, **152**, 453-466.

10. Ling, J.P., Chhabra, R., Merran, J.D., Schaughency, P.M., Wheelan, S.J., Corden, J.L. and Wong, P.C. (2016) PTBP1 and PTBP2 Repress Nonconserved Cryptic Exons. *Cell Rep*, **17**, 104-113.

11. Yeo, G., Hoon, S., Venkatesh, B. and Burge, C.B. (2004) Variation in sequence and organization of splicing regulatory elements in vertebrate genes. *Proc Natl Acad Sci U S A*, **101**, 15700-15705.

12. McCullough, A.J. and Berget, S.M. (1997) G triplets located throughout a class of small vertebrate introns enforce intron borders and regulate splice site selection. *Mol Cell Biol*, **17**, 4562-4571.

13. Martinez-Contreras, R., Fisette, J.F., Nasim, F.U., Madden, R., Cordeau, M. and Chabot, B. (2006) Intronic binding sites for hnRNP A/B and hnRNP F/H proteins stimulate pre-mRNA splicing. *PLoS Biol*, **4**, e21.

14. Nguyen, H., Das, U., Wang, B. and Xie, J.Y. (2018) The matrices and constraints of GT/AG splice sites of more than 1000 species/lineages. *Gene*, **660**, 92-101.

15. Sohail, M. and Xie, J. (2015) Evolutionary emergence of a novel splice variant with an opposite effect on the cell cycle. *Mol Cell Biol*, **35**, 2203-2214.

16. Sohail, M. and Xie, J. (2015) Diverse regulation of 3' splice site usage. *Cell Mol Life Sci*, **72**, 4771-4793.

17. Sohail, M., Cao, W., Mahmood, N., Myschyshyn, M., Hong, S.P. and Xie, J. (2014) Evolutionarily emerged G tracts between the polypyrimidine tract and 3' AG are splicing silencers enriched in genes involved in cancer. *BMC Genomics*, **15**, 1143.

18. Nguyen, H. and Xie, J. (2018) Widespread Separation of the Polypyrimidine Tract From 3' AG by G Tracts in Association With Alternative Exons in Metazoa and Plants. *Front Genet*, **9**, 741.

19. Romano, M. (2010) G runs in cystathionine beta-synthase c.833C/c.844_845ins68 mRNA are splicing silencers of pathogenic 3' splice sites. *Biochimica et biophysica acta*, **1799**, 568-574.

20. Xie, J. (2014) Differential evolution of signal-responsive RNA elements and upstream factors that control alternative splicing. *Cell Mol Life Sci*, **71**, 4347-4360.

21. Barbosa-Morais, N.L., Irimia, M., Pan, Q., Xiong, H.Y., Gueroussov, S., Lee, L.J., Slobodeniuc,
V., Kutter, C., Watt, S., Colak, R. *et al.* (2012) The evolutionary landscape of alternative
splicing in vertebrate species. *Science*, **338**, 1587-1593.

22. Sullivan, P.J., Gayevskiy, V., Davis, R.L., Wong, M., Mayoh, C., Mallawaarachchi, A., Hort, Y.,
McCabe, M.J., Beecroft, S., Jackson, M.R. *et al.* (2023) Introme accurately predicts the impact
of coding and noncoding variants on gene splicing, with clinical applications. *Genome Biol*,
24, 118.

23. Jaganathan, K., Kyriazopoulou Panagiotopoulou, S., McRae, J.F., Darbandi, S.F., Knowles, D.,
Li, Y.I., Kosmicki, J.A., Arbelaez, J., Cui, W., Schwartz, G.B. *et al.* (2019) Predicting Splicing
from Primary Sequence with Deep Learning. *Cell*, **176**, 535-548 e524.

24. Dawes, R., Bournazos, A.M., Bryen, S.J., Bommireddipalli, S., Marchant, R.G., Joshi, H. and
Cooper, S.T. (2023) SpliceVault predicts the precise nature of variant-associated mis-splicing.
Nat Genet, **55**, 324-332.

25. Zeng, T. and Li, Y.I. (2022) Predicting RNA splicing from DNA sequence using Pangolin.
Genome Biol, **23**, 103.

26. Yeo, G. and Burge, C.B. (2004) Maximum entropy modeling of short sequence motifs with
applications to RNA splicing signals. *J Comput Biol*, **11**, 377-394.

27. Nassar, L.R., Barber, G.P., Benet-Pages, A., Casper, J., Clawson, H., Diekhans, M., Fischer, C.,
Gonzalez, J.N., Hinrichs, A.S., Lee, B.T. *et al.* (2023) The UCSC Genome Browser database:
2023 update. *Nucleic Acids Res*, **51**, D1188-D1195.

28. Sullivan, P.F., Meadows, J.R.S., Gazal, S., Phan, B.N., Li, X., Genereux, D.P., Dong, M.X.,
Bianchi, M., Andrews, G., Sakthikumar, S. *et al.* (2023) Leveraging base-pair mammalian
constraint to understand genetic variation and human disease. *Science*, **380**, eabn2937.

29. Consortium, G.T. (2013) The Genotype-Tissue Expression (GTEx) project. *Nat Genet*, **45**, 580-
585.

30. Ma, J., Huang, A., Yan, K., Li, Y., Sun, X., Joehanes, R., Huan, T., Levy, D. and Liu, C. (2023)
Blood transcriptomic biomarkers of alcohol consumption and cardiovascular disease risk
factors: the Framingham Heart Study. *Hum Mol Genet*, **32**, 649-658.

31. Merendino, L., Guth, S., Bilbao, D., Martinez, C. and Valcarcel, J. (1999) Inhibition of msl-2
splicing by Sex-lethal reveals interaction between U2AF35 and the 3' splice site AG. *Nature*,
402, 838-841.

32. Kato, G.J., Lee, W.M., Chen, L.L. and Dang, C.V. (1992) Max: functional domains and
interaction with c-Myc. *Genes Dev*, **6**, 81-92.

33. Makela, T.P., Koskinen, P.J., Vastrik, I. and Alitalo, K. (1992) Alternative forms of Max as
enhancers or suppressors of Myc-ras cotransformation. *Science*, **256**, 373-377.

34. Mathsyaraja, H., Freie, B., Cheng, P.F., Babaeva, E., Catchpole, J.T., Janssens, D., Henikoff, S.
and Eisenman, R.N. (2019) Max deletion destabilizes MYC protein and abrogates Emicro-Myc
lymphomagenesis. *Genes Dev*, **33**, 1252-1264.

35. Arsura, M., Deshpande, A., Hann, S.R. and Sonenshein, G.E. (1995) Variant Max protein,
derived by alternative splicing, associates with c-Myc in vivo and inhibits transactivation. *Mol
Cell Biol*, **15**, 6702-6709.

36. Babic, I., Anderson, E.S., Tanaka, K., Guo, D., Masui, K., Li, B., Zhu, S., Gu, Y., Villa, G.R.,
Akhavan, D. *et al.* (2013) EGFR mutation-induced alternative splicing of Max contributes to
growth of glycolytic tumors in brain cancer. *Cell Metab*, **17**, 1000-1008.

37. Zhang, H., Fan, S. and Prochownik, E.V. (1997) Distinct roles for MAX protein isoforms in
proliferation and apoptosis. *J Biol Chem*, **272**, 17416-17424.

38. Vuckovic, D., Bao, E.L., Akbari, P., Lareau, C.A., Mousas, A., Jiang, T., Chen, M.H., Raffield,
L.M., Tardaguila, M., Huffman, J.E. *et al.* (2020) The Polygenic and Monogenic Basis of Blood
Traits and Diseases. *Cell*, **182**, 1214-1231 e1211.

39.   Caillard, A., Sadoune, M., Cescau, A., Meddour, M., Gandon, M., Polidano, E., Delcayre, C.,
Da Silva, K., Manivet, P., Gomez, A.M. *et al.* (2018) QSOX1, a novel actor of cardiac
protection upon acute stress in mice. *J Mol Cell Cardiol*, **119**, 75-86.
40.   Giri, A., Hellwege, J.N., Keaton, J.M., Park, J., Qiu, C., Warren, H.R., Torstenson, E.S., Kovesdy,
C.P., Sun, Y.V., Wilson, O.D. *et al.* (2019) Trans-ethnic association study of blood pressure
determinants in over 750,000 individuals. *Nat Genet*, **51**, 51-62.
41.   Bacolla, A., Zhu, X., Chen, H., Howells, K., Cooper, D.N. and Vasquez, K.M. (2015) Local DNA
dynamics shape mutational patterns of mononucleotide repeats in human genomes. *Nucleic
Acids Res*, **43**, 5065-5080.
42.   Landrum, M.J., Chitipiralla, S., Brown, G.R., Chen, C., Gu, B., Hart, J., Hoffman, D., Jang, W.,
Kaur, K., Liu, C. *et al.* (2020) ClinVar: improvements to accessing data. *Nucleic Acids Res*, **48**,
D835-D844.
43.   Li, Z., Huang, Z., Liu, Y., Cao, Y., Li, Y., Fang, Y., Huang, M., Liu, Z., Lin, L. and Jiang, L. (2023)
Genotypic and phenotypic characterization of glucose-6-phosphate dehydrogenase (G6PD)
deficiency in Guangzhou, China. *Hum Genomics*, **17**, 26.
44.   Bancone, G., Chu, C.S., Somsakchaicharoen, R., Chowwiwat, N., Parker, D.M.,
Charunwatthana, P., White, N.J. and Nosten, F.H. (2014) Characterization of G6PD genotypes
and phenotypes on the northwestern Thailand-Myanmar border. *PLoS One*, **9**, e116063.
45.   Chiu, D.T., Zuo, L., Chao, L., Chen, E., Louie, E., Lubin, B., Liu, T.Z. and Du, C.S. (1993)
Molecular characterization of glucose-6-phosphate dehydrogenase (G6PD) deficiency in
patients of Chinese descent and identification of new base substitutions in the human G6PD
gene. *Blood*, **81**, 2150-2154.
46.   Wei, X., Kixmoeller, K., Baltrusaitis, E., Yang, X. and Marmorstein, R. (2022) Allosteric role of
a structural NADP(+) molecule in glucose-6-phosphate dehydrogenase activity. *Proc Natl
Acad Sci U S A*, **119**, e2119695119.
47.   Lallena, M.J., Chalmers, K.J., Llamazares, S., Lamond, A.I. and Valcarcel, J. (2002) Splicing
regulation at the second catalytic step by Sex-lethal involves 3' splice site recognition by
SPF45. *Cell*, **109**, 285-296.
48.   Dybkov, O., Preussner, M., El Ayoubi, L., Feng, V.Y., Harnisch, C., Merz, K., Leupold, P.,
Yudichev, P., Agafonov, D.E., Will, C.L. *et al.* (2023) Regulation of 3' splice site selection after
step 1 of splicing by spliceosomal C* proteins. *Sci Adv*, **9**, eadf1785.
49.   Burnight, E.R., Gupta, M., Wiley, L.A., Anfinson, K.R., Tran, A., Triboulet, R., Hoffmann, J.M.,
Klaahsen, D.L., Andorf, J.L., Jiao, C. *et al.* (2017) Using CRISPR-Cas9 to Generate Gene-
Corrected Autologous iPSCs for the Treatment of Inherited Retinal Degeneration. *Mol Ther*,
25, 1999-2013.
50.   Martelly, W., Fellows, B., Kang, P., Vashisht, A., Wohlschlegel, J.A. and Sharma, S. (2021)
Synergistic roles for human U1 snRNA stem-loops in pre-mRNA splicing. *RNA Biol*, **18**, 2576-
2593.
51.   Taliun, D., Harris, D.N., Kessler, M.D., Carlson, J., Szpiech, Z.A., Torres, R., Taliun, S.A.G.,
Corvelo, A., Gogarten, S.M., Kang, H.M. *et al.* (2021) Sequencing of 53,831 diverse genomes
from the NHLBI TOPMed Program. *Nature*, **590**, 290-299.
52.   Bailey, T.L., Johnson, J., Grant, C.E. and Noble, W.S. (2015) The MEME Suite. *Nucleic Acids
Res*, **43**, W39-49.
53.   Buniello, A., MacArthur, J.A.L., Cerezo, M., Harris, L.W., Hayhurst, J., Malangone, C.,
McMahon, A., Morales, J., Mountjoy, E., Sollis, E. *et al.* (2019) The NHGRI-EBI GWAS Catalog
of published genome-wide association studies, targeted arrays and summary statistics 2019.
Nucleic Acids Res, **47**, D1005-D1012.
54.   Paul, D.S., Albers, C.A., Rendon, A., Voss, K., Stephens, J., HaemGen, C., van der Harst, P.,
Chambers, J.C., Soranzo, N., Ouwehand, W.H. *et al.* (2013) Maps of open chromatin highlight
cell type-restricted patterns of regulatory sequence variation at hematological trait loci.
Genome Res, **23**, 1130-1141.

55.   Becker, K.G., Barnes, K.C., Bright, T.J. and Wang, S.A. (2004) The genetic association
database. *Nat Genet*, **36**, 431-432.
56.   Sherman, B.T., Hao, M., Qiu, J., Jiao, X., Baseler, M.W., Lane, H.C., Imamichi, T. and Chang,
W. (2022) DAVID: a web server for functional enrichment analysis and functional annotation
of gene lists (2021 update). *Nucleic Acids Res*, **50**, W216-W221.
57.   Liu, C., Joehanes, R., Ma, J., Wang, Y., Sun, X., Keshawarz, A., Sooda, M., Huan, T., Hwang,
S.J., Bui, H. *et al.* (2022) Whole genome DNA and RNA sequencing of whole blood elucidates
the genetic architecture of gene expression underlying a wide range of diseases. *Sci Rep*, **12**,
20167.
58.   Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M.
and Gingeras, T.R. (2013) STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*, **29**, 15-
21.
59.   Lun, A.T., Chen, Y. and Smyth, G.K. (2016) It's DE-licious: A Recipe for Differential Expression
Analyses of RNA-seq Experiments Using Quasi-Likelihood Methods in edgeR. *Methods Mol
Biol*, **1418**, 391-416.
60.   Robinson, J.T., Thorvaldsdottir, H., Wenger, A.M., Zehir, A. and Mesirov, J.P. (2017) Variant
Review with the Integrative Genomics Viewer. *Cancer Res*, **77**, e31-e34.
61.   Liu, L., Das, U., Ogunsola, S. and Xie, J. (2022) Transcriptome-Wide Detection of Intron/Exon
Definition in the Endogenous Pre-mRNA Transcripts of Mammalian Cells and Its Regulation
by Depolarization. *Int J Mol Sci*, **23**, 10157.
62.   Khodor, Y.L., Menet, J.S., Tolan, M. and Rosbash, M. (2012) Cotranscriptional splicing
efficiency differs dramatically between Drosophila and mouse. *RNA*, **18**, 2174-2186.
63.   Liao, Y., Smyth, G.K. and Shi, W. (2014) featureCounts: an efficient general purpose program
for assigning sequence reads to genomic features. *Bioinformatics*, **30**, 923-930.
64.   Liu, C., Joehanes, R., Ma, J., Xie, J., Yang, J., Wang, M., Huan, T., Hwang, S.J., Wen, J., Sun, Q.
et al. (2025) Integrating Whole Genome and Transcriptome Sequencing to Characterize the
Genetic Architecture of Isoform Variation and its Implications for Health and Disease. *Nature
Communications*, **10.1101/2024.12.04.24318434**.
65.   Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O., Willmore,
L., Ballard, A.J., Bambrick, J. *et al.* (2024) Accurate structure prediction of biomolecular
interactions with AlphaFold 3. *Nature*, **630**, 493-500.
66.   Berman, H.M., Westbrook, J., Feng, Z., Gilliland, G., Bhat, T.N., Weissig, H., Shindyalov, I.N.
and Bourne, P.E. (2000) The Protein Data Bank. *Nucleic Acids Res*, **28**, 235-242.
67.   Magala, P., Klevit, R.E., Thomas, W.E., Sokurenko, E.V. and Stenkamp, R.E. (2020) RMSD
analysis of structures of the bacterial protein FimH identifies five conformations of its lectin
domain. *Proteins*, **88**, 593-603.
68.   Gibson, D.G., Young, L., Chuang, R.Y., Venter, J.C., Hutchison, C.A., 3rd and Smith, H.O.
(2009) Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat Methods*,
6, 343-345.
69.   Liu, L., Nguyen, H., Das, U., Ogunsola, S., Yu, J., Lei, L., Kung, M., Pejhan, S., Rastegar, M. and
Xie, J. (2024) Epigenetic control of adaptive or homeostatic splicing during interval-training
activities. *Nucleic Acids Res*, **52**, 7211-7224.
70.   Schneider, C.A., Rasband, W.S. and Eliceiri, K.W. (2012) NIH Image to ImageJ: 25 years of
image analysis. *Nat Methods*, **9**, 671-675.

Figure Legends

Figure 1. Distribution of G-tract-AG signatures in the human genome and GWAS-mapped regions, and in mammals. **A**. Diagram of the genome-wide search for G-tract-AG signatures. (Upper) A successful search within the Ensembl-annotated 3´SS with the criteria: the consensus motifs G-tract and AG of identified REPAG-3´SS (n = 1000). (Lower) Search criteria for the G-tract, AG positions/ranges within the 23nt 3´SS-like sequences in the genome, with the result summary and G-tract consensus motifs at the bottom. #: n = 700, E-value =1.9e-478; ##: n = 419, E-value =3.6e-122. **B**. Enrichment index of SNVs within G-tract-3´SSL among GWAS SNVs (input n = 128, 991, MAF $\geq$ 0.05, $p <$ 5E-08). Random SNVs from the TOPMed (n = 1000 sets, each with 128, 991 SNVs (RND), matched by MAF bins and TSS distance, excluding GWAS SNVs), were used as background (Index = 1.0) for calculating the enrichment ratios of the abundance (GWAS/RND). p-value: by one-sample t-test. NS: non-significant. **C**. Genomic location profile of the GWAS SNVs within G-tract-3´SSL (n = 41,353), showing their mainly intron and intergenic locations. **D**. The G-tract-3´SSL of the *MAX* (Left) or *QSOX1* (Right) aligned among mammalian species, with human SNV variant sequences as the top two. 3´SSL MaxENT scores are to the right of each sequence. Black dot: location of potential branch point. Dotted box: exon, and heavy black line: intron. Shaded box: exon already emerged in humans as a tissue-restricted alternative exon. **E**. PhyloP scores (470 mammals) of the motifs and SNVs from the GWAS database, at the human 3´SSL of G-tract-AG signatures. Scores are Means $\pm$ SEM, n = 3113, 3102, 3105, 3094, 3136 (GWAS); 104,705, 104,613, 104,486, 104,174, 105,758 (non-GWAS), for the G-tract, AG, SNV, G-Tract-AG and 3´SSL groups, respectively. [****]: $p <$ 3.5E-8.

Figure 2. Functional validation of the NCR G-tract-AG signatures and SNVs: G-tract repression of splicing from the AG and disruption of the repression by SNV. **A**. Diagram of the SNV rs762810 in the genomic structure of the *MAX* gene transcripts, with the mammal-specific SNV region in the enlarged

snapshots of GTEx RNA-Seq peaks and multiz alignment of 100 vertebrates from the UCSC Genome

Browser. *: transcripts containing E4 (ENST00000557746.5 and NM_001407112.1). None of the other

53 GTEx tissues reaches the same level of E4 RNA-Seq peaks as in the blood. **B**. Splicing reporter assay

of the G-tract-3´SS of the SNV allele in human embryonic kidney HEK293T cells. Intron/exon structure

and sequence around the rs762810 loci with either the allele of G or T are above the agarose gels, which

show the RT-PCR of spliced products. **C**. The SNV effect in genotype-RNA-Seq matched human

samples of white blood cells from the FHS cohort, demonstrating the RNA-Seq reads at the G-tract-

3´SS with different genotypes of the SNV alleles. **D**. Relationship between the 3′ AG usage (measured

by the RNA-Seq reads, Mean ± SEM) and number of Gs in the G-tract (left panel, [a]:7 or 8 Gs) or distance

(nt) of the last G of $(G)_3$ tracts to the 3′ AG (right panel) in 100 FHS participants of G/G genotypes. The

corresponding 3′ AG usage among variant genotypes G/H or H/H (H: non-G nucleotides, A, C or T) is

plotted in the same graphs for comparison. N = 171, 37, 19, 8, 8 (left panel), and 63, 25, 12, 13, 19, 7,

10, 15, 6, 5, 7 (right panel) participants. [***]: $p < 0.005$, between G/G and variant genotype of the same

number of Gs. [****]: $p < 0.002$ between $(G)_3$ or $(G)_4$ and their respective longer G-tract G/G genotypes.

[*****]: $p < 3.5E-27$ between G/G and non-G variant genotypes of the $(G)_3$-tracts. Note: at 2nt or 6nt

upstream (right panel), $(G)_3$ or the H/H homozygote, respectively, exhibited stronger effects than that

observed at their flanking positions ($p < 0.05$). However, these position effects need a larger sample size

to validate.

Figure 3. REPAG and SNV effects on the two transesterification steps of splicing. **A**. Left, diagram

of the *in vitro* splicing reporters, AdML vector, REPAG-WT and REPAG-Mut and their 3´SS sequences.

Right, examples of the *in vitro* splicing lariat products and pre-mRNA. N = 3-6 experiments of the three

reporters. **B**. Time course of lariat product generation in the 1[st] and 2[nd] steps, after normalizing each

product to its peak band intensities during the time course of the experiment. [***]$p < 0.001$, [*]$p < 0.05$, n

= 8 pairs in total at two (10 and 15min) time points. **C**. Time course of the conversion of the 1st step

lariat intermediate to the 2nd step lariat, after normalizing the lariat intermediate percentages of total

lariats (1st step lariat intermediate plus 2nd step lariat) in each reaction to the highest percentage during

the time course of the experiment. $^{a}p < 9E\text{-}09$, n = 12 pairs in total; $^{b}p < 4E\text{-}06$, $^{c}p < 0.001$, n = 9 pairs

in total, at three (15, 30 and 45min) time points of 3-6 independent experiments. Blue dots and SEM

(standard error of the mean) error bars: half conversion time (min). The best fitted curve (with minimal

Chi-squared values, $X^2 < 0.007$) was used to calculate the mean half conversion time of the 1st step lariat

intermediates in each experiment by their equations in Igor Pro (Wavemetrics).

**Figure 4. Disruption of the G-tracts of the signatures by human genetic variants *in cis* among

GWAS-mapped genes, protein-coding genes and genetic disease genes**. **A**. Significantly enriched

GAD (Genetic Association Database) disease classes among the 3202 GWAS-mapped genes that harbor

G-tract-AG signatures *in cis* with their G-tracts directly disrupted by GWAS SNVs, by DAVID

functional clustering analysis. Random: mean of the -log10(*p*-value) of 3 sets of the same number of

randomly picked human genes analyzed using the same parameters. 1. Chemdependency, 2. Metabolic,

3. Cardiovascular, 4. Pharmacogenomic, 5. Hematological, 6. Immune, 7. Renal, 8. Developmental, 9.

Cancer, 10. Unknown. **B**. Percentages of the GWAS-mapped or genome-wide genes that harbor such

signatures and SNVs. N = 3,202 and 23,479 genes for GWAS and the genome, respectively. **C**. Total

number of such SNVs *in cis* within the host genes among the databases screened. **D**. A reasonable

explanation provided by the G-tract-AG signature and its disruption to account for the unexplainable

protein deficiency and genetic disease caused by a rare genetic variant, the SNV c.392G->T of the *G6PD*

gene. Disruption of the G-tract motif by the variant causes partial exon skipping and protein truncation

at the N-terminal. The AlphaFold3.0-predicted structure of the wild type or missense variant protein

aligned well (RMSD = 0.925 or 0.885, respectively), but the truncated protein failed to do so (green,

RMSD = 14.500) with the corresponding N-terminal region (HQSD……FREDQI) of the electron microscopy structure (magenta, PDB #7UAG).

A

GWAS rs762810, Eosinophil percentage of white blood cells, *p* = 6E-10, SpliceAI = 0.33, in the *MAX* gene

A

B

C

D

- *G6PD*, NM_001042351.1, c.392G>T, rs137852341, chrX:154535261, G131V, < 20% activity in hemizygotes

- Mutation in the (G)$_4$ of the G-tract-AG Signature in Exon 5

- More dramatic change explained by G-tract mutation: increasing 3′SSL MaxENT score from 5 to 6.9 and aberrant splicing, deletion of an N-terminal peptide region critical for NADP$^+$ binding and catalysis.

In-Frame Deletion
(…PFFK-(del44aa)-ANRL…)

Aberrant splicing -> 132nt deletion -> 44aa deletion (90 – 133, Y112) and structural change

Supplementary Figures (S_Figs.)

S_Figure 1. **168 millions of AG/GT dinucleotides as potential cryptic splice sites in the human genome. A**. Total number of AG/GT dinucleotides within SS-like sequences with MaxENT scores > 0.1 on both strands of the human genome hg38, with the relatively much smaller numbers of the annotated AG/GT 3′SS and 5′SS in the Ensembl for comparison. **B**. Index ratios of the MaxENT scores (Mean ± SEM) of sequences harboring the dinucleotides on either strand of chromosomes, assuming the AG/GT as the SS dinucleotides, relative to a subset of annotated constitutive exon's SSs (n = 5,010 exons) in the Chr. 1 of the human genome hg38. The means of the 3′SS and 5′SS scores are 6.7 (± 0.07, SEM, minimal -33.57 – maximal 15.33, >0.1: 92.6%) and 6.96 (± 0.07, SEM, minimal -26.61 – maximal 11.81, >0.1: 94.0%), respectively. The scores of these AG/GT dinucleotide sites in the genome are half of the constitutive SS's on average, posing significant challenges with the potential for cryptic splicing under appropriate conditions (e.g. *trans*-acting factors, signaling or cell/tissue types, etc. spatial or temporal changes).

S_Figure 2. **Length (nt), location and phyloP score of the G-tracts in the genomic signatures. A**. Length distribution of G-tracts. n = 7916 G-tracts, from 6114 GWAS (*p* < 5E-08) SNV's G-tract-AG-3′SSL. **B**. MEME consensus motifs of G-tract-AG 3′SSL of 1998 sQTL within human genes, E value = 1.4E-442, showing the enrichment of Gs at the 3′SSL. Black dots indicate the phyloP scores from 470 mammalian species at each position of >107K sequences within human genes.

S_Figure 3. The human G-tract-AG signature is conserved in mammals only. Shown are 10 examples of the manually examined 37 SNVs that disrupt the G-tracts of the signatures in the MultiZ alignment of 100 vertebrates. Screenshots, taken from the UCSC Genome Browser, indicate their <u>conservation in mammals only</u>, in contrast to their flanking exon regions that are also in the other vertebrates like birds or fish. *: The rs10788958 alignment in chicken does not contain the G-tract nor AG.

S_Figure 4. **Divergent sequence and 3´SS strength changes of the G-tract-3´SS of the *MAX* gene in some mammalian species**. **A**. Percentages of species containing the G-tract, its disruption and predicted exon 4 (E4) inclusion among 6 of 62 mammalian species, based on: higher 3´ and 5´ SS-combined MaxENT scores than the human reference sequence, presence of 3' AG and branch point motifs, shortened $(G)_4$- tract and intact ORF. H: A, C or T. **B**. In comparison with the human reference variant G (Left), 3´SS MaxENT scores of E4 of representative mammalian species that are higher (Middle) or lower (Right), in comparison with those of the upstream E3. **C**. Additional examples of 3´SS sequences showing the divergent changes of the *MAX* G-tract-3´SS motifs, besides the SNV-disruption of the G-tract among mammals, with human and guinea pig sequences for comparison. Upper panel: G-tract-3´SS alignment. Lower panel: the less divergent upstream 3´SS of exon 3 of the same species for comparison. Nucleotides different from the human reference genome sequences are shaded in black. Black dots: positions of potential branch points in some species. Py: polypyrimidine tract. Splicing prediction: by MaxENT scores, essential 3´SS motifs and/or transcript evidence. 1: positive or more likely; 0: negative or less likely. Note that here we use 3´SS since it is an authentic splice site in humans (see details on functional validation later).

S_Figure 5. Validation of the effect of SNV rs3767199-T on splicing through the intronic G-tract-AG signature of the *QSOX1* gene by minigene splicing reporter assay. **A**. Diagram of the SNV region in the intron 1 of the *QSOX1* gene, adapted from the UCSC genome browser. **B**. Left, repression of the SNV-3′SSL from cryptic splicing in human tissues: snapshot of the SNV region in GTEx showing RNA-Seq peaks of the upstream exon 1 in several tissues but none downstream the SNV, nor in the rest of the 54 tissues, with the aligned 100 vertebrate conservation profile below, both from the UCSC genome browser. Note: the GGG-AG signature is not in *X. tropicalis* though it is also aligned locally. Right, percentages of 67 vertebrate species with potential 3′SS usage or the 3′SSL motifs. The GGG and GGT are in only a small percentages of the species and all are in mammals only. 3′SS usage (only with human SNV) based on: presence of 3′AG, shortened G-tract and higher MaxENT 3′SSL score than the human′s reference sequence. **C**. Left, diagram of the minigene splicing reporter, derived from the DUPST1 harboring the STREX exon (Xie & Black, *Nature*'01). Right, agarose gel of the RT-PCR products of cytoplasmic RNA of the splicing reporters transiently expressed in HEK293T cells. C: negative control with untransfected-cellular RNA. The $(G)_3$ repression was also observed when in front of a 51nt beta-globin exon.

S_Figure 6. SNV-association with *MAX* exon 4 usage in blood cells in GTEx. **A**. Snapshot of the IGV view of the RNA-Seq reads peaks of the rs762810-T allele-associated exon 4 in the intron 3 of MAX ENST00000557746.5, among a number of GTEx tissues in the UCSC Genome Browser. The red arrowhead points to the location of the SNV and the upstream 3´SS of exon 4. Black arrows: direction of the *MAX* gene. **B**. GTEx result of the intron 3 excision ratios with different genotypes of the SNV rs762810.

S_Figure 7. AlphaFold 3.0-predicted structural conformation changes in the interaction between the G-tract-G or -U RNA variants and the trans-acting splicing repressor hnRNP H1 qRRM1 and 2 domains. Shown are the MAX G-tract 5´-UGGGGCU-3´ (Gs in red) or its rs762810-T variant 5´-UGGGUCU-3´(SNV U in green) in complex with the repressor. RMSD = 9.211 when aligned between the two complexes, indicating significant structural differences induced by the variants (p < 0.0001), in contrast to an average RMSD of 0.559 (±0.225, SEM) of 21 protein-RNA (≤ 30nt) complexes aligned with their experimental structures from the PDB in PyMol. The high-confidence RNA-protein complex show that the U-variant dramatically relaxed the G-tract RNA interaction with the trans-acting splicing repressor hnRNP H1. The complex switched from a compact key-lock pocket for the G-variant to a wide open configuration for the U-variant, in a manner similar to the crystal structure changes for different G-tracts (S. R. Penumutchu et al., *J Am Chem Soc* 140, 11661-11673 (2018)), suggesting destabilization of the G-tract-repressor interaction by the U-variant.

A

MAX — bHLH_SF Superfamily — NLS — Growth-Suppressive, NM_002382 (a)

deltaMAX — bHLH_SF Superfamily — Growth-Promoting, ENST00000555667.5 (b)

MAX + Exon 4 (T-allele) — bHLH_SF Superfamily — NLS — C/N, ENST00000557746.5 (c)

— NLS — N, NM_001407111.1 (d)

Products from Transcript Variants c & d: Growth-Promoting as well?

B Validated MAX NLS peptide by Kato, G&D'92   Peptide from Exon 4-Included ENST00000557746.5

DGGSDSSSESEPEEPQSRKKLRMEASAAAAAAAAAGALPAVGITPGIRCLKRALRSLEWNVLWESES

17SRKKLR22

0.08 (2 State HMM Dynamic Model)

1.25

0.25

http://www.moseslab.csb.utoronto.ca/NLStradamus/; Nguyen Ba AN, Pogoutse A, Provart N, Moses AM. **NLStradamus: a simple Hidden Markov Model for nuclear localization signal prediction.** *BMC Bioinformatics.* 2009;**10(1)**:202.

S_Figure 8. Domain loss of the MAX protein isoforms generated from the transcripts with the SNV-T allele-promoted exon 4 usage. **A**. Truncation of the COOH or NH$_2$ terminals in the isoforms from the exon 4-included transcripts, in comparison with the representative full length or short (delta) isoforms with known effects on cell growth. **B**. Significant loss of the NLS signal strength from the transcript ENST00000557746.5 containing exon 4. Prediction based on Nguyen et al.: a simple Hidden Markov Model for nuclear localization signal prediction. *BMC Bioinformatics.* 2009;10(1):202, with the known NLS peptide and ENST00000557746.5-encoded NLS-candidate peptide input together with a poly(Alanine) linker.

S_Figure 9. MAX variant genotype distribution among cancer patients of different tissues and exon 4 ectopic inclusion in melanoma cells with the T-variant. A. Tissue-dependent ratio of cancer prevalence among the T/T over G/G genotypes of SNV rs762810 of the *MAX* gene. Shown is the log2 ratios of the cancer prevalence among different tissues or histological categories over that among the cohort after normalization to (divided by) the allele frequencies in the cohort. From left to right: 1. hematopoietic and reticuloendothelial systems, 2. trachea bronchus, and lung, 3. only one cancer, 4. colon, 5. lymph nodes, 6. rectum, rectosigmoid junction, anal canal and anus, not otherwise specified, 7. control: all FHS participants with genotype data for SNP rs762810, 8. healthy controls from FHS with genotype data for SNP rs762810, 9. prostate gland, 10. urinary bladder, 11. adenocarcinoma, 12. female breast (excludes skin of breast), 13. basal cell carcinoma, 14. corpus uteri or endometrium, 15. skin, excludes skin of labia majora, skin of vulva, skin of penis and skin of scrotum, 16. with multiple cancers, 17. squamous cell carcinoma, 18. duct carcinoma, 19. nevi & melanomas, in an FHS cohort with sex- and age-matched genotype data for the SNV rs762810 (***: $p < 4E\text{-}11$, hypergeometric density test, n = 8,481 participants). **B**. Exon 4 ectopic inclusion (>50%) in patient melanoma-derived cells with the T variant only. Raw RNA-Seq sequence and BAM file derived from SRA #SRR31631252. Another sample with the T variant only and similar inclusion is #SRR31631251. The extra AG at the exon start is likely from the last two nts of exon 3 based on the gene sequence and RNA-Seq reads.

S_Figure 10. A group of G-tract mutations/variants identified in human genetic diseases in the HGMD database. Shown are the changes of the consensus motif sequences by G-tract mutations in comparison with the wild type sequences, as analyzed using the MEME. The mutations are from that published by Bacolla, et al., NAR'15 (43)5065.

S_Figure 11. Underrepresentation of the G-tract-AG signatures in current splicing-prediction software. Shown is the rarity of the G-tract-3´SSL SNVs among SpliceAI output results by comparison of the percentages of G-tract-3´SSL-overlapping SNVs with the percentages of the SpliceAI output among GWAS SNVs (n = 128,991 and 153,477 input GWAS SNVs in total, respectively). Of the 166 SpliceAI-scored SNVs above the confidence level of >0.5, only 7 are G-tract-disrupting SNVs (0.005% of GWAS). The G-tract-disrupting SNVs at 3´SSLs, 5.62% GTEx sQTLs reported here, are thus expected to add significantly to the list of splicing-regulating SNVs without the limitation of SNV distance to annotated splice sites by SpliceAI or other programs.