Title: Alpha-synuclein overexpression reduces neural activity within a basal ganglia vocal

nucleus in a zebra finch model

Running title: Alpha-synuclein overexpression reduces neural activity in a finch song

nucleus

Brian R. Dominguez[1], Gabriel Holguin[2], Madeleine S. Daly[1], Reed T. Bjork[1], Stephen L.
Cowen[2,3], and Julie E. Miller*[1,4]
9

[1]Department of Neuroscience, University of Arizona, Tucson, Arizona, United States of
America.
[2]Department of Psychology, University of Arizona, Tucson, Arizona, United States of
America.
[3]Evelyn F. McKnight Brain Institute, University of Arizona, Tucson, Arizona, United States
of America
[4]Departments of Speech, Language and Hearing Sciences, Neurology, and the BIO5
Institute, Tucson, Arizona, United States of America.

*Corresponding Author:

Julie E. Miller, Ph.D.; E-mail: juliemiller@arizona.edu

Declaration of Competing Interest: J.E. Miller is a 2025 member of the "T2 Targets to

Therapies" advisory working group for the Michael J. Fox Foundation for Parkinson's

Research.

Other author emails:
Dominguezbrian2@gmail.com
gholguin@scripps.edu
madeleinedaly@arizona.edu
bjork2@arizona.edu
scowen@arizona.edu

Abstract

Vocal motor impairments, including changes in pitch, loudness, and timing are prevalent in Parkinson's Disease (PD) and a target for early intervention and treatment, yet the neural mechanisms underlying these impairments are not understood, motivating work in animal models. The adult male zebra finch songbird is uniquely poised for these studies given vocally-dedicated brain nuclei and a quantifiable output (birdsong). Our prior publication revealed that injection of an adeno-associated virus (AAV5) expressing the human (h) alpha-synuclein (h*SNCA*, a-syn) gene into basal ganglia vocal nucleus Area X results in higher levels of insoluble a-syn protein and changes reminiscent of human PD including softer, shorter, and reduced number of vocalizations compared to AAV5 controls. Here, we test the hypothesis that AAV-h*SNCA* overexpression reduces the firing rate of specific neuronal sub-types in Area X using *in vivo* recordings in anesthetized finches. Five classes of neurons were differentiated in AAV-treated finches based on waveform width (narrow vs. wide) and firing rates (low vs. fast). Wide-waveform/low-rate activity is a consistent feature of striatal medium spiny neurons (MSNs), a dominant cell type in Area X and in mammalian basal ganglia. Reduced firing rates and enhanced post-peak rebound were detected in the AAV-h*SNCA* group for putative MSN neurons compared to AAV controls. No differences in firing rate nor waveform shape were detected for the narrow waveform neurons. Our findings provide the first characterization of early a-syn-driven neural activity changes in vocal control neurocircuitry.

Introduction:

Pathological accumulation of alpha-synuclein (a-syn) is believed to contribute to multiple neurodegenerative diseases, including Parkinson's disease (PD). Accumulation of toxic a-syn species culminates in the development of Lewy Bodies and Lewy Neurites that are detected in inherited and sporadic cases of PD, dementia with Lewy Body disease, and Multiple Systems Atrophy (1-5). How abnormal increases in a-syn protein expression and its aggregation disrupts ongoing neuronal activity in the basal ganglia (BG) is not understood, yet this information is critical to developing cell-specific therapeutic targets (6-8). One strategy is to examine how elevated a-syn protein interferes with the ongoing firing activity of striatal neurons.

Elevated levels of a-syn protein at cortico-striatal synapses can result in synaptic failure and loss of cortico-striatal plasticity based on electrophysiological recordings from rodent brain slices (6-11). Striatal Medium Spiny Neurons (MSNs) show alterations in dendritic spine density, interactions between a-syn aggregates and their NMDA receptors, and reduced efficacy of cortical pre-synaptic glutamatergic input that affects their firing rate (12-15). Furthermore, burst firing is a characteristic feature of MSNs (16), and dysregulated MSN burst firing may contribute to PD as shown in 6-hydroxydopamine (6-OHDA) models (17). A recent study used induced human pluripotent stem cells from patients with Multiple System Atrophy of the Parkinsonian type that were differentiated into MSNs. These MSNs showed hypoexcitability in the form of reduced spontaneous firing rates and reduced frequency and amplitude of miniature post-synaptic currents. These changes were also associated with elevated a-syn release (18). How *in vivo* cell-specific neuronal firing patterns in the basal ganglia relates to the behavioral output is

unclear and a critical gap. The scant literature on *in vivo* recordings in rodent h*SNCA*

overexpression models comes from nigral, thalamic, and hippocampal neurons

evaluating movement and cognition (19-21). Pairing electrophysiological changes in

basal ganglia circuitry with vocal behavior offers an early and critical entry-point into

furthering our understanding of the effects of alpha-synucleinopathies on neural circuits.

Voice and speech deficits are prevalent in PD, impacting communication and

quality of life for at least 90% of individuals (22, 23). Early stage vocal deficits (a soft,

monotonous voice, altered speaking rate) often arise years before the limb motor

symptoms, offering strong potential as an early diagnostic tool, strengthened by insights

using machine learning algorithms and smartphone applications (24-37). Our group and

others have modeled PD-related vocal deficits using pharmacological and genetic

approaches in rodents and songbirds with data from these experiments implicating the

h*SNCA* gene in driving early vocal dysfunction (38-46).

Songbird models offer distinct advantages over rodents in exploring how the brain

encodes sensory and motor information, shaping vocal learning and production in

response to feedback (47). In the zebra finch species, the males sing and the females do

not. The male song control system in the brain **(Fig 1A)** has strikingly similar neural

organization and physiology of the basal ganglia to mammalian species (48-50). Several

reviews   detail   the   anatomical,   developmental,   genetic,   functional,   and

electrophysiological features of this system and their close similarity to human speech

and auditory centers in the human brain (51-57). Vocal learning and on-going song

modification in the zebra finch model can be investigated through the anterior forebrain

pathway (AFP), a structure homologous to the cortico-basal ganglia-thalamo-cortical loop

in humans and other mammals (50). Song-dedicated brain region Area X in the basal

ganglia shows high genetic similarity to the striatum, a major human speech control area,

and to mammalian neuronal populations (48, 54, 58, 59).

Fig 1A. Finch song control circuitry. Song-dedicated brain nuclei in male finch are

shown (HVC and Area X: proper names; RA: robust nucleus of the Arcopallium, lMAN:

lateral magnocellular nucleus of the anterior nidopallium). Solid lines: cortico-brainstem

circuits for vocal production (finch: HVC to RA to brainstem); dashed lines: cortico-basal

ganglia-thalamo-cortico circuits for on-going vocal learning and refinement (lMAN-Area

X-DLM-lMAN). Cx – cortex. BG – basal ganglia. DLM – dorsal medial nucleus of the

thalamus. Th – thalamus. MB – midbrain. HB – hindbrain. Cb – cerebellum. OB – olfactory

bulb. Area X (violet) contains striatal, globus pallidus, and interneurons whereas human

BG has separate striatal and pallidal areas. Auditory input (not shown) to finch vocal

nuclei is also well-studied (60, 61). Figure modified from Medina et al. (2022).

Electrophysiological recordings in singing finches and anatomical tracings show

that MSNs are the *dominant cell type* in Area X and drive the responses of globus pallidus-

like (PAL) projection neurons (50, 58, 62). As in mammals, MSNs receive glutamatergic

(Glu) inputs from cortex, and globus-pallidus (GP, PAL) type neurons project to the

thalamus which then relays input back to the cortex. In the song control system (**Fig 1A**),

cortical song nuclei HVC and lMAN provide Glu input to GABAergic MSNs and PAL

neurons within Area X (63). While studies of MSN activity in animal models and humans

are difficult due to low spontaneous activity (14, 64), finch MSNs are dynamically active

during singing and show temporally precise relationships to song structure (62, 65).

The PAL nerve terminals end in the dorsal medial thalamic nucleus (DLM) (63).

DLM projections to a neuronal sub-population within the premotor nucleus, HVC, can also

modulate syllable transitions (66). High tonic firing in these PAL neurons inhibits DLM,

reducing excitation onto lMAN and then song nucleus RA, leading to altered song output

including variations in pitch (55, 62). Area X and lMAN neurons exhibit firing variability

that is related to birdsong features (55, 62). The well-characterized neuronal populations

in these brain nuclei and circuitry connections make this a powerful model to study the

impact of alpha-synucleinopathies on vocalizations.

In our prior work, we showed that targeting Area X with an adeno-associated virus

(AAV) drives increased expression of the human (h) wild-type *SNCA* gene leading to PD-

like changes in the song – including decreased vocalizations (singing), duration, and a

softer, more monotonous vocal output (45). We also showed that a-syn overexpression

results in protein aggregates in Area X cell bodies (46). Here, as a critical first step, we

categorize cell-specific firing patterns and waveforms in Area X that are affected by a-syn

overexpression compared to controls using tetrode recordings in anesthetized finches

immediately following singing behavior. Our working hypothesis (**Fig. 1B**) is that h*SNCA*

overexpression in Area X will decrease firing rates and alter burst-firing in putative Area

X MSNs. We tested this hypothesis by recording from Area X neurons in anesthetized

zebra finches.

Fig 1B. **Hypothetical model for a-syn driven changes in neural activity in Area X and consequences on song control circuitry.** Cortical vocal nuclei (lMAN, HVC) provide glutamatergic (Glu) input to Area X neurons. Striatal Area X MSNs provide GABAergic input to globus-pallidus like (GP, PAL) neurons including those that inhibit thalamic nucleus DLM. Here, we *predict* that abnormally elevated levels of AAV-driven a-syn protein in Area X could alter intrinsic membrane properties of MSNs and/or reduce Glu input to MSNs. Consequently, MSNs would reduce their firing rate (red arrow), leading to increased PAL projection neuron firing and increased inhibition of DLM. Increased DLM inhibition could be responsible for reduced singing and changes in amplitude (loudness) and duration of the song syllables in finches with the AAV-h*SNCA* phenotype (45).

Materials and Methods:

2.1. Finch subjects, song recording and analyses. All animal use was approved by the Institutional Care and Use Committee at the University of Arizona. **Fig 1C** shows our experimental workflow and tetrode array.

Fig 1C. Tetrode array and experimental timeline. Schematic of AAV constructs and coronal section showing bilateral song nucleus Area X (purple) from Medina et al. (2022). Song was collected within three days of the bilateral injection into Area X and then again at two months post-virus injection followed by tetrode recordings in anesthetized finches within five days. At the end of the recording morning, the finch was humanely euthanized,

and the brain removed post-mortem for cryosectioning followed by immunohistochemistry

to confirm AAV and a-syn protein expression along with tetrode placement.

Animal procedures followed Medina et al. (2022). Adult non-breeding male zebra finches

(approx. 197-365 days post-hatch at time of euthanasia) were moved into individual

sound attenuation chambers (Eckel Noise Control Technologies) and acclimated under a

14:10 h light-dark cycle for at least two days prior to song recordings and neurosurgery.

Behavioral experiments were conducted in the morning from lights-on for two hours to

minimize effects of circadian rhythms on gene expression within Area X and to match our

prior study (45) along with ensuring singing-driven neuronal activity just prior to

conducting the tetrode recordings in anesthetized finches. Songs were acquired from

individual males singing alone (known as undirected song) using free-field omnidirectional

microphones (Shure 93) suspended at set points over the center of the cage, connected

to an audiobox (Audiobox: 44.1 kHz sampling rate per 24-bit depth; Niles, IL) and

recorded with Sound Analysis Pro 2011 (SAP 2011, (67) http://soundanalysispro.com/).

Song data collected and used for this study can be found at the University of Arizona

ReData repository upon publication (10.25422/azu.data.27868266). A basic unit of

birdsong is defined as a motif; a motif is comprised of a sequence of repeated syllables

(**S1 Fig A**). Wav files from the two-hour recording period were joined (Shuang's audio

joiner) and then the entire file was viewed in Audacity (https://www.audacityteam.org/).

The most frequently occurring motif was then identified in the joined wav file for two pre-

surgical mornings (time point 0) and then two post-surgical mornings (time point, 2

months, **Fig 1C**) by an investigator blind to the AAV condition. The two-month timepoint

was selected as the endpoint for tetrode recordings because of our prior publication

showing the most prominent song and protein changes occurred at this time point in the

AAV-h*SNCA* finches (45, 46). Following morning lights-on, the first motif was identified in

the wav file and then 25 consecutive renditions of the individual syllable wav files were

exported from Audacity from the four mornings. Up to five unique syllables within a motif

for each bird were included in the analyses.

Syllable wav files were processed in Matlab R2014 using the SAP SAT Tools box

with code provided by Dr. Nancy Day (Whitman College) to extract individual acoustic

features as a feature batch function. Raw amplitude (intensity) measurements were

calibrated based on sound chamber location as in our prior study (68). Mean, standard

deviation, and coefficient of variation (CV, standard deviation/mean) were then calculated

by averaging 50 scores (25 syllable copies x 2 recording sessions) across time points 0

(pre-AAV injection) and 2 months post-AAV injection for each syllable within the bird's

motifs to account for natural day to day fluctuations in song. For group level comparisons,

a normalization score was obtained per bird by dividing timepoint two months (2) post-

injection by pre-injection (0). After verifying that scores were not normally distributed,

scores were then compared using the Mann Whitney U test in GraphPad Prism (version

10) between AAV-h*SNCA* and AAV-GFP groups for all syllable types combined and then

sorted by syllable type (**S1 Table**) given prior findings on a-syn differential effects (45).

Syllables were sorted into noisy, mixed, or harmonic categories based on visual

inspection and Wiener Entropy (WE) scores following our prior publications (**S1 Fig**) (45,

69). We focused our analyses on comparing duration, amplitude, and WE scores between

the two AAV groups, and their correlations with the electrophysiological analyses for an

individual bird (see section on **Statistics** and **S1-2 Figs**). A subset of these birds was

used in our separate publication evaluating regional distribution of a-syn protein in finch

song nuclei and linear correlations to syllable features in Area X and lMAN (46).

As described in Medina et al. (2022), duration is the length of each syllable (in seconds),

and amplitude is the intensity or loudness of the syllable (in decibels-dB). WE is a

measure of uniformity in the power spectra (i.e., structure of the syllable), where scores

are calculated on a logarithmic scale; noisy syllables have values approaching zero while

harmonic syllables have more negative scores (approaching infinity). The coefficient of

variation (CV) is calculated by dividing the standard deviation of 50 syllable renditions by

the mean score and represents variability across motifs for an acoustic feature,

modulated by the lMAN-Area X-DLM-lMAN loop (70).

2.2 Viral injections: An adeno-associated virus (AAV) encoding the h*SNCA* gene

(AAV5-CBA-*SNCA*, N = 6) was bilaterally injected into Area X while finches were under

isoflurane anesthesia with oxygenation (2-3%, 0.5 liters/min oxygen). A control AAV

encoding for green fluorescent protein (AAV5-CBA-eGFP, N = 6) was also bilaterally

injected into Area X of a separate control group. Both viruses were under the control of

the broad promoter, chicken beta-actin (CBA). For the injections, stereotaxic coordinates

were used from the bifurcation of the mid-sagittal sinus: 3.1 mm rostral-caudal, 1.62 mm

medio-lateral, a depth of 3.1 mm, and head angle of 40 degrees. These coordinates avoid

indirect targeting of song nucleus lMAN with virus as previously described (45). Viruses

(AAV5) were made and obtained from the University of North Carolina Viral Vector Core

through the support of the Michael J. Fox Foundation for Parkinson's Research

(https://www.michaeljfox.org/research-tools).

A glass pipette was fitted into a Nanoject II pressure-injector and back-filled with mineral

oil, then loaded with either the control or the a-syn virus. Approximately 500 nL of virus

was injected at a rate of 27.6 nL/injection every 15 seconds for a total of 18 injections.

After five minutes, the pipette was slowly retracted and the tip was visually inspected for

clogging. Birds were then returned to their sound chambers following post-operative

monitoring and resumed singing within a few hours after surgery.

2.3 Tetrode Implantation and Electrophysiology

Surgical procedure and implantation: Approximately two months post-virus

injection, finches were anesthetized with isoflurane (2-3%, constant flow rate of 0.5-0.8 L

of oxygen/min) and underwent acute electrophysiological recordings of right hemisphere

Area X. Breathing rate was monitored throughout the experiment, and an infrared heating

pad (Kent Scientific) maintained body temperature at 37ºC. The same stereotaxic

coordinates were used for the AAV injections targeting Area X. **Ground wire insertion:**

A ground/reference wire was inserted into the left hemisphere, contralateral to the virus

injection site (mid-sagittal sinus: 3.1 mm rostral-caudal, 1.62 mm medio-lateral). After

performing the craniotomy, the dura was pierced with a 30G needle, and the end of the

ground wire was lowered to a depth of 2-3 mm. Gel superglue was used to secure the

wire to the acid-etched skull (acid from the Metabond kit). Zip Kicker (Pacer Technology)

was used to rapidly cure the superglue. **Electrode array insertion:** The existing

craniotomy on the right hemisphere (used for virus injection) was expanded to a diameter

of ~1.5 mm to better accommodate the electrode array and to have access to dura that

was undamaged during virus injection. Tetrode position was confirmed in Area X post-

mortem following electrophysiological recordings (**Fig 2**).

Electrophysiological recordings: Neurons from Area X were recorded using a

custom-made 800 μm diameter tetrode array composed of 8 twisted-wire tetrodes.

Tetrodes were constructed from polyamide-insulated nickel chromium steel wire (~12 μm

in diameter; see **Fig 1C**). Prior to insertion, tetrodes were coated with DiI to allow

anatomical tracing of the electrode path (**Fig 2A**). The tetrode array was lowered in ~100

μm increments until the tip reached Area X at ~3100 μm (right hemisphere) and allowed

to stabilize for 5-10 minutes prior to recording. Neural data was acquired at 20 kHz using

the INTAN neural recording system (Intan Technologies, Inc.). Neural recordings from

Area X neurons occurred in 5-minute blocks. After each 5-min block of data was acquired,

the electrode array was lowered ~50 μm and another recording block was acquired. All

recordings were restricted to Area X (3100-4100 μm). Data was spike sorted using

Kilosort   v2.5   (https://zenodo.org/records/4482749)   (71)   and   Phy

(https://github.com/cortex-lab/phy). Methods for cell-type estimation are described in

Results.

Data/Statistical Analysis:

All single-unit data was analyzed using Matlab (R2025a). All Matlab analysis code is

available upon publication at: https://github.com/CowenLab/ZebraFinch.

Measurement of burst activity: The temporal structure of the firing activity of individual

neurons was quantified using "local variance" (LV), a measure that assesses the

variability of inter-spike intervals (ISIs) and is robust to changes in firing rate (72). LV is

computed from the vector of ISIs for each neuron as shown in the formula below:

$$LV = \frac{3}{n-1}\sum_{i=1}^{n-1}\left(1+\frac{4I_iI_{i+1}}{(I_i+I_{i+1})^2}\right)\left(1-\frac{4R}{I_i+I_{i+1}}\right), \qquad (1)$$

where I_i and I_{i+1} are the i-th and $i+1$ ISI and n is the number of ISIs. LV scores ~0

indicate tonic firing, scores ~1 show Poisson firing, and scores >1 indicate burst-like firing

patterns.

Parametric and non-parametric tests were used for hypothesis testing. For group-level

analysis of firing-rate data (n = 201 categorized neurons from n = 12 finches) non-

parametric tests were used as firing rates do not follow normal distributions (73).

Consequently, rank-sum tests were used for two-group comparisons and the Kruskal-

Wallis test was used for multi-group data. The Holm-Bonferroni correction was used to

correct for multiple comparisons. For normally distributed data, ANOVA and t-tests were

used followed by Tukey or Holm-Bonferroni post-hoc corrections. Alpha was set to 0.05.

Relationships between neural data and each bird's song features (e.g., mean Entropy,

Duration, and Amplitude) were investigated using a generalized linear regression model

where the outcome variable was neural activity (e.g., firing rate) and the independent

variables were song features and group category (ASYN or GFP) as described in **Results**

and S2 Fig. Multicollinearity between independent variables can impair the ability to

interpret regression models. We observed that some song features did indeed covary (r

> 0.4 for 5 song feature pairs). To address this, we repeated the regression using the first

three principal components of the independent variable matrix and compared results to

those from the original regression model. As reported in **S2 Fig**, similar results were

obtained from the original and dimensionality-reduced model.

2.4 Tissue processing and immunohistochemistry

Immediately following the end of the neural recording session, finches were humanely

euthanized by isoflurane inhalant overdose and transcardially perfused with warmed

saline followed by chilled 4% paraformaldehyde in Dulbecco's Phosphate Buffer Saline.

Fixed brains were cryoprotected in 20% sucrose overnight then cryosectioned in the

coronal plane at 30 μm on a Microtome cryostat through Area X. Tissue was processed,

following our previously published methods (45, 46) and summarized here: Hydrophobic

borders were drawn on the slides, using a pap pen (ImmEdge, Vector Labs) followed by

3 X 5 minute washes in TBS with 0.3% Triton X (Tx). To block non-specific antibody

binding, the tissue was then incubated for one hour at room temperature with 5% goat

serum (Sigma #G-9023) in TBS/0.3% Tx then 3 x 5-minute washes in 1% goat serum in

TBS/0.3% Tx were performed. A primary antibody was used against a-syn (1:250, rabbit,

Proteintech 10842-1-AP, RRID: AB_2192672). GFP expression was robust enough to be

visualized without an antibody.

The primary antibody was incubated in a solution of 1% goat serum in TBS/0.3% Tx

overnight at 4°C. One tissue section per slide was used as a 'no primary' control and a

pre-absorption control experiment was performed to validate antibody specificity in a

companion study evaluating the distribution of a-syn protein following AAV injection (46).

The next day, sections were washed 5 x 5 minutes in TBS/0.3% Tx and incubated for

three hrs at room temperature in a fluorescently conjugated secondary antibody (1:1000:

goat anti-rabbit 647 A-21245, RRID:AB_2535813). After secondary incubation, sections

were washed 3 x 10 minutes in TBS followed by 2 x 5 washes in filtered TBS. Slides were

then cover-slipped in Pro-Long Anti-Fade Gold mounting medium (Molecular Probes,

P36930) and imaged using a Zeiss Axio Observer 7 with Apotome III Microscope under

the University of Arizona Imaging Cores - Optical Core Facility (RRID:SCR_023355).

Results:

Virally-driven expression is confirmed in song nucleus Area X

Fig 2 is a representative example of AAV expression in Area X. Two months post-virus

injection, we detected GFP cell bodies in Area X (**Fig 2A,C**) accompanied by low a-syn

protein expression (**Fig 2B**). By contrast, a finch overexpressing a-syn showed staining

in processes and cell bodies (**Fig 1D**). These results are consistent with our two prior

publications (45, 46).

Fig 2. AAV expression in finch song nucleus Area X. The Area X region is represented

by the dotted line. Small arrowheads denote the border between the striatum and the

nidopallium. Staining for alpha-synuclein (a-syn) protein is shown in red. **A)** An AAV

control expressing finch has GFP labeled cell bodies (green) in Area X and large

arrowhead denotes the last recorded tetrode position (yellow). **B)** A-syn protein

expression is low in Area X compared to surrounding basal ganglia in this AAV-GFP finch.

C) View of (A) showing the GFP channel only. **D)** An AAV-h*SNCA* expressing finch shows

widespread expression of a-syn in processes throughout Area X. Inset in bottom left

corner shows a zoomed in view of bead-like a-syn protein in the processes (inset scale

bar = 10 µm).

Extracellular identification of neuronal subtypes in anesthetized

finches

Area X neurons share much in common with neurons in the mammalian striatum

and pallidum. For example, the most abundant Area X neuron type is homologous to the

mammalian medium spiny neuron (MSN,(58)), with Area X and mammalian MSNs

sharing wide waveforms, low firing rates, and bursting activity (74-76). There is also

evidence that Area X contains striatal fast-firing interneurons, cholinergic interneurons,

and low-threshold spiking neurons (48, 74, 77). Furthermore, Area X contains two classes

of fast-firing pallidal-like neurons: local-circuit (HF-1) and thalamus-projecting (HF-2) (59,

78). These six cell classes were identified by their firing rates and waveform features

using multiple approaches from this literature, including correlating neuronal activity *in*

vivo with birdsong and patch-clamp recording in brain slices. Our present study is

constrained by the fact that all data was acquired from extracellular electrodes implanted

in anesthetized finches. We found no prior publications with recordings from Area X of

adult anesthetized finches; prior work includes some anesthetized recordings from lMAN

and Area X in the context of juvenile song learning (79). Consequently, our classification

was informed by criteria developed from *in vivo* recordings from Area X in singing finches

and restricted to waveform shape, firing rate, and assessment of bursting activity in the

following five identified categories (74, 78, 80).

Neuron categorization and criteria (results shown in **Fig 3**):

1) **Wide-Low-Rate (WLR) Criteria**: Neurons firing ≤ 6 Hz with a peak half width > 0.25

ms or a peak-to-trough width of > 0.35 ms (**Fig 3 A,D,E**). WLR neurons were also

more 'bursty'' relative to other neuronal types (**Fig 3F**) as measured by Local Variance

(LV, see Methods (81)) and were the most abundant of the five identified cell types

(**Fig 3C**). Together, these features suggest that WLR cells were MSN-like given

evidence that MSNs are the most prominent cell type in the striatum, exhibit wide

waveforms, have low firing rates, and express bursting activity (74) (75, 76).

2) **Wide-High-Rate (WHR) Criteria**: Neurons with a firing rate between 20 and 50 Hz

with a peak half width > 0.25 ms or a peak-to-trough width of > 0.35 ms (**Fig 3A,D,E**).

Post-classification analysis also showed that these neurons fired tonically (LV < 1, see

Fig 3F). Tonically active WHR cells have features consistent with striatal cholinergic

interneurons (74).

3) **Narrow-Low-Rate (NLR) Criteria**: Narrow waveform neurons firing ≤ 6 Hz (74) having

a peak half width < 0.25 ms or a peak-to-trough width of < 0.35 ms (**Fig 3A,D,E**).

4) **Narrow-High-Rate (NHR) Criteria**: Narrow waveform neurons with rates ≥ 8 Hz but

< 50Hz (**Fig 3A,D,E**). Narrow waveforms are characteristic of local circuit fast-firing

inhibitory neurons.

5) **Very-High-Rate (VHR) Criteria**: Neurons firing ≥ 50 Hz (**Fig 3A,D,E**). Area X pallidal-

like projection neurons are characterized by exceedingly high firing rates (74, 78).

Consequently, we established a category for neurons with rates ≥ 50 Hz. Waveform

features were not used to further subdivide neurons in this group given its small size

(n = 18 neurons, **Fig 3C**). Analysis of VHR firing activity revealed that these neurons

exhibited decidedly tonic activity, with a mean LV of 0.51 (**Fig 3F**). These data suggest

that the identified VHR neurons are type-2 pallidal-like projection neurons (HF-2 in

(78) (82)) given the presence of tonic firing and relatively wide waveforms.

Furthermore, the small size of this group is consistent with other reports that suggest

PAL neurons comprise approximately just 5.4% of the Area X cell population (58).

Fig 3. Identification of five classes of Area X neurons in anesthetized finches. A)

Mean shape for each of the five identified classes of neurons (n = 201 neurons from n =

12 finches). Neurons were categorized based on spike width and firing rate (see Results).

Amplitude is normalized such that the minimum (trough) = -1. Colors indicate neuron

subtype (see key). **B)** Inter-spike interval distributions for each neuronal subtype. Y-axis

indicates probability (p). **C)** Histogram indicating the number of identified neurons in each

category. **D)** Box plot of the half-width (ms-milliseconds) for each neuronal group. Each

dot indicates an individual neuron. Horizontal red lines indicate the median. Hypothesis

tests were not performed as half-width was a key criterion for classification. **E)** Firing rates

for each neuronal group (H-Hertz). Each dot indicates an individual neuron. Hypothesis

tests were not performed as firing rate was a key criterion for classification. **F)** Burst firing

differed between groups. While neurons were not categorized based on burst-firing, the

measure of bursting (Local Variance-LV) indicated that Wide-Low-Rate neurons had

increased burst firing (LV>1) relative to other groups (ANOVA $p < 0.000001$, Tukey post-

hoc comparison. Thick red lines indicate significant post-hoc differences (*$p < 0.05$, **$p <$

0.01 (not shown), ***$p < 0.001$). The red lines in the box plots indicate the median and the

whiskers indicate 1.5X the Interquartile Range.

Reduced Area X firing rates in ASYN overexpressing finches

We predicted that a-syn expression would reduce putative MSN (WLR) neuron
activity, thus releasing putative PAL neurons (VHR) from inhibition. Consequently, we
investigated whether firing rates of WLR neurons were reduced in the ASYN group and
whether rates in VHR neurons were increased. Visual inspection of raster plots of WLR
activity (**Fig 4A-B**) suggested reduced WLR firing activity in the ASYN group. Statistical
comparison of firing rates of GFP and ASYN neurons identified a significant difference
(**Fig 4C**, rank-sum, $p = 0.005$ before correction and $p = 0.026$ after Holm-Bonferroni
correction). No significant between-group differences were identified in the other neuronal
classes, including the VHR neurons.

Fig 4. Reduced firing rates of WLR neurons in the ASYN group. A-B) Raster plots of
WLR in the GFP and AAV-h*SNCA* (ASYN) groups. The x-axis shows time in seconds.
Each row is a raster for a distinct neuron. **C)** Between-group comparison (GFP vs. ASYN)
for each of the five neuron types. WLR neurons were only found in 4/6 finches. Firing
rates for WLR neurons were lower in the ASYN group relative to the GFP control (rank-
sum test, $p = 0.005$ prior to correction, *$p = 0.026$ after Holm-Bonferroni correction).

Burst firing was not changed in ASYN overexpressing finches

While burst firing is a canonical property of MSNs, MSN burst firing may become
dysregulated in PD. Indeed, altered MSN burst firing has been reported in the 6-OHDA
mouse model of PD (83), non-human MPTP primate models (84), and in human PD

patients (64). Consequently, we predicted that burst firing will be affected by ASYN

overexpression in finch Area X. To investigate this, the local variance (LV) measure (72),

see Methods) was used to evaluate the pattern of inter-spike intervals and quantify the

degree that neuronal activity was either tonic (low LV), Poisson (LV near 1), or bursty (LV

> 1). Contrary to our prediction, analysis of spiking statistics with LV did not identify any

difference between GFP and ASYN groups in LV values for any neuronal subtype (**Fig 5**,

t-test, all p > 0.05 after Holm-Bonferroni post-hoc correction). Thus, while ASYN

significantly reduced firing rates of WLR neurons, these changes were not accompanied

by a change in burst-firing.

434
435

Fig 5. Local Variance (LV), a measure of bursting, is not different between ASYN

and GFP control groups. Between-group comparisons (GFP vs. ASYN) were made for

each of the five neuron types, and no between-group differences were identified in any of

the neuron classes (rank-sum test, all p > 0.05 after Holm-Bonferroni post-hoc correction).

Extracellular waveform shape was altered in wide-waveform neurons

While waveform features and firing rates are commonly used to classify neurons

into putative subtypes (74, 80, 85), inferring physiological properties of neurons from

extracellular recordings is highly problematic as waveform features vary as a function of

recording location relative to the morphology of the neuron (86-88). Even so, we did

observe consistent differences in extracellular waveform shapes between ASYN and GFP

animals that encourage future investigations. Specifically, we found that WLR and WHR

neurons showed a large positive extracellular 'rebound' following the initial trough (**Fig**

6A). This was determined by first normalizing the waveform of each neuron such that the

trough = -1 and baseline was zero. Consequently, positive values indicate proportional

changes above baseline. The "Rebound Peak" for each neuron was measured as the

normalized amplitude of the largest peak following the time of the trough (t = 0 ms). Values

for each neuron type are shown in **Fig 6B** for the GFP and ASYN groups. Rank-sum tests

were performed, and significant between-group differences were identified in the WLR

and WHR groups (p_{WLR} = 0.003, p_{WHR} = 0.0002 after Bonferroni-Holm correction).

Fig 6. **ASYN group shows enhanced post-peak rebound for wide-waveform**

neurons. A) Mean waveforms for the ASYN and GFP groups for each neuron type. For

each waveform, data was normalized such that the distance from baseline to the trough

= -1. Inspection of these plots suggested that wide-waveform (WLR, WHR) ASYN

neurons had a larger rebound following the initial trough. **B)** Statistical analysis of the

rebound was performed by measuring the normalized amplitude of the largest peak

following t=0 (rebound peak) for each neuron (dots). These values were compared

between groups using rank-sum tests and p values were adjusted using the Holm-

Bonferroni correction. **p<0.01, ***p<0.001. Error bars in box plots indicate ±SEM.

No observed correlation between song features and neuronal firing

properties

Next, we investigated whether ASYN and GFP birds differed in their song features

and if variation in mean WLR neuronal activity in each bird correlated with song features.

Analysis of syllable-level features failed to identify any between-group difference (Mann-

Whitney U test, $p > 0.05$, **S1 Fig, S1 Table**).

Investigation of whether firing rates correlated with song features was performed

by first calculating the mean firing rate of WLR neurons (the largest group and the putative

MSNs) for each bird. Two birds had to be excluded from the analysis as no WLR neurons

were identified in these animals. Mean WLR firing rates were used as the outcome

variable for a generalized linear model where the independent variables were song

features (mean amplitude, duration, and entropy and CV amplitude, duration, and

entropy) and the group (ASYN or GFP). No significant relationship between song features

and firing rate was identified ($F(10,2) = 11.7$, $p = 0.082$);**S2 Fig**.

4. Discussion

The goal of this study was to obtain new information about the cellular and network-

level activity affected by overexpression of human a-syn protein in an adult male zebra

finch model. To carry out this goal, we performed extracellular recordings from song

nucleus Area X of the basal ganglia in anesthetized adult male finches and identified five

distinct cell types. Our main finding was that wide, low firing rate (WLR) neurons, putative

MSNs, show reduced firing activity and enhanced post-inhibitory rebound in the a-syn

overexpression condition compared to controls. This aligns with our working hypothesis

that abnormally elevated a-syn protein levels in Area X will decrease MSNs firing rate.

We also predicted that GP-projecting neurons would show reduced firing in Area X. VHR

neurons may represent these GP-projecting neurons, but they did not show a difference

in firing rate between the ASYN and GFP control groups. This finding may reflect low

statistical power given the small number of VHR neurons recorded.

A benefit of anesthetized recordings is that they allow for high data throughput,

enabling the recording of a substantial proportion of neurons without confounds due to

animal movement. A limitation of this approach in our finch model is that on-going real

time modification of neural activity related to song production cannot be evaluated. It is

also likely that neuronal activity was dampened by isoflurane anesthesia. While

information is scarce on the effects of isoflurane on *in vivo* recordings from the striatum

of naïve finches or rodents, patch clamp studies in rat brain slices show that brief

isoflurane exposure can dampen sodium currents in hippocampal neurons (89) and

reduce spiking and bursting modes in thalamic neurons (90). Our future investigations will

compare isoflurane-induced anesthesia on Area X of naïve finches versus awake, singing

finches where we can examine how a-syn modulates neuronal dynamics under more

physiologically relevant conditions.

The observed reduction in WLR (putative MSN) firing rates observed in Area X of

the ASYN finches contrasts with some measures of MSN activity reported in human PD

patients and primate PD models. For example, striatal projection neurons in the human

basal ganglia of PD patients undergoing deep brain stimulation had high frequency

neuronal firing patterns resembling the advanced parkinsonism of the Non-Human

Primate model, with hyperactivity and "bursty" neuronal firing activity (14, 64). In a mouse

model of bilateral dopamine depletion (via the neurotoxin 6-OHDA) in the dorsolateral

striatum, single unit recordings from MSNs showed reduced bursting activity during

movement. Intriguingly, they classified MSNs into two types with type 1 showing

increased firing during movement and type 2, decreased firing with no difference in firing

rate when the mouse was stationary (83). In freely moving rats, 6-OHDA treatment

increased oscillatory synchronization in the striatum (91). Future recordings in singing,

ambulatory finches with the ASYN phenotype will determine if hyperactivity of MSNs and

differential sub-types exist, evidence for dopamine loss, and how these various sub-

components contribute to changes in real-time song output.

The reduced firing rates of WLR neurons in ASYN expressing finches may reflect

reduced mean frequency of spontaneous excitatory postsynaptic glutamatergic currents

(EPSCs) from cortical song nucleus lMAN onto Area X MSNs and GP neurons. Striatal

recordings from h*SNCA* overexpressing mice show reduced glu currents (12). Future

work using electrophysiological recordings from Area X brain slices *in vitro* can determine

whether glu input is perturbed in our ASYN condition.

ASYN-driven abnormal changes in synaptic input to Area X may also be

accompanied by altered intrinsic membrane properties. Both the WLR and WHR neuronal

classes showed greater positive rebound potential following the initial spike in the ASYN

expressing group compared to GFP controls. While extracellular recordings of action

potentials cannot reliably identify after-hyperpolarization (AHP) currents, the observed

rebound potential occurs during the typical time interval of the AHP, suggesting that

membrane properties in the ASYN neurons are altered. The AHP is regulated by K^+

currents, and several classes of K^+ channels are implicated in altered neuronal excitability

in PD models (92). The significantly larger rebound potential in the ASYN expressing

group may reflect an enhanced AHP-related conductance. Patch-clamp

electrophysiology in brain slices is a necessary approach to confirm true fluctuations in

membrane properties during the AHP period. Changes in AHP-related conductance could

result from increased SK/BK channel activity (93) (94)) or a reduction in I_h current (95,

96), leading to greater hyperpolarization following spikes and dampened excitability. In

nigral DA neurons, abnormal a-syn oligo aggregates increase the conductance of K^+_{ATP}

channels leading to reduced firing (97). In our finch model, a-syn overexpression could

progressively decrease intrinsic excitability in MSNs via these channels, and/or others.

To evaluate the specific role of a-syn expression on MSNs, future directions will employ

a custom-made AAV containing a CAMKII promoter that has been previously used in

naïve finches to evaluate MSNs firing activity (98). By selectively overexpressing a-syn in

MSNs, we can evaluate the relationship between the extent of protein pathology and the

neural activity changes underlying song. In a recent publication, we found that regional

changes in virally-expressed human a-syn protein within Area X correlated with changes

in syllable level features. For example, we detected a positive correlation between right

hemisphere a-syn proteinopathy and a reduction in the variation of harmonic syllable

duration, among other features correlated to the right hemisphere Area X (46). Our neural

recordings here are also from right hemisphere Area X; the contribution of the left Area X

is not known. Prior studies suggest a transition between right-side dominance of song

nucleus HVC in adult birdsong (99) while the left-side dominates during juvenile

learning(100). Thus, when assessing cell-type changes in electrical activity within brain

nuclei aligned with real-time singing behavior, factors to take into account include the

recording hemisphere, the distribution of protein expression, and the presence of

pathology, e.g. a-syn aggregates in individual cell types.

Conclusions

Our study takes an important first step forward in evaluating the consequences of

a-syn overexpression on the activities of neuronal subtypes involved in vocal behavior. A

growing body of research has elucidated the role that neurons in the cortico-striatal-

thalamic circuit play in male zebra finch song production. By introducing human a-syn to

the zebra finch model, this study supports a systems- and cellular-level investigation of

human vocal deficits in PD. This is important given that current medications and brain

stimulation used in the treatment of PD fail to resolve the vocal deficits and can worsen

them (101-103). Data from pre-clinical animal models such as the songbird can provide

important mechanistic insight needed for the development of new brain-targeted cellular

therapies for the vocal dysfunction.

CRediT Contributions
580
CRediT authorship contribution statement: Brian R. Dominguez: Data curation,

Investigation, Methodology, Validation, Visualization, Writing-Original Draft, Writing-

review & editing. Gabriel Holguin: Data curation, Formal Analysis, Investigation,

Methodology, Software, Validation, Visualization, Writing-Original Draft, Writing-review &

editing. Madeleine S. Daly: Data curation, Formal Analysis, Investigation, Methodology,

Validation, Writing-review & editing. Reed T. Bjork: Investigation, Methodology,

Validation, Visualization, Writing-review & editing. Stephen Cowen: Conceptualization,

Data Curation, Formal Analysis, Investigation, Methodology, Software, Supervision,

Validation, Visualization, Writing-Original Draft, Writing-review & editing. Julie E. Miller:

Conceptualization, Data curation, Formal Analysis, Funding acquisition, Investigation,

Methodology, Project administration, Resources, Supervision, Validation, Visualization,

Writing-original draft, Writing-review & editing.

Acknowledgements

597   We thank Kathy Dillon, Michelle Gordon, and Mauricio Serna for technical contributions

598   along with University of Arizona Animal Care staff and Mr. Douglas Cromey of the

599   University of Arizona Imaging Cores - Optical Core Facility (RRID:SCR_023355).

References

1.   Polymeropoulos MH, Lavedan C, Leroy E, Ide SE, Dehejia A, Dutra A, et al.

Mutation in the alpha-synuclein gene identified in families with Parkinson's disease.

Science. 1997;276(5321):2045-7.

2.   Chartier-Harlin M-C, Kachergus J, Roumier C, Mouroux V, Douay X, Lincoln S, et

al. α-synuclein locus duplication as a cause of familial Parkinson's disease. The Lancet.

2004;364(9440):1167-9.

3.   Singleton A, Farrer M, Johnson J, Singleton A, Hague S, Kachergus J, et al. α-

Synuclein locus triplication causes Parkinson's disease. Science. 2003;302(5646):841-.

4.   Mak E, Su L, Williams GB, O'Brien JT. Neuroimaging characteristics of dementia

with Lewy bodies. Alzheimer's research & therapy. 2014;6:1-10.

5.   Koga S, Dickson DW. Recent advances in neuropathology, biomarkers and

therapeutic approach of multiple system atrophy. Journal of Neurology, Neurosurgery &

Psychiatry. 2018;89(2):175-84.

6.   Calo L, Wegrzynowicz M, Santivanez-Perez J, Grazia Spillantini M. Synaptic

failure and alpha-synuclein. Movement disorders: official journal of the Movement

Disorder Society. 2016;31(2):169-77.

7.   Marino G, Calabresi P, Ghiglieri V. Alpha-synuclein and cortico-striatal plasticity in

animal models of Parkinson disease. Handbook of clinical neurology. 2022;184:153-66.

8.   Calabresi P, Mechelli A, Natale G, Volpicelli-Daley L, Di Lazzaro G, Ghiglieri V.

Alpha-synuclein in Parkinson's disease and other synucleinopathies: from overt

neurodegeneration back to early synaptic dysfunction. Cell death & disease.

2023;14(3):176.

9.   Watson JB, Hatami A, David H, Masliah E, Roberts K, Evans CE, et al. Alterations

in corticostriatal synaptic plasticity in mice overexpressing human alpha-synuclein.

Neuroscience. 2009;159(2):501-13.

10.   Tozzi A, de Iure A, Bagetta V, Tantucci M, Durante V, Quiroga-Varela A, et al.

Alpha-Synuclein Produces Early Behavioral Alterations via Striatal Cholinergic Synaptic

Dysfunction by Interacting With GluN2D N-Methyl-D-Aspartate Receptor Subunit. Biol

Psychiat. 2016;79(5):402-14.

11.   Tozzi A, Sciaccaluga M, Loffredo V, Megaro A, Ledonne A, Cardinale A, et al.

Dopamine-dependent early synaptic and motor dysfunctions induced by α-synuclein in

the nigrostriatal circuit. Brain. 2021;144(11):3477-91.

12.   Wu N, Joshi PR, Cepeda C, Masliah E, Levine MS. Alpha-synuclein

overexpression in mice alters synaptic communication in the corticostriatal pathway.

Journal of neuroscience research. 2010;88(8):1764-76.

13.   Lam HA, Wu N, Cely I, Kelly RL, Hean S, Richter F, et al. Elevated tonic

extracellular dopamine concentration and altered dopamine modulation of synaptic

activity precede dopamine loss in the striatum of mice overexpressing human α-synuclein.

Journal of neuroscience research. 2011;89(7):1091-102.

14.   Beck G, Singh A, Papa SM. Dysregulation of striatal projection neurons in

Parkinson's disease. Journal of Neural Transmission. 2018;125(3):449-60.

15.   Durante V, de Iure A, Loffredo V, Vaikath N, De Risi M, Paciotti S, et al. Alpha-

synuclein targets GluN2A NMDA receptor subunit causing striatal synaptic dysfunction

and visuospatial memory alteration. Brain. 2019;142(5):1365-85.

16.   Wilson CJ, Groves PM. Spontaneous firing patterns of identified spiny neurons in

the rat neostriatum. Brain research. 1981;220(1):67-80.

17.   Kita H, Kita T. Role of striatum in the pause and burst generation in the globus

pallidus of 6-OHDA-treated rats. Frontiers in systems neuroscience. 2011;5:42.

18. Henkel LM, Kankowski S, Moellenkamp TM, Smandzich NJ, Schwarz S, Di Fonzo

A, et al. iPSC-derived striatal medium spiny neurons from patients with multiple system

atrophy show hypoexcitability and elevated α-synuclein release. Cells. 2023;12(2):223.

19. Lobb CJ, Zaheer AK, Smith Y, Jaeger D. In vivo electrophysiology of nigral and

thalamic neurons in alpha-synuclein-overexpressing mice highlights differences from

toxin-based models of parkinsonism. Journal of neurophysiology. 2013;110(12):2792-

805.

20. Brys I, Nunes J, Fuentes R. Motor deficits and beta oscillations are dissociable in

an alpha-synuclein model of Parkinson's disease. European Journal of Neuroscience.

2017;46(3):1906-17.

21. Robson E, Tweedy C, Manzanza N, Taylor J-P, Atkinson P, Randall F, et al.

Impaired fast network oscillations and mitochondrial dysfunction in a mouse model of

alpha-synucleinopathy (A30P). Neuroscience. 2018;377:161-73.

22. Sapir S. Multiple factors are involved in the dysarthria associated with Parkinson's

disease: a review with implications for clinical practice and research. Journal of speech,

language, and hearing research : JSLHR. 2014;57(4):1330-43.

23. Ma A, Lau KK, Thyagarajan D. Voice changes in Parkinson's disease: What are

they telling us? Journal of Clinical Neuroscience. 2020;72:1-7.

24. Stewart C, Winfield L, Hunt A, Bressman SB, Fahn S, Blitzer A, et al. Speech

dysfunction in early Parkinson's disease. Movement disorders: official journal of the

Movement Disorder Society. 1995;10(5):562-5.

25. Ho AK, Iansek R, Marigliani C, Bradshaw JL, Gates S. Speech impairment in a

large sample of patients with Parkinson's disease. Behav Neurol. 1998;11(3):131-7.

26.   Harel B, Cannizzaro M, Snyder PJ. Variability in fundamental frequency during

speech in prodromal and incipient Parkinson's disease: a longitudinal case study. Brain

Cogn. 2004;56(1):24-9.

27.   Skodda S, Schlegel U. Speech rate and rhythm in Parkinson's disease. Movement

disorders: official journal of the Movement Disorder Society. 2008;23(7):985-92.

28.   Skodda S, Rinsche H, Schlegel U. Progression of dysprosody in Parkinson's

disease over time--a longitudinal study. Movement disorders: official journal of the

Movement Disorder Society. 2009;24(5):716-22.

29.   Rusz J, Cmejla R, Ruzickova H, Ruzicka E. Quantitative acoustic measurements

for characterization of speech and voice disorders in early untreated Parkinson's disease.

The Journal of the Acoustical Society of America. 2011;129(1):350-67.

30.   Defazio G, Guerrieri M, Liuzzi D, Gigante AF, di Nicola V. Assessment of voice

and speech symptoms in early Parkinson's disease by the Robertson dysarthria profile.

Neurol Sci. 2016;37(3):443-9.

31.   V. Berisha JL, T. Huston, A. Wisler, Y. Jiao, and J. Eig. Float Like a Butterfly Sting

Like a Bee: Changes in Speech Preceded Parkinsonism Diagnosis for Muhammad Ali.

Proceedings of the 2017 Interspeech Conference.

32.   Rusz J, Tykalová T, Novotný M, Růžička E, Dušek P. Distinct patterns of speech

disorder in early-onset and late-onset de-novo Parkinson's disease. npj Parkinson's

Disease. 2021;7(1):98.

33.   Šimek M, Rusz J. Validation of cepstral peak prominence in assessing early voice

changes of Parkinson's disease: Effect of speaking task and ambient noise. The Journal

of the Acoustical Society of America. 2021;150(6):4522-33.

34.   Berus L, Klancnik S, Brezocnik M, Ficko M. Classifying Parkinson's disease based

on acoustic measures using artificial neural networks. Sensors. 2018;19(1):16.

35.   Solana-Lavalle G, Galán-Hernández J-C, Rosas-Romero R. Automatic Parkinson

disease detection at early stages as a pre-diagnosis tool by using classifiers and a small

set of vocal features. Biocybernetics and Biomedical Engineering. 2020;40(1):505-16.

36.   Suppa A, Costantini G, Asci F, Di Leo P, Al-Wardat MS, Di Lazzaro G, et al. Voice

in Parkinson's disease: a machine learning study. Frontiers in Neurology.

2022;13:831428.

37.   Lim WS, Chiu S-I, Wu M-C, Tsai S-F, Wang P-H, Lin K-P, et al. An integrated

biometric voice and facial features for early detection of Parkinson's disease. npj

Parkinson's Disease. 2022;8(1):145.

38.   Ciucci MR, Ma ST, Fox C, Kane JR, Ramig LO, Schallert T. Qualitative changes

in ultrasonic vocalization in rats after unilateral dopamine depletion or haloperidol: a

preliminary study. Behav Brain Res. 2007;182(2):284-9.

39.   Gombash SE, Manfredsson FP, Kemp CJ, Kuhn NC, Fleming SM, Egan AE, et al.

Morphological and behavioral impact of AAV2/5-mediated overexpression of human

wildtype alpha-synuclein in the rat nigrostriatal system. PloS one. 2013;8(11):e81426.

40.   Miller JE, Hafzalla GW, Burkett ZD, Fox CM, White SA. Reduced vocal variability

in a zebra finch model of dopamine depletion: implications for Parkinson disease. Physiol

Rep. 2015;3(11).

41.   Grant LM, Kelm-Nelson CA, Hilby BL, Blue KV, Paul Rajamanickam ES, Pultorak

JD, et al. Evidence for early and progressive ultrasonic vocalization and oromotor deficits

in a PINK1 gene knockout rat model of Parkinson's disease. J Neurosci Res.
2015;3:1713–27.

42.   Yang KM, Blue KV, Mulholland HM, Kurup MP, Kelm-Nelson CA, Ciucci MR.
Characterization of oromotor and limb motor dysfunction in the DJ1-/-model of Parkinson
disease. Behavioural brain research. 2018;339:47-56.

43.   Kelm-Nelson CA, Gammie S. Gene expression within the periaqueductal gray is
linked to vocal behavior and early-onset parkinsonism in Pink1 knockout rats. BMC
genomics. 2020;21:1-13.

44.   Grant LM, Richter F, Miller JE, White SA, Fox CM, Zhu C, et al. Vocalization deficits
in mice over-expressing alpha-synuclein, a model of pre-manifest Parkinson's disease.
Behavioral neuroscience. 2014;128(2):110-21.

45.   Medina CA, Vargas E, Munger SJ, Miller JE. Vocal changes in a zebra finch model
of Parkinson's disease characterized by alpha-synuclein overexpression in the song-
dedicated anterior forebrain pathway. PloS one. 2022;17(5):e0265604.

46.   Bjork RT, Patel FZ, Daly MS, Miller JE. Evaluating alpha-synuclein proteinopathy
and consequences for birdsong in zebra finch basal ganglia Area X. Behavioural Brain
Research. 2025:115698.

47.   Brainard MS, Doupe AJ. Translating birdsong: songbirds as a model for basic and
applied medical research. Annu Rev Neurosci. 2013;36:489-517.

48.   Farries MA, Perkel DJ. Electrophysiological properties of avian basal ganglia
neurons recorded in vitro. Journal of neurophysiology. 2000;84(5):2502-13.

49.   Person AL, Gale SD, Farries MA, Perkel DJ. Organization of the songbird basal
ganglia, including area X. J Comp Neurol. 2008;508(5):840-66.

50.   Gale SD, Perkel DJ. Anatomy of a songbird basal ganglia circuit essential for vocal

learning and plasticity. Journal of chemical neuroanatomy. 2010;39(2):124-31.

51.   Reiner A, Perkel DJ, Bruce L, Butler AB, Csillag A, Kuenzel W, et al. Revised

nomenclature for avian telencephalon and some related brainstem nuclei. *Journal of*

Comparative Neurology. 2004;473:377-414.

52.   White SA. Genes and vocal learning. Brain Lang. 2010;115(1):21-8.

53.   Simonyan K, Horwitz B, Jarvis ED. Dopamine regulation of human speech and bird

song: A critical review. Brain Lang. 2012;122(3):142-50.

54.   Pfenning AR, Hara E, Whitney O, Rivas MV, Wang R, Roulhac PL, et al.

Convergent transcriptional specializations in the brains of humans and song-learning

birds. Science. 2014;346(6215):1256846.

55.   Woolley SC, Kao MH. Variability in action: Contributions of a songbird cortical-

basal ganglia circuit to vocal motor learning and control. Neuroscience. 2015;296:39-47.

56.   Lovell PV, Wirthlin M, Kaser T, Buckner AA, Carleton JB, Snider BR, et al. ZEBrA:

Zebra finch Expression Brain Atlas—A resource for comparative molecular

neuroanatomy and brain evolution studies. Journal of Comparative Neurology.

2020;528(12):2099-131.

57.   Sakata JT, Birdsong D. Vocal Learning and Behaviors in Birds and Human

Bilinguals: Parallels, Divergences and Directions for Research. Languages. 2021;7(1):5.

58.   Reiner A, Laverghetta AV, Meade CA, Cuthbertson SL, Bottjer SW. An

immunohistochemical and pathway tracing study of the striatopallidal organization of area

X in the male zebra finch. J Comp Neurol. 2004;469(2):239-61.

59. Farries MA, Perkel DJ. A telencephalic nucleus essential for song learning

contains neurons with physiological characteristics of both striatum and globus pallidus.

The Journal of neuroscience: the official journal of the Society for Neuroscience.

2002;22(9):3776-87.

60. Coleman MJ, Roy A, Wild JM, Mooney R. Thalamic gating of auditory responses

in telencephalic song control nuclei. Journal of Neuroscience. 2007;27(37):10024-36.

61. Peh WY, Roberts TF, Mooney R. Imaging auditory representations of song and

syllables in populations of sensorimotor neurons essential to vocal communication.

Journal of Neuroscience. 2015;35(14):5589-605.

62. Woolley SC, Rajan R, Joshua M, Doupe AJ. Emergence of context-dependent

variability across a basal ganglia network. Neuron. 2014;82(1):208-23.

63. Farries MA, Ding L, Perkel DJ. Evidence for "direct" and "indirect" pathways

through the song system basal ganglia. J Comp Neurol. 2005;484(1):93-104.

64. Singh A, Mewes K, Gross RE, DeLong MR, Obeso JA, Papa SM. Human striatal

recordings reveal abnormal discharge of projection neurons in Parkinson's disease.

Proceedings of the National Academy of Sciences. 2016;113(34):9629-34.

65. Kojima S, Kao MH, Doupe AJ. Task-related "cortical" bursting depends critically

on basal ganglia input and is linked to vocal plasticity. Proceedings of the National

Academy of Sciences of the United States of America. 2013;110(12):4756-61.

66. Moll FW, Kranz D, Corredera Asensio A, Elmaleh M, Ackert-Smith LA, Long MA.

Thalamus drives vocal onsets in the zebra finch courtship song. Nature. 2023:1-5.

67. Tchernichovski O, Nottebohm F, Ho CE, Pesaran B, Mitra PP. A procedure for an

automated measurement of song similarity. Animal Behaviour. 2000;59:1167-76.

68.   Badwal A, Poertner J, Samlan RA, Miller JE. Common terminology and acoustic

measures for human voice and birdsong. Journal of Speech, Language, and Hearing

Research. 2018;61(12).

69.   Badwal A, Borgstrom M, Samlan RA, Miller JE. Middle age, a key time point for

changes in birdsong and human voice. Behavioral neuroscience. 2020;134(3):208.

70.   Sober SJ, Wohlgemuth MJ, Brainard MS. Central contributions to acoustic

variation in birdsong. The Journal of neuroscience: the official journal of the Society for

Neuroscience. 2008;28(41):10370-9.

71.   Pachitariu M, Steinmetz NA, Kadir SN, Carandini M, Harris KD. Fast and accurate

spike sorting of high-channel count probes with KiloSort. Advances in neural information

processing systems. 2016;29.

72.   Shinomoto S, Kim H, Shimokawa T, Matsuno N, Funahashi S, Shima K, et al.

Relating neuronal firing patterns to functional differentiation of cerebral cortex. PLoS

computational biology. 2009;5(7):e1000433.

73.   Buzsáki G, Mizuseki K. The log-dynamic brain: how skewed distributions affect

network operations. Nature reviews neuroscience. 2014;15(4):264-78.

74.   Goldberg JH, Fee MS. Singing-related neural activity distinguishes four classes of

putative striatal neurons in the songbird basal ganglia. Journal of neurophysiology.

2010;103:2002-14.

75.   Berke JD, Okatan M, Skurski J, Eichenbaum HB. Oscillatory entrainment of striatal

neurons in freely moving rats. Neuron. 2004;43(6):883-96.

76.   Wilson CJ. The generation of natural firing patterns in neostriatal neurons.

Progress in brain research. 1993;99:277-97.

77. Budzillo A, Duffy A, Miller KE, Fairhall AL, Perkel DJ. Dopaminergic modulation of

basal ganglia output through coupled excitation-inhibition. Proceedings of the National

Academy of Sciences of the United States of America. 2017;114(22):5713-8.

78. Goldberg JH, Adler A, Bergman H, Fee MS. Singing-related neural activity

distinguishes two putative pallidal cell types in the songbird basal ganglia: comparison to

the primate internal and external pallidal segments. The Journal of neuroscience: the

official journal of the Society for Neuroscience. 2010;30(20):7088-98.

79. Doupe AJ, Solis MM, Kimpo R, Boettiger CA. Cellular, circuit, and synaptic

mechanisms in song learning. Annals of the New York Academy of Sciences.

2004;1016(1):495-523.

80. Barthó P, Hirase H, Monconduit L, Zugaro M, Harris KD, Buzsaki G.

Characterization of neocortical principal cells and interneurons by network interactions

and extracellular features. Journal of neurophysiology. 2004;92(1):600-8.

81. Shinomoto S, Shima K, Tanji J. Differences in spiking patterns among cortical

neurons. Neural computation. 2003;15(12):2823-42.

82. Bugaysen J, Bronfeld M, Tischler H, Bar-Gad I, Korngreen A. Electrophysiological

characteristics of globus pallidus neurons. PloS one. 2010;5(8):e12001.

83. Chen H, Lei H, Xu Q. Neuronal activity pattern defects in the striatum in awake

mouse model of Parkinson's disease. Behavioural brain research. 2018;341:135-45.

84. Singh A, Liang L, Kaneoke Y, Cao X, Papa SM. Dopamine regulates distinctively

the activity patterns of striatal output neurons in advanced parkinsonian primates. Journal

of neurophysiology. 2015;113(5):1533-44.

85.   Vishwanath A, Bartlett MJ, Falk T, Cowen SL. Decoupling of motor cortex to

movement in Parkinson's dyskinesia rescued by sub-anaesthetic ketamine. Brain.

2025;148(6):2135-50.

86.   Gold C, Henze DA, Koch C, Buzsaki G. On the origin of the extracellular action

potential waveform: a modeling study. Journal of neurophysiology. 2006;95(5):3113-28.

87.   Teleńczuk M, Brette R, Destexhe A, Teleńczuk B. Contribution of the axon initial

segment to action potentials recorded extracellularly. Eneuro. 2018;5(3).

88.   Bakkum DJ, Obien MEJ, Radivojevic M, Jäckel D, Frey U, Takahashi H, et al. The

axon initial segment is the dominant contributor to the neuron's extracellular electrical

potential landscape. Advanced biosystems. 2019;3(2):1800308.

89.   Zhao W, Zhang M, Liu J, Liang P, Wang R, Hemmings HC, et al. Isoflurane

modulates hippocampal CA1 pyramidal neuron excitability by inhibition of both transient

and persistent sodium currents in mice. Anesthesiology. 2019;131(1):94.

90.   Ries CR, Puil E. Isoflurane prevents transitions to tonic and burst firing modes in

thalamic neurons. Neuroscience letters. 1993;159(1-2):91-4.

91.   Dejean C, Gross CE, Bioulac B, Boraud T. Dynamic changes in the cortex-basal

ganglia network after dopamine depletion in the rat. Journal of neurophysiology.

2008;100(1):385-96.

92.   Choudhury SP, Bano S, Sen S, Suchal K, Kumar S, Nikolajeff F, et al. Altered

neural cell junctions and ion-channels leading to disrupted neuron communication in

Parkinson's disease. npj Parkinson's Disease. 2022;8(1):66.

93.   Deister CA, Chan CS, Surmeier DJ, Wilson CJ. Calcium-activated SK channels

influence voltage-gated ion channels to determine the precision of firing in globus pallidus

neurons. Journal of Neuroscience. 2009;29(26):8452-61.

94.   Faber EL, Sah P. Physiological role of calcium-activated potassium currents in the

rat lateral amygdala. Journal of Neuroscience. 2002;22(5):1618-28.

95.   Robinson RB, Siegelbaum SA. Hyperpolarization-activated cation currents: from

molecules to physiological function. Annual review of physiology. 2003;65(1):453-80.

96.   Storm JF. Potassium currents in hippocampal pyramidal cells. Progress in brain

research. 1990;83:161-87.

97.   Hill E, Gowers R, Richardson M, Wall M. α-Synuclein aggregates increase the

conductance of substantia nigra dopamine neurons, an effect partly reversed by the

KATP channel inhibitor glibenclamide. eneuro. 2021;8(1).

98.   Singh Alvarado J, Goffinet J, Michael V, Liberti W, Hatfield J, Gardner T, et al.

Neural dynamics underlying birdsong practice and performance. Nature. 2021:1-5.

99.   Williams H, Crane LA, Hale TK, Esposito MA, Nottebohm F. Right-side dominance

for song control in the zebra finch. Journal of Neurobiology. 1992;23(8):1006-20.

100.   Frank SY, Hunt JL, Bae AJ, Chirathivat N, Lotfi S, Raja SC, et al. Hemispheric

dominance in HVC is experience-dependent in juvenile male zebra finches. Scientific

reports. 2024;14(1):5781.

101.   Plowman-Prine E, Okun M, Sapienza C, Shrivastav R, Fernandez H, Foote K, et

al. Perceptual characteristics of Parkinsonian speech: a comparison of the

pharmacological effects of levodopa across speech and non-speech motor systems.

NeuroRehabilitation. 2009;24(2):131-44.

102. Skodda S. Effect of deep brain stimulation on speech performance in Parkinson's

disease. Parkinson's Disease. 2012;2012.

103. Fabbri M, Guimaraes I, Cardoso R, Coelho M, Guedes LC, Rosa MM, et al. Speech

and voice response to a levodopa challenge in late-stage Parkinson's Disease. Front

Neurol. 2017;8:432.

Supporting Information

S1 Table 1. Song data and statistics. For further information, see legends embedded

into the tabs of this excel file.

S1 Fig. Exemplar song motif and group level comparisons of song syllable data.

A) Spectrogram of song motif from a pre-injected GFP control bird (as in **Fig 2A-C**) with

time in seconds on the x-axis and frequency in kilo (k) Hertz on the y-axis. Unique

syllables are assigned a letter (A-E) and classified into harmonic, noisy, or mixed syllables

(68). **B-E)** The mean and SEM scores shown for syllable level acoustic features (duration,

amplitude, and entropy) for the ASYN (red) and GFP (black) control groups. Raw scores

can be found in the **S1 Table**. The y-axis is the normalized score calculated from dividing

the post-AAV injection scores by the pre-injection scores. Each dot represents an average

score per finch. **B)** Comparisons of all syllable types combined or by individual syllable

types **(C-E)** reveal no group differences (Mann Whitney U, p>0.05). Individual variation in

CV scores is detected.

S2 Fig. No significant correlation between song features and firing rate. Scatter plots

showing the mean firing rate of WLR neurons averaged for each animal (dot) organized

by GFP (black) and ASYN (red) groups. Plots were generated for each song feature,

representing all syllable types combined. Generalized linear regression (see Results) did

not identify a relationship between firing rate and song features or group ($p = 0.0815$).

Figure 1B

Figure 1C

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A

Figure 1A