Title: Alpha-synuclein overexpression reduces neural activity within a basal ganglia

vocal nucleus in a zebra finch model

Running title: Alpha-synuclein overexpression reduces neural activity in a finch song

nucleus

Brian R. Dominguez[1], Gabriel Holguin[2], Madeleine S. Daly[1], Reed T. Bjork[1], Stephen L.
Cowen[2,3], and Julie E. Miller*[1,4]
9

[1]Department of Neuroscience, University of Arizona, Tucson, Arizona, United States of America.
[2]Department of Psychology, University of Arizona, Tucson, Arizona, United States of America.
[3]Evelyn F. McKnight Brain Institute, University of Arizona, Tucson, Arizona, United States of America
[4]Departments of Speech, Language and Hearing Sciences, Neurology, and the BIO5 Institute, Tucson, Arizona, United States of America.

*Corresponding Author:

Julie E. Miller, Ph.D.; E-mail: juliemiller@arizona.edu

Declaration of Competing Interest: J.E. Miller is a 2025 member of the "T2 Targets to

Therapies" advisory working group for the Michael J. Fox Foundation for Parkinson's

Research.

Other author emails:
Dominguezbrian2@gmail.com
gholguin@scripps.edu
madeleinedaly@arizona.edu
bjork2@arizona.edu
scowen@arizona.edu

Abstract

Changes in vocal pitch, loudness, and timing are prevalent in Parkinson's Disease (PD) and a target for early intervention and treatment. The neural mechanisms underlying these impairments are not understood, motivating work in animal models. The adult male zebra finch songbird is uniquely poised for these studies given vocally-dedicated brain nuclei and a quantifiable output (birdsong). Our prior publication revealed that injection of an adeno-associated virus (AAV5) expressing the human (h) alpha-synuclein (h*SNCA*, a-syn) gene into basal ganglia vocal nucleus Area X results in elevated insoluble a-syn protein and parkinsonian-like changes including softer, shorter, and reduced vocalizations compared to controls. Here, we test the hypothesis that AAV-h*SNCA* overexpression reduces the firing rate of specific neuronal sub-types in Area X using *in vivo* recordings in anesthetized finches. Five classes of neurons were differentiated in AAV-treated finches based on waveform width (narrow vs. wide) and firing rates (low vs. fast). Wide-waveform/low-rate activity is a consistent feature of striatal medium spiny neurons (MSNs), a dominant cell type in Area X and in mammalian basal ganglia. Reduced firing rates and enhanced post-peak rebound were detected in the AAV-h*SNCA* group for putative MSN neurons compared to AAV controls. No differences in firing rate nor waveform shape were detected for the narrow waveform neurons. Our findings provide the first characterization of early a-syn-driven neural activity changes in vocal control neurocircuitry.

Key words: Parkinson's, alpha-synuclein, zebra finch, songbird, vocal, basal ganglia

Introduction:

Pathological accumulation of alpha-synuclein (a-syn) is believed to contribute to multiple neurodegenerative diseases, including Parkinson's disease (PD). Accumulation of toxic a-syn species culminates in the development of Lewy Bodies and Lewy Neurites that are detected in inherited and sporadic cases of PD, dementia with Lewy Body disease, and Multiple Systems Atrophy (1-5). How abnormal increases in a-syn protein expression and its aggregation disrupts ongoing neuronal activity in the basal ganglia (BG) is not understood, yet this information is critical to developing cell-specific therapeutic targets (6-8). One experimental approach is to examine how elevated a-syn protein interferes with the ongoing firing activity of striatal neurons.

Elevated levels of a-syn protein at cortico-striatal synapses can result in synaptic failure and loss of cortico-striatal plasticity based on electrophysiological recordings from rodent brain slices (6-11). Striatal Medium Spiny Neurons (MSNs) show alterations in dendritic spine density, interactions between a-syn aggregates and their NMDA receptors, and reduced efficacy of cortical pre-synaptic glutamatergic input that affects their firing rate (12-15). Furthermore, burst firing is a characteristic feature of MSNs (16), and dysregulated MSN burst firing may contribute to PD as shown in 6-hydroxydopamine (6-OHDA) models (17). A recent study used induced human pluripotent stem cells from patients with Multiple System Atrophy of the Parkinsonian type that were differentiated into MSNs. These MSNs showed reduced spontaneous firing rates, frequency, and amplitude of miniature post-synaptic currents. These changes were also associated with increased a-syn release (18). How *in vivo* cell-specific neuronal firing patterns in the basal ganglia relates to the behavioral output is

unclear and a critical gap. The scant literature on *in vivo* recordings in rodent h*SNCA*

overexpression models comes from nigral, thalamic, and hippocampal neurons

evaluating movement and cognition (19-21). Pairing electrophysiological changes in

basal ganglia circuitry with vocal behavior offers an early and critical entry-point into

furthering our understanding of the effects of alpha-synucleinopathies on neural circuits.

Voice and speech deficits are prevalent in PD and affect quality of life (22, 23).

Early stage vocal deficits (e.g. a soft, monotonous voice, altered speaking rate) can

appear years before the movement symptoms, highlighting their utility as an early

diagnostic tool. The use of machine learning algorithms and smartphone applications

also support using vocal changes as an early biomarker of PD (24-37). Our group and

others have modeled PD-related vocal deficits using pharmacological and genetic

approaches in rodents and songbirds with data from these experiments implicating the

h*SNCA* gene in driving early vocal dysfunction (38-46).

Songbird models offer distinct advantages over rodents in exploring how the

brain encodes sensory and motor information, shaping vocal learning and production in

response to feedback (47). Males of the zebra finch species sing but the females do

not. The male finch song control system in the brain **(Fig 1A**) has similar neural

organization, genetics, and physiology of the basal ganglia to mammalian species (48-

50). The song control system has strong similarities to human speech and auditory

centers in the brain (51-57). Vocal learning and on-going song modification in the zebra

finch model can be investigated through the anterior forebrain pathway, a structure

homologous to the cortico-basal ganglia-thalamo-cortical loop in humans and other

mammals (50). Song-dedicated brain region Area X in the basal ganglia shares similar

genetics to human striatum, a major area involved in speech initiation and learning, and

has cell types characteristic of mammalian neuronal populations (48, 54, 58, 59).

Fig 1A. Finch song control circuitry. Brain nuclei in the adult male finch are

specialized for song learning and production (HVC and Area X: proper names; RA:

robust nucleus of the Arcopallium, lMAN: lateral magnocellular nucleus of the anterior

nidopallium). The solid lines represent cortico-brainstem circuits involved in vocal

production (finch: HVC to RA to brainstem). The dashed lines represent cortico-basal

ganglia-thalamo-cortico circuits for on-going vocal learning and modification (lMAN-Area

X-DLM-lMAN). Abbreviations: Cx – cortex. BG – basal ganglia. DLM – dorsal medial

nucleus of the thalamus. Th – thalamus. MB – midbrain. HB – hindbrain. Cb –

cerebellum. OB – olfactory bulb. Area X (violet) contains striatal, globus pallidus, and

interneurons but in the human BG, striatal and pallidal areas are separate. Several

auditory areas provide input to these vocal nuclei not shown here) (60, 61). Figure and

legend modified from finch brain image in Fig 1 of Medina et al. (2022).

Electrophysiological recordings in singing finches and anatomical tracings show

that MSNs are the dominant cell type in Area X and drive firing activity of globus

pallidus-like (PAL) projection neurons (50, 58, 62). As in mammals, GABAergic MSNs

and Globus-pallidus (GP, PAL) type neurons receive glutamatergic (Glu) inputs from

cortex (in finch-song nuclei HVC and lMAN, **Fig 1A**). GP neurons project to the dorsal

medial thalamic nucleus (DLM) which then relays its output to the cortex (63). While

studies of MSN activity in animal models and humans are difficult due to low

spontaneous activity (14, 64), finch MSNs are dynamically active during singing and

show temporally precise relationships to song structure (62, 65). DLM projections to

HVC can also modulate syllable transitions (66). High tonic firing in these PAL neurons

inhibits DLM, reducing excitation onto lMAN and then subsequently onto song nucleus

RA, leading to altered song output including variations in pitch (55, 62). Area X and

lMAN neurons exhibit firing variability that is related to birdsong features (55, 62). The

well-characterized neuronal populations in these brain nuclei and circuitry connections

make this a powerful model to study the impact of alpha-synucleinopathies on

vocalizations.

In our prior work, we showed that targeting Area X with an adeno-associated

virus (AAV) driving increased expression of the human (h) wild-type *SNCA* gene leads

to PD-like changes in the song including reduced duration and intensity. (45). We also

showed that a-syn overexpression results in protein aggregates in Area X cell bodies

(46). Here, as a critical first step, we categorize cell-specific firing patterns and

waveforms in Area X that are affected by a-syn overexpression compared to controls

using tetrode recordings in anesthetized finches immediately following singing behavior.

Our working hypothesis (**Fig. 1B**) is that h*SNCA* overexpression in Area X will decrease

firing rates and alter burst-firing in putative Area X MSNs. We tested this hypothesis by

recording from Area X neurons in anesthetized zebra finches.

Fig 1B. **Hypothetical model for a-syn driven changes in neural activity in Area X**

and consequences on song control circuitry. Cortical vocal nuclei (lMAN, HVC)

provide glutamatergic (Glu) input to Area X neurons. Striatal Area X MSNs provide

GABAergic input to globus-pallidus like (GP, PAL) neurons including those that inhibit

thalamic nucleus DLM. Here, we *predict* that abnormally elevated levels of AAV-driven

a-syn protein in Area X could alter intrinsic membrane properties of MSNs and/or

reduce Glu input to MSNs. Consequently, MSNs would reduce their firing rate (red

arrow), leading to increased GP PAL projection neuron firing and increased inhibition of

DLM. Increased DLM inhibition could be responsible for changes in amplitude

(loudness) and duration of the song syllables in finches with the AAV-h*SNCA* phenotype

(45).

Materials and Methods:

2.1. Finch subjects, song recording and analyses. All animal use was

approved by the Institutional Care and Use Committee at the University of Arizona.

Details on finch numbers, criteria for study inclusion/exclusion, and experimental

endpoints are provided in the following sub-sections. **Fig 1C** shows our experimental

timeline, workflow, and tetrode array.

Fig 1C. Tetrode array and experimental timeline. Schematic of AAV constructs and

coronal section showing bilateral song nucleus Area X (purple) modified from Fig 3b in

Medina et al. (2022). Song was collected within three days of the bilateral injection into

Area X and then again at two months post-virus injection followed by tetrode recordings

in anesthetized finches within five days, for a total experimental duration of

approximately two and a half months. At the end of the recording morning (≤3 hrs), the

finch was humanely euthanized, and the brain removed post-mortem for cryosectioning

followed by immunohistochemistry to confirm AAV and a-syn protein expression along

with tetrode placement.

Animal procedures followed Medina et al. (2022) and are summarized here. Only

finches that showed normal eating/drinking, perching, and movement in their cages

were selected for the study. Finches that did not sing prior to the pre-injection day were

excluded from the study and returned to the general aviary population. Adult non-

breeding male zebra finches (approx. 197-365 days post-hatch at time of euthanasia)

were moved into individual sound attenuation chambers (Eckel Noise Control

Technologies) and acclimated under a 14:10 h light-dark cycle for at least two days prior

to song recordings and neurosurgery. Behavioral experiments were conducted in the

morning from lights-on for two hours to match our prior study (45) along with ensuring

singing-driven neuronal activity just prior to conducting the tetrode recordings in

anesthetized finches.

Song recordings and analyses: Songs were acquired from individual males singing

alone (known as undirected song) using free-field omnidirectional microphones (Shure

93) suspended at set points over the center of the cage, connected to an audiobox

(Audiobox: 44.1 kHz sampling rate per 24-bit depth; Niles, IL) and recorded with Sound

Analysis Pro 2011 (SAP 2011, (67) http://soundanalysispro.com/). Song data collected

and used for this study can be found at the University of Arizona ReData repository

upon publication (10.25422/azu.data.27868266).

A basic unit of birdsong is defined as a motif; a motif is comprised of a sequence of

repeated syllables (**S1 Fig A**). Wav files from the two-hour recording period were joined

(Shuang's audio joiner) and then the entire file was viewed in Audacity

(https://www.audacityteam.org/). The most frequently occurring motif was then identified

in the joined wav file for two pre-surgical mornings (time point 0) and then two post-

surgical mornings (time point, 2 months, **Fig 1C**) by an investigator blind to the AAV

condition. The two-month timepoint was selected as the experimental endpoint for song

analyses and tetrode recordings in a non-survival surgery followed by humane

euthanasia. Our prior publications show that the most prominent song and protein

changes occurred at this time point in the AAV-h*SNCA* finches (45, 46). Following

morning lights-on, the first motif was identified in the wav file and then 25 consecutive

renditions of the individual syllable wav files were exported from Audacity from the four

mornings. Up to five unique syllables within a motif for each bird were included in the

analyses.

Syllable wav files were processed in Matlab R2014 using the SAP SAT Tools box

using code (provided by Dr. Nancy Day, Whitman College) to extract individual acoustic

features as a feature batch function. Raw amplitude (intensity) measurements were

calibrated based on sound chamber location as in our prior study (68). Mean, standard

deviation, and coefficient of variation (CV, standard deviation/mean) were then

calculated by averaging 50 scores (25 syllable copies x 2 recording sessions) across

time points 0 (pre-AAV injection) and 2 months post-AAV injection for each syllable

within the bird's motifs to account for natural, day to day fluctuations in song. For group

level comparisons, a normalization score was obtained per bird by dividing timepoint

two months (2) post-injection by pre-injection (0). After verifying that scores were not

normally distributed, scores were then compared using the Mann Whitney U test in

GraphPad Prism (version 10) between AAV-h*SNCA* and AAV-GFP groups for all

syllable types combined and then sorted by syllable type (**S1 Table**) given prior findings

on a-syn differential effects (45). Syllables were sorted into noisy, mixed, or harmonic

categories based on visual inspection and Wiener Entropy (WE) scores following our

prior publications (**S1 Fig**) (45, 69). We focused our analyses on comparing duration,

amplitude, and WE scores between the two AAV groups, and their correlations with the

electrophysiological analyses for an individual bird (see section on **Statistics** and **S1-2**

Figs). A subset of these birds was used in our separate publication evaluating regional

distribution of a-syn protein in finch song nuclei and linear correlations to syllable

features in Area X and lMAN (46).

As described in Medina et al. (2022), duration is the length of each syllable (in

seconds), and amplitude is the intensity or loudness of the syllable (in decibels-dB). WE

is a measure of uniformity in the power spectra (i.e., structure of the syllable), where

scores are calculated on a logarithmic scale; noisy syllables have values approaching

zero while harmonic syllables have more negative scores. The coefficient of variation

(CV) is calculated by dividing the standard deviation of 50 syllable renditions by the

mean score and represents variability across motifs for an acoustic feature, modulated

by the lMAN-Area X-DLM-lMAN loop (70).

2.2 Surgical procedures: viral injections: A total of 19 finches were originally

used for adeno-associated virus (AAV) bilateral injections into Area X (n=11, AAV5-

CBA-h*SNCA* [human alpha-synuclein]; n=8, AAV5-CBA-eGFP controls [Green

Fluorescent Protein]; CBA: chicken beta-actin promoter). One bird in each group
spontaneously died prior to the two month post-AAV injection experimental endpoint
due to unknown causes and therefore, they were removed from the study. We
alternated the order of the AAV experimental vs. control virus surgeries when two
surgeries were done on the same day.

The surgery suite and instruments were prepared for aseptic survival surgery under PI
Miller's approved IACUC Protocol 13-489. Following two hours of morning song
collection, finches were put under isoflurane anesthesia with oxygenation (2-3%, 0.5
liters/min oxygen) and set up in the stereotaxis. The analgesic lidocaine (0.05ml) was
subcutaneously injected across the planned incision site of the scalp at four locations.
For the bilateral injections into Area X, stereotaxic coordinates were used from the
bifurcation of the mid-sagittal sinus: 3.1 mm rostral-caudal, 1.62 mm medio-lateral, a
depth of 3.1 mm, and head angle of 40 degrees. These coordinates avoid indirect
targeting of song nucleus lMAN with virus as previously described (45). Viruses (AAV5)
were obtained from the University of North Carolina Viral Vector Core through financial
support of the Michael J. Fox Foundation for Parkinson's Research
(https://www.michaeljfox.org/research-tools).

A glass pipette was fitted into a Nanoject II pressure-injector and back-filled with mineral
oil, then loaded with either AAV. Approximately 500 nL of virus was injected at a rate of
27.6 nL/injection every 15 seconds for a total of 18 injections followed by five minutes of
wait time then inspection of pipette tip for clogging. Upon removal of isoflurane
anesthesia, birds were monitored for any signs of pain and distress while on a
homeothermic heating blanket. Within a half-hour, birds were noted to be alert, upright,

and vocalizing and did not require any post-operative analgesia. They were returned to

their sound chambers and monitored for any surgical complications, abnormal postural

or movement, eating and drinking behavior for at least three days post-surgery by

research personnel fully trained in bird care. Their drinking water was supplemented

with an oral antibiotic dose of TMS (Sulfamethoxazole and Trimethoprim Oral

Suspension, USP grade; 1ml/100 ml deionized water). University Animal Care staff

performed daily checks on animal welfare with Miller lab personnel providing oversight

throughout the duration of the experiment.

2.3 Tetrode Implantation and Electrophysiology

Surgical procedure and implantation: Approximately two months post-virus

injection, 17 finches, out of the 19 originally included (two birds died post-virus injection

and prior to tetrode implantation), were anesthetized with isoflurane (2-3%, constant

flow rate of 0.5-0.8 L of oxygen/min). While under anesthesia, they underwent acute

electrophysiological recordings of right hemisphere Area X as a non-survival surgery

approved under PI Miller's IACUC Protocol #13-489. Breathing rate was monitored

throughout the experiment, and an infrared heating pad (Kent Scientific) maintained

body temperature at 37ºC. The same stereotaxic coordinates were used for the AAV

injections targeting Area X. **Ground wire insertion:** A ground/reference wire was

inserted into the left hemisphere, contralateral to the virus injection site (mid-sagittal

sinus: 3.1 mm rostral-caudal, 1.62 mm medio-lateral). After performing the craniotomy,

the dura was pierced with a 30G needle, and the end of the ground wire was lowered to

a depth of 2-3 mm. Gel superglue was used to secure the wire to the acid-etched skull

(acid from the Metabond kit). Zip Kicker (Pacer Technology) was used to rapidly cure

the superglue. **Electrode array insertion:** The existing craniotomy on the right

hemisphere (used for virus injection) was expanded to a diameter of ~1.5 mm to better

accommodate the electrode array and to have access to dura that was undamaged

during virus injection. Tetrode position was confirmed in Area X post-mortem following

electrophysiological recordings (**Fig 2**).

Electrophysiological recordings: Neurons from Area X were recorded using a

custom-made 800 µm diameter tetrode array composed of 8 twisted-wire tetrodes.

Tetrodes were constructed from polyamide-insulated nickel chromium steel wire (~12

µm in diameter; see **Fig 1C**). Prior to insertion, tetrodes were coated with DiI to allow

anatomical tracing of the electrode path (**Fig 2A**). The tetrode array was lowered in

~100 µm increments until the tip reached Area X at ~3100 µm (right hemisphere) and

allowed to stabilize for 5-10 minutes prior to recording. Neural data was acquired at 20

kHz using the INTAN neural recording system (Intan Technologies, Inc.). Neural

recordings from Area X neurons occurred in 5-minute blocks. After each 5-min block of

data was acquired, the electrode array was lowered ~50 µm and another recording

block was acquired. All recordings were restricted to Area X (3100-4100 µm). Data was

spike sorted using Kilosort v2.5 (https://zenodo.org/records/4482749) (71) and Phy

(https://github.com/cortex-lab/phy). Methods for cell-type estimation are described in

Results.

Data/Statistical Analysis:

All single-unit data was analyzed using Matlab (R2025a). All Matlab analysis code is

available upon publication at: https://github.com/CowenLab/ZebraFinch. Out of the 17

finches implanted with the tetrode array, four finches in the ASYN group yielded no

recordings (e.g. neurons were quiet), and one GFP control finch showed no virus

expression detected in Area X post-mortem. These five finches were therefore excluded

from our data analyses but humanely euthanized with the other finches at the end of

that day's electrophysiological recording period. Thus, n=12 finches (n=6/per AAV

group) were used for electrophysiological and song analyse presented in the Results

section.

Measurement of burst activity: The temporal structure of the firing activity of individual

neurons was quantified using "local variance" (LV), a measure that assesses the

variability of inter-spike intervals (ISIs) and is robust to changes in firing rate (72). LV is

computed from the vector of ISIs for each neuron as shown in the formula below:

$$LV = \frac{3}{n-1}\sum_{i=1}^{n-1}\left(1 + \frac{4I_i I_{i+1}}{(I_i+I_{i+1})^2}\right)\left(1 - \frac{4R}{I_i+I_{i+1}}\right), \qquad (1)$$

where I_i and I_{i+1} are the i-th and $i+1$ ISI and n is the number of ISIs. LV scores ~0

indicate tonic firing, scores ~1 show Poisson firing, and scores >1 indicate burst-like

firing patterns.

Parametric and non-parametric tests were used for hypothesis testing. For group-level

analysis of firing-rate data (n = 201 categorized neurons from n = 12 finches) non-

parametric tests were used as firing rates do not follow normal distributions (73).

Consequently, rank-sum tests were used for two-group comparisons and the Kruskal-

Wallis test was used for multi-group data. The Holm-Bonferroni correction was used to

correct for multiple comparisons. For normally distributed data, ANOVA and t-tests were

used followed by Tukey or Holm-Bonferroni post-hoc corrections. Alpha was set to 0.05.

Relationships between neural data and each bird's song features (e.g., mean Entropy,

Duration, and Amplitude) were investigated using a generalized linear regression model

where the outcome variable was neural activity (e.g., firing rate) and the independent

variables were song features and group category (ASYN or GFP) as described in

Results and S2 Fig. Multicollinearity between independent variables can impair the

ability to interpret regression models. We observed that some song features did indeed

covary (r > 0.4 for 5 song feature pairs). To address this, we repeated the regression

using the first three principal components of the independent variable matrix and

compared results to those from the original regression model. As reported in **S2 Fig**,

similar results were obtained from the original and dimensionality-reduced model.

2.4 Tissue processing and immunohistochemistry

Immediately following the end of the neural recording session, finches were humanely

euthanized by isoflurane inhalant overdose and transcardially perfused with warmed

saline followed by chilled 4% paraformaldehyde in Dulbecco's Phosphate Buffer Saline.

Fixed brains were cryoprotected in 20% sucrose overnight then cryosectioned in the

coronal plane at 30 µm on a Microtome cryostat through Area X. Tissue was processed,

following our previously published methods (45, 46) and summarized here: Hydrophobic

borders were drawn on the slides, using a pap pen (ImmEdge, Vector Labs) followed by

3 X 5 minute washes in TBS with 0.3% Triton X (Tx). To block non-specific antibody

binding, the tissue was then incubated for one hour at room temperature with 5% goat

serum (Sigma #G-9023) in TBS/0.3% Tx then 3 x 5-minute washes in 1% goat serum in

TBS/0.3% Tx were performed. A primary antibody was used against a-syn (1:250,

rabbit, Proteintech 10842-1-AP, RRID: AB_2192672). GFP expression was robust

enough to be visualized without an antibody.

The primary antibody was incubated in a solution of 1% goat serum in TBS/0.3% Tx

overnight at 4°C. One tissue section per slide was used as a 'no primary' control and a

pre-absorption control experiment was performed to validate antibody specificity in a

companion study evaluating the distribution of a-syn protein following AAV injection

(46). The next day, sections were washed 5 x 5 minutes in TBS/0.3% Tx and incubated

for three hrs at room temperature in a fluorescently conjugated secondary antibody

(1:1000: goat anti-rabbit 647 A-21245, RRID:AB_2535813). After secondary incubation,

sections were washed 3 x 10 minutes in TBS followed by 2 x 5 washes in filtered TBS.

Slides were then cover-slipped in Pro-Long Anti-Fade Gold mounting medium

(Molecular Probes, P36930). Tissue was imaged using a Zeiss Axio Observer 7 with

Apotome III Microscope under the University of Arizona Imaging Cores - Optical Core

Facility (RRID:SCR_023355).

Results:

Virally-driven expression is confirmed in song nucleus Area X

Fig 2 is a representative example of AAV expression in Area X. Two months post-virus

injection, we detected GFP cell bodies in Area X (**Fig 2A,C**) accompanied by low a-syn

protein expression (**Fig 2B**). By contrast, a finch overexpressing a-syn showed staining

in processes and cell bodies (**Fig 1D**). These results are consistent with our two prior

publications (45, 46).

Fig 2. AAV expression in finch song nucleus Area X. The Area X region is

represented by the dotted line. Small arrowheads denote the border between the

striatum and the nidopallium. Staining for alpha-synuclein (a-syn) protein is shown in

red. **A)** An AAV control expressing finch has GFP labeled cell bodies (green) in Area X,

and the large arrowhead denotes the last recorded tetrode position (yellow). **B)** A-syn

protein expression is low in Area X compared to surrounding basal ganglia in this AAV-

GFP finch. **C)** View of (A) showing the GFP channel only. **D)** An AAV-h*SNCA*

expressing finch shows widespread expression of a-syn in processes throughout Area

X. Inset in bottom left corner shows a zoomed in view of bead-like a-syn protein in the

processes (inset scale bar = 10 μm).

Extracellular identification of neuronal subtypes in anesthetized

finches

Area X neurons share much in common with neurons in the mammalian striatum

and pallidum. For example, the most abundant Area X neuron type is homologous to

the mammalian medium spiny neuron (MSN,(58)), with Area X and mammalian MSNs

sharing wide waveforms, low firing rates, and bursting activity (74-76). There is also

evidence that Area X contains striatal fast-firing interneurons, cholinergic interneurons,

and low-threshold spiking neurons (48, 74, 77). Furthermore, Area X contains two

classes of fast-firing pallidal-like neurons: local-circuit (HF-1) and thalamus-projecting

(HF-2) (59, 78). These six cell classes were identified by their firing rates and waveform

features using multiple approaches from this literature, including correlating neuronal

activity *in vivo* with birdsong and patch-clamp recording in brain slices. Our present

study is constrained by the fact that all data was acquired from extracellular electrodes

implanted in anesthetized finches. We found no prior publications with recordings from

Area X of adult anesthetized finches; prior work includes some anesthetized recordings

from lMAN and Area X in the context of juvenile song learning (79). Consequently, our

classification was informed by criteria developed from *in vivo* recordings from Area X in

singing finches and restricted to waveform shape, firing rate, and assessment of

bursting activity in the following five identified categories (74, 78, 80).

Neuron categorization and criteria (results shown in **Fig 3**):

1) **Wide-Low-Rate (WLR) Criteria**: Neurons firing ≤ 6 Hz with a peak half width > 0.25

ms or a peak-to-trough width of > 0.35 ms (**Fig 3 A,D,E**). WLR neurons were also

more 'bursty'' relative to other neuronal types (**Fig 3F**) as measured by Local

Variance (LV, see Methods (81)) and were the most abundant of the five identified

cell types (**Fig 3C**). Together, these features suggest that WLR cells were MSN-like

given evidence that MSNs are the most prominent cell type in the striatum, exhibit

wide waveforms, have low firing rates, and express bursting activity (74) (75, 76).

2) **Wide-High-Rate (WHR) Criteria**: Neurons with a firing rate between 20 and 50 Hz

with a peak half width > 0.25 ms or a peak-to-trough width of > 0.35 ms (**Fig

3A,D,E**). Post-classification analysis also showed that these neurons fired tonically

(LV < 1, see **Fig 3F**). Tonically active WHR cells have features consistent with

striatal cholinergic interneurons (74).

3) **Narrow-Low-Rate (NLR) Criteria**: Narrow waveform neurons firing ≤ 6 Hz (74)

having a peak half width < 0.25 ms or a peak-to-trough width of < 0.35 ms (**Fig**

3A,D,E).

4) **Narrow-High-Rate (NHR) Criteria**: Narrow waveform neurons with rates ≥ 8 Hz but

< 50Hz (**Fig 3A,D,E**). Narrow waveforms are characteristic of local circuit fast-firing

inhibitory neurons.

5) **Very-High-Rate (VHR) Criteria**: Neurons firing ≥ 50 Hz (**Fig 3A,D,E**). Area X

pallidal-like projection neurons are characterized by exceedingly high firing rates (74,

78). Consequently, we established a category for neurons with rates ≥ 50 Hz.

Waveform features were not used to further subdivide neurons in this group given its

small size (n = 18 neurons, **Fig 3C**). Analysis of VHR firing activity revealed that

these neurons exhibited decidedly tonic activity, with a mean LV of 0.51 (**Fig 3F**).

These data suggest that the identified VHR neurons are type-2 pallidal-like

projection neurons (HF-2 in (78) (82)) given the presence of tonic firing and relatively

wide waveforms. Furthermore, the small size of this group is consistent with other

reports that suggest PAL neurons comprise approximately just 5.4% of the Area X

cell population (58).

Fig 3. Identification of five classes of Area X neurons in anesthetized finches. A)

Mean shape for each of the five identified classes of neurons (n = 201 neurons from n =

12 finches). Neurons were categorized based on spike width and firing rate (see

Results). Amplitude is normalized such that the minimum (trough) = -1. Colors indicate

neuron subtype (see key). **B)** Inter-spike interval distributions for each neuronal

subtype. Y-axis indicates probability (p). **C)** Histogram indicating the number of

identified neurons in each category. **D)** Box plot of the half-width (ms-milliseconds) for

each neuronal group. Each dot indicates an individual neuron. Horizontal red lines

indicate the median. Hypothesis tests were not performed as half-width was a key

criterion for classification. **E)** Firing rates for each neuronal group (H-Hertz). Each dot

indicates an individual neuron. Hypothesis tests were not performed as firing rate was a

key criterion for classification. **F)** Burst firing differed between groups. While neurons

were not categorized based on burst-firing, the measure of bursting (Local Variance-LV)

indicated that Wide-Low-Rate neurons had increased burst firing (LV>1) relative to other

groups (ANOVA $p < 0.000001$, Tukey post-hoc comparison. Thick red lines indicate

significant post-hoc differences (*$p < 0.05$, **$p < 0.01$ (not shown), ***$p < 0.001$). The

red lines in the box plots indicate the median and the whiskers indicate 1.5X the

Interquartile Range.

Reduced Area X firing rates in ASYN overexpressing finches

We predicted that a-syn expression would reduce putative MSN (WLR) neuron

activity, thus releasing putative PAL neurons (VHR) from inhibition. Consequently, we

investigated whether firing rates of WLR neurons were reduced in the ASYN group and

whether rates in VHR neurons were increased. Visual inspection of raster plots of WLR

activity (**Fig 4A-B**) suggested reduced WLR firing activity in the ASYN group. Statistical

comparison of firing rates of GFP and ASYN neurons identified a significant difference

(**Fig 4C**, rank-sum, p = 0.005 before correction and p = 0.026 after Holm-Bonferroni

correction). No significant between-group differences were identified in the other

neuronal classes, including the VHR neurons.

Fig 4. Reduced firing rates of WLR neurons in the ASYN group. A-B) Raster plots

of WLR in the GFP and AAV-h*SNCA* (ASYN) groups. The x-axis shows time in

seconds. Each row is a raster for a distinct neuron. **C)** Between-group comparison (GFP

vs. ASYN) for each of the five neuron types. WLR neurons were only found in 4/6

finches. Firing rates for WLR neurons were lower in the ASYN group relative to the GFP

control (rank-sum test, p = 0.005 prior to correction, *p = 0.026 after Holm-Bonferroni

correction).

Burst firing was not changed in ASYN overexpressing finches

While burst firing is a canonical property of MSNs, MSN burst firing may become

dysregulated in PD. Indeed, altered MSN burst firing has been reported in the 6-OHDA

mouse model of PD (83), non-human MPTP primate models (84), and in human PD

patients (64). Consequently, we predicted that burst firing will be affected by ASYN

overexpression in finch Area X. To investigate this, the local variance (LV) measure

(72), see Methods) was used to evaluate the pattern of inter-spike intervals and quantify

the degree that neuronal activity was either tonic (low LV), Poisson (LV near 1), or

bursty (LV > 1). Contrary to our prediction, analysis of spiking statistics with LV did not

identify any difference between GFP and ASYN groups in LV values for any neuronal

subtype (**Fig 5**, t-test, all p > 0.05 after Holm-Bonferroni post-hoc correction). Thus,

while ASYN significantly reduced firing rates of WLR neurons, these changes were not

accompanied by a change in burst-firing.

475
476

Fig 5. Local Variance (LV), a measure of bursting, is not different between ASYN

and GFP control groups. Between-group comparisons (GFP vs. ASYN) were made for

each of the five neuron types, and no between-group differences were identified in any

of the neuron classes (rank-sum test, all p > 0.05 after Holm-Bonferroni post-hoc

correction).

Extracellular waveform shape was altered in wide-waveform neurons

While waveform features and firing rates are commonly used to classify neurons

into putative subtypes (74, 80, 85), inferring physiological properties of neurons from

extracellular recordings is highly problematic as waveform features vary as a function of

recording location relative to the morphology of the neuron (86-88). Even so, we did

observe consistent differences in extracellular waveform shapes between ASYN and

GFP animals that encourage future investigations. Specifically, we found that WLR and

WHR neurons showed a large positive extracellular 'rebound' following the initial trough

(**Fig 6A**). This was determined by first normalizing the waveform of each neuron such

that the trough = -1 and baseline was zero. Consequently, positive values indicate

proportional changes above baseline. The "Rebound Peak" for each neuron was

measured as the normalized amplitude of the largest peak following the time of the

trough (t = 0 ms). Values for each neuron type are shown in **Fig 6B** for the GFP and

ASYN groups. Rank-sum tests were performed, and significant between-group

differences were identified in the WLR and WHR groups (p_{WLR} = 0.003, p_{WHR} = 0.0002

after Bonferroni-Holm correction).

Fig 6. **ASYN group shows enhanced post-peak rebound for wide-waveform**

neurons. A) Mean waveforms for the ASYN and GFP groups for each neuron type. For

each waveform, data was normalized such that the distance from baseline to the trough

= -1. Inspection of these plots suggested that wide-waveform (WLR, WHR) ASYN

neurons had a larger rebound following the initial trough. **B)** Statistical analysis of the

rebound was performed by measuring the normalized amplitude of the largest peak

following t=0 (rebound peak) for each neuron (dots). These values were compared

between groups using rank-sum tests and p values were adjusted using the Holm-

Bonferroni correction. **p<0.01, ***p<0.001. Error bars in box plots indicate ±SEM.

No observed correlation between song features and neuronal firing properties

513   Next, we investigated whether ASYN and GFP birds differed in their song

514   features and if variation in mean WLR neuronal activity in each bird correlated with song

515   features. Analysis of syllable-level features failed to identify any between-group

516   difference (Mann-Whitney U test, p > 0.05, **S1 Fig, S1 Table**).

517   Investigation of whether firing rates correlated with song features was performed

518 by first calculating the mean firing rate of WLR neurons (the largest group and the

519 putative MSNs) for each bird. Two birds had to be excluded from the analysis as no

520 WLR neurons were identified in these animals. Mean WLR firing rates were used as the

521 outcome variable for a generalized linear model where the independent variables were

522 song features (mean amplitude, duration, and entropy and CV amplitude, duration, and

523 entropy) and the group (ASYN or GFP). No significant relationship between song

524 features and firing rate was identified ($F_{(10,2)}$ = 11.7, p = 0.082);**S2 Fig**.

527 **4. Discussion**

528   The goal of this study was to obtain new information about the cellular and

529 network-level activity affected by overexpression of human a-syn protein in an adult

530 male zebra finch model. To carry out this goal, we performed extracellular recordings

531 from song nucleus Area X of the basal ganglia in anesthetized adult male finches and

532 identified five distinct cell types. Our main finding was that wide, low firing rate (WLR)

533 neurons, putative MSNs, show reduced firing activity and enhanced post-inhibitory

534 rebound in the a-syn overexpression condition compared to controls. This aligns with

535 our working hypothesis that abnormally elevated a-syn protein levels in Area X will

536 decrease MSNs firing rate. We also predicted that GP-projecting neurons would show

537 reduced firing in Area X. VHR neurons may represent these GP-projecting neurons, but

538 they did not show a difference in firing rate between the ASYN and GFP control groups.

This finding may reflect low statistical power given the small number of VHR neurons

recorded.

A benefit of anesthetized recordings is that they allow for high data throughput,

enabling the recording of a substantial proportion of neurons without confounds due to

animal movement. A limitation of this approach in our finch model is that on-going real

time modification of neural activity related to song production cannot be evaluated. It is

also likely that neuronal activity was dampened by isoflurane anesthesia. While

information is scarce on the effects of isoflurane on *in vivo* recordings from the striatum

of naïve finches or rodents, patch clamp studies in rat brain slices show that brief

isoflurane exposure can dampen sodium currents in hippocampal neurons (89) and

reduce spiking and bursting modes in thalamic neurons (90). Our future investigations

will compare isoflurane-induced anesthesia on Area X of naïve finches versus awake,

singing finches where we can examine how a-syn modulates neuronal dynamics under

more physiologically relevant conditions.

The observed reduction in WLR (putative MSN) firing rates observed in Area X of

the ASYN finches and lack of bursting activity contrasts with some measures of MSN

activity reported in human PD patients and primate PD models. For example, striatal

projection neurons in the human basal ganglia of PD patients undergoing deep brain

stimulation had high frequency neuronal firing patterns resembling the advanced

parkinsonism of the Non-Human Primate model, with hyperactivity and "bursty"

neuronal firing activity (14, 64). In a mouse model of bilateral dopamine depletion (via

the neurotoxin 6-OHDA) in the dorsolateral striatum, single unit recordings from MSNs

showed reduced bursting activity during movement. Intriguingly, they classified MSNs

into two types with type 1 showing increased firing during movement and type 2,

decreased firing with no difference in firing rate when the mouse was stationary (83). In

freely moving rats, 6-OHDA treatment increased oscillatory synchronization in the

striatum (91). Future recordings in singing, ambulatory finches with the ASYN

phenotype will determine if hyperactivity of MSNs and differential sub-types exist,

evidence for dopamine loss, and how these various sub-components contribute to

changes in real-time song output.

The reduced firing rates of WLR neurons in ASYN expressing finches may reflect

reduced mean frequency of spontaneous excitatory postsynaptic glutamatergic currents

from cortical song nucleus lMAN onto Area X MSNs and GP neurons. Striatal

recordings from h*SNCA* overexpressing mice show reduced glu currents (12). Future

work using electrophysiological recordings from Area X brain slices *in vitro* can

determine whether glu input is perturbed in our ASYN condition.

ASYN-driven abnormal changes in synaptic input to Area X may also be

accompanied by altered intrinsic membrane properties. Both the WLR and WHR

neuronal classes showed greater positive rebound potential following the initial spike in

the ASYN expressing group compared to GFP controls. While extracellular recordings

of action potentials cannot reliably identify after-hyperpolarization (AHP) currents, the

observed rebound potential occurs during the typical time interval of the AHP,

suggesting that membrane properties in the ASYN neurons are altered. The AHP is

regulated by K^+ currents, and several classes of K^+ channels are implicated in altered

neuronal excitability in PD models (92). The significantly larger rebound potential in the

ASYN expressing group may reflect an enhanced AHP-related conductance. Patch-

clamp electrophysiology in brain slices is a necessary approach to confirm true

fluctuations in membrane properties during the AHP period. Changes in AHP-related

conductance could result from increased SK/BK channel activity (93) (94)) or a

reduction in I_h current (95, 96), leading to greater hyperpolarization following spikes and

dampened excitability. In nigral DA neurons, abnormal a-syn oligo aggregates increase

the conductance of K^+_{ATP} channels leading to reduced firing (97). In our finch model, a-

syn overexpression could progressively decrease intrinsic excitability in MSNs via these

channels, and/or others. To evaluate the specific role of a-syn expression on MSNs,

future directions will employ a custom-made AAV containing a CAMKII promoter that

has been previously used in naïve finches to evaluate MSNs firing activity (98). By

selectively overexpressing a-syn in MSNs, we can evaluate the relationship between

the extent of protein pathology and the neural activity changes underlying song. In a

recent publication, we found that regional changes in virally-expressed human a-syn

protein within Area X correlated with changes in syllable level features. For example, we

detected a positive correlation between right hemisphere a-syn proteinopathy and a

reduction in the variation of harmonic syllable duration, among other features, correlated

to the right hemisphere Area X (46). Our neural recordings here are also from right

hemisphere Area X; the contribution of the left Area X is not known. Prior studies

suggest a transition in the unilateral activation of the hemispheres: right-side dominance

of song nucleus HVC is detected in adult birdsong (99) while the left-side dominates

during juvenile learning (100). Thus, when assessing cell-type changes in electrical

activity within brain nuclei aligned with real-time singing behavior, factors to take into

account include the recording hemisphere, the distribution of protein expression, and

the presence of pathology, e.g. a-syn aggregates in individual cell types.

Conclusions

Our study takes an important first step forward in evaluating the consequences of

a-syn overexpression on the activities of neuronal subtypes involved in vocal behavior.

A growing body of research has elucidated the role that neurons in the cortico-striatal-

thalamic circuit play in male zebra finch song production. By introducing human a-syn to

the zebra finch model, this study supports a systems- and cellular-level investigation of

human vocal deficits in PD. This is important given that current medications and brain

stimulation used in the treatment of PD fail to resolve the vocal deficits and can worsen

them (101-103). Data from pre-clinical animal models such as the songbird can provide

important mechanistic insight needed for the development of new brain-targeted cellular

therapies for the vocal dysfunction.

CRediT Contributions
626
CRediT authorship contribution statement: Brian R. Dominguez: Data curation,

Investigation, Methodology, Validation, Visualization, Writing-Original Draft, Writing-

review & editing. Gabriel Holguin: Data curation, Formal Analysis, Investigation,

Methodology, Software, Validation, Visualization, Writing-Original Draft, Writing-review

& editing. Madeleine S. Daly: Data curation, Formal Analysis, Investigation,

Methodology, Validation, Writing-review & editing. Reed T. Bjork: Investigation,

Methodology, Validation, Visualization, Writing-review & editing. Stephen Cowen:

Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology,

Software, Supervision, Validation, Visualization, Writing-Original Draft, Writing-review &

editing. Julie E. Miller: Conceptualization, Data curation, Formal Analysis, Funding

acquisition, Investigation, Methodology, Project administration, Resources, Supervision,

Validation, Visualization, Writing-original draft, Writing-review & editing.

639
640

Acknowledgements

643 We thank Kathy Dillon, Michelle Gordon, and Mauricio Serna for technical contributions

644 along with University of Arizona Animal Care staff and Mr. Douglas Cromey of the

645 University of Arizona Imaging Cores - Optical Core Facility (RRID:SCR_023355).

References

1.   Polymeropoulos MH, Lavedan C, Leroy E, Ide SE, Dehejia A, Dutra A, et al.

Mutation in the alpha-synuclein gene identified in families with Parkinson's disease.

Science. 1997;276(5321):2045-7.

2.   Chartier-Harlin M-C, Kachergus J, Roumier C, Mouroux V, Douay X, Lincoln S, et

al. α-synuclein locus duplication as a cause of familial Parkinson's disease. The Lancet.

2004;364(9440):1167-9.

3.   Singleton A, Farrer M, Johnson J, Singleton A, Hague S, Kachergus J, et al. α-

Synuclein locus triplication causes Parkinson's disease. Science. 2003;302(5646):841-.

4.   Mak E, Su L, Williams GB, O'Brien JT. Neuroimaging characteristics of dementia

with Lewy bodies. Alzheimer's research & therapy. 2014;6:1-10.

5.   Koga S, Dickson DW. Recent advances in neuropathology, biomarkers and

therapeutic approach of multiple system atrophy. Journal of Neurology, Neurosurgery &

Psychiatry. 2018;89(2):175-84.

6.   Calo L, Wegrzynowicz M, Santivanez-Perez J, Grazia Spillantini M. Synaptic

failure and alpha-synuclein. Movement disorders: official journal of the Movement

Disorder Society. 2016;31(2):169-77.

7.   Marino G, Calabresi P, Ghiglieri V. Alpha-synuclein and cortico-striatal plasticity

in animal models of Parkinson disease. Handbook of clinical neurology. 2022;184:153-

66.

8.   Calabresi P, Mechelli A, Natale G, Volpicelli-Daley L, Di Lazzaro G, Ghiglieri V.

Alpha-synuclein in Parkinson's disease and other synucleinopathies: from overt

neurodegeneration back to early synaptic dysfunction. Cell death & disease.

2023;14(3):176.

9.   Watson JB, Hatami A, David H, Masliah E, Roberts K, Evans CE, et al.

Alterations in corticostriatal synaptic plasticity in mice overexpressing human alpha-

synuclein. Neuroscience. 2009;159(2):501-13.

10.   Tozzi A, de Iure A, Bagetta V, Tantucci M, Durante V, Quiroga-Varela A, et al.

Alpha-Synuclein Produces Early Behavioral Alterations via Striatal Cholinergic Synaptic

Dysfunction by Interacting With GluN2D N-Methyl-D-Aspartate Receptor Subunit. Biol

Psychiat. 2016;79(5):402-14.

11.   Tozzi A, Sciaccaluga M, Loffredo V, Megaro A, Ledonne A, Cardinale A, et al.

Dopamine-dependent early synaptic and motor dysfunctions induced by α-synuclein in

the nigrostriatal circuit. Brain. 2021;144(11):3477-91.

12.   Wu N, Joshi PR, Cepeda C, Masliah E, Levine MS. Alpha-synuclein

overexpression in mice alters synaptic communication in the corticostriatal pathway.

Journal of neuroscience research. 2010;88(8):1764-76.

13.   Lam HA, Wu N, Cely I, Kelly RL, Hean S, Richter F, et al. Elevated tonic

extracellular dopamine concentration and altered dopamine modulation of synaptic

activity precede dopamine loss in the striatum of mice overexpressing human

α-synuclein. Journal of neuroscience research. 2011;89(7):1091-102.

14.   Beck G, Singh A, Papa SM. Dysregulation of striatal projection neurons in

Parkinson's disease. Journal of Neural Transmission. 2018;125(3):449-60.

15.   Durante V, de Iure A, Loffredo V, Vaikath N, De Risi M, Paciotti S, et al. Alpha-

synuclein targets GluN2A NMDA receptor subunit causing striatal synaptic dysfunction

and visuospatial memory alteration. Brain. 2019;142(5):1365-85.

16.   Wilson CJ, Groves PM. Spontaneous firing patterns of identified spiny neurons in

the rat neostriatum. Brain research. 1981;220(1):67-80.

17.   Kita H, Kita T. Role of striatum in the pause and burst generation in the globus

pallidus of 6-OHDA-treated rats. Frontiers in systems neuroscience. 2011;5:42.

18.   Henkel LM, Kankowski S, Moellenkamp TM, Smandzich NJ, Schwarz S, Di

Fonzo A, et al. iPSC-derived striatal medium spiny neurons from patients with multiple

system atrophy show hypoexcitability and elevated α-synuclein release. Cells.

2023;12(2):223.

19.   Lobb CJ, Zaheer AK, Smith Y, Jaeger D. In vivo electrophysiology of nigral and

thalamic neurons in alpha-synuclein-overexpressing mice highlights differences from

toxin-based models of parkinsonism. Journal of neurophysiology. 2013;110(12):2792-

805.

20.   Brys I, Nunes J, Fuentes R. Motor deficits and beta oscillations are dissociable in

an alpha-synuclein model of Parkinson's disease. European Journal of Neuroscience.

2017;46(3):1906-17.

21.   Robson E, Tweedy C, Manzanza N, Taylor J-P, Atkinson P, Randall F, et al.

Impaired fast network oscillations and mitochondrial dysfunction in a mouse model of

alpha-synucleinopathy (A30P). Neuroscience. 2018;377:161-73.

22.   Sapir S. Multiple factors are involved in the dysarthria associated with

Parkinson's disease: a review with implications for clinical practice and research.

Journal of speech, language, and hearing research : JSLHR. 2014;57(4):1330-43.

23.   Ma A, Lau KK, Thyagarajan D. Voice changes in Parkinson's disease: What are

they telling us? Journal of Clinical Neuroscience. 2020;72:1-7.

24.   Stewart C, Winfield L, Hunt A, Bressman SB, Fahn S, Blitzer A, et al. Speech

dysfunction in early Parkinson's disease. Movement disorders: official journal of the

Movement Disorder Society. 1995;10(5):562-5.

25. Ho AK, Iansek R, Marigliani C, Bradshaw JL, Gates S. Speech impairment in a

large sample of patients with Parkinson's disease. Behav Neurol. 1998;11(3):131-7.

26. Harel B, Cannizzaro M, Snyder PJ. Variability in fundamental frequency during

speech in prodromal and incipient Parkinson's disease: a longitudinal case study. Brain

Cogn. 2004;56(1):24-9.

27. Skodda S, Schlegel U. Speech rate and rhythm in Parkinson's disease.

Movement disorders: official journal of the Movement Disorder Society. 2008;23(7):985-

92.

28. Skodda S, Rinsche H, Schlegel U. Progression of dysprosody in Parkinson's

disease over time--a longitudinal study. Movement disorders: official journal of the

Movement Disorder Society. 2009;24(5):716-22.

29. Rusz J, Cmejla R, Ruzickova H, Ruzicka E. Quantitative acoustic measurements

for characterization of speech and voice disorders in early untreated Parkinson's

disease. The Journal of the Acoustical Society of America. 2011;129(1):350-67.

30. Defazio G, Guerrieri M, Liuzzi D, Gigante AF, di Nicola V. Assessment of voice

and speech symptoms in early Parkinson's disease by the Robertson dysarthria profile.

Neurol Sci. 2016;37(3):443-9.

31. V. Berisha JL, T. Huston, A. Wisler, Y. Jiao, and J. Eig. Float Like a Butterfly

Sting Like a Bee: Changes in Speech Preceded Parkinsonism Diagnosis for

Muhammad Ali. Proceedings of the 2017 Interspeech Conference.

32. Rusz J, Tykalová T, Novotný M, Růžička E, Dušek P. Distinct patterns of speech

disorder in early-onset and late-onset de-novo Parkinson's disease. npj Parkinson's

Disease. 2021;7(1):98.

33. Šimek M, Rusz J. Validation of cepstral peak prominence in assessing early

voice changes of Parkinson's disease: Effect of speaking task and ambient noise. The

Journal of the Acoustical Society of America. 2021;150(6):4522-33.

34. Berus L, Klancnik S, Brezocnik M, Ficko M. Classifying Parkinson's disease

based on acoustic measures using artificial neural networks. Sensors. 2018;19(1):16.

35. Solana-Lavalle G, Galán-Hernández J-C, Rosas-Romero R. Automatic Parkinson

disease detection at early stages as a pre-diagnosis tool by using classifiers and a small

set of vocal features. Biocybernetics and Biomedical Engineering. 2020;40(1):505-16.

36. Suppa A, Costantini G, Asci F, Di Leo P, Al-Wardat MS, Di Lazzaro G, et al.

Voice in Parkinson's disease: a machine learning study. Frontiers in Neurology.

2022;13:831428.

37. Lim WS, Chiu S-I, Wu M-C, Tsai S-F, Wang P-H, Lin K-P, et al. An integrated

biometric voice and facial features for early detection of Parkinson's disease. npj

Parkinson's Disease. 2022;8(1):145.

38. Ciucci MR, Ma ST, Fox C, Kane JR, Ramig LO, Schallert T. Qualitative changes

in ultrasonic vocalization in rats after unilateral dopamine depletion or haloperidol: a

preliminary study. Behav Brain Res. 2007;182(2):284-9.

39. Gombash SE, Manfredsson FP, Kemp CJ, Kuhn NC, Fleming SM, Egan AE, et

al. Morphological and behavioral impact of AAV2/5-mediated overexpression of human

wildtype alpha-synuclein in the rat nigrostriatal system. PloS one. 2013;8(11):e81426.

40. Miller JE, Hafzalla GW, Burkett ZD, Fox CM, White SA. Reduced vocal variability

in a zebra finch model of dopamine depletion: implications for Parkinson disease.

Physiol Rep. 2015;3(11).

41. Grant LM, Kelm-Nelson CA, Hilby BL, Blue KV, Paul Rajamanickam ES, Pultorak

JD, et al. Evidence for early and progressive ultrasonic vocalization and oromotor

deficits in a PINK1 gene knockout rat model of Parkinson's disease. J Neurosci Res.

2015;3:1713–27.

42. Yang KM, Blue KV, Mulholland HM, Kurup MP, Kelm-Nelson CA, Ciucci MR.

Characterization of oromotor and limb motor dysfunction in the DJ1-/-model of

Parkinson disease. Behavioural brain research. 2018;339:47-56.

43. Kelm-Nelson CA, Gammie S. Gene expression within the periaqueductal gray is

linked to vocal behavior and early-onset parkinsonism in Pink1 knockout rats. BMC

genomics. 2020;21:1-13.

44. Grant LM, Richter F, Miller JE, White SA, Fox CM, Zhu C, et al. Vocalization

deficits in mice over-expressing alpha-synuclein, a model of pre-manifest Parkinson's

disease. Behavioral neuroscience. 2014;128(2):110-21.

45. Medina CA, Vargas E, Munger SJ, Miller JE. Vocal changes in a zebra finch

model of Parkinson's disease characterized by alpha-synuclein overexpression in the

song-dedicated anterior forebrain pathway. PloS one. 2022;17(5):e0265604.

46. Bjork RT, Patel FZ, Daly MS, Miller JE. Evaluating alpha-synuclein proteinopathy

and consequences for birdsong in zebra finch basal ganglia Area X. Behavioural Brain

Research. 2025:115698.

47. Brainard MS, Doupe AJ. Translating birdsong: songbirds as a model for basic

and applied medical research. Annu Rev Neurosci. 2013;36:489-517.

48. Farries MA, Perkel DJ. Electrophysiological properties of avian basal ganglia

neurons recorded in vitro. Journal of neurophysiology. 2000;84(5):2502-13.

49. Person AL, Gale SD, Farries MA, Perkel DJ. Organization of the songbird basal

ganglia, including area X. J Comp Neurol. 2008;508(5):840-66.

50. Gale SD, Perkel DJ. Anatomy of a songbird basal ganglia circuit essential for

vocal learning and plasticity. Journal of chemical neuroanatomy. 2010;39(2):124-31.

51. Reiner A, Perkel DJ, Bruce L, Butler AB, Csillag A, Kuenzel W, et al. Revised

nomenclature for avian telencephalon and some related brainstem nuclei. *Journal of*

Comparative Neurology. 2004;473:377-414.

52. White SA. Genes and vocal learning. Brain Lang. 2010;115(1):21-8.

53. Simonyan K, Horwitz B, Jarvis ED. Dopamine regulation of human speech and

bird song: A critical review. Brain Lang. 2012;122(3):142-50.

54. Pfenning AR, Hara E, Whitney O, Rivas MV, Wang R, Roulhac PL, et al.

Convergent transcriptional specializations in the brains of humans and song-learning

birds. Science. 2014;346(6215):1256846.

55. Woolley SC, Kao MH. Variability in action: Contributions of a songbird cortical-

basal ganglia circuit to vocal motor learning and control. Neuroscience. 2015;296:39-47.

56. Lovell PV, Wirthlin M, Kaser T, Buckner AA, Carleton JB, Snider BR, et al.

ZEBrA: Zebra finch Expression Brain Atlas—A resource for comparative molecular

neuroanatomy and brain evolution studies. Journal of Comparative Neurology.

2020;528(12):2099-131.

57. Sakata JT, Birdsong D. Vocal Learning and Behaviors in Birds and Human

Bilinguals: Parallels, Divergences and Directions for Research. Languages. 2021;7(1):5.

58. Reiner A, Laverghetta AV, Meade CA, Cuthbertson SL, Bottjer SW. An

immunohistochemical and pathway tracing study of the striatopallidal organization of

area X in the male zebra finch. J Comp Neurol. 2004;469(2):239-61.

59. Farries MA, Perkel DJ. A telencephalic nucleus essential for song learning

contains neurons with physiological characteristics of both striatum and globus pallidus.

The Journal of neuroscience: the official journal of the Society for Neuroscience.

2002;22(9):3776-87.

60. Coleman MJ, Roy A, Wild JM, Mooney R. Thalamic gating of auditory responses

in telencephalic song control nuclei. Journal of Neuroscience. 2007;27(37):10024-36.

61. Peh WY, Roberts TF, Mooney R. Imaging auditory representations of song and

syllables in populations of sensorimotor neurons essential to vocal communication.

Journal of Neuroscience. 2015;35(14):5589-605.

62. Woolley SC, Rajan R, Joshua M, Doupe AJ. Emergence of context-dependent

variability across a basal ganglia network. Neuron. 2014;82(1):208-23.

63. Farries MA, Ding L, Perkel DJ. Evidence for "direct" and "indirect" pathways

through the song system basal ganglia. J Comp Neurol. 2005;484(1):93-104.

64. Singh A, Mewes K, Gross RE, DeLong MR, Obeso JA, Papa SM. Human striatal

recordings reveal abnormal discharge of projection neurons in Parkinson's disease.

Proceedings of the National Academy of Sciences. 2016;113(34):9629-34.

65. Kojima S, Kao MH, Doupe AJ. Task-related "cortical" bursting depends critically

on basal ganglia input and is linked to vocal plasticity. Proceedings of the National

Academy of Sciences of the United States of America. 2013;110(12):4756-61.

66.   Moll FW, Kranz D, Corredera Asensio A, Elmaleh M, Ackert-Smith LA, Long MA.

Thalamus drives vocal onsets in the zebra finch courtship song. Nature. 2023:1-5.

67.   Tchernichovski O, Nottebohm F, Ho CE, Pesaran B, Mitra PP. A procedure for

an automated measurement of song similarity. Animal Behaviour. 2000;59:1167-76.

68.   Badwal A, Poertner J, Samlan RA, Miller JE. Common terminology and acoustic

measures for human voice and birdsong. Journal of Speech, Language, and Hearing

Research. 2018;61(12).

69.   Badwal A, Borgstrom M, Samlan RA, Miller JE. Middle age, a key time point for

changes in birdsong and human voice. Behavioral neuroscience. 2020;134(3):208.

70.   Sober SJ, Wohlgemuth MJ, Brainard MS. Central contributions to acoustic

variation in birdsong. The Journal of neuroscience: the official journal of the Society for

Neuroscience. 2008;28(41):10370-9.

71.   Pachitariu M, Steinmetz NA, Kadir SN, Carandini M, Harris KD. Fast and

accurate spike sorting of high-channel count probes with KiloSort. Advances in neural

information processing systems. 2016;29.

72.   Shinomoto S, Kim H, Shimokawa T, Matsuno N, Funahashi S, Shima K, et al.

Relating neuronal firing patterns to functional differentiation of cerebral cortex. PLoS

computational biology. 2009;5(7):e1000433.

73.   Buzsáki G, Mizuseki K. The log-dynamic brain: how skewed distributions affect

network operations. Nature reviews neuroscience. 2014;15(4):264-78.

74.   Goldberg JH, Fee MS. Singing-related neural activity distinguishes four classes

of putative striatal neurons in the songbird basal ganglia. Journal of neurophysiology.

2010;103:2002-14.

75.   Berke JD, Okatan M, Skurski J, Eichenbaum HB. Oscillatory entrainment of

striatal neurons in freely moving rats. Neuron. 2004;43(6):883-96.

76.   Wilson CJ. The generation of natural firing patterns in neostriatal neurons.

Progress in brain research. 1993;99:277-97.

77.   Budzillo A, Duffy A, Miller KE, Fairhall AL, Perkel DJ. Dopaminergic modulation

of basal ganglia output through coupled excitation-inhibition. Proceedings of the

National Academy of Sciences of the United States of America. 2017;114(22):5713-8.

78.   Goldberg JH, Adler A, Bergman H, Fee MS. Singing-related neural activity

distinguishes two putative pallidal cell types in the songbird basal ganglia: comparison

to the primate internal and external pallidal segments. The Journal of neuroscience: the

official journal of the Society for Neuroscience. 2010;30(20):7088-98.

79.   Doupe AJ, Solis MM, Kimpo R, Boettiger CA. Cellular, circuit, and synaptic

mechanisms in song learning. Annals of the New York Academy of Sciences.

2004;1016(1):495-523.

80.   Barthó P, Hirase H, Monconduit L, Zugaro M, Harris KD, Buzsaki G.

Characterization of neocortical principal cells and interneurons by network interactions

and extracellular features. Journal of neurophysiology. 2004;92(1):600-8.

81.   Shinomoto S, Shima K, Tanji J. Differences in spiking patterns among cortical

neurons. Neural computation. 2003;15(12):2823-42.

82.   Bugaysen J, Bronfeld M, Tischler H, Bar-Gad I, Korngreen A.

Electrophysiological characteristics of globus pallidus neurons. PloS one.

2010;5(8):e12001.

83.   Chen H, Lei H, Xu Q. Neuronal activity pattern defects in the striatum in awake

mouse model of Parkinson's disease. Behavioural brain research. 2018;341:135-45.

84.   Singh A, Liang L, Kaneoke Y, Cao X, Papa SM. Dopamine regulates distinctively

the activity patterns of striatal output neurons in advanced parkinsonian primates.

Journal of neurophysiology. 2015;113(5):1533-44.

85.   Vishwanath A, Bartlett MJ, Falk T, Cowen SL. Decoupling of motor cortex to

movement in Parkinson's dyskinesia rescued by sub-anaesthetic ketamine. Brain.

2025;148(6):2135-50.

86.   Gold C, Henze DA, Koch C, Buzsaki G. On the origin of the extracellular action

potential waveform: a modeling study. Journal of neurophysiology. 2006;95(5):3113-28.

87.   Teleńczuk M, Brette R, Destexhe A, Teleńczuk B. Contribution of the axon initial

segment to action potentials recorded extracellularly. Eneuro. 2018;5(3).

88.   Bakkum DJ, Obien MEJ, Radivojevic M, Jäckel D, Frey U, Takahashi H, et al.

The axon initial segment is the dominant contributor to the neuron's extracellular

electrical potential landscape. Advanced biosystems. 2019;3(2):1800308.

89.   Zhao W, Zhang M, Liu J, Liang P, Wang R, Hemmings HC, et al. Isoflurane

modulates hippocampal CA1 pyramidal neuron excitability by inhibition of both transient

and persistent sodium currents in mice. Anesthesiology. 2019;131(1):94.

90.   Ries CR, Puil E. Isoflurane prevents transitions to tonic and burst firing modes in

thalamic neurons. Neuroscience letters. 1993;159(1-2):91-4.

91.   Dejean C, Gross CE, Bioulac B, Boraud T. Dynamic changes in the cortex-basal

ganglia network after dopamine depletion in the rat. Journal of neurophysiology.

2008;100(1):385-96.

92.   Choudhury SP, Bano S, Sen S, Suchal K, Kumar S, Nikolajeff F, et al. Altered

neural cell junctions and ion-channels leading to disrupted neuron communication in

Parkinson's disease. npj Parkinson's Disease. 2022;8(1):66.

93.   Deister CA, Chan CS, Surmeier DJ, Wilson CJ. Calcium-activated SK channels

influence voltage-gated ion channels to determine the precision of firing in globus

pallidus neurons. Journal of Neuroscience. 2009;29(26):8452-61.

94.   Faber EL, Sah P. Physiological role of calcium-activated potassium currents in

the rat lateral amygdala. Journal of Neuroscience. 2002;22(5):1618-28.

95.   Robinson RB, Siegelbaum SA. Hyperpolarization-activated cation currents: from

molecules to physiological function. Annual review of physiology. 2003;65(1):453-80.

96.   Storm JF. Potassium currents in hippocampal pyramidal cells. Progress in brain

research. 1990;83:161-87.

97.   Hill E, Gowers R, Richardson M, Wall M. α-Synuclein aggregates increase the

conductance of substantia nigra dopamine neurons, an effect partly reversed by the

KATP channel inhibitor glibenclamide. eneuro. 2021;8(1).

98.   Singh Alvarado J, Goffinet J, Michael V, Liberti W, Hatfield J, Gardner T, et al.

Neural dynamics underlying birdsong practice and performance. Nature. 2021:1-5.

99.   Williams H, Crane LA, Hale TK, Esposito MA, Nottebohm F. Right-side

dominance for song control in the zebra finch. Journal of Neurobiology.

1992;23(8):1006-20.

100.   Frank SY, Hunt JL, Bae AJ, Chirathivat N, Lotfi S, Raja SC, et al. Hemispheric

dominance in HVC is experience-dependent in juvenile male zebra finches. Scientific

reports. 2024;14(1):5781.

101. Plowman-Prine E, Okun M, Sapienza C, Shrivastav R, Fernandez H, Foote K, et

al. Perceptual characteristics of Parkinsonian speech: a comparison of the

pharmacological effects of levodopa across speech and non-speech motor systems.

NeuroRehabilitation. 2009;24(2):131-44.

102. Skodda S. Effect of deep brain stimulation on speech performance in Parkinson's

disease. Parkinson's Disease. 2012;2012.

103. Fabbri M, Guimaraes I, Cardoso R, Coelho M, Guedes LC, Rosa MM, et al.

Speech and voice response to a levodopa challenge in late-stage Parkinson's Disease.

Front Neurol. 2017;8:432.

Supporting Information

S1 Table 1. Song data and statistics. For further information, see legends embedded

into the tabs of this excel file.

S1 Fig. Exemplar song motif and group level comparisons of song syllable data.

A) Spectrogram of song motif from a pre-injected GFP control bird (as in **Fig 2A-C**) with

time in seconds on the x-axis and frequency in kilo (k) Hertz on the y-axis. Unique

syllables are assigned a letter (A-E) and classified into harmonic, noisy, or mixed

syllables (68). **B-E)** The mean and SEM scores shown for syllable level acoustic

features (duration, amplitude, and entropy) for the ASYN (red) and GFP (black) control

groups. Raw scores can be found in the **S1 Table**. The y-axis is the normalized score

calculated from dividing the post-AAV injection scores by the pre-injection scores. Each

dot represents an average score per finch. **B)** Comparisons of all syllable types

combined or by individual syllable types **(C-E)** reveal no group differences (Mann

Whitney U, p>0.05). Individual variation in CV scores is detected.

S2 Fig. No significant correlation between song features and firing rate. Scatter

plots showing the mean firing rate of WLR neurons averaged for each animal (dot)

organized by GFP (black) and ASYN (red) groups. Plots were generated for each song

feature, representing all syllable types combined. Generalized linear regression (see

Results) did not identify a relationship between firing rate and song features or group (p

= 0.0815).

A

Zebra Finch

Glutamate
GABA
Dopamine

HVC
IMAN
X
RA
DLM
Song

Cx
BG
Th
MB
HB
Cb
OB

B

C

5' ITR CBA eGFP WPRE ITR 3'

OR

5' ITR CBA SNCA WPRE ITR 3'

rAAV5

in vivo

Area X

Day 1-3
Birdsong
recording

Day 3
Virus injections

2 Months
Birdsong
recording

≤ 5 days
Anesthetized
electrophysiological recordings