A morphometric approach to the taxonomic dilemma of *Zonozoe drabowiensis* Barrande, 1872

and *Zonoscutum solum* Chlupáč, 1999 (Upper Ordovician, Czech Republic)

Lorenzo Lustri[1, 2]*, Lukáš Laibl[3], Luis Collantes[1,2], Jana Bruthansová[4], Martina Nohejlová[5], Yu

Liu[1, 2] and Stephen Pates[6 7]

[1]Yunnan Key Laboratory for Palaeobiology, Institute of Palaeontology, Yunnan University,

650091, Kunming, China

[2]MEC International Joint Laboratory for Palaeobiology and Palaeoenvironment, Yunnan

University, 650500 Kunming, China

[3]Czech Academy of Sciences, Institute of Geology, Rozvojová 269, 165 00, Prague 6, Czech

Republic

[4]Department of Palaeontology, National Museum, Cirkusová 1740, 193 00, Prague 9, Czech

Republic

[5]Czech Geological Survey, Klárov 3, 11800, Prague, Czech Republic

[6]Department of Earth Sciences, University College London, Gower Street, London, WC1E 6BT,

United Kingdom

[7]Department of Genetics, Evolution and Environment, University College London, Gower Street,

London, WC1E 6BT, United Kingdom

* Corresponding author

ORCID

LLu, 0000-0002-8951-5160; LLa 0000-0001-9049-3811; LC 0000-0001-9571-4433; JB 0000-

0002-6218-2255; MN 0000-0001-7972-8165; YL 0000-0002-2346-740X; SP 0000-0001-8063-

9469

ABSTRACT

Zonozoe drabowiensis Barrande, 1872 and *Zonoscutum solum* Chlupáč, 1999 are rare and

incompletely preserved arthropods from the Upper Ordovician of the Czech Republic. Their

classification has been a subject of debate for over a century due to the limited number of

specimens and the absence of clear diagnostic features. They were previously thought to belong to

Aglaspidida, an extinct group of arthropods from the Cambrian and Ordovician, within

Vicissicaudata, a branch of the larger arthropod clade Artiopoda that also includes trilobites. Herein,

we used analyses of cephalic outline to quantify whether *Zonozoe* and *Zonoscutum* are

morphologically more similar to vicissicaudates than other early Palaeozoic arthropods. We

assembled a dataset of twenty-six cephalic outlines representing twenty-six early Palaeozoic

species, including *Zonozoe*, *Zonoscutum*, five euchelicerates, six aglaspidids, three cheloniellids,

and a selection of other artiopodans. We quantified their shape using elliptical Fourier analysis.

Principal component analysis (PCA) and linear discriminant analysis (LDA) place *Zonozoe* and

Zonoscutum within and close to the vicissicaudatan morphospace, clearly distinguishing them from

euchelicerates and non-vicissicaudatan artiopodans. These data support the most conservative

classification of *Zonozoe* and *Zonoscutum* within Artiopoda, while strengthening the case for a more

specific affinity with Vicissicaudata, helping to resolve a 150-year-old taxonomic uncertainty. More

broadly, this study demonstrates the value of outline-based morphometrics in testing systematic

hypotheses when discrete characters are unavailable or scarce, offering a reproducible tool for re-

evaluating other problematic fossils.

Key words: *Zonozoe drabowiensis*, *Zonoscutum solum,* elliptical Fourier analysis, Vicissicaudata,

Artiopoda

47   Key words: *Zonozoe drabowiensis*, *Zonoscutum solum,* elliptical Fourier analysis, Vicissicaudata,

48   Artiopoda

INTRODUCTION

Zonozoe drabowiensis Barrande, 1872 and *Zonoscutum solum* are enigmatic arthropods known from the Letná and the Libeň Formations of the upper Ordovician of the Czech Republic (Chlupáč, 1965, 1999b; Rak et al., 2009; Vokáč & Grigar, 2024). Barrande (1872) originally described *Zonozoe* as an arthropod, specifically as a large carapace of an ostracod (Barrande, 1872). Later on, Novák in his unpublished manuscript (Novák, 1887) considered *Zonozoe* to be a merostomate arthropod close to Synziphosurina (Chlupáč, 1963, 1965, 1999b; Rak, 2009) . Since then, these fossils have not received much attention until the second half of the twentieth century. Chlupáč, 1963 assigned *Zonozoe* into the group Aglaspidida (Walcott, 1911) – a group of extinct marine arthropods, known from the Cambrian and Ordovician strata. However, this assignment remains uncertain (Chlupáč, 1963). It should also be noted that at that time Aglaspidida was considered as one of the basal-most groups within Xiphosura. Subsequent studies have revisited the taxon, reiterating its general association with Aglaspidida and/or the Merostomata, but always with a low degree of confidence (Chlupáč, 1965; Hou, 1997; Chlupáč, 1999a; Rak et al., 2009). As for *Zonoscutum*, the only fossil ascribed to this species was described in Chlupáč (1999a), who tentatively assigned it to Aglaspidida.

During the one hundred and fifty years since the discovery of *Zonozoe*, and twenty-five years for *Zonoscutum,* their classification has been complicated by two main factors. Firstly, the material ascribed to both *Zonozoe* and *Zonoscutum* is scarce, incomplete and not well preserved (Ortega Hernández et al., 2013; Van Roy et al., 2022). There are currently only eight known specimens of *Zonozoe* and one of *Zonoscutum,* all of which are isolated moulds of cephala (Chlupáč, 1999b; Rak et al., 2009; Vokáč & Grigar, 2024), preserved in quartzose sandstones of the Libeň and Letná Formations. In most cases, the specimens preserve only the general vaulted cephalon outline with a raised longitudinal region terminating with a ridge bearing what is a vague indication of the eyes (Rak et al., 2009; Van Roy et al., 2022). The only exception is a single specimen of *Zonozoe* preserving what has been referred to as glabellar furrows (Rak et al., 2009)

and a specimen of *Zonoscutum* preserving similar structures but in a different position (Chlupáč,

1999a). Secondly, the classification of *Zonozoe* and *Zonoscutum* has been further complicated by

the complex history of the classification of aglaspidid arthropods themselves, especially during the

second half of the twentieth century. Their overall dorsal morphology led several authors to

consider the aglaspidids as an order of the subclass Xiphosura (Størmer, 1955), grouping

Aglaspidida with Xiphosura and Eurypterida, as part of the Merostomata (Raasch, 1939; Størmer,

1944; Eldredge & Smith, 1974) or to consider Aglaspidida as the sister group to chelicerates

(Weygoldt and Paulus, 1979). This is in contrast to more recent aglaspidid phylogenies, which

emphasize ventral anatomical features as key to understanding their true affinities (Briggs et al.,

1979; Hesselbo, 1992; Ortega☐Hernández et al., 2013; Lerosey-Aubril, Paterson, et al., 2017;

Lerosey-Aubril, Zhu, et al., 2017). Currently, they have been placed within the broader clade

Artiopoda, as part of the clade Vicissicaudata (Ortega☐Hernández et al., 2013; Lerosey-Aubril,

Paterson, et al., 2017; Lerosey-Aubril, Zhu, et al., 2017). Until very recently, the absence of strong

defining characteristics for the clade has made Aglaspidida prone to being used as a wastebasket

taxon (Plotnick & Wagner, 2006). This could be the case for the poorly preserved *Zonozoe* and

Zonoscutum, which lack strong anatomical evidence supporting placement within aglaspidids. Other

factors that led previous workers to assign *Zonozoe* and *Zonoscutum* to Aglaspidida are likely not

only the general similarity between their cephala, but also the known (or suspected) presence of

aglaspidids in the same paleogeographic region (Chlupáč, 1963, 1965; Van Roy, 2005; Ortega-

Hernandez et al., 2010) and the weak resemblance with other arthropod groups (e.g. position of the

eyes and glabella if compared with trilobites).

Later authors commenting on *Zonozoe* and *Zonoscutum* decided to have a more parsimonious

approach. (Ortega☐Hernández et al. 2013) opted for the exclusion of those taxa from an extensive

phylogenetic analysis of Vicissicaudata due to the lack of compelling taxonomically relevant

features. (Van Roy et al., 2022), instead, revised *Zonozoe* and *Zonoscutum* to critically assess their

possible affinity with *Triopus draboviensis* (Barrande, 1872). In their re-examination of the

material, the authors did more than simply dismiss the affinity with *Triopus*; they advanced a

conservative and more parsimonious taxonomic classification for both *Zonozoe* and *Zonoscutum*.

They suggested that, at most, the fossils could be broadly classified as Artiopoda, and argued that

any attempt to refine the classification beyond that level would be unwarranted (Van Roy et al.,

2022). Known only from cephalic moulds, both *Zonozoe* and *Zonoscotum* lack the diagnostic

features of Vicissicaudata, all of which are post-cephalic (Lerosey-Aubril, Zhu, et al., 2017; Briggs

et al., 2023). They also lack several defining traits of Aglaspidida, including all post-cephalic

diagnostic characters, most notably the modified posterior appendages, as well as evidence of a

biomineralized cuticle and dorsal ecdysial sutures (Ortega☐Hernández et al., 2013; Lerosey-Aubril,

Zhu, et al., 2017). As a result, they cannot be confidently classified through standard phylogenetic

or comparative analysis (Ortega☐Hernández et al., 2013; Van Roy et al., 2022), a conclusion fully

supported by this study. While the enigmatic *Zonozoe* and *Zonoscutum* are unlikely to be key taxa

in contributing to resolving the phylogeny of Aglaspidida, a more accurate assignment of this

species to this group may yield valuable palaeoecological and paleobiogeographical insights.

In order to resolve the taxonomic dilemma surrounding *Zonozoe* and *Zonoscutum* more effectively,

we adopted an alternative approach to phylogenetic analysis and classical comparative anatomy,

focusing instead on the only reliably preserved feature in these fossils: the overall cephalic outline.

We compared the outline of the cephalon of *Zonozoe* and *Zonoscutum* with twenty-four other

arthropods. We implemented a principal component analysis (PCA) and linear discriminant analysis

(LDA) on the study of the cephalic outline to extract the maximum morphological information from

the limited material of *Zonozoe* and *Zonoscutum*. This methodology has been used in paleobiology

to compare the outline of different fossils and can provide information on their taxonomy, ecology,

and development (Gevirtz, 1976; Healy-Williams & Williams, 1981; Burke et al., 1987; Foote,

1989; Pates et al., 2021; Ma et al., 2023; Braig et al., 2024; Drage & Pates, 2024; Collantes & Pates,

in press). A significant advantage of this methodology over landmark-based approaches is that it

does not rely on discrete morphological characters, facilitating quantification of shape variation

when such discrete characters are absent. Ultimately, our analysis provides a novel line of evidence

toward supporting the systematic placement of problematic species. Both *Zonozoe* and *Zonoscutum*

plotted within the Vicissicaudata area of morphospace, and thus the shape of their cephala concurs

with a vicissicaudate affinity.

MATERIALS AND METHODS

Species involved in the study and the criteria of selection

In addition to one *Zonozoe* and one *Zonoscutum* specimen (Figure 1), five euchelicerates, five non-

vicissicaudatans artiopodans and fourteen vicissicaudatans arthropods (including six aglaspidids

and three cheloinellids), representing twenty-six species, were included in our study. The most

complete and horizontally preserved specimen of *Zonozoe* (Lectotype NM L 23586) was chosen,

while *Zonoscutum* is known from only one specimen (Holotype NM L 33021). The selection from

the literature of the other taxa, was made following three main criteria: 1) to include Early

Palaeozoic diversity from various geographic regions, but with particular emphasis on species

coeval and sympatric to *Zonozoe* and *Zonoscutum* (e.g. *Paraeurypterus anatoliensis* (Lamsdell et

al., 2013)); 2) to exclude taxa with aberrant cephalic outlines and favour those potentially related to

Zonozoe and *Zonoscutum* (e.g. *Eozetetes gemmelli* (Edgecombe et al., 2017) and *Chlupacaris dubia*

(Van Roy, 2005)); and 3) to maximize higher-level taxonomic diversity (e.g. *Carimersa neptuni*

(Briggs et al., 2023) and *Lunataspis borealis* (Lamsdell et al., 2022)). A summary of the species and

specimens included in the analysis, along with inventory numbers, geological age, geographic

provenance, and references to the original sources from which raw images were obtained, is

provided in Table 1, while the specimen sample, one for each high taxonomic group, is shown in

Figure 1.

Material and photography

The studied material of *Zonozoe* and *Zonoscutum* is housed in the palaeontological collections of

the National Museum, Prague (prefix NM L). The photographs of *Zonozoe* were taken using a Sony

ILCE-7RM5 digital camera with associated DT 100mm F2.8 SAM lens under normal lighting

conditions, using a stand and a sandbox to hold the sample in place and ensure the horizontality of

both the camera and the specimen. Specimens were coated by ammonium chloride prior

photography to enhance morphological details.

Geological settings of Zonozoe *and* Zonoscutum

The examined specimens of *Zonozoe* and *Zonoscutum* (Figure 1) come from the Ordovician

deposits of the Prague Basin (Czech Republic, Barrandian area). All *Zonozoe* and *Zonoscutum*

specimens represent internal moulds preserved as imprints in coarse quartzose sandstones, showing

limited anatomical details. The majority of them were collected at the at Děd Hill near Beroun (five

specimens) or in the fields between Trubín and Trubská villages near Králův Dvůr (one specimen,

see (Drage et al., 2018) for details of the location), in the upper part of the Letná Formation

(Chlupáč, 1965). One specimen (NM L 37385) was recovered from the Řevnice Quartzite Member

within the Libeň Formation near Kařez (Rak et al., 2009). (Vokáč & Grigar, 2024) recently reported

additional cephala of *Zonozoe* from the Řevnice Quartzite Member (Libeň Formation) near Starý

Plzenec. The only known cephala of *Zonoscutum* comes also from the Letná Formation within the

Barrandian area (Chlupáč, 1999a). It has been discovered in the fields between Trubín and Trubská

Villages near Králův Dvůr.

The Libeň Formation is developed in three different members – the shallow water R□evnice

Quartzite, the sequence of black shales deposited deeper sea below the wave base in anoxic

conditions, and the pyroclastic material (Havlíček, 1998). The Letná Formation is generally

composed of alternating quartzose sandstones, greywackes, siltstones, and occasional shales, with

sandstones representing a shallow-water environment (Chlupác & Kukal, 1988; Kraft et al., 2023).

Zonozoe and *Zonoscutum* have so far been discovered only in the shallow water facies of Libeň and

Letná Formations. Both Letná and Libeň Formations are of Sandbian age (Upper Ordovician) (Kraft

et al., 2023). From the palaeogeographical perspective, these formations within the Prague Basin

were deposited in high-latitude area near the western margin of Gondwana (Kraft et al., 2023;

Scotese, 2023).

Silhouette drawing, Principal component analysis (PCA), linear discriminant analysis (LDA)

We extracted silhouette drawings from photographs of selected specimens, either taken by the

authors in museum collections or sourced from the literature (see Table 1 and Figure 1). Using

Adobe Illustrator 2021, we manually traced each specimen's outline over the photograph, then

exported the silhouette as a JPG file (Figure 3A). These image files served as input for the Momocs

package in R (Bonhomme et al., 2014; RCoreTeam, 2024), which we used to perform the

subsequent outline analyses. JPG files were imported and converted to outlines, scaled, centred, and

sampled at 64 point resolution, and subjected to elliptical Fourier analysis. The function

calibrate_harmonicpower_efourier was used to determine the number of harmonics required for

99.9% of the harmonic power. Elliptical Fourier coefficients were visualised using Principal

Component Analysis. Linear Discriminant Analysis was used to model differences between the

major groupings in the dataset (Artiopoda (non-Vicissicaudata), Euchelicerata, Vicissicaudata,

Zonozoe, Zonoscutum).

RESULTS

EFA, Principal component analysis (PCA), linear discriminant analysis (LDA)

Fifteen harmonics were retained, providing 99.9% of the harmonic power for the dataset. Principal

component analysis reduced the dimensionality of elliptical Fourier analysis results, with PC1 and

PC2 explaining 88.3% of the variation (67.4% and 20.9% respectively). Higher principal

components (PC3–PC26) explain <5% of the variation each, and <15% overall. Three slightly

overlapping groups are observed in the PCA space: Euchelicerata, Vicissicaudata, and Artiopoda

(not including Vicissicaudata) (Figure 3B). Euchelicerates cluster toward the lower left (negative

PC1, with PC2 values from close to zero to negative), while non☐Vicissicaudatan artiopodans

occupy the area of the space with PC2 close to zero and positive PC1 values. Within Vicissicaudata,

aglaspidids fall near the origin of morphospace (PC1, PC2 close to zero), except for *Tremaglaspis*

unite (Fortey & Rushton, 2003), which plots with a negative PC1 value (Figures☐3B and C).

Cheloniellids form a tight cluster with positive PC2 values. Both *Zonozoe* and *Zonoscutum* lie

within the upper region of the vicissicaudatan space. *Zonozoe* plots between the cheloniellid

Neostrabops martini (Caster & Macke, 1952) and the aglaspidid *Aglaspis barrandei* (Hall,

1862)(Figures☐3B and C), whereas *Zonoscutum* lies close to the cheloniellids *Triopus* and

Drabovaspis complexa (Barrande, 1872)(Figures☐3B and C).

Linear discriminant analysis is dominated by LD1 (83.1% of the trace), with LD2 (9.9%), LD3

(5.2%) and LD4 (1.9%) providing only a relatively small amount of information. Our LDA

(Figure☐3D) separates Euchelicerata from all other taxa, but cannot separate vicissicaudates from

non-vicissicaudate artiopodans (Table 2). Leave-one-out cross-validation was 45.7% (11/24).

Non☐Vicissicaudata artiopodans fall into the vicissicaudatans space while both *Zonozoe* and

Zonoscutum plot at the upper part of the LDA space (Figure 3D).

DISCUSSION

Our outline analyses provide morphological support for an Artiopoda, most likely Vicissicaudata,

affinity for both *Zonozoe and Zonoscutum*. This provides the first quantitative evidence placing

these enigmatic taxa not only within Artiopoda, previously inferred through taxonomically weak

morphological traits, but more specifically within Vicissicaudata. However, the PCA does not

explicitly support an affinity with Aglaspidida, particularly in the case of *Zonoscutum*. Instead, both

taxa are placed in an intermediate position between cheloniellids and aglaspidids (Figures☐3B and

C).

Although our analysis places *Zonozoe and Zonoscutum* near the cheloniellid cluster, we interpret

this proximity cautiously and do not suggest it as evidence of a closer affinity with cheloniellids

than with other vicissicaudatans. Key differences in cephalic morphology, which are most apparent

in three-dimensional views, are not fully captured by our two-dimensional dataset, and both

Zonoscutum and cheloniellids follow in this case. They show, respectively, a highly vaulted and

extremely flattened cephalon, which may have been more affected by this methodological

constraint, respectively to what was retrieved for *Zonozoe* and the other involved taxa. We believe

this is the explanation behind the results of our PCA, concerning *Zonoscutum* clustering closer to

the well-defined cheloniellids *Triopus* and *Drabovaspis* than to any other taxa. Based on this, we

consider our method less effective for taxa with extremely vaulted or flattened cephala, such as

cheloniellids and *Zonoscutum*. This limitation might have been addressed by using 3D volume data,

but such data are difficult to obtain for fossils held in collections across multiple continents, and

will also be impacted by flattening during or after fossilisation. Thus, we broadly agree with (Van

Roy et al., 2022), who rejected a close relationship between *Zonozoe*, *Zonoscutum*, and

cheloniellids, especially with the coeval and sympatric *Triopus*.

The closest non-cheloniellid vicissicaudatan to *Zonoscutum,* on the other hand, is *Chlupacaris*,

which we regard as a more likely relative, also due to both general morphological similarities and

geographic context. Both *Zonoscutum* and *Chlupacaris* exhibit a highly vaulted cephalon, again, a

feature not fully accounted for in outline-based 2D morphometric methods, yet they have been

recovered closely. Moreover, during the Upper Ordovician, the Prague Basin was situated near the

western margin of Gondwana (Kraft et al., 2023; Scotese, 2023) and, thus, shared a close position to

Morocco from a paleogeographic point of view. For *Zonozoe*, the closest taxa in our PCA are the

cheloniellid *Neostrabops* and the aglaspidid *A. barrandei*. Ortega-Hernández et al. (2013) assigned

Neostrabops as the less derived Cheloniellida, on the basis of a shared "presence of rounded genal

angles, the anterior boundaries of the trunk reflexed anterolaterally and a reduced head shield".

However, the absence of a well-developed thoracic axis, already led Van Roy (2006) to suggest it

may represent the less derived form of the group. Other studies (e.g., Legg et al., 2013; Lerosey-

Aubril, Zhu, et al., 2017) also placed *Neostrabops* as a basal cheloniellid, but again relied on limited

synapomorphies. The lack of preservation of key traits, such as furcal rami or the morphology of

terminal tergites, hinders reliable comparison with better-known forms like *Triopus* or *Cheloniellon*

calmani (Broili, 1932). The proximity of *Zonozoe* and *Neostrabops* in our PCA may therefore

reflect more on the uncertain taxonomy of *Neostrabops* than on *Zonozoe* itself. Both fall within the

vicissicaudatan morphospace, closer to aglaspidids than to other cheloniellids. Based on our results,

we suggest that the taxonomical assignment of *Neostrabops* could be reconsidered towards

Aglaspidida after new investigations or, in a more parsimonious way, towards Xenopoda.

On the other hand, the close PCA placement of *Zonozoe* (and *Neostrabops*) with *A. barrandei*

supports a possible affinity with aglaspidids. While we do not draw a definitive conclusion, this

result is consistent with previous observations, such as the position of the eyes and glabellar

furrows behind them, which, while not strongly diagnostic (Van Roy et al., 2022), align with our

morphometric findings and the exclusion of a possible euchelicerate affinity retrieved in our

analyses (Fig. 3).

Our LDA results, though more influenced by predefined categories, provide additional insight.

Euchelicerates are distinguished from Artiopoda, however vicissicaudatans and non-

vicissicaudatans cannot be clearly separated (Figure 3D; Table 2). This was an unexpected finding

for our outline data, given the superficial similarity in dorsal morphology between some early

euchelicerates and vicissicaudatans like aglaspidids (e.g. *P. anatoliensis* and *T. unite*). Previous

authors have even suggested the affinities of euchelicerates with aglaspidids based on such

similarities, but our LDA analyses were able to discriminate the two groups. Both taxa of interest,

Zonoscutum and *Zonozoe,* are positioned closer to vicissicaudatans than to non-vicissicaudatan

artiopods, and are clearly separated from Euchelicerata. These results reinforce the more recent

interpretation of aglaspidids, and by extension *Zonozoe* and *Zonoscutum*, as artiopodans, most

likely vicissicaudatans.

CONCLUSION

Our outline□based morphometric analyses provide quantitative support for placing *Zonozoe* and

Zonoscutum within Artiopoda, most likely as Vicissicaudata. Although they plot between

cheloniellids and aglaspidids in the morphospace of two-dimensional carapace outlines, reflecting

methodological limits when three□dimensional cephalic shape is compressed, both PCA and LDA

consistently separate them from euchelicerates. Within PCA space, they plot closest to

vicissicaudatan artiopods. Taken together, these results support a vicissicaudatan affinity for these

taxa. Overall, our method has proven to be a useful alternative for classifying arthropod fossils

when discrete characters required for phylogenetic methods are absent. This is often the case with

moulds, which rarely preserve fine anatomical details. While our approach is less precise than

formal phylogenetic analysis, it allows for the inclusion of specimens that would otherwise be

excluded from broader evolutionary or paleoecological studies. In our case, it provides evidence for

the presence of non-cheloniellid vicissicaudatans in the Letná Formation, based on poorly preserved

specimens that would likely be overlooked using only traditional qualitative methods.

ACKNOWLEDGMENTS

We are thankful to the reviewers for their valuable comments and suggestions, which greatly

improved this manuscript. This study was supported by the grant "Borse di studio SPI" from

"Società Paleontologica Italiana" (SPI) awarded to L. Lu. in 2024. This study was also supported by

the grants of PostDoctoral Research Fund to L.Lu. and L.C at Yunnan University. Y.L. is currently

supported by the Yunnan Revitalization Talent Support Program (Grant number:

202401BC070012) from the Natural Science Foundation of Yunnan Province. SP acknowledges

support from a NERC IRF NE/X017745/1. L.La is supported by the Czech Science Foundation (25-

16610S) and by the institutional support RVO 67985831 of the Institute of Geology of the Czech

Academy of Sciences. J.B. was financially supported by Ministry of Culture of the Czech Republic

through the DKRVO 2024–2028/2.III.b, National Museum, 00023272. M.N. was supported by

internal grant of the Czech Geological Survey no. 311 630, which contributes to the Strategic

Research Plan of the Czech Geological Survey (DKRVO/CGS 2023-2027). This paper is a

contribution to IGCP project 735 "Rocks and the Rise of Ordovician Life: Filling knowledge gaps

in the Early Palaeozoic Biodiversification.

AUTHOR CONTRIBUTIONS

L.Lu. conceptualized the original idea and wrote the first draft of the manuscript. L.Lu., L.La. and

L.C. collected the data. S.P. analysed the data. L.La. and L.C., and S.P. prepared the figures and

wrote sections of the material and methods, results, and discussion. Y.L., J.B. and M.N. were

involved in the discussion and final editing of the manuscript. All authors read and approved the

final version of the manuscript.

Figure 1. Representative cephala of *Zonozoe drabowiensis*, *Zonoscutum solum*, euchelicerate arthropods, vicissicaudatan cheloniellids, vicissicaudatan aglaspidids, and non-vicissicaudatan artiopodans featured in this study. A1, *Zonozoe drabowiensis*, dorsal view; A2, *Zonozoe drabowiensis*, frontal view; B, *Zonoscutum solum*, dorsal view, modified from Van Roy et al. (2022); C, unnamed Xiphosura from the Fezouata biota, modified from Lamsdell & Ocon (2025). D, *Eozetetes gemmelli*, modified from (Edgecombe et al., 2017); E, *Neostrabops martini*, modified from (Caster & Macke, 1952); F, *Aglaspis barrandei*, modified from (Hesselbo, 1992). Scale bar equals 5 mm.

322

323 **Figure 2. geographical and stratigraphical position of the localities where *Zonozoe***

324 ***drabowiensis* and *Zonoscutum solum* were collected.** A; map of the Czech republic, B;

325 stratigraphical column of the upper Ordovician (modify after (Van Roy et al., 2022)).

Figure 3. Data and Results of Principal Component Analysis and Linear Discriminant

Analysis of elliptical Fourier coefficients. A, Cephalic outline of *Zonozoe drabowiensis*,

Zonoscutum solum and the other specimens involved in the study; B, position of *Zonoscutum* and

Zonozoe in relation to major taxonomic groupings: Euchelicerata, Artiopoda (not including

Vicissicaudata), and Vicissicaudata. Grey convex hulls show subgroups within Vicissicaudata,

specifically Aglaspidida and Cheloniellida. Coloured marks along x and y axis indicate PC1 and

PC2 values of individual specimens plotted in the space, with positions of specimens shown by

filled circles. Grey outlines show variation in cephalic outline across PC1 and PC2. Histogram in

bottom right shows amount of variation explained by PC1, PC2, and PC3-5; C, results by individual

outline of PCA analyses. Abbreviation 'unnamedfez' refers to an unnamed Xiphosura from the

Fezouata biota (Van Roy et al., 2010; Lamsdell & Ocon, 2025); D, Position of *Zonoscutum* and

Zonozoe in relation to major taxonomic groupings: Euchelicerata, Artiopoda (not including

Vicissicaudata), and Vicissicaudata in Linear Discriminant space. Coloured circles show positions

of individual specimens. Histogram in B, C and D the shows the proportion of the trace for each

linear discriminant.

Table 1. List of specimens used for the outline analyses.

Species	Number	Age and origin	Clade	Reference	Figures
Zonoscutum solum	NM L 33021	Upper Ordovician, Czech Republic	Suspected Aglaspidida	(Van Roy et al., 2022)	Fig. 8C
Zonozoe drabowiensis	NM L 23586	Upper Ordovician, Czech Republic	Suspected Aglaspidida	Present work	Fig. 1E
Beckwithia typa	BPM 1060	Late Cambrian, Utah	Aglaspidida	(Lerosey-Aubril, Paterson, et al., 2017)	Fig. 9F
Chlupacaris dubia	NMSG.2005.103.1	Upper Ordovician, Morocco	Aglaspidida	(Van Roy, 2005)	Fig. 3A
Tremaglaspis unite	NHMIA56	Upper Ordovician, Wales	Aglaspidida	(Fortey & Rushton, 2003)	Fig. 1B
Glypharthrus trispinicaudatus	NIGPAS 165044	Late Cambrian, South China	Aglaspidida	(Lerosey-Aubril, Paterson, et al., 2017)	Fig. 3 G
Chraspedops modesta	MPM 11938	Late Cambrian, US	Aglaspidida	(Hesselbo, 1992)	Fig. 18, 4
Aglaspis barrandei	AMNH 39193	Late Cambrian, Wisconsin	Aglaspidida	(Hesselbo, 1992)	Fig. 2, 5
Xiphosura from Fezouata	YPM IP 532152	Lower Ordovician, , Morocco	Euchelicerata	(Lamsdell & Ocon, 2025)	Fig. 2, 7
Paraeurypterus anatoliensis	MTANHMSETR 10-İZ-01-1	Upper Ordovician, Turkey	Euchelicerata	(Lamsdell et al., 2013)	Fig. 3A
Lunataspis borealis	MM I-4583	Upper Ordovician, Canada	Euchelicerata	(Lamsdell et al., 2022)	Fig. 2, 1
Pseudoniscus falcatus	UK. NHMUK PI	Early Silurian, Scotland	Euchelicerata	(Bicknell & Pates, 2020)	Fig. 9A
Kasibelinurus amicorum	AM F68969	Devonian, Australia	Euchelicerata	(Bicknell & Pates, 2020)	Fig. 11A
Kwanyinaspis maotianshanensis	ELI-12004001	Early Cambrian, South China	Artiopoda	(Zhang & Shu, 2005)	Fig. 2A-D
Pygmaclypeatus daziensis	YKLP 11427	Early Cambrian, South China	Artiopoda	(Schmidt et al., 2025)	Fig. 1A
Thulaspis tholops	MGUH 34172a	Early Cambrian, North Greenland	Artiopoda	(Berks et al., 2023)	Fig. 1A
Retifacies abnormalis	JS-840	Early Cambrian, South China	Artiopoda, Trilobitomorpha	(Lin et al., 2024)	Fig. 1A
Kuamaia lata	YKLP 17296	Early Cambrian, South China	Artiopoda, Helmetiidae	(O'Flynn et al., 2024)	Fig. 2A
Emeraldella brocki	USNM 136641	Middle Cambrian, Canada	Artiopoda, Vicissicaudata	(Stein & Selden, 2012)	Fig. 3B
Tardisia broedeae	FMNH PE 88856	Late Carboniferous, Illinois	Artiopoda, Vicissicaudata	(McCoy et al., 2025)	Fig. 1D
Eozetetes gemmelli	SAM P48369a	Early Cambrian, Australia	Artiopoda, Vicissicaudata	(Edgecombe et al., 2017)	Fig. 2A
Carimersa neptuni	OUMNH PAL-C.376503	Early Silurian, England	Artiopoda, Vicissicaudata	(Briggs et al., 2023)	Fig. 1A
Sidneyia minor	YKLP 12435	Early Cambrian, South China	Artiopoda, Vicissicaudata	(Du et al., 2023)	Fig. 2A
Triopus draboviensis	NM L 16736	Upper Ordovician, Czech Republic	Cheloniellida	(Van Roy et al., 2022)	Fig. 2B
Drabovaspis complexa	NM L 23577	Upper Ordovician, Czech Republic	Cheloniellida	(Van Roy et al., 2022)	Fig. 9A
Neostrabops martini	25560	Late Cambrian, Missuri	Cheloniellida	(Caster & Macke, 1952)	Fig. 2

346   **Table 2.** Cross validation table following Linear Discriminant Analysis, showing the classification

347   of groups and the class accuracy. Class accuracy and classification not performed for *Zonoscutum*

348   and *Zonozoe* as they are represented only by a single specimen.

	Euchelicerata	Non-Vicissicadata	Vicissicaudata	*Zonoscutum*	*Zonozoe*	Class accuracy
Euchelicerata	4	0	0	0	1	80%
Non-Vicissicadata	1	0	4	0	0	0%
Vicissicaudata	0	4	7	2	1	50%

References

Barrande J. (1872). *Systême silurien du centre de la Bohême: trilobites, crustacés divers et*

poissons: supplément (Vol. 1). chez l'auteur et l'éditeur.

Berks H. O., Lunde Nielsen M., Flannery☐Sutherland J., Thorshøj Nielsen A., Park T. Y. S., &

Vinther J. (2023). A possibly deep branching artiopodan arthropod from the lower Cambrian

Sirius Passet Lagerstätte (North Greenland). *Papers in Palaeontology*, *9*(3), e1495.

Bicknell R. D., & Pates S. (2020). Pictorial atlas of fossil and extant horseshoe crabs, with focus on

Xiphosurida. *Frontiers in Earth Science*, *8*, 98.

Bonhomme V., Picq S., Gaucherel C., & Claude J. (2014). Momocs: outline analysis using R.

Journal of Statistical Software, *56*, 1-24.

Braig F., Torres G., Giménez L., & Haug J. T. (2024). Outline analysis as a new method for

investigating development in fossil crabs. *Palaeontologia Electronica*, *27*(1), 1-13.

Briggs D., Bruton D., & Whittington H. (1979). Appendages of the arthropod. *Aglaspis spinifer*.

Briggs D. E., Siveter D. J., Siveter D. J., Sutton M. D., Legg D., & Lamsdell J. C. (2023). A

vicissicaudatan arthropod from the Silurian Herefordshire Lagerstätte, UK. *Royal Society*

Open Science, *10*(8), 230661.

Broili F. (1932). Eine neue Crustacee aus dem rheinischen Unterdevon. *Sitzungsberichte der*

Mathematisch-Naturwissenschaftlichen (Abteilung) Klasse der Bayerischen Akademie der

Wissenschaften zu Munchen, 27–38.

Burke C. D., Full W. E., & Gernant R. E. (1987). Recognition of fossil fresh water ostracodes:

Fourier shape analysis. *Lethaia*, *20*(4), 307-314.

Caster K. E., & Macke W. B. (1952). An aglaspid merostome from the Upper Ordovician of Ohio.

Journal of Paleontology, *26*(5), 753-757.

Chlupáč I. (1963). Report on the merostomes from the Ordovician of Central Bohemia. *Věstník*

Ústředního ústavu geologického, *38*(6), 399-403.

Chlupáč I. (1965). Xiphosuran merostomes from the Bohemian Ordovician. *Sborník geologických*

věd, Paleontologie, *5*, 7-38.

Chlupáč I. (1999a). Some problematical arthropods from the Upper Ordovician Letná formation of

Bohemia. *Journal of Geosciences*, *44*(1-2), 79-92.

Chlupáč I. (1999b). Unusual arthropods from the Bohemian Ordovician-a review. *ACTA-*

UNIVERSITATIS CAROLINAE GEOLOGICA(1/2), 393-396.

Chlupác I., & Kukal Z. (1988). Possible global events and the stratigraphy of the Palaeozoic of the

Barrandian (Cambrian-Middle Devonian, Czechoslovakia). *Sborník geologických věd.*

Geologie(43), 83-146.

Collantes L., & Pates S. (in press). *Isoxys* carbonelli and the palaeoenvironmental disparity of

Isoxys during Cambrian Stage 3. *Historical Biology*.

Drage H. B., Laibl L., & Budil P. (2018). Postembryonic development of Dalmanitina, and the

evolution of facial suture fusion in Phacopina. *Paleobiology*, *44*(4), 638-659.

Drage H. B., & Pates S. (2024). Distinct causes underlie double-peaked trilobite morphological

disparity in cephalic shape. *Communications Biology*, *7*(1), 1490.

Du K., Bruton D. L., Yang J., & Zhang X. (2023). An early Cambrian *Sidneyia* (Arthropoda)

resolves the century-long debate of its head organization. *Science China Earth Sciences*,

66(3), 521-527.

Edgecombe G. D., Paterson J. R., & Garcia-Bellido D. C. (2017). A new aglaspidid-like

euarthropod from the lower Cambrian Emu Bay Shale of South Australia. *Geological*

Magazine, *154*(1), 87-95.

Eldredge N., & Smith L. (1974). Revision of the suborder Synziphosurina (Chelicerata,

Merostomata): with remarks on merostome phylogeny. *American Museum novitates*, *no.*

2543, 1– 41.

Foote M. (1989). Perimeter-based Fourier analysis: a new morphometric method applied to the

trilobite cranidium. *Journal of Paleontology*, *63*(6), 880-885.

Fortey R. A., & Rushton A. W. (2003). A new aglaspidid arthropod from the Lower Ordovician of

Wales. *Palaeontology*, *46*(5), 1031-1038.

Gevirtz J. L. (1976). Fourier analysis of bivalve outlines: implications on evolution and autecology.

Journal of the International Association for Mathematical Geology, *8*, 151-163.

Hall J. (1862). *On a new crustacean from the Potsdam Sandstone*.

Havlíček V. (1998). In: Chlupác□, I., Havlíc□ek, V., Kr□íž, J., Kukal, Z. and Štorch, P. (eds)

Palaeozoic of the Barrandian (Cambrian to Devonian). Czech Geological Survey, Prague,

41–79.. 41–79.

Healy-Williams N., & Williams D. F. (1981). Fourier analysis of test shape of planktonic

foraminifera. *Nature*, *289*(5797), 485-487.

Hesselbo S. P. (1992). Aglaspidida (Arthropoda) from the upper Cambrian of Wisconsin. *Journal of*

Paleontology, 885-923.

Hou X.-G. (1997). Arthropods of the lower Cambrian Chengjiang fauna, southwest China. *Fossils*

and Strata, *45*, 1-116.

Kraft P., Linnemann U., Mergl M., Bruthansová J., Laibl L., & Geyer G. (2023). Ordovician of the

Bohemian Massif.

Lamsdell J. C., Hoşgör İ., & Selden P. A. (2013). A new Ordovician eurypterid (Arthropoda:

Chelicerata) from southeast Turkey: evidence for a cryptic Ordovician record of

Eurypterida. *Gondwana Research*, *23*(1), 354-366.

Lamsdell J. C., Isotalo P. A., Rudkin D. M., & Martin M. J. (2022). A new species of the Ordovician

horseshoe crab Lunataspis. *Geological Magazine*, 1-5.

Lamsdell J. C., & Ocon S. B. (2025). Segmentation in early Xiphosura and the evolution of the

thoracetron. *Journal of Paleontology*, 1-20.

Legg D. A., Sutton M. D., & Edgecombe G. D. (2013). Arthropod fossil data increase congruence

of morphological and molecular phylogenies. *Nature communications*, *4*(1), 1-7.

Lerosey-Aubril R., Paterson J. R., Gibb S., & Chatterton B. D. (2017). Exceptionally-preserved late

Cambrian fossils from the McKay Group (British Columbia, Canada) and the evolution of

tagmosis in aglaspidid arthropods. *Gondwana Research*, *42*, 264-279.

Lerosey-Aubril R., Zhu X., & Ortega-Hernández J. (2017). The Vicissicaudata revisited–insights

from a new aglaspidid arthropod with caudal appendages from the Furongian of China.

Scientific Reports, *7*(1), 1-18.

Lin W., Pates S., Losso S. R., & Fu D. (2024). Intraspecific variation of early Cambrian (stage 3)

arthropod *Retifacies abnormalis* revealed by morphometric analyses. *Frontiers in Ecology*

and Evolution, *12*, 1336365.

Ma J., Pates S., Wu Y., Lin W., Liu C., Wu Y., Zhang M., & Fu D. (2023). Ontogeny and brooding

strategy of the early Cambrian arthropod *Isoxys minor* from the Qingjiang biota. *Frontiers in*

Ecology and Evolution, *11*, 1174564.

McCoy V. E., Herrera F., Wittry J., Mayer P., & Lamsdell J. C. (2025). A possible vicissicaudatan

arthropod from the Late Carboniferous Mazon Creek Lagerstätte. *Geological Magazine*,

162, e3.

Novák O. (1887). Merostomata. *Archive of the National Museum, Unpublished manuscript, Prague*.

O'Flynn R. J., Williams M., Liu Y., Hou X., Guo J., & Edgecombe G. D. (2024). The early

Cambrian *Kuamaia lata*, an artiopodan euarthropod with a raptorial frontal appendage.

Journal of Paleontology, *98*(5), 808-820.

Ortega-Hernandez J., Braddy S. J., & Rak Š. (2010). Trilobite and xiphosuran affinities for putative

aglaspidid arthropods Caryon and Drabovaspis, Upper Ordovician, Czech Republic.

Lethaia, *43*(3), 427-431.

Ortega‐Hernández J., Legg D. A., & Braddy S. J. (2013). The phylogeny of aglaspidid arthropods

and the internal relationships within Artiopoda. *Cladistics*, *29*(1), 15-45.

Pates S., Daley A. C., Legg D. A., & Rahman I. A. (2021). Vertically migrating *Isoxys* and the early

Cambrian biological pump. *Proceedings of the Royal Society B*, *288*(1953), 20210464.

Plotnick R. E., & Wagner P. J. (2006). Round up the usual suspects: common genera in the fossil

record and the nature of wastebasket taxa. *Paleobiology*, *32*(1), 126-146.

Raasch G. O. (1939). Cambrian merostomata. *Special Papers. Geological Society of America*, *19*,

1-146.

Rak S. (2009). The occurrence of non– trilobite arthropods from the quartzites of Letná Formation

from the vicinity of Beroun. *Geologie, paleonto– logie, speleologie–Ceský kras*, *35*, 14-25.

Rak Š., Bergström J., Fatka O., & Budil P. (2009). The Upper Ordovician arthropod *Zonozoe*

drabowiensis Barrande (Libeň and Letná formations, Sandbian, Barrandian area, Czech

Republic). *Bulletin of Geosciences*, *84*, 185-188.

RCoreTeam. (2024). R: A Language and Environment for Statistical Computing. R Foundation for

Statistical Computing. In. Vienna, Austria.

Schmidt M., Schoenemann B., Hou X., Melzer R. R., & Liu Y. (2025). *Pygmaclypeatus daziensis*, a

unique lower Cambrian arthropod with two different compound eye systems.

Communications Biology, *8*(1), 317.

Scotese C. R. (2023). Ordovician plate tectonic and palaeogeographical maps.

Stein M., & Selden P. A. (2012). A restudy of the Burgess Shale (Cambrian) arthropod *Emeraldella*

brocki and reassessment of its affinities. *Journal of Systematic Palaeontology*, *10*(2), 361-

383.

Størmer L. (1944). On the relationships and phylogeny of fossil and recent Arachnomorpha: a

comparative study on Arachnida, Xiphosura, Eurypterida, Tribolita, and other fossil

Arthropoda. *Skrifter Utgitt av Det Norske Videnskaps-Academi I Oslo. IMatematisk-*

Naturvidenskapelig Klasse, *5*, 1–158.

Størmer L. (1955). Merostomata in RC Moore, ed., Treatise on invertebrate paleontology, Pt. P,

Arthropoda 2: Geol. Soc. America and Univ. In: Kansas Press.

Van Roy P. (2005). An aglaspidid arthropod from the Upper Ordovician of Morocco with remarks

on the affinities and limitations of Aglaspidida. *Earth and Environmental Science*

Transactions of the Royal Society of Edinburgh, *96*(4), 327-350.

Van Roy P., Orr P. J., Botting J. P., Muir L. A., Vinther J., Lefebvre B., El Hariri K., & Briggs D. E.

(2010). Ordovician faunas of Burgess Shale type. *Nature*, *465*(7295), 215-218.

Van Roy P., Rak Š., Budil P., & Fatka O. (2022). Redescription of the cheloniellid euarthropod

Triopus draboviensis from the Upper Ordovician of Bohemia, with comments on the

affinities of *Parioscorpio venator. Geological Magazine*, *159*(9), 1471-1489.

Vokáč V., & Grigar L. (2024). New finding of the fossil arthropod *Zonozoe drabowiensis* Barrande,

1872 in the Řevnice Quartzite (Libeň Formation, Upper Ordovician, Sandbian) bear Starý

Plzenec (Prague Basin, Czech Republic). *Erica*, *31*, 83-87.

Walcott C. D. (1911). Cambrian geology and paleontology II. Middle Cambrian Merostomata.

Smithsonian Miscellaneous Collections, *57*(2), 17–40.

Zhang X., & Shu D. (2005). A new arthropod from the Chengjiang Lagerstätte, Early Cambrian,

southern China. *Alcheringa*, *29*(2), 185-194.

492