Extracellular vesicles from morphine-exposed prefrontal cortex carry transcriptomic and proteomic signatures of synaptic dysfunction

Pedro H. Gobira[1]; Savio Bastos[1]; Rachele Rossi [2]; Christian Bjerggaard Vægter[2]; Fenghua Chen[1]; Nicole Rodrigues da Silva[2]; Samia Joca[2]; Thomas Boesen [3]; Thiago Wendt Viola [4]; Yan Yan [5]; Martin R. Larsen[6]; Victor Corasolla Carregar [6]; Rodrigo Grassi-Oliveira[1,4]

[1] Translational Neuropsychiatry Unit, Department of Clinical Medicine, Aarhus University, Aarhus, Denmark; [2] Aarhus University, Biomedicine, Aarhus, Denmark; [3] Interdisciplinary Nanoscience Center -iNANO, Aarhus University, Aarhus; [4] Nanopsychiatry Lab, School of Medicine, Pontifical Catholic University of Rio Grande do Sul (PUCRS), Brazil; [5] EVGenomics ApS, Aarhus, Denmark; [6] Department of Biochemistry and Molecular Biology, University of Southern Denmark, Odense, Denmark.

Abstract

Extracellular vesicles (EVs) released by neurons and glial cells mediate intercellular communication in the brain and regulate synaptic function, neuronal survival, and neuropathological processes. Although chronic opioid exposure induces widespread neuroadaptations, the role of brain-derived EVs (BDEVs) in these processes remains poorly understood. We isolated BDEVs from the prefrontal cortex of rats subjected to chronic morphine exposure and performed integrative analyses of their RNA and protein cargo. Total RNA sequencing was combined with unbiased proteomics to define morphine-induced alterations. Functional validation was conducted by exposing primary cortical neurons to BDEVs from morphine- or vehicle-treated animals, followed by transcriptional profiling. Morphine significantly reprogrammed the transcriptome and proteome of BDEVs, with enrichment of pathways related to synaptic plasticity, endoplasmic reticulum stress, mitochondrial dysfunction, and neurodegeneration. Among the most robust changes, the synaptic regulator **ARC** and the ER stress marker **Hspa5** were consistently modulated at both mRNA and protein levels. In functional assays, morphine-derived BDEVs altered neuronal expression of genes associated with synaptic remodeling and excitability, including **Ncam1** and **Kcnn1**. These findings provide the first evidence that chronic opioid exposure reprograms BDEV cargo in a brain region critical for addiction, and that these vesicles are sufficient to induce transcriptional changes in recipient neurons. BDEVs emerge as active mediators of morphine-induced neuroadaptations and hold promise as biomarkers and therapeutic targets in opioid-related neuropathologies.

extracellular vesicles; morphine; prefrontal cortex; transcriptomics; proteomics; synaptic plasticity

Introduction

Extracellular vesicles (EVs) are nanosized particles secreted by virtually all cell types, serving as mediators of intercellular communication by transferring proteins, nucleic acids (DNA, mRNA, microRNAs, and long noncoding RNAs), and lipids to recipient cells (1, 2). Within the central nervous system, neurons, astrocytes, and microglia secrete EVs that play essential roles in brain function, including synaptic transmission, trophic support, and dendritic spine remodeling (3-7). Disruptions in these processes have been increasingly implicated in psychiatric conditions such as substance use disorder (SUD) (8, 9).

Emerging evidence suggests that brain-derived extracellular vesicles (BDEVs) may act as conduits through which addictive substances reshape neural circuitry. Exposure to drugs of abuse alters the molecular cargo of BDEVs, thereby influencing neurobiological responses to substances (8-12). To date, most studies have centered on small RNA species, particularly microRNAs, based on the long-standing assumption that EVs are enriched for this RNA class (13, 14). Yet recent findings challenge this view, showing that BDEVs can selectively package and transfer intact mRNAs that are translated in recipient neurons (15). One striking example is the activity-regulated cytoskeleton-associated protein (ARC) mRNA, which has been shown to be selectively incorporated into EVs and delivered functionally to neurons in vivo (16).

Despite this progress, a major knowledge gap remains: it is unknown whether chronic opioid exposure reprograms the full transcriptomic and proteomic composition of BDEVs, and whether these vesicles are sufficient to drive transcriptional reorganization in recipient neurons. Addressing this gap is critical, given the escalating opioid crisis and the urgent need to identify novel molecular mechanisms underpinning opioid-induced neuroadaptations. Here, we directly test the hypothesis that chronic morphine exposure remodels BDEV cargo and that these vesicles, once secreted, are sufficient to reshape neuronal gene expression in ways relevant to opioid-evoked behavioral phenotypes.

We isolate BDEVs from the prefrontal cortex (PFC)—a region crucial for addiction-related plasticity—and comprehensively profile their proteome and transcriptome. To assess functional impact, we expose primary cortical neurons to BDEVs from vehicle- and

morphine-treated donors and quantify transcriptional responses. This approach establishes a mechanistic framework for understanding EV-mediated contributions to opioid-induced neuroplasticity, with implications for both biomarker discovery and therapeutic innovation.

Methods and Materials

Methods are summarized below. A complete description of all experimental procedures is available in Supplementary Materials.

Animals and drug treatment

Male Sprague–Dawley rats (8 weeks old, Janvier Labs) were pair-housed under standard conditions with ad libitum access to food and water. After two weeks of habituation, animals received intraperitoneal injections of saline or escalating doses of morphine (10–50 mg/kg, twice daily for 5 days). Four hours after the final injection, rats were sacrificed, and bilateral prefrontal cortex (PFC) was dissected and frozen at −80 °C. All procedures were approved by the Danish Animal Experiments Inspectorate and complied with EU Directive 2010/63/EU.

Isolation and characterization of brain-derived extracellular vesicles (BDEVs)

BDEVs were isolated from PFC tissue by enzymatic dissociation, differential centrifugation, filtration, and ultracentrifugation, as previously described (18). Vesicle purity and morphology were assessed by BCA protein assay, Western blotting, transmission electron microscopy (TEM), and nanoparticle tracking analysis (NTA). Full protocols are provided in Supplementary Methods.

RNA extraction, sequencing, and analysis

RNA was extracted from BDEVs, DNA was removed, and total RNA libraries were prepared using a stranded RNA-seq kit. Libraries were sequenced on an Illumina NovaSeq X Plus platform (150 bp, paired-end). Quality control, differential gene expression analysis (DESeq2), and pathway enrichment (Metascape, GO, and Reactome) followed standard procedures, with significance defined at FDR-adjusted $p < 0.1$ and log2FC ±0.5.

Proteomic analysis

BDEV proteins were extracted, digested with trypsin, and analyzed by LC–MS/MS on an Orbitrap Q Exactive HF system. Data were searched against the Rattus norvegicus UniProt database and filtered at 1% FDR. Proteins were quantified based on unique peptides, and enrichment analyses were conducted using Cytoscape, IPA, and R-based workflows.

Primary neuronal cultures and BDEV treatment

Cortical neurons were prepared from embryonic day 17 (E17) C57BL/6 embryos and plated in Neurobasal medium with B27 supplement. On DIV7, neurons were treated with BDEVs isolated from saline- or morphine-exposed rats, and RNA was extracted 24 h later.

NanoString gene expression profiling

Gene expression in primary neurons was measured using the NanoString nCounter® Mouse Neuropathology Panel (770 genes). Data were normalized to housekeeping genes using nSolver™ 4.0, and pathway enrichment was conducted in Reactome.

Results

Characterization of EVs from the prefrontal cortex of chronically morphine-treated rats

BDEVs were isolated from the prefrontal cortex of rats via ultracentrifugation as previously described (17). We used NTA to determine the size distribution and particle number, with the latter normalized to the individual prefrontal cortex weight. Results revealed no significant differences between animals treated with morphine and those treated with vehicle **(Fig. 1B).** Using TEM, we characterized the BDEVs, revealing their characteristic morphology (**Fig. 1C**). We also analyzed BDEVs via specific positive and negative markers via Western blotting. As expected, the positive markers flotillin-1 and Hsp70 were detected in the BDEV fractions, whereas the negative marker GM130 was absent from the BDEV fractions but present in the whole cortex lysates (**Fig. 1D**). Together, these findings confirm the reliability of our isolation technique.

Effect of morphine treatment on the RNA profile of BDEVs

The RNA-seq analysis identified a total of 12,224 RNA molecules that surpassed the quality control threshold for downstream analysis, comprising 49 long noncoding RNAs (0.40%), 94 small noncoding RNAs (0.77%), 11,948 protein-coding mRNA molecules

(97.74%), and other RNAs (e.g., ribosomal RNA, transfer RNA, mitochondrial RNA, and pseudogenes—1.09%). Prior to conducting the differential gene expression (DGE) analysis, the RNA-seq data were deconvoluted from publicly available single-cell RNA sequencing data from rodent brains. We analyzed the proportions of microglia, glutamatergic corticothalamic (CT), intratelencephalic (IT), and pyramidal tract (PT) neurons, as well as GABAergic interneurons. The proportions of oligodendrocytes, astrocytes, and glutamatergic NP neurons were negligible. No significant group differences in cell-type proportions were observed (all p values > 0.05; see **Fig.** 2A). However, we observed that almost 40% of the RNA-seq data were from BDEVs originating from GABAergic interneurons.

A comprehensive analysis revealed 625 RNA transcripts with unadjusted p values less than 0.05 (Supplemental Table 1), indicating an effect of morphine treatment on the EV transcriptomic cargo. Four hits were identified by DGE analysis as having significant group differences that surpassed the FDR correction thresholds: Arc, Hspa5, Hivep3, and RT1-CE16 **(Fig. 2B)**. These four molecules are mRNAs derived from protein-coding genes. Arc, Hspa5, and Hivep3 were upregulated in EVs from morphine-treated animals, whereas RT1-CE16 was downregulated. Additionally, we cross-referenced our RNA-seq data with a large-scale human genome-wide association study (GWAS) meta-analysis of opioid use disorder (18) to identify differentially regulated transcripts originating from previously identified genes associated with the disorder, such as Oprm1, Cdkal1, Btnl2, Rabepk, Fbxw4, Ncam1, Furin, Kcnn1, and Rnf114. Among these, two transcripts, Ncam1 (log2FC = 0.45; p value = 0.02) and Kcnn1 (log2FC = 0.49; p value = 0.04), had unadjusted p values below 0.05, indicating a greater abundance of EVs in morphine-treated animals than in control animals **(Fig. 2C).**

Pathway enrichment analysis was conducted using all RNAs with an unadjusted p value < 0.05 on the basis of the KEGG 2019 mouse database (*Rattus norvegicus* gene codes were converted to mice before enrichment). Among the enriched pathways, only Lysosome, Protein Processing in the Endoplasmic Reticulum, and Axon Guidance were significantly represented, predominantly comprising upregulated transcripts (Figure y1 a-b). Notably, the enriched RNAs within each pathway were mutually exclusive, with no individual RNA shared across multiple pathways. This suggests a limited overlap and low interpathway correlation among the transcriptional alterations observed (Figure y2a-b).

Effect of morphine treatment on the protein profile of BDEVs

A total of 1007 proteins were reliably quantified, 379 of which passed the quality control criteria for downstream analysis. Among these, 82 proteins were found to be dysregulated on the basis of unadjusted p values < 0.05 (Supplemental Table 2), including 44 downregulated and 38 upregulated proteins. However, none of these changes remained statistically significant after false discovery rate (FDR) correction. Consequently, we adopted a more permissive threshold, selecting proteins with unadjusted p values < 0.01 and an absolute log2-fold change (log2FC) ≥ 0.5 to highlight potentially relevant alterations, which is consistent with prior proteomic studies applying exploratory thresholds in complex biological models (19, 20). Using these criteria, ten proteins were identified as differentially regulated in brain-derived extracellular vesicles (BDEVs) following morphine exposure (Fig. 3). Among these proteins, Krt33a, Fabp5, Krt72, Sprr1a, and Pkp1 were downregulated, whereas Stx1b, Ywhag, Grm8, Hspa5, and Pdia3 were upregulated.

A volcano plot was used to visualize the overall distribution of protein abundance changes, clearly highlighting the most significantly dysregulated proteins (p value < 0.05). Furthermore, dynamic range analysis revealed a broad spectrum of protein abundances, indicating that both low- and high-abundance proteins were successfully identified and quantified, an important aspect of exosome proteomics that enables the detection of diverse cargo molecules (Figure 3a-b) (21, 22).

To elucidate the biological implications of these morphine-associated protein alterations, pathway enrichment analysis was performed via the KEGG Mouse 2019 database (23). The dysregulated proteins were significantly associated with pathways involved in neurodegenerative diseases, oxidative phosphorylation, nonalcoholic fatty liver disease, the citrate (TCA) cycle, retrograde endocannabinoid signaling, thermogenesis, and protein processing in the endoplasmic reticulum (Figure 3 C).

Inspection of the dysregulation patterns revealed notable enrichment of mitochondrial proteins, particularly members of the NDUF family, as well as COX1 and COX7A2L (Figure 3 D-E), which play essential roles in oxidative phosphorylation and energy metabolism, mechanisms implicated in neuropsychiatric disorders (24-26). Additionally, the upregulation of the GRM5-encoded metabotropic glutamate receptor in the exosomal cargo of morphine-exposed subjects suggests potential activation of the phosphatidylinositol–calcium second messenger signaling pathway, which is known to influence synaptic plasticity and drug-related neuroadaptation (27, 28).

Integrative analysis of differentially regulated RNAs and proteins

Proteomic analysis identified 1,084 proteins, of which 1,007 were quantifiable. Notably, more than 85% overlapped with transcripts detected at the mRNA level, demonstrating substantial concordance between the transcriptomic and proteomic datasets. However, this overlap was not preserved within statistically dysregulated molecules were considered (p value < 0.05, unadjusted). Only five genes—Hspa5, S100a6, Plp1, Slc24a2, and Uggt1—were commonly dysregulated at both the mRNA and protein levels (Figure 4 A-B). In addition to the complexity of EVs, this limited concordance likely reflects the influence of posttranscriptional and posttranslational regulatory mechanisms, including mRNA stability, translational efficiency, and protein turnover, which are well documented contributors to the discordance between transcriptomic and proteomic data.

Among the enriched pathways identified independently from the transcriptomic and proteomic datasets, protein processing in the endoplasmic reticulum was the only pathway common to both analyses. Interestingly, within this pathway, Hspa5 is upregulated at both the mRNA and protein levels in response to morphine exposure in BDEVs, suggesting conserved stress response activation (29, 30). In contrast, other transcripts contributing to this pathway were upregulated at the mRNA level, while their corresponding proteins were downregulated, further supporting the role of translational regulation or differential protein degradation in shaping the exosomal proteome under the influence of morphine (Figure 4 C-D) (31, 32).

Morphine-BDEVs modulate neurodegenerative gene expression and activate opioid signaling in cross-species recipient neurons

To investigate the functional role of BDEVs and their potential to modulate neurodegenerative processes, we treated primary neurons with morphine-BDEVs and control-BDEVs, followed by a comprehensive multiplex gene expression analysis targeting 770 neuropathology-associated genes **(Fig. 5A)**. After controlling for multiple comparisons, we identified significant differential expression of 15 genes in primary neuronal cells following treatment with morphine-derived BDEVs (FDR q value < 0.1). Among these genes, *Mnat1*, a gene implicated in blood brain barrier (BBB) damage in Alzheimer's disease (AD), was downregulated, whereas *PSEN1*, *NSF*, and *NRG1*, which are associated with neurodegenerative processes, particularly in AD, were upregulated **(Fig. 5B)**. This evidence supports previous findings suggesting that extracellular vesicles (EVs) may contribute to the neurodegeneration and dementia observed with long-term opioid use(9).

Pathway analysis revealed a selective enrichment of non-canonical WNT signaling components in neurons exposed to BDEVs derived from morphine-treated rats, despite the absence of direct morphine exposure. The top-ranked pathway was "Ca2+ pathway" (FDR = 2.75 × 10□□; ratio = 0.005), followed by "Beta-catenin independent WNT signaling" (FDR = 5.71 × 10□□; ratio = 0.01), and "Signaling by WNT" (FDR = 1.64 × 10□□; ratio = 0.02). These non-canonical pathways are critical for cytoskeleton remodeling, intracellular calcium flux, and neuronal polarity. Remarkably, neurons also showed enrichment for "DARPP-32 events" and "Opioid signalling" (both FDR = 0.004), indicating activation of dopaminergic integration and GPCR-related neuromodulatory cascades. These results suggest that morphine-induced BDEVs can transfer biologically active signals capable of reprogramming recipient neurons toward a phenotype consistent with drug exposure, engaging both polarity-regulating pathways and classical neuroadaptive circuits **(Fig. 5C)**.

Discussion

This study demonstrates that chronic morphine induces significant transcriptomic and proteomic reprograming in the cargo of BDEVs isolated from the rat prefrontal cortex. wich are capable of dysregulating neuronal function in in vitro. Through transcriptomic, proteomic, and integrative analyses, we identified BDEV-enriched gene and protein signatures involved in synaptic plasticity, neurodegenerative pathways, and ER stress, including upregulation of key modulators such as ARC and Hspa5. Functional assays further revealed that morphine-altered BDEVs actively reshape neuropathology-associated gene expression in recipient neurons, implicating these vesicles as mediators of opioid-induced neuroadaptation and contributors to long-term neuropathological processes.

EVs play critical roles in intercellular communication, and alterations in their cargo have been increasingly linked to opioid responses (3, 9). Previous studies have shown that the EV cargo changes, mainly in microRNAs in blood-derived EVs in adult individuals and animals exposed to opioids (33, 34). Similarly, BDEVs from animals exposed to opioids in utero and postnatally showed an upregulation of miR-504 and a downregulation of miR-30c-1 (10). However, these studies profiled small RNAs, limiting insights into the long transcript diversity of EVs (e.g. mRNAs and lncRNAs) (35). In this study, we employed total RNA sequencing with rRNA removal, allowing the identification of mRNAs in BDEVs, which is consistent with recent advances showing functional mRNAs in EVs (15).

Our RNA sequencing data revealed an increase in the expression of the ARC gene in BDEVs from animals treated with morphine. ARCs play a central role in synaptic plasticity and neuronal adaptation to chronic drug exposure, with previous studies reporting elevated ARC expression in the bulk prefrontal cortex of animals with a history of opioid exposure (36-38). Interestingly, ARC has also been implicated in the biogenesis and release of EVs. It can self-assemble into mRNA-containing, virus-like capsids that are packaged into EVs and released by neurons, where they can be endocytosed by recipient cells at postsynaptic sites (16, 39, 40). These ARC-containing EVs are proposed to modulate synaptic plasticity by delivering ARC mRNA to neighboring neurons, where local translation facilitates AMPA-type glutamate receptor (AMPAR) endocytosis and synaptic remodeling (40). Therefore, the observed upregulation of the ARC gene in BDEVs following morphine treatment may reflect an EV-mediated mechanism contributing to opioid-induced synaptic plasticity, a pivotal process underlying addiction-related neuroadaptation.

This interpretation aligns with our pathway enrichment analysis, which revealed significant enrichment of axon guidance, lysosome, and protein processing in the endoplasmic reticulum (ER) pathways, which were driven primarily by upregulated transcripts. These pathways are relevant to neuronal adaptations triggered by opioids (41). For example, previous studies have demonstrated altered expression of specific axon guidance genes in the adult mouse brain following chronic self-administration of oxycodone (42), which may reflect alterations in axon–target connectivity and synaptogenesis, potentially contributing to disrupted neural circuits and the behavioral changes associated with opioid addiction. Together, these findings suggest that morphine exposure provokes transcriptional changes in BDEVs that may underline core molecular mechanisms associated with opioid-related neuroplasticity.

We applied deconvolutional analysis to estimate the cell type origin of BDEVs by aligning the EV transcriptome to establish expression profiles of specific brain cell types (43). We found that approximately 40% of BDEV RNA originated from GABAergic interneurons, which are known to modulate reward functions affected by opioids (44). These findings, together with recent evidence showing the behavioral effects of GABAergic EVs (45), point to a role for interneuron-derived EVs in opioid-related neuroplasticity.

To increase our understanding of the functional implications of morphine-induced changes in BDEV cargo, we cross-referenced our RNA-seq data with a large-scale

human GWAS meta-analysis for opioid use disorder (OUD) (18). This integrative approach allowed us to identify differentially expressed transcripts originating from genes previously associated with OUD susceptibility, such as Oprm1, Furin, Ncam1 Kcnn1, and Rnf114 (18). We observed that Ncam1 and Kcnn1 transcripts were significantly more abundant in BDEVs from morphine-treated animals than in those from control animals. These genes are functionally relevant to opioid effects, with Ncam1 playing a role in synaptic plasticity, whereas Kcnn1 is involved in regulating neuronal excitability, highlighting a potential role of BDEVs in morphine-induced synaptic remodeling (46-48).

Our proteomic analyses r further demonstrated significant changes in the cargo of BDEVs following chronic morphine treatment by identifying dysregulated proteins implicated in morphine-induced neuroadaptation. Specifically, we observed increased levels of proteins associated with synaptic plasticity and neurodegenerative processes, including PDIA3, STX1B, YWHAG, and HSPA5, n previouslylinked to synaptic dysfunction and pathways relevant to neurodegenerative diseases (49-52). These findings align with previous evidence suggesting that chronic opioid exposure may increase the risk of neurodegeneration through mechanisms involving EVs (53). For example, increased levels of protein biomarkers of neurodegenerative processes have been detected in EVs isolated from the plasma of monkeys exposed to oxycodone (34). Similarly, morphine-treated astrocytes have been shown to secrete EVs enriched with amyloid-related proteins, contributing to neurodegenerative processes in vitro (54).

Consistent with these findings, our pathway enrichment analysis revealed significant associations between the dysregulated proteins and pathways implicated in neurodegenerative diseases. Notably, inspection of the dysregulation patterns revealed an enrichment of mitochondrial proteins, including COX1, COX7A2L, and members of the NDUF family, such as NDUFV1 and NDUFC2. These proteins play critical roles in mitochondrial function and are important for ATP production and cellular energy metabolism (55-57). Chronic opioid use has been linked to mitochondrial dysfunction, potentially driven by increased production of reactive oxygen species, which can damage mitochondrial DNA and disrupt cellular energy metabolism, processes closely associated with neurodegenerative pathologies (25, 58). Therefore, the observed alterations in mitochondrial proteins in BDEVs may contribute to increased neuronal vulnerability and the progression of opioid-induced neurodegenerative processes. Similarly, abnormal levels of mitochondrial proteins in EVs

from plasma have also been reported in neurodegenerative conditions, including Parkinson's disease and Alzheimer's disease (59, 60)

Moreover, our proteomic analysis also demonstrated the upregulation of the GRM5-encoded metabotropic glutamate receptor in BDEVs from morphine-exposed animals. This observation suggests potential alterations in glutamatergic signaling pathways, possibly affecting synaptic plasticity mechanisms relevant to neurodegenerative processes. Aberrant neuronal signaling mediated by mGluR5, which leads to synaptic dysfunction, has been recognized as an emerging mechanism in the biology of both drug addiction and neurodegeneration(61, 62). In support of this concept, a recent study reported elevated mGluR5 expression in the hippocampus of individuals with Alzheimer's disease, with expression levels positively correlated with disease progression in both human patients and mouse models (63). Although the presence of mGluR5 has not previously been described specifically in extracellular vesicles isolated from brain tissue after morphine treatment, its occurrence in EVs has been reported in animal models of Alzheimer's disease (64). Our findings align with these observations and further suggest that mGluR5 in BDEVs could represent a novel mechanism contributing to opioid-induced neurodegenerative changes.

We also performed an integrative analysis, which revealed substantial overlap (>85%) between proteins identified via proteomics and their corresponding transcripts detected via RNA sequencing, supporting the overall consistency and quality of our datasets. However, when statistically dysregulated molecules are specifically evaluated, this concordance notably diminishes. Five genes, *Hspa5*, *S100a6*, *Plp1*, *Slc24a2*, and *Uggt1*, exhibited common dysregulation at both the mRNA and protein levels. This limited overlap highlights the inherent complexity involved in selectively packaging cargo into EVs. Cargo loading into EVs is governed by an interplay of mechanisms, including several posttranslational modifications, such as ubiquitination, and interactions with RNA-binding proteins (65). Although these mechanisms have not been fully elucidated in BDEVs, it has been suggested that chronic opioid exposure may alter some of these selective cargo-loading processes, potentially influencing the BDEV cargo of both proteins and mRNAs (66, 67). Moreover, beyond selective cargo packaging, discrepancies between transcriptomic and proteomic profiles may be further influenced by factors such as mRNA stability, translational efficiency, and protein turnover, as previously described, including in the neural context (68-70).

Among the biological pathways independently enriched in both datasets, protein processing in the endoplasmic reticulum (ER) emerged as the common pathway, suggesting a convergent stress response signature in BDEVs following morphine exposure. Notably, Hspa5 was upregulated at both the transcript and protein levels, highlighting its potential as a biomarker for ER stress induced by chronic opioid treatment. Previous studies have demonstrated that the activation of opioid receptors triggers cellular events leading to ER stress, which significantly contributes to common opioid-related phenomena such as tolerance (71). Indeed, substantial upregulation of Hspa5, a well-established ER stress marker, has been reported in the spinal cord and several mesolimbic brain regions, including the prefrontal cortex, following opioid exposure (71-74). Taken together, these findings indicate that elevated Hspa5 expression in BDEVs is associated with morphine-induced cellular responses, suggesting that chronic opioid exposure modulates EV cargo composition through ER stress pathways. Thus, BDEVs represent promising platforms for investigating and potentially monitoring the neurobiological changes associated with opioid use.

Finally, to explore whether the molecular reprogramming observed in the BDEVs from morphine-treated rats would have functional consequences for recipient neurons, we evaluated the transcriptional response of primary neuronal cells to BDEVs isolated from morphine- and vehicle-exposed animals. A targeted multiplex gene expression analysis of 770 neuropathology-associated genes revealed significant differential expression of 15 genes following morphine-BDEV treatment. Among these genes, Mnat1, a gene involved in blood brain barrier integrity and linked to Alzheimer's disease, was downregulated (75). Conversely, Psen1, Nsf, and Negr1, all of which are associated with neurodegenerative processes, were upregulated (76-78). These findings indicate that morphine-associated BDEVs are sufficient to induce transcriptional programs in naïve neurons that mimic canonical features of opioid exposure. Notably, the selective enrichment of pathways such as DARPP-32 events and Opioid Signalling emerged in the absence of any direct pharmacological stimulus, suggesting that BDEVs carry bioactive cargo capable of reprogramming neuronal signaling cascades. In addition, the co-enrichment of non-canonical WNT and calcium-related pathways points to intracellular remodeling processes linked to neuronal polarization, cytoskeletal plasticity, and synaptic modulation, processes associated with neuroadaptive responses typically observed during the initial stages of drug exposure or sensitization (79-81).

Crucially, transcriptomic profiling of the BDEV cargo revealed robust overexpression of ARC mRNA, a key regulator of synaptic scaling and long-term depression (LTD) (82, 83) . ARC mRNA is known to be selectively packaged into neuronal EVs and has been shown to transfer plasticity-related signals between cells(84). Its presence in morphine-altered BDEVs likely contributes to the observed activation of non-canonical WNT pathways and calcium flux components, both of which underlie actin cytoskeleton rearrangement, spine destabilization, and polarity shifts. Because the recipient cultures were never exposed to morphine, these results suggest that morphine BDEVs act as vectors of neuroplasticity, reprogramming target neurons toward a transcriptional and structural state resembling that of drug-exposed circuits. This supports a model in which BDEVs mediate a trans-synaptic form of memory encoding and adaptation, potentially contributing to the systems-level persistence of drug-induced neuroadaptations.

In summary, our findings demonstrate that chronic morphine exposure significantly alters the molecular cargo of BDEVs, especially those associated with pathways involved in synaptic function, cellular stress responses, and neurodegeneration. Functional assays confirmed that these vesicles can modulate gene expression in recipient neurons, including genes implicated in neurodegeneration. Together, these results highlight BDEVs as crucial mediators of the persistent neural adaptations induced by chronic opioid use and suggest their potential as biomarkers or therapeutic targets in opioid-related neuropathologies.

In summary, our findings are the first demonstrating that BDEVs derived from the brain of rats chronically exposed to morphine present significant changes in their molecular content reflecting pathways associated with synaptic (dys)function, cellular stress responses, and neurodegeneration. Furthermore, our functional assays originally demonstrate that these vesicles trigger transcriptional reprogramming in recipient neurons, including the regulation of genes implicated in neurodegeneration. Together, these results highlight BDEVs content as crucial mediators of the persistent neural adaptations induced by chronic opioid use and suggest their potential as biomarkers and/or therapeutic targets in opioid-related neuropathologies.

Acknowledgements

This work was supported by the Independent Research Fund Denmark (DFF, grant no. 3166-00139B) and by the Brazilian National Council for Scientific and Technological Development (CNPq, grant nos. 444358/2024-2, 401678/2022-9, and 306580/2022-5), all awarded to Dr. Rodrigo Grassi-Oliveira (RGO).

Disclosures

All authors report no biomedical financial interests or potential conflicts of interest.

References

1.	Gruner HN, McManus MT (2021): Examining the evidence for extracellular RNA function in mammals. *Nat Rev Genet*. 22:448–458.
2.	van Niel G, Carter DRF, Clayton A, Lambert DW, Raposo G, Vader P (2022): Challenges and directions in studying cell-cell communication by extracellular vesicles. *Nat Rev Mol Cell Biol*. 23:369–382.
3.	Kim Y, D'Acunzo P, Levy E (2024): Biogenesis and secretion of mitovesicles, small extracellular vesicles of mitochondrial origin at the crossroads between brain health and disease. *Curr Opin Physiol*. 40.
4.	Filannino FM, Panaro MA, Benameur T, Pizzolorusso I, Porro C (2024): Extracellular Vesicles in the Central Nervous System: A Novel Mechanism of Neuronal Cell Communication. *Int J Mol Sci*. 25.
5.	Korkut C, Ataman B, Ramachandran P, Ashley J, Barria R, Gherbesi N, et al. (2009): Trans-synaptic transmission of vesicular Wnt signals through Evi/Wntless. *Cell*. 139:393–404.
6.	Antonucci F, Turola E, Riganti L, Caleo M, Gabrielli M, Perrotta C, et al. (2012): Microvesicles released from microglia stimulate synaptic activity via enhanced sphingolipid metabolism. *EMBO J*. 31:1231–1240.
7.	Li K, Wang K, Xu SX, Xie XH, Tang Y, Zhang L, et al. (2024): Investigating Neuroplasticity Changes Reflected by BDNF Levels in Astrocyte-Derived Extracellular Vesicles in Patients with Depression. *Int J Nanomedicine*. 19:8971–8985.
8.	Rao PSS, O'Connell K, Finnerty TK (2018): Potential Role of Extracellular Vesicles in the Pathophysiology of Drug Addiction. *Mol Neurobiol*. 55:6906–6913.
9.	Morris DC, Zacharek A, Zhang ZG, Chopp M (2023): Extracellular vesicles-Mediators of opioid use disorder? *Addict Biol*. 28:e13353.
10.	Chand S, Gowen A, Savine M, Moore D, Clark A, Huynh W, et al. (2021): A comprehensive study to delineate the role of an extracellular vesicle-associated microRNA-29a in chronic methamphetamine use disorder. *J Extracell Vesicles*. 10:12177.
11.	Shahjin F, Guda RS, Schaal VL, Odegaard K, Clark A, Gowen A, et al. (2019): Brain-Derived Extracellular Vesicle microRNA Signatures Associated with In Utero and Postnatal Oxycodone Exposure. *Cells*. 9.
12.	Landfield Q, Saito M, Hashim A, Canals-Baker S, Sershen H, Levy E, et al. (2021): Cocaine Induces Sex-Associated Changes in Lipid Profiles of Brain Extracellular Vesicles. *Neurochem Res*. 46:2909–2922.

13.	O'Brien K, Breyne K, Ughetto S, Laurent LC, Breakefield XO (2020): RNA delivery by extracellular vesicles in mammalian cells and its applications. *Nat Rev Mol Cell Biol*. 21:585–606.

14.	Dellar ER, Hill C, Melling GE, Carter DRF, Baena-Lopez LA (2022): Unpacking extracellular vesicles: RNA cargo loading and function. *J Extracell Biol*. 1:e40.

15.	Ransom LS, Liu CS, Dunsmore E, Palmer CR, Nicodemus J, Ziomek D, et al. (2024): Human brain small extracellular vesicles contain selectively packaged, full-length mRNA. *Cell Rep*. 43:114061.

16.	Pastuzyn ED, Day CE, Kearns RB, Kyrke-Smith M, Taibi AV, McCormick J, et al. (2018): The Neuronal Gene Arc Encodes a Repurposed Retrotransposon Gag Protein that Mediates Intercellular RNA Transfer. *Cell*. 172:275–288 e218.

17.	D'Acunzo P, Kim Y, Ungania JM, Perez-Gonzalez R, Goulbourne CN, Levy E (2022): Isolation of mitochondria-derived mitovesicles and subpopulations of microvesicles and exosomes from brain tissues. *Nat Protoc*. 17:2517–2549.

18.	Kember RL, Vickers-Smith R, Xu H, Toikumo S, Niarchou M, Zhou H, et al. (2022): Cross-ancestry meta-analysis of opioid use disorder uncovers novel loci with predominant effects in brain regions associated with addiction. *Nat Neurosci*. 25:1279–1287.

19.	Zhang B, Wang J, Wang X, Zhu J, Liu Q, Shi Z, et al. (2014): Proteogenomic characterization of human colon and rectal cancer. *Nature*. 513:382–387.

20.	Cox J, Mann M (2008): MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat Biotechnol*. 26:1367–1372.

21.	Thery C, Witwer KW, Aikawa E, Alcaraz MJ, Anderson JD, Andriantsitohaina R, et al. (2018): Minimal information for studies of extracellular vesicles 2018 (MISEV2018): a position statement of the International Society for Extracellular Vesicles and update of the MISEV2014 guidelines. *J Extracell Vesicles*. 7:1535750.

22.	Li M, Zeringer E, Barta T, Schageman J, Cheng A, Vlassov AV (2014): Analysis of the RNA content of the exosomes derived from blood serum and urine and its potential as biomarkers. *Philos Trans R Soc Lond B Biol Sci*. 369.

23.	Kanehisa M, Sato Y, Furumichi M, Morishima K, Tanabe M (2019): New approach for understanding genome variations in KEGG. *Nucleic Acids Res*. 47:D590–D595.

24.	Lin MT, Beal MF (2006): Mitochondrial dysfunction and oxidative stress in neurodegenerative diseases. *Nature*. 443:787–795.

25.	Federico A, Cardaioli E, Da Pozzo P, Formichi P, Gallus GN, Radi E (2012): Mitochondria, oxidative stress and neurodegeneration. *J Neurol Sci*. 322:254–262.

26.	Wallace DC (2018): Mitochondrial genetic medicine. *Nat Genet*. 50:1642–1649.

27.	Niswender CM, Conn PJ (2010): Metabotropic glutamate receptors: physiology, pharmacology, and disease. *Annu Rev Pharmacol Toxicol*. 50:295–322.

28.	Trantham-Davidson H, Chandler LJ (2015): Alcohol-induced alterations in dopamine modulation of prefrontal activity. *Alcohol*. 49:773–779.

29.	Lee AS (2001): The glucose-regulated proteins: stress induction and clinical applications. *Trends Biochem Sci*. 26:504–510.

30.	Chen X, Shi C, He M, Xiong S, Xia X (2023): Endoplasmic reticulum stress: molecular mechanism and therapeutic targets. *Signal Transduct Target Ther*. 8:352.

31.	Tyanova S, Temu T, Cox J (2016): The MaxQuant computational platform for mass spectrometry-based shotgun proteomics. *Nat Protoc*. 11:2301–2319.

32.	Jung M, Lee EK (2021): RNA-Binding Protein HuD as a Versatile Factor in Neuronal and Non-Neuronal Systems. *Biology (Basel)*. 10.

33. Chen F, Zou L, Dai Y, Sun J, Chen C, Zhang Y, et al. (2021): Prognostic plasma exosomal microRNA biomarkers in patients with substance use disorders presenting comorbid with anxiety and depression. *Sci Rep*. 11:6271.

34. Kumar A, Kim S, Su Y, Sharma M, Kumar P, Singh S, et al. (2021): Brain cell-derived exosomes in plasma serve as neurodegeneration biomarkers in male cynomolgus monkeys self-administrating oxycodone. *EBioMedicine*. 63:103192.

35. Miceli RT, Chen TY, Nose Y, Tichkule S, Brown B, Fullard JF, et al. (2024): Extracellular vesicles, RNA sequencing, and bioinformatic analyses: Challenges, solutions, and recommendations. *J Extracell Vesicles*. 13:e70005.

36. Shepherd JD, Bear MF (2011): New views of Arc, a master regulator of synaptic plasticity. *Nat Neurosci*. 14:279–284.

37. Garcia-Perez D, Ferenczi S, Kovacs KJ, Milanes MV (2019): Distinct regulation pattern of Egr-1, BDNF and Arc during morphine-withdrawal conditioned place aversion paradigm: Role of glucocorticoids. *Behav Brain Res*. 360:244–254.

38. Li M, Liu WJ, Lu B, Wang YH, Liu JG (2013): Differential expression of Arc in the mesocorticolimbic system is involved in drug and natural rewarding behavior in rats. *Acta Pharmacol Sin*. 34:1013–1024.

39. Ashley J, Cordy B, Lucia D, Fradkin LG, Budnik V, Thomson T (2018): Retrovirus-like Gag Protein Arc1 Binds RNA and Traffics across Synaptic Boutons. *Cell*. 172:262–274 e211.

40. Hantak MP, Einstein J, Kearns RB, Shepherd JD (2021): Intercellular Communication in the Nervous System Goes Viral. *Trends Neurosci*. 44:248–259.

41. Dacher M, Nugent FS (2011): Opiates and plasticity. *Neuropharmacology*. 61:1088–1096.

42. Yuferov V, Zhang Y, Liang Y, Zhao C, Randesi M, Kreek MJ (2018): Oxycodone Self-Administration Induces Alterations in Expression of Integrin, Semaphorin and Ephrin Genes in the Mouse Striatum. *Front Psychiatry*. 9:257.

43. Larsen JH, Jensen IS, Svenningsen P (2024): Benchmarking transcriptome deconvolution methods for estimating tissue- and cell-type-specific extracellular vesicle abundances. *J Extracell Vesicles*. 13:e12511.

44. Hosseinzadeh Sahafi O, Sardari M, Alijanpour S, Rezayof A (2023): Shared Mechanisms of GABAergic and Opioidergic Transmission Regulate Corticolimbic Reward Systems and Cognitive Aspects of Motivational Behaviors. *Brain Sci*. 13.

45. Smith SM, Ranjan K, Hoover BM, Drayson OGG, Acharya MM, Kramar EA, et al. (2024): Extracellular vesicles from GABAergic but not glutamatergic neurons protect against neurological dysfunction following cranial irradiation. *Sci Rep*. 14:12274.

46. Johnson ECB, Bian S, Haque RU, Carter EK, Watson CM, Gordon BA, et al. (2023): Cerebrospinal fluid proteomics define the natural history of autosomal dominant Alzheimer's disease. *Nat Med*. 29:1979–1988.

47. Li L, Cao J, Zhang S, Wang C, Wang J, Song G, et al. (2014): NCAM signaling mediates the effects of GDNF on chronic morphine-induced neuroadaptations. *J Mol Neurosci*. 53:580–589.

48. Kshatri AS, Gonzalez-Hernandez A, Giraldez T (2018): Physiological Roles and Therapeutic Potential of Ca(2+) Activated Potassium Channels in the Nervous System. *Front Mol Neurosci*. 11:258.

49. Cassano T, Giamogante F, Calcagnini S, Romano A, Lavecchia AM, Inglese F, et al. (2023): PDIA3 Expression Is Altered in the Limbic Brain Regions of Triple-Transgenic Mouse Model of Alzheimer's Disease. *Int J Mol Sci*. 24.

50.		Mishima T, Fujiwara T, Kofuji T, Saito A, Terao Y, Akagawa K (2021): Syntaxin 1B regulates synaptic GABA release and extracellular GABA concentration, and is associated with temperature-dependent seizures. *J Neurochem*. 156:604–613.

51.		Enogieru AB, Omoruyi SI, Hiss DC, Ekpo OE (2019): GRP78/BIP/HSPA5 as a Therapeutic Target in Models of Parkinson's Disease: A Mini Review. *Adv Pharmacol Sci*. 2019:2706783.

52.		Oh HS, Urey DY, Karlsson L, Zhu Z, Shen Y, Farinas A, et al. (2025): A cerebrospinal fluid synaptic protein biomarker for prediction of cognitive resilience versus decline in Alzheimer's disease. *Nat Med*. 31:1592–1603.

53.		Venkatesan S, Binko MA, Mielnik CA, Ramsey AJ, Lambe EK (2023): Deficits in integrative NMDA receptors caused by Grin1 disruption can be rescued in adulthood. *Neuropsychopharmacology*. 48:1742–1751.

54.		Sil S, Singh S, Chemparathy DT, Chivero ET, Gordon L, Buch S (2021): Astrocytes & Astrocyte derived Extracellular Vesicles in Morphine Induced Amyloidopathy: Implications for Cognitive Deficits in Opiate Abusers. *Aging Dis*. 12:1389–1408.

55.		Mimaki M, Wang X, McKenzie M, Thorburn DR, Ryan MT (2012): Understanding mitochondrial complex I assembly in health and disease. *Biochim Biophys Acta*. 1817:851–862.

56.		Mick DU, Fox TD, Rehling P (2011): Inventory control: cytochrome c oxidase assembly regulates mitochondrial translation. *Nat Rev Mol Cell Biol*. 12:14–20.

57.		Perez-Perez R, Lobo-Jarne T, Milenkovic D, Mourier A, Bratic A, Garcia-Bartolome A, et al. (2016): COX7A2L Is a Mitochondrial Complex III Binding Protein that Stabilizes the III2+IV Supercomplex without Affecting Respirasome Formation. *Cell Rep*. 16:2387–2398.

58.		Sun E, Torices S, Osborne OM, Toborek M (2025): Microvascular Dysfunction, Mitochondrial Reprogramming, and Inflammasome Activation as Critical Regulators of Ischemic Stroke Severity Induced by Chronic Exposure to Prescription Opioids. *J Neurosci*. 45.

59.		Yao PJ, Eren E, Goetzl EJ, Kapogiannis D (2021): Mitochondrial Electron Transport Chain Protein Abnormalities Detected in Plasma Extracellular Vesicles in Alzheimer's Disease. *Biomedicines*. 9.

60.		Picca A, Guerra F, Calvani R, Marini F, Biancolillo A, Landi G, et al. (2020): Mitochondrial Signatures in Circulating Extracellular Vesicles of Older Adults with Parkinson's Disease: Results from the EXosomes in PArkiNson's Disease (EXPAND) Study. *J Clin Med*. 9.

61.		Piers TM, Kim DH, Kim BC, Regan P, Whitcomb DJ, Cho K (2012): Translational Concepts of mGluR5 in Synaptic Diseases of the Brain. *Front Pharmacol*. 3:199.

62.		Cleva RM, Olive MF (2011): Positive allosteric modulators of type 5 metabotropic glutamate receptors (mGluR5) and their therapeutic potential for the treatment of CNS disorders. *Molecules*. 16:2097–2106.

63.		Yang T, Zhang D, Huang H, Liu F, Wu J, Ma X, et al. (2025): Astrocytic mGluR5-dependent calcium hyperactivity promotes amyloid-beta pathology and cognitive impairment. *Brain*.

64.		Sanchez-Melgar A, Albasanz JL, Grinan-Ferre C, Pallas M, Martin M (2020): Adenosine and Metabotropic Glutamate Receptors Are Present in Blood Serum and Exosomes from SAMP8 Mice: Modulation by Aging and Resveratrol. *Cells*. 9.

65.		Chen Y, Zhao Y, Yin Y, Jia X, Mao L (2021): Mechanism of cargo sorting into small extracellular vesicles. *Bioengineered*. 12:8186–8201.

66.   Caputi FF, Rullo L, Stamatakos S, Candeletti S, Romualdi P (2019): Interplay between the Endogenous Opioid System and Proteasome Complex: Beyond Signaling. *Int J Mol Sci*. 20.

67.   Song KY, Choi HS, Law PY, Wei LN, Loh HH (2017): Post-Transcriptional Regulation of the Human Mu-Opioid Receptor (MOR) by Morphine-Induced RNA Binding Proteins hnRNP K and PCBP1. *J Cell Physiol*. 232:576–584.

68.   Buccitelli C, Selbach M (2020): mRNAs, proteins and the emerging principles of gene expression control. *Nat Rev Genet*. 21:630–644.

69.   Liu Y, Beyer A, Aebersold R (2016): On the Dependency of Cellular Protein Levels on mRNA Abundance. *Cell*. 165:535–550.

70.   Lindhout FW, Kooistra R, Portegies S, Herstel LJ, Stucchi R, Snoek BL, et al. (2020): Quantitative mapping of transcriptome and proteome dynamics during polarization of human iPSC-derived neurons. *Elife*. 9.

71.   Liu D, Zhou Y, Peng Y, Su P, Li Z, Xu Q, et al. (2018): Endoplasmic Reticulum Stress in Spinal Cord Contributes to the Development of Morphine Tolerance. *Front Mol Neurosci*. 11:72.

72.   Looschen K, Jeffers A, Chakraborty S, Salisbury C, Dodge A, Hochstetler M, et al. (2025): Adolescent stress avoidance influences cue-induced heroin seeking and chaperonin gene expression in the dorsal striatum of adult female rats. *Am J Drug Alcohol Abuse*.1–9.

73.   Lefevre EM, Pisansky MT, Toddes C, Baruffaldi F, Pravetoni M, Tian L, et al. (2020): Interruption of continuous opioid exposure exacerbates drug-evoked adaptations in the mesolimbic dopamine system. *Neuropsychopharmacology*. 45:1781–1792.

74.   Ammon S, Mayer P, Riechert U, Tischmeyer H, Hollt V (2003): Microarray analysis of genes expressed in the frontal cortex of rats chronically treated with morphine and after naloxone precipitated withdrawal. *Brain Res Mol Brain Res*. 112:113–125.

75.   Zou C, Mifflin L, Hu Z, Zhang T, Shan B, Wang H, et al. (2020): Reduction of mNAT1/hNAT2 Contributes to Cerebral Endothelial Necroptosis and Abeta Accumulation in Alzheimer's Disease. *Cell Rep*. 33:108447.

76.   Bagaria J, Bagyinszky E, An SSA (2022): Genetics, Functions, and Clinical Impact of Presenilin-1 (PSEN1) Gene. *Int J Mol Sci*. 23.

77.   Prikas E, Paric E, Asih PR, Stefanoska K, Stefen H, Fath T, et al. (2022): Tau target identification reveals NSF-dependent effects on AMPA receptor trafficking and memory formation. *EMBO J*. 41:e10242.

78.   Ni H, Xu M, Zhan GL, Fan Y, Zhou H, Jiang HY, et al. (2018): The GWAS Risk Genes for Depression May Be Actively Involved in Alzheimer's Disease. *J Alzheimers Dis*. 64:1149–1161.

79.   Oliva CA, Vargas JY, Inestrosa NC (2013): Wnts in adult brain: from synaptic plasticity to cognitive deficiencies. *Front Cell Neurosci*. 7:224.

80.   Robison AJ, Vialou V, Mazei-Robison M, Feng J, Kourrich S, Collins M, et al. (2013): Behavioral and structural responses to chronic cocaine require a feedforward loop involving DeltaFosB and calcium/calmodulin-dependent protein kinase II in the nucleus accumbens shell. *J Neurosci*. 33:4295–4307.

81.   DePoy LM, Gourley SL (2015): Synaptic Cytoskeletal Plasticity in the Prefrontal Cortex Following Psychostimulant Exposure. *Traffic*. 16:919–940.

82.   Shepherd JD, Rumbaugh G, Wu J, Chowdhury S, Plath N, Kuhl D, et al. (2006): Arc/Arg3.1 mediates homeostatic synaptic scaling of AMPA receptors. *Neuron*. 52:475–484.

83.   Zhang H, Bramham CR (2021): Arc/Arg3.1 function in long-term synaptic plasticity: Emerging mechanisms and unresolved issues. *Eur J Neurosci*. 54:6696–6712.

84.   Sullivan KR, Ravens A, Walker AC, Shepherd JD (2025): "Arc - A viral vector of memory and synaptic plasticity". *Curr Opin Neurobiol*. 91:102979.

Figure 1. Characterization of BDEVs from rat prefrontal cortex.

(A) Experimental design. (B) Nanoparticle tracking analysis (NTA) showing no significant changes in concentration or size distribution of isolated BDEVs between groups.(C) Representative electron micrographs of EVs. (D) Western blot analysis confirming the presence of EV markers Flotillin and HSP70, and absence of the exclusion marker GM130 in the EV samples.

Figure 2. Effect of morphine treatment on the RNA profile of BDEVs

(A) Volcano plot of differentially expressed mRNAs in BDEVs from morphine-treated versus control animals. (B) KEGG pathway enrichment analysis of morphine-induced RNA alterations. (C) Dot plot showing upregulated and downregulated transcripts within enriched pathways. (D) Heatmap of significantly dysregulated mRNAs across enriched pathways (log₂ fold-change). (E) Network visualization of enriched pathways and associated transcripts. (F) Integration with a human GWAS for opioid use disorder highlights risk-associated genes differentially expressed in BDEVs. (G) RNA deconvolution estimates brain cell-type composition, with nearly 40% of transcripts inferred from GABAergic interneuron-derived BDEVs.

Figure 3. Effect of morphine treatment on the protein profile of BDEVs

(A) Volcano plot showing differentially expressed proteins in BDEVs from morphine-treated versus control animals. (B) Dynamic range of protein abundance (left) and MA plot (right), highlighting significantly dysregulated proteins. (C) KEGG pathway enrichment analysis identifying pathways associated with morphine-induced protein alterations. (D) Network visualization of enriched pathways and associated proteins. Gray nodes represent KEGG pathways; colored nodes represent proteins, with node color indicating log₂ fold-change (red = upregulated, blue = downregulated). (E) Heatmap of log₂ fold-change values for dysregulated proteins across enriched KEGG pathways.

Figure 4. Integrated transcriptomic and proteomic analysis of BDEVs from morphine-treated rats.

(A) Circos plot linking significantly dysregulated mRNAs (left) and proteins (right). Node colors represent log□ fold-change (red/orange = upregulated, blue = downregulated). (B) UpSet plot showing the overlap between significantly altered RNAs and proteins; bar height represents the number of molecules in each set. (C) Comparative KEGG pathway enrichment analysis of dysregulated mRNAs and proteins. Bubble size reflects the number of molecules per pathway, and color indicates −log□□(p). (D) Chord diagram of the protein processing in endoplasmic reticulum pathway, depicting concordance and divergence between RNA and protein levels. Ribbons connect matched RNA–protein pairs, with ribbon color indicating log□ fold-change.

Figure 5. Effect of morphine-derived BDEVs on primary neuronal cultures.
(A) Experimental design: BDEVs from morphine- and vehicle-treated rats were applied for 24 h to primary cortical neurons, followed by NanoString analysis. (B) Volcano plot showing differentially expressed transcripts in neurons exposed to morphine-derived BDEVs; selected significant genes are labeled. (C) Reactome pathway enrichment of dysregulated genes, highlighting Ca²□, WNT, opioid, and DARP 32-related signaling pathways.

(A)

(B)

(C)