Flexible Self-Protection as Evidence of Pain-Like States in House Crickets

Oscar Manzi[1,] Kate E. Lynch[2], Daniel M. Allman[1], Tanya Latty[1], Thomas E. White[1*]

[1]School of Life and Environmental Sciences, The University of Sydney, Camperdown, NSW, Australia.

[2]School of History and Philosophy of Science, The University of Sydney, Camperdown, NSW, Australia.

*Corresponding author: thomas.white@sydney.edu.au

Word count (excluding references): 3605

Number of figures: 2

Number of tables: 1

Keywords: *Acheta domesticus,* insect welfare, nociception, sentience, affect, entomophagy, precautionary principle

Abstract

The possibility that insects experience pain is a frontier question at the intersection of animal behaviour, comparative cognition, and philosophy of mind. Interest has been fuelled not only by anatomical discoveries but also by expanding behavioural and comparative evidence. Under leading frameworks, the most informative behavioural indicators of pain-like experience include flexible, targeted responses to harm that suggest more than stereotyped, reflexive withdrawal. Here we tested for such responses in the house cricket (*Acheta domesticus*), a species of both evolutionary and commercial importance. Using a fully blinded, within-subjects design, we applied either noxious heat, tactile contact, or no contact to a single antenna under both lower- and higher-stress environmental conditions and recorded subsequent grooming behaviour. Crickets were significantly more likely to groom the noxiously stimulated antenna, and did so for longer durations, than under control or tactile treatments. Grooming also showed a distinct temporal profile, with elevated activity sustained across the early observation period. Environmental condition and sex had no effect, indicating that self-protective grooming was expressed consistently across contexts and sexes. These findings provide robust evidence of flexible, site-directed self-protection in Orthoptera, addressing a key gap in the evidence base for pain-like states outside vertebrates. Such evidence strengthens the case for precautionary consideration of insect welfare, while also bearing on the broader scientific question of how felt experience is distributed across the animal kingdom.

Introduction

Animals navigate a world of shifting threats and fleeting opportunities, where thriving depends not just on reacting, but on evaluating. Sentience is commonly defined as the capacity for subjective, valenced experience, and is increasingly recognised as a core, widely-distributed biological capacity that shapes how animals respond to the world (Low et al., 2012). Under adaptive accounts, it is variously hypothesised to support deep forms of learning (Ginsburg & Jablonka, 2019), inform complex decision-making (Veit, 2023), and enable flexible, goal-directed behaviour (Elwood, 2011). While its neural substrates likely differ widely across taxa, its functional signature—experience-guided evaluation of internal and external conditions—appears broadly conserved (Birch et al., 2021; Brown, 2015; Gibbons, Crump, et al., 2022). In both humans and non-human animals, felt experiences are suggested to enable responses that go beyond 'mere reflex' or stereotyped output into behaviours marked by complexity, context sensitivity, and/or specificity (Ginsburg & Jablonka, 2019; Veit, 2023). Yet because experience is inherently private its presence is difficult to infer, especially in invertebrates where brains lack close homology to mammals and behavioural profiles can appear equally unfamiliar, resisting easy comparison to vertebrate models.

Pain presents a critical test case. Distinct from nociception—the unconscious detection of damaging stimuli—pain is typically (though not uniformly) defined as a negatively valenced experience of harm (Raja et al., 2020; Sneddon et al., 2014). Unlike nociceptive reflexes, which can occur without central processing, pain supports persistent behavioural change, lasting avoidance, and context-sensitive self-protection (Elwood, 2019; Sneddon et al., 2014). Nociception is widespread across the animal kingdom and confers clear evolutionary value, but pain is thought to involve additional costs such as the metabolic demands of centralised processing and the risk of behavioural distraction (Bateson, 1991). Such costs imply that pain would not evolve without compensating benefits. From this perspective, pain can be seen as an adaptation that builds upon nociception to enable richer forms of behaviour and learning and, ultimately, support long-term fitness.

Efforts to assess pain and sentience in animals have changed markedly over time. Historically,

the capacity for subjective experience was tightly coupled to anatomical similarity with humans, and

the scope of morally considerable animals was narrowly drawn. That boundary has since shifted

outward, with birds, fish, and more recently cephalopod molluscs and decapod crustaceans recognised

as potentially sentient (Birch et al., 2021; Gibbons, Crump, et al., 2022; Sneddon, 2019). This creeping

frontier reflects a broader shift in thinking: rather than asking which brains are most like ours, a question

with clear failings (Panksepp, 2016), researchers now ask which behavioural and physiological features

justify the inference of experience (Crump et al., 2022; Jablonka & Ginsburg, 2022; Sneddon et al.,

2014). Frameworks such as that proposed by Crump et al. (2022) have codified this shift by outlining

eight empirically testable criteria for pain-like states, including motivational trade-offs, associative

learning, and flexible self-protection. While the framework, understandably, does not provide a formal

method for balancing these criteria, it highlights how different lines of evidence can be assembled to

guide judgement about the likelihood of pain.

In this context insects have emerged as a critical frontier. Once dismissed as too small-brained

or simple to support experience (Eisemann et al., 1984), they are now known as capable of remarkably

complex tasks, including associative learning, context-sensitive decision-making, and cross-modal

sensory integration (Gibbons et al., 2024; Gibbons, Versace, et al., 2022; Giurfa, 2013). Recent studies

have also identified brain regions such as the mushroom bodies and central complex that appear to

support evaluative processing functionally analogous to that seen in vertebrates (Barron & Klein, 2016).

Yet the question of pain in insects cannot be settled by neural architecture alone. Given the diversity of

nervous systems across phyla, and the sheer creative power of adaptive evolution showcased via

multiple realisability, behaviour remains our most direct and inclusive route to inferring experience

(Elwood, 2011). That is, rather than asking whether an animal has the same hardware, the more relevant

question is whether it shows a comparable behavioural profile under similar conditions.

Among insects, much of the extant behavioural evidence concerns bees. Honeybees (*Apis*

mellifera) and bumblebees (*Bombus terrestris*) display hallmarks of pain-like states under the Crump

et al. (2022) framework: they form associative memories linking noxious stimuli to conditioned cues

(Vergoz et al., 2007), regulate foraging based on trade-offs between reward and harm (Gibbons, Versace, et al., 2022), and perform localised self-grooming at injured body parts (Gibbons et al., 2024). These behaviours are persistent, targeted, and sensitive to internal state—features that argue against stereotypy and in favour of evaluative processing. More broadly, there is growing evidence that insects exhibit affective processes beyond pain, ranging from optimism–pessimism biases to play-like behaviour (Perry & Baciadonna, 2017; Triphan et al., 2025). Such findings suggest that insects can generate and regulate a wider suite of valenced states than once assumed. Still, the bulk of evidence remains taxonomically narrow. In a recent review, Gibbons et al. (2022) assessed six major insect orders and found that while Diptera and Blattodea satisfied multiple pain-related criteria, Orthoptera— grasshoppers, locusts, and crickets—remain underrepresented.

This gap is significant. Orthopterans are both ecologically and evolutionarily important, and central to the edible insect industry: *Acheta domesticus*, the house cricket, is among the most widely farmed insects worldwide, with billions reared annually (Rowe et al., 2024). Orthopterans also represent one of the more basal branches of the winged insects, diverging early within the Polyneoptera (Tihelka et al., 2021). Evidence of sophisticated processing in this group therefore bears directly on questions of how widespread the capacity for felt states may be across insects. Once thought too simple to support sophisticated processing, orthopterans have since been shown to possess nociceptive ion channels (Ylla et al., 2021) and evidence of sensory integration in grasshopper brains (Gupta & Stopfer, 2012). Behavioural data, however, remain scarce. Only two studies have directly tested associative learning with noxious stimuli, both indicating that grasshoppers can learn to avoid harm through limb movement (Forman, 1984; Punzo, 1980), while a third demonstrated crickets' ability to learn to use visual cues to seek relief from noxious heat (Wessnitzer et al., 2008). Indirectly related efforts have shown behavioural effects of morphine, which increase individuals' latencies to escape from heated substrates in a dose- and time-dependent manner (Zabala & Gómez, 1991). Given the scale of their use and the ethical stakes of pain, further targeted behavioural tests are urgently needed. Without them, welfare assessments will remain speculative, and our ability to prevent unnecessary suffering limited (Birch et al., 2021; Broom, 2011).

Here we tested a key behavioural criterion for pain in the house cricket *Acheta domesticus*:

flexible self-protection, expressed as persistent and directed grooming of a specific site after noxious

stimulation (Crump et al., 2022). We applied either a heated, tactile, or no stimulus to a single antenna

and measured whether crickets groomed the treated antenna more frequently, for longer, and with a

distinct temporal profile. If pain-like experience underlies protective grooming, then theory predicts

that noxious stimulation should elicit greater, more sustained, and more targeted grooming than either

tactile or control conditions.

Methods

Study System

We used commercially reared *Acheta domesticus* (house crickets), sourced from Petstock

(www.petstock.com.au), and housed them in a laboratory at The University of Sydney, Camperdown,

Sydney. We housed crickets in shared containers (40 x 40 x 100 cm) under a 12:12 hour light–dark

cycle at 23 ± 1 °C and provided them with free access to wheatgerm and peaches in juice for food and

hydration. We conducted all trials under artificial lighting between 10:00 and 15:00 maintaining an

ambient light level of approximately 1000 and 100 lux in the higher and lower stress test environments,

respectively (detailed below).

We designed this experiment to test whether crickets exhibit flexible, site-specific self-

protective behaviour consistent with an evaluative response to noxious stimuli. Our design was broadly

inspired by Gibbons et al. (2024), who tested self-directed grooming responses in bumblebees. To

examine how environmental context influences such behaviour, we included an additional manipulation

comparing responses under higher- and lower-stress conditions; an approach motivated by existing

evidence that ambient stressors can modulate the expression of affective behaviour (Bateson, 1991).

 Experimental Treatments

We tested 80 adult crickets (40 males, 40 females) in a fully within-subjects design. We assigned

each individual to one of two housing conditions, higher-stress or lower-stress (N = 40 each), and

exposed them to three treatment conditions in randomised order: a noxious stimulus (heated probe), a

innocuous stimulus (unheated probe), and a control condition involving handling only. The innocuous

and control conditions served as procedural and restraint controls, respectively.

For each trial, we gently immobilised the cricket using a sponge and contacted a randomly

chosen antenna, with side not predetermined or counter-balanced across trials. In the noxious condition,

we applied a soldering iron (Hakko FX-888D, 6 mm chisel tip) heated to 65 °C for five seconds. In the

innocuous condition we applied the same probe without heating. While in the control condition we

handled the cricket in the same manner but did not apply any stimulus. We selected the temperature of

65 °C in our noxious treatment to provide a stimulus hot enough to be noxious, while avoiding lasting

tissue damage. This set point also reflects that used in a comparable test with bumblebees, where it

elicited robust avoidance without causing visible injury (Gibbons et al. 2024). We confirmed probe

temperature prior to each application using a digital infrared thermometer (Sovarcate HS960D). After

the application of a given treatment, we immediately transferred the cricket into the observation arena.

We constructed two environmental contexts to coarsely manipulate background stress levels.

In the 'higher-stress' condition, we placed crickets into open-top plastic arenas (30 × 22 × 10.5 cm)

wrapped in red cellophane on the base and sides. This design removed cover, limited substrate for

digging, and supplied strong illumination, which elevates vigilance and stress in orthopterans (Hedrick,

2000). In the 'lower-stress' condition, we used closed-top arenas with a sand substrate and reduced

ambient light by enclosing them in a brown paper shelter (Fig. 1), thereby providing structural refuge

and a more naturalistic, darkened environment. We balanced sex and treatment order across

environmental conditions. We filmed all behavioural trials for ten minutes using a Canon 1500D camera

fit with a Canon 18-55 mm lens (fixed at 18 mm) at 30 fps, and allowed a ten-minute intertrial interval

during which we returned each cricket to its individual holding container.

Behavioural analysis

Three treatment-blind observers coded all grooming behaviour from video recordings using predefined criteria. We defined grooming as either a "touch" (when a cricket brushed the antenna using the prothoracic leg) or "to mouth" (when it drew the antenna toward the mandibles using the same leg), and we recorded which of their two antenna was groomed. Observers recorded timestamps for the start and end of each grooming bout to calculate both frequency and duration.

Statistical analysis

We ran four analyses to test our core predictions. First, we asked whether crickets allocated a greater proportion of grooming time to the antenna that had been stimulated. Second, we tested whether they were more likely to groom the stimulated antenna at all. Third, we asked whether grooming of the stimulated antenna lasted longer under noxious conditions than under control or innocuous contact. Fourth, we examined how the temporal profile of grooming evolved across the trial under different treatment conditions.

To test whether crickets preferentially directed grooming to the treated antenna within trials, we fit a beta regression model with the proportion of total antennal grooming duration allocated to the focal side as the response. We included treatment (noxious, innocuous, control), environmental condition (higher-stress vs. lower-stress), and sex as fixed effects, and included individual ID as a random intercept. To satisfy model assumptions, we rescaled proportions to fall strictly within the $(0,1)$ interval. We initially tested a more complex model with a treatment $\times$ environment interaction, but this did not improve model fit ($\chi^2 = 0.59$, df = 2, p = 0.744), so we retained the simpler additive model to maximise power. We then used one-sided post hoc tests on the estimated marginal means to assess whether grooming under each treatment condition was significantly biased toward the stimulated side. We report means and confidence intervals on the response scale.

To test the probability of any grooming being directed toward the treated antenna, we used a

binomial GLMM. The binary response was coded as 1 if the focal antenna was groomed and 0

otherwise. Fixed effects included treatment, environment, and sex, with individual ID as a random

effect. A full model including the treatment × environment interaction again showed no improvement

in fit ($\chi^2 = 1.22$, df = 2, p = 0.544) and was simplified accordingly.

To test the duration of directed grooming, we used a Tweedie GLMM with a log link, which

allowed us to model the right-skewed data with many zero values. The model included treatment,

environmental condition, and sex as fixed effects, with individual ID as a random intercept. The more

complex model including the treatment × environment interaction did not improve fit ($\chi^2 = 0.37$, df =

2, p = 0.830) and was likewise simplified.

Finally, to explore the temporal dynamics of grooming, we fitted generalised additive models

(GAMs) to the durations of focal grooming bouts over the ten-minute trial. Each model included

treatment as a parametric term, treatment-specific smooths for time since stimulus, and a random

intercept for individual identity. We used a Tweedie distribution with a log link to accommodate zero-

inflated, right-skewed bout durations. Predictions were generated marginal over individuals by

excluding the random-effect term, and we back-transformed estimates to the response scale with 95%

confidence intervals derived from the standard errors. These models allowed us to visualise whether

grooming in response to noxious stimulation was transient or sustained across time, and whether its

temporal profile differed significantly from control and tactile treatments.

We conducted all statistical analyses in RStudio (v4.3.2; RStudio Team, 2024), using the

packages lme4 for mixed models (Scheipl, 2011), glmmTMB for Tweedie models (Brooks et al., 2017),

mgcv for GAMs (Wood, 2018), DHARMa for residual diagnostics (Hartig, 2022), and emmeans for

post hoc comparisons (Russell et al., 2018). We adopted the conventional threshold of statistical

significance at $\alpha = 0.05$, and report all summary statistics with their standard errors.

 Ethical note

Although invertebrate research is not subject to formal ethics approval in Australia, we followed

high standards of care throughout. We kept crickets in stable environmental conditions, provided

continuous access to food and water, and ensured that no procedures caused injury or lasting harm.

After completing the study, we returned all individuals to their housing containers, where they lived out

their natural lifespans under continued laboratory care with free access to food and water.

Data and code availability

All underlying data and code are freely available via GitHub (https://github.com/SEEC-

[lab/cricket_protection](https://github.com/SEEC-lab/cricket_protection)) and will be persistently archived upon acceptance.

241 Results
242

Specificity of grooming

Within trials, crickets allocated a significantly greater proportion of antennal grooming to the

stimulated side following noxious treatment compared to control ($\beta = 0.62 \pm 0.29$, $z = 2.14$, $p = 0.032$;

Fig. 2a; Table 1). On average, 59% (95% CI: 0.505–0.674) of grooming time was directed to the focal

antenna after noxious stimulation, compared with 44% (0.337–0.548) in control trials and 52% (0.418–

0.615) in innocuous contact trials. A one-sided test against chance confirmed that the noxious

proportion was significantly greater than 0.5 ($z = 2.07$, $p = 0.019$), while control and innocuous

treatments did not differ from chance. There was no effect of environmental condition ($\beta = -0.14 \pm 0.23$,

$z = -0.62$, $p = 0.538$) or sex ($\beta = -0.37 \pm 0.24$, $z = -1.56$, $p = 0.118$).

 Directed-grooming probability

Crickets were significantly more likely to groom the antenna that received the noxious stimulus

compared to the control (log-odds $= 1.52 \pm 0.38$, $z = 3.96$, $p < 0.001$), with grooming also elevated—but

not significantly so—under the innocuous treatment (log-odds $= 0.69 \pm 0.36$, $z = 1.91$, $p = 0.056$). On

the response scale, we estimated that grooming occurred on the focal side in 63% of noxious trials (95%

CI: 0.50–0.74), 43% of innocuous trials (0.31–0.55), and 27% of control trials (0.18–0.40). Post hoc

comparisons confirmed a robust noxious–vs-control difference (Δlog-odds $= -1.52 \pm 0.38$, $z = -3.96$,

$p < 0.001$), with weaker support for a noxious–vs-innocuous difference (Δlog-odds $= -0.83 \pm 0.36$,

$z = -2.34$, $p = 0.051$) and no significant difference between innocuous and control (Δlog-

odds $= -0.69 \pm 0.36$, $z = -1.91$, $p = 0.136$). While there was a trend toward increased grooming under

the higher-stress condition ($\beta = 0.66 \pm 0.34$, $p = 0.051$), this was not statistically significant,

Directed-grooming duration

Crickets groomed for significantly longer following noxious stimulation than under control (*est.*

$= 1.39 \pm 0.27$, $z = 5.15$, $p < 0.001$) or innocuous (est.$= 0.60 \pm 0.30$, $z = 2.04$, $p = 0.042$) treatment

conditions (Fig. 2c; Table 1). The average estimated grooming duration was 13.48 ± 2.39 s after noxious

treatment, compared to 6.12 ± 1.32 s after innocuous and 3.35 ± 0.82 s in the control. Post hoc pairwise

comparisons confirmed a significant increase in grooming duration under noxious treatment compared

to control (ratio $= 0.25 \pm 0.07$, $z = -5.15$, $p < 0.001$) and innocuous (ratio $= 0.45 \pm 0.11$, $z = -3.29$,

$p = 0.0029$). Grooming duration did not differ significantly between innocuous and control treatments

(ratio $= 0.55 \pm 0.16$, $z = -2.04$, $p = 0.104$). We detected no effect of environmental condition

(*est.* $= 0.10 \pm 0.24$, $z = 0.43$, $p = 0.67$) or sex (est. $= 0.11 \pm 0.24$, $z = 0.46$, $p = 0.64$) on grooming

duration.

 Temporal dynamics of directed-grooming

We found that including treatment-specific smooth terms significantly improved model fit

($F = 2.64$, $df = 9.8$, residual $df = 405$, $p = 0.036$), confirming that the duration of antennal grooming

varied meaningfully over time in a treatment-specific manner. In plain terms, we detected clear

treatment effects on the timecourse of directed grooming: the noxious stimulus led to sustained, elevated

grooming that declined across the observation period; innocuous stimulation produced a delayed peak

in grooming around minute 8; and control responses remained stable and low throughout (Fig. 2d).

Smooth terms were statistically significant for both noxious ($edf = 1.00$, $F = 6.996$, $p = 0.009$)

and innocuous ($edf = 3.39$, $F = 2.813$, $p = 0.024$) treatments, indicating reliable temporal variation in

grooming under those conditions. In contrast, grooming under control conditions showed no significant

change over time ($edf = 1.00$, $F = 0.003$, $p = 0.954$), consistent with a flat response.

Together, these results support our prediction that noxious stimulation elicits both more

frequent and more persistent grooming — a behavioural profile suggestive of flexible, experience-

guided self-protection. The innocuous treatment also produced an extended grooming response, though

with a delayed and less consistent timecourse, potentially indicating residual mechanical irritation or

procedural artefacts.

Discussion

Pain remains one of the most elusive and consequential frontiers in animal cognition, and

insects provide a particularly demanding test case. These animals display complex behavioural

repertoires, yet their subjective capacities remain uncertain. While anatomical and physiological

findings have renewed interest in insect sentience (Barron & Klein, 2016; Crump et al., 2022; Gibbons,

Crump, et al., 2022), it is behavioural evidence—especially flexible, site-specific responses to harm—

that offer an essential route to inference. Frameworks such as that of Crump et al. (2022) emphasise

patterns of persistence, context sensitivity, and goal-directedness as criteria distinguishing pain-like

states from 'mere' reflexive nociception. Our study provides new evidence bearing on this question,

demonstrating site-directed self-protection in *Acheta domesticus*, a species of both evolutionary and

commercial importance.

Crickets in our experiment groomed a noxiously stimulated antenna more frequently and for

longer than under tactile or no-contact control conditions. The response was both site-specific and

persistent, suggesting that crickets monitor injury location and adjust their behaviour in ways not

reducible to simple reflexes. Grooming intensity followed a clear temporal trajectory (Fig. 2c):

noxiously stimulated individuals showed an elevated, sustained phase that gradually declined, a pattern

reminiscent of findings in bees and rodents (Gibbons et al., 2024; Mittal et al., 2009). By contrast, the

innocuous contact treatment produced an early dip in grooming before a later rise. One possibility is

that sudden, non-noxious touch was initially treated as a potential predatory threat, eliciting vigilance

or an orienting response that temporarily displaced grooming (Zacarias et al., 2018). Alternatively, the

tactile stimulus may have prompted competing behaviours such as scanning or locomotion, which

delayed antennal maintenance until the salience of the stimulus subsided (Oram & Card, 2022). Similar

context-dependent processing has been demonstrated in crickets: when antennae were contacted with a

glass rod, a hunting spider leg, or an orb-weaver leg, responses varied from aggression to aversion to

search behaviour, with glass-rod contact especially likely to elicit search or no response (Okada &

Akamine, 2012). This parallel suggests that our probe, like the glass rod, was interpreted flexibly rather

than as a simple mechanical trigger. Overall, the observed patterns align with the criterion of flexible

self-protection and add Orthoptera to the short list of insect orders for which such evidence exists

(Gibbons, Crump, et al., 2022).

We also manipulated environmental stress to test its influence on behavioural expression, given

prior evidence that stress can suppress affectively mediated responses (Perry & Baciadonna, 2017).

Grooming likelihood and duration did not differ significantly between 'higher-' and 'lower-stress'

contexts, and we likewise found no effect of sex in any model. This may indicate that our environmental

manipulation lacked sufficient salience, and/or that self-protective responses are robust to background

stress and consistent across sexes. Either way, the persistent expression of grooming across

environmental and sex conditions supports the view that these behaviours are internally regulated and

functionally significant, rather than fragile or reflexively triggered.

Taken together, our findings further erode the once-dominant view of insects as behaviourally

rigid and insentient. Crickets have already been shown to possess nociceptors (Ylla et al., 2021),

centralised sensory integration (Gupta & Stopfer, 2012), and have the capacity to learn from aversive events (Forman, 1984; Punzo, 1980), and moderate their responses to noxious stimuli following analgesia (Zabala & Gómez, 1991). By demonstrating flexible self-protection, we add a fifth pillar to this growing body of evidence in Orthoptera and strengthen coverage of the core criteria for pain-like states outlined by Crump et al. (2022). Together, these lines of evidence suggest that orthopterans exhibit the same conjunction of features—nociception, integrative processing, learning, and targeted self-protection—that are taken to warrant serious consideration of sentience. While this does not resolve the question of insect pain, it meaningfully narrows the evidential gap and underscores the need for further comparative and taxonomically broad investigations.

The implications of this work extend beyond theory. *Acheta domesticus* is reared by the millions for food, feed, and research, often under conditions that assume an absence of felt experience. If insects can respond to injury in a way consistent with pain—as our findings suggest—then the ethical landscape shifts. Under a precautionary approach (Birch, 2024), the burden moves toward reducing harm in farming, handling, and experimentation. This shift is not only moral, it reflects the accelerating recognition, across fields, that sentience may be evolutionarily deep and taxonomically widespread.

The expanding frontier of behavioural evidence is not just pushing disciplinary boundaries, it is redrawing ethical ones. As more insect taxa exhibit traits once thought exclusive to vertebrates, the phylogenetic perimeter of likely sentience continues to creep. Our findings join a broader movement toward reassessing long-held assumptions, urging not premature conclusions but a recalibration of care in the face of uncertainty; one that may prove transformative for how we regard the smallest of animals.

Acknowledgements

This work was supported by the Australia and Pacific Science Foundation (APSF240014).

References

Barron, A. B., & Klein, C. (2016). What insects can tell us about the origins of consciousness. *Proceedings of the National Academy of Sciences*, *113*(18), 4900–4908. https://doi.org/10.1073/pnas.1520084113

Bateson, P. (1991). Assessment of pain in animals. *Animal Behaviour*, *42*(5), 827–839. https://doi.org/10.1016/S0003-3472(05)80127-7

Birch, J. (2024). *The edge of sentience: Risk and precaution in humans, other animals, and AI*. Oxford University Press.

Birch, J., Burn, C., Schnell, A., Browning, H., & Crump, A. (2021). Review of the Evidence of Sentience in Cephalopod Molluscs and Decapod Crustaceans. *General - Animal Feeling*. https://www.wellbeingintlstudiesrepository.org/af_gen/2

Brooks, M., E., Kristensen, K., Benthem, K., J. ,van, Magnusson, A., Berg, C., W., Nielsen, A., Skaug, H., J., Mächler, M., & Bolker, B., M. (2017). glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. *The R Journal*, *9*(2), 378. https://doi.org/10.32614/RJ-2017-066

Broom, D. M. (2011). A History of Animal Welfare Science. *Acta Biotheoretica*, *59*(2), 121–137. https://doi.org/10.1007/s10441-011-9123-3

Brown, C. (2015). Fish intelligence, sentience and ethics. *Animal Cognition*, *18*(1), 1–17. https://doi.org/10.1007/s10071-014-0761-0

Crump, A., Browning, H., Schnell, A., Burn, C., & Birch, J. (2022). Sentience in decapod crustaceans: A general framework and review of the evidence. *Animal Sentience*, *7*(32). https://doi.org/10.51291/2377-7478.1691

Eisemann, C. H., Jorgensen, W., Merritt, D., Rice, M., Cribb, B., Webb, P., & Zalucki, M. (1984). Do insects feel pain? A biological view. *Experientia*, *40*(2), 164–167.

Elwood, R. W. (2011). Pain and Suffering in Invertebrates? *ILAR Journal*, *52*(2), 175–184. https://doi.org/10.1093/ilar.52.2.175

Elwood, R. W. (2019). Discrimination between nociceptive reflexes and more complex responses

consistent with pain in crustaceans. *Philosophical Transactions of the Royal Society B:*

Biological Sciences, *374*(1785), 20190368. https://doi.org/10.1098/rstb.2019.0368

Forman, R. R. (1984). Leg position learning by an insect. I. A heat avoidance learning paradigm.

Journal of Neurobiology, *15*(2), 127–140.

Gibbons, M., Crump, A., Barrett, M., Sarlak, S., Birch, J., & Chittka, L. (2022). Can insects feel pain?

A review of the neural and behavioural evidence. In *Advances in Insect Physiology* (Vol. 63,

pp. 155–229). Elsevier. https://doi.org/10.1016/bs.aiip.2022.10.001

Gibbons, M., Pasquini, E., Kowalewska, A., Read, E., Gibson, S., Crump, A., Solvi, C., Versace, E., &

Chittka, L. (2024). Noxious stimulation induces self-protective behavior in bumblebees.

iScience, 110440. https://doi.org/10.1016/j.isci.2024.110440

Gibbons, M., Versace, E., Crump, A., Baran, B., & Chittka, L. (2022). Motivational trade-offs and

modulation of nociception in bumblebees. *Proceedings of the National Academy of Sciences*

of the United States of America, *119*(31), e2205821119.

https://doi.org/10.1073/pnas.2205821119

Ginsburg, S., & Jablonka, E. (2019). *The evolution of the sensitive soul: Learning and the origins of*

consciousness. MIT press.

Giurfa, M. (2013). Cognition with few neurons: Higher-order learning in insects. *Trends in*

Neurosciences, *36*(5), 285–294. https://doi.org/10.1016/j.tins.2012.12.011

Gupta, N., & Stopfer, M. (2012). Functional analysis of a higher olfactory center, the lateral horn.

Journal of Neuroscience, *32*(24), 8138–8148.

Hartig, F. (2022). *DHARMa: Residual diagnostics for hierarchical (multi-level / mixed) regression*

models [Manual]. https://CRAN.R-project.org/package=DHARMa

Hedrick, A. V. (2000). Crickets with extravagant mating songs compensate for predation risk with

extra caution. *Proceedings of the Royal Society of London. Series B: Biological Sciences*,

267(1444), 671–675. https://doi.org/10.1098/rspb.2000.1054

Jablonka, E., & Ginsburg, S. (2022). Pain sentience criteria and their grading. *Animal Sentience*,

7(32). https://doi.org/10.51291/2377-7478.1713

Low, P., Panksepp, J., Reiss, D., Edelman, D., Van Swinderen, B., & Koch, C. (2012). The Cambridge

declaration on consciousness. *Francis Crick Memorial Conference*, *7*.

Mittal, N., Joshi, R., Hota, D., & Chakrabarti, A. (2009). Evaluation of antihyperalgesic effect of

curcumin on formalin-induced orofacial pain in rat. *Phytotherapy Research*, *23*(4), 507–512.

https://doi.org/10.1002/ptr.2662

Okada, J., & Akamine, S. (2012). Behavioral response to antennal tactile stimulation in the field

cricket Gryllus bimaculatus. *Journal of Comparative Physiology A: Neuroethology, Sensory,*

Neural, and Behavioral Physiology, *198*(7), 557–565. https://doi.org/10.1007/s00359-012-

0729-y

Oram, T. B., & Card, G. M. (2022). Context-dependent control of behavior in Drosophila. *Current*

Opinion in Neurobiology, *73*, 102523. https://doi.org/10.1016/j.conb.2022.02.003

Panksepp, J. (2016). Brain processes for "good" and "bad" feelings: How far back in evolution?

Animal Sentience, *1*(3). https://doi.org/10.51291/2377-7478.1050

Perry, C. J., & Baciadonna, L. (2017). Studying emotion in invertebrates: What has been done, what

can be measured and what they can provide. *Journal of Experimental Biology*, *220*(21),

3856–3868. https://doi.org/10.1242/jeb.151308

Punzo, F. (1980). Neurochemical changes associated with learning in Schistocerca americana

(Orthoptera: Acrididae). *Journal of the Kansas Entomological Society*, 787–796.

Raja, S. N., Carr, D. B., Cohen, M., Finnerup, N. B., Flor, H., Gibson, S., Keefe, F. J., Mogil, J. S.,

Ringkamp, M., Sluka, K. A., & others. (2020). The revised International Association for the

Study of Pain definition of pain: Concepts, challenges, and compromises. *Pain*, *161*(9), 1976–

1982.

Rowe, E., Robles López, K. Y., Robinson, K. M., Baudier, K. M., & Barrett, M. (2024). Farmed

cricket (Acheta domesticus, Gryllus assimilis, and Gryllodes sigillatus; Orthoptera) welfare

considerations: Recommendations for improving global practice. *Journal of Insects as Food*

and Feed, *10*(8), 1253–1311. https://doi.org/10.1163/23524588-00001087

Russell, L., Henrik, S., Jonathon, L., Paul, B., & Maxime, H. (2018). Estimated marginal means, aka

least-squares means. *The American Statistician*, *34*, 216–221.

Scheipl, F. (2011). *amer: Using lme4 to fit generalized additive mixed models*.

Sneddon, L. U. (2019). Evolution of nociception and pain: Evidence from fish models. *Philosophical*

Transactions of the Royal Society B: Biological Sciences, *374*(1785), 20190290.

https://doi.org/10.1098/rstb.2019.0290

Sneddon, L. U., Elwood, R. W., Adamo, S. A., & Leach, M. C. (2014). Defining and assessing animal

pain. *Animal Behaviour*, *97*, 201–212. https://doi.org/10.1016/j.anbehav.2014.09.007

Tihelka, E., Cai, C., Giacomelli, M., Lozano-Fernandez, J., Rota-Stabelli, O., Huang, D., Engel, M.

S., Donoghue, P. C. J., & Pisani, D. (2021). The evolution of insect biodiversity. *Current*

Biology, *31*(19), R1299–R1311. https://doi.org/10.1016/j.cub.2021.08.057

Triphan, T., Ferreira, C. H., & Huetteroth, W. (2025). Play-like behavior exhibited by the vinegar fly

Drosophila melanogaster. *Current Biology*, *35*(5), 1145-1155.e2.

https://doi.org/10.1016/j.cub.2025.01.025

Veit, W. (2023). *A philosophy for the science of animal consciousness*. Routledge.

Vergoz, V., Roussel, E., Sandoz, J.-C., & Giurfa, M. (2007). Aversive Learning in Honeybees

Revealed by the Olfactory Conditioning of the Sting Extension Reflex. *PLoS ONE*, *2*(3),

e288. https://doi.org/10.1371/journal.pone.0000288

Wessnitzer, J., Mangan, M., & Webb, B. (2008). Place memory in crickets. *Proceedings of the Royal*

Society B: Biological Sciences, *275*(1637), 915–921. https://doi.org/10.1098/rspb.2007.1647

Wood, S. (2018). *Mixed GAM computation vehicle with automatic smoothness estimation. R package*

version 1.8–12. Vienna: RStudio.

Ylla, G., Nakamura, T., Itoh, T., Kajitani, R., Toyoda, A., Tomonari, S., Bando, T., Ishimaru, Y.,

Watanabe, T., Fuketa, M., Matsuoka, Y., Barnett, A. A., Noji, S., Mito, T., & Extavour, C. G.

(2021). Insights into the genomic evolution of insects from cricket genomes. *Communications*

Biology, *4*(1), 733. https://doi.org/10.1038/s42003-021-02197-9

Zabala, N. A., & Gómez, M. A. (1991). Morphine analgesia, tolerance and addiction in the cricket

pteronemobius sp. (Orthoptera, insecta). *Pharmacology Biochemistry and Behavior*, *40*(4),

887–891. https://doi.org/10.1016/0091-3057(91)90102-8

Zacarias, R., Namiki, S., Card, G. M., Vasconcelos, M. L., & Moita, M. A. (2018). Speed dependent

descending control of freezing behavior in Drosophila melanogaster. *Nature Communications*,

9(1), 3697. https://doi.org/10.1038/s41467-018-05875-1

Figures and Tables

Figure 1: Experimental setup for treatment application and environmental conditions. (a) Noxious treatment: contact of a heated soldering iron tip (65 °C) with a randomly selected antenna; innocuous treatment: contact with an unheated tip; control: no contact applied. (b) 'Higher-stress' condition: open-top plastic container (30 × 22 × 10.5 cm) wrapped with red cellophane on the base and sides. (c) 'Lower-stress' condition: closed-top, cellophane-wrapped container with a sand layer on the arena floor, placed within a sheltered brown paper structure.

Figure 2: Directed grooming responses following antennal stimulation in crickets. (a) Proportion of grooming time allocated to the stimulated antenna, shown as estimated marginal means (± 95% CI) from a beta regression. (b) Probability that the stimulated antenna was groomed at all, with grey points indicating individual responses (1 = groomed, 0 = not groomed) and black points ± 95% CI from a binomial GLMM. (c) Duration of grooming directed to the stimulated antenna, with dots representing individuals and boxplots showing distributions by treatment. (d) Temporal dynamics of grooming over the 12-min trial, with lines depicting treatment-specific smooths from a GAM and shaded ribbons showing standard errors. Letters above plots (a) through (c) denote grouping by statistical significance of post-hoc tests.

Table 1: Summary of model results for directed grooming responses following antennal stimulation.
Results are presented for three models: (i) the proportion of grooming directed to the stimulated antenna
(beta regression), (ii) the probability that any grooming was directed toward the stimulated antenna
(binomial GLMM), and (iii) the duration of directed grooming bouts (Tweedie GLMM). Fixed effects
include treatment (control, innocuous contact, noxious heat) and environmental condition (higher- vs.
lower-stress); all models included individual ID as a random effect. Estimates (est.) are shown with
standard errors (s.e.), test statistics (z), and *p*-values.

Effect	Est.	SE	z	p
Specificity of grooming				
intercept	-0.24	0.26	-0.94	0.35
treatment (innocuous)	0.33	0.30	1.10	0.27
treatment (noxious)	0.68	0.28	2.40	**0.016**
environment (higher-stress)	-0.12	0.23	-0.54	0.59
sex (male)	0.01	0.23	0.04	0.97
Grooming probability				
intercept	-1.53	0.40	-3.8	**< 0.01**
treatment (innocuous)	0.69	0.36	1.91	0.06
treatment (noxious)	1.52	0.38	3.96	**< 0.01**
environment (higher-stress)	0.66	0.34	1.95	0.051
sex (male)	0.42	0.34	1.25	0.21
Grooming duration				
intercept	1.10	0.31	3.52	**< 0.01**
treatment (innocuous)	0.60	0.30	2.03	**0.04**
treatment (noxious)	1.39	0.27	5.14	**< 0.01**
environment (higher-stress)	0.10	0.24	0.43	0.67
sex (male)	0.11	0.24	0.46	0.64

518
519