PlasmoFP: leveraging deep learning to predict protein function of uncharacterized proteins across the malaria parasite genus

Harsh R. Srivastava[1,2], Daniel Berenberg[3], Omar Qassab[1], Ziyi Wang[2,4], Richard Bonneau[5*], and Jane M. Carlton[1,2,4*]

[1] Center for Genomics and Systems Biology, Department of Biology, New York University, New York, NY, USA

[2] Johns Hopkins Malaria Research Institute, Department of Molecular Microbiology and Immunology, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD, USA

[3] Courant Institute, Department of Computer Science, New York University, New York, NY, USA

[4] Department of Biomedical Engineering, Johns Hopkins University, Baltimore, MD 21218 USA

[5] Prescient Design, gRED Computational Sciences, Genentech, New York, NY 10010, USA

* Correspondence should be addressed to JaneCarlton@jhu.edu and bonneau.richard@gene.com

Keywords: Protein function prediction, deep learning, protein language models, uncertainty, *Plasmodium*, malaria

Abstract

Malaria, caused by parasites in the genus *Plasmodium*, remains a global health burden, with thousands of deaths and millions of infections each year. Sequencing of the *Plasmodium falciparum* genome in 2002 jump-started functional studies, but a large fraction of all predicted proteins remain poorly characterized. Here, we introduce Plasmodium Function Predictor (PlasmoFP), deep learning models designed specifically for the genus *Plasmodium*. PlasmoFP models are trained on structure-function relationships of proteins of the phylogenetically relevant SAR (Stramenopiles, Alveolate, and Rhizarians) supergroup to predict Gene Ontology terms for partially annotated and 'unknown function' proteins of 19 *Plasmodium* species. Our structure-focused approach addresses long-standing challenges to annotating *Plasmodium* proteins due to their low sequence similarity to other well-

characterized proteins. PlasmoFP models estimate epistemic uncertainty, control false discovery rates in model predictions, and outperform existing methods. By integrating PlasmoFP false discovery rate-controlled predictions with existing annotations for *Plasmodium* proteins, we reduced the proportion of unannotated proteins from 15-59% across species to 3-28% and improved the proportion of fully annotated proteins from 7-42% to 36-68%, improving proteome-wide annotation completeness across the genus. Combined, PlasmoFP predictions help advance *Plasmodium* basic research which can aid progress towards global malaria elimination.

Introduction

Malaria remains a significant global health burden, with an estimated 263 million cases reported in 2023 across 83 endemic countries[1]. Infections are caused by protozoan parasites of genus *Plasmodium*, a member of phylum Apicomplexa within clade Alveolata, which together with Stramenopiles and Rhizaria forms the Stramenopiles–Alveolates-Rhizaria (SAR) eukaryotic supergroup[2]. Five major species are known to cause most human malaria: *P. falciparum*, *P. vivax*, *P. malariae, P. ovale,* and *P. knowlesi*[1]. Other *Plasmodium* species that are used as human disease models infect mammals include African tree-dwelling rodents (e.g., *P. berghei*) and non-human primates (e.g., *P. coatneyi*), while more evolutionarily distant species infect birds and lizards (e.g., *P. gallinaceum*)[3]. Since the publication of the first *Plasmodium* genomes in 2002 (*P. falciparum*[4] and *P. yoelii*[5]), successive efforts have generated assemblies of species from virtually every major clade within the genus. This information is consolidated in PlasmoDB, a resource that provides standardized genomic, transcriptomic, and functional annotations across *Plasmodium* species[6].

Plasmodium genome sequencing has outpaced gene functional annotation, with many predicted gene products labeled simply as "proteins of unknown function" (PUFs). Even in the extensively studied *P. falciparum* proteome, 16% of protein coding genes lack any annotation from the tripartite Gene Ontology (GO) of molecular function (MF), biological process (BP), and cellular component (CC), while 42% are partially GO annotated, and 42% have complete GO annotations. In the less well characterized *P. ovale curtisi*, ~50% of proteins are unannotated, 38% are partially

annotated, and just 12% possess full GO annotations[7]. Inadequately characterized *Plasmodium*

proteins constitute a major bottleneck to malaria parasite basic research, impeding efforts to elucidate

parasite survival strategies and host–parasite interactions.

    The functions of 'unknown' proteins have traditionally been elucidated through resource-

intensive wet-lab experiments[8], which are an impractical method for high-throughput proteome-wide

studies in challenging non-model organisms like *Plasmodium*[9]. Automated function prediction (AFP)

techniques offer an alternative by leveraging the relationship between protein sequence, structure,

and function to infer annotations. AFP annotations often encompass family- or ontology-based

classifications such as Pfam domains, InterPro signatures, and GO terms[10, 11, 12]; the latter are widely

used in multi-omics analyses to uncover mechanistic processes underlying experimental results[11, 13].

    Many AFP methods use sequence similarity approaches such as BLAST, PSI-BLAST, and

HMMER, which infer protein function by comparing query sequences to protein family models

associated with characterized functions, or well-curated proteins in databases like SwissProt[10, 14, 15, 16,

17]. In parallel, early AFP approaches employed genetic interaction networks derived from large scale

genomic interrogation of model species and well-studied organisms chosen for their low cost and

tractable genetics [18, 19]. Not surprisingly, these methods are relatively ineffective for proteins with low

sequence similarity to known proteins[20, 21, 22]. A significant proportion of *Plasmodium* proteomes is

either unique to the genus, or is shared only with other poorly-annotated apicomplexan parasites, or is

only remotely homologous to well-characterized proteins[4]. Additionally, *Plasmodium* proteins

frequently contain long intrinsically disordered regions which do not align well, further reducing the

performance of sequence-based approaches[23]. Such limitations explain why many *Plasmodium*

proteins remain functionally uncharacterized and underscore the need for alternative strategies.

    Recent advances in deep learning have transformed AFP by enabling models to extract signal

directly from protein sequences and capture heuristics correlated with protein sequence, structure,

and function[17, 24, 25, 26, 27, 28]. Foundational protein language models, such as ProtT5 and ESM, have

shown great promise in encoding sequence- and structure-based features into information-rich,

relatively low dimensional vector embeddings[29, 30]. These advances provide groundwork for exploring

alternative strategies beyond traditional sequence similarity approaches. We hypothesize that leveraging the direct relationship between protein structure and function while jointly conditioning learning on phylogenetically relevant relationships offers a more effective strategy for predicting the functions of highly divergent proteins like those found in *Plasmodium*.

Here, we introduce *Plasmodium* Function Predictor (PlasmoFP), a family of deep learning-based AFP models trained separately on each of the three GO subontologies to predict GO terms for *Plasmodium* PUFs and partially annotated proteins. PlasmoFP explicitly models phylogenetically relevant relationships between structure and function within the SAR taxonomic supergroup. Through comprehensive evaluation of model performance and ablations of model architectures, we demonstrate that PlasmoFP outperforms both baseline AFP approaches and existing AFP methods. In turn, PlasmoFP represents the most effective function prediction model developed specifically for the genus *Plasmodium*. We show that integrating PlasmoFP predictions with existing annotations across the MF, BP, and CC GO subontologies significantly reduces the proportion of *Plasmodium* PUFs and increases the proportion of proteins annotated with all three GO subontologies. Collectively, PlasmoFP predictions offer a critical resource for advancing our understanding of *Plasmodium* biology and can serve as a foundation for developing novel therapeutic and diagnostic strategies for malaria.

Results

PlasmoFP models are trained to predict GO terms for each subontology

We developed PlasmoFP, three independently trained deep-learning models on protein structure–function relationships, one for each GO subontology (MF, BP, and CC). Our PlasmoFP models use TM-Vec embeddings[31], structure-imbued vectors of fixed dimensionality, to predict protein function. TM-Vec fine-tunes an underlying sequence encoder (e.g., ProtT5) with a structural similarity learning objective and has been shown to perform well in several tasks related to AFP[29, 31]. Motivated by the evident expressive power and versatility of TM-Vec embeddings, we followed previous work in AFP by choosing to model protein function by $P_\theta(y \mid x_s)$, where θ is the set of learnable parameters of a multi-

layer perceptron (MLP), $y \in \{0, 1\}^c$ is a vector of GO term annotations, and $x_s \in R^d$ is the TM-Vec

embedding of a protein sequence s **(Fig 1)**. PlasmoFP models are trained to optimize the binary

cross-entropy objective, with no gradient updates applied to the TM-Vec encoder. Our choice to keep

the TM-Vec model frozen during training forces PlasmoFP to learn structure-function relationships

from a fixed-sequence structure prior with a relatively small number of parameters rather than

requiring large-scale updates to the underlying protein language model. PlasmoFP models are trained

on automatic and manual assertion GO terms (see Methods), with sequences partitioned by similarity

within the SAR taxonomic supergroup to prevent information leakage. We evaluated models on a

SAR test set and a curated *Plasmodium* holdout set consisting of manually curated GO annotations

held out entirely from model training **(Fig S1)**.

PlasmoFP models quantify uncertainty with deep ensembles

For each of the three GO subontologies (MF, BP, and CC), we used four PlasmoPF models to

capture epistemic uncertainty, the reducible component of error due to limited model knowledge[32]. We

evaluated four approximations to the posterior predictive, *i.e.*, the distribution of outcomes after

averaging over many plausible models learned from the data: (1) Monte Carlo Dropout (MCD)[33]; (2) a

k-member temporal ensemble (predictions from the last k checkpoints); (3) a k-fold ensemble (k

independently initialized models); and (4) a deterministic baseline with no uncertainty **(Fig S3, 1C)**.

For each method, we then calculated uncertainty-adjusted model probabilities, *i.e.*, central estimate

minus dispersion, to curb overconfident errors **(Fig. S4)**.

Across all subontologies, the performance of the temporal ensemble behaved nearly

identically to the deterministic baseline **(Fig 2A)**. This indicates that successive checkpoints are highly

correlated and therefore fail to provide meaningful epistemic uncertainty across k checkpoints. In

contrast, increasing the number of MCD passes led to a gradual drop in (false discovery rate) FDR_{micro}

and sharpness, and an increase in $Precision_{Macro}$ **(Fig 2A)**. This suggests additional stochastic

samples help the model approximate the posterior predictive distribution more faithfully while

producing less overconfident predictions. Deep ensembles exhibited a similar trend; as k increased,

FDR$_{micro}$ decreased and Precision$_{Macro}$ increased **(Fig 2A)**. Notably, at k = 20, we observed the largest

gain in Precision$_{Macro}$ without substantial decrease in Recall$_{Macro}$. This implies that independent weight

initializations explore diverse local optima, providing a richer posterior approximation that yields fewer

false positives. Given the results of this analysis, we selected k = 20 deep ensembles for all

downstream inference.

We next examined the correlation between uncertainty and each PlasmoFP model's softmax

confidence and found that predictions with softmax probability around 0.5 exhibited the highest

uncertainty **(Fig 2B)**. We also observed an inverse correlation between the log count of true positives

per GO term in the test set and epistemic uncertainty, not only suggesting that rarer terms with fewer

positive examples are ascribed higher uncertainty, but also indicating that our estimates capture data

scarcity and underrepresentation during model training **(Fig S5)**. For PlasmoFP GO MF predictions,

the highest uncertainty was observed for enzyme-related functions, including lyases, ligases, and

transferases **(Fig S6)**. These functions also tended to have low to medium information accretion

values, suggesting that the model struggles most with functions requiring fine-grained, residue-level

features that are not fully captured by per-protein embeddings. To further validate our uncertainty

estimates, we perturbed TM-Vec embeddings with Gaussian noise and found that uncertainty initially

rose as inputs drifted from the training manifold, before declining at higher noise levels when

ensemble members converged on uniformly low-confidence predictions **(Fig 2C)**.

Most existing function-prediction methods select a single decision threshold to maximize F$_{max}$

on a given dataset. We introduce an alternative approach that uses empirically observed per-term

FDR on the SAR test set to assemble predicted GO term sets at specified FDR thresholds for each

PlasmoFP model. We adopt the conservative term "expected FDR" (eFDR) at inference time.

Plasmodium PUFs are included in the SAR supergroup, so these empirically-derived eFDR thresholds

should generalize to *Plasmodium* proteins. This method directly controls the expected FDR of

prediction sets by tailoring uncertainty-adjusted threshold cutoffs to each term rather than balancing

overall recall and precision with a one-size-fits-all threshold. We find that the per-term variability in

uncertainty-adjusted probabilities required to meet an FDR cutoff is substantial **(Fig S7)**. At stringent

FDR values (e.g., 1–5% eFDR), some GO terms require uncertainty-adjusted probability cutoffs above 0.9, while others can be called at around 0.5. Notably, CC terms are the most conservative, demanding higher uncertainty-adjusted probabilities, whereas MF terms can be predicted at much lower uncertainty-adjusted probabilities for the same error rate **(Fig S7).**

By performing a grid search over eFDR levels **(Fig. S8)**, we identified an optimal range of 10-20% eFDR within which $Recall_{micro}$ remains high without large decreases in $Precision_{micro}$. Beyond 30% eFDR, recall gains plateau at the cost of precision. $S_{min,}$ an information-content weighted metric combining *remaining uncertainty* (ground truth terms not predicted) *(ru)* and *misinformation* (false positive predictions) *(mi),* reveals a convex trade-off as *ru* falls rapidly as cutoffs relax, while *mi* climbs linearly. This yields an S_{min} minimum also at 10–20% eFDR across all subontologies.

Phylogenetic relevance and TM-Vec embeddings improve function prediction in SAR

We evaluated the impact of conditioning model training on phylogenetically related sequences and incorporating structural information, on function-prediction performance within the SAR supergroup. For each GO subontology, we compared baseline methods along two orthogonal axes. First, to isolate the data effect, we held the algorithm constant while varying the training set, comparing (1) CAFA-CNN, trained on CAFA-5 SwissProt sequences, versus SAR-CNN, trained on SAR-specific sequences; (2) CAFA-TM-Vec trained on TM-Vec embeddings for CAFA-5 SwissProt sequences set versus PlasmoFP, trained on TM-Vec embeddings for SAR-specific sequences; (3) BLAST against SwissProt versus BLAST against SAR-specific sequences; (4) Foldseek against SwissProt versus Foldseek against SAR-specific sequences[14, 34] (**Fig S9)**. In all comparisons, identical train and test splits were used to ensure that observed differences in F_{max} and S_{min} reflect purely the impact of phylogenetic conditioning during training or the use of structure-imbued embeddings.

The inclusion of phylogenetically relevant sequences during training improved performance across architectures: SAR-CNN achieved higher F_{max} and lower S_{min} than CAFA-CNN; PlasmoFP models outperformed CAFA-TM-Vec models; and $BLAST_{Train}$ outperformed $BLAST_{SwissProt}$ in all subontologies **(Table 1)**. We observed no significant hits when querying SAR test sets with

Foldseek$_{SwissProt}$, underscoring the scarcity of detectable structure-based homologs to the broader SwissProt set. Taken together, these results indicate that restricting training or search data to phylogenetically relevant sequences improves model accuracy.

Using a fixed training set to isolate the model effect, we varied the method comparing (1) CAFA-CNN versus CAFA-TM-Vec; (2) SAR-CNN versus PlasmoFP; and (3) BLAST versus Foldseek on SAR databases. We found that training or searching using structure-imbued representations enhanced performance, as reflected by higher F_{max} and lower S_{min} when comparing CAFA-TM-Vec to CAFA-CNN, and PlasmoFP to SAR-CNN across all subontologies **(Table 1)**. When comparing Foldseek$_{Train}$ to BLAST$_{Train}$ we observed higher F-score$_{Macro}$ results for the MF GO ontology and comparable performance for the BP and CC ontologies, consistent with MF annotations being more tightly linked to local structure-based motifs than the broader, protein-level biology captured by BP and CC. Taken together, these results indicate that explicitly leveraging structure–function relationships helps address remote homology in sequence space, consistent with prior work showing structure is highly informative for functional characterization in *P. falciparum*[35]. Additionally, the larger performance gain of PlasmoFP models over Foldseek$_{Train}$ implies that structure-imbued protein embeddings can offset the impact of lower-quality predicted structures during structure–structure searches for function prediction.

To assess whether PlasmoFP models learned SAR-specific structure–function relationships, we evaluated performance on CAFA-5 proteins filtered for terms present in the PlasmoFP training data. We hypothesized that, if PlasmoFP models had specialized on SAR features, performance would decline on the broader CAFA-5 set. Consistent with this, PlasmoFP MF showed a decrease in F_{micro} from 0.908 to 0.485 at 10% eFDR and an increase in S_{min} by ~7 points on the CAFA-5 MF subset compared to the SAR test set **(Table 2)**. This drop was even more pronounced for BP, where F_{micro} fell from 0.831 to 0.332 and S_{min} rose by ~20 points at 20% eFDR, indicating that BP annotations are the most SAR-restricted. By contrast, CC was the most transferable: F_{micro} declined from 0.852 to 0.537 and S_{min} by ~5 points on CAFA-5. Together, these patterns suggest that PlasmoFP captures SAR-specific structure–function signals, most tightly for BP, moderately for MF, and least for CC,

reflecting the deeper conservation of enzymes and core metabolism functions in the MF subontology and cellular component annotations across phylogenetic groups.

PlasmoFP models outperform existing protein function prediction models for *Plasmodium*, and sustain performance with intrinsically disordered proteins

On the *Plasmodium* holdout set, PlasmoFP exhibits a consistent increase in F_{micro} and a decrease in S_{min} as the eFDR threshold is relaxed from 5% to 30% **(Table 3)**. The magnitude of S_{min} reduction is greater than the relative gain in F_{micro}, indicating that improved coverage is accompanied by a disproportionate drop in semantic distance to the true term. This reduction is largely driven by decreasing *ru* rather than *mi*, reflecting the intended design of PlasmoFP to recover additional true annotations while minimizing false positives at higher eFDR thresholds. When compared against DeepGOPlus and ProteInfer[36, 37], at the F_{max} threshold, PlasmoFP outperforms both baselines on F_{micro} and S_{min} across all subontologies, including under the strictest 5% eFDR **(Table 3)**.

Since intrinsic disorder is widespread in *Plasmodium* proteins, we stratified SAR test sets into low (0.0–0.3), medium (0.3–0.6) and high (≥0.6) predicted disorder[38] and evaluated PlasmoFP at 10% eFDR **(Table 4)**. We found that overall, performance is stable: F_{Macro} declines by 0.04 in MF and 0.05 in BP, while CC decreases by 0.03 at high disorder. S_{min} rises modestly by 0.37 in MF and 0.59 in CC but shows a larger increase of ~1.55 in BP from low to high disorder. Precision is largely maintained across bins, decreasing by at most 0.044 in BP and 0.043 in CC, and remaining within 0.045 of the low-disorder baseline in MF while recall decreases slightly with disorder by 0.036 in MF, 0.063 in BP, and 0.038 in CC indicating only mild sensitivity to highly disordered proteins. Together, these results suggest that PlasmoFP models learn relevant structure–function features from proteins with regions of intrinsic disorder, enabling function prediction even in scenarios where traditional sequence-based methods often fail.

PlasmoFP models predict GO terms for partially annotated proteins

Given that PlasmoFP models were trained independently on each ontology, we evaluated their ability to predict meaningful GO terms for partially annotated SAR proteins using a cross-ontology imputation analysis (see Methods). This analysis specifically tests the model's ability to complete missing annotations in one ontology using existing annotations from other subontologies across all SAR test sets as the background. At 20% eFDR, PlasmoFP achieves a mean imputation efficiency of 82.3% across 10,731 proteins, with ontology-specific efficiencies of 85% (95% CI 84–86) for MF, 83% (81.7–84.14) for BP, and 80.4% (79–81.7) for CC. Corresponding pairwise quality scores are 82.3% (95% CI 81.5–83.2) for MF, 80.2% (79.2–81.2) for BP, and 79.8% (78.6–81.0) for CC **(Fig S10)**. Relative to a cross-protein shuffle control, pairwise quality improves by +7.99 (MF, paired-permutation p ≤ 0.001), +6.97 (BP, p ≤ 0.001), and +1.3 (CC, p=0.1512) percentage points (pp); versus a vocabulary-matched random baseline, gains are +7.91, +9.07, and +9.21 pp (p ≤ 0.001).

Results at 5% and 10% eFDR closely mirrored the 20% findings for MF and BP, while CC remained statistically indistinguishable from shuffle at all thresholds (20%: Δ=1.3 pp, p=0.15; 10%: Δ=0.67 pp, p=0.57; 5%: Δ=−2.44 pp, p=0.08) **(Fig S10)**. The lack of separation from the shuffle control across all eFDRs suggests that PlasmoFP CC's gap to the control is minimal, likely because lowering the eFDR reduces the CC prediction vocabulary and per-protein outputs (e.g., 114 terms and 1.6 predictions/protein at 20%; 98 terms and 1.4 at 5%). Therefore, this concentrates predictions on high-frequency, generic localization terms that co-annotate broadly with many MF/BP contexts. Shuffling generic terms across proteins produces similar cross-ontology co-occurrence patterns, compressing the difference to the model. In addition, CC–BP/MF coupling is intrinsically weaker in this evaluation, hence protein-level improvements over shuffle are muted even though performance remains above random. Together, these results demonstrate that PlasmoFP models can effectively predict meaningful, contextual GO terms for partially annotated proteins, extending their benefits from *Plasmodium* PUFs (no GO annotation) to proteins lacking annotation in one or two GO subontologies.

New functional annotations for partially annotated and proteins of unknown function in
Plasmodium

We utilized PlasmoFP models to predict GO terms for PUFs and partially annotated proteins (excluding multigene families) across 19 *Plasmodium* species, over a range of eFDR thresholds. By integrating existing PlasmoDB proteome annotations with PlasmoFP predictions at ≤30% eFDR, we increased the fraction of proteins annotated with all three GO subontologies by an average of 35.3% and reduced the unannotated fraction by 25.3% across all species **(Fig 3)**. Relaxing the eFDR threshold drove a steady increase in full annotation. By 10% eFDR, the unannotated fraction had fallen by a 73.2% average relative to baseline. Beyond 20% eFDR, the unannotated fraction plateaued. Meanwhile, the fully annotated set climbed by an additional 11.1% between 20% and 30% eFDR. The partially annotated category spans ~32.9–57.8% at the existing baseline, peaks at 67.8% around 10% eFDR, and then declines as those proteins transition into the fully annotated pool at higher eFDR thresholds. Notably, PlasmoFP predictions elevated particularly poorly annotated species such as *P. cynomolgi*, *P. yoelii*, and *P. ovale wallikeri* to levels approaching the more richly annotated *P. falciparum* proteome. Even at a stringent 5% eFDR, these species achieve fully annotated fractions of 16.2–19.2%, a 90.5% average relative increase over the existing baseline. Conversely, *P. falciparum*, already at 42.3% fully annotated coverage, showed the smallest absolute gain, reflecting the comprehensive annotation coverage already provided by community efforts[39].

To test whether PlasmoFP's predictions are biologically plausible, we evaluated how novel GO terms relate to existing annotations and functional groupings. We first computed semantic-distance clusters using Resnik's measure and then applied UMAP for dimensionality reduction[40]. We found that newly predicted terms consistently clustered alongside existing annotations across all species and subontologies, indicating strong biological coherence when extant sequences are encoded into the model's latent space, suggesting this featurization is useful and relevant in an AFP setting **(Fig 4)**. Next, we grouped GO terms into functional clusters **(Fig S11)**. In the BP subontology, the largest proportions of PlasmoFP BP predictions across partially annotated and PUF proteins fell into three clusters: (1) a "RNA Processing & Modification" cluster, (2) an "Amino Acid & Fatty Acid Metabolism" cluster, and (3) a "Regulation of Biological Processes" cluster **(Fig S12)**. This suggests that PlasmoFP fills gaps in pathways that are potentially critical for parasite development. In the MF subontology,

most predictions mapped to the "DNA/RNA Binding & Polymerase activity" cluster, followed by "Redox Activity" and "Hydrolases & Phosphatases" clusters **(Fig S12)**. An observed enrichment of transporter-related predictions in both BP and MF subontologies is noteworthy, given recent reports on the challenges of characterizing membrane proteins with standard proteomic methods[41].

When restricting our analysis to PUFs alone, the clustering distribution across subontologies closely mirrored that of the combined set of PUFs and partially annotated proteins, indicating that PUFs do not preferentially associate with any single functional cluster but rather distribute broadly across biological roles **(Fig S12)**. This also implies that, since PUFs mirror the overall distribution, adding PlasmoFP predictions with existing annotations will not bias enrichment analyses or network reconstructions toward any particular functional category. Comparing the integrated dataset of PlasmoDB-existing BP annotations and PlasmoFP BP predictions against existing PlasmoDB BP annotations alone, we observed a 282.66% increase in proteins annotated within the "Regulation of Biological Processes" cluster **(Fig 4)**. In the MF subontology, "Redox Activity" annotations grew by 400.6%, and in the CC subontology, the "Gene-Expression Complexes" cluster saw a 434.1% increase **(Fig 4)**.

Experimental validation of expanded PlasmoFP RNA-associated protein repertoire across the genus *Plasmodium*

Predicting RNA-associated protein functions and RNA-binding remains a major challenge for AFP across diverse contexts[42]. To demonstrate the utility of PlasmoFP predictions, we explored whether they could recapitulate and extend experimentally defined RNA-associated protein repertoires in *P. falciparum* across the genus *Plasmodium*. *P. falciparum* has a limited transcription factor repertoire, suggesting a greater reliance on epigenetic and translation-level regulation of gene expression [43, 44, 45]. Previous bioinformatic surveys catalogued 189 putative RNA-binding proteins spanning 13 families, including RRM (RNA recognition motif), KH (K homology), and zinc-finger domains,[46] and an HMM-based screen of 793 RNA-related domains annotated 18.1% of the proteome as RNA-associated[45]. More recently, proteomic profiling identified 898 RNA-dependent proteins, 545 of which had not been

previously linked to RNA[47]. However, these studies were limited to *P. falciparum*, leaving the RNA-associated landscape of other *Plasmodium* species unexplored.

In PlasmoDB, *P. falciparum* annotations catalogue 920 RNA-associated proteins, comprising approximately 20.03% of the annotated proteome, while other species show substantially lower raw counts. PlasmoFP predictions in all functional categories added 121 new RNA-associated candidates in *P. falciparum*, refining the RNA-associated fraction of the annotated proteome to 19.90% from 20.03%. This slight decrease indicates that, even as PlasmoFP predictions expand overall annotation coverage, the relative proportion of RNA-associated proteins to other functional groups of proteins remains unchanged. This stability also supports findings from previous experimental and computational surveys in *P. falciparum* [45, 46, 47].

Across the genus, PlasmoFP predictions uncovered considerable annotation gains, with the largest increases observed in the non-human primate species *P. cynomolgi* (+330 proteins), *P. inui* (+322 proteins), and *P. fragile* (+284 proteins) **(Fig S13)**. Consistent with *P. falciparum*, despite these large additions, the normalized fraction of RNA-associated proteins in the annotated proteome remains stable at ~15-20%, across all 19 species, changing at most by 1.65% in *P. yoelli* **(Fig 5A)**. Additionally, we find the RNA-associated fraction of the PlasmoFP integrated proteome to be two-to three-fold higher than the DNA-associated fraction, demonstrating that PlasmoFP increases coverage without distorting functional balance across species. Collectively, these results suggest that a large RNA-associated proteome is a conserved, genus-wide feature, underscoring the importance of post-transcriptional control mechanisms across *Plasmodium*[48].

PlasmoFP predictions expand existing functional classes in human-infecting malaria parasites
Proteases in *P. falciparum* are essential for parasite survival, mediating hemoglobin digestion (via aspartic, cysteine, and metalloproteases) and merozoite invasion/egress (via serine and metalloproteases)[49]. However, ~85% of *P. falciparum* proteases remain uncharacterized, and many protease-like proteins in other *Plasmodium* species lack any annotation[49]. Protein transport and localization (protein sorting) constitutes another important aspect of malaria parasite biology,

facilitating processes ranging from organelle targeting to nutrient uptake, but transport-related

functions are under-annotated outside of *P. falciparum* and *P. vivax* [41, 50, 51]. To assess how PlasmoFP

fills annotation gaps in these two critical functional categories (protease-like enzymes and protein

sorting), we investigated PlasmoFP 20% eFDR MF and 30% eFDR BP predictions across all human-

infecting *Plasmodium* species.

      For proteases, we filtered PlasmoFP MF-predicted terms for aspartic, cysteine, serine and

metallo- protease activities. PlasmoFP predictions added seven metalloproteases, five serine

proteases, and two previously unannotated aspartic proteases in *P. falciparum*, boosting the total

count of protease-like functions by 28% **(Fig 5C)**. (Because our MF models could not be trained on

threonine-type proteases due to insufficient training data for this term, that category showed no

change.) When this approach was applied across all human-infecting *Plasmodium* species, the

relative proportions of metalloproteases, serine, cysteine, and aspartic proteases remained similar,

but less-studied species realized larger absolute gains. For example, *P. vivax* and *P. ovale curtisi*

gained 5 and 10 additional serine proteases, an 83% and 100% increase, respectively. Overall,

predicted protease-like proteins increased by 34% on average across all human infecting *Plasmodium*

species.

      For transport/localization proteins, we took the subset of proteins found in the "Protein

Transport & Localization" cluster, classified GO terms in this cluster into Transporter Classification

Database[52] inspired subclusters and counted the fraction of proteins in each subcluster **(Fig 5B)**. The

largest gain occurred in the "localization/targeting without direct transmembrane step" cluster, which

rose from 1–5 proteins to 148–231 proteins across species. GO terms found in this cluster such as

"establishment of protein localization to organelle," "establishment of protein localization to

membrane," and "organelle localization by membrane tethering" that were previously under-annotated

are now broadly represented in all five species. The "generic/not clearly defined transmembrane

transport" cluster more than doubled in size on average while the "vesicle-mediated transport" cluster

increased by up to 60 proteins in *P. ovale curtisi and P. ovale wallikeri*. This is noteworthy since

vesicle-mediated transport is vital for protein export, nutrient uptake, and organelle maintenance[53].

"Cytoskeletal-based transport" annotations, previously absent in the genus *Plasmodium*, appeared in all species with 2-6 proteins. While the "unclassified" cluster also increased with the PlasmoFP integrated dataset, the total number of unique proteins annotated with terms from the "Protein Transport & Localization" increased between 155-273 proteins across all species. Although *P. falciparum* and *P. vivax* remain highest in total transport annotations, *P. malariae*, *P. ovale curtisi*, and *P. ovale* wallikeri each gained ~twofold more transport-related proteins.

Discussion

Despite considerable progress in deep learning-based protein function prediction frameworks, no end-to-end methodologies exist to annotate uncharacterized proteins that have low sequence homology to well-characterized proteins within a phylogenetically defined subset of species. Existing models, such as DeepGOPlus and ProteInfer, are trained on large, phylogenetically agnostic datasets such as SwissProt, which consist primarily of well-characterized sequences from model organisms[36, 37]. This approach often overlooks the unique characteristics of non-model organisms, where deviations from expected model performance are common due to remote homology instances. To address these challenges, we developed PlasmoFP, protein function prediction models tailored to a subset of non-model eukaryotic species found within the Stramenopiles–Alveolates-Rhizaria (SAR) eukaryotic supergroup. We address challenges posed by both remote homology and non-model organism representation during training by leveraging protein structure representations using a fine-tuned structure-imbued protein language model, TM-Vec[31], and conditioning model learning on the phylogenetically relevant SAR structure-function relationships. By focusing on structure-function relationships, our approach to building a species-specific function prediction model can be generalized to non-model proteomes across the tree of life, provided their lineages have sufficient representation in annotation databases.

Although existing AFP techniques often provide confidence scores, few directly leverage uncertainty quantification to generate calibrated GO term prediction sets with explicit error-rate control. We find that by incorporating uncertainty using deep ensembles, we lower the observed FDR

in prediction sets across all subontologies in comparison to point models. Since PlasmoFP targets a subset of species rather than a broad spectrum, we also developed an FDR threshold-selection method that directly controls the false positive rate in prediction sets. Complementary structure-guided screens in *P. falciparum* using DALI on AlphaFold models have mapped folds to hundreds of PUFs[35]; however, they do not produce calibrated GO term prediction sets or provide explicit error-rate control. PlasmoFP addresses this gap while retaining the benefits of structural signal. To evaluate PlasmoFP predictions against existing *Plasmodium* annotations, we developed a score to quantify biologically plausible GO term pairs across subontologies, and find that PlasmoFP predictions generated annotation sets with co-occurrence patterns aligning closely with SwissProt proteins. PlasmoFP models were also able to maintain high performance on proteins with regions of intrinsic disorder, which are challenging to annotate using sequence-based deep learning techniques. This suggests that the intrinsic disorder content of a protein has little impact on learned structure-function relationships during model training, and highlights the potential for developing a function prediction model for disordered proteins. Collectively, these findings establish PlasmoFP models as an effective protein function predictor for the genus *Plasmodium*.

At the genus level, incorporating PlasmoFP predictions increased annotation completeness across all three GO subontologies and harmonized coverage in less-studied species. Semantic-distance clustering and functional cluster analyses indicate that the newly predicted terms are biologically plausible and fill pathway gaps spanning metabolism, regulation, and membrane transport. Together, these additions broaden and balance the functional landscape across 19 *Plasmodium* species and provide a stronger and richer foundation for downstream multi-omics analyses and hypothesis generation. We show that PlasmoFP model predictions reduced the proportion of unannotated proteins in the genus *Plasmodium* at controlled expected false discovery rates and use PlasmoFP predictions to estimate the fraction of proteins in functional subclasses relevant to ongoing drug-target discovery efforts.

These analyses show PlasmoFP captures both canonical and structurally conserved functional domains that traditional sequence-based methods overlook. Importantly, PlasmoFP predictions for

Plasmodium PUFs show no spurious enrichment in any single GO process, helping to prevent false-positive biases in downstream tasks such as GO term enrichment. The underlying framework used to train PlasmoFP is also compatible with additional information streams, including co-evolutionary features from MSAs[54, 55], per-residue sequence–structure embeddings, phylogenetic weighting, and calibrated uncertainty strategies. This positions PlasmoFP as a scalable, uncertainty-aware foundation for predicting annotation classes beyond GO across the genus *Plasmodium*.

Methods

Datasets

Train, validation, and test set construction: We assembled training, validation, test sets by collecting 907,145 proteins from the Uniprot release (2024_04): '(taxonomy_id:2698737) AND ((existence:3) OR (existence:2) OR (existence:1)) AND (length:[0 TO 1200]) NOT (taxonomy_id:5820) AND (active:true) AND (go:*) + '(taxonomy_id:5820) AND ((existence:3) OR (existence:2) OR (existence:1)) AND (length:[0 TO 1200]) AND (go_automatic:*) AND (active:true)'. Our criteria included proteins within the SAR supergroup, a protein existence score between 1-3, a total protein length ≤ 1200 amino acids, and exclusion of sequences with manually asserted GO terms in the genus *Plasmodium*. The dataset was partitioned by GO subontology into 716,735 MF, 573,293 BP, and 630,944 CC proteins.

   Redundant sequences were eliminated using the MMseqs2 clustering function with parameters `–seq-id=0.9` and `–cov=1`[56]. Cluster representatives with the greatest number of GO terms were retained for maximum informational value. GO terms associated with each ontology were propagated to the respective root node using the 'ancestors' and 'is_a' queries from the QuickGO REST API for each GO term[57]. Information Accretion for was calculated using the method described in and implemented via [https://github.com/claradepaolis/InformationAccretion], with release 2024-09-08 of the go-basic.obo[58].

   Each ontology dataset was processed with MMseqs2 using `–seq-id=0.3` and `–cov=1` to form clusters of sequences sharing 30% similarity. We allocated whole clusters to the train, validation, or

test sets using 80/10/10 partitions to prevent data leakage and removed terms with a frequency ≤ 50 across all sets to avoid training on low-frequency terms. This resulted in 289,930, 247,286, and 227,756 training sequences for the MF, BP, and CC subontologies, respectively; 29,188, 27,602, and 24,531 sequences for validation; and 33,343, 29,679, and 26,746 sequences for testing. The training label spaces comprised 744 MF terms, 1,548 BP terms, and 255 CC terms. For evaluation, GO terms in the validation and test sets were restricted to those present in the corresponding training label spaces.

Plasmodium holdout set composition: We collected 1,527 sequences from Uniprot (2024_04): '(taxonomy_id:5820) AND ((existence:3) OR (existence:2) OR (existence:1)) AND (length:[0 TO 1200]) AND (go_manual:*) AND (active:true)'. Our criteria included proteins in the genus *Plasmodium*, possessing manually asserted terms and a total protein length ≤ 1200 amino acids. This dataset was also divided into three subsets, one for each GO ontology. GO terms for sequences in this set were not filtered to create the most representative version of the current annotation in the genus.

Plasmodium PUFs and partially annotated set: We processed *Plasmodium* genomes from Release 68 of PlasmoDB by parsing Annotated_Proteins.fasta files for each species and applying five sequential filters: (1) gene products longer than 1,200 amino acids were removed due to TM-Vec embedding limitations; (2) entries annotated as pseudogenes or partial genes were excluded based on description fields; (3) only primary-isoform transcripts ending in ".1", "_1" or "_t1" were retained; (4) sequences belonging to large, variable multigene families were excluded by screening for the keywords "emp", "msp", "rifin", "stevor", "vir", "fam-", "pir", "cir", "yir" and "surf"; and (5) any gene annotated as part of the apicoplast genome was removed.

TM-Vec embeddings: We used the TM-Vec model to compute embeddings for sequences in all datasets as input for PlasmoFP and ALL-TM-Vec models[31]. Each sequence was encoded using the 'CATH' model, resulting in a flattened embedding with a dimensionality of (1,512) per protein sequence.

PlasmoFP training and hyperparameter tuning

For each model in each ontology, we conducted a grid search over the following parameters: number of cascading hidden layers ([256], [256,128], [256,128,64], [256,128,64,32]), learning rate (0.01–0.00001), and number of epochs (10, 15, 20, 30, 40), using a standard binary cross-entropy loss function. During this grid search, the dropout parameter was fixed at 0.3 in the last layer before the output and a ReLU activation function was used for each layer. Each trained model was evaluated on the respective ontology validation set using the F_{max} and S_{min} CAFA metrics, with the chosen model exhibiting the lowest scores for both (**Fig S2**)[20].

To address potential class imbalance, we adjusted the binary cross-entropy loss function by weighting terms according to their depth and inverse frequency. Additionally, we explored alternative loss functions, including Focal, Jaccard, Hinge, and Asymmetric loss functions, but found that the unweighted binary cross-entropy consistently produced the lowest F_{max} and S_{min} scores across all PlasmoFP models (**Fig S2**) on the validation set.

Model inference, evaluation, and output

For each GO subontology and protein–term pair, we infer a probability from our k-member deep ensemble by computing the median prediction minus its median absolute deviation (MAD). We then preform a parameter sweep over decision thresholds from 0.01 to 0.99, linearly spaced by 0.01, and for each GO term calculate the empirical FDR on the test set. If a term never appears in validation, we borrow the closest observed ancestor or descendant term (by the ontology graph) to estimate its FDR. This lets us map each binarization threshold to an observed FDR for every GO term. Finally, when we predict annotation sets for partially annotated or unknown proteins, we choose thresholds that guarantee a user-specified expected FDR. If a threshold is unavailable, we use the threshold corresponding to the maximum FDR ≤ target FDR.

We evaluated our models using standard CAFA metrics[20] reporting F_{max}, S_{min}, remaining uncertainty (*ru*), and misinformation (*mi*) in "partial mode" analysis as outlined in[20, 58]. For our intrinsic disorder evaluation, disorder was predicted using AIUPRED [38], with predicted disorder content for each protein calculated as the proportion of residues predicted to be disordered relative to the total

protein length. We measured model performance using F_{max} and S_{min} on bins of SAR test set sequences containing varying levels of predicted intrinsic disorder.

Model uncertainty estimation

We approximate the posterior predictive with an ensemble $p(y \mid x_s, D) \approx \frac{1}{M}\sum_{j=1}^{M} p(y \mid x_s, \theta_j)$.

For each protein–GO pair we obtain M predictive outputs $\left\{\widehat{y^{(j)}}\right\}_{j=1}^{M}$, where j indexes the ensemble member produced by one of four approaches: (1) Deterministic baseline: M=1; (2) Monte Carlo Dropout (MCD): M stochastic forward passes with dropout enabled at inference; (3) Temporal ensemble: predictions from the last k=5 checkpoints from model training (M=k); (4) Deep ensemble: k independently initialized models *(M=k)* on different folds of the training data. We compute an uncertainty-adjusted probability by taking the central estimate (ensemble mean or median of the softmax outputs) minus a dispersion measure (MAD or standard deviation). Models are trained on the full training set except for the deep ensemble method where each k member is trained independently with a distinct initialization.

Training baseline models

BLAST: For each GO sub-ontology, protein sequence databases were built with makeblastdb (BLAST+ v2.15.0) from the SAR train set, SAR test set, and UniProtKB/SwissProt (accessed August 29, 2025). We performed BLASTp searches with an E-value threshold of 1×10^{-5} and returned up to 100 target sequences per query. Searches were run in two phases: first, testing sequences against the Swiss-Prot database, and second, testing sequences against the SAR training sequences. Baseline predictions were produced via annotation transfer from significant BLAST hits, using percent identity rescaled to a 0–1 confidence score. For each query, GO terms were gathered from all significant hits within the corresponding sub-ontology. When multiple hits supported the same GO term, we kept the maximum confidence score.

FoldSeek: Structural databases for the SAR train and test sets were created with FoldSeek's createdb, using ProstT5 model weights to convert amino-acid sequences into 3Di structural alphabet representations. We then ran easy-search with an E-value filter of 1×10^{-5} and returned up to 100 sequences per query, searching both the Swiss-Prot structural database (AlphaFold DB representations) and the SAR training-set structural databases. Confidence scores for GO terms were derived by applying a sigmoid transformation to the bit scores of significant hits, scaled by the global standard deviation of all bit scores (with a minimum scale of 10.0 to avoid overly steep curves). As with BLAST, when multiple hits mapped to the same GO term, we retained the maximum confidence value.

SAR CNN: The SAR CNN models were trained and evaluated using the train and validation sets described earlier, with a one-dimensional convolutional neural network (CNN) architecture. The CNN architecture consists of two one-dimensional convolutional layers with kernel size 8 and padding of 4 to preserve sequence length. The first convolutional layer takes 21 input channels (corresponding to the one-hot encoded amino acids), while the number of output channels for both layers was optimized during hyperparameter tuning. Each convolutional layer is followed by ReLU activation functions and max-pooling operations with kernel size 2, resulting in a 4-fold reduction in sequence length. After the convolutional layers, the output is flattened and passed through two fully connected layers with dropout regularization (dropout rate optimized between 0.1-0.5). The architecture and training hyperparameters were optimized using Optuna with 20 trials per ontology. The optimization search space included: first convolutional layer output channels (32-512), second convolutional layer output channels (32-512), fully connected layer size (128-1,024), dropout rate (0.1-0.5), and learning rate (1×10^{-5} to 1×10^{-2} on log scale). Each trial was trained for 10 epochs, and the configuration maximizing F_{max} on the validation set was selected for final model training. Final models were trained for 20 epochs using the Adam optimizer with binary cross-entropy. Training was performed with batch size 32, and sequences were processed in batches of 1,000.

CAFA-CNN based on DeepGOPlus: The CAFA-CNN models employed identical architecture and training procedures as the SAR CNN models but utilized the unified CAFA5 protein dataset from

the CAFA5 competition [https://www.kaggle.com/competitions/cafa-5-protein-function-prediction]. The dataset was filtered in two stages: (1) retain only proteins with target ontology annotations (2) include GO terms present in the corresponding SAR training set vocabulary. Optuna hyperparameter optimization was performed with 20 trials on a 5% validation split randomly selected from the training data. The same optimization search space was used (convolutional channels 32-512, fully connected size 128-1,024, dropout 0.1-0.5, learning rate 1×10^{-5} to 1×10^{-2}), with each trial trained for 10 epochs to maximize F_{max}.

Final models were trained on the complete dataset using optimal hyperparameters for 20 epochs, maintaining the same technical specifications as SAR CNN (Adam optimizer, binary cross-entropy loss, batch size 32).

CAFA-TM-Vec: The CAFA-TM-Vec models utilized the same CAFA5 dataset as CAFA-CNN but employed the PlasmoFP neural network architecture instead of convolutional layers. Input consisted of 512-dimensional TM-Vec protein embeddings rather than one-hot encoded sequences. Hyperparameter optimization used Optuna with 50 trials on a 90/10 train-validation split, optimizing four predefined architectures ([256], [256,128], [256,128,64], [256,128,64,32]), five learning rates (0.01 to 0.00001), five epoch counts (10-40), and dropout rates (0.1-0.5). Models were trained with batch size 256 using Adam optimizer and binary cross-entropy loss, with final models trained on the complete dataset using optimal hyperparameters.

Pairwise score

We assessed PlasmoFP's ability to predict missing GO annotations in one ontology using existing annotations from the other two subontologies via a cross-ontology imputation analysis. For each protein annotated in ≥1 of the three subontologies, we masked its target-ontology annotations, generated PlasmoFP predictions for the target ontology from sequence, and then evaluated predicted versus real (masked) annotations against the protein's existing cross-ontology annotations.

Term coherence was quantified with a pairwise quality score:

$$\text{Pairwise Quality Score} = \frac{|\{(p,e) \mid \text{cooccurrence}(p,e) \geq 3\}|}{|\{(p,e) \mid \text{cooccurrence}(p,e) > 0\}|} \times 100$$

where *(p,e)* pairs each predicted target-ontology term *p* with each existing cross-ontology term *e* for

the same protein. Co-occurrence counts were taken from the SwissProt unpropagated co-occurrence

matrix (28,505 terms; \~6.9M non-zero entries). Pairs with count ≥3 were counted as "good"; pairs

with count >0 were "defined." We applied deepest-term filtering to target-ontology terms (predictions,

and we applied the same filtering to ground truth and baselines) to reduce parent–child artifacts

before scoring.

Imputation efficiency was defined as:

$$\text{Imputation Efficiency} = \frac{\text{Prediction Pairwise Score}}{\text{Ground-Truth Pairwise Score}} \times 100$$

where the ground-truth score is computed identically after substituting the real (masked) target-

ontology annotations for predictions. Pairwise scores (and the efficiencies derived from them) were

computed as ratios of sums across proteins—i.e., we first sum "good" and "defined" pairs over

proteins and then take their ratio. Analyses were run at eFDR thresholds of 20%, 10%, and 5%. To

contextualize performance, we used two baselines evaluated with the same filtering pipeline: (1) a

cross-protein shuffle, a uniform permutation (without replacement) that redistributes each model-

predicted term set across proteins (testing protein-specific assignment while preserving term

frequencies and set sizes); and (2) a pure random control that samples terms uniformly from the

target-ontology prediction vocabulary, matching each protein's predicted set size. Overall efficiency is

reported as the unweighted mean of ontology-specific efficiencies.

For the four scores (prediction, ground-truth, shuffle, random), we computed 95% CIs via a

nonparametric bootstrap over proteins, resampling proteins with replacement and recomputing the

ratio-of-sums score within each replicate; intervals are percentile-based. For efficiencies (ratios of

scores), we used a paired bootstrap over proteins (B=5,000), resampling the same indices for numerator (prediction, shuffle, or random) and denominator (ground truth).

Significance testing: To assess model-versus-control differences in pairwise score, we performed a two-sided paired permutation test across proteins (N=10,000 permutations). In each permutation, we randomly swapped (per protein) the two conditions being compared (e.g., model vs shuffle), recomputed the aggregate difference in pairwise score (ratio-of-summed "good" to "defined" pairs), and formed a null distribution. P-values used add-one smoothing,

$$p = \frac{k + 1}{N + 1}$$

where *k* is the number of permutations whose absolute aggregate difference in pairwise score (computed as the ratio of summed "good" to summed "defined" pairs across proteins) was at least as large as the observed absolute difference, and *N* is the total number of permutations.

Quality control

We performed quality control on the predicted annotations from all PlasmoFP models. Across all eFDR thresholds, we filtered out terms listed in the Gene Ontology Consortium's *gocheck_do_not_annotate.obo* file. These are terms which are considered general and uninformative.

GO term clustering by semantic distance

A GO term similarity matrix using Resnik for existing annotations combined with PlasmoFP predicted GO terms was calculated using biopython[59]. An elbow plot was used to determine the optimal number of clusters for k-means clusters and generate word clouds using term names and manually labeled.

Comparison with other models

To benchmark the performance of the PlasmoFP model, we compared it against two state-of-the-art models: ProteInfer and DeepGOPlus[36, 37]. These models were selected for their demonstrated effectiveness in GO term prediction tasks. Both models were installed and configured locally in accordance with the instructions provided in respective repositories. Predictions were generated for the *Plasmodium* holdout set. For both ProteInfer and DeepGOPlus, we set the prediction threshold to zero, allowing us to retrieve all predicted probabilities associated with each GO term. This ensured that no potential predictions were excluded prior to post-processing. To enable a fair comparison, we applied the same evaluation criteria across all models. Using the *Plasmodium* holdout set predictions, we determined the threshold that optimized the F_{max} metric for each model individually.

Data availability

Training, validation and testing data splits for PlasmoFP can be found at https://doi.org/10.6084/m9.figshare.30100396.v1. PlasmoFP predictions for all 19 *Plasmodium* species can be explored at https://janecarltonlab.org/plasmofp. A Google Colab Jupyter notebook can be accessed at: http://tiny.cc/PlasmoFP

Code availability

The source code for all key findings in this study are available on Github at https://github.com/harshstava/PlasmoFP_public. Raw dictionaries containing model predictions and the source code for the interactive Python webapp can be found at https://github.com/harshstava/PlasmoFP_Explorer.

References

1.   World Health Organization. World Malaria Report 2024. Geneva (2024).

2.   Burki F, Roger AJ, Brown MW, Simpson AGB. The New Tree of Eukaryotes. *Trends Ecol Evol* **35**, 43-55 (2020).

3.   Gozalo AS, Robinson CK, Holdridge J, Mahecha OFL, Elkins WR. Overview of Plasmodium spp. and Animal Models in Malaria Research. *Comp Med* **74**, 205-230 (2024).

4.   Gardner MJ*, et al.* Genome sequence of the human malaria parasite Plasmodium falciparum. *Nature* **419**, 498-511 (2002).

5.   Carlton JM*, et al.* Genome sequence and comparative analysis of the model rodent malaria parasite Plasmodium yoelii yoelii. *Nature* **419**, 512-519 (2002).

6.   Harb OS, McDowell MA, Roos DS. VEuPathDB Resources: A Platform for Free Online Data Exploration, Integration, and Analysis. *Methods Mol Biol* **2802**, 573-586 (2024).

7.   Aurrecoechea C*, et al.* PlasmoDB: a functional genomic database for malaria parasites. *Nucleic Acids Res* **37**, D539-543 (2009).

8.   Schnoes AM, Ream DC, Thorman AW, Babbitt PC, Friedberg I. Biases in the experimental annotations of protein function and their effect on our understanding of protein function space. *PLoS Comput Biol* **9**, e1003063 (2013).

9.   Kimmel J*, et al.* Gene-by-gene screen of the unknown proteins encoded on Plasmodium falciparum chromosome 3. *Cell Syst* **14**, 9-23 e27 (2023).

10. UniProt C. UniProt: the Universal Protein Knowledgebase in 2025. *Nucleic Acids Res* **53**, D609-D617 (2025).

11. Gene Ontology C*, et al.* The Gene Ontology knowledgebase in 2023. *Genetics* **224**, (2023).

12. Blum M*, et al.* InterPro: the protein sequence classification resource in 2025. *Nucleic Acids Res* **53**, D444-D456 (2025).

13. Maleki F, Ovens K, Hogan DJ, Kusalik AJ. Gene Set Analysis: Challenges, Opportunities, and Future Research. *Front Genet* **11**, 654 (2020).

14. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol* **215**, 403-410 (1990).

15. Altschul SF*, et al.* Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* **25**, 3389-3402 (1997).

16. Finn RD, Clements J, Eddy SR. HMMER web server: interactive sequence similarity searching. *Nucleic Acids Res* **39**, W29-37 (2011).

17. Lin B, Luo X, Liu Y, Jin X. A comprehensive review and comparison of existing computational methods for protein function prediction. *Brief Bioinform* **25**, (2024).

18. Costanzo M*, et al.* A global genetic interaction network maps a wiring diagram of cellular function. *Science* **353**, (2016).

19. Li S*, et al.* A map of the interactome network of the metazoan C. elegans. *Science* **303**, 540-543 (2004).

20. Zhou N, *et al.* The CAFA challenge reports improved protein function prediction and new functional annotations for hundreds of genes through experimental screens. *Genome Biol* **20**, 244 (2019).

21. Jiang Y, *et al.* An expanded evaluation of protein function prediction methods shows an improvement in accuracy. *Genome Biol* **17**, 184 (2016).

22. Radivojac P, *et al.* A large-scale evaluation of computational protein function prediction. *Nat Methods* **10**, 221-227 (2013).

23. Guy AJ, *et al.* Insights into the Immunological Properties of Intrinsically Disordered Malaria Proteins Using Proteome Scale Predictions. *PLoS One* **10**, e0141729 (2015).

24. Bepler T, Berger B. Learning the protein language: Evolution, structure, and function. *Cell Syst* **12**, 654-669 e653 (2021).

25. Kryshtafovych A, Schwede T, Topf M, Fidelis K, Moult J. Critical assessment of methods of protein structure prediction (CASP)-Round XIV. *Proteins* **89**, 1607-1617 (2021).

26. Ofer D, Brandes N, Linial M. The language of proteins: NLP, machine learning & protein sequences. *Comput Struct Biotechnol J* **19**, 1750-1758 (2021).

27. Rives A, *et al.* Biological structure and function emerge from scaling unsupervised learning to 250 million protein sequences. *Proc Natl Acad Sci U S A* **118**, (2021).

28. Schmirler R, Heinzinger M, Rost B. Fine-tuning protein language models boosts predictions across diverse tasks. *Nat Commun* **15**, 7407 (2024).

29. Elnaggar A, *et al.* ProtTrans: Toward Understanding the Language of Life Through Self-Supervised Learning. *IEEE Trans Pattern Anal Mach Intell* **44**, 7112-7127 (2022).

30. Lin Z, *et al.* Evolutionary-scale prediction of atomic-level protein structure with a language model. *Science* **379**, 1123-1130 (2023).

31. Hamamsy T, *et al.* Protein remote homology detection and structural alignment using deep learning. *Nat Biotechnol* **42**, 975-985 (2024).

32. Gawlikowski J, *et al.* A survey of uncertainty in deep neural networks. *Artificial Intelligence Review* **56**, 1513-1589 (2023).

33. Gal Y, Ghahramani Z. Dropout as a bayesian approximation: Representing model uncertainty in deep learning. In: *international conference on machine learning*). PMLR (2016).

34. van Kempen M, *et al.* Fast and accurate protein structure search with Foldseek. *Nat Biotechnol* **42**, 243-246 (2024).

35. Behrens HM, Spielmann T. Identification of domains in Plasmodium falciparum proteins of unknown function using DALI search on AlphaFold predictions. *Sci Rep* **14**, 10527 (2024).

36. Sanderson T, Bileschi ML, Belanger D, Colwell LJ. ProteInfer, deep neural networks for protein functional inference. *Elife* **12**, (2023).

37. Kulmanov M, Hoehndorf R. DeepGOPlus: improved protein function prediction from sequence. *Bioinformatics* **36**, 422-429 (2020).

38. Erdos G, Dosztanyi Z. AIUPred: combining energy estimation with deep learning for the enhanced prediction of protein disorder. *Nucleic Acids Res* **52**, W176-W181 (2024).

39. Bohme U, Otto TD, Sanders M, Newbold CI, Berriman M. Progression of the canonical reference malaria parasite genome from 2002-2019. *Wellcome Open Res* **4**, 58 (2019).

40. Supek F, Bosnjak M, Skunca N, Smuc T. REVIGO summarizes and visualizes long lists of gene ontology terms. *PLoS One* **6**, e21800 (2011).

41. Wunderlich J. Updated List of Transport Proteins in Plasmodium falciparum. *Front Cell Infect Microbiol* **12**, 926541 (2022).

42. Li X, Quon G, Lipshitz HD, Morris Q. Predicting in vivo binding sites of RNA-binding proteins using mRNA secondary structure. *RNA* **16**, 1096-1107 (2010).

43. Rios KT*, et al.* Widespread release of translational repression across Plasmodium's host-to-vector transmission event. *PLoS Pathog* **21**, e1012823 (2025).

44. Hollin T, Gupta M, Lenz T, Le Roch KG. Dynamic Chromatin Structure and Epigenetics Control the Fate of Malaria Parasites. *Trends Genet* **37**, 73-85 (2021).

45. Reddy BP*, et al.* A bioinformatic survey of RNA-binding proteins in Plasmodium. *BMC Genomics* **16**, 890 (2015).

46. Bunnik EM, Batugedara G, Saraf A, Prudhomme J, Florens L, Le Roch KG. The mRNA-bound proteome of the human malaria parasite Plasmodium falciparum. *Genome Biol* **17**, 147 (2016).

47. Hollin T*, et al.* Proteome-Wide Identification of RNA-dependent proteins and an emerging role for RNAs in Plasmodium falciparum protein complexes. *Nat Commun* **15**, 1365 (2024).

48. Simantov K, Goyal M. Chapter 17 - Post-transcriptional regulation of gene expression in human malaria parasite Plasmodium falciparum. In: *Post-Transcriptional Gene Regulation in Human Disease* (eds Jain BP, Goswami SK, Sharma T). Academic Press (2022).

49. Mahanta PJ, Lhouvum K. Plasmodium falciparum proteases as new drug targets with special focus on metalloproteases. *Mol Biochem Parasitol* **258**, 111617 (2024).

50. Martin RE, Henry RI, Abbey JL, Clements JD, Kirk K. The 'permeome' of the malaria parasite: an overview of the membrane transport proteins of Plasmodium falciparum. *Genome Biol* **6**, R26 (2005).

51. Taraschi TF, Trelka D, Martinez S, Schneider T, O'Donnell ME. Vesicle-mediated trafficking of parasite proteins to the host cell cytosol and erythrocyte surface membrane in Plasmodium falciparum infected erythrocytes. *Int J Parasitol* **31**, 1381-1391 (2001).

52. Saier MH*, et al.* The Transporter Classification Database (TCDB): 2021 update. *Nucleic Acids Res* **49**, D461-D467 (2021).

53. Abida*, et al.* Extracellular vesicles in malaria pathogenesis and Therapy: Emerging insights and future perspectives. *Microb Pathog* **207**, 107902 (2025).

54.   de Oliveira S, Deane C. Co-evolution techniques are reshaping the way we do structural bioinformatics. *F1000Res* **6**, 1224 (2017).

55.   Hopf TA*, et al.* Sequence co-evolution gives 3D contacts and structures of protein complexes. *Elife* **3**, (2014).

56.   Steinegger M, Soding J. MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. *Nat Biotechnol* **35**, 1026-1028 (2017).

57.   Binns D, Dimmer E, Huntley R, Barrell D, O'Donovan C, Apweiler R. QuickGO: a web-based tool for Gene Ontology searching. *Bioinformatics* **25**, 3045-3046 (2009).

58.   Clark WT, Radivojac P. Information-theoretic evaluation of predicted ontological annotations. *Bioinformatics* **29**, i53-61 (2013).

59.   Cock PJ*, et al.* Biopython: freely available Python tools for computational molecular biology and bioinformatics. *Bioinformatics* **25**, 1422-1423 (2009).

Acknowledgments

We thank Nathan Frey and Francisco Callejas-Hernandez for helpful discussions around project ideation. We thank Steven Sullivan, Mari Shiratori, and Kavya Phadke for editorial assistance. Research reported in this publication was partially supported by the NYU IT High Performance Computing resources, services, and staff expertise.

Author contributions

H.R.S collected data, performed the research, implemented the method, and conducted analysis. H.R.S, D.B. and O.Q. designed experiments. Z.W. assisted in downstream applications for the method and technical advice. J.M.C. and R.B. advised the research. J.M.C provided the funding. All authors contributed to writing the manuscript.

Competing interests

The following author wishes to disclose his industry relations although there are no competing interests to the work published here: R.B. is an employee of Genentech/Roche. The remaining authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at …

Correspondence and requests for materials should be addressed to Jane M. Carlton and Richard

Bonneau.

Table 1: PlasmoFP performance on SAR test set

	Model	F_{Macro}	S_{min}
MF	**PlasmoFP 10% eFDR**	**0.900**	**2.050**
	CAFA-TM-Vec	0.632	7.425
	SAR-CNN	0.721	5.514
	CAFA-CNN	0.456	10.062
	$BLASTp_{SwissProt}$	0.639	24.525
	$BLASTp_{Train}$	0.765	9.328
	$Foldseek_{Train}$	0.844	10.613
BP	**PlasmoFP 10% eFDR**	**0.851**	**3.780**
	CAFA-TM-Vec	0.593	8.826
	SAR-CNN	0.644	7.466
	CAFA-CNN	0.396	12.415
	$BLASTp_{SwissProt}$	0.514	57.571
	$BLASTp_{Train}$	0.782	9.827
	$Foldseek_{Train}$	0.764	12.329
CC	**PlasmoFP 10% eFDR**	**0.875**	**1.235**
	CAFA-TM-Vec	0.698	2.864
	SAR-CNN	0.766	2.175
	CAFA-CNN	0.687	2.918
	$BLASTp_{SwissProt}$	0.501	20.343
	$BLASTp_{Train}$	0.777	3.008
	$Foldseek_{Train}$	0.771	3.900

Bolded values indicate best performance in each subontology; F_{Macro} values are $F_{Macro \mid max}$ for all

models except PlasmoFP 10% eFDR. S_{min} values are reported at the threshold corresponding to

$$F_{Macro \mid max}$$

Table 2: PlasmoFP performance on SAR test set and CAFA test set

	Test set (eFDR)	F_{micro}	S_{min}	Precision$_{micro}$	Recall$_{micro}$	*ru*	*mi*
MF	PlasmoFP 5%	0.903	2.629	0.958	0.855	2.601	0.379
	PlasmoFP 10%	**0.908**	**2.050**	**0.918**	**0.899**	**1.855**	**0.873**
	PlasmoFP 20%	0.894	2.208	0.858	0.933	1.308	1.779
	PlasmoFP 30%	0.878	2.838	0.817	0.949	1.037	2.642
	CAFA-5 5%	0.478	9.106	0.509	0.451	8.69	2.722
	CAFA-5 10%	0.485	9.006	0.468	0.503	8.145	3.842
	CAFA-5 20%	0.478	9.322	0.422	0.551	7.582	5.423
	CAFA-5 30%	0.472	9.867	0.398	0.579	7.233	6.711
BP	PlasmoFP 5%	0.777	4.716	0.959	0.653	4.708	0.273
	PlasmoFP 10%	0.821	3.780	0.92	0.742	3.721	0.668
	PlasmoFP 20%	**0.831**	**3.140**	**0.844**	**0.818**	**2.66**	**1.669**
	PlasmoFP 30%	0.814	3.587	0.778	0.854	2.173	2.854
	CAFA-5 5%	0.278	24.874	0.587	0.182	24.8	1.908
	CAFA-5 10%	0.311	24.368	0.526	0.22	24.195	2.896
	CAFA-5 20%	0.332	23.941	0.458	0.26	23.495	4.6
	CAFA-5 30%	0.340	23.827	0.415	0.288	22.981	6.293
CC	PlasmoFP 5%	0.780	1.623	0.966	0.654	1.622	0.071
	PlasmoFP 10%	0.830	1.235	0.926	0.753	1.22	0.195
	PlasmoFP 20%	**0.852**	**0.984**	**0.859**	**0.845**	**0.839**	**0.514**
	PlasmoFP 30%	0.839	1.121	0.792	0.892	0.647	0.915
	CAFA-5 5%	0.454	6.462	0.88	0.306	6.45	0.399
	CAFA-5 10%	0.492	6.305	0.831	0.35	6.269	0.668
	CAFA-5 20%	0.537	6.118	0.759	0.415	6.005	1.169
	CAFA-5 30%	0.564	6.022	0.704	0.47	5.782	1.684

Bolded values indicate best performance in each subontology, and optimal eFDR thresholds for each PlasmoFP model

Table 3: PlasmoFP performance on *Plasmodium* holdout set

	Model	F_{micro}	S_{min}	Precision$_{micro}$	Recall$_{micro}$	*ru*	*mi*
MF	PlasmoFP 5% eFDR	0.885	3.822	0.982	0.805	3.816	0.213
	PlasmoFP 10% eFDR	0.900	3.067	0.957	0.850	3.022	0.518
	PlasmoFP 20% eFDR	**0.902**	**2.676**	**0.920**	**0.884**	**2.464**	**1.044**
	PlasmoFP 30% eFDR	0.899	2.567	0.892	0.906	2.007	1.600
	ProteInfer	0.639	8.363	0.746	0.559	7.705	3.252
	DeepGOPlus	0.570	12.240	0.548	0.594	3.691	11.671
BP	PlasmoFP 5% eFDR	0.736	8.157	0.954	0.599	8.146	0.417
	PlasmoFP 10% eFDR	0.783	7.179	0.931	0.675	7.146	0.697
	PlasmoFP 20% eFDR	**0.804**	**6.258**	**0.884**	**0.738**	**6.119**	**1.312**
	PlasmoFP 30% eFDR	0.804	5.899	0.841	0.770	5.537	2.034
	ProteInfer	0.439	12.295	0.598	0.346	10.606	6.220
	DeepGOPlus	0.495	14.831	0.453	0.545	6.760	13.201
CC	PlasmoFP 5% eFDR	0.668	3.623	0.921	0.524	3.619	0.180
	PlasmoFP 10% eFDR	0.717	3.168	0.877	0.606	3.149	0.341
	PlasmoFP 20% eFDR	0.754	2.758	0.838	0.685	2.702	0.551
	PlasmoFP 30% eFDR	**0.774**	**2.493**	**0.810**	**0.741**	**2.364**	**0.793**
	ProteInfer	0.437	4.456	0.599	0.344	4.026	1.908
	DeepGOPlus	0.420	9.561	0.327	0.589	2.757	9.155

Bolded values indicate best performance in each subontology, and optimal eFDR thresholds for each PlasmoFP model

Table 4: PlasmoFP performance (10% eFDR) with predicted intrinsically disordered proteins

	Disorder Content	# Proteins	F_{macro}	S_{min}	Precision$_{macro}$	Recall$_{macro}$	*ru*	*mi*
MF	Low	8911	0.901	2.087	0.910	0.892	1.892	0.880
	Medium	1752	0.897	2.090	0.907	0.886	1.925	0.812
	High	368	0.861	2.457	0.866	0.856	2.226	1.035
BP	Low	6905	0.853	3.679	0.917	0.798	3.626	0.622
	Medium	1512	0.832	3.829	0.898	0.775	3.755	0.744
	High	372	0.803	5.230	0.887	0.735	5.159	0.834
CC	Low	7466	0.880	1.147	0.939	0.828	1.132	0.182
	Medium	1869	0.863	1.384	0.922	0.810	1.363	0.237
	High	443	0.850	1.733	0.922	0.789	1.715	0.234

Metrics are averaged per bin; Low, medium, and high disorder are (0.0-0.3), (0.3-0.6), (≥0.6) respectively across the entire protein

sequence

Fig 1 | PlasmoFP models train on structure representations and utilize deep ensembles (A) Schematic representation of the workflow for training PlasmoFP models using protein language model embeddings. Protein sequences (e.g., X and Y) are processed through the pre-trained ProtT5 language model to generate embeddings (H_X and H_Y). These embeddings are then fine-tuned through the TM-Vec twin network (Φ) to produce sequence-specific representations (H'_X and H'_Y), which serve as inputs to the PlasmoFP models during model training and inference. (B) PlasmoFP models predict GO term probabilities from the processed sequence embeddings (H') via fully connected (FC) layers. (C) Each PlasmoFP model for each subontology is configured as a deep ensemble trained on different k folds of the training dataset. We calculate per-term epistemic uncertainty and create a conservative uncertainty adjusted probability for each input-GO term pair.

Fig 2 | Deep ensembles improve model precision, FDR and sharpness (A) FDR (micro), sharpness, precision (macro) and recall (macro) curves for across all the subontologies for all models (B) Boxplots showing the distribution of MAD for true positive predictions at binned softmax probabilities (C) Average MAD of true positive predictions at differing levels of Gaussian noise added to input TM-Vec embeddings.

Fig 3 | PlasmoFP models predict new functional annotations for *Plasmodium* proteins (A) Fraction of proteins in each species which have annotations in all three GO subontologies after integrating PlasmoFP predictions with existing PlasmoDB annotations at specified eFDR thresholds. The "Existing in PlasmoDB" column refers to curated + IEA annotations found in the current .GAF for the species from PlasmoDB. (B) Fraction of proteins in each species which have annotations in at least one GO subontology in the PlasmoFP + PlasmoDB dataset (C) Fraction of proteins in each species which lack annotations in all three GO subontologies in the PlasmoFP + PlasmoDB dataset.

Fig. 4 | PlasmoFP predicted GO terms co-cluster with existing annotations and expand functional categories across subontologies (A) UMAP projections of Resnik semantic-distance embeddings for GO terms in each subontology (MF: blue, BP: red, CC: green). Gray points are existing annotations; colored points terms predicted by PlasmoFP not found in the existing PlasmoDB annotation across all 19 *Plasmodium* species (B) Heatmaps showing the increase in annotated proteins per functional cluster in the PlasmoFP + PlasmoDB set verses the PlasmoDB annotati

Fig. 5 | PlasmoFP predictions uncover new *Plasmodium* biology (A) RNA- vs. DNA-associated proteome fractions in the *Plasmodium* genus. For each species, bars show the fraction of the annotated proteome labeled as RNA-associated (blue) or DNA-associated (green), before PlasmoFP integration (pale shading) and after adding PlasmoFP predictions to PlasmoDB annotations(solid shading). Normalized to each species' total annotated proteome after integrating PlasmoFP 30% eFDR predictions. (B) Transport & localization counts across human-infecting *Plasmodium* species. Stacked bars for the five major human-infecting species show counts of proteins in Transport & Localization BP clusters mapped to TCDB-inspired categories. Proteins can belong to more than one cluster. Solid fill = PlasmoFP predictions; hatched fill = existing PlasmoDB annotations. (C) Protease-like protein counts across human-infecting *Plasmodium* species. Stacked bars for the five major human-infecting species show numbers of proteins annotated as each protease class (cysteine, serine, metalloprotease, aspartic peptidase, threonine peptidase, and aspartic-type endopeptidase). Solid fill = PlasmoFP additions; hatched fill = existing PlasmoDB annotations.