A comparison of movement-related neuronal activities in cerebellar- and basal ganglia-recipient regions of the macaque thalamus

Daisuke Kase[1], Andrew J. Zimnik[2,3], Thomas Pearce[4], Robert S. Turner[1]

1. *Department of Neurobiology, Center for Neuroscience and The Center for the Neural Basis of Cognition, University of Pittsburgh, Pittsburgh, PA, USA*
2. *Department of Neuroscience, Columbia University Medical Center, New York, NY*
3. *Zuckerman Institute, Columbia University, New York, NY, USA*
4. *Division of Neuropathology, University of Pittsburgh, Pittsburgh, PA, USA*

Address correspondence to:

Robert S Turner, PhD
Department of Neurobiology
University of Pittsburgh
4047 BST-3, 3501 Fifth Avenue
Pittsburgh, PA 15261

Phone: 412-383-5395
Fax: 412-383-9061
email: rturner@pitt.edu

Abstract

The ventral lateral (VL) nucleus of the thalamus relays signals from the cerebellum (Cb) and basal ganglia (BG) to primary motor cortex (M1). In primates, glutamatergic Cb efferents from the deep cerebellar nuclei and GABAergic BG efferents from the internal segment of the globus pallidus (GPi) terminate in distinct subregions of VL: the posterior (VLp) and anterior (VLa) divisions, respectively. This anatomical segregation suggests that Cb- and BG-thalamocortical circuits may play distinct roles in motor control, which could be revealed by comparing movement-related activity in VLp and VLa. Here, we recorded single-unit activity from VLp and VLa, identified via electrical stimulation of superior cerebellar peduncle and GPi, during a choice reaction time reaching task. We also recorded from M1, which maintains bidirectional connections with both VLp and VLa. VLa neurons exhibited a significantly higher proportion of decrease-type responses compared with VLp and M1, consistent with inhibitory GPi input. Time-resolved general linear model analysis showed dynamic encoding of task parameters, particularly movement direction, in all three regions. Direction encoding was strongest in M1, moderate in VLp, and weakest in VLa. Direction encoding in VLa also lagged behind that in M1 and VLp. Clustering analysis of direction encoding strength and timing revealed a subpopulation of VLp neurons that encoded direction particularly strongly during the reaction time period. These results highlight a limitation of traditional assumptions that activity characteristics are distributed homogeneously across neural populations and point to a novel functional organization within VLp neurons.

Introduction

The learning and production of skilled movement is thought to depend on dynamic interactions between primary motor cortex (M1) (21, 23, 49, 67) and multiple subcortical reentrant loop circuits. Chief among these are circuits through the cerebellum (Cb) (22) and basal ganglia (BG) (1); (32). The Cb circuit has been hypothesized to be involved in predictive control, especially of the timing of movement (39, 56) and correction of sensorimotor error (24, 26, 33, 43, 69) whereas the BG circuit may be involved in reward-related selection of action (51, 61), formation of habitual skills (9, 30), and modulation of motor vigor (50, 77, 84). Both subcortical circuits relay signals to primary motor cortex (M1) by way of neurons in the ventral lateral (VL) nucleus of the thalamus. In primates, Cb efferents from the deep cerebellar nuclei (DCN) and BG efferents from the internal segment of the globus pallidus (GPi) terminate in distinct posterior and anterior parts of VL (VLp and VLa, respectively) (7, 20, 34, 57) such that, in primates, convergence of inputs from the DCN and GPi onto individual VL neurons is rare (3, 68, 82), but see (44, 62, 65). This anatomical separation of the DCN- and GPi-recipient thalamic neurons suggests that the distinct roles in motor control hypothesized for Cb- and BG-thalamocortical circuits may be evident through a comparison of the movement-related activities of neurons in VLp and VLa.

The DCN- and GPi-thalamic circuits share many similarities. The synapses of both onto their respective VL thalamo-cortical neurons have morphologic features characteristic of giant "driver-type" synapses (45, 64, 70). Electrical stimulation applied to either of the two projections evokes strong responses in recipient VL neurons, effects interpreted as evidence of a driver-type synaptic function (4, 42, 54, 79); however, see (41). In addition, projection neurons in both DCN and GPi have high firing rates at rest (both typically 50 spikes/sec or higher in primates)(18, 52,

73). One critical difference, however, is that DCN→VLp synapses are glutamatergic whereas GPi→VLa synapses are GABAergic (58, 64, 78, 79).

A default perspective would be that the thalamocortical neurons of VLp and VLa simply relay to cortex the input signals they receive from the Cb and BG, respectively. That view predicts that single-unit activity in VLp should reflect the combined effects of high tonic excitation received from DCN projection neurons and modulations away from that high baseline reflecting the contributions of Cb output to motor cortical function. In contrast, activity in VLa should show the combined effects of high frequency resting inhibition from GPi and phasic modulations in firing rate that reflect BG contributions to motor cortex. It is quite surprising then that the few studies to have investigated this question found only minor differences between the two nuclei with respect to single-unit baseline firing rates or patterns (59), but see (81), and the general sign and timing of movement-related modulations in activity (3, 80). In the rodent motor thalamus, segregation of cerebellar- and basal ganglia-recipient neurons into distinct anatomic regions is less obvious or complete than in primates (13, 44, 60, 62) although see (46). As a result, few neurophysiologic studies of motor thalamus in the rodent have distinguished cerebellar- and basal ganglia-recipient populations (14, 27, 31, 38, 66).

The conventional view of motor thalamus as a relay for cortex-bound subcortical inputs sidesteps the abundant anatomic and physiological evidence that thalamic nuclei are also influenced strongly by inputs from cortex mediated via mono-synaptic excitatory inputs (28, 63), via disynaptic inhibitory inputs through the reticular nucleus of the thalamus (35, 40) and, in primates, via GABAergic intra-thalamic interneurons (5, 28, 36). The anatomy and physiology of these circuits suggest that VLp and VLa are likely to integrate their respective ascending inputs with descending signals received from cortex. While the segregation in the primate thalamus of

inputs from the Cb and BG suggest distinct computational roles for VLp and VLa, the extent to which these pathways are engaged in different dimensions of behavior has seldom been investigated and, as mentioned above, the extant studies described surprisingly minor differences between VLp and VLa (3, 55, 80).

To clarify the roles in motor control of cerebello- and basal ganglia-thalamocortical circuits, we compared the activity of neurons in VLp and VLa during performance of a choice reaction time reaching task. Following existing evidence (3, 55, 80), we predicted that general characteristics, such as baseline firing rates, and the timing and sign of movement-related modulations (i.e. increases or decreases in firing rate), would not differ significantly between VLp and VLa. We suspected, however, that a more detailed examination of the relationships between peri-movement activity and measures of task performance would reveal differences between VLp and VLa populations consistent with the computational roles proposed for their respective Cb and BG inputs. To address these hypotheses, we studied the single-unit activity of electrophysiologically-identified neurons in the VLp and VLa of two non-human primates during performance of a simple well-learned reaching task. We evaluated the dynamic encoding of individual parameters of task performance using a time-resolved general linear model (GLM) analysis. For added context, we compared those results against single-unit data collected from the proximal arm region of M1. In most respects, our results confirm previous reports of overall similarity in firing properties between the two nuclei and extend those findings to general features of the time-resolved encoding of movement parameters. Those results suggest that common inputs to both thalamic regions from motor cortical areas dominate many aspects of the activity of both VLp and VLa, at least in the context of a well-learned highly stereotyped reaching behavior. The regression analysis did reveal a subpopulation of VLp neurons that

encoded movement direction particularly strongly during the reaction time interval. In that VLp population, direction-related partial R^2 reached higher values than those seen in VLa and M1 neurons. In addition, we did find that encoding of direction in VLa lagged behind what was observed in VLp and M1. These results provide clues to the different roles in motor control played by cerebello-thalamic and basal ganglia-thalamic circuits.

Materials and Methods

Animals and behavioral task

Two monkeys (Macaca mulatta; G, female 7.1kg; I, female 7.5kg) were used in this study. All aspects of animal care were in accord with the National Institutes of Health Guide for the Care and Use of Laboratory Animals, the PHS Policy on the Humane Care and Use of Laboratory Animals, and the American Physiological Society's Guiding Principles in the Care and Use of Animals. All procedures were approved by the Institutional Animal Care and Use Committee. The animals performed a choice reaction time reaching task (Fig. 1A) that has been described in detail previously (25, 85)(DOI: 10.5281/zenodo.11397910). In brief, the animal faced a vertical response panel that contained two target LEDs, positioned 7 cm to the left and right of midline. To begin each trial, the animal's left hand rested at a "home-position", a platform located lateral to the left hip at waist height and equipped with a proximity sensor. The animal was trained to hold the home-position (2–10s, uniform random distribution) until illumination of one of the target LEDs, selected in pseudo-random order trial-to-trial, served as both a "Go" signal and a directional instruction. To successfully complete the trial, the animal was required to move its hand from the home-position to the target within 1s of appearance of the Go cue, and hold the

hand at the target position for 0.6-1.2s (random uniform distribution). After the target hold condition was met, a drop of pureed food was delivered via a sipper tube and computer-controlled peristaltic pump. Following reward delivery, the animal was allowed, with no time constraints, to return its hand to the home-position and thereby initiate the next trial. A trial was marked as an error and reward was not delivered if the animal failed to complete the hold period, reached to the incorrect target, or if the 1 sec response window was exceeded. The presence of the hand at the home-position and at targets was detected by infrared proximity sensors (Takex, GS20N).

Surgery

General surgical procedures have been described previously (19, 85). The chamber implantation surgery was performed under sterile conditions with ketamine induction followed by Isoflurane anesthesia. Vital signs (pulse rate, blood pressure, respiration, end-tidal pCO_2, and EKG) were monitored continuously to ensure proper anesthesia. A cylindrical titanium recording chamber was affixed to the skull at stereotaxic coordinates to allow access to the right globus pallidus and ventrolateral thalamus via a parasagittal approach. A second chamber was positioned over the right hemisphere in the coronal plane to allow access to the arm area of primary motor cortex (M1) and, in the midbrain, the decussation of the superior cerebellar peduncle (SCP). The chambers and head stabilization devices were fastened to the skull via bone screws and methyl methacrylate polymer. Prophylactic antibiotics and analgesics were administered post-surgically. We have uploaded a more detailed description of procedures to protocols.io (implantation of head fixation and recording chamber https://dx.doi.org/10.17504/protocols.io.5jyl8p9yrg2w/v1 and craniotomy https://dx.doi.org/10.17504/protocols.io.n92ldmz48l5b/v1).

Localization of stimulation sites and implantation of indwelling macroelectrodes

To allow electrical stimulation-based identification of the cerebellar-recipient VLp and the GPi-recipient VLa (3), we implanted stimulation electrodes chronically in the SCP at its decussation (Fig. 1B top) and the arm-related region of the primary motor cortex. The anatomic locations of sites for implantation were estimated initially from structural MRI scans (Siemens 3T Allegra Scanner, voxel size of 0.6mm) using an interactive 3D software system (Cicerone) to visualize MRI images and predict trajectories for microelectrode penetrations (53). Precise coordinates for each target were then determined using microelectrode mapping methods. And then, custom-built stimulating electrodes were implanted at those sites using methods described previously (74). The custom macroelectrodes consisted of two Teflon-insulated Pt-Ir microwires (50µm) glued inside a short stainless-steel cannula with ~0.5mm separation between the distal ends of the microwires. Insulation was stripped from ~0.2mm of the distal ends of the microwire to achieve an impedance of ~10kΩ. Electrode assemblies for SCP and M1 targets were implanted transdurally via the coronal chamber using a protective guide cannula and stylus mounted in the microdrive. In the months following implantation, the location and integrity of macroelectrodes were monitored by comparing electrode impedances and the current thresholds for activation and the specific muscle contractions evoked by stimulation through each electrode.

Localization of subcortical target regions for recording and stimulation

Many of the target localization methods have been described previously (68). In brief, the chamber coordinates for candidate regions were estimated initially from structural MRIs as described above. Single-unit microelectrode recording was then performed in combination with electrical stimulation (single biphasic pulses <200µA, 0.2ms-duration at 2Hz max.; Model 2100, A-M Systems) and proprioceptive stimulation.

Globus Pallidus interna: To locate the boundaries of the GPi-recipient VLa nucleus of thalamus, we first needed to identify an appropriate location in GPi for electrical stimulation. The target location in GPi was identified by the presence of typical high firing rate single-units, many of which responded briskly to a) proprioceptive stimulation of the forelimb (76) and b) electrical stimulation in the arm region of primary motor cortex (83). We placed a stimulating electrode acutely at this GPi location during the subsequent mapping of VLa boundaries (Fig. 1B bottom).

VL thalamus: The general boundaries of VL thalamus were identified using stereotactic planning under MR guidance and then electrophysiological confirmation as described in (3, 15, 81). Subregions of the VL that were arm-related and likely to be connected with M1 were defined by a) the presence of unit activity typical of thalamus that was modulated by voluntary or passive movements of the animal's proximal arm and b) observation, in a subset of those units, of short latency responses to electrical stimulation of one or more of the macroelectrodes implanted in M1.

VLp thalamus: Within the VL nucleus, as defined above, VLp was identified by its position (34) and the presence of neuronal discharge that: a) responded at short-latency to SCP stimulation (Fig. 1C), and b) did not respond to GPi stimulation (Fig. 1F).

VLa thalamus: VLa was identified by its characteristic position in the anterior-lateral part of VL and the presence of single-unit discharge that: a) responded to GPi stimulation with a short latency pause in firing (Fig. 1G)(3, 55), often followed by a rebound increase in firing probability, and b) showed no response to SCP stimulation (Fig. 1D).

Thalamic microstimulation: Also, we tested the electrical excitability of VL recording sites in accord with previous reports that low current microstimulation evoked muscle contractions only

from locations within VLp and not from VLa (15, 81). We delivered short trains of high

frequency electrical stimulation (glass-insulated tungsten microelectrodes; 10 biphasic pulses

<200µA, 0.2ms-duration at 300Hz; Model 2100, A-M Systems) at sites classified as VLa and

VLp according to the criteria above. Consistent with previous reports, stimulation at many

locations classified as VLp evoked movement at 40-60 µA thresholds (15, 81), while stimulation

at locations classified as VLa rarely evoked movement. On occasion, however, we evoked

muscle contractions from VLa sites that were located near the border with VLp, making

microstimulation an imprecise marker of the boundary between nuclei. Therefore,

microstimulation findings were not used as a primary criterion for identification of the VLa/VLp

border.

Primary motor cortex (M1): Single-unit recordings were obtained from the M1 of one animal

(monkey I). M1 sites for single-unit recording were localized by identification of the principal

arm territory along the anterior bank and gyrus of the pre-central sulcus. As reported extensively,

typical neuronal discharge responded to active and/or passive movement of the arm, and

microstimulation at low current (<60 µA, 10 biphasic pulses at 300 Hz) evoked contraction of

forelimb muscles (8). Single-unit data from the M1 of the other animal were not available due to

technical problems.

Recording protocol

The extracellular spiking activity of neurons in VLp, VLa and M1 was sampled using multiple

glass-insulated tungsten microelectrodes (0.5–1.5MΩ, Alpha Omega Co.) or 16-contact linear

probes (0.5–1.0MΩ, V-probe, Plexon Inc.). Probes were introduced transdurally protected within

custom-sharpened cannulae (Vita Needle Company, Inc.). Voltage signals from each electrode contact were amplified, band-pass filtered (4×, 2Hz–7.5kHz), and digitized at 24 kHz (16-bit resolution; Tucker Davis Technologies). To enhance the mechanical stability of recordings, the 16-contact linear probes were lowered initially to a depth 1 mm below the target depth, then withdrawn and left at the target depth for 30min before initiating data collection. When stable single-unit isolation was available from one or more single-units in either VLp or VLa, as determined by online spike sorting, neuronal data and behavioral events were collected while the animal performed the behavioral task. On protocols.io, we have posted protocols relevant to electrophysiological recording (https://dx.doi.org/10.17504/protocols.io.bp2l6xx91lqe/v1) and recording chamber maintenance (https://dx.doi.org/10.17504/protocols.io.5jyl8ppm9g2w/v1).

Offline analysis

Behavior

Task performance was screened and error trials and outliers in task performance were excluded from further analyses. Error trials included reaches to the incorrect target and failure to reach the target within the allowed 1s interval after the go cue. Two behavioral metrics, reaction time and movement duration, were used for outlier detection. Reaction time (RT) was defined as the interval between the go cue and the lift of the hand off the home position as detected by the infrared sensor, termed movement onset. Movement duration (MD) was defined as the interval between movement onset and target capture, which was detected by an infrared sensor at the target. Trials were classified as outliers if either behavioral metric exceeded a threshold of 6 median absolute deviations from the median (Matlab: ISOUTLIER). For trials that met the above inclusion criteria, average movement speed was computed by dividing the distance from the home position to the target (left: 35cm; right: 42cm) by the MD.

Spike sorting and detection of peri-movement discharge

The stored wide-band neuronal signal was high-pass filtered (Fpass: 300Hz, Matlab FIRPM), manually thresholded, and candidate action potentials were sorted into clusters in principal components space (Off-line Sorter, Plexon Inc.). Clusters were accepted as well-isolated single-units only if the unit's action potentials were of a consistent shape and could be separated reliably from the waveforms of other neurons as well as from background noise throughout the period of recording. Single-units from VLp, VLa and M1 were included for further analysis if they exceeded minimum firing rate thresholds (0.9Hz for VLp and VLa, 0.1Hz for M1, averaged over the entire recording session). A minimum of 20 valid behavioral trials (minimum of 10 in each direction) was required to include a unit in task-based analyses.

Measures of baseline activity were calculated from neural activity sampled during start position hold period before the presentation of the go cue. Measurements of the spike waveform were calculated from spike data that were up-sampled to 244 kHz using linear interpolation (Matlab INTERP1) and then averaged. Action potential width was calculated as the full width at half-maximum of the first trough. Burst-like events were detected using the Poisson surprise method as described previously ((47); initial filtering threshold factor: 1.5, minimum spike number of the burst: 3, surprise cutoff: 5).

We tested for peri-movement changes in single-unit spike rate using a standard rate change detection method (2, 85). First, spike density functions (SDFs) were constructed by convolving each unit's spike time stamps (1 kHz resolution) with a Gaussian kernel ($\sigma = 25$ ms). SDFs were then aligned to the time of movement onset for each trial and across-trial mean SDFs were constructed separately for reaches to each target. The baseline activity for each unit was calculated as the mean and standard deviation of the SDF over a 700ms window ending at the

median time of target LED onset, correcting for linear trends in the SDF over this window. This baseline activity was then used to detect movement-related changes in the SDF during a peri-movement period that extended from the median time of target LED onset (go-cue) until the median time of target capture, thereby encompassing both reaction time and movement duration intervals. We defined a movement-related change in activity as a statistically-significant elevation or depression in the SDF from baseline that lasted at least 60ms (e.g., Fig.2 A, solid vertical lines; t-test, one-sample vs. baseline; omnibus p < 0.001 after Bonferroni correction for multiple comparisons). All movement-related changes in firing rate were classified as increases or decreases in discharge rate and the time of onset of that change was defined as the first time point in the SDF that crossed the threshold for significance. Note that this approach enabled detection of biphasic changes (e.g., an increase followed by a decrease), but further analysis was based only on the earliest-occurring peri-movement change in firing.

Time-resolved general linear model (GLM) regression

We used general linear model regression to determine if single-unit firing rate encoding of measures of task performance differed between VLp and VLa neurons. We used the Matlab command 'lassoglm' with a normal regression model, 10-fold cross validation (to estimate the reliability of results) and an 'alpha' set to 0.5, thereby implementing elastic-net regularization of potential covariation between regressors. The regressors included trial-by-trial values for parameters controlled by the task (target direction and duration of the start position hold period before go-cue presentation), measures of how the animal performed the task (reaction time and mean movement speed), and the elapsed time since the beginning of the recording session. Target direction was coded as +1 (right, i.e., contralateral to the working arm) and -1 (left), and the remaining regressors were z-scored to facilitate the comparison of results between regressors.

We assessed collinearity using Belsley's collinearity diagnostics with the Matlab command 'collintest' (11). A condition index threshold of 10 was used to indicate acceptable level of low collinearity (10). To ensure that the resulting model was sparse, the regularization penalty value lambda was set to the greatest value within one standard deviation of the lambda that minimized cross-validation error. The neural response function to be predicted was a time series of one unit's spike counts across peri-movement time for each trial. More specifically, we calculated spike counts within a 200ms wide window at 1200 time points starting centered at 700ms before movement onset and swept in 1ms steps to 500ms after movement onset. An independent lassoglm was run on the trial-by-trial spike counts at each 1ms time step of the response function. Coefficients of partial determination (partial R^2 values, pR^2) for each regressor were defined as the proportion of across-trials variation in spike counts explained by the full regression (sum of squared errors) that could not be explained by a reduced model that excluded that regressor.

Then, for each single-unit, we estimated the timing of peri-movement changes in the encoding of individual regressors. We tested for significant increases in a regressor's pR^2 during a peri-movement test period that extended from the latest time across trials of go-cue presentation relative to movement onset (i.e., beginning at the shortest RT prior to movement onset) to the earliest end of reach (i.e., ending at the shortest MD following movement onset). If the maximum partial R^2 within the test period reached a value greater than 0.01 then it was classified as a significant increase in partial R^2. The time of onset of that increase was defined as the time immediately prior to the maximum when the partial R^2 crossed a 5% of maximum threshold.

Cluster analysis with Gaussian mixture model

We tested the GLM results for sub-groups (clusters) using Gaussian mixture models (MatLab function: fitgmdist). Those models used full covariance matrices, 10000 iterations, 0.01 of the regularization value, and the number of clusters tested for ranged from one through five. The best-fit number of clusters was identified as the number associated with a minimum in the Bayes Information Criterion (BIC) calculated in the MatLab function (fitgmdist).

Results

Basic approach and task performance

We studied the single-unit activity of neurons sampled from VLp and VLa of two macaque monkeys and from M1 of one macaque monkey during skilled performance of a choice reaction time reaching task for food reward (16, 25, 68, 85) (Fig. 1A). Animals performed the reaching task in a highly stereotyped fashion. Reaction times (248.1 ± 33.9 ms and 237.8 ± 42.7 ms for monkey G and I respectively; mean ± SD) and movement durations (249.8 ± 36.0 ms and 199.6 ± 32.8 ms for monkey G and I respectively; mean ± SD) were short and showed a low level of variability. Animals made very few errors in task performance (success rates of 97.5% and 96.3% for monkey G and I, respectively), and the rare error trials were excluded from further analysis.

Identification of VLp and VLa

One or two microelectrodes, or one 16-contact linear probe, were positioned acutely in the arm-related regions of one or the other thalamic nucleus or, occasionally, both nuclei (32). We used a combination of electrophysiological techniques to classify the recording site as described previously (see Methods, (68)). In brief, VLp regions were identified by the presence of neurons that were excited at short-latency in response to electrical stimulation of the SCP (Fig. 1C) and were not inhibited at short-latency following electrical stimulation of GPi (Figs. 1F). In contrast, VLa regions were identified by the presence of neurons that were inhibited at short-latency following stimulation of GPi (Fig. 1G) and were not excited by stimulation of SCP (Figs. 1D). For most recording tracks, the locations of borders between the VLp and VLa regions were

determined during preliminary mapping penetrations performed prior to data collection (supplemental Fig. S1). The sub-regions of VLp and VLa involved in arm movement were identified by the presence of short-latency spiking responses to stimulation of arm-related areas of M1. Based on their physiologic responses during mapping, neurons assigned to VLp and VLa categories were found in separate regions of thalamus. Anatomic intermixing of neurons categorized as VLp and VLa was exceedingly rare, consistent with previous reports (20, 34, 57, 68). In rare cases, a neuron encountered close to the electrophysiologically-defined boundary of VLp and VLa territories was both excited by SCP stimulation and inhibited by GPi stimulation. Similar observations have been described previously (55). To avoid potential ambiguities associated with studying such "boundary neurons," we restricted all subsequent analyses to cells located at least 50um away from the electrophysiologically defined border.

Neuronal recordings and activity at rest

A total of 83 single-units were sampled from the VLp and 193 single-units were sampled from VLa according to the criteria defined above (see *Methods* for more detailed descriptions). 172 neurons were sampled from the proximal arm-related region of M1 of one animal and are included here for comparison. Units were studied over the course of 132.8 ± 77.3 trials of the behavioral task (mean ± SD) with a mean of 66 trials for each of the two movement directions.

Contrary to expectations based on the distinct synaptic inputs received by VLp and VLa neurons, past studies of the baseline activities of those two populations found negligible differences (3, 59). To revisit that question, we quantified neuronal firing properties across each trial's start position hold period, which amounted to multisecond-long periods of attentive immobility. Consistent with some (3, 59) but not all (81) previous reports, we found that mean firing rates during the hold period were statistically indistinguishable between VLp and VLa

neurons (Wilcoxon rank test; p=0.29; Table 1). Measures of firing rate variability such as the coefficient of variation of inter-spike intervals (CV), proportion of spikes in the burst activity, and the normalized intra-burst firing rate also did not differ significantly between VLp and VLa neurons (Wilcoxon rank test; p=0.38, 0.48, and 0.14 respectively; Table 1). In addition, two measures of action potential duration (half width of the first trough and trough to peak time) did not differ between nuclei (Wilcoxon rank test; p=0.29 and 0.58, respectively; Table 1), again contrary to expectation (60). The ratio of trough magnitudes to peak magnitudes did not differ between nuclei (Wilcoxon rank test; p=0.94).

Relative to the activity of M1 neurons, neither thalamic nucleus differed significantly in resting firing rates (Wilcoxon rank test; p=0.42 and 0.80 for VLp and VLa respectively). Thalamic neurons, however, were more likely than M1 neurons to fire action potentials in irregular and bursty firing patterns, as evidenced by elevations of the CV of inter-spike intervals and the fraction of spikes in the burst (Wilcoxon rank test; p<0.01; Table 1). Finally, consistent with previous work (71, 72), the action potentials of thalamic neurons were prolonged in duration as compared with M1 neurons (half width of the first trough and trough-to-peak times; Wilcoxon rank test; p<0.01). Considering data from NHP I separately (ie., the animal M1 neurons were recorded from) the trough-to-peak magnitude ratio was larger in thalamic neurons than in M1 (Wilcoxon rank test; p<0.01 and p<0.05 for VLp and VLa respectively; Table 1).

In summary, the baseline activities of neurons in VLp and VLa were indistinguishable from each other in nearly every respect, despite the markedly different subcortical inputs the two thalamic nuclei receive. The baseline activity found in both thalamic nuclei differed from that of M1 across multiple measures in ways that were consistent with previously described generic differences between cortical and thalamic extracellular activity (71, 72).

Peri-movement decreases in firing are more common and longer-lasting in VLa than in VLp

Next, we compared the general features of peri-movement modulations in firing of neurons in VLp and VLa. We predicted that the sign and timing of movement-related modulations in activity would not differ substantially between nuclei, consistent with previous reports (3, 80), but contrary to the common assumption that task-related activity in VLp and VLa is determined primarily by their respective ascending glutamatergic inputs from DCN and GABAergic inputs from GPi. Any differences in response prevalence, form or timing may provide clues into the influences in VL thalamus of those distinct subcortical inputs.

To that end, we tested mean spike density functions (SDFs; see *Methods*) for peri-movement changes in firing relative to a unit's baseline activity (i.e., linear trend ± SD of activity prior to go-cue presentation, *blue slanted lines* in Fig. 2A). Significant peri-movement changes were detected in large fractions of all neural populations (89%, 89% and 96% in VLp, VLa and M1, respectively (chi-square test; $\chi^2(2, n=448) = 0.74$, p=0.69)). The shape and timing of those responses differed widely between neurons and occasionally between directions of movement. We observed both monophasic and poly-phasic responses in all areas. Monophasic responses were composed of a single increase or decrease in firing (Fig. 2 B-D; *Incr* and *Decr*, respectively). Poly-phasic responses were classified according to the sign of the initial response (i.e., an initial increase or decrease in firing rate; Fig. 2 B-D; *Poly (+/-)* and *Poly (-/+)*, respectively). All detected responses are shown in Fig. 2B sorted according to brain area, response type, and latency of response onset relative to the time of movement initiation (time 0). Population averages of these response types for each brain area are shown in Fig. 2C. The rate of occurrence of the four general response types differed between the three brain areas (chi-square

test; $\chi^2(3, n=604) = 32.78$, p<0.01; Fig. 2D). Mono-phasic decreases were more common in VLa than in VLp or M1 whereas the poly-phasic decrease-increase response type was less common in VLa than in VLp or M1 (p<0.01, post-hoc adjusted standardized residual analysis, Haberman 1973). In addition, the poly-phasic increase-decrease type of response was less common in VLa than in M1 (p<0.05, adjusted standardized residual analysis).

We measured the properties of individual monophasic responses and then compared them between VLp, VLa and M1 populations (Table 2). (The small numbers of polyphasic responses detected, particularly in VLa, prevented a similar comparison of those response types.)

For increase-type responses, only one measure differed significantly between VLp and VLa populations: responses began earlier relative to movement onset in VLp than in VLa (mean 30 msec earlier; Wilcoxon rank test; p<0.01; Table 2). Increase-type responses in VLp and in M1 began at similar pre-movement onset times. The magnitudes and durations of increase-type responses were statistically indistinguishable between VLp and VLa (Wilcoxon rank test; ps≥0.1), while those in M1 were markedly larger and shorter duration (Wilcoxon rank test; ps<0.01; Table 2, *Magnitude, Slope and Half-width*).

For decrease-type responses, onset latencies did not differ significantly between the two thalamic nuclei or relative to M1. Similar to what was seen for increase-type responses, response magnitudes were indistinguishable between VLp and VLa while those in M1 were larger (Wilcoxon rank test; all p<0.05; Table 2, *Magnitude* and *Slope*). Finally, the duration of decrease-type responses was significantly longer in VLa than in M1 (Wilcoxon rank test; p<0.01), while in VLp the duration of decreases was intermediate between VLa and M1 (Wilcoxon rank test; p=0.16).

To summarize, we found two notable exceptions to our general prediction that peri-movement responses do not differ between VLp and VLa (3, 55, 80). First, increase-type responses began earlier in VLp than they did in VLa. That result is consistent with the idea that movement-related signals in the cerebello-thalamocortical circuit have an early influence on the timing of movement initiation (17, 39, 56), whereas signals in the basal ganglia-thalamocortical circuit do not (29, 68). We also found that decrease-types responses were more common in VLa than in VLp. That difference might be a product of the strong inhibitory input that VLa receives from GPi. For the most part, however, peri-movement responses in VLp and VLa were statistically indistinguishable, contrary to what might be predicted from the categorically-different sub-cortical inputs that the two nuclei receive. We next tested whether the activity of VLp and VLa might be more readily differentiated by the dimensions of task performance encoded by their neuronal activity or by the timing of that neural encoding.

Time-resolved regression identified differences in encoding

We used time-resolved linear regression analyses to determine if peri-movement activity differed between VLp, VLa, and M1 with respect to the dimensions of task performance encoded or the timing of that encoding. Between-trial variations in a unit's peri-movement firing rate were regressed against the following trial-by-trial measures of task performance: 1) duration of the start position hold period (prior to presentation of the go-cue), 2) reaction time, 3) mean movement speed, 4) reach direction, and 5) elapsed time from the beginning of the recording session ("session time"; see *Methods*). The collinearity between predictors was acceptably low (Belsey's collinearity: 3.39, 4.18 and 2.49 for predictors from VLp, VLa and M1 recording sessions, respectively (10). Trial-by-trial time-evolving estimates of firing rate were regressed

against those five predictors. All units were included in this analysis regardless of the response type observed.

The regressions identified relationships between one or more predictors in large proportions of all neural populations (63%, 53% and 74% of units in VLp, VLa and M1, chi-square test; χ^2(2, n=448) = 17.33, p<0.01). Typical regression results are shown for a single VLp unit in Fig. 3A. During the start position hold period, prior to go-cue presentation (Fig. 3A; median shown by *vertical solid line* and IQR by *dark gray* shading), the full model accounted for only a small fraction of the variance in firing rate (5% approx.; R^2 for full model, *black trace* Fig. 3A top), essentially all of which was attributable to a significant relationship between firing rate and the time elapsed in the recording session (partial R^2 , "pR^2", for session time, *red trace* Fig. 3A top). Small but consistently negative beta values for that relationship between firing rate and session time (*red trace* Fig. 3A bottom) indicated that firing rates during this period declined slowly across the whole recording session. Other regressors accounted for negligible fractions of firing rate variance during the delay period. Immediately following the median time of go-cue presentation (240 +/- 32 ms prior to movement onset) the full model R^2 increased sharply and reached a peak of 26% of variance at 154 ms prior to movement onset. That increase was attributable to a strengthening of the relationship between firing rate and session time and the appearance of a strong relationship to movement direction (pR^2 for direction; *blue trace* Fig. 3A top). A second, long-lasting elevation in R^2, which began 50 ms before movement onset, was composed of successive peaks in pR^2 for movement direction, movement speed (*yellow trace*), delay period (*gray trace*) and then a third large peak for direction (*blue*).

We analyzed the number and combination of predictors that showed significant increases in pR^2 values during the peri-movement test period ("test period"; see *Methods*). The magnitudes

of peak pR^2 values of all predictors for all units are summarized in heat maps (Fig. 3B), where each row represents the pR^2 coefficients for a single unit for each of the five predictors (columns). Target direction was the most common predictor for all neural populations. Similar fractions of neurons encoded target direction in VLp and VLa populations (47% and 40% respectively; chi-square test; χ^2(1, n=276) = 1.20, p=0.27) although those fractions were smaller than the fraction of M1 neurons found to encode direction (66%; chi-square test; χ^2(2, n=448) = 26.11, p<0.01). Progressively smaller fractions of neurons in each population encoded session time and movement speed. Only a small number of neurons showed significant effects for RT and delay predictors. We found no consistent pattern to the way encodings of multiple predictors were distributed across populations. Therefore, from here-on we will focus primarily on results for the full model and the target direction and movement speed regressors.

To compare the timing and strength of task encoding, we constructed time-resolved population averages of overall R^2 values for the three neural populations. Those curves began to rise at approximately 200ms before movement onset for all neural populations (Fig. 3C top). Encoding was stronger in VLp than in VLa throughout the reaction time and move execution periods. Interestingly, task encoding was stronger in the M1 population than in either of the thalamic areas (Fig. 3C top). Indeed, the times of onset of significant encoding averaged across individual neurons confirmed that the onset of encoding did not differ between populations (Wilcoxon rank test; between all pairs; ps>0.23; Fig. 3C middle). Neuron-by-neuron measures of the peak magnitude of overall R^2 also confirmed the observation that overall encoding of task parameters was weaker in VLp and VLa than in the M1 population (Wilcoxon rank test; ps<0.01; Fig. 3C bottom). Interestingly, we found no significant difference between VLp and VLa populations in the mean magnitude of peak R^2 (Wilcoxon rank test; p=0.24), contrary to what the

population average curves in Fig. 3C top might suggest. We revisit this disparity in the next section.

Next, to identify potential differences between brain areas in their relationship to the encoding of target direction, we computed pR^2 and beta values separately for each neural population and compard the onsets and peak magnitudes of significant encoding between VLp, VLa and M1 populations (Fig. 3D).

Encoding of direction appears earlier and is stronger in VLp than in VLa

For both VLp and M1 populations, the population averages of the pR² values for target direction were similar to the population averages of overall R² values in terms of time course and magnitude of change. The curves of pR² values began to rise approximately 200 ms before movement onset (Figs. 3C top and D top, *magenta* and *yellow*). In contrast, for VLa the rise in direction encoding was more gradual, lagging well behind that of overall R² values (Figs. 3C top and D top, green). Comparison of the mean time of onset of direction encoding across individual neurons confirmed that in VLa the appearance of direction encoding was delayed relative to that in VLp or M1 (Wilcoxon rank test; $p < 0.05$ for VLp, $p < 0.01$ for M1).

The strength of direction encoding was also similar for VLp and M1.

When comparing overall R² values and partial R² values for target direction to evaluate the contribution of target direction encoding to overall encoding, we found that VLp and VLa showed similar magnitudes of change between these metrics (Fig. 3D top, magenta and green). In contrast, in M1, the changes in partial R² values were smaller than those in overall R² values, suggesting that other predictors had a greater influence on overall R² in M1 compared to VLp and VLa (Fig. 3D top, yellow). The close alignment of overall and partial R² magnitudes in VLp

and VLa suggests that target direction encoding was the primary contributor to the overall R^2 in these regions. Neuron-by-neuron measurements of peak partial R^2 magnitudes revealed significant differences across brain areas (Fig. 2D second from bottom). Specifically, the partial R^2 magnitudes in both VLp and VLa populations were significantly smaller than in the M1 population (Wilcoxon rank test; $p < 0.05$ for VLp, $p < 0.01$ for VLa). However, despite the difference in population averages of partial R^2 values between VLp and VLa, this difference was not statistically significant at the individual neuron level (Wilcoxon rank test; $p = 0.06$).

Encoding of mean speed begins early in VLa

The population averages of the partial R^2 values for mean speed, derived from the speed-encoding neuron population, were different from the population averages of overall R^2 values in VLp, VLa and M1. The curves of partial R^2 values began to rise approximately 200 ms before movement onset, but the mean magnitude was smaller in VLp compared to VLa and M1 (Fig. 3E top, magenta). In VLa, the partial R^2 value for mean speed began increasing immediately after go-cue presentation, with strong encoding observed in the reaction time (Fig. 3E top, green). In contrast, mean speed encoding in M1 was stronger during the reach phase (Fig. 3E top, yellow). Quantitative analyses of the onset of significant encoding and the magnitude of partial R^2 value in individual neurons aligned with trends observed in the population averages of partial R^2 and beta values (Fig. 3E middle to bottom).

Clustering analysis identified two types of target direction encoding in VLp

In the population-averaged curves of both overall and partial R^2 values for target direction, we observed clearly distinct encoding patterns between VLp and VLa populations (Figs. 3C top and 2D top). However, neuron-by-neuron analysis of peak encoding magnitudes did not show statistically significant differences (Figs. 4A). We hypothesized that this discrepancy might arise from large variability in encoding magnitudes within the VLp population. To test this hypothesis, we compared the distributions of peak magnitudes for partial R^2 values of target direction across the three populations (Fig. 4A). The cumulative distribution functions (CDFs) for partial R^2 values in VLa and M1 populations displayed smooth, logarithmic curves (Fig. 4A, green and yellow). In contrast, the CDFs in the VLp population showed a plateau in the middle, suggesting that the magnitude distribution in VLp may arise from multiple underlying sources (Fig. 4A, magenta).

The presence of multiple subpopulations with varying peak magnitudes of partial R^2 in VLp may help explain the high p-values observed in our statistical tests. We focused on target direction encoding because most of the overall R^2 values in VLp and VLa arise from the partial R^2 values associated with target direction (Figs. 3C top and 3D top, magenta and green). In addition to the cumulative distribution function (CDF) of partial R^2 magnitude, we also observed a plateau in the CDF of encoding onset in VLp (Fig. 4B, magenta).

To further analyze the distribution of peak partial R^2 magnitudes within the VLp population, we created a scatter plot showing the relationship between peak partial R^2 magnitude and encoding onset (Fig. 4C left). This plot suggested two clusters within VLp: one with high peak partial R^2 values and early encoding onset, and another with low peak magnitudes and a broad range of encoding onsets. To test this, we fitted Gaussian mixture models to the

distributions of both peak magnitudes and encoding onsets, testing models with 1 to 5 clusters. The optimal number of clusters was determined by comparing the Bayesian Information Criterion (BIC), with a smaller BIC indicating a better fit. In VLp, the two-cluster model yielded the lowest BIC (Fig. 4C left inset). In the scatter plot, markers are color-coded to indicate cluster membership (Fig. 4C). One VLp cluster demonstrated early and strong target direction encoding (Cluster 1: dark magenta markers), while the other displayed a wide range of onset times and weaker encoding (Cluster 2: light magenta markers). Similar clustering analyses for VLa and M1 populations suggested single, unified clusters in both regions (Fig. 4C middle and right).

Finally, we analyzed the metrics of clusters 1 and 2 in the VLp population and compared them with those in VLa and M1. The magnitude of encoding in VLp cluster 1 neurons was significantly greater than those in all other populations (Wilcoxon rank test; all $p < 0.01$; Fig. 4D). The onset of encoding in VLp cluster 1 neurons was also significantly earlier than in VLa (Wilcoxon rank test; $p < 0.05$; Fig. 4E). Conversely, the magnitude of encoding in VLp cluster 2 neurons did not differ significantly from that in VLa but was significantly smaller than in M1 (Wilcoxon rank test; $p = 0.89$ for VLa and $p < 0.01$ for M1; Fig. 4D). Similarly, the encoding onset of VLp cluster 2 neurons was not significantly different from that in either VLa or M1 (Wilcoxon rank test; $p = 0.15$ for VLa and $p = 0.27$ for M1; Fig. 4E).

We observed similar plateau in the CDF of magnitude of partial R^2 values for mean speed in VLa (Fig. 4B, magenta). However, clustering analyses for mean speed encoding did not reveal a multi-cluster structure in any of the regions analyzed.

Although VLp and VLa did not significantly differ in peri-movement SDF, linear regression analysis indicated that VLp neurons tended to encode target direction earlier and more strongly than VLa neurons. Our clustering analysis using Gaussian model fitting showed that a

two-cluster model produced the smallest BIC for target direction-encoding VLp neurons. This result suggests that target direction-encoding neurons in VLp can be divided into two groups: one group that begins encoding target direction during the reaction time with strong encoding, and another group that encodes target direction weakly, with encoding onset distributed throughout the response time. This clustering pattern was not observed in VLa or M1 and appears to be a unique property of VLp.

Discussion

Strong inhibitory input in VLa

Although peri-movement activity patterns in VLp and VLa were previously reported to be similar (3, 80), our analysis revealed several indications of strong inhibitory input in VLa. Specifically, the proportion of neurons exhibiting a monophasic decrease in firing rate around movement was significantly higher in VLa compared to VLp (Fig. 2D). Quantitative analysis of the spike density function (SDF) revealed a prolonged decrease in VLa, although the duration was not significantly longer than that observed in VLp (Table 2). These findings suggest that VLa neurons receive stronger inhibitory input than VLp neurons.

The globus pallidus internus (GPi) is the most likely source of this inhibitory input to VLa. This hypothesis is supported by the unique giant synaptic structures formed by inhibitory projections from the basal ganglia output nuclei to the motor thalamus (12, 37, 64) and by studies

demonstrating inhibitory effects of GPi electrical stimulation on thalamic activity (4, 42, 54, 79). Other GABAergic sources, such as the reticular thalamic nucleus (nRT) and local interneurons, may also contribute to the stronger inhibitory input in VLa. However, these GABAergic projections are known to target VLp as well, and previous studies suggest that nRT and interneuron inputs may actually be stronger in VLp than in VLa. For instance, axons from motor nRT neurons predominantly form synapses on the distal dendrites of thalamocortical neurons in VLa, whereas in VLp, nRT axons synapse directly on the soma and proximal dendrites (35). Additionally, cortico-thalamic projections from M1 form synapses on the dendrites of interneurons in VLa but form presynaptic terminals of interneurons in VLp (28).

Thus, given the relatively weaker inhibitory contributions from nRT and local interneurons in VLa, the greater prevalence of decrease-type peri-movement responses is most likely a product of the strong inhibitory input to VLa from the GPi.

Small fraction of VLp neuron encode the target direction strongly

Previous studies reported similar activity patterns in VLp and VLa during movement. Those studies, however, did not test for differences between nuclei in the neural encoding of task parameters. To address this, we performed time-resolved linear regression analyses using behavioral parameters. Among these, target direction emerged as the most influential predictor across all examined target areas (Fig. 3B). This result aligns with prior findings of target direction encoding in the regions upstream of VLp and VLa [e.g., the deep cerebellar nuclei (26) and GPi (76)].

The distribution of target direction encoding differed, however, between VLp and VLa (Fig. 4C). Clustering analysis revealed that the best-fitting model for VLp target direction

encoding units was a two-cluster model (based on the lowest BIC), whereas the same analysis for VLa units supported a single-cluster model. Using the clustering results, we recalculated the metrics of target direction encoding and found that the subgroup of VLp neurons with smaller partial R^2 values for target direction were indistinguishable from VLa neurons in their encoding metrics (Fig. 4D-E).

These findings suggest that only a subset of VLp neurons receives strong target-direction-related input from their upstream brain areas.

Potential encoding of behavioral parameter by a few impactful thalamic neurons

Similar to the distribution of partial R^2 values for target direction, we observed an irregular CDF for partial R^2 values associated with mean speed in VLa (Supplemental Fig. S2B, top). The encoding of mean speed in the VLa aligns with previous studies that demonstrated the effect of transient GPi inactivation on movement vigor ((77) and references therein). Additionally, the observation that only a small fraction of VLa neurons strongly encode mean speed is consistent with a prior report of peak speed encoding in a subset of GP neurons (75). However, our clustering analysis did not identify multi-cluster models as the best fit for this dataset. This result of clustering analysis may be due to the small number of VLa neurons that strongly encoded mean speed, despite a total sample size of 193 VLa neurons, which is not insubstantial.

One potential implication of these findings is that both strong target direction encoding in VLp and strong speed encoding in VLa are represented in small subpopulations of neurons within these recipient VL nuclei. This hypothesis is supported by histological evidence showing that axons from both the dentate nucleus and GPi form focal terminal fields in their target

regions (6, 37, 48). These studies suggest that individual axons may selectively influence small clusters of neurons.

This model could also explain the consistently small differences between VLp and VLa when compared at the whole population level. Future research using high-density probes capable of capturing neural activity from neurons in these small clusters may provide a more definitive understanding of these patterns.

Conclusion

The anatomical separation of Cb- and BG-recipient thalamic neurons suggests that the distinct roles of Cb- and BG-thalamocortical circuits in motor control should be evident when comparing the movement-related activities of neurons in VLp and VLa. Surprisingly, however, previous studies reported similar peri-movement activity in these nuclei during behavioral tasks (3, 80). Our analysis revealed that the fraction of neurons showing a decrease-only response was significantly larger in VLa than in VLp. While quantitative analyses of peri-movement activity indicated a trend toward stronger peri-movement increase in firing rate in VLp and stronger peri-movement decrease in firing rate in VLa, these differences were not statistically significant.

In contrast, our regression and clustering analyses highlighted differences in the encoding of the behavioral parameter associated with target direction between VLp and VLa. Specifically, we identified two groups of VLp neurons with differing degrees of target direction encoding. In contrast, the same clustering analysis suggested a single-cluster distribution for VLa neurons. One group of VLp neurons exhibited strong early target direction encoding during the reaction

time, while the other group's encoding was indistinguishable from the target direction encoding observed in VLa.

Future studies employing higher spatial resolution recording techniques and pharmacological inactivation of upstream brain areas will be critical for further elucidation of the functional differences between VLp and VLa within the motor control system.

Data Availability

Both electrophysiological data and behavioral data are available on the DANDI Archive at DANDI:000947/0.240510.2211. All MatLab code and the processed source data sufficient to reproduce all figures and tables are in preparation and will be available on Zenodo (<u>TO</u> BE FILLED).

Author Contributions

RST conceived and designed research; DK, AJZ, and RST performed experiments; DK analyzed data; DK, TP, and RST interpreted results of experiments; DK prepared figures; DK, and RST drafted manuscript; DK, AJZ, TP, and RST edited and revised manuscript and approved final version of manuscript.

Acknowledgment

We thank Lisa Nieman-Vento and Cherie Lee Cornmesser for their contributions to animal care.

Research reported in this publication was supported by the National Institutes of Health under award number R01NS113817. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Disclosures

The Authors declare no interest.

Figure and Figure Legend

Figure 1

A. Schematic of the task. B and E. Schematic illustration of the stimulation electrode tracks targeting the superior cerebellar peduncle (SCP) (B) and GPi (E). Scale bar: 10 mm. C-D and F-G. Exemplar peri-stimulus time histograms of neuronal spike activity averaged around the times of stimulation (time 0) for a single unit located in VLp (C, F) and a unit located in VLa (D, G).

The top panels (C-D) show responses to SCP stimulation, and the bottom panels (F-G) show responses to GPi stimulation.

Figure 2

A. Peri-movement activity of exemplar single units sampled from VLp (left), VLa (middle), and

M1 (right). Across-trial mean spike density functions (SDFs; top) and raster plots (bottom) are

aligned to movement onset. *Sloped blue lines:* the unit's mean baseline trendline ± confidence

interval. *Vertical cyan line*: the estimated onset of the mean response. In rasters, single-trial spike

trains are sorted by reaction time (shortest at top). Red circles indicate go-cue onset for each trial.

B. Peri-movement SDFs of all recorded single units, sorted by response type and onset latency

(earlier responses at the top within each category) in VLp (left), VLa (middle), and M1 (right).

SDFs were z-scored relative to mean baseline activity prior to the go-cue onset. C. Population

averages of z-scored SDFs. Top: monophasic responses (red: increases only; blue: decreases only). Bottom: polyphasic responses (red: increase–decrease [+/-]; blue: decrease–increase [-/+]). Data shown for VLp (left), VLa (middle), and M1 (right). *Shaded areas*: ±SEM. D. Proportions of peri-movement response types in VLp (magenta), VLa (green), and M1 (cyan). The proportion of decrease-only responses in VLp and M1 was significantly smaller than the proportion estimated from all responses, whereas the proportion of polyphasic responses in VLp and M1 was significantly larger than expected (*p<0.05, **p<0.01; standardized residual analysis). Magenta and cyan asterisks denote significant effects in VLp and M1, respectively.

Figure 3

A. Exemplar regression results for activity of a single VLp neurons influenced by multiple

predictors. Top: R² values. Bottom: beta values. Black: full model; orange: mean speed;

magenta: target direction; blue: delay period; green: session time. Vertical black dashed lines and

gray rectangles mark the median and interquartile range (IQR) of go-cue onset times (left) and

movement end times (right), aligned to movement onset. B. Peak partial R^2 values for all single units, sorted according to the most effective predictor and peak magnitude (larger partial R^2 values at the top within each predictor category). C. Population averages of R^2 values for VLp (magenta), VLa (green), and M1 (yellow). *Shaded areas*: ±SEM. D–E. Population averages of partial R^2 values (top) and beta values (bottom) associated with target direction (D) and mean speed (E). Vertical black dashed lines indicate the median times of go-cue onset (left) and movement end (right). Shaded areas represent ±SEM. C–E. Horizontal lines above each panel indicate Wilcoxon rank test results for early, middle, and late epochs of the response interval (thin black line: $p < 0.05$; thick black line: $p < 0.01$; Wilcoxon test with p-values corrected for multiple comparisons using false discovery rate; horizongal dotted line: not significant).

Figure 4

A. Cumulative distributions of the peak magnitudes of partial R² values associated with target

direction in VLp (magenta), VLa (green), and M1 (yellow). Peak magnitudes in M1 were

significantly larger than those in VLp and VLa (** $p < 0.01$, Wilcoxon rank test), whereas no significant difference was observed between VLp and VLa. B. Cumulative distributions of the onsets of change in partial R^2 values associated with target direction in VLp (magenta), VLa (green), and M1 (yellow). Onsets in VLa were significantly delayed relative to VLp and M1 (* $p < 0.05$, ** $p < 0.01$, Wilcoxon rank test), whereas no significant difference was found between VLp and M1. C. Scatter plots of the onset of change in partial R^2 values associated with target direction (x-axis) versus the magnitude of change (y-axis) for VLp (left), VLa (middle), and M1 (right). Insets: Bayesian Information Criterion (BIC) values from clustering analyses. X-axis indicates the number of clusters used. D–E. Mean magnitude (D) and mean onset (E) of the significant increase in partial R^2 values associated with target direction (** $p < 0.01$, Wilcoxon rank test)

Table 1

Activity at rest

	NHP G			NHP I			Total		
	VLp	VLa	M1	VLp	VLa	M1	VLp	VLa	M1
Number of Units	35	59	-	48	134	172	83	193	172
Firing rate at rest (Hz)	6.50 ± 5.4	7.6 ± 5.4	-	8.5 ± 5.8	7.1 ± 5.2	6.2 ± 10.1	7.8 ± 5.7	7.2 ± 5.3	6.2 ± 10.1
Coefficient of variation of ISI	1.6 ± 0.3	1.5 ± 0.3	-	1.3 ± 0.4	1.5 ± 0.4 ** 1.2 ± 0.5		1.5 ± 0.3	1.5 ± 0.3 ** 1.2 ± 0.5 **	
Fraction of spike in the burst	0.36 ± 0.13	0.33 ± 0.11	-	0.21 ± 0.11	0.26 ± 0.12 ** 0.11 ± 0.13		0.25 ± 0.13	0.27 ± 0.12 ** 0.11 ± 0.13	
Normalized intra-burst firing rate (Hz)	9.1 ± 25.2	8.1 ± 61.0	-	6.3 ± 15.3	7.5 ± 74.2	8.9 ± 140.0	7.1 ± 19.8	7.7 ± 70.3	8.9 ± 140.0
Half trough width (msec)	0.19 ± 0.11	0.26 ± 0.10	-	0.29 ± 0.08	0.30 ± 0.08 ** 0.16 ± 0.08 **		0.26 ± 0.10	0.30 ± 0.09 ** 0.16 ± 0.08	
Trough-Peak time (msec)	0.42 ± 0.23	0.23 ± 0.34	-	0.60 ± 0.16	0.60 ± 0.23 ** 0.23 ± 0.30 **		0.57 ± 0.20	0.57 ± 0.29 ** 0.23 ± 0.30	
Trough/Peak magnitude ratio	2.5 ± 1.6	2.7 ± 1.6	-	3.6 ± 1.5	3.5 ± 1.3	3.4 ± 1.7	3.4 ± 1.6	3.3 ± 1.4 **	3.4 ± 1.7
Ramping of firing rate	32 (91%)	51 (86%)	-	43 (90%)	120 (90%)	157 (91%)	75 (90%)	171 (89%)	157 (91%)
Positive slope	17 (53%)	25 (49%)	-	20 (47%)	60 (50%)	69 (44%)	37 (49%)	85 (50%)	69 (44%)
Negative slope	15 (47%)	26 (51%)	-	23 (53%)	60 (50%)	88 (56%)	38 (51%)	86 (50%)	88 (56%)
Behavior encoding units	18	36	-	34	67	128	52	103	128
Target direction	13	29	-	26	48	114	39	77	114
Session time	9	26	-	14	41	70	23	67	70
Mean speed	5	9	-	10	17	73	15	26	73
Reaction time	1	9	-	8	17	41	9	26	41
Delay period	4	8	-	7	13	22	11	21	22
Median (maximum) condition index of Belsey's collinearity diagnosis	1.31 (2.54)	1.34 (3.60)	-	1.43 (3.39)	1.44 (4.18)	1.34 (2.49)	1.39 (3.39)	1.41 (4.18)	1.34 (2.49)

* p<0.05, ** p<0.01 rank-sum test

Table 2

Metrics of peri-movement SDF

		VLp		VLa		M1
Increase only	Latency (msec)	112.5 ± 9.3	**	82.4 ± 7.0	*	104.3 ± 6.6
	Magnitude (Hz)	16.7 ± 1.4	**	13.2 ± 0.8	**	34.2 ± 2.2
	Slope (Hz/msec)	0.16 ± 0.01	**	0.15 ± 0.01	**	0.35 ± 0.02
	Half width (msec)	248.2 ± 26.1	**	257.0 ± 22.0	**	199.5 ± 18.1
Decrease only	Latency (msec)	58.6 ± 23.7		71.9 ± 8.6		87.6 ± 13.7
	Magnitude (Hz)	9.1 ± 1.2	*	9.4 ± 0.5	**	12.8 ± 1.0
	Slope (Hz/msec)	0.13 ± 0.02	*	0.11 ± 0.01	**	0.18 ± 0.02
	Half width (msec)	342.8 ± 98.4		472.8 ± 54.9	**	333.5 ± 64.6

* p<0.05 ** p<0.01 Wilcoxon rank test

Supplemental Figure S1

Reconstructed locations of all VLp (magenta) and VLa (green) single units included in the database. The VLp–VLa boundary (black dashed lines) was defined by responses to SCP and GPi stimulation during border-mapping sessions prior to data collection. Locations are projected onto parasagittal sections at 1 mm intervals for animals G and I separately. The boundary of thalamus was taken from a standard atlas and aligned with the mapping results from each animal.

Supplemental Figure S2

A–D. Top and second rows: cumulative distributions of the peak magnitudes (top row) and onsets (second row) of partial R^2 values associated with session time, mean speed, reaction time, and delay period (from left to right). Partial R^2 values associated with mean speed in VLp were significantly smaller than those in M1 (* $p < 0.05$, Wilcoxon rank test). Third, fourth, and bottom rows: scatter plots of the onset of change in partial R^2 values associated with predictors (x-axis, session time, mean speed, reaction time, and delay period, from left to right) versus the magnitude of change (y-axis) for VLp (third row), VLa (fourth row), and M1 (bottom row). Insets: BIC values from clustering analyses. X-axis indicates the number of clusters used.

References

1.   **Alexander GE, Crutcher MD**. Functional architecture of basal ganglia circuits: neural substrates of parallel processing. *Trends in neurosciences* 13: 266-271, 1990.
2.   **Alexander GE, Crutcher MD**. Preparation for movement: neural representations of intended direction in three motor areas of the monkey. *Journal of neurophysiology* 64: 133-150, 1990.
3.   **Anderson ME, Turner RS**. Activity of neurons in cerebellar-receiving and pallidal-receiving areas of the thalamus of the behaving monkey. *Journal of neurophysiology* 66: 879-893, 1991.
4.   **Ando N, Izawa Y, Shinoda Y**. Relative contributions of thalamic reticular nucleus neurons and intrinsic interneurons to inhibition of thalamic neurons projecting to the motor cortex. *Journal of neurophysiology* 73: 2470-2485, 1995.
5.   **Arcelli P, Frassoni C, Regondi MC, De Biasi S, Spreafico R**. GABAergic neurons in mammalian thalamus: a marker of thalamic complexity? *Brain research bulletin* 42: 27-37, 1997.
6.   **Arecchi-Bouchhioua P, Yelnik J, Francois C, Percheron G, Tande D**. 3-D tracing of biocytin-labelled pallido-thalamic axons in the monkey. *Neuroreport* 7: 981-984, 1996.
7.   **Asanuma C, Thach WT, Jones EG**. Cytoarchitectonic delineation of the ventral lateral thalamic region in the monkey. *Brain research* 286: 219-235, 1983.
8.   **Asanuma H, Rosen I**. Topographical organization of cortical efferent zones projecting to distal forelimb muscles in the monkey. *Experimental brain research Experimentelle Hirnforschung* 14: 243-256, 1972.

9.   **Ashby FG, Turner BO, Horvitz JC**. Cortical and basal ganglia contributions to habit learning and automaticity. *Trends in cognitive sciences* 14: 208-215, 2010.

10.   **Belsley DA**. A Guide to using the collinearity diagnostics. *Computer Science in Economics and Management* 4: 33-50, 1991.

11.   **Belsley DA, Kuh, B.E., Welsch, R.E.** *Regression Diagnostics: Identifying Influential Data and Sources of Collinearity*. Wiley Online Library, 1980.

12.   **Bodor AL, Giber K, Rovo Z, Ulbert I, Acsady L**. Structural correlates of efficient GABAergic transmission in the basal ganglia-thalamus pathway. *J Neurosci* 28: 3090-3102, 2008.

13.   **Bosch-Bouju C, Hyland BI, Parr-Brownlie LC**. Motor thalamus integration of cortical, cerebellar and basal ganglia information: implications for normal and parkinsonian conditions. *Frontiers in computational neuroscience* 7: 163, 2013.

14.   **Bosch-Bouju C, Smither RA, Hyland BI, Parr-Brownlie LC**. Reduced reach-related modulation of motor thalamus neural activity in a rat model of Parkinson's disease. *J Neurosci* 34: 15836-15850, 2014.

15.   **Buford JA, Inase M, Anderson ME**. Contrasting locations of pallidal-receiving neurons and microexcitable zones in primate thalamus. *Journal of neurophysiology* 75: 1105-1116, 1996.

16.   **Cox KM, Kase D, Znati T, Turner RS**. Detecting rhythmic spiking through the power spectra of point process model residuals. *Journal of neural engineering* 21: 2024.

17.   **Dacre J, Colligan M, Clarke T, Ammer JJ, Schiemann J, Chamosa-Pino V, Claudi F, Harston JA, Eleftheriou C, Pakan JMP, Huang CC, Hantman AW, Rochefort NL, Duguid I**. A cerebellar-thalamocortical pathway drives behavioral context-dependent movement initiation. *Neuron* 109: 2326-2338 e2328, 2021.

18.   **DeLong MR**. Activity of basal ganglia neurons during movement. *Brain Res* 40: 127-135, 1972.

19.   **Desmurget M, Turner RS**. Testing Basal Ganglia motor functions through reversible inactivations in the posterior internal globus pallidus. *Journal of neurophysiology* 99: 1057-1076, 2008.

20.   **DeVito JL, Anderson ME**. An autoradiographic study of efferent connections of the globus pallidus in Macaca mulatta. *Experimental brain research Experimentelle Hirnforschung* 46: 107-117, 1982.

21.   **Evarts E**. Motor cortex reflexes associated with learned movement. *Science (New York, NY* 179: 501-503, 1973.

22.   **Evarts E, Thach W**. Motor mechanism of the CNS: Cerebrocerebellar interrelations. *Ann Rev Physiol* 31: 451-498, 1969.

23.   **Floyer-Lea A, Matthews PM**. Distinguishable brain activation networks for short- and long-term motor skill learning. *Journal of neurophysiology* 94: 512-518, 2005.

24.   **Fortier PA, Kalaska JF, Smith AM**. Cerebellar neuronal activity related to whole-arm reaching movements in the monkey. *JNeurophysiol* 62(1): 198-211, 1989.

25.   **Franco V, Turner RS**. Testing the contributions of striatal dopamine loss to the genesis of parkinsonian signs. *Neurobiol Dis* 47: 114-125, 2012.

26.   **Fu Q-G, Flament D, Coltz J, Ebner T**. Relationship of cerebellar purkinje cell simple spike discharge to movement kinematics in the monkey. *Journal of neurophysiology* 78: 478-491, 1997.

27.   **Gaidica M, Hurst A, Cyr C, Leventhal DK**. Distinct Populations of Motor Thalamic Neurons Encode Action Initiation, Action Selection, and Movement Vigor. *J Neurosci* 38: 6563-6573, 2018.

28.   **Galvan A, Hu X, Smith Y, Wichmann T**. Effects of Optogenetic Activation of Corticothalamic Terminals in the Motor Thalamus of Awake Monkeys. *J Neurosci* 36: 3519-3530, 2016.

29.   **Goldberg JH, Fee MS**. A cortical motor nucleus drives the basal ganglia-recipient thalamus in singing birds. *Nature neuroscience* 15: 620-627, 2012.

30.   **Graybiel AM**. Habits, rituals, and the evaluative brain. *Annual review of neuroscience* 31: 359-387, 2008.

31.   **Guo ZV, Inagaki HK, Daie K, Druckmann S, Gerfen CR, Svoboda K**. Maintenance of persistent activity in a frontal thalamocortical loop. *Nature* 545: 181-186, 2017.

32.   **Hoover JE, Strick PL**. The organization of cerebellar and basal ganglia outputs to primary motor cortex as revealed by retrograde transneuronal transport of herpes simplex virus type 1. *J Neurosci* 19: 1446-1463, 1999.

33.   **Horne MK, Butler EG**. The role of the cerebello-thalamo-cortical pathway in skilled movement. *Prog Neurobiol* 46: 199-213, 1995.

34.   **Ilinsky I, Kultas-Ilinsky K**. Sagittal cytoarchitectonic maps of the macaca mulatta thalamus with a revised nomenclature of the motor-related nuclei validated by observations on their connectivity. *J Comp Neurol* 262: 331-364, 1987.

35.   **Ilinsky IA, Ambardekar AV, Kultas-Ilinsky K**. Organization of projections from the anterior pole of the nucleus reticularis thalami (NRT) to subdivisions of the motor thalamus: light and electron microscopic studies in the rhesus monkey. *J Comp Neurol* 409: 369-384, 1999.

36.   **Ilinsky IA, Kultas-Ilinsky K**. Fine structure of the magnocellular subdivision of the ventral anterior thalamic nucleus (VAmc) of Macaca mulatta: I. Cell types and synaptology. *J Comp Neurol* 294: 455-478, 1990.

37.   **Ilinsky IA, Yi H, Kultas-Ilinsky K**. Mode of termination of pallidal afferents to the thalamus: a light and electron microscopic study with anterograde tracers and immunocytochemistry in Macaca mulatta. *J Comp Neurol* 386: 601-612, 1997.

38.   **Inagaki HK, Chen S, Daie K, Finkelstein A, Fontolan L, Romani S, Svoboda K**. Neural Algorithms and Circuits for Motor Planning. *Annual review of neuroscience* 45: 249-271, 2022.

39.   **Ivanusic JJ, Bourke DW, Xu ZM, Butler EG, Horne MK**. Cerebellar thalamic activity in the macaque monkey encodes the duration but not the force or velocity of wrist movement. *Brain research* 1041: 181-197, 2005.

40.   **Jones EG**. *The Thalamus*. New York: Cambridge University Press, 2007.

41.   **Kase D, Uta D, Ishihara H, Imoto K**. Inhibitory synaptic transmission from the substantia nigra pars reticulata to the ventral medial thalamus in mice. *Neurosci Res* 97: 26-35, 2015.

42.   **Kim J, Kim Y, Nakajima R, Shin A, Jeong M, Park AH, Jeong Y, Jo S, Yang S, Park H, Cho SH, Cho KH, Shim I, Chung JH, Paik SB, Augustine GJ, Kim D**. Inhibitory Basal Ganglia Inputs Induce Excitatory Motor Signals in the Thalamus. *Neuron* 95: 1181-1196 e1188, 2017.

43.   **Kitazawa S, Kimura T, Yin PB**. Cerebellar complex spikes encode both destinations and errors in arm movements. *Nature* 392: 494-497, 1998.

44.	**Koster KP, Sherman SM**. Convergence of inputs from the basal ganglia with layer 5 of motor cortex and cerebellum in mouse motor thalamus. *Elife* 13: 2024.

45.	**Kultas-Ilinsky K, Sivan-Loukianova E, Ilinsky IA**. Reevaluation of the primary motor cortex connections with the thalamus in primates. *J Comp Neurol* 457: 133-158, 2003.

46.	**Kuramoto E, Fujiyama F, Nakamura KC, Tanaka Y, Hioki H, Kaneko T**. Complementary distribution of glutamatergic cerebellar and GABAergic basal ganglia afferents to the rat motor thalamic nuclei. *The European journal of neuroscience* 33: 95-109, 2011.

47.	**Legendy CR, Salcman M**. Bursts and recurrences of bursts in the spike trains of spontaneously active striate cortex neurons. *Journal of neurophysiology* 53: 926-939, 1985.

48.	**Mason A, Ilinsky IA, Maldonado S, Kultas-Ilinsky K**. Thalamic terminal fields of individual axons from the ventral part of the dentate nucleus of the cerebellum in Macaca mulatta. *J Comp Neurol* 421: 412-428, 2000.

49.	**Matsuzaka Y, Picard N, Strick PL**. Skill representation in the primary motor cortex after long-term practice. *Journal of neurophysiology* 97: 1819-1832, 2007.

50.	**Mazzoni P, Hristova A, Krakauer JW**. Why don't we move faster? Parkinson's disease, movement vigor, and implicit motivation. *J Neurosci* 27: 7105-7116, 2007.

51.	**Mink J**. The basal ganglia: focused selection and inhibition of competing motor programs. *Prog Neurobiol* 50: 381-425, 1996.

52.	**Mink J, Thach W**. Basal ganglia motor control. III. pallidal ablation: normal reaction time, muscle cocontraction, and slow movement. *Journal of neurophysiology* 65: 330-351, 1991.

53.	**Miocinovic S, Zhang J, Xu W, Russo GS, Vitek JL, McIntyre CC**. Stereotactic neurosurgical planning, recording, and visualization for deep brain stimulation in non-human primates. *J Neurosci Methods* 162: 32-41, 2007.

54.	**Nambu A, Yoshida S, Jinnai K**. Movement-related activity of thalamic neurons with input from the globus pallidus and projection to the motor cortex in the monkey. *Experimental brain research Experimentelle Hirnforschung* 84: 279-284, 1991.

55.	**Nambu A, Yoshida S, Jinnai K**. Projection on the motor cortex of thalamic neurons with pallidal input in the monkey. *Experimental brain research Experimentelle Hirnforschung* 71: 658-662, 1988.

56.	**Nashef A, Cohen O, Harel R, Israel Z, Prut Y**. Reversible Block of Cerebellar Outflow Reveals Cortical Circuitry for Motor Coordination. *Cell Rep* 27: 2608-2619 e2604, 2019.

57.	**Nauta HJW, Mehler WR**. Projections of the lentiform nucleus in the monkey. *Brain Res* 1: 3-42, 1966.

58.	**Penney JB, Young AB**. GABA as the pallidothalamic neurotransmitter: implications for basal ganglia function. *Brain Res* 207: 195-199, 1981.

59.	**Pessiglione M, Guehl D, Rolland AS, Francois C, Hirsch EC, Feger J, Tremblay L**. Thalamic neuronal activity in dopamine-depleted primates: evidence for a loss of functional segregation within basal ganglia circuits. *J Neurosci* 25: 1523-1531, 2005.

60.	**Phillips JW, Schulmann A, Hara E, Winnubst J, Liu C, Valakh V, Wang L, Shields BC, Korff W, Chandrashekar J, Lemire AL, Mensh B, Dudman JT, Nelson SB, Hantman AW**. A repeated molecular architecture across thalamic pathways. *Nature neuroscience* 22: 1925-1935, 2019.

61.	**Redgrave P, Prescott TJ, Gurney K**. The basal ganglia: a vertebrate solution to the selection problem? *Neuroscience* 89: 1009-1023, 1999.

62.   **Roth RH, Muniak MA, Huang CJ, Hwang FJ, Sun Y, Min C, Mao T, Ding JB**. Thalamic integration of basal ganglia and cerebellar circuits during motor learning. *bioRxiv* 2024.

63.   **Rouiller EM, Tanne J, Moret V, Kermadi I, Boussaoud D, Welker E**. Dual morphology and topography of the corticothalamic terminals originating from the primary, supplementary motor, and dorsal premotor cortical areas in macaque monkeys. *J Comp Neurol* 396: 169-185, 1998.

64.   **Rovo Z, Ulbert I, Acsady L**. Drivers of the primate thalamus. *J Neurosci* 32: 17894-17908, 2012.

65.   **Sakai ST, Inase M, Tanji J**. Comparison of cerebellothalamic and pallidothalamic projections in the monkey (Macaca fuscata): a double anterograde labeling study. *J Comp Neurol* 368: 215-228, 1996.

66.   **Sauerbrei BA, Guo JZ, Cohen JD, Mischiati M, Guo W, Kabra M, Verma N, Mensh B, Branson K, Hantman AW**. Cortical pattern generation during dexterous movement is input-driven. *Nature* 577: 386-391, 2020.

67.   **Schieber MH, Rivlis G**. Partial reconstruction of muscle activity from a pruned network of diverse motor cortex neurons. *Journal of neurophysiology* 97: 70-82, 2007.

68.   **Schwab BC, Kase D, Zimnik A, Rosenbaum R, Codianni MG, Rubin JE, Turner RS**. Neural activity during a simple reaching task in macaques is counter to gating and rebound in basal ganglia-thalamic communication. *PLoS biology* 18: e3000829, 2020.

69.   **Sedaghat-Nejad E, Herzfeld DJ, Hage P, Karbasi K, Palin T, Wang X, Shadmehr R**. Behavioral training of marmosets and electrophysiological recording from the cerebellum. *Journal of neurophysiology* 122: 1502-1517, 2019.

70.   **Sherman SM, Guillery RW**. Functional organization of thalamocortical relays. *Journal of neurophysiology* 76: 1367-1395, 1996.

71.   **Sherman SM, Guillery RW**. The role of the thalamus in the flow of information to the cortex. *Philosophical transactions of the Royal Society of London* 357: 1695-1708, 2002.

72.   **Steriade M, Llinas RR**. The functional states of the thalamus and the associated neuronal interplay. *Physiol Rev* 68: 649-742, 1988.

73.   **Thach WT**. Discharge of cerebellar neurons related to two maintained postures and two prompt movements. I. Nuclear cell output. *Journal of neurophysiology* 33: 527-536, 1970.

74.   **Turner R, DeLong M**. Corticostriatal activity in primary motor cortex of the macaque. *J Neurosci* 20: 7096-7108., 2000.

75.   **Turner RS, Anderson ME**. Context-dependent modulation of movement-related discharge in the primate globus pallidus. *J Neurosci* 25: 2965-2976, 2005.

76.   **Turner RS, Anderson ME**. Pallidal discharge related to the kinematics of reaching movements in two dimensions. *Journal of neurophysiology* 77: 1051-1074, 1997.

77.   **Turner RS, Desmurget M**. Basal ganglia contributions to motor control: a vigorous tutor. *Current opinion in neurobiology* 20: 704-716, 2010.

78.   **Uno M, Ozawa N, Yoshida M**. The mole of pallido-thalamic transmission investigated with intracellular recording from cat thalamus. *Experimental brain research Experimentelle Hirnforschung* 33: 493-507, 1978.

79.   **Uno M, Yoshida M, Hirota I**. The mode of cerebello-thalamic relay transmission investigated with intracellular recording from cells of the ventrolateral nucleus of cat's thalamus. *Experimental brain research Experimentelle Hirnforschung* 10: 121-139, 1970.

80.   **van Donkelaar P, Stein JF, Passingham RE, Miall RC**. Neuronal activity in the primate motor thalamus during visually triggered and internally generated limb movements. *Journal of neurophysiology* 82: 934-945, 1999.

81.   **Vitek JL, Ashe J, DeLong MR, Alexander GE**. Physiologic properties and somatotopic organization of the primate motor thalamus. *Journal of neurophysiology* 71: 1498-1513, 1994.

82.   **Yamamoto T, Hassler R, Huber C, Wagner A, Sasaki K**. Electrophysiologic studies on the pallido- and cerebellothalamic projections in squirrel monkeys (Saimiri sciureus). *Experimental brain research Experimentelle Hirnforschung* 51: 77-87, 1983.

83.   **Yoshida S, Nambu A, Jinnai K**. The distribution of the globus pallidus neurons with input from various cortical areas in the monkeys. *Brain Res* 611: 170-174, 1993.

84.   **Yttri EA, Dudman JT**. Opponent and bidirectional control of movement velocity in the basal ganglia. *Nature* 533: 402-406, 2016.

85.   **Zimnik AJ, Nora GJ, Desmurget M, Turner RS**. Movement-related discharge in the macaque globus pallidus during high-frequency stimulation of the subthalamic nucleus. *J Neurosci* 35: 3978-3989, 2015.