Exploring the multifactorial causes and therapeutic strategies for anabolic

resistance in sarcopenia: A systems modeling study

Taylor J. McColl[1,2]* Daniel R. Moore[3], Eldon Emberly[4], David D. Church[5], and David C.

Clarke[1,2,]*^

[1]Department of Biomedical Physiology and Kinesiology and [2]Centre for Cell Biology,

Development and Disease, Simon Fraser University, Burnaby BC, V5A 1S6, Canada; [3]Faculty of

Kinesiology and Physical Education, University of Toronto, Toronto, ON, M5S 2C9, Canada;

[4]Department of Physics, Simon Fraser University, Burnaby BC, V5A 1S6, Canada; [5]Department

of Geriatrics, Donald W. Reynolds Institute of Aging, Center for Translational Research in Aging

and Longevity, University of Arkansas for Medical Sciences, Little Rock, AR, United States.

*^ Co-corresponding author and lead contact:

Email: dcclarke@sfu.ca

Tel: 778-782-9777

Fax: 778-782-3040

* Co-corresponding author:

Email: tmccoll@sfu.ca

Running title: multifactorial cause of anabolic resistance

Word count:

Abstract

Background: Sarcopenia is the progressive loss of skeletal muscle mass, strength, and function with age, driven by dysregulation in the rates of muscle protein synthesis (MPS) and breakdown (MPB). Although MPB contributes to net protein balance (NB), a primary contributor of sarcopenia is anabolic resistance, defined as the blunted MPS response to anabolic stimuli such as feeding. While candidate mechanisms of anabolic resistance have been identified, none singularly explains the reduction in MPS. Instead, multiple mechanisms likely act simultaneously and interactively to suppress MPS. Studying these interactions experimentally is challenging. Alternatively, mathematical modeling is well suited to analyzing complex biological phenomena such as anabolic resistance.

Methods: We analyzed a previously developed kinetic model of leucine-mediated signaling and protein metabolism in human skeletal muscle to systematically investigate potential mechanisms contributing to anabolic resistance. A global sensitivity analysis identified key controllers of MPS, MPB, and NB. We then simulated an essential amino acid feeding intervention in older adults, classified the responses as either anabolic sensitive or resistant, and compared the resulting parameter distributions of the two groups. A targeted analysis evaluated the effects of individual and combined dysregulated putative mechanisms of anabolic resistance on muscle metabolism. Finally, we simulated therapeutic interventions aimed at restoring muscle metabolism.

Results: Sensitivity analysis revealed that MPS and MPB are primarily controlled by their proximal signaling controllers, while NB is largely driven by MPS dynamics. Exploratory simulations showed that several parameters and signaling protein concentrations, particularly those controlling MPS, differed significantly between anabolic sensitive and resistant groups. Targeted simulations indicated that multiple dysregulated mechanisms were required to account for the experimentally observed reductions in MPS in older adults. Therapeutic simulations showed that while single-target interventions could largely restore MPS when isolated mechanisms were perturbed (e.g., enhancing p70S6K levels, increasing mTORC1 sensitivity), a multifactorial approach was required to recover muscle metabolism when all anabolic resistance mechanisms were present.

Conclusion: This study demonstrates the multifactorial nature of anabolic resistance. We provide a framework for the experimental manipulation of independent and related factors that could underpin age-related anabolic resistance. The results motivate new hypotheses regarding the mechanisms most likely to be impaired and support the development of multi-target therapeutic strategies to help restore muscle protein metabolism in aging.

57   **Keywords**

58   Skeletal muscle, protein synthesis, mTOR, signal transduction, mathematical model,

59   computational biology

Introduction

Sarcopenia is the progressive decline in skeletal muscle mass, strength, and function that

accompanies aging [1]. Those with sarcopenia experience higher hospitalization rates [2] and

contribute to increased public healthcare costs [3–5]. Given the impact of sarcopenia on health and

quality of life, further research is needed to enhance our understanding of age-related muscle loss

and to develop effective strategies for its prevention and management.

Skeletal muscle mass is largely determined by its protein content, which undergoes

continuous turnover through concurrent synthesis and degradation processes [6]. The balance

between the rates of muscle protein synthesis (MPS) and muscle protein breakdown (MPB)

governs overall muscle protein levels, with the difference between MPS and MPB termed *net*

protein balance (NB). MPS increases in response to anabolic stimuli such as nutrient intake [7],

with amino acid availability – particularly leucine [8–10] – playing a key stimulatory role. In

contrast, the attenuation of MPB during the postprandial period is primarily mediated via an

increase in plasma insulin levels [11]. In the postabsorptive state, MPS and MPB rates are

generally similar between young and older adults [12–15]. However, aging muscle exhibits

anabolic resistance, characterized by a blunted stimulation of MPS and impaired suppression of

MPB following anabolic stimuli [16–18]. Over time, this anabolic resistance drives the gradual

loss of skeletal muscle.

Anabolic resistance of aging has been suggested to be underpinned by a range of factors

including reduced amino acid availability to muscle due to increased splanchnic extraction [19–

21], impaired transport from blood into muscle [20,22], impaired insulin-mediated protein

signaling [23,24], reduced mTOR and p70S6K protein levels [18,25], and diminished mTORC1

signaling sensitivity to protein intake [18,26]. Sustained MPB rates are due to reduced insulin-

mediated blunting of proteolysis or other age-related factors, such as chronic inflammation [27–

29]. Effectively understanding and treating anabolic resistance therefore requires quantifying the

relative contributions of these mechanisms and identifying targets for intervention.

The limited long-term success of treatments for anabolic resistance may stem from the

simultaneous dysregulation of multiple mechanisms affecting muscle protein metabolism.

Nutritional and pharmacological interventions tend to target individual pathways rather than

addressing the broader complexity of these metabolic disruptions. Therefore, strategies that

consider multiple interacting factors influencing muscle protein metabolism may be needed to
develop more effective treatment strategies.

Mathematical models are valuable tools for studying complex biological systems. They
could be applied to identify the most likely combination of dysregulated mechanisms underlying
anabolic resistance in older adults, without relying on logistically challenging and costly human
clinical research needed to fully characterize the myriad potential biological mediators. We
recently developed a kinetic model of leucine-mediated skeletal muscle signaling and protein
metabolism, incorporating key leucine- and insulin-mediated pathways that control muscle
metabolism [10]. Although originally developed using data from healthy younger adults, these
molecular pathways are also present in older individuals but likely operate with altered reaction
rates. Therefore, this kinetic model provides a framework to investigate the multifactorial nature
of anabolic resistance and identify key physiological mechanisms driving impaired feeding-
induced muscle metabolism.

The purpose of this study was to investigate the multifactorial causes of anabolic resistance
by quantifying how dysregulated mechanisms in muscle protein metabolism contribute to impaired
muscle mass in older adults. Specifically, we applied sensitivity analyses to the McColl and Clarke
kinetic model [10] to identify key parameters controlling muscle metabolism. We pursued two
strategies: 1) a "naïve" global sensitivity analysis to identify parameters potentially responsible for
reduced MPS, and 2) a targeted sensitivity analysis that simulated previously proposed anabolic
resistance mechanisms. These targeted simulations were then used to evaluate therapeutic
strategies aimed at restoring MPS and NB. Collectively, our findings provide insight into the
principal physiological drivers of anabolic resistance and identify promising targets for therapeutic
intervention to mitigate muscle loss in older adults.

Methods

Kinetic model of leucine-mediated signaling and muscle protein metabolism

The McColl and Clarke kinetic model of leucine-mediated signaling and muscle protein
metabolism in human skeletal muscle [10] (henceforth, the kinetic model) served as the foundation
for this study (Figure 1). This model simulates skeletal muscle protein metabolism in response to
leucine ingestion in healthy adults and contains four interconnected modules: 1) the mTOR
signaling module, 2) the leucine kinetic module, 3) the digestive system module, and 4) the insulin

secretion module [10]. Together, these modules simulate the ingestion of leucine, leucine-

mediated stimulation of insulin-and leucine-dependent signaling pathways, and muscle protein

metabolism. While originally developed using data from healthy adults, the model can be applied

to study anabolic resistance in sarcopenia because the biological mechanisms it represents are

conserved in young and older individuals. Differences between young and older adults manifest

as differences in parameter values. Further details of the kinetic model are provided in the

Supplementary Information.

Defining anabolic resistance

We established an anabolic resistance "threshold" based on two studies by Mitchell et al.

[30,31]. Although these studies measured myofibrillar protein synthesis specifically, this protein

fraction, which constitutes ~60% of muscle proteins, demonstrates a similar nutrient sensitivity

and age-related resistance as sarcoplasmic proteins [18,32]. Both studies used the same

intervention: young [30] and older adults [31] received a 15-gram mixed essential amino acid

(EAA) bolus (3.59 grams of leucine), with muscle biopsies collected over the subsequent four

hours to calculate MPS, measured as fractional synthetic rate (FSR) [30,31]. Using the "total

leucine synthesized" calculations from McColl and Clarke [10], the experimentally measured MPS

response in young adults was 0.461 ± 0.111 grams of leucine synthesized in muscle over four hours,

while in older adults it was 0.267 ± 0.092 grams of leucine. Despite the older adults being classified

as healthy [31], the reduced MPS response to EAA feeding indicated the presence of anabolic

resistance. To define the anabolic resistance threshold, we used the lower 95% confidence interval

of the younger adult MPS response (0.461 ± 0.111 grams of leucine), yielding a threshold of 0.251

g leucine synthesized in muscle. Simulated MPS responses below this threshold were classified as

anabolically resistant, whereas those above it were considered *anabolically sensitive*. This value

closely approximates the mean MPS response for the older adult group.

Multi-parametric sensitivity analysis

We applied a multi-parametric sensitivity analysis (MPSA) to generate physiologically

plausible parameter sets, with the purpose of identifying model parameters that most influence

system-level behaviors representative of anabolic resistance. MPSA is a global sensitivity analysis

approach that systematically samples parameter sets and evaluates their relative contribution to

model dynamics [33]. Complete details for the MPSA, including the criteria for parameter filtering

and analyses applied to the filtered parameter sets, are provided in the Supplementary Information.

MPSA: Kolmogorov-Smirnov test

We applied the two-sample Kolmogorov-Smirnov (KS) test to assess whether parameter

sets from the anabolic-sensitive and anabolic-resistant groups originated from the same

distribution or represented unique distributions. We performed the test using the R function *ks.test*,

which calculated the KS statistic and the associated p-value. To account for multiple comparisons,

we applied false discovery rate (FDR) correction using the Benjamini-Hochberg method (R

function: *p.adjust*) to compute the *q*-value. A *q*-value less than 0.05 indicated a statistically

significant difference between parameter distributions.

Targeted approach to evaluate the contributions of anabolic resistance mechanisms

to muscle protein metabolism

We used a targeted approach to evaluate how the putative mechanisms underlying anabolic

resistance in sarcopenia independently influence muscle protein metabolism and how these

mechanisms might operate synergistically to further impair muscle protein metabolism. As noted

in the Introduction and outlined in Table 1, several mechanisms contribute to anabolic resistance

(Figure 1). Of note, elevated postabsorptive levels of phospho-p70S6K have been observed in

older adults, which we hypothesized represents a compensatory response to dysregulated

mechanisms [10,34] – a phenomenon we term "compensatory signaling". For each mechanism,

we curated experimental data quantifying the differences between young and older adults,

identified the model parameter(s) corresponding to each mechanism, and estimated both a

consensus fold change and a *worst-case fold change* in the values of these parameters between

young and old based on a subset of relevant studies (Table 1). While several of the studies

informing each mechanism did not directly assess the modeled process, they reported

measurements of critical pathway components that we used to guide the changes representing

anabolic resistance.

Table 1. Modeling the putative anabolic resistance mechanisms. Summary of putative mechanisms contributing to anabolic resistance in sarcopenia, along with quantitative evidence for the consensus (i.e., average) and worst-case effects identified across a subset of studies. The worst-case effect was selected from the study reporting the greatest deviation in fold-change relative to the young group. The number of studies informing each consensus estimate is indicated by n. The kinetic model parameters that were altered to represent the corresponding mechanism are listed in the rightmost column.

Putative anabolic resistance mechanism	Consensus change	Worst-case change	Integration into model
First-pass splanchnic extraction	1.76±0.25 F.C. extraction in elderly (n=3) [19–21]	2.20±0.52 F.C. extraction in elderly [19]	Increased k68 (first-pass splanchnic extraction)
Reduced delivery of amino acids to muscle	0.73±0.30 F.C. $F_{m,a}$ (n=2) [20,22]	0.69±0.39 F.C. $F_{m,a}$ [22]	Reduced k6 (leucine transport into muscle)
Impaired insulin signalling	0.80±0.24 F.C. Akt activity in elderly (n=3) [17,27,35]	0.66±0.41 F.C. Akt activity in elderly [27]	Reduced k16 (insulin binding mediated insulin signalling)
mTOR protein level	0.79±0.15 F.C. mTOR levels in elderly (n=2) [18,25]	0.50±0.18 F.C. mTOR levels in elderly [18,25]	Reduced both mTORC1 and mTORC2 initial concentrations
Reduced mTORC1 sensitivity[#]	0.57±0.16 F.C. phospho-p70S6K in elderly (n=3) [17,18,35]	0.43±0.36 F.C. phospho-p70S6K in elderly [18]	Reduced k40 (mTORC1-mediated p70S6K phosphorylation)
p70S6K protein level[#]	0.78±0.09 F.C. p70S6K levels in elderly (n=2) [18,25]	0.50±0.18 F.C. p70S6K levels in elderly [18,25]	Reduced the p70S6K initial concentration
Elevated postabsorptive phospho-p70S6K (compensatory signaling)[#]	1.87±0.55 F.C. phospho-p70S6K per total p70S6K in elderly (n=2) [17,18]	1.70±0.60 F.C. phospho-p70S6K per total p70S6K in elderly [17]	Added a constitutive p70S6K phosphorylation reaction to the ODEs; p70S6K concentrations were unchanged

[#]The combined simulation of reduced mTORC1 sensitivity, reduced p70S6K protein levels, and elevated postabsorptive phospho-p70S6K represents the ***compensatory protein signalling mechanism*** in older adults.

F.C., fold change; $F_{m,a}$, inward amino acid transport from artery to muscle.

The consensus fold change for each mechanism was calculated in two steps. First, the fold change within each study was calculated as the ratio between the mean response of the young and older adults, with the corresponding standard error (SE) derived using propagation of error. Second, a consensus mean fold change for each mechanism was calculated across studies using an inverse variance-weighted average, which assigns greater weight to estimates with higher precision (Equations 7-9) [36]:

$$w_i = \frac{1}{SE_i^2} \tag{7}$$

$$FC_{consensus} = \frac{\sum_{i=1}^{k} w_i \cdot FC_i}{\sum_{i=1}^{k} w_i} \tag{8}$$

$$SE_{consensus} = \sqrt{\frac{1}{\sum_{i=1}^{k} w_i}} \tag{9}$$

where i is each of the k independent studies, FC_i and SE_i are the fold change (FC) and
corresponding SE for study i, respectively, w_i is the inverse variance weight of study i, and
$FC_{consensus}$ and $SE_{consensus}$ are the consensus fold change and SE across k studies for the
respective putative mechanism of anabolic resistance. The worst-case fold change was assigned as
the value from the study within the group of studies that reported the greatest deviation from the
young adult population.

We then used the kinetic model to simulate the effects of each putative mechanism
individually, as well as the combined effect of all mechanisms together, in both the presence and
absence of elevated postabsorptive phospho-p70S6K (i.e., compensatory signaling), as follows.
First, we created a virtual population of 100 anabolic sensitive (i.e., "healthy") models by
randomly sampling 100 parameter sets from the anabolic sensitive group of the MPSA (Naïve
approach). Performing the simulations for a virtual population was done to foster the
generalizability of the results. Second, the anabolic resistance mechanisms were simulated within
each of the 100 models by multiplying the relevant model kinetic parameter or initial protein
concentration by the fold change corresponding consensus or worst-case estimate. For the
simulated compensatory phospho-p70S6K signaling mechanism, an additional reaction was added
to the model to simulate the elevated postabsorptive phosphorylation rate, with its kinetic
parameter informed by the estimates reported in Table 1. We then simulated the anabolic resistant
models in response to a 3.59 g leucine bolus and extracted the MPS, MPB, and NB responses over
the 4-hour postprandial period. This dose was chosen to allow comparison with the Mitchell et al.
studies, which provided a 15 g EAA bolus containing 3.59 g leucine and informed the anabolic
resistance thresholds.

Simulation of therapeutic strategies to mitigate anabolic resistance

Therapeutic strategies to mitigate anabolic resistance were simulated using models

developed through the Targeted Approach, which had quantified the effects of individual or

combined anabolic resistance mechanisms on MPS, MPB, and NB. Each therapeutic strategy

involved restoring one or more dysregulated parameters to their "healthy" value by multiplying

the parameters by the inverse of its consensus or worst-case fold-change estimate. Each therapeutic

intervention was applied across all anabolic resistance simulation from the Targeted Analysis –

including those in which the targeted parameter was not originally perturbed. This approach

attempts to mimic the scenario in which one or more druggable targets are treated when the causal

mechanism itself is not druggable (cf. [37,38]). The efficacy of each therapeutic intervention was

assessed against its ability to recover MPS and NB relative to the healthy response.

Numerical methods and software

The 95% confidence interval was calculated using one-tail and the t-value corresponding

to the appropriate degrees of freedom. MATLAB version R2022a was used to perform the MPSA,

model simulations, and multiple regression analyses. R (version 4.3.2) was used for statistical

testing and the visualization of results using the ggplot2 package [39]. WebPlotDigitizer was used

to extract published data (i.e., means, SE) that were only reported in figures [40]. The code

generated   in   this   study   is   freely   available   in   the   GitHub   repository:

https://github.com/tjmccoll/MultifactorialCauseOfAnabolicResistance/.

Results

Multi-parametric sensitivity analysis of the kinetic model

The MPSA resulted in 2,663 physiologically plausible parameter sets for healthy adults.

Regression models using $\log_{10}$-transformed outcome measures and $\log_{10}$-transformed parameter

values were the most predictive for MPS and MPB, whereas using raw outcomes with $\log_{10}$-

transformed parameter values were most predictive for NB. The best-performing regression

models for MPS, MPB, and NB include the top 35, 45, and 35 parameters from the main-effect-

only models as interaction terms, respectively (Table S2). The p-1 sensitivity analysis of MPS and

MPB revealed that parameters most proximal to their respective reactions had the greatest

influence on their outcomes (Table S3). For NB, the most influential parameters were a

combination of those affecting MPS and MPB, but parameters more proximal to MPS had a greater

influence on the NB response (Table S3).

Naïve approach reveals mechanisms contributing to anabolic resistance

We used an unbiased, exploratory approach to assess mechanisms that may contribute to

anabolic resistance in sarcopenia and influence muscle protein metabolism. To do this, we repeated

the MPSA but with the following modifications. First, we excluded the MPS criterion because the

original MPSA was constrained to generate physiologically reasonable MPS responses in healthy,

young adults. This criterion likely restricted the range of MPS responses observed in older adults

with anabolic resistance (see "Defining Anabolic Resistance" in Methods), thereby removing

relevant parameter sets. To better simulate theoretical MPS responses in older adults, we removed

the constraint to allow the full breadth of MPS variability to be captured. Second, the original

MPSA featured fixed values for the parameters controlling the model's input function. However,

one of these parameters controls the rate of splanchnic extraction (k_{68}), which literature suggests

is elevated in older adults [19–21]. To comprehensively assess mechanisms contributing to

anabolic resistance, we allowed k_{68} to vary as part of the MPSA. We first performed an MPSA in

which only k_{68} was varied within a two-fold range to determine its physiologically plausible range

(MPS criterion removed). This range was then used to constrain k_{68} in the subsequent MPSA to

ensure sufficient passing parameter sets. Simulating 10,000 values for k_{68} we found its plausible

range to be $0.0025 - 0.0087$ min^{-1}, corresponding to a 2-fold decrease to a 1.75-fold increase from

its calibrated value.

This MPSA generated 4,054 physiologically plausible parameter sets, including parameter

sets producing MPS responses within the healthy range (Table S1), above it, and below it. We

simulated the kinetic model using these parameter sets following a 3.59-gram leucine bolus and

calculated MPS over the 4-hour postprandial period. The parameter sets were then classified as

"anabolic sensitive" (n = 3,328) or "anabolic resistant" (n = 726) based on the anabolic resistance

threshold (see "Defining Anabolic Resistance" in Methods). The Kolmogorov-Smirnov tests

revealed significant differences in the distributions of ten parameters (Figure 2, Figures S2-4,

Table S4). These ten parameters were largely consistent with those identified in the p-1 sensitivity

analysis of the MPS regression model (see Supplementary Results and Table S2). Two exceptions

were splanchnic extraction, which was not varied in the regression analysis, and hepatic glucose

production, which only marginally met the $q<0.05$ threshold for inclusion. Of the ten parameters,

eight were proximal to the MPS reaction, while the remaining two reflected systemic processes:

first-pass splanchnic extraction and hepatic glucose production (from the insulin secretion

module).

Targeted simulations reveal a multifactorial basis of anabolic resistance

We then simulated the model by integrating the putative mechanisms underlying anabolic

resistance – individually and in combination, with or without compensatory signaling – using both

consensus and worst-case estimates to evaluate their contributions to blunted MPS responses

(Table 1). Simulations using the consensus estimates revealed that increased first-pass splanchnic

extraction and reduced mTORC1 sensitivity (as indicated by p70S6K phosphorylation status) each

moderately decreased MPS (Figure 3A). In contrast, reducing mTOR and p70S6K concentrations

modestly decreased MPS, while reduced blood flow and impaired insulin signaling had minimal

effects. When all mechanisms were simulated together, the model predicted a substantial decline

in MPS, closely approximating the MPS response in older adults as calculated from Mitchell et al.

(2017) [31] (0.267±0.092 grams of leucine synthesized). However, when compensatory signaling

was included – simulating the elevated postabsorptive phospho-p70S6K levels observed in older

adults – the loss of MPS was largely recovered, though still below the healthy level. Changes in

NB closely mirrored those in MPS, as MPB responses remained relatively stable compared to the

healthy simulations (Figures 3B and S5).

Simulations using the worst-case estimates of the anabolic resistance mechanisms revealed

that increased first-pass splanchnic extraction, reduced mTORC1 sensitivity, and reduced mTOR

and p70S6K levels each meaningfully reduced MPS, with those of the latter three mechanisms

approaching the MPS response calculated from Mitchell et al. (2017) [31] (Figure 3C). Reduced

blood flow and impaired insulin-mediated signaling had little effect on MPS. When all

mechanisms were simulated together under worst-case conditions, MPS was substantially reduced

– falling below the MPS response calculated from Mitchell et al. (2017) [31]. Although

compensatory signaling restored much of this loss, the MPS response remained lower than that

observed in the older adult data. Similar to the consensus simulations, MPB responses were largely

unaffected, such that changes in NB closely reflected the changes in MPS (Figures 3D and S5).

Next, we simulated the effect of elevated MPB rates on muscle protein metabolism.
Although accurately estimating MPB in older adults is technically challenging and statistically
significant age-related increases in MPB have not been consistently observed in healthy aging
[12], factors such as chronic low-grade inflammation, which contributes to increased MPB [29],
and reduced insulin-mediated suppression of MPB [27,28] have been observed in older adults. To
evaluate the potential effects of increased MPB, we simulated the model with a 50% increase in
the kinetic parameters controlling Akt- and mTORC1-mediated MPB, both independently and in
combination with all anabolic resistance mechanisms, with or without compensatory signaling.
Increased MPB led to a slight rise in MPS due to greater availability of free intramuscular leucine,
but this effect was offset by a larger increase in MPB, resulting in a net reduction in protein balance
(Figure S6). When these elevated MPB effects were added to the combined anabolic resistance
simulations – with or without compensatory signaling – MPS increased slightly, but the larger rise
in MPB further reduced NB.

Therapeutic strategies to overcome the mechanisms of anabolic resistance

Next, we simulated therapeutic interventions by restoring each dysregulated mechanism to
its healthy value under three scenarios: 1) when the mechanism itself was individually
dysregulated, 2) when the same magnitude of recovery was applied to other individually
dysregulated mechanisms, even though these parameters were not themselves impaired, to
evaluate potential compensatory effects, and 3) when all mechanisms were simultaneously
dysregulated. The effectiveness of each intervention was evaluated against its ability to recovery
of MPS and NB to the healthy responses.

Under the *consensus estimates*, recovering mTORC1 sensitivity fully restored MPS and
NB when applied to individually simulated mechanisms, while recovery of p70S6K levels also
substantially restored both outcomes (Figure 4A-C). However, when all anabolic resistance
mechanisms were simulated together without compensatory signaling, no single therapeutic target
was sufficient to restore MPS or NB. But when we simulated the combined recovery of splanchnic
extraction, mTORC1 sensitivity, and both mTOR and p70S6K levels, near-complete recovery of
MPS and NB was achieved.

When therapeutic strategies were simulated using the *worst-case estimates* (i.e., the
measured values deviating most from the healthy state), recovery of p70S6K levels fully restored

MPS and NB when anabolic resistance mechanisms were simulated individually, while recovery

of either mTOR levels or mTORC1 sensitivity also largely restored both responses (Figure 4D-F).

As with the consensus simulations, no single therapeutic target could overcome the combined

effects of all anabolic resistance mechanisms. However, combined recovery of splanchnic

extraction, mTORC1 sensitivity, and mTOR and p70S6K levels produced a near-complete

restoration of MPS and NB.

We then simulated the therapeutic strategies in the presence of compensatory signaling.

This analysis was based on the hypothesis that compensatory signaling – reflected by elevated

postabsorptive phospho-p70S6K levels in older adults [10,34] – may offset the effects of reduced

mTOR and p70S6K concentrations and decreased mTORC1 sensitivity. Accordingly, we

simulated these three mechanisms together ("[mTOR] + [p70S6K] + mTORC1 sensitivity"), as

well as all anabolic resistance mechanisms together, using the worst-case estimates, in the presence

of compensatory signaling and the applied therapeutic strategies (Figure 5). Compensatory

signaling alone partially restored MPS and NB toward healthy levels in both simulated conditions,

but neither fully reached the healthy level (healthy values: MPS = 0.47 g leucine synthesized, NB

= 0.27 g leucine integrated into muscle; Figure 5). When the three mechanisms were simulated

together, several therapeutic strategies in the presence of compensatory signaling successfully

restored MPS and NB: recovery of p70S6K levels fully restored the MPS response but not NB,

while combining p70S6K recovery with either restored mTOR levels or mTORC1 sensitivity was

sufficient to recover both responses. When all six mechanisms were simulated with compensatory

signaling, additional interventions were required: recovery of mTOR and p70S6K levels along

with mTORC1 sensitivity fully restored MPS but fell slightly short of restoring NB. However, the

further addition of recovered splanchnic extraction fully restored both MPS and NB.

Discussion

In this study we used a computational modeling approach to systematically evaluate the

mechanisms that most likely contribute to anabolic resistance. We first conducted a global

sensitivity analysis of a previously developed kinetic model of leucine-mediated signaling and

muscle protein metabolism [10] to identify mechanisms that most strongly control MPS, MPB,

and NB. The analysis revealed that MPS and MPB are primarily controlled by their proximal

regulatory mechanisms, whereas NB is largely influenced by MPS regulation. Next, we performed

an exploratory analysis simulating an EAA feeding intervention to evaluate mechanisms

contributing to anabolic resistance in older adults [31], classifying simulated responses as either

anabolic sensitive or resistant. Several parameters and signaling protein concentrations differed

significantly between groups, particularly those controlling MPS. We then conducted a targeted

analysis to evaluate the extent to which putative mechanisms underlying anabolic resistance could

explain the reduced MPS observed with aging. Model simulations showed that, when considered

individually, consensus and worst-case estimates of individual mechanisms were unlikely to

account for anabolic resistance. However, when these mechanisms were simulated in combination,

they produced substantial reductions in MPS consistent with anabolic resistance. Finally, we

simulated therapeutic interventions aimed at restoring muscle metabolism in anabolic resistance.

While single-target approaches (e.g., enhancing mTOR or p70S6K levels, or increasing mTORC1

sensitivity) were generally effective when only individual mechanisms were impaired, they proved

insufficient when multiple dysregulated processes were simultaneously present. In contrast,

restoring MPS when all dysregulated processes were simultaneously present required a

multifactorial therapeutic strategy targeting several pathways. These findings highlight the

complex, multifactorial nature of anabolic resistance and motivate novel hypotheses about the

most likely dysregulated mechanisms, providing a foundation for future experimental validation

and multi-targeted therapeutic development.

Main findings and implications

This study examined strategies to restore the feeding-mediated MPS response in older

adults, with a focus on *individual feedings* rather than adjustments to feeding variables such as

dose or scheduling. In our simulations, a single 15 g EAA feeding (containing 3.59 g leucine) was

modelled. While increasing the simulated dose or changing the scheduling of feeding could be

considered as alternatives to improving the MPS responses in older adults, evidence from human

studies suggests these strategies provide limited benefits: in older adults, a 10 g EAA feeding

maximizes myofibrillar MPS rates with no further increases at 20 g or 40 g EAA [18]. Similarly,

studies report that more frequent, smaller feedings (4 × 3.75 g EAA every 45 minutes) [42] or a

leucine top-up after an initial EAA bolus (15 g EAA + 3 g leucine at 90 minutes) [31] do not

meaningfully enhance MPS rates in older adults compared to a single 15 g EAA bolus. Together,

these findings suggest that overcoming anabolic resistance in older adults will require targeting

mechanisms that control the response to individual feedings, rather than modifying nutritional
feeding strategies alone.

In contrast, some studies indicate that increasing protein dose relative to body weight [14]
or enriching the EAA dose with leucine [43] can restore MPS in older adults to levels observed in
younger adults. While these compensatory strategies may offset the blunted anabolic response,
they do not explain *why* higher protein or leucine doses are necessary. The underlying factors
driving this reduced sensitivity are likely contributing to muscle loss, and directly addressing them
offers a more sustainable solution for overcoming anabolic resistance. Accordingly, this study
focused on identifying the mechanisms responsible for the diminished MPS response to individual
feedings and proposes therapeutic strategies to restore this response.

A well-documented mechanism of anabolic resistance is the diminished sensitivity of
mTORC1 signaling to protein or amino acid ingestion [17,18,26]. The findings from our naïve
approach support this notion because we observed widespread dysregulation in leucine-mediated
signaling, including leucine-mediated mTORC1 activity, mTORC1-mediated p70S6K
phosphorylation, p70S6K dephosphorylation, and p70S6K-mediated negative feedback on
mTORC1. While experimental studies have yet to pinpoint the specific mechanisms responsible
for reduced mTORC1 signaling sensitivity in anabolic resistance [42], our results suggest that a
broad dysregulation of the mTORC1 signaling machinery may underlie this phenomenon.

One factor that contributes to reduced mTORC1 signaling sensitivity is physical activity.
A synergistic relationship exists between nutrient- and exercise-mediated mTORC1 signaling and
MPS, in which exercise enhances muscle sensitivity to protein feeding [44] and improves insulin-
mediated amino acid delivery to muscle, thereby supporting mTORC1 signaling [45]. Conversely,
physical inactivity desensitizes muscle to feeding-mediated MPS responses [46]. While our
simulations focused solely on the feeding response, reduced levels of physical activity in older
adults may contribute to the diminished mTORC1 signaling sensitivity seen in anabolic resistance
[47–49]. This combination contributes to the progressive desensitization of mTORC1 signaling to
anabolic stimuli, exacerbating anabolic resistance.

Our simulations from the naïve approach revealed that mTORC1 and p70S6K
concentrations were significantly reduced in the anabolic-resistant group, suggesting a potential
mechanism for the blunted MPS response in aging. Consistent with these findings, previous

research shows that older adults have 50% lower mTOR and p70S6K levels compared to younger

adults that was concomitant with a diminished EAA feeding-induced myofibrillar FSR [18].

Despite these reductions in protein content, older adults exhibit increased postabsorptive phospho-

p70S6K^{T389}, a potential compensatory mechanism. This increased postabsorptive phospho-

p70S6K is seen both in experimental studies [17,18,27,50] and in our previous modeling

simulations [10]. This compensatory elevation likely occurs to partially offset the reduced

mTORC1 and p70S6K concentrations. Indeed, our targeted analysis found that compensatory

increases in postabsorptive phospho-p70S6K can restore a large portion of the MPS and NB

response; however, this compensatory signaling response was insufficient to fully restore muscle

metabolism in our simulations. This persistent postabsorptive phosphorylation reduces the

dynamic range for further postprandial increases [17,18]. This impaired signaling response may

contribute to the diminished MPS observed in older adults following amino acid ingestion.

The compensations in muscle metabolism induced by elevated p70S6K phosphorylation in

aging could be offset by two negative feedback mechanisms. First, phosphorylated p70S6K

inhibits insulin signaling via IRS1 phosphorylation, reducing insulin sensitivity [51–53]. Second,

phospho-p70S6K phosphorylates mTORC1 at Ser2448, further suppressing mTORC1 activation

[54–57]. These feedback pathways likely act together to decrease the sensitivity of p70S6K to

anabolic stimuli and may in part lead to anabolic resistance. Aged rat models similarly exhibit

elevated postabsorptive phosphorylation at both p70S6K and its downstream target, ribosomal

protein S6 (rpS6), further supporting this mechanism [58]. Interestingly, inhibition of mTORC1

signaling with rapamycin in aged rats reduced phoshpo-p70S6K and phospho-rpS6 levels,

resulting in improved muscle mass maintenance and reduced age-related muscle loss [58]. Given

that rapalogs inhibiting mTOR activity have been proposed to improve muscle health with age

[59], our modeling suggests this could further restore skeletal muscle sensitivity to anabolic

stimuli. However, careful titration of doses is needed, particularly in older adults with co-

morbidities and/or low physical activity, to avoid excessive suppression of mTOR signaling and

anabolism [60]. Nevertheless, our results suggest an alternative intervention to inhibiting

mTORC1 – which we hypothesize primarily targets the compensatory response to reduced

p70S6K levels – could be increasing absolute mTORC1 and/or p70S6K. Our model suggests that

increasing mTOR or p70S6K levels, individually, were two of the most efficacious therapeutic

approaches in recovering muscle metabolism. In particular, therapeutic targeting of p70S6K levels

was highly effective in restoring MPS and NB in the presence of compensatory signaling. This
approach of targeting mTOR or p70S6K protein levels could restore the capacity for
phosphorylation increases, reduce postabsorptive mTORC1 signaling, and ultimately mitigate
anabolic resistance.

While this study has identified key potential therapeutic targets to overcome anabolic
resistance, a critical next step is to determine which of these is *druggable*, i.e., can be modified by
a pharmacological agent (cf. [37]). Pharmacotherapies for sarcopenia have a long history of
targeting cellular signaling pathways by binding to intracellular or membrane receptors [38]. These
therapies primarily focus on modulating signaling pathway activity to either increase MPS or
decrease MPB [38]. However, our results suggest that restoring mTORC1 sensitivity and
recovering mTOR and p70S6K protein levels may represent more efficacious strategies to
overcome anabolic resistance – strategies that differ fundamentally from conventional approaches
targeting anabolic or catabolic signalling pathways in muscle. Pharmacological agents such as
MHY1485 [61] and rapamycin [58] can enhance or repress mTOR activity, respectively, offering
opportunities to modulate mTORC1 sensitivity. In contrast, pharmacological strategies aimed at
restoring signaling protein levels in older adult muscle represents a novel and promising avenue.

Determining effective pharmacological strategies to modulate mTOR or p70S6K protein
levels can be informed by bioinformatics-based approaches. Specifically, these proteins are
encoded by the *MTOR* and *RPS6KB1* genes, such that plausible strategies include identifying
factors that upregulate their expression, such as transcription factors (TFs) and cofactors, and
targeting these to increase activity. Conversely, repressors of gene expression, such as microRNAs
(miRNA) and long non-coding RNAs (lncRNAs), could be inhibited to relieve suppression. For
example, miR-199a-3p is a miRNA known to repress *MTOR* expression [62] and could be
inhibited to enhance *MTOR* expression [63]. One issue with targeting miRNAs is that multiple
MTOR-specific miRNAs have been identified, such that targeting just one might be insufficient.
Regarding *RPS6KB1*, estrogen receptor alpha (*ESR1*) is one of the few TFs studied for its role in
controlling *RPS6KB1* gene expression [64,65], making it a potential target to increase *RPS6KB1*
mRNA levels. Additionally, miR-200b [66] and miR-223-3p [67] represent candidate miRNAs
whose inhibition may upregulate *RPS6KB1* expression. While these targets for modulating mTOR
and p70S6K protein levels are speculative, pharmacological approaches that inhibit TF-DNA
interactions, modulate enhancer activity, or repress miRNAs have been developed and successfully

applied in other biological contexts [68–70]. It is important to note that these interventions
primarily affect mRNA levels; therefore, proteomic analyses are essential to confirm
corresponding changes at the protein level and assess the functional impact of these strategies.

Our study features the following noteworthy limitations. First, we evaluated simulated
responses and their approximation to anabolic resistance against two studies [30,31]. Given the
substantial heterogeneity in anabolic resistance and dysregulated muscle protein metabolism with
aging [72,73], future studies should further validate and refine our model using larger, individual-
level datasets. Additionally, future human clinical trials could benefit from measuring the critical
contributors to anabolic resistance identified here to generate robust datasets. Second, our model
was developed using data from males. Some [74], but not all [75], studies suggest that aging-
related dysregulation of muscle protein metabolism may exhibit sexual dimorphism. Therefore,
future studies should include females to confirm the generalizability of our findings. Finally, our
results focus on muscle protein metabolism, which most likely affects muscle mass. While losses
of muscle mass, driven by dysregulations in protein turnover, often precedes declines in muscle
function [73], the relationship between mass and function is not always direct [75].

Conclusions

We report a novel modeling approach to experimentally manipulate key variables
associated with age-related anabolic resistance, without the need for logistically challenging and
costly clinical trials. The initial sensitivity analysis revealed widespread dysregulation of reactions
controlling mTORC1 activity in the anabolic resistant group, along with reduced levels of
mTORC1 and p70S6K. Simulating the putative mechanisms underlying anabolic resistance
suggested that these mechanisms do not act in isolation; rather, they act synergistically to suppress
MPS. Based on these results, we simulated therapeutic interventions to restore muscle metabolism.
Targeting individual mechanisms proximal to mTORC1 signalling (e.g., mTOR or p70S6K levels,
mTORC1 sensitivity) effectively restored MPS and NB when only individual impairments were
simulated. However, these single-target strategies failed to fully restore MPS when all mechanisms
were impaired simultaneously. Under this more realistic condition – reflecting what we suggest is
likely occurring in older adults – a multifactorial therapeutic approach was required to recover
anabolic responses to feeding. These findings motivate new hypotheses regarding the
combinatorial mechanisms underlying anabolic resistance and offer insight for future experimental

studies and therapeutic strategies to mitigate feeding-mediated anabolic resistance. Furthermore,

our framework could also be applied to other models of muscle loss characterized by similar

dysregulations, such as inactivity-induced anabolic resistance [46], or distinct mechanisms, such

as increased MPB in cachexia [71].

Acknowledgements

This work was supported by a Natural Sciences and Engineering Research Council of Canada

(NSERC) Collaborative Research and Training Experience scholarship to T.J.M. and a NSERC

Discovery Grant to D.C.C. (RGPIN 06004-2014). The funders had no role in study design, data

collection and analysis, decision to publish, or preparation of the manuscript.

Conflicts of interest

The authors declare no competing interests.

Figure titles and legends

Figure 1. Kinetic model of leucine-mediated signalling and muscle protein metabolism. The

topology of the McColl and Clarke kinetic model is shown, with hatched red boxes highlighting

the investigated mechanisms of anabolic resistance: 1) increased first-pass splanchnic extraction,

2) reduced transport of amino acids into muscle, 3) reduced insulin-mediated signalling due to

insulin resistance, 4) reduced mTOR concentration, 5) reduced sensitivity of mTORC1 to leucine

signaling, 6) reduced p70S6K concentration, and 7) elevated muscle protein breakdown rates.

Figure 2. Parameter distributions that significantly differ between the anabolic sensitive and

resistant groups. Each panel displays a violin plot showing the parameter distribution density

across the range, accompanied by boxplots indicating the median, first and third quartile, and the

minimum and maximum values, as well as a scatterplot of the data. The y-axis displays the $\log_{10}$-

transformed values of the indicated kinetic model variable.

Figure 3. Simulation of putative mechanisms underlying anabolic resistance: Impact on muscle

protein synthesis and net balance. Muscle protein synthesis (MPS) and net balance (NB) responses

were simulated following a 3.59-gram leucine bolus using either consensus (A, B) or worst-case

(C, D) estimates for six putative mechanisms of anabolic resistance (Table 1). Each mechanism

was individually simulated by multiplying the relevant model parameter by the value estimated

from older adult data. In the "Combined" condition, all six putative mechanisms were

simultaneously simulated. The "Combined (comp. sig.)" condition additionally includes

compensatory signaling (comp. sig.) to simulate elevated postabsorptive phospho-p70S6K levels.

The "Healthy" simulation represents a subset of 100 plausible muscle protein metabolism profiles

that met a conservative MPSA criterion for a healthy individual. This same subset was used to

simulate each anabolic resistance condition, enabling visualization of the potential distribution of

responses across mechanisms.

Each simulated condition displays a violin plot showing the distribution density, overlaid with a

boxplot (median, first and third quartile, and minimum and maximum values), and individual data

points. In the MPS panels, the grey dashed line denotes the MPS response in young adults, as

calculated from Mitchell et al. (2015) [30]. The red dot-dashed line and surrounding red gradient

represent the mean MPS response and 95% confidence interval for older adults, as calculated from

Mitchell et al. (2017) [31]. In the NB panels, the grey dashed line indicates the median NB value

of the healthy simulation.

Figure 4. Simulated therapeutic strategies to overcome anabolic resistance and restore muscle

protein metabolism. Each therapeutic target was simulated across a subset of 100 model parameter

sets by restoring the relevant parameter to its healthy value (i.e., multiplying by the inverse of the

pathological estimate from Table 1). Bolded entries in therapeutic targets 1 to 6 indicate cases

where restoring a single mechanism, together with the simulation of the respective anabolic

resistance mechanism, returned the muscle metabolism response to the healthy level.

Combined therapeutic strategies (denoted by "+") reflect the co-simulation of multiple therapeutic

targets. All individual and combined therapeutic strategies were simulated against each anabolic

resistance mechanism individually and against all six mechanisms combined. The "Pathological

baseline" column shows the median muscle metabolism response across the 100 parameter sets

for each anabolic resistance condition without intervention (Figure 3).

A heatmap displays the simulated median responses for muscle protein synthesis, muscle protein

breakdown, and net balance (NB) across conditions, using either consensus (A-C) or worst-case

(D-F) estimates. Gold indicates recovery to the healthy value; blue indicates a response that would

cause a reduction in NB compared to the healthy value; red indicates a response that cause produce

an increase in NB compared to the healthy value. The numerical value in each cell represents the

median muscle metabolism response – expressed in units of grams of leucine synthesized (MPS),

degraded (MPB), or net incorporated into muscle (NB) – calculated across the subset of 100

simulated model parameter sets.

Figure 5. Simulated therapeutic strategies to overcome anabolic resistance and restore protein

metabolism in muscle cells with compensatory signaling. The worst-case estimates for reduced

mTOR concentration, reduced p70S6K concentration, and reduced mTORC1 sensitivity were

simulated simultaneously ("[mTOR] + [p70S6K] + mTORC1 sensitivity"), along with the

combined simulation of all six anabolic resistance mechanisms ("Combined"). These simulations

were performed without compensatory signaling ("Pathological baseline"), with compensatory

signaling to simulate elevated postabsorptive phospho-p70S6K levels, and with compensatory

signaling plus individual or combined therapeutic strategies (combinations denoted by "+").

A heatmap displays the simulated median responses for muscle protein synthesis, muscle protein

breakdown, and net balance (NB) across conditions. Gold indicates recovery to the healthy value;

blue indicates a response that would produce a reduction in NB compared to the healthy value; red

indicates a response that would produce an increase in NB compared to the healthy value. The

numerical value in each cell represents the median muscle metabolism response – expressed in

units of grams of leucine synthesized (MPS), degraded (MPB), or net incorporated into muscle

(NB) – calculated across the subset of 100 simulated model parameter sets. For reference, the

healthy values for MPS, MPB, and NB were 0.47 g synthesized, 0.154 g degraded, and 0.27 g net

incorporated, respectively.

References

[1] Kirk B, Cawthon PM, Arai H, et al. The Conceptual Definition of Sarcopenia: Delphi Consensus from the Global Leadership Initiative in Sarcopenia (GLIS). Age and Ageing 2024;53:afae052. https://doi.org/10.1093/ageing/afae052.

[2] Beaudart C, Zaaria M, Pasleau F, et al. Health Outcomes of Sarcopenia: A Systematic Review and Meta-Analysis. PLoS One 2017;12:e0169548. https://doi.org/10.1371/journal.pone.0169548.

[3] Steffl M, Sima J, Shiells K, et al. The increase in health care costs associated with muscle weakness in older people without long-term illnesses in the Czech Republic: results from the Survey of Health, Ageing and Retirement in Europe (SHARE). Clin Interv Aging 2017;12:2003–7. https://doi.org/10.2147/CIA.S150826.

[4] Janssen I, Shepard DS, Katzmarzyk PT, et al. The healthcare costs of sarcopenia in the United States. J Am Geriatr Soc 2004;52:80–5. https://doi.org/10.1111/j.1532-5415.2004.52014.x.

[5] Sousa AS, Guerra RS, Fonseca I, et al. Financial impact of sarcopenia on hospitalization costs. Eur J Clin Nutr 2016;70:1046–51. https://doi.org/10.1038/ejcn.2016.73.

[6] Atherton PJ, Smith K. Muscle protein synthesis in response to nutrition and exercise. The Journal of Physiology 2012;590:1049–57. https://doi.org/10.1113/jphysiol.2011.225003.

[7] Phillips BE, Hill DS, Atherton PJ. Regulation of muscle protein synthesis in humans. Current Opinion in Clinical Nutrition and Metabolic Care 2012;15:58–63. https://doi.org/10.1097/MCO.0b013e32834d19bc.

[8] Anthony JC, Yoshizawa F, Anthony TG, et al. Leucine stimulates translation initiation in skeletal muscle of postabsorptive rats via a rapamycin-sensitive pathway. The Journal of Nutrition 2000;130:2413–9. https://doi.org/10.1093/jn/130.10.2413.

[9] Atherton PJ, Smith K, Etheridge T, et al. Distinct anabolic signalling responses to amino acids in C2C12 skeletal muscle cells. Amino Acids 2010;38:1533–9. https://doi.org/10.1007/s00726-009-0377-x.

[10] McColl TJ, Clarke DC. Kinetic modeling of leucine-mediated signaling and protein metabolism in human skeletal muscle. iScience 2024;27. https://doi.org/10.1016/j.isci.2023.108634.

[11] Greenhaff PL, Karagounis LG, Peirce N, et al. Disassociation between the effects of amino acids and insulin on signaling, ubiquitin ligases, and protein turnover in human muscle. Am J Physiol Endocrinol Metab 2008;295:E595-604. https://doi.org/10.1152/ajpendo.90411.2008.

[12] Volpi E, Sheffield-Moore M, Rasmussen BB, et al. Basal muscle amino acid kinetics and protein synthesis in healthy young and older men. JAMA 2001;286:1206–12. https://doi.org/10.1001/jama.286.10.1206.

[13] Dickinson JM, Gundermann DM, Walker DK, et al. Leucine-enriched amino acid ingestion after resistance exercise prolongs myofibrillar protein synthesis and amino acid transporter expression in older men. The Journal of Nutrition 2014;144:1694–702. https://doi.org/10.3945/jn.114.198671.

[14] Moore DR, Churchward-Venne TA, Witard O, et al. Protein ingestion to stimulate myofibrillar protein synthesis requires greater relative protein intakes in healthy older versus younger men. J Gerontol A Biol Sci Med Sci 2015;70:57–62. https://doi.org/10.1093/gerona/glu103.

[15] Bukhari SSI, Phillips BE, Wilkinson DJ, et al. Intake of low-dose leucine-rich essential amino acids stimulates muscle anabolism equivalently to bolus whey protein in older women at rest and after exercise. American Journal of Physiology Endocrinology and Metabolism 2015;308:E1056-65. https://doi.org/10.1152/ajpendo.00481.2014.

[16] Rennie MJ. Anabolic resistance: the effects of aging, sexual dimorphism, and immobilization on human muscle protein turnover. Appl Physiol Nutr Metab 2009;34:377–81. https://doi.org/10.1139/H09-012.

[17] Guillet C, Prod'homme M, Balage M, et al. Impaired anabolic response of muscle protein synthesis is associated with S6K1 dysregulation in elderly humans. FASEB J 2004;18:1586–7. https://doi.org/10.1096/fj.03-1341fje.

[18] Cuthbertson D, Smith K, Babraj J, et al. Anabolic signaling deficits underlie amino acid resistance of wasting, aging muscle. The FASEB Journal 2005;19:1–22. https://doi.org/10.1096/fj.04-2640fje.

[19] Boirie Y, Gachon P, Beaufrère B. Splanchnic and whole-body leucine kinetics in young and elderly men. The American Journal of Clinical Nutrition 1997;65:489–95. https://doi.org/10.1093/ajcn/65.2.489.

[20] Volpi E, Mittendorfer B, Wolf SE, et al. Oral amino acids stimulate muscle protein anabolism in the elderly despite higher first-pass splanchnic extraction. The American Journal of Physiology 1999;277:E513-20. https://doi.org/10.1152/ajpendo.1999.277.3.E513.

[21] Moreau K, Walrand S, Boirie Y. Protein redistribution from skeletal muscle to splanchnic tissue on fasting and refeeding in young and older healthy individuals. J Am Med Dir Assoc 2013;14:696–704. https://doi.org/10.1016/j.jamda.2013.06.005.

[22] Volpi E, Mittendorfer B, Rasmussen BB, et al. The response of muscle protein anabolism to combined hyperaminoacidemia and glucose-induced hyperinsulinemia is impaired in the elderly. J Clin Endocrinol Metab 2000;85:4481–90. https://doi.org/10.1210/jcem.85.12.7021.

[23] Rasmussen BB, Fujita S, Wolfe RR, et al. Insulin resistance of muscle protein metabolism in aging. FASEB J 2006;20:768–9. https://doi.org/10.1096/fj.05-4607fje.

[24] Fujita S, Glynn EL, Timmerman KL, et al. Supraphysiological hyperinsulinaemia is necessary to stimulate skeletal muscle protein anabolism in older adults: evidence of a true age-related insulin resistance of muscle protein metabolism. Diabetologia 2009;52:1889–98. https://doi.org/10.1007/s00125-009-1430-8.

[25] Markofski MM, Dickinson JM, Drummond MJ, et al. Effect of age on basal muscle protein synthesis and mTORC1 signaling in a large cohort of young and older men and women. Exp Gerontol 2015;65:1–7. https://doi.org/10.1016/j.exger.2015.02.015.

[26] Drummond MJ, Dreyer HC, Pennings B, et al. Skeletal muscle protein anabolic response to resistance exercise and essential amino acids is delayed with aging. Journal of Applied Physiology 2008;104:1452–61. https://doi.org/10.1152/japplphysiol.00021.2008.

[27] Wilkes EA, Selby AL, Atherton PJ, et al. Blunting of insulin inhibition of proteolysis in legs of older subjects may contribute to age-related sarcopenia. Am J Clin Nutr 2009;90:1343–50. https://doi.org/10.3945/ajcn.2009.27543.

[28] Hirsch KR, Church DD, Kim I-Y, et al. Comparison of basal whole-body protein kinetics and muscle protein synthesis between young and older adults. Physiol Rep 2020;8:e14633. https://doi.org/10.14814/phy2.14633.

[29] Dalle S, Rossmeislova L, Koppo K. The Role of Inflammation in Age-Related Sarcopenia. Front Physiol 2017;8:1045. https://doi.org/10.3389/fphys.2017.01045.

[30] Mitchell WK, Phillips BE, Williams JP, et al. A dose- rather than delivery profile-dependent
mechanism regulates the "muscle-full" effect in response to oral essential amino acid intake
in   young   men.   The   Journal   of   Nutrition   2015;145:207–14.
https://doi.org/10.3945/jn.114.199604.
[31] Mitchell WK, Phillips BE, Hill I, et al. Human skeletal muscle is refractory to the anabolic
effects of leucine during the postprandial muscle-full period in older men. Clinical Science
(London, England: 1979) 2017;131:2643–53. https://doi.org/10.1042/CS20171230.
[32] Mittendorfer B, Andersen JL, Plomgaard P, et al. Protein synthesis rates in human muscles:
neither anatomical location nor fibre-type composition are major determinants. The Journal
of Physiology 2005;563:203–11. https://doi.org/10.1113/jphysiol.2004.077180.
[33] Zi Z. Sensitivity analysis approaches applied to systems biology models. IET Syst Biol
2011;5:336–336. https://doi.org/10.1049/iet-syb.2011.0015.
[34] McColl TJ. Computational modelling of skeletal muscle protein metabolism and its control.
PhD Dissertation. Simon Fraser University, 2025.
[35] Francaux M, Demeulder B, Naslain D, et al. Aging Reduces the Activation of the mTORC1
Pathway after Resistance Exercise and Protein Intake in Human Skeletal Muscle: Potential
Role   of   REDD1   and   Impaired   Anabolic   Sensitivity.   Nutrients   2016;8:47.
https://doi.org/10.3390/nu8010047.
[36] Lee CH, Cook S, Lee JS, et al. Comparison of Two Meta-Analysis Methods: Inverse-
Variance-Weighted Average and Weighted Sum of Z-Scores. Genomics Inform
2016;14:173–80. https://doi.org/10.5808/GI.2016.14.4.173.
[37] Owens   J.   Determining   druggability.   Nat   Rev   Drug   Discov   2007;6:187–187.
https://doi.org/10.1038/nrd2275.
[38] Yoon JH, Kwon K-S. Receptor-Mediated Muscle Homeostasis as a Target for Sarcopenia
Therapeutics.   Endocrinol   Metab   (Seoul)   2021;36:478–90.
https://doi.org/10.3803/EnM.2021.1081.
[39] Wickham H. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York; 2016.
[40] Rohatgi A. WebPlotDigitizer 2022.
[41] Burd NA, Gorissen SH, van Loon LJC. Anabolic resistance of muscle protein synthesis with
aging.   Exerc   Sport   Sci   Rev   2013;41:169–73.
https://doi.org/10.1097/JES.0b013e318292f3d5.
[42] Mitchell WK, Phillips BE, Williams JP, et al. The impact of delivery profile of essential
amino acids upon skeletal muscle protein synthesis in older men: clinical efficacy of pulse
vs. bolus supply. American Journal of Physiology-Endocrinology and Metabolism
2015;309:E450–7. https://doi.org/10.1152/ajpendo.00112.2015.
[43] Katsanos CS, Kobayashi H, Sheffield-Moore M, et al. A high proportion of leucine is
required for optimal stimulation of the rate of muscle protein synthesis by essential amino
acids in the elderly. American Journal of Physiology Endocrinology and Metabolism
2006;291:E381-7. https://doi.org/10.1152/ajpendo.00488.2005.
[44] Moore DR, Williamson EP, Hodson N, et al. Walking or body weight squat "activity snacks"
increase dietary amino acid utilization for myofibrillar protein synthesis during prolonged
sitting.   J   Appl   Physiol   (1985)   2022;133:777–85.
https://doi.org/10.1152/japplphysiol.00106.2022.
[45] Fujita S, Rasmussen BB, Cadenas JG, et al. Aerobic exercise overcomes the age-related
insulin resistance of muscle protein metabolism by improving endothelial function and

Akt/mammalian target of rapamycin signaling. Diabetes 2007;56:1615–22. https://doi.org/10.2337/db06-1566.

[46] Deane CS, Ely IA, Wilkinson DJ, et al. Dietary protein, exercise, ageing and physical inactivity: interactive influences on skeletal muscle proteostasis. Proc Nutr Soc 2021;80:106–17. https://doi.org/10.1017/S0029665120007879.

[47] Smeuninx B, Mckendry J, Wilson D, et al. Age-Related Anabolic Resistance of Myofibrillar Protein Synthesis Is Exacerbated in Obese Inactive Individuals. J Clin Endocrinol Metab 2017;102:3535–45. https://doi.org/10.1210/jc.2017-00869.

[48] English KL, Paddon-Jones D. Protecting muscle mass and function in older adults during bed rest. Current Opinion in Clinical Nutrition & Metabolic Care 2010;13:34. https://doi.org/10.1097/MCO.0b013e328333aa66.

[49] Sun F, Norman IJ, While AE. Physical activity in older people: a systematic review. BMC Public Health 2013;13:449. https://doi.org/10.1186/1471-2458-13-449.

[50] Horwath O, Moberg M, Hodson N, et al. Anabolic Sensitivity in Healthy, Lean, Older Men Is Associated With Higher Expression of Amino Acid Sensors and mTORC1 Activators Compared to Young. J Cachexia Sarcopenia Muscle 2025;16:e13613. https://doi.org/10.1002/jcsm.13613.

[51] Bertuzzi A, Conte F, Mingrone G, et al. Insulin Signaling in Insulin Resistance States and Cancer: A Modeling Analysis. PloS One 2016;11:e0154415. https://doi.org/10.1371/journal.pone.0154415.

[52] Magnuson B, Ekim B, Fingar DC. Regulation and function of ribosomal protein S6 kinase (S6K) within mTOR signalling networks. The Biochemical Journal 2012;441:1–21. https://doi.org/10.1042/BJ20110892.

[53] Sedaghat AR, Sherman A, Quon MJ. A mathematical model of metabolic insulin signaling pathways. American Journal of Physiology Endocrinology and Metabolism 2002;283:E1084-101. https://doi.org/10.1152/ajpendo.00571.2001.

[54] Ben-Hur V, Denichenko P, Siegfried Z, et al. S6K1 Alternative Splicing Modulates Its Oncogenic Activity and Regulates mTORC1. Cell Reports 2013;3:103–15. https://doi.org/10.1016/j.celrep.2012.11.020.

[55] Figueiredo VC, Markworth JF, Cameron-Smith D. Considerations on mTOR regulation at serine 2448: implications for muscle metabolism studies. Cellular and Molecular Life Sciences : CMLS 2017;74:2537–45. https://doi.org/10.1007/s00018-017-2481-5.

[56] Pearce LR, Alton GR, Richter DT, et al. Characterization of PF-4708671, a novel and highly specific inhibitor of p70 ribosomal S6 kinase (S6K1). Biochem J 2010;431:245–55. https://doi.org/10.1042/BJ20101024.

[57] von Walden F, Liu C, Aurigemma N, et al. mTOR signaling regulates myotube hypertrophy by modulating protein synthesis, rDNA transcription, and chromatin remodeling. Am J Physiol Cell Physiol 2016;311:C663–72. https://doi.org/10.1152/ajpcell.00144.2016.

[58] Joseph GA, Wang SX, Jacobs CE, et al. Partial Inhibition of mTORC1 in Aged Rats Counteracts the Decline in Muscle Mass and Reverses Molecular Signaling Associated with Sarcopenia. Mol Cell Biol 2019;39:e00141-19. https://doi.org/10.1128/MCB.00141-19.

[59] Tang H, Shrager JB, Goldman D. Rapamycin protects aging muscle. Aging (Albany NY) 2019;11:5868–70. https://doi.org/10.18632/aging.102176.

[60] Lees MJ, Hodson N, Tinline-Goodfellow CT, et al. Challenges of rapamycin repurposing as a potential therapeutic candidate for COVID-19: implications for skeletal muscle metabolic

health in older persons. Am J Physiol Endocrinol Metab 2022;322:E551–5.
https://doi.org/10.1152/ajpendo.00064.2022.

[61] Choi YJ, Park YJ, Park JY, et al. Inhibitory Effect of mTOR Activator MHY1485 on
Autophagy: Suppression of Lysosomal Fusion. PLoS One 2012;7:e43418.
https://doi.org/10.1371/journal.pone.0043418.

[62] Shen L, Sun C, Li Y, et al. MicroRNA-199a-3p suppresses glioma cell proliferation by
regulating the AKT/mTOR signaling pathway. Tumor Biol 2015;36:6929–38.
https://doi.org/10.1007/s13277-015-3409-z.

[63] Zhang Y, Huang B, Wang H-Y, et al. Emerging Role of MicroRNAs in mTOR Signaling.
Cell Mol Life Sci 2017;74:2613–25. https://doi.org/10.1007/s00018-017-2485-1.

[64] Berman AY, Manna S, Schwartz NS, et al. ERRα regulates the growth of triple-negative
breast cancer cells via S6K1-dependent mechanism. Sig Transduct Target Ther 2017;2:1–9.
https://doi.org/10.1038/sigtrans.2017.35.

[65] Maruani DM, Spiegel TN, Harris EN, et al. Estrogenic regulation of S6K1 expression creates
a positive regulatory loop in control of breast cancer cell proliferation. Oncogene
2012;31:5073–80. https://doi.org/10.1038/onc.2011.657.

[66] Tao C, Ren H, Xu P, et al. Adipocyte miR-200b/a/429 ablation in mice leads to high-fat-diet-
induced obesity. Oncotarget 2016;7:67796–807. https://doi.org/10.18632/oncotarget.12080.

[67] Dai G-H, Ma P-Z, Song X-B, et al. MicroRNA-223-3p Inhibits the Angiogenesis of Ischemic
Cardiac Microvascular Endothelial Cells via Affecting RPS6KB1/hif-1a Signal Pathway.
PLOS ONE 2014;9:e108468. https://doi.org/10.1371/journal.pone.0108468.

[68] Lovén J, Hoke HA, Lin CY, et al. Selective Inhibition of Tumor Oncogenes by Disruption of
Super-Enhancers. Cell 2013;153:320–34. https://doi.org/10.1016/j.cell.2013.03.036.

[69] Blume SW, Snyder RC, Ray R, et al. Mithramycin inhibits SP1 binding and selectively
inhibits transcriptional activity of the dihydrofolate reductase gene in vitro and in vivo. 1991.
https://doi.org/10.1172/JCI115474.

[70] Janssen HLA, Reesink HW, Lawitz EJ, et al. Treatment of HCV Infection by Targeting
MicroRNA. New England Journal of Medicine 2013;368:1685–94.
https://doi.org/10.1056/NEJMoa1209026.

[71] Setiawan T, Sari IN, Wijaya YT, et al. Cancer cachexia: molecular mechanisms and treatment
strategies. Journal of Hematology & Oncology 2023;16:54. https://doi.org/10.1186/s13045-
023-01454-0.

[72] Paulussen KJM, McKenna CF, Beals JW, et al. Anabolic Resistance of Muscle Protein
Turnover Comes in Various Shapes and Sizes. Front Nutr 2021;8.
https://doi.org/10.3389/fnut.2021.615849.

[73] Moore DR. Keeping Older Muscle "Young" through Dietary Protein and Physical Activity.
Advances in Nutrition 2014;5:599S-607S. https://doi.org/10.3945/an.113.005405.

[74] Smith GI, Atherton P, Villareal DT, et al. Differences in muscle protein synthesis and
anabolic signaling in the postabsorptive state and in response to food in 65-80 year old men
and women. PLoS One 2008;3:e1875. https://doi.org/10.1371/journal.pone.0001875.

[75] Cegielski J, Brook MS, Phillips BE, et al. The Combined Oral Stable Isotope Assessment of
Muscle (COSIAM) reveals D-3 creatine derived muscle mass as a standout cross-sectional
biomarker of muscle physiology vitality in older age. Geroscience 2022;44:2129–38.
https://doi.org/10.1007/s11357-022-00541-3.