Title: **Bottom-up effects of a megaherbivore alter plant growth**

and competition regimes, promoting vegetation heterogeneity

Authors: Hansraj Gautam[*1,2], Thanikodi M[3], and Mahesh Sankaran[1]

Affiliations: 1) National Centre for Biological Sciences, Bengaluru, India, 2) Department of

Biology, University of Turku, Finland. 3) Jawaharlal Nehru Centre for Advanced Scientific

Research, Bengaluru, India.

Email of authors: Hansraj Gautam: gautamhansraj@yahoo.com, hansraj.gautam@utu.fi,

Thanikodi M: thaninivi2@gmail.com, Mahesh Sankaran: mahesh@ncbs.res.in

Keywords: megafauna, large herbivores, functional role, bottom-up effects, nutrient cycling,

heterogeneity, neighborhood competition, nitrogen-fixers, zoogeochemistry

*Corresponding author: Hansraj Gautam, Department of Biology, University of Turku,

Turku 20014, Finland. ORCID: https://orcid.org/0000-0001-8289-5562, Contact:

gautamhansraj@yahoo.com, hansraj.gautam@utu.fi, Telephone: +358 041 742 1966

AUTHOR CONTRIBUTIONS

HG: funding acquisition, conceptualization and research design, fieldwork and laboratory analyses, formal statistical analyses, writing the first draft, review and editing; MT: fieldwork; MS: research design, review and editing. All authors read and approved the manuscript for submission.

Conflict of interest: Authors do not have any conflict of interest.

Data Availability Statement: All data files and scripts used to prepare plots and analyses tables are attached as supplementary files. The data will be uploaded on a publicly accessible repository platform (Zenodo) after acceptance.

ABSTRACT

1. Megaherbivores are known to strongly influence multiple ecological processes, but their

bottom-up impacts on vegetation via nutrient redistribution remain poorly understood,

particularly in mesic ecosystems. Here, we investigated how woody plant communities are

influenced by large nutrient inputs in the form of dung deposited by Asia's largest herbivore,

the Asian elephant (*Elephas maximus*), in the tropical forests of southern India.

2. We conducted field and mesocosm experiments on woody saplings to examine three

mechanisms through which dung deposition by elephants can alter plant community

assembly. Specifically, we tested if elephant dung input 1) creates hotspots of plant growth,

and shapes plant communities by altering 2) the negative density-dependent effects of

neighborhood competition, and 3) interference competition between plant functional types

differing in nutrient limitation, namely nitrogen-fixers and non-nitrogen-fixers.

3. Our findings show that dung deposition by elephants can generate fine-scale spatial

differences in woody sapling communities by creating local growth hotspots and altering

competitive interactions. We analyzed relative growth rate and final sapling size in the field

experiment, and found that average-sized and large saplings receiving dung inputs were

buffered against the negative density-dependent effects of neighborhood competition. In the

mesocosm experiment, non-nitrogen-fixing species (which are nitrogen-limited) outcompeted

nitrogen-fixers in accumulating biomass when supplied with elephant dung. These outcomes

were associated with changes in their relative competitive strength which was stronger for

non-nitrogen fixers under dung treatment and for nitrogen-fixers under control. Such bottom-

up effects on plant growth and competition can be of substantially large magnitude, as we

estimated that elephants in these forests create a total of 11000 such nutrient-rich sites

/km^2/year, with each elephant redistributing ~130 kg Nitrogen/year through this pathway and

each site receiving ~20 g Nitrogen.

Synthesis: Our findings on the outcomes of sapling competition highlight the role of nutrient

redistribution by megaherbivores as an underappreciated driver of species interactions that

can alter plant communities at fine scales, effects that are widespread across megaherbivore

habitats. Such bottom-up effects of megaherbivores, along with their top-down effects, have

important conservation implications and can help in restoring species interactions and spatial

heterogeneity in plant communities in defaunated habitats.

INTRODUCTION

Large herbivores are integral to the functioning of many ecosystems as they regulate multiple ecosystem processes (Pringle et al., 2023). They have pronounced impacts on biodiversity through their effects on vegetation structure and composition (Jia et al., 2018; Van Der Plas et al., 2016), nutrient cycling (Augustine et al., 2003; Coetsee et al., 2023; Pastor et al., 2006; Villar et al., 2021) and fire regimes (Archibald & Hempson, 2016; Karp et al., 2024), in addition to downstream effects on other organisms (McCleery et al., 2024; Trepel et al., 2024). Worryingly, while ecosystems worldwide have become functionally degraded due to the body-size-downgrading of herbivore communities (Dirzo et al., 2014), our current understanding of their functional roles is strongly biased, both taxonomically and geographically (Pringle et al., 2023; Ripple et al., 2015; Sankaran & Ahrestani, 2016; Trepel et al., 2024). Such regional and taxonomic blind-spots are particularly evident for megaherbivores i.e., herbivores with >1000 kg body weight (Hyvarinen et al., 2021). Megaherbivores are thought to have disproportionate impacts on ecosystem functioning (Owen-Smith, 1988) and are considered keystone species (Malhi et al., 2016). Yet, empirical understanding of their various ecological roles is scarce for most species (systematic review in Hyvarinen et al., 2021), which hinders a generalized understanding of the functional importance of a very large body size (Pringle et al., 2023). This limitation impedes locally informed application of such knowledge in conservation, particularly in rewilding and restoration of ecological functions lost from defaunated ecosystems (Geremia et al., 2025; Ripple et al., 2015; Svenning et al., 2016).

Among large herbivores, the disproportionate ecological impacts of a very large body size stem from its association with unique functional traits (Owen-Smith, 1988; Pringle et al.,

2023). For instance, the large daily forage requirement, generalist diet and destructive

foraging behaviour of megaherbivores exert strong top-down control on plants, while their

wide-ranging behaviour promotes long-distance transport of nutrients and seeds (Berzaghi et

al., 2018; Malhi et al., 2016; Trepel et al., 2024). Being largely immune as adults to top-down

control by non-human predators, megaherbivores are instead bottom-up limited by resource

availability – such bottom-up limitation amplifies their effects on vegetation (Hyvarinen et

al., 2021). In recent reviews, their ecological roles have been emphasized as so uniquely

important for conservation that their per-capita impact cannot be compensated for by

managing densities of smaller herbivores (Hyvarinen et al., 2021; Pringle et al., 2023).

Recent meta-analyses of herbivore exclusion experiments have concluded that megafauna

alter the composition of plant communities (e.g. species richness and evenness, Jia et al.,

2018) and enhance spatial heterogeneity in vegetation and soil nutrients (Trepel et al., 2024).

Such megafauna-induced changes in community composition and heterogeneity can arise via

top-down (consumptive effects) and/or bottom-up (alteration of resource availability)

regulatory pathways (Ferraro et al., 2024; Hobbs, 1996; Knapp et al., 1999; Pringle et al.,

2023). Past studies on megaherbivores have predominantly focused on the top-down

regulation of vegetation through consumption and trampling, with most research concentrated

in African savannas (Hyvarinen et al., 2021). In contrast, their bottom-up effects are poorly

understood and have been studied primarily through the lens of seed dispersal (Campos-

Arceiz & Blake, 2011). A critical bottom-up impact of megaherbivores is nutrient distribution

and cycling (through dung, urine and carcasses, Wolf et al., 2013), but this function remains

largely understudied for megaherbivores other than hippos and savanna elephants (Hyvarinen

et al., 2021). Studying the impacts of this key ecological role can shed light on the

coexistence dynamics (Sitters & Olde Venterink, 2021), structure and heterogeneity of plant

communities.

Megaherbivores are strong agents of nutrient cycling as they return large amounts of nutrient-

rich faeces and urine (Le Roux et al., 2020; Pastor et al., 2006; Wolf et al., 2013) and

facilitate long-distance transport of nutrients, like "arteries" of ecosystems (Malhi et al.,

2016; Wolf et al., 2013). The patchy deposition of nutrient-rich faeces by megaherbivores

creates a heterogeneous matrix of nutrient availability (Afzal & Adams, 1992; Ferraro &

Lienau, 2025; Le Roux et al., 2020), which in turn, can generate fine-scale mosaics of

vegetation in multiple ways. These nutrient-mediated effects could alter plant growth and

survival (Tandalla et al., 2024), as well as changes in ecological interactions among plants

such as competition (Van der Waal et al., 2009). For instance, herbivore dung can alter the

competitive outcomes between species differing in nutrient limitation, thereby shaping

community composition (Sitters & Olde Venterink, 2018, 2021). Because such responses are

mediated by the quantity of nutrient inputs via dung (Sitters & Olde Venterink, 2021; Valdés-

Correcher et al., 2019), megaherbivores are expected to exert disproportionate effects since

they deposit large amounts of dung. However, empirical examinations of such bottom-up

impacts remain scarce (Kalbitzer et al., 2019; Sitters & Olde Venterink, 2021). Therefore, our

study aimed to elucidate the mechanisms through which nutrient deposition by

megaherbivores influences plant communities, using elephants as a model system.

Elephants, the world's largest herbivores, are often referred to as "ecosystem engineers" or

"megagardeners" due to their large impacts on ecosystem processes (Campos-Arceiz &

Blake, 2011; Poulsen et al., 2018). Existing knowledge on their ecological roles largely

derives from studies on the African savanna elephant (Guldemond et al., 2017; for example,

Sukumar, 2003 pp. 223-252) that occupies arid and semi-arid open habitats. In contrast, the

African forest (*Loxodonta cyclotis*) and Asian elephants remain understudied, with research

primarily focused on their role as seed dispersers (Campos-Arceiz & Blake, 2011; Sekar et

al., 2017; but see Berzaghi et al., 2023; Blake, 2002; Omeja et al., 2014; Sivasubramaniyan &

Sivaganesan, 1996). Studying these two species can elucidate the impacts of megaherbivores

in wetter niches, a key gap identified in Hyvarinen et al.'s systematic review (2021).

Currently, the impacts of nutrient distribution by elephants on vegetation are poorly

understood for all three species. Experimental manipulation of dung input can reveal its

effects on vegetation (Kalbitzer et al., 2019; Sitters & Olde Venterink, 2021; Valdés-

Correcher et al., 2019), but to our knowledge, only one such study exists. Interestingly, this

study by Kalbitzer et al. (2019) did not find growth-enhancing effects of elephant dung on

plants in central African rainforests. Given the lack of other field experiments, further

evaluation of the supposedly large bottom-up impacts of elephants on vegetation is required.

Here, we conducted field and ex-situ mesocosm experiments manipulating elephant dung

input to woody saplings in the mesic tropical forests of southern India to understand bottom-

up regulation of vegetation by megaherbivores. We hypothesized that elephants create fine-

scale differences in nutrient availability through dung deposition, which in turn, alters plant

growth and competition regimes, with implications for local heterogeneity in vegetation (Fig

1). We tested three predictions linked to the distinct mechanistic responses by plants. First,

we expected dung-deposition sites to act as growth hotspots compared to adjacent control

sites (*Prediction 1*). Second, we predicted that such nutrient input via dung provides a

localized buffer against negative density-dependent effects on growth (also known as the

neighborhood effect or scramble competition, *Prediction 2*). We then examined how dung

input affects competitive balance between plant functional types (PFT) differing in nutrient

limitation (inter-PFT interference competition). To this end, we chose nitrogen-fixing (N-

fixers) and non-nitrogen-fixing (non-N-fixers) woody plants; these two PFTs present an

excellent contrast model as they fundamentally differ in their ability to acquire nitrogen,

which often limits plant growth in many ecosystems (Vitousek & Howarth, 1991). Non-N-

fixers rely solely on available forms of nitrogen in soils and thus benefit more from external

nitrogen inputs, whereas N-fixers can additionally also access atmospheric di-nitrogen

through symbionts (although this N-fixing ability can be expensive and is typically limited by

other nutrients like Phosphorous, Tandalla et al., 2024; Varma et al., 2018). Localized

deposition of nitrogen-rich dung is expected to alter the coexistence dynamics between these

two plant types by alleviating nitrogen-limitation for non-N-fixers. While a previous study

showed that herbivore dung alters tree-grass coexistence by shaping competition between

grasses and woody N-fixers (Sitters & Olde Venterink, 2021), understanding dung-impacts

on the competition between woody non-N-fixers and woody N-fixers is important in mesic

ecosystems where woody non-N-fixers are more abundant. We predicted that woody non-N-

fixers, being nitrogen-limited, would be able to better compete with N-fixers when supplied

with nitrogen-rich dung, such that dung addition would lower the relative competitive ability

of N-fixers (relative to non-N-fixers) as compared to control conditions (*Prediction 3*).

Finally, to understand the overall magnitude of these bottom-up effects, we estimated the

annual production of such nutrient-rich sites by elephants.

Figure 1. a) Top-down and bottom-up effects of megaherbivores as distinct regulatory

pathways that can enhance local heterogeneity by promoting fine-scale spatial differences in

vegetation, **b)** Hypothesized mechanisms through which nutrient input via dung deposition

can create fine-scale differences between vegetation that receives such nutrient input and the

adjacent control sites.

MATERIALS AND METHODS

MATERIALS AND METHODS

We conducted this study in and around the tropical forests of Nagarahole National Park

(henceforth called Nagarahole). Tropical moist deciduous forests, tropical dry deciduous

forests and teak plantations are the major vegetation types in the park, with mesic and hot

conditions characterized by clear seasonality in rainfall and temperature (Fig. 2b). In the two

decades preceding this study, mean annual rainfall ranged from ~1000-2300 mm and ambient

temperature ranged from ~11 °C to ~36 °C (Fig. 2a, Table S8, Supplement text B). The soil is

loamy, clay-loamy and deep (Reddy et al., 2016).

Figure 2: Climatic variation in Nagarahole National Park during 2012-2023 (years

overlapping this study are marked). a) Inter-annual variation in total rainfall and ambient

temperature (min-max range shown in orange). b) Seasonal variation in ambient temperature

and total rainfall (thick lines represent long-term mean across years).

Field experiment on woody plant growth and neighborhood competition

To examine if elephant dung deposition creates spatial differences in plant growth and

neighborhood competition, we selected 19 experimental sites in September 2022 in different

locations in the park, representing a range of ecological conditions spanning the three forest

types. These sites were at least 2km apart from each other and were in randomly selected

2km x 2km cells, with their GPS locations pre-determined based on sampling in a previous

study (*Reference removed for double-blind review*). At each location, we established two

pairs of plots, 20m apart; each pair consisted of a control and a dung treatment plot ($1m^2$) that

were 10m apart (plot-design in Fig S1). We first sampled plots to quantify the initial state of

the vegetation before applying the dung treatment. All woody individuals ≥30cm tall with

diameter at breast height less than 10cm (henceforth called saplings) in plots were marked

with uniquely numbered aluminum tags, and their basal diameter at 10cm aboveground

measured using an INSIZE 1112-150 digital caliper (0.01mm resolution). For individuals

with multiple stems, we separately measured the diameter of all stems at 10cm height, and

then obtained the cumulative sum of stem diameters (henceforth called 'basal diameter'). We

also measured their height up to 1m using a scale (saplings taller than 1m were categorized

into 1-2m or >2m bins).

After the initial vegetation inventory, we experimentally added dung to one plot (henceforth

called treatment plot) in each pair. The treatment involved placement of an entire pile (5-

10kg) of fresh dung in each plot. We collected dung piles from fresh (or overnight)

defecations by adult wild elephants and transferred them to the treatment plot on the same

day. We re-sampled the basal diameter and height of marked saplings after ~6 months during

April and May 2023. We also noted the status (dead/alive) of saplings; in case of saplings

with multiple stems, if the main stem had died and only the smaller stem(s) survived, we

classified it as 'partially dead'. The total duration between the initial and final measurements

was 162 days on average, but varied due to logistics (143-215 days). One site burned in a

forest fire, leaving us with 18 sites or 36 plot-pairs.

Ex-situ mesocosm experiment on competition between plant functional types (PFT)

To examine if dung input affected the competitive dynamics between nitrogen-fixers and

non-fixers, we set up a semi-natural mesocosm experiment at a site ~3km from the

southeastern boundary of Nagarahole, in September-October 2022. The site had naturally

regrowing vegetation with an overstory of deciduous trees and a dense understory. At the site,

we enclosed a flat rectangular area ~600m^2 using a wire mesh (one-inch gap). We further

covered the enclosure with a plastic net to prevent the entry of birds and mammals or falling

tree branches while retaining the effects of sunlight and canopy. We selected seedlings (8-10

weeks) of native N-fixing (*Dalbergia latifolia*, *Pterocarpus marsupium*, *Butea monosperma*,

Senegalia ferruginea (*Acacia ferruginea*)) and non-N-fixing tree species (*Terminalia*

tomentosa, *Terminalia anogeissiana* (*Anogeissus latifolia*), *Tectonia grandis*, *Zizyphus*

oenoplia, *Cassia fistula*, *Terminalia bellirica* and *Syzigium cumini*). These seedlings were

obtained from nearby forestry nurseries (Tithimati, Beechanahalli) where seeds collected

from natural forests were germinated for plantation/restoration purposes. For *Terminalia*

anogeissiana and *Zizyphus oenoplia*, which are widespread in this landscape but were not

available in nurseries, we collected similar sized seedlings from vegetation outside the park

boundary. Together, these species represent >50% of the naturally occurring community of

tree seedlings in Nagarahole forests (based on data in Gautam et al., 2019). We used a total of

132 seedlings in this experiment. Seedlings were grown in large, rectangular pots (~18"

length, 12" width, 11" depth, i.e., with surface area of 0.14 m^2 on top) and local soil from

uncultivated land adjacent to the eastern border of the park. Pebbles and plant debris in soils

were first removed, after which the soil was mixed with sand in a 2:1 ratio to ensure water

percolation and prevent clogging. Pots were then filled with this soil-sand mixture up to

~90% of their heights (~16kg), and planted with two seedlings, as described below.

We planted two seedlings (of comparable basal diameter, height) ~10 inches apart in each pot

to examine how elephant dung input influences seedling competition. We created two

competition scenarios for each plant functional type (PFT): i) inter-PFT or mixture

competition, i.e., competition between one N-fixer and one non-N-fixer sapling, and ii)

monoculture or intraspecific competition, i.e., competition within each PFT. This resulted in

two types of monocultures, one each for N-fixers and non-N-fixers. We maintained two sets

of each monoculture and mixture regime pots, one being the control while the other received

dung treatments, in each block (N=11 blocks of six pots containing two seedlings each). The

dung treatment involved placing 500g (wet weight, ~2/3rd of a bolus) of fresh elephant dung

between the two seedlings (adding whole dung pile is not possible due to limit pot size).

Since we were interested in inter-PFT competition, we ensured a balanced sample size for

each PFT (N=66); however, the competition design resulted in twice the number of seedlings

under monoculture than the mixture (inter-PFT) competition. Replication at the species level

was limited by logistical constraints (availability of seedlings in nurseries, soil requirements

for large pots, and enclosure size). At the seedling level, this experiment had a 2

(dung/control treatment) × 2 (PFT: N-fixers/non-N-fixers) × 2 (competition type: inter-PFT/

within-PFT (or intra-specific i.e., monoculture) design. All pots were watered 2-3 times a

week to prevent sapling mortality, with every pot receiving similar amount of water. After the

experiment ended in April 2022 (after ~5-6 months), all saplings were harvested for dry

biomass measurements. We separated shoots from roots (at the soil mark) to quantify shoot

and root biomass separately, and then added them to obtain final sapling biomass. Any soil

sticking to the saplings was brushed off, after which the saplings were dried in an oven at 60

°C for 60 hours and their dry weight measured in the laboratory.

Statistical analyses of field experiment data

Our analysis of the field experiment was based on all saplings alive at the end of the

experiment (N=418 saplings in 18 plot-pairs, raw mortality rates provided in Results). The

treatment and control plots did not seem to differ in initial diameter of saplings (raw plot-

wise means in Fig. S1). We obtained two response variables for basal diameter: relative

growth rate ($RGR_{diameter}$) and final basal diameter. RGR was calculated as [ln (final size) – ln

(initial size)]/ *t*, where *t* is the duration of experiment. We analyzed RGR, which reflects

biomass production per unit existing size, as it is an effective metric for quantifying growth

efficiency at the seedling/sapling stage when plants are growing exponentially. As both

nutrient supply and competition can modify this efficiency independent of plant size, RGR is

useful in capturing performance under different ecological conditions. In contrast, final size

represents the outcome of competition and nutrient input given an initial size. Therefore,

these two variables inform about two complementary aspects of sapling stage, performance

(RGR) and outcome (size).

We first examined if dung treatment influenced $RGR_{diameter}$, by computing Gaussian general

linear mixed effects models with maximum likelihood (*lmer* function of the R package *lmer4*

Bates et al., 2015). We included treatment (dung/control), sapling density and forest type as

fixed effects, while also controlling for the effect of initial basal diameter by including it as a

fixed effect because RGR of smaller saplings tends to be greater. We further included the

random intercepts for species and plot-pair identity to account for potential non-independence

of individuals within sites and species. All continuous predictors were standardized (mean =

0, standard deviation = 1) to make their effect sizes comparable. Further, we included the

three-way and two-way interactions involving dung treatment, sapling density and initial

diameter, since one of our objectives was to examine if elephant dung mediates neighborhood

competition which could depend on the focal sapling's size-dependent nutrient requirements

and performance. A positive effect of the interaction involving dung treatment and sapling

density would suggest that dung treatment buffers saplings against negative-density

dependence. We did not simplify the full three-way interaction between treatment, sapling

density and size, as all these interactions with treatment were of our *apriori* interest. Upon

inspection of the Gaussian GLMM (identity link), we found strong heteroscedasticity, with

residual variance increasing with initial size and differing across forest type (assessed using

simulated residual diagnostics, *DHARMa* package Hartig & Hartig, 2017). We therefore

fitted generalized linear mixed effects models (GLMMs) with a Student t family of error

distribution and a dispersion structure dependent on forest type and initial diameter, as

summarized in Table 1 (using *glmmTMB* package Brooks et al., 2017). To understand the

overall effect of dung treatment, we estimated the marginal means for the two treatment

levels using the *emmeans* package and tested their difference using a Wald z-test on the

treatment-control contrast (*emmeans* averaged over forest types). Similarly, we examined the

treatment-control contrast at specified levels of density and initial size, and obtained the

conditional-effects plot for the 3-way interaction using *ggpredict* function. We also present

the summary table of conditional effects of the 3-way interaction for dry deciduous forest

which dominated our sample. Further, since RGR takes very small values, we used predicted

$RGR_{diameter}$ to derive predicted values for two more intuitive variables, i.e., percentage change

and absolute change in diameter. Percent change was calculated as $100 \times (\exp (emmean \times t)$

$-1)$, and absolute change as $(\exp (emmean \times t) -1) \times$ initial diameter), for t = 215 days (full

duration of the experiment) (derivations in Supplement text A). These were then visualized

for three discrete initial sapling sizes, i.e., average, small and large (mean initial diameter,

mean $- 0.5SD$, mean $+1SD$), for treatment and density (mean and $\pm 1SD$) contrasts. For small

sapling size, we chose mean $- 0.5$ SD because in our dataset initial diameter is right-skewed

and mean $- 1SD$ does not exist.

Similarly, we developed GLMMs for final diameter and final height, while taking into

account effects of initial size by including it as a fixed effect, as specified in Table 1. In the

case of sapling height, since we had measured height only up to 1m, we excluded all saplings

whose initial and final height was >1m and modelled final height for this filtered subset of

308 saplings.

We analyzed sapling mortality (binary response variable) in our full dataset (N=432) using a

binomial GLMM with a logit link. We also inspected net change in the count of stems (stem-

count) between initial and final measurement of saplings. Net change in stem-count can result

from a) natural growth trajectory where new stems sprout and grow, and b) partial mortality

i.e., mortality of some of the stems. Finally, we selected the subset of saplings whose stem-

count remained unchanged (N=316) and repeated the above analyses on $RGR_{diameter}$ and final

basal diameter, with models specified in Table 1, to check if the trends of growth in this

subset reflected the full dataset.

Table 1. Generalized linear mixed model specifications for different response variables. PFT:

plant functional type (nitrogen-fixer/non-nitrogen-fixer).

Response Variable	Fixed effects	Random effects	Error distribution family	Dispersion structure (in heteroscedastic models)
1. Field experiment				
1.1. RGR_diameter (mm/mm/day)	Dung $\times$ density (std) $\times$ initial diameter (std)	Species + Plot-pair	Student t	initial diameter (std) + forest type
1.2. Final diameter (mm)	Dung $\times$ density (std) $\times$ initial diameter (std)	Species + Plot-pair	Student t	log (initial diameter) + forest type
1.3. Final height (cm)	Dung $\times$ density (std) $\times$ ploy(initial height (std), 2)	Species + Plot-pair	Student t	initial height (std)
1.4. Mortality (1/0)	Dung	-	Binomial	-
2) Mesocosm experiment				
2.1. Final biomass (g)	Dung $\times$ PFT $\times$ Competition type + Initial size index (std)	Species + Pot ID + Block	Student t	PFT
2.2. Relative Competitive Strength (ratio based on biomass)	Dung $\times$ PFT + Initial size index ratio (std)	Species + Pot ID	Gamma (link = log)	PFT

Statistical analyses of mesocosm experiment data

To examine how dung treatment affects competition between the two plant functional types

(PFT), i.e., N-fixers and non-N-fixers (henceforth inter-PFT competition), we analyzed the

final sapling biomass data from the mesocosm experiment in two ways. First, we tested the

effect of elephant dung on final sapling biomass, using a GLMM containing the fixed effects

arising from a full interaction between treatment (control/dung), PFT (N-fixer/non-N-fixer)

and competition type (mixture/monoculture). To account for any variation due to initial size,

initial size index (standardized, mean = 0, SD = 1) was included as a fixed effect, while

species identity and block were used as random effects. The initial size index was based on

an allometric equation (diameter^2 × height). We inspected the outcomes under inter-PFT

competition to infer the effect of elephant dung on the competitive ability of each PFT. We

considered whether N-fixers or non-N-fixers attained greater final biomass under the dung

treatment or control, when in inter-PFT competition. For this, we obtained the estimated

marginal means for further comparison.

Second, we quantified the relative competitive strength (RCS) of each plant functional type

(PFT), using observed sapling total biomass as a measure of performance under each

competition type within each block

$Relative\ competitive\ strength = \dfrac{final\ biomass\ under\ inter-PFT\ competition}{final\ biomass\ under\ intraspecific\ competition}$ *Equation 2*

We considered a PFT to be a better competitor under inter-PFT competition if its RCS

exceeded 1.0 or if it reduced the RCS of the competing PFT (Sitters & Olde Venterink,

2021). To examine this, we used a GLMM (Gamma distribution) to model RCS as a function

of the fixed effects arising from a full interaction between treatment (control/dung) and PFT

of the focal saplings (N-fixer/non-N-fixer). We also accounted for variation in RCS due to

the fixed effects of the initial size differences between the saplings being compared

(allometric size index in mixture pot/size index in monoculture pot) and the random effects of

species identity. Further, we included mixture pot ID as a random effect because for the focal

sapling of each PFT competing in each mixture pot (i.e. inter-PFT competition pot), two RCS

values were obtained due to comparison of the focal sapling with two saplings of the same

species in the corresponding monoculture pot within the block. We inferred a PFT's

competitive ability in two ways. We first tested the estimated effects of dung treatment to

examine if it significantly shifted the RCS value of the focal PFT, compared to control

conditions. Second, we examined the 95% confidence intervals of the estimated marginal

means (using *emmeans*) of each group in the treatment × PFT interaction to test if each PFT's

RCS (95% confidence interval) was different from the null expectation of 1.0 under control

and dung treatment. RCS = 1.0 implies similar performance in inter-PFT and conspecific

competition. We then checked if each PFT's "RCS>1.0 or not" status changed from control

to dung treatment conditions.

Quantifying nitrogen redistribution by elephants

To understand the broader implications of our experiments, we estimated annual rates of

nitrogen redistribution per km^2 by elephants in Nagarahole following an earlier approach to

quantify carbon stocks in elephant dung (Sandhage-Hofmann et al., 2021) as follows:

$\text{Nitrogen (kg km}^{-2}\text{y}^{-1}) = \text{avg. wet weight of dung pile (kg)} \times$

$\text{dry weight as proportion} \times \text{N\% in dung (mg/g)} \times \text{dung deposition rates (day}^{-1}) \times$

$\text{elephant density (km}^{-2}) \times 365 \text{ (days)}$ *Equation 1*

We obtained average wet weight by measuring whole fresh dung piles (N=42) in the field. To

estimate dry weight, we collected ~200-400g of sample for each dung pile, sun dried them

initially, and then dried them further in the lab at 60° C for ~60 hours. We then weighed them

to obtain dry weight as a fraction of the original wet weight. Nitrogen content in elephant

dung was quantified for three seasons using LECO CN analyzer (LECO Corporation, St.

Joseph, MI) in a parallel study on elephant diet (*Reference removed for double-blind review*),

to which we added data from another season (Table S7). Elephant density in Nagarahole and

dung deposition rates of Asian elephants were taken from previous studies (Table S7).

RESULTS

1. Field experiment:

1a) *Elephant dung increases sapling growth and alters neighborhood competition*: Spatial

differences in sapling growth in response to dung deposition were clearly evident, supporting

Prediction 1. The raw percentage increase in the basal diameter of saplings under dung

treatment was 16% (n=194 saplings, SD=33.7), while it was only 9% in the control plots 10m

away (n=224, SD=25.5; plot-wise means in Fig. S1). Accordingly, the GLMM of relative

growth rate (RGR) showed that dung addition enhanced $RGR_{diameter}$ by over 70% relative to

control saplings (comparison of estimated marginal means, *P*=0.040, Fig. 3a). Density had a

negative effect on the $RGR_{diameter}$ of larger saplings under control conditions ($P = 0.033$) and

while dung treatment appeared to reduce this negative effect of density on large saplings, this

3-way interaction effect was statistically not significant ($P = 0.12$; Table S1). We obtained

the estimated marginal means at different sapling densities for three discrete initial sizes and

found a size-dependent buffering against the negative effects of high density, consistent with

Prediction 2. Dung treatment provided a significant advantage at average and high sapling

densities for average-sized (mean initial diameter) and large (mean + 1SD) saplings, whereas

this advantage was not seen at lower densities and for small saplings (Fig. 3b and 3c). Fig 3c

shows these effects in terms of percentage and absolute growth in diameter derived from

predicted $RGR_{diameter}$ for discrete values of sapling density and initial size.

The positive effects of dung treatment were statistically more evident in the GLMMs of final

sapling size (Table 1). In addition to the overall positive marginal effect of dung treatment

(comparison of *emmeans*: P=0.035, Fig 4a), this effect was greater for large saplings which

were also buffered against the negative effects of high sapling density under dung treatment,

consistent with *Prediction 2* (Fig 4b, Table 1). In contrast, the GLMM of final height

revealed that the positive marginal effect of dung treatment was statistically not significant

(based on *emmeans*, *P*=0.063, Fig 4c). Saplings increased their height under crowding, but

this was not influenced by dung treatment (Table S2).

a) Marginal effect (overall)   b) Treatment effect at specified initial sizes and densities

c) Percent and absolute change derived from predicted RGR for specified sizes and densities

Figure 3: Dung treatment buffers larger saplings against the negative effects of sapling density on growth. a) Overall effect of dung treatment on $RGR_{diameter}$ (based on estimated marginal means or *emmeans* across densities, sizes and forest types), b) Predicted $RGR_{diameter}$ based on *emmeans* at specified densities for three discrete initial sizes (obtained using *emmeans*, raw data plotted in Fig S2), and b) The same effects expressed as percent change and absolute change in basal diameter, derived from the *emmeans* of $RGR_{diameter}$ at specified densities and initial sizes (see Supplement text A). Error bars represent 95% confidence intervals. The discrete value of initial diameter for small sapling is defined as mean – 0.5 SD instead of mean – 1SD which is non-existent in our dataset due to its right skew. Significance codes correspond to tests of treatment-control contrasts in the predicted RGR at specified density/size (based on *emmeans*): ns: p>0.05; *: p <0.05; **: p <0.01; ***: p<0.001.

472 **Table 2**: Conditional effects from a GLMM (with Student t error distribution, and variance

473 structure) showing final sapling basal diameter (mm) increasing with dung input, with this

474 effect being especially evident at high density and for larger saplings. Terms of interest are

475 marked blue and significant *P* values are marked bold.

Conditional effects	Estimate	S.E.	CI lower	CI upper	Pr(>|z|)
Intercept (ref: control, DDF, mean sapling size and density)	12.97	0.197	12.584	13.355	0.000
Treatment (Dung)	**0.55**	0.259	0.040	1.053	**0.035**
Sapling density	**-0.48**	0.188	-0.852	-0.116	**0.010**
Initial girth	**13.16**	0.287	12.602	13.725	**0.000**
Forest type – MDF	-0.18	0.136	-0.443	0.090	0.195
Forest type – Teak	**-0.72**	0.339	-1.386	-0.056	**0.034**
Treatment (Dung) × Sapling density	**0.63**	0.269	0.103	1.158	**0.019**
Treatment (Dung) × Initial girth	**0.76**	0.386	0.002	1.513	**0.049**
Sapling density × Initial girth	-0.53	0.275	-1.070	0.006	0.053
Treatment (Dung) × Sapling density × Initial girth	0.66	0.397	-0.118	1.439	0.096

Figure 4. a & c. Estimated marginal means of final sapling diameter and final height for

saplings under dung treatment and control (across the distribution of sizes, densities and

forest types). **b & d.** Estimated marginal means at specified densities for discrete initial sizes,

under dung treatment and control. Predicted values are shown for three discrete initial sizes.

In panel b, small sapling diameter is specified as mean – 0.5 SD initial diameter instead of

mean – 1SD which is non-existent in our dataset due to its right skew.

*1b) Sapling mortality and change in stem-count***:** Mortality during the ~6-month long field

experiment was low in the full dataset, with mortality rate being 3.96% under the control

(8/202) and 2.61% under the dung treatment (6/230), although a binomial GLMM revealed

that these differences were not significant (P=0.43). We also inspected the net change in

stem-count for the filtered dataset containing only the saplings alive at the end of the

experiment (N=418). We found that a net increase in stem-count was more common in

saplings receiving the dung treatment (13.4%) than control saplings (8.8%), whereas a net

loss of stem-count was marginally more common in control saplings (14.4% vs 12.1% for

dung treatment, Figure S4, Supplement text A).

We selected the 76% saplings whose stem-count remained the same between the initial and

final measurement (N=316), and repeated our analyses of the effects of dung treatment on

RGR$_{diameter}$ and final basal diameter for this subset, with models specified in Table 1. In this

smaller dataset, while the directions of effects were broadly consistent with the findings in

the full dataset, many effects were statistically not significant (Supplement Table S3 and

Table S4). Nevertheless, for both RGR$_{diameter}$ and final diameter, we found clear statistical

support for dung input providing a buffer against the negative effects of density for average-

sized saplings (Fig. S3).

2. Ex-situ mesocosm experiment: *Non-N-fixers outcompete N-fixers under dung input*

Both analyses of final sapling biomass and relative competitive strength (RCS) suggested that

elephant dung affected competition between plant functional types (inter-PFT competition),

supporting *Prediction 3*. When in inter-PFT competition, there was no difference in the

biomass of N-fixers and non-N-fixers under control conditions ($P = 0.178$ for *emmeans*

contrast, Figure 5a, Table S6b). In comparison, dung addition resulted in non-nitrogen fixers

accumulating over twice the biomass as nitrogen-fixers (*emmeans* contrast: $P=0.003$).

GLMM of relative competitive strength (RCS) revealed that the dung treatment substantially

increased the RCS of non-N-fixers (i.e., improved performance under inter-PFT vis-à-vis

intraspecific competition) relative to control conditions, after accounting for initial size and

random effects (Figure 5b, Table 3b). Dung treatment appeared to reduce the RCS of N-

fixers, but not significantly so. Next, we inspected the 95% confidence intervals to test for

differences in the competitor status of the two PFTs with respect to the null expectation of

1.0. In the absence of dung input, N-fixers performed better under inter-PFT competition than

under intra-specific competition (mean RCS= 1.762, $P=0.009$, null RCS=1.0) (Figure 5b,

Table 3a). In contrast, this better performance of N-fixers under inter-PFT competition did

not hold when dung was added (RCS did not differ from 1.0). Contrastingly, the non-N-fixers

showed better performance under inter-PFT competition under the dung treatment (mean

RCS =1.649, $P=0.013$), which was not the case under control conditions.

Figure 5: Outcomes of competition between the plant functional types N-fixers and non-N-

fixers in the mesocosm experiment. **a)** Predicted total sapling biomass under competition

between plant functional types (based on *ggpredict*): when supplied with dung, non-N-fixers

accumulate greater biomass than N-fixers under mixture competition. **b)** Predicted relative

competitive strength (RCS, based on *emmeans*): RCS is significantly >1.0 for N-fixers under

control and for non-N-fixers under dung treatment (raw data plotted in Fig. S4). RCS =

biomass under inter-PFT competition / intraspecific competition. Error bars represent 95%

confidence intervals around estimated means.

Table 3: Results from the GLMM of relative competitive strength (RCS). **a)** Estimated mean

RCS: N-fixers were more competitive (RCS>1.0) under while non-N-fixers were more

competitive under dung treatment, **b)** Conditional effects form GLMM: non-N-fixers

increased their RCS under dung treatment relative to control (rows of interest are marked in

blue). Rows with P<0.05 are marked in bold.

a) Difference between estimated marginal means of RCS and its null expected value

Treatment: PFT	*emmean* (log scale)	SE (log scale)	z	*P*	RCS (back-transformed)	CI lower	CI upper	Null expectation
Control: N-fixer	0.566	0.216	2.627	**0.009**	1.762	**1.155**	**2.688**	1
Dung: N-fixer	0.277	0.220	1.263	0.206	1.320	0.858	2.029	1
Control: Non-N-fixer	0.200	0.205	0.977	0.329	1.221	0.818	1.824	1
Dung: Non-N-fixer	0.500	0.201	2.491	**0.013**	1.649	**1.113**	**2.444**	1

b) Conditional effects from GLMM

	Estimate (log scale)	SE (log scale)	z	P	RCS (back-transformed)	CI lower	CI upper
Intercept (ref: Control, N-fixers)	0.566	0.216	2.627	**0.009**	1.762	1.155	2.688
Standardized initial size ratio	0.209	0.093	2.260	**0.024**	1.233	1.028	1.478
Treatment (Dung)	-0.289	0.260	-1.111	0.267	0.749	0.450	1.247
PFT (Non-N-fixer)	-0.366	0.254	-1.444	0.149	0.693	0.422	1.140
Treatment (Dung) × PFT (non-N-fixer)	0.589	0.285	2.064	**0.039**	**1.802**	**1.030**	**3.154**

3. The overall magnitude of nitrogen redistribution by elephants

We estimated that elephants annually produce 11038 such nutrient-rich patches/ km^2 on

average in these tropical forests, amounting to an annual cycling of 218.88 kg N/km^2 through

this pathway alone (Table S6). On a per capita basis, each adult elephant redistributes

~130.29 kg N/year throughout its home range by creating 6570 dung sites/year (Table S6).

Elephant dung piles are fertility hotspots as their nitrogen content (annual average = 0.971%)

is substantially higher than the upper soil layers in this landscape's deciduous forests (0.14 to

0.31%; Mani et al., 2018, 0.40% in Swamy and Proctor, 1994; 0.26% in Ramachandra et al.,

2012).

DISCUSSION

Due to their exceptional body size and wide-ranging behaviours, megaherbivores are expected to have a profound role in ecosystem functioning through both top-down and bottom-up impacts. Our experiments offer important insights into their bottom-up effects via large-scale redistribution of nutrients and its associated impacts on woody plant communities in mesic ecosystems. We found that elephant dung deposition sites 1) act as hotspots of plant growth (Fig 3a, 4a), 2) weaken the negative effects of high sapling density on average-sized and large saplings (Fig 3c, 4b), and 3) shift the competitive balance between plant functional types differing in nutrient limitation, by favoring non-N-fixers over N-fixers (Fig 5). Importantly, 4) the magnitude of such effects is substantial; we estimate that elephants create >11000 nutrient-rich sites annually in these mesic tropical forests (Table S7). These findings highlight the role of megaherbivores as modifiers of species interactions key to plant community assembly, and as promoters of fine-scale vegetation heterogeneity across their vast home ranges.

Elephant dung deposition creates hotspots of plant growth

We found that the large inputs of nitrogen (~19.83g N, Table S7) make elephant dung deposition sites hotspots for plant growth. Elephant dung strongly enhanced growth in the diameter of woody saplings compared to adjacent control plots. However, this growth-enhancing effect was not statistically significant for plant height (P=0.063), possibly because height responses may take longer to manifest than diameter (eg. Noyer et al., 2019; Seymour et al., 2022), potentially reflecting preferential allocation to bark thickening before height as a strategy to reduce mortality (eg. from fire Lawes et al., 2011). Such potentially delayed effects on height could explain why the only previous experiment of this nature could not

detect growth-enhancing effects of elephant dung in central Africa's rainforests (Kalbitzer et

al., 2019). Kalbitzer et al. (2019) analyzed height, and not diameter, and did not find growth-

enhancing effects of elephant dung one year after experimental dung input. However, shade-

tolerant saplings in their study enhanced leaf production, suggesting that the allocation of this

productivity to increases in shoot length may be delayed but potentially detectable with

longer monitoring. Our inspection of stem-count data suggests that the observed increase in

diameter partly ensued from a net increase in stem-count in saplings receiving elephant dung

input (Fig S3). Even for saplings whose stem-count remained unchanged, basal diameter

appears to have grown under elephant dung treatment, although such effects were not

statistically significant in this down-sized dataset. Furthermore, sapling survival appears to be

enhanced by dung treatment (2.61% mortality vs. 3.96% mortality in control saplings) in our

dataset, although this difference was not statistically significant, whereas Kalbitzer et al.

(2019) reported significantly lower mortality under dung treatment.

Elephants modify ecological interactions key to plant community assembly

We found that nutrient inputs via dung can locally alter plant competition in two ways. First,

elephant dung can buffer saplings against neighborhood competition or negative density-

dependence. This was especially evident in the case of average-sized and large saplings

whose growth under control conditions declined sharply with increasing sapling density (Fig.

3-4). In contrast, dung addition was associated with a near consistent ecological release from

neighborhood competition for all sapling sizes (see flattish slopes in Fig 3, Fig 4b). Since

larger saplings have greater nutrient requirements, their dung-mediated release from

neighborhood competition suggests that nutrient inputs from dung can be substantial. In

contrast, dung input did not influence the effects of density on sapling height; increased

vertical growth was only observed in response to increased sapling density, which is a well-

known response to crowding, possibly mediated by light competition (Gruntman et al., 2017).

Second, our mesocosm experiment revealed that dung can modify the outcome of

interference competition between plant functional types differing in nutrient limitation. Non-

N-fixers accumulated more biomass and outcompeted N-fixers when amended with dung but

not otherwise. That elephant dung favored non-N-fixers was also clearly evident in the

analyses of relative competitive strength (RCS). The otherwise better performance of N-

fixers in control conditions (RCS>1.0) did not hold in the presence of dung. Dung input

implied net gains for non-N-fixers in two ways: by enhancing the competitive status of non-

N-fixers (RCS > 1.0) and by lowering it for N-fixers which had high RCS in control

conditions. Herbivore dung has been previously shown to alter the competitive balance

between grasses and woody N-fixers in African savannas (Sitters & Olde Venterink, 2021).

Taken together, these findings suggest that such bottom-up effects of herbivores can be

critical in regulating the abundance of different plant types, including woody N-fixers, non-

fixers and grasses.

Dung-mediated differences in local competitive outcomes can determine which saplings

succeed in transitioning to the recruitment stage, thereby influencing community

composition. Our findings indicate that dung deposition can locally accelerate competitive

exclusion along at least two axes: 1) size and 2) nutrient limitation among plant functional

types (PFTs). First, although larger saplings were negatively affected by neighborhood

density under control conditions, they were buffered against neighborhood competition in the

presence of dung (Fig. 3, Fig. 4). This similarity in RGR between large and small saplings,

especially at high densities (Fig. S2), suggests that herbivores can reinforce or even amplify

existing size differences through their dung inputs (Fig. 4). Such dung-mediated ecological

release from competition can reinforce the beginners' advantage for larger saplings (for

example, taller understory plants have an advantage in light competition), thus potentially

excluding smaller saplings. Second, along the nutrient-limitation axis, herbivore dung

addition is likely to favor nutrient-limited plant species. In our mesocosm experiment,

nitrogen-rich dung shifted the competitive balance between non-N-fixers and N-fixers (Fig

4), suggesting that dung deposition could bring about fine-scale spatial differences in the

relative abundances of these two plant types by favoring non-N-fixers. It must be noted that

the observed effects on inter-PFT competition in our mesocosm experiment may be an

underestimate as we used only <10% (500g) of the amount of an average elephant defecation

(6.88 kg) in our treatments. Importantly, such effects on the coexistence among plant species

may be widespread along other axes of nutrient limitation, for example, fast- vs- slow-

growing and shade-tolerant vs. light-demanding species. Localized competitive exclusion

along these axes could thus change plant communities in multiple ways. Such dung-mediated

gains by large, nutrient-limited plants at the expense of smaller, non-nutrient-limited saplings

can potentially contribute to the decline in tree community richness often reported in

herbivore presence (Jia et al., 2018).

Conservation implications

Our study has implications for understanding how defaunation may impact ecosystem

functioning through effects on species interactions shaping plant communities. Poulsen et al.

(2018) speculated that the loss of nutrient cycling associated with megaherbivores would

result in the loss of light-demanding species as they are more nutrient-limited. While we did

not focus on the light-demanding/shade-tolerant distinction, elephant dung indeed favored

nutrient-limited plant types in our experiment (non-N-fixers) and thus elephant presence can

substantially shift the composition of plant communities, possibly even at the landscape scale

through cumulative generation of numerous temporary growth hotspots over multiple years.

Nevertheless, we note the contradictory earlier findings of Kalbitzer et al. (2019) where

elephant dung had small positive effects on shade-tolerant species that are less nutrient

limited, although these results were based only on leaf production. In addition to the effects at

the landscape level, we propose that enhanced fine-scale heterogeneity in community

composition is another key consequence of such nutrient redistribution by megaherbivores.

The large-scale annual production of nutrient hotspots by elephants (totally >11000/km^2/year

in Nagarahole) likely amplifies vegetation heterogeneity through fine-scale differences in

growth and competition, which is in line with earlier findings that herbivore dung deposition

increases soil heterogeneity (Afzal & Adams, 1992; Ferraro & Lienau, 2025; Hobbs, 1996).

Importantly, the functional importance of megaherbivores is underscored by the fact that the

amount of dung input is a key modulator of vegetation responses (Valdés-Correcher et al.,

2019), and this input is exceptionally large in the case of megaherbivores. It is likely that

such oversized nutrient-mediated impacts could also have downstream effects on other

organisms. Thus, protecting megaherbivore populations is critical to conserving such

important functional roles.

Future directions

Our experiments open several possibilities to further understand bottom-up effects of large

herbivores. As these dung sites are temporary nutrient hotspots, they would only induce

short-term spatial differences in plant growth. To what extent these pulses of altered growth

trajectories shape the recruitment fate to mature stages can only be understood through

longer-term studies, or models that consider such pulsed nutrient effects on plant growth and

competition. Second, we encourage investigations on competition between plant functional

types along other axes of differentiation, including fast- vs slow-growth and shade-tolerant vs

light-demanding species. Third, stoichiometric ratios of different growth-limiting elements in

dung can be important for plant growth and interactions. Megaherbivores have high

phosphorous requirements and thus their faeces tend to be P-depleted i.e., with high N:P

ratios (Le Roux et al., 2020) that could favour non-N-fixers over N-fixers (Sitters & Olde

Venterink, 2021).

In conclusion, our study highlights large-scale nutrient redistribution by megaherbivores as

an underappreciated driver of plant growth and species interactions that generate fine-scale

vegetation heterogeneity. Such functional roles of megafauna remain understudied in the

mesic tropics (Hyvarinen et al., 2021), particularly in South and Southeast Asia (Sankaran &

Ahrestani, 2016). These regions have faced extensive defaunation and deforestation in both

modern and prehistoric times (Svenning et al., 2016). Localized restoration efforts in such

ecosystems could benefit from a mechanistic understanding of herbivore impacts (see

Geremia et al., 2025), including the relative impacts of top-down and bottom-up regulation.

Our findings suggest that large herbivore rewilding could benefit conservation actions by

restoring species interactions and fine-scale vegetation heterogeneity, although such

ecological roles should be seen more as an outcome of well-preserved ecosystems. We call

for further investigations to advance a more comprehensive understanding of herbivore

functions in these understudied ecosystems.

ACKNOWLEDGEMENTS

We are grateful to the Principal Chief Conservator of Forests, Karnataka Forest Department, the office of the Conservator of Forests and Director, Rajiv Gandhi (Nagarahole) National Park, Hunsur, for field research permits and other forest officials for facilitating our fieldwork. We are also grateful to the Manager, Jungle Lodges and Resorts, Kabini, for providing us the space to conduct the ex-situ experiment, and to the forestry nurseries in Tithimati and Beechanahalli for providing us seedlings. We especially thank Prof. T.N.C. Vidya and the Kabini Elephant Project for logistical support during fieldwork as well as for the collaboration under which the fieldwork was carried out. We are thankful to our field assistants Krishna, Pramod and Shankar for assisting with fieldwork and ensuring our safety. We thank Abhirami Ravichandran for help during nutrient analyses of fecal samples. We are thankful to the two reviewers for the very helpful critical feedback on the first draft and to Mayank Kohli for feedback on a pre-submission draft. HG analyzed and wrote this paper while working at the University of Turku.

FUNDING STATEMENT

The fieldwork and laboratory analyses for this study were funded by the National Postdoctoral Fellowship (PDF/2021/002320) granted to HG by the Science and Engineering Research Board, India. Institutional support and laboratory facilities for processing fecal and plant samples was provided by the National Centre for Biological Sciences.

REFERENCES

Afzal, M., & Adams, W. A. (1992). Heterogeneity of Soil Mineral Nitrogen in Pasture Grazed by Cattle. *Soil Science Society of America Journal*, *56*(4), 1160–1166. https://doi.org/10.2136/sssaj1992.03615995005600040026x

Archibald, S., & Hempson, G. P. (2016). Competing consumers: Contrasting the patterns and impacts of fire and mammalian herbivory in Africa. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *371*(1703), 20150309. https://doi.org/10.1098/rstb.2015.0309

Augustine, D. J., McNaughton, S. J., & Frank, D. A. (2003). Feedbacks between soil nutrients and large herbivores in a managed savanna ecosystem. *Ecological Applications*, *13*(5), 1325–1337. https://doi.org/10.1890/02-5283

Bates, D., Maechler, Martin, Bolker, B., Walker, S., & Christensen, R. H. B. (2015). Package 'lme4'. *Convergence*, *12*(1), 2.

Berzaghi, F., Bretagnolle, F., Durand-Bessart, C., & Blake, S. (2023). Megaherbivores modify forest structure and increase carbon stocks through multiple pathways. *Proceedings of the National Academy of Sciences*, *120*(5), e2201832120. https://doi.org/10.1073/pnas.2201832120

Berzaghi, F., Verbeeck, H., Nielsen, M. R., Doughty, C. E., Bretagnolle, F., Marchetti, M., & Scarascia-Mugnozza, G. (2018). Assessing the role of megafauna in tropical forest ecosystems and biogeochemical cycles – the potential of vegetation models. *Ecography*, *41*(12), 1934–1954. https://doi.org/10.1111/ecog.03309

Blake, S. (2002). *The Ecology of Forest Elephant Distribution and its implications for conservation*. University of Edinburgh.

Brooks, M., E., Kristensen, K., Benthem, K., J. ,van, Magnusson, A., Berg, C., W., Nielsen, A., Skaug, H., J., Mächler, M., & Bolker, B., M. (2017). glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. *The R Journal*, *9*(2), 378. https://doi.org/10.32614/RJ-2017-066

Campos-Arceiz, A., & Blake, S. (2011). Megagardeners of the forest – the role of elephants in seed dispersal. *Acta Oecologica*, *37*(6), 542–553. https://doi.org/10.1016/j.actao.2011.01.014

Coetsee, C., Wigley, B. J., Sankaran, M., Ratnam, J., & Augustine, D. J. (2023). Contrasting Effects of Grazing vs Browsing Herbivores Determine Changes in Soil Fertility in an East African Savanna. *Ecosystems*, *26*(1), 161–173. https://doi.org/10.1007/s10021-022-00748-7

Dirzo, R., Young, H. S., Galetti, M., Ceballos, G., Isaac, N. J. B., & Collen, B. (2014). Defaunation in the Anthropocene. *Science*, *345*(6195), 401–406.

Ferraro, K. M., Albrecht, D., Hendrix, J. G., Wal, E. V., Schmitz, O. J., Webber, Q. M. R., & Bradford, M. A. (2024). The biogeochemical boomerang: Site fidelity creates nutritional hotspots that may promote recurrent calving site reuse. *Ecology Letters*, *27*(8), e14491. https://doi.org/10.1111/ele.14491

Ferraro, K. M., & Lienau, J. R. (2025). Of all shapes and sizes: A theoretical framework for animal-mediated terrestrial heterogeneity across scales. *Ecography*, e08006. https://doi.org/10.1002/ecog.08006

Gautam, H., Arulmalar, E., Kulkarni, M. R., & Vidya, T. N. C. (2019). NDVI is not reliable as a surrogate of forage abundance for a large herbivore in tropical forest habitat. *Biotropica*, *51*(3), 443–456. https://doi.org/10.1111/btp.12651

Geremia, C., Hamilton, E. W., & Merkle, J. A. (2025). Yellowstone's free-moving large bison herds provide a glimpse of their past ecosystem function. *Science*, *389*(6763), 904–908.

Gruntman, M., Groß, D., Májeková, M., & Tielbörger, K. (2017). Decision-making in plants under competition. *Nature Communications*, *8*(1), 2235. https://doi.org/10.1038/s41467-017-02147-2

Guldemond, R. A. R., Purdon, A., & Van Aarde, R. J. (2017). A systematic review of elephant impact across Africa. *PLOS ONE*, *12*(6), e0178935. https://doi.org/10.1371/journal.pone.0178935

Hartig, F., & Hartig, M. F. (2017). *Package 'DHARMa' residual diagnostics for hierarchical (multi-level/mixed) regression models.* (Version 0.1.5) [Computer software].

Hobbs, N. T. (1996). Modification of Ecosystems by Ungulates. *The Journal of Wildlife Management*, *60*(4), 695. https://doi.org/10.2307/3802368

Hyvarinen, O., Te Beest, M., Le Roux, E., Kerley, G., De Groot, E., Vinita, R., & Cromsigt, J. P. G. M. (2021). Megaherbivore impacts on ecosystem and Earth system functioning: The current state of the science. *Ecography*, *44*(11), 1579–1594. https://doi.org/10.1111/ecog.05703

Jia, S., Wang, X., Yuan, Z., Lin, F., Ye, J., Hao, Z., & Luskin, M. S. (2018). Global signal of top-down control of terrestrial plant communities by herbivores. *Proceedings of the National Academy of Sciences*, *115*(24), 6237–6242. https://doi.org/10.1073/pnas.1707984115

Kalbitzer, U., McInnis, V., Omeja, P. A., Bortolamiol, S., & Chapman, C. A. (2019). Does the presence of elephant dung create hotspots of growth for existing seedlings? *Journal of Tropical Ecology*, *35*(03), 132–139. https://doi.org/10.1017/S0266467419000051

Karp, A. T., Koerner, S. E., Hempson, G. P., Abraham, J. O., Anderson, T. M., Bond, W. J., Burkepile, D. E., Fillion, E. N., Goheen, J. R., Guyton, J. A., Kartzinel, T. R., Kimuyu, D. M., Mohanbabu, N., Palmer, T. M., Porensky, L. M., Pringle, R. M., Ritchie, M. E., Smith, M. D., Thompson, D. I., … Staver, A. C. (2024). Grazing herbivores reduce herbaceous biomass and fire activity across African savannas. *Ecology Letters*, *27*(6), e14450. https://doi.org/10.1111/ele.14450

Knapp, A., Blair, J. M., Briggs, J. M., Collins, S. L., & Hartnett, D. C. (1999). The Keystone Role of Bison in North American Tallgrass PrairieBison increase habitat heterogeneity and alter a broad array of plant, community, and ecosystem processes. *BioScience*, *49*(1), 39–50. https://doi.org/10.1525/bisi.1999.49.1.39

Lawes, M. J., Adie, H., Russell-Smith, J., Murphy, B., & Midgley, J. J. (2011). How do small
savanna trees avoid stem mortality by fire? The roles of stem diameter, height and bark
thickness. *Ecosphere*, *2*(4), art42. https://doi.org/10.1890/ES10-00204.1

Le Roux, E., Van Veenhuisen, L. S., Kerley, G. I. H., & Cromsigt, J. P. G. M. (2020). Animal
body size distribution influences the ratios of nutrients supplied to plants. *Proceedings of the*
National Academy of Sciences, *117*(36), 22256–22263.
https://doi.org/10.1073/pnas.2003269117

Malhi, Y., Doughty, C. E., Galetti, M., Smith, F. A., Svenning, J.-C., & Terborgh, J. W.
(2016). Megafauna and ecosystem function from the Pleistocene to the Anthropocene.
Proceedings of the National Academy of Sciences, *113*(4), 838–846.
https://doi.org/10.1073/pnas.1502540113

Mani, S., Merino, A., García-Oliva, F., Riotte, J., & Sukumar, R. (2018). Soil properties and
organic matter quality in relation to climate and vegetation in southern Indian tropical
ecosystems. *Soil Research*, *56*(1), 80. https://doi.org/10.1071/SR16262

McCleery, R. A., Kruger, L., Monadjem, A., Tye, D., Mhlava, P., Zwane, E., Bijl, A., Jones,
M., Hartfelder, J., Coetsee, C., & Fletcher, R. (2024). *Megaherbivores Facilitate Large*
Grazing Herbivores and Suppress Small Mammals Through Vegetation Structure and
Cryptic Pathways. https://doi.org/10.2139/ssrn.5069873

Noyer, E., Ningre, F., Dlouhá, J., Fournier, M., & Collet, C. (2019). Time shifts in height and
diameter growth allocation in understory European beech (Fagus sylvatica L.) following
canopy release. *Trees*, *33*(2), 333–344. https://doi.org/10.1007/s00468-018-1779-8

Omeja, P. A., Jacob, A. L., Lawes, M. J., Lwanga, J. S., Rothman, J. M., Tumwesigye, C., &
Chapman, C. A. (2014). Changes in Elephant Abundance Affect Forest Composition or
Regeneration? *Biotropica*, *46*(6), 704–711. https://doi.org/10.1111/btp.12154

Owen-Smith, R. N. (1988). Megaherbivores: The Influence of Very Large Body Size on
Ecology. *Cambridge University Press, Cambridge Studies in Ecology*.

Pastor, J., Cohen, Y., & Hobbs, N. T. (2006). The role of large herbivores in ecosystem
nutrient cycles. In *Large herbivore ecology, ecosystem dynamics and conservation* (pp. 289–
319). Cambridge University Press.

Poulsen, J. R., Rosin, C., Meier, A., Mills, E., Nuñez, C. L., Koerner, S. E., Blanchard, E.,
Callejas, J., Moore, S., & Sowers, M. (2018). Ecological consequences of forest elephant
declines for Afrotropical forests. *Conservation Biology*, *32*(3), 559–567.
https://doi.org/10.1111/cobi.13035

Pringle, R. M., Abraham, J. O., Anderson, T. M., Coverdale, T. C., Davies, A. B., Dutton, C.
L., Gaylard, A., Goheen, J. R., Holdo, R. M., Hutchinson, M. C., Kimuyu, D. M., Long, R.
A., Subalusky, A. L., & Veldhuis, M. P. (2023). Impacts of large herbivores on terrestrial
ecosystems. *Current Biology*, *33*(11), R584–R610. https://doi.org/10.1016/j.cub.2023.04.024

Ramachandra, T. V., Chandran M.D., S., Joshi, D., & Kumar, M. (2012). *Soil Quality across*
Diverse Landscapes in Central Western Ghats, India (Sahyadri Conservation Series 16,
ENVIS Technical Report 42, p. 72). Indian Institute of Science.

Reddy, G. V., Karanth, K. U., Kumar, N. S., Krishnaswamy, J., & Karanth, K. K. (2016).
Recovering Biodiversity in Indian Forests. Springer Singapore. https://doi.org/10.1007/978-
981-10-0911-2

Ripple, W. J., Newsome, T. M., Wolf, C., Dirzo, R., Everatt, K. T., Galetti, M., Hayward, M.
W., Kerley, G. I. H., Levi, T., Lindsey, P. A., Macdonald, D. W., Malhi, Y., Painter, L. E.,
Sandom, C. J., Terborgh, J., & Van Valkenburgh, B. (2015). Collapse of the world's largest
herbivores. *Science Advances*, *1*(4), e1400103. https://doi.org/10.1126/sciadv.1400103

Sandhage-Hofmann, A., Linstädter, A., Kindermann, L., Angombe, S., & Amelung, W.
(2021). Conservation with elevated elephant densities sequesters carbon in soils despite
losses of woody biomass. *Global Change Biology*, *27*(19), 4601–4614.
https://doi.org/10.1111/gcb.15779

Sankaran, M., & Ahrestani, F. (2016). The Ecology of Large Herbivores of South and
Southeast Asia: Synthesis and Future Directions. In *The Ecology of Large Herbivores in*
South and Southeast Asia. Springer.

Sekar, N., Lee, C., & Sukumar, R. (2017). Functional nonredundancy of elephants in a
disturbed tropical forest. *Conservation Biology*, *31*(5), 1152–1162.
https://doi.org/10.1111/cobi.12907

Seymour, C. L., Joseph, G. S., Calitz, W., Henschel, J. R., Ramaswiela, T., & Van Der
Merwe, H. (2022). Mean height increase in saplings of a keystone woody savanna species
over 15 years similar to that over a single season. *Ecosphere*, *13*(9), e4173.
https://doi.org/10.1002/ecs2.4173

Sitters, J., & Olde Venterink, H. (2018). A stoichiometric perspective of the effect of
herbivore dung on ecosystem functioning. *Ecology and Evolution*, *8*(2), 1043–1046.
https://doi.org/10.1002/ece3.3666

Sitters, J., & Olde Venterink, H. (2021). Herbivore dung stoichiometry drives competition
between savanna trees and grasses. *Journal of Ecology*, *109*(5), 2095–2106.
https://doi.org/10.1111/1365-2745.13623

Sivasubramaniyan, G., & Sivaganesan, N. (1996). Role of elephants in Sujalkuttai-Bannari
Corridor in Sathyamangalam Forest Division, Tamil Nadu, southern India. *Gajah*, *15*, 9–27.

Sukumar, R. (2003). *The living elephants: Evolutionary ecology, behavior, and conservation*.
Oxford Univ. Press.

Svenning, J.-C., Pedersen, P. B. M., Donlan, C. J., Ejrnæs, R., Faurby, S., Galetti, M., Hanse,
D. M., Sandel, B., Sandom, C. J., Terborgh, J., & Vera, F. W. M. (2016). Science for a wilder
Anthropocene: Synthesis and future directions for trophic rewilding research. *Proceedings of*
the National Academy of Sciences, *113*(4), 898–906.

Tandalla, D. C., Homeier, J., & Batáry, P. (2024). Responses of Tropical Tree Seedlings to
Nutrient Addition: A Meta-analysis to understand future changes in Tropical Forest
Dynamics. *Current Forestry Reports*, *11*(1), 3. https://doi.org/10.1007/s40725-024-00240-6

Trepel, J., Le Roux, E., Abraham, A. J., Buitenwerf, R., Kamp, J., Kristensen, J. A., Tietje,
M., Lundgren, E. J., & Svenning, J.-C. (2024). Meta-analysis shows that wild large

herbivores shape ecosystem properties and promote spatial heterogeneity. *Nature Ecology & Evolution*, *8*(4), 705–716. https://doi.org/10.1038/s41559-024-02327-6

Valdés-Correcher, E., Sitters, J., Wassen, M., Brion, N., & Olde Venterink, H. (2019). Herbivore dung quality affects plant community diversity. *Scientific Reports*, *9*(1), 5675. https://doi.org/10.1038/s41598-019-42249-z

Van Der Plas, F., Howison, R. A., Mpanza, N., Cromsigt, J. P. G. M., & Olff, H. (2016). Different-sized grazers have distinctive effects on plant functional composition of an A frican savannah. *Journal of Ecology*, *104*(3), 864–875. https://doi.org/10.1111/1365-2745.12549

Van der Waal, C., De Kroon, H., De Boer, W. F., Heitkönig, I. M. A., Skidmore, A. K., De Knegt, H. J., Van Langevelde, F., Van Wieren, S. E., Grant, R. C., Page, B. R., Slotow, R., Kohi, E. M., Mwakiwa, E., & Prins, H. H. T. (2009). Water and nutrients alter herbaceous competitive effects on tree seedlings in a semi-arid savanna. *Journal of Ecology*, *97*(3), 430–439. https://doi.org/10.1111/j.1365-2745.2009.01498.x

Varma, V., Catherin, A. M., & Sankaran, M. (2018). Effects of increased N and P availability on biomass allocation and root carbohydrate reserves differ between N-fixing and non-N-fixing savanna tree seedlings. *Ecology and Evolution*, *8*, 8467–8476.

Villar, N., Paz, C., Zipparro, V., Nazareth, S., Bulascoschi, L., Bakker, E. S., & Galetti, M. (2021). Frugivory underpins the nitrogen cycle. *Functional Ecology*, *35*(2), 357–368. https://doi.org/10.1111/1365-2435.13707

Vitousek, PeterM., & Howarth, RobertW. (1991). Nitrogen limitation on land and in the sea: How can it occur? *Biogeochemistry*, *13*(2). https://doi.org/10.1007/BF00002772

Wolf, A., Doughty, C. E., & Malhi, Y. (2013). Lateral Diffusion of Nutrients by Mammalian Herbivores in Terrestrial Ecosystems. *PLoS ONE*, *8*(8), e71352. https://doi.org/10.1371/journal.pone.0071352

SUPPLEMENTARY INFORMATION

Figure S1: a) Representative arrangement of the sampling regime in the field experiment on

sapling response to elephant dung treatment. b) Raw plot-wise means of initial basal diameter

of saplings in the plot pairs (averaged across all individuals within each plot), c) Raw plot-

wise means of observed percentage change in the basal diameter of saplings in 36 plot-pairs

during the course of the field experiment (i.e., change between initial and final measurement).

An increased growth in response to dung treatment is evident in the majority of plot pairs,

relative to control plots 10m away.

Figure S2. Size-dependent effect of neighborhood competition on sapling growth

($RGR_{diameter}$) under dung treatment and control conditions, with raw data points shown. 17

data points were omitted due to y-axis truncation for better visualization. The buffering

provided by dung input against negative density dependence for larger saplings is evident in

contrast with control saplings. Estimates derived from the GLMM using *ggpredict*. Small

sapling diameter is specified as mean – 0.5 SD initial diameter instead of mean – 1SD which

is non-existent in our dataset due to its right skew.

a) Loss/gain in stemcount
during the experiment

b) Marginal effects (overall)

c) Conditional effects on RGRdiameter (in dry deciduous forest)

d) Conditional effects on final diameter (in dry deciduous forest)

Figure S3: a) Net change in the stem-count of individuals, b) marginal effects and c)

conditional effects of dung treatment on $RGR_{diameter}$, for a subset of 316 individuals that did

not show any change in stem-count. While the overall growth enhancing effect of dung was

statistically not significant (P=0.063), the conditional effects (for the deciduous forest which

dominated the sample) show that dung treatment significantly buffered the saplings against

the negative density-dependent effectsstem-count. This buffering was statistically significant

for average-sized saplings (density x treatment interaction) but not larger saplings. d) Similar

trends were observed for final basal diameter. See Supplement text A for more details.

Figure S4: Predicted relative competitive strength (RCS, based on *emmeans*) for N-fixers

and non-N-fixers under dung treatment and control in the mesocosm experiment: RCS is

significantly >1.0 for N-fixers under control and for non-N-fixers under dung treatment. RCS

= biomass under inter-PFT competition / intraspecific competition. Error bars represent 95%

confidence intervals around estimated marginal means. This figure is an extension of Figure

4b in the main text, with only raw data points added. One data point for N-fixers has been

omitted from control group for better visualization.

948 **Table S1**: Summary table from a GLMM (with variance structure) of $RGR_{diameter}$

949 (mm/mm/day) showing conditional effects in dry deciduous forest. Terms of interest are

950 marked blue and significant *P* values are marked bold.

Conditional effects	Estimate	S.E.	CI lower	CI upper	Pr(>|z|)
Intercept (ref: control, DDF, mean sapling size and density)	0.00047	0.00010	0.00027	0.00067	**0.000**
Treatment (Dung)	0.00025	0.00012	0.00001	0.00049	**0.040**
Sapling density	-0.00018	0.00010	-0.00037	0.00002	0.072
Initial diameter	-0.00036	0.00012	-0.00060	-0.00013	**0.003**
Forest type – MDF	-0.00006	0.00015	-0.00035	0.00023	0.677
Forest type – Teak	-0.00031	0.00021	-0.00072	0.00010	0.143
Treatment (Dung) × Sapling density	0.00021	0.00014	-0.00006	0.00048	0.120
Treatment (Dung) × Initial diameter	0.00017	0.00015	-0.00013	0.00047	0.267
Sapling density × Initial diameter	-0.00029	0.00014	-0.00056	-0.00002	**0.033**
Treatment (Dung) × Sapling density × Initial diameter	0.00028	0.00018	-0.00008	0.00064	0.123

 Table S2: Conditional effects from GLMM showing final height being not significantly

 increased by dung input. Terms of interest are marked blue and significant effects marked

 bold (based on 95% CIs).

Conditional effects	Estimate	S.E.	CI lower	CI upper	Pr(>|z|)
Intercept (ref: control, DDF, mean sapling height and density)	49.58	0.979	47.666	51.502	**0.000**
Treatment (Dung)	2.07	1.197	-0.272	4.420	0.083
Sapling density	1.75	0.840	0.104	3.398	**0.037**
poly (Initial height) 1	21.23	0.932	19.408	23.062	**0.000**
poly (Initial height) 2	-0.78	0.788	-2.319	0.768	0.325
Forest type – MDF	0.99	0.813	-0.601	2.588	0.222
Forest type – Teak	1.84	1.565	-1.230	4.905	0.240
Treatment (Dung) × Sapling density	-1.29	1.249	-3.736	1.160	0.302
Treatment (Dung) × poly (Initial height) 1	0.80	1.214	-1.576	3.184	0.508
Treatment (Dung) × poly (Initial height) 2	-3.24	1.264	-5.717	-0.764	**0.010**
Sapling density × poly (Initial height) 1	1.01	0.767	-0.495	2.511	0.189
Sapling density × poly (Initial height) 2	-1.77	0.689	-3.124	-0.422	**0.010**
Treatment (Dung) × Sapling density × poly (Initial height) 1	0.11	1.159	-2.163	2.379	0.926

Treatment (Dung) × Sapling density × poly (Initial height) 2	1.41	1.351	-1.242	4.055	0.298

Table S3: Results based on the final sapling basal diameter (mm) of the subset of 316

saplings whose stem-count remained the same. Conditional effects from a GLMM (Student t

family of error distribution, with variance structure: dispersion ~ std. initial diameter)

showing dung input buffers saplings of average size against the negative effects of density,

for saplings in the dry deciduous forest (which dominated the sample). The marginal positive

effect of dung treatment is not significant (P=0.157, see Fig S3). Terms of interest are

marked blue and significant P values are marked bold.

Conditional effects	Estimate	S.E.	CI lower	CI upper	Pr(>|z|)
Intercept (ref: control, DDF, mean sapling size and density)	13.34	0.163	13.020	13.658	0.000
Treatment (Dung)	0.30	0.205	-0.100	0.704	0.140
Sapling density	-0.37	0.137	-0.637	-0.100	**0.007**
Initial girth	13.76	0.248	13.272	14.246	**0.000**
Forest type – MDF	-0.12	0.115	-0.351	0.101	0.280
Forest type – Teak	-0.83	0.238	-1.297	-0.365	0.000
Treatment (Dung) × Sapling density	0.42	0.206	0.013	0.822	**0.043**
Treatment (Dung) × Initial girth	0.30	0.332	-0.349	0.955	0.362
Sapling density × Initial girth	-0.38	0.224	-0.816	0.061	0.091
Treatment (Dung) × Sapling density × Initial girth	0.36	0.334	-0.295	1.015	0.282

981 **Table S4**: Results based on RGR$_{diameter}$ of the subset of 316 saplings whose stem-count

982 remained the same. Conditional effects from a GLMM (Student t family of error distribution)

983 showing dung input buffers saplings of average size against the negative effects of density,

984 for saplings in the dry deciduous forest (which dominated the sample). The marginal positive

985 effect of dung treatment is not significant (based on *emmeans*, *P*=0.184, see Fig S3). Terms

986 of interest are marked blue and significant *P* values are marked bold.

Conditional effects	Estimate	S.E.	CI lower	CI upper	Pr(>|z|)
Intercept (ref: control, DDF, mean sapling size and density)	0.00060	0.00010	0.00040	0.00080	**0.000**
Treatment (Dung)	0.00016	0.00011	-0.00006	0.00037	0.150
Sapling density	-0.00024	0.00009	-0.00042	-0.00006	**0.009**
Initial girth	-0.00032	0.00012	-0.00055	-0.00008	**0.008**
Forest type – MDF	-0.00009	0.00013	-0.00035	0.00017	0.502
Forest type – Teak	-0.00055	0.00022	-0.00098	-0.00013	**0.011**
Treatment (Dung) × Sapling density	0.00029	0.00015	0.00000	0.00058	**0.047**
Treatment (Dung) × Initial girth	0.00002	0.00016	-0.00029	0.00033	0.888
Sapling density × Initial girth	-0.00023	0.00012	-0.00046	0.00001	0.058
Treatment (Dung) × Sapling density × Initial girth	0.00014	0.00019	-0.00023	0.00051	0.461

989 **Table S5:** Conditional effects from the GLMM of total sapling biomass under control and

990 dung treatment, and under monoculture and mixture (i.e., inter-PFT) competition in the

991 mesocosm experiment. Terms of *a priori* interest are marked blue and effects different from 0

992 are marked bold.

	Estimate	SE	Lower CI	Upper CI	
Intercept (reference: control, N-fixer, monoculture)	7.614	1.604	4.470	10.758	**0.000**
Treatment (Dung)	-0.325	0.513	-1.331	0.680	0.526
Competition type (Mix)	0.434	0.751	-1.038	1.906	0.564
PFT (Non-N-fixer)	5.420	2.703	0.121	10.718	**0.045**
Initial size index (standardized)	2.177	0.492	1.213	3.142	**0.000**
Treatment (Dung) × Competition type (Mix)	-0.075	1.164	-2.358	2.207	0.948
Treatment (Dung) × PFT (Non-N-fixer)	0.515	2.630	-4.639	5.670	0.845
PFT (Non-N-fixer) × Competition type (Mix)	-0.592	3.455	-7.364	6.180	0.864
Treatment (Dung) × Competition type (Mix) × PFT (Non-N-fixer)	9.225	5.876	-2.291	20.741	0.116

994　**Table S6**: a) Estimated marginal mean sapling biomass (g) of all groups in the mesocosm

995　experiment. b) Pair-wise comparisons for contrasts between plant-functional types within

996　each competition type. Groups/contrasts of interest are marked with blue text.

a) Estimated marginal means of groups

Competition type	PFT	Treatment	*emmean*	SE	CI (lower)	CI (upper)
Monoculture	N-fixer	Control	7.61	1.60	4.47	10.76
Mixture	N-fixer	Control	8.05	1.69	4.74	11.36
Monoculture	Non-N-fixer	Control	13.03	2.27	8.59	17.48
Mixture	Non-N-fixer	Control	12.87	3.23	6.54	19.21
Monoculture	N-fixer	Dung	7.29	1.59	4.17	10.40
Mixture	N-fixer	Dung	7.65	1.76	4.19	11.10
Monoculture	Non-N-fixer	Dung	13.22	2.46	8.40	18.05
Mixture	Non-N-fixer	Dung	22.21	4.55	13.30	31.13

b) Pair-wise contrasts

	Competition type	Treatment	Difference (based on *emmeans*)	SE	*z* ratio	*P*
(N-fixer) - (Non-N-fixer)	Monoculture	Control	-5.420	2.703	-2.005	0.045
(N-fixer) - (Non-N-fixer)	Mixture	Control	-4.827	3.588	-1.345	0.178
(N-fixer) - (Non-N-fixer)	Monoculture	Dung	-5.935	2.864	-2.073	0.038
(N-fixer) - (Non-N-fixer)	Mixture	Dung	-14.568	4.863	-2.996	0.003

1000

1001

1002

1003

Table S7: Quantification of annual nitrogen redistribution via dung deposition by elephants

in Nagarahole National Park, southern India. N% was averaged from N% quantified at every

3 months: October 2022 (1.151%), January 2023 (1.027%), April 2023 (0.738) based on

Gautam et al. 2024, and August 2023 (0.967%) quantified in this study. Estimates of elephant

density (~1.68/km^2) in Nagarahole were obtained from the Karnataka Elephant Survey 2010

(Baskaran and Sukumar 2011[1]). We used 18/day as the daily defecation rate as reported for

Mudumalai in this landscape (Santosh and Sukumar 1995[2]) as well as another study with a

large sample size (Hedges 2012 pp. 82[3]).

Variable	Mean (SD, N)	Source
Nitrogen % in dung	0.9707 (4 seasons, 113 samples)	This study and *Reference removed for double-blind review*
Wet weight of a dung pile (kg)	6.881 (2.333, 42)	This study
Proportion of dry mass in dung	0.297 (0.031, 29)	This study
Defecation rate (day^{-1})	18	Santosh and Sukumar 1995, Hedges 2012, pp.82
Elephant density (km^{-2})	1.68	Baskaran and Sukumar 2011
Total annual dung deposition sites (km^{-2})	11038	-
N deposited at each defecation site	19.831g	This study
Total annual N deposition (kg km^{-2})	218.882	This study

Per capita annual N deposition	130.287	This study
(throughout the range)		

[1]Baskaran, N., Sukumar, R., 2011. Karnataka Elephant Census 2010. Technical Report submitted

to Karnataka Forest Department – March 2011. Asian Nature Conservation Foundation and Centre

for Ecological Sciences, Bangalore, Karnataka.

[2]Santosh, J.A., Sukumar, R., 1995. Some solutions to estimation of elephant densities. In: Daniel,

J.C., Datye, H.S. (Eds.), A Week with Elephants. Bombay Natural History Society and Oxford

University Press, Bombay, pp. 394–404.

[3]Hedges, S. (2012). Estimating absolute density from dung pile density. *Monitoring elephant*

populations and assessing threats, 61.

1020

1021

1022

1023

1024

1025

1026

1027

1028

1029

 Supplement text: A

 Using predicted RGR$_{diameter}$ to derive percentage and absolute change in diameter

 As RGR takes very small values and thus may appear less intuitive, we used RGR$_{diameter}$ to

 derive two intuitive variables, i.e., percentage change and absolute change in diameter, for

 different contrasts of treatment, sapling size and density. This conversion is based on

 following logic.

 a) Deriving absolute change in diameter from predicted RGR$_{diameter}$

 The RGR's formula, RGR$_{diameter}$ = [ln (final diameter) – ln (initial diameter)]/ t

 can be rearranged as:

 ln (final diameter) = ln (initial diameter) + RGR$_{diameter}$ × t,

 Both sides can be exponentiated to get:

 Final diameter = Initial diameter × e$^{RGRdiameter \times t}$

 Absolute change in diameter = Final diameter - Initial diameter,

 which can be derived from above as

 Absolute change in diameter = Initial diameter × (e$^{RGRdiameter \times t}$ – 1)

 We used a constant $t = 215$ days for all size contrasts as it represented the full duration of our

 field experiment.

b) Deriving percentage change in diameter from predicted RGR$_{diameter}$

From the rearrangement,

Final diameter = Initial diameter $\times$ e$^{RGRdiameter \times t}$,

we can derive relative change as:

(Final diameter - Initial diameter)/ Initial diameter = (e$^{RGRdiameter \times t}$ – 1),

And thus, percentage change in diameter = $100 \times$ (e$^{RGRdiameter \times t}$ – 1).

1055

1056 We present in Fig 3 in main text these two derived measures for three discrete densities:

1057 mean density, mean – 1SD and mean + 1SD.

1058

1059

<u>*Effect of dung treatment on net loss and gain in stem-count*</u>

We also investigated if the dung-mediated changes in the total diameter and RGR$_{diameter}$, as

presented in the main text, came largely from the changes in stem-count of individuals. Such

changes in the initial and final stem-count that can arise, for example, from a) natural

sprouting/growth trajectory where new stems sprout and grow, and b) partial mortality i.e.,

mortality of some of the stems. We first quantified the net change in the stem-count of

individuals under control and dung treatment for the dataset analyzed in the maintext. Figure

S3 presents the % of individuals under each change class. 75% or more individuals

experienced no change in stem-count under both control (n=149) and dung treatment

(n=167). Compared to control treatment, more individuals experienced net gain in stem-count

under dung treatment (30 vs. 17) while fewer individuals showed a net loss in stem-count (27

vs 28). There was no statistical difference in the frequency distributions (Chi squared =

2.499, P=0.287, df=2).

1073

Since even these minute changes in stem-counts could have contributed to the overall trends

presented in the main text, we further used GLMMs (similar to main text) to analyze final

size (total diameter) and $RGR_{diameter}$ for only those individuals who showed no change in

stem-count (316 individuals). Our findings were largely consistent with results based on the

full dataset, as presented in the main text. In the case of final basal diameter, the control

individuals were more negatively impacted by sapling density whereas dung treatment

provided a buffer against these negative density-dependent effects, althoug this buffering was

statistically significant for only average-sized saplings (Figure S3). Similarly, $RGR_{diameter}$ of

control individuals was more sensitive to density than the individuals receiving dung input.

However, this effect of dung treatment was statistically significant only for average-sized

individuals but not for larger individuals, although a trend for buffering was visually evident

for large saplings (Figure S3). See supplement Table S3 and Table S4 for more details.

1086

1087

1088

1089

1090

1091

 Supplement text: B

We characterized the climatic space of Nagarahole National Park by extracting temperature

data from ERA 5 Land Daily Aggregates (temperature_2m, temperature_2m_min, and

temperature_2m_max) and monthly total rainfall data from CHIRPS (total precipitation).

These values were calculated for each pixel within Nagarahole National Park and then a

mean across all pixels was obtained to capture average conditions. The seasonality of climate

is presented in Fig 2 in the main text, whereas annual variation is provided in Table S8

below.

1100

1101 **Table S8**: Inter-annual variation in temperature, rainfall and primary productivity of

1102 vegetation measured by normalized difference vegetation index during 2003-2023 in

1103 Nagarahole National Park. The years overlapping with this study are marked in grey.

Year	Mean temp (°C)	Min temp (°C)	Max temp (°C)	Total annual rainfall (mm)
2003	22.63	12.16	34.28	1621.57
2004	22.35	12.63	35.47	1893.71
2005	22.51	12.00	34.78	2274.65
2006	22.33	12.55	34.08	1909.18
2007	22.35	11.77	35.67	2255.60
2008	22.15	12.09	34.27	1839.26
2009	22.57	11.57	34.95	1912.78
2010	22.66	12.31	35.72	2301.46
2011	22.07	11.88	33.12	1901.65
2012	22.52	11.31	34.75	1748.98
2013	22.41	12.53	34.96	1407.30
2014	22.56	13.29	35.23	1525.38
2015	22.59	12.15	34.02	1454.92
2016	23.19	13.55	35.80	1004.71
2017	22.74	13.12	35.22	1663.78
2018	22.62	12.27	33.95	1776.06
2019	22.85	12.20	35.34	1619.18
2020	22.86	13.58	34.92	1692.06
2021	22.30	12.86	34.38	1857.65
2022	22.41	13.08	34.50	1957.72
2023	22.98	11.62	35.47	1145.62