1 ***Bacillus subtilis* reprograms host transcriptome and rhizosphere microbiome via systemic**
2 **signaling to confer alkaline stress tolerance in garden pea**
3

4 **Ahmad H. Kabir[1,][*], Asha Thapa[2], Md Rokibul Hasan[3], Mohammad G. Mostofa[4]**
5
[1]Department of Biology, Lamar University, Beaumont, TX 77705, USA
[2]Evolution, Behavior and Ecology Program, Department of Biology, Boston University, MA
02215, USA
[3]School of Sciences, University of Louisiana at Monroe, Monroe, Louisiana, LA 71209, USA
[4]Department of Chemistry, State University of New York College of Environmental Science and
Forestry, Syracuse, NY 13210, USA
12
13 [*]Correspondence: akabir2@lamar.edu

14
Abstract

Soil alkalinity severely limits legume growth, but the role of *Bacillus subtilis* in alkaline stress
tolerance remains unclear in garden pea. We found that multiple garden pea genotypes
inoculated with *B. subtilis* under alkaline stress showed host-specific improvements in growth
parameters. Mechanistic analysis conducted on Sugar Snap showed improved nodulation,
mineral status, and photosystem efficiency, while split-root assays revealed that *B. subtilis*
triggered systemic signaling underlying alkaline tolerance. Further, FeEDDHA partially reduced
alkaline stress symptoms but did not fully restore nodulation. In contrast, *B. subtilis* enhanced
siderophore availability and improved nodulation, leading to stronger symbiotic recovery than
inorganic Fe alone. This indicates that nodulation recovery is driven primarily by *B. subtilis*–
mediated stimulation of *Rhizobium leguminosarum*, not by Fe availability alone. This is further
supported by *in vitro* co-culture experiments that showed increased growth of both *R.*
leguminosarum and *B. subtilis*, pointing to their complementary interactions that promote mutual
fitness under alkaline stress. RNA-seq analysis identified 958 upregulated and 1,134
downregulated genes in roots inoculated with *B. subtilis* under alkaline conditions. The
upregulated genes were mostly involved in the sugar-mediated symbiotic association (*SWEET*
and *GLUT*), pH homeostasis (*cation/H+ exchanger* and *ATPase*), and nutrient assimilation
(*Ammonium transporter* and *Zn/Fe permease*). Furthermore, *B. subtilis* reshaped the rhizosphere
by restoring microbial community structure and enriching beneficial taxa such as *Pseudomonas*,
Pseudorhizobium, and *Chaetomium*, which may act as helper microbes to promote pea survival
under alkalinity. Taken together, microbial interventions such as *B. subtilis* offer an effective
strategy to boost legume tolerance to alkaline soils.

37
38 **Keywords:** Helper microbes; nutrient assimilation; microbiome dynamics; soil alkalinity; sugar
39 transport; symbiosis.

INTRODUCTION

Soil alkalinity is a widespread abiotic stress that severely constrains crop productivity by limiting the solubility and availability of essential nutrients, particularly iron (Fe), manganese (Mn), and phosphorus (P) (Wang et al., 2025; Rengasamy, 2010). In alkaline soil, high pH disrupts root membrane transport systems, impairs chlorophyll (Chl) biosynthesis, and reduces photosynthetic efficiency, resulting in stunted growth and yield reduction (Thapa et al., 2015). Legume crops are especially sensitive to alkaline stress due to their reliance on symbiotic nitrogen fixation, a process that is highly dependent on optimal nutrient uptake and balanced rhizosphere conditions (Tang et al., 2024). The increasing prevalence of alkaline soils in arid and semi-arid agricultural zones, exacerbated by irrigation with bicarbonate-rich water, underscores the urgent need for sustainable strategies to mitigate their impact (Gamalero et al., 2020).

To cope with soil alkalinity, plants employ a suite of adaptive mechanisms to maintain nutrient acquisition and cellular homeostasis. These include selective ion transport and proton pump activity to regulate the uptake of essential nutrients and cellular pH, respectively (Ku et al., 2022; Kabir et al., 2012). Another strategy involves the exudation of organic acids from roots, which acidify the rhizosphere and improve the solubility of otherwise inaccessible nutrients (Barrow & Hartemink, 2023). Morphological adaptations, such as the proliferation of deep and extensive root networks and the formation of microbial symbioses, further enhance nutrient uptake efficiency under alkaline conditions (Zhang et al., 2020; Chen et al., 2016).

Microbiome-based interventions represent a promising, environmentally sustainable strategy to strengthen plant resilience against alkalinity. Beneficial microbes influence not only local root physiology but also systemic signaling networks that coordinate stress responses at the whole-plant level (Pantigoso et al., 2022; Ortíz-Casro et al., 2009). Such signals can reprogram the root transcriptome, activate defense mechanisms, and restructure microbial communities, making plants more adaptable to stressful conditions (Jamil et al., 2022; Pantigoso et al., 2022; Ruffel et al., 2008). Among plant growth-promoting rhizobacteria (PGPR), *Bacillus subtilis* has gained significant attention for its ability to withstand environmental stresses (Sharma et al., 2023), rapid colonization of roots, and production of diverse bioactive metabolites (Mahapatra et al., 2022; Hashem et al., 2019). It facilitates the secretion of organic acids to solubilize minerals, production of siderophores to chelate Fe, synthesis of phytohormones, and modulation of antioxidant defenses (Hazarika et al., 2023; Chandwani et al., 2023). Despite these known benefits, little is understood about the roles of *B. subtills* in regulating transcriptomes and root microbiomes in the context of soil alkalinity in legumes.

The garden pea (*Pisum sativum* ssp. *hortense*), an important source of dietary micronutrients, is highly sensitive to alkaline stress (Thapa et al., 2025b; Dahl et al., 2012). Elevated soil pH can severely reduce yields by disrupting photosynthetic efficiency, which manifests as leaf chlorosis and growth penalty (Meisrimler et al., 2016). However, the potential of *B. subtilis* to alleviate alkaline stress in garden pea remains poorly understood. In this study, we investigated how *B. subtilis* inoculation modulates defense-signaling pathways, transcriptome reprogramming, and rhizosphere microbiome shifts to enhance alkaline stress tolerance in garden pea. Our findings

90 provide novel insights into the potential of *B. subtilis*-based bioinoculants for improving legume
91 productivity on alkaline soils, contributing to sustainable agriculture in marginal environments.
92
MATERIALS AND METHODS
Plant cultivation and growth conditions
Several garden pea cultivars, such as Sugar Snap, Oregon Sugar Pod, Sugar Ann, Little Marvel,
Green Arrow, and Lincoln, were used in the initial screening of genotypes. Following
preliminary screening, Sugar Snap was chosen for subsequent experiments. Seeds were surface-
sterilized in 2% sodium hypochlorite for 5 min, rinsed three times with sterile water, and placed
in trays for germination at 25 °C for 48 h. Uniform seedlings were transplanted into pots
containing 500 g of a soil mixture (field soil from a pea-growing area combined with commercial
potting mix at a 1:2 ratio). The treatment groups include four conditions using lime [7.5 g
$NaHCO_3$ and 4.5 g $CaCO_3$ per pot] and *B. subtilis* (1×10^8 CFU/mL; ARS Culture Collection–
USDA, NRRL B-14596): (i) control (no lime, pH ~6.5), (ii) alkaline (lime-amended soil, pH
~7.8), (iii) alkaline+BS (lime-amended soil, pH ~7.8, plus *B. subtilis* inoculum), and (iv) BS+
(no lime, pH ~6.5, plus *B. subtilis* inoculum). For plant inoculation, 1 mL of the 1×10^8 CFU
mL^{-1} inoculum was applied to the base of each seedling once at the time of transplantation. This
NRRL B-14596 strain was selected based on a pilot study in which different *B. subtilis* strains
were tested for their strong potential to enhance tolerance in pea exposed to alkalinity. In
addition to the $NaHCO_3$ mixed with the soil from day 1, a 15 mM $NaHCO_3$ solution was
applied weekly throughout the 5-week growth period before data collection (Supplementary Fig.
S1). For a targeted experiment, two additional treatments were included: FeEDDHA+ (lime-
amended soil, pH ~7.8, supplemented with FeEDDHA at 0.5 g/500 g soil) and FeEDDHA
control (non-lime soil, pH ~6.5, with FeEDDHA at 0.5 g/500 g soil). Plants were grown in a
randomized complete block design, with blocks organized by treatment group, under a 10 h light
/ 14 h dark cycle (250 µmol m^{-2} s^{-1}) at ~25 °C in greenhouse conditions. Soil physicochemical
properties are provided in Supplementary Table S1.
117
Molecular detection of *B. subtilis* colonization in roots
DNA-based qPCR analysis was performed to study the root colonization in the roots of Sugar
Snap under different growth conditions. Root samples were rinsed twice in sterile phosphate-
buffered saline, briefly vortexed, and washed twice with sterile water. Genomic DNA was
extracted from approximately 0.2 g of root tissue using the cetyltrimethylammonium bromide
(CTAB) method (Clarke, 2009). DNA quantity and purity were verified using a NanoDrop ND-
1000 (Wilmington, USA), and all samples were normalized prior to qPCR. Amplification was
conducted on a CFX96 Touch Real-Time PCR Detection System (Bio-Rad, USA) using *B.*
subtilis-specific primers (forward: 5′-TCTGCTCGTGAACGGTGCT-3′; reverse: 5′-
TTTCGCCTTATTTACTTGG-3′) (IARRPCAAS, 2013). Reactions were set up with iTaq™
Universal SYBR® Green Supermix (Bio-Rad, USA), with *P. sativum* GAPDH (Fw
GTGGTCTCCACTGACTTTATTGGT, Rv TTCCTGCCTTGGCATCAAA) serving as the
reference genes. Relative quantification of *B. subtilis* abundance was calculated using the
$2^{-\Delta\Delta CT}$ method (Livak & Schmittgen, 2001). This assay was conducted using three biological
replicates and three technical replicates per treatment. Since the goal of this assay was to
compare relative differences in *B. subtilis* colonization among treatments rather than to
determine absolute bacterial numbers, quantification was performed using the $2^{-\Delta\Delta Ct}$ method
normalized to the pea housekeeping gene *GAPDH*.

Measurement of morphological features and chlorophyll fluorescence kinetics

Shoot height and root length were measured for each plant using a measuring tape. Shoots and roots were then harvested and oven-dried at 70 °C for 3 days to determine dry weight. For nodule assessment, roots were gently washed to remove soil debris, and nodules were carefully detached, counted under a magnifying lens, and stored at −80 °C for subsequent analyses. Chlorophyll content was determined with a handheld SPAD meter (AMTAST, USA). Furthermore, chlorophyll fluorescence kinetics (OJIP), including the maximal photochemical efficiency of PSII (F_v/F_m) and the photosynthetic performance index (Pi_ABS), were recorded on the uppermost fully expanded leaves at three distinct points using a portable FluorPen FP 110 (Photon Systems Instruments, Czech Republic). For OJIP measurements, leaves were dark-adapted for 1 h before data acquisition. SPAD and OJIP measurements were taken once at the final harvest. All readings were collected between 10:00am and 12:00pm to minimize the effects of diurnal variation on photosynthetic parameters.

Nutrient analysis in pea roots and leaves

Nutrient analysis was conducted on roots and young leaves. Root samples were excised, rinsed under running tap water, immersed in 0.1 mM $CaSO_4$ for 10 min, and subsequently washed with deionized water. Young leaves were washed separately with deionized water. All samples were placed in labeled envelopes and oven-dried at 75 °C for 3 days. Elemental concentrations were determined using inductively coupled plasma mass spectrometry (ICP-MS) at the Agricultural and Environmental Services Laboratories, University of Georgia.

Determination of ammonium in roots

Ammonia concentration in root tissues was measured using Nessler's reagent spectrophotometric method (Jeong et al., 2013). Fresh root tissue (100 mg) was homogenized in 5 mL of 2% (w/v) potassium chloride, and the homogenate was centrifuged at 12,000 rpm for 10 min. One milliliter of the resulting supernatant was mixed with 1 mL of Nessler's reagent and incubated at room temperature for 15 min. Absorbance was recorded at 420 nm using a spectrophotometer. Ammonia content was calculated from a standard curve generated with known concentrations of ammonium sulfate and expressed on a fresh weight basis (μg g^{-1}FW).

Split-root assay

Plants were first grown for 2 weeks in sterile vermiculite prior to initiating the split-root assay in soil pots. Following transplant, split-root plants were cultivated for an additional 4 weeks before measurements were taken. The split-root setup was established using a single pot with a partitioned assembly, with modifications to the method described by Thilakarathna and Cope (2021). The taproot of each seedling was severed at the midpoint, and the lateral roots were evenly distributed between two compartments within the same pot, each filled with a 1:2 mixture of natural field soil and commercial potting mix. A solid plastic barrier was used to separate the compartments. The plastic barrier prevented soil and solution exchange between compartments. All plants were maintained under identical growth conditions, with soil alkalinity applied according to the treatment design. In each split-root system, one compartment received *B. subtilis* inoculation, while the other remained uninoculated.

182
183
RNA-sequencing and bioinformatics analysis
RNA-seq analysis was performed on root tissues. Before RNA extraction, roots were rinsed twice with sterile water and vortexed for 10 s in sterile phosphate-buffered saline to remove surface debris. Cleaned roots were flash-frozen in liquid nitrogen and ground to a fine powder using a pre-chilled mortar and pestle. Total RNA was extracted using the SV Total RNA Isolation System (Promega Corporation, USA). Samples with RNA integrity number (RIN) values greater than 8 were selected, and 1 µg of RNA per sample was used for library preparation. Libraries were generated with the KAPA RNA HyperPrep Kit with Poly-A Selection (Kapa Biosystems, USA) and amplified using the KAPA HiFi HotStart Library Amplification Kit (Kapa Biosystems, USA). Sequencing was conducted on an Illumina NovaSeq 6000 platform (0.2 Shared S4, 150 bp paired-end) at the RTSF Genomics Core, Michigan State University. Overall, 90.0% of reads passed quality filtering with scores ≥ Q30.

Bioinformatics processing of raw FastQ files was performed using Partek Flow genomic analysis software (Partek, St. Louis, MO, USA). Adapter sequences and low-quality reads were removed with Trimmomatic (Bolger et al., 2014) to obtain high-quality clean reads. The filtered reads were aligned to the *Pisum sativum* reference genome (Ensembl Genomes 62) using HISAT2 (Kim et al., 2015). Furthermore, read counts for each gene were quantified with HTSeq (Anders et al., 2015). Differentially expressed genes (DEGs) were identified using the Limma-Voom method, applying a false discovery rate (FDR) < 0.05 and an absolute log☐ fold change ≥ 2 as significance thresholds. Genes with low expression (<10 normalized counts across all samples) were excluded from further analysis. Expression levels were calculated as fragments per kilobase of exon per million mapped fragments (FPKM). Venn diagrams were generated using VennDetail (Hinder et al., 2017), and heatmaps were produced with the *pheatmap* package in R (Hu et al., 2021). Functional enrichment analysis of DEGs was conducted using ShinyGO 0.76.3 (Ge et al., 2020), with gene ontology (GO) annotations obtained from EnsemblPlants (Yates et al., 2022).

Amplicon-based 16S and ITS sequencing in the rhizosphere
Rhizosphere microbial populations were characterized using Illumina amplicon sequencing, targeting the ITS region for fungi and the 16S rRNA gene for bacteria. Root samples were vortexed in sterile phosphate-buffered saline (PBS) for 10 s, rinsed twice with sterile water, and ~0.2 g of cleaned tissue was used for DNA extraction with the CTAB method (Clarke, 2009). RNase and Proteinase K treatments were included to remove RNA and protein contaminants, respectively. Amplicon libraries were generated using primer pairs 341F (CCTACGGGNGGCWGCAG) and 805R (GACTACHVGGGTATCTAATCC) for the 16S rRNA gene, and ITS3 (GATGAAGAACGYAGYRAA) and ITS4 (TCCTCCGCTTATTGATATGC) for the ITS region. Sequencing was conducted on the Illumina NovaSeq 6000 platform (PE250). Raw reads were processed and trimmed with Cutadapt (Martin, 2011) and analyzed using the DADA2 pipeline (Callahan et al., 2016). Sequences identified as mitochondrial and chloroplast origin were removed. Taxonomic classification of amplicon sequence variants (ASVs) was performed using the UNITE database (Nilsson et al., 2019) for fungal ITS sequences and SILVA 138 (Quast et al., 2013) for bacterial 16S sequences. Microbial community structure was assessed by calculating Bray-Curtis distance

matrices from Hellinger-transformed abundance data, followed by Principal Coordinate Analysis (PCoA) using the *phyloseq* and *vegan* packages in R. Diversity analyses included alpha and beta diversity metrics, relative abundance profiling, and non-parametric Kruskal–Wallis tests, with significance determined at P < 0.05 (McMurdie & Holmes, 2013).

Microbial co-culture method

Interactions between *B. subtilis* and *R. leguminosarum* were evaluated on nutrient agar under control and alkaline conditions, with the latter supplemented with 15 mM NaHCO₃ to adjust the pH to 7.8. *B. subtilis* NRRL B-14596 and *R. leguminosarum* viciae 3841 were first grown overnight in nutrient broth at 28 °C with shaking at 200 rpm. Cultures were then adjusted to $1 \times 10^\square$ CFU mL^{-1}, as determined by dilution plating and OD–CFU calibration curves established for each species. A 5 μL inoculum of each microbial strain was cultured in three configurations: *B. subtilis* alone, *R. leguminosarum* alone, and a mixed culture (1:1 ratio) of *B. subtilis* + *R. leguminosarum*. Plates were incubated for 7 days, after which colony growth was quantified using ImageJ software (National Institutes of Health, USA). The radius of each colony was measured at three independent points, and the mean value was calculated for subsequent analysis.

Data analysis

Statistical analyses were performed in SPSS Statistics 20.0 (IBM Corp., Armonk, NY, USA) and R using the ggplot2 package for visualization. Prior to ANOVA, data were tested for normality (Shapiro–Wilk test) and homogeneity of variances (Levene's test). When assumptions were violated, data were log or square root transformed; however, non-transformed values are shown when results remain unchanged. For experiments involving multiple pea genotypes and treatments, data were analyzed using a two-way ANOVA with genotype, treatment, and their interaction (genotype × treatment) as fixed factors. For analysis within a single genotype (Sugar Snap), one-way ANOVA was used with treatment as the independent factor. When ANOVA indicated significant effects ($p < 0.05$), post hoc mean separation was performed using Duncan's Multiple Range Test (DMRT). In this study, the physiological assays were performed with five independent replications, whereas the RNA-seq and amplicon sequencing analyses were conducted with three independent replications.

RESULTS

Screening of genotypes for tolerance to soil alkalinity with *B. subtilis*

Across genotypes, alkaline stress consistently reduced shoot height and SPAD scores compared to control plants, though the extent of recovery with *B. subtilis* varied (Table 1). In Sugar Snap and Oregon Sugar Pod, both shoot height and SPAD scores decreased significantly under alkaline stress but were similar to control levels when inoculated with *B. subtilis* with or without alkaline stress (Table 1). In Sugar Ann and Little Marvel, alkaline stress reduced both traits. Plants exposed to alkaline stress inoculated with *B. subtilis* improved shoot height, while BS+ produced the highest values in Little Marvel and Sugar Ann, exceeding the control (Table 1). In Little Marvel, SPAD scores improved under alkaline+BS and BS+ conditions, but BS+ matched control plants (Table 1). For the cultivar Sugar Ann, BS+ showed higher SPAD values than the control, but the difference was not statistically significant. Green Arrow and Lincoln displayed a different pattern, with shoot height reduced under alkaline stress and remaining low under alkaline+BS conditions but were found similar to control levels by *B. subtilis*. Similarly, SPAD scores declined under alkaline conditions, remained unchanged in response to *B. subtilis* under

soil alkalinity. Green Arrow and Lincoln inoculated with *B. subtilis* showed similar SPAD values to those of controls (Table 1).

Effect of *B. subtilis* on physiological parameters and root colonization

Phenotypic observation showed that plants under alkaline had smaller shoots and roots compared to control (Fig. 1A). Plants inoculated with *B. subtilis* showed an improvement in shoot and root growth compared to alkaline stress, while plants inoculated with *B. subtilis* without stress appeared similar to control (Fig. 1A). The relative abundance of *B. subtilis* in roots was low in control and alkaline stress but significantly increased only in alkaline+BS conditions (Fig. 1B). Plant fresh weight was significantly reduced in alkaline compared to control, increased under soil alkalinity with *B. subtilis* to values comparable to control, and remained similar to the plants solely inoculated with *B. subtilis* (Fig. 1C). Further, shoot height followed the same trend (Fig. 1D). Furthermore, root length (Fig. 1E) and root FW (Fig. 1F) were significantly reduced under alkaline stress compared to control. However, both parameters were restored to control levels under alkaline+BS conditions and remained comparable to control in BS+. We also determined the root ammonium content in the roots. Root ammonium content was significantly affected by treatments. Control and BS+ plants showed the highest values, which were not significantly different from each other, while alkaline treatment had the lowest ammonium content (Fig. 1G). Plants inoculated with *B. subtilis* under soil alkalinity partially restored ammonium levels, resulting in significantly higher values than alkaline alone but still lower than control and BS+ conditions (Fig. 1G). We also found that F_v/F_m and PI_ABS declined significantly in alkaline conditions compared to control, increased under alkaline stress inoculated with *B. subtilis* conditions to control levels, and remained unchanged in BS+ relative to control (Fig. 1H and 1I).

Effect of *B. subtilis* on nutrient concentrations in roots and leaves

ICP-MS analysis of nutrient contents showed a variation in their concentrations in the root and leaf tissues of garden peas grown with or without *B. subtilis* under alkaline conditions (Table 2). In alkaline-stressed plants, there was a decrease in concentrations of Fe, Mn, and N in both root and leaf tissues. However, the addition of *B. subtilis* to alkaline-stressed plants recovered these nutrient contents in both root and leaf tissues and was similar to control plants. Additionally, *B. subtilis* inoculated with control plants showed Fe, Mn, and N concentrations in the roots and leaves of garden peas that were similar to those in the control and alkaline + BS treatments. However, there were no significant changes in the amount of S in the root tissues under alkaline conditions compared to the control. There was a decrease in S content in leaf tissues in such conditions, and it was recovered when *B. subtilis* was added (Table 2). Similarly, there was a significant decrease in magnesium (Mg) content in roots of alkaline-stressed plants; however, there was no change in Mg content in leaf tissues under all four treatment conditions. Additionally, under alkaline conditions, calcium (Ca) concentrations in both the roots and leaves of garden peas decreased significantly, while inoculation of alkaline-stressed plants with *B. subtilis* increased Ca content in both tissues (Table 2).

Comparative performance of *B. subtilis* and inorganic Fe in alkaline soil

The effects of *B. subtilis* inoculation and inorganic Fe supplementation on pea growth under alkaline conditions were investigated and compared (Fig. 2A). The relative abundance of *B.*

subtilis in roots was low in control, alkaline, alkaline+Fe, and inorganic Fe plants (Fig. 2B). In contrast, alkaline+BS conditions showed a significant increase in *B. subtilis* abundance as compared to other treatments without *B. subtilis* additions (Fig. 2B). Rhizosphere siderophore production differed markedly among treatments (Fig. 2C). Before applying the inoculum to plants, we first tested whether *B. subtilis* maintained consistent siderophore production under control and alkaline nutrient conditions. Siderophore production by the *B. subtilis* inoculum showed no statistically significant differences between treatments (Supplementary Fig. S2). Alkaline conditions caused a significant decline in siderophore in the rhizosphere compared to controls. When *B. subtilis* was introduced under alkaline conditions, siderophore production increased significantly compared to alkaline conditions, reaching levels statistically comparable to the control (Fig. 2C). Interestingly, supplementation with inorganic Fe under alkaline conditions resulted in no significant increase compared to alkaline stress, which remained significantly lower than *B. subtilis*-treated plants (Fig. 2C). Inorganic Fe under alkaline conditions did not restore siderophore production to control levels, whereas alkaline+BS fully recovered siderophore activity (Fig. 2C).

Alkaline stress had a significant negative effect on nodule formation in pea roots compared to the control (Fig. 2D). Inoculation with *B. subtilis* under alkaline conditions partially rescued nodulation, resulting in significantly higher nodule numbers than alkaline-stressed plants without inoculation. In contrast, supplementation with inorganic Fe under alkaline conditions provided only a modest improvement, and the increase remained lower than that observed in *B. subtilis*–inoculated alkaline plants (Fig. 2D). Furthermore, plants inoculated with *B. subtilis* under non-alkaline conditions or supplemented with inorganic Fe exhibited nodule numbers comparable to the control (Fig. 2D). Under alkaline stress, plant height decreased significantly compared to the control (Fig. 2E). Inoculation with BS or supplementation with inorganic Fe under alkaline stress significantly increased shoot height relative to alkaline conditions, with values approaching those of the control. Plants inoculated with *B. subtilis* exhibited plant height similar to controls, whereas inorganic Fe supplementation increased plant height slightly above control values (Fig. 2E). Furthermore, SPAD scores under alkaline stress significantly reduced in relation to that in control (Fig. 2F). Both alkaline+BS and alkaline+Fe treatments restored SPAD scores to levels significantly higher than alkaline-stressed plants. Plants solely inoculated with *B. subtilis* or inorganic Fe showed similar SPAD scores to those of control plants (Fig. 2F).

Split-root assay for identifying systemic signaling
A split-root system was used to distinguish between local and systemic effects of *B. subtilis* inoculation underlying alkaline stress tolerance in pea plants (Fig. 3A–F). In split-root experiments, root length and root fresh weight did not differ significantly between compartments across all treatments (Fig. 3A–B). However, plants exposed to alkaline stress in both compartments (SR2) showed no significant reduction in root length and root fresh weight compared with the control treatment (SR1) (Fig. 3A-B). In contrast, SR3 (alkaline/alkaline+BS) and SR4 (alkaline+BS/alkaline+BS) plants showed substantial improvements in root length and root fresh weight relative to SR2 (alkaline/alkaline) plants (Fig. 3.A–B). Further, shoot traits responded strongly to treatment combinations (Fig. 3C–D). Among all treatments, SR2 consistently exhibited the lowest shoot growth responses compared to SR1. Plants exposed to *B. subtilis* in both compartments (SR4) or in one compartment under alkaline stress (SR3) showed significantly greater shoot height and shoot fresh weight than plants under alkaline stress without

inoculation (Fig. 3C–D). Furthermore, leaf SPAD and Fv/Fm were significantly reduced under alkaline stress without inoculation (SR2), whereas plants inoculated with *B. subtilis* in one or two compartments with or without alkaline stress (SR3–SR6) showed a significant increase in SPAD and Fv/Fm levels than SR2 plants (Fig. 3E-F).

Interaction between *B. subtilis* and *R. leguminosarum* in co-culture
The effect of co-culturing *B. subtilis* with *R. leguminosarum* was assessed under both control and alkaline conditions over three, five, and seven days (Fig. 4A-4B; Supplementary Fig. S3). Under non-alkaline conditions, colony sizes of *R. leguminosarum* and *B. subtilis* increased progressively over time in both single and co-culture treatments. On day three, no significant differences in colony size were observed between single and co-culture for either species. By day five and day seven, both species exhibited larger colony diameters compared to day three, but differences between single and co-culture remained statistically non-significant (Fig. 4A). Under alkaline conditions, colony sizes of both *R. leguminosarum* and *B. subtilis* were consistently smaller compared to control pH across all time points. On day three, growth reduction was evident in both species relative to their respective control pH counterparts, and coculture significantly altered the colony size of *R. leguminosarum* with an increase in colony area (Fig. 4B). Similar patterns were observed on days five and seven, with gradual increases in the colony diameter of *R. leguminosarum* over time, and significant differences between the single- and co-culture treatments under alkaline stress. However, there were no changes in colony growth of *B. subtilis* over three, five, and seven growing days between single and co-culture conditions (Fig. 4B). Overall, colony expansion for both species was slower in alkaline conditions than in control pH, and co-culturing confers a measurable growth advantage for only *R. leguminosarum* under alkaline conditions (Fig. 4B).

Transcriptomic changes in roots
Principal coordinate analysis (PCoA) revealed four distinct clusters, indicating that each treatment generated a unique transcriptomic profile (Fig. 5A). In Venn diagram analysis, alkaline+BS treatment had a unique set of upregulated (958) and downregulated (1134) genes in the roots compared to alkaline stress (Fig. 5B-5C). Specifically, alkaline+BS exhibited a large number of uniquely upregulated genes not shared with alkaline or BS+, along with a smaller subset of genes overlapping with control-related profiles (Fig. 5A). Heatmap visualization (Fig. 5D–G) highlighted key functional categories of differentially expressed genes (DEGs). In the alkaline+BS treatment, numerous genes related to symbiotic association (e.g., sugar phosphate transporter, glucose transporter GLUT, Sugar transporter SWEET, MF sugar transporter–like, Sugar transport protein STP, UDP–sugar pyrophosphorylase) were significantly upregulated compared to alkaline stress (Fig. 5D; Supplementary Table S2). Genes associated with nutrient transport and assimilation (e.g., ammonium transporter, nitrite and sulphite reductase, inorganic pyrophosphatase, amino acid transporter, Cys/Met metabolism, Iron/zinc purple acid phosphatase, malic oxidoreductase, SulP transporter, Zinc/iron permease, MFS transporter) were also more highly expressed in alkaline+BS compared to alkaline stress (Fig. 5E). Under alkaline conditions without *B. subtilis*, these genes generally exhibited reduced expression compared to control plants (Fig. 5E). Furthermore, redox and osmotic adjustment genes (e.g., glutaredoxin, oxidative stress regulator, glutathione synthase, thioredoxin, glutathione *S*-transferase, ferritin-like superfamily, dehydrin, aquaporin transporter) were induced in alkaline+BS relative to the plant solely stressed with alkaline (Fig. 5F). Lastly, several genes (*isoprenoid synthase*, *ethylene-*

responsive TF, *cation/H+ exchanger*, *ABC-2 type transporter*, *ABC transporter type 1*, *root meristem growth factor 3*, *AP2/ERF ethylene responsive factor*, *oxoglutarate*, *small auxin-up RNA*, *ATPase*) related to growth and stress response showed significant upregulation due to B. subtilis under soil alkalinity related to alkaline soil (Fig. 5G; Supplementary Table S2).

Gene ontology enrichment analysis

Gene ontology (GO) enrichment analysis of the alkaline+BS versus alkaline comparison revealed distinct functional categories associated with upregulated and downregulated genes (Fig. 6A-B). Among the upregulated genes, the most significantly enriched GO terms were related to responses to water, responses to acid chemicals, and responses to inorganic substances, indicating activation of stress-responsive pathways under *B. subtilis* treatment. Additional enriched terms included oxidoreductase activity, protein processing in the endoplasmic reticulum, and several membrane- and transporter-associated functions, suggesting enhanced cellular transport and redox regulation (Fig. 6A). Conversely, downregulated genes were strongly enriched for GO categories associated with oxygen binding, heme and tetrapyrrole binding, ion binding, and defense responses, indicating repression of oxidative and defense-related processes in the alkaline+BS treatment. Several broad metabolic and regulatory categories including regulation of cellular processes, cellular metabolic processes, nitrogen (N) compound metabolism, and macromolecule biosynthesis were also significantly enriched among the downregulated genes, reflecting a reorganization of metabolic priorities (Fig. 6B).

Rhizosphere microbial community structure

Principal coordinate analysis (PCoA) based on Bray–Curtis dissimilarity indicated that plants grown under non-stressed conditions, with or without *B. subtilis*, occupied a similar region of ordination space (Fig. 7A). In contrast, alkaline and alkaline+BS samples were clearly separated from the non-stressed groups and from each other (Fig. 7A). Alpha diversity analysis showed modest variation across treatments, but overall differences were limited. Particularly, soil alkalinity altered bacterial richness, while B. subtilis modestly shifted richness patterns; however, these trends were not strongly treatment-specific (Fig. 7B). In contrast, Simpson diversity did not differ significantly among treatments (Fig. 7B). At the family level, *Pseudomonadaceae*, *Gemmatiomonadaceae*, and *Flavobacteriaceae* showed statistically significant differences in relative abundance among the treatments (Fig. 7C). *Pseudomonadaceae* and *Flavobacteriaceae* were markedly enriched in the rhizosphere of plants inoculated with *B. subtilis* under soil alkalinity compared to alkaline stress only. Other dominant families such as *Gemmatiomonadaceae*, *Rhizobiaceae*, and *Sphingomonadaceae* exhibited treatment-dependent shifts but without statistical significance (Fig. 7C). At the genus level, *Variovorax*, *Shinella*, *Pseudorhizobium*, *Pseudomonas*, *Novosphingobium*, *Flavobacterium*, *Brevundimonas*, and *Bosea* displayed statistically significant variation across treatments (Fig. 7D). Particularly, *Pseudomonas* and *Pseudorhizobium* showed a significant enrichment in alkaline-exposed conditions inoculated with *B. subtilis* relative to the conditions solely exposed to soil alkalinity. In contrast, *Shinella* was enriched under alkaline stress, while *Bosea* and *Variovorax* were predominant in control plants (Fig. 7D).

In this study, PCoA of ITS showed distinct clustering of rhizosphere fungal communities among the treatment groups (Fig. 8A). Particularly, control and BS+ grouped closely, indicating minimal differences in fungal composition under non-stress conditions. In contrast, alkaline and

alkaline+BS occupied separate positions from control, with alkaline+BS partially shifting toward the control/BS+ cluster, indicating that *B. subtilis* inoculation under alkaline stress altered fungal community structure (Fig. 8A). In our analysis, observed richness and Simpson diversity index showed no significant changes due to B. subtilis in the presence or absence of soil alkalinity (Fig. 8B). Among the families, *Halosphaeriaceae* and *Cladosporiaceae* exhibited lower relative abundance, whereas *Chaetomiaceae* showed higher abundance under soil alkalinity inoculated with *B. subtilis* compared to alkaline stress (Fig. 8C). In contrast, the dominant family *Aspergillaceae* displayed broadly similar abundance between the two treatments. Among significant genera, *Pseudallescheria* and *Chaetomium* showed significant enrichment in the rhizosphere of plants inoculated with *B. subtilis* under soil alkalinity compared to alkaline stress (Fig. 8D).

DISCUSSION

Soil alkalinity is a major constraint to legume productivity, primarily due to reduced availability of essential micronutrients (Gong et al., 2020). In this study, alkaline stress significantly disrupted pea performance across multiple physiological scales from nutrient acquisition to canopy chlorophyll status and photosystem function, leading to reduced biomass accumulation. Inoculation with *B. subtilis* effectively alleviated these negative impacts, restoring nutrient uptake, improving photosynthetic efficiency, and enhancing biomass accumulation. The underlying resilience mechanism appears to be multifactorial, encompassing direct nutrient mobilization, systemic signaling, and rhizosphere microbiome restructuring, suggesting *B. subtilis* as a holistic modulator of pea performance under high pH conditions.

***B. subtilis* rebalances physiological responses under alkaline stress**

The present study highlights substantial genotype-dependent variation in alkaline stress tolerance and responsiveness to *B. subtilis* (Table 1). Although all cultivars experienced reduced growth and Chl content under alkaline conditions, the magnitude of recovery following inoculation differed markedly across genotypes. Such variability is consistent with the previous reports, suggesting host genotype as a major player in shaping rhizomicrobial colonization, signal perception, and downstream transcriptional reprogramming (Thapa et al., 2025a; Jacoby et al., 2021). Genotype-specific differences in host response modulation and symbiotic compatibility likely influence the extent to which *B. subtilis* helps pea plants cope with stress. Alkaline stress reduced Fe and Mn concentrations in both roots and leaves in Sugar Snap, aligning with the reduced solubility and impaired Strategy I Fe acquisition in high-pH soils (Kim and Guerinot, 2007). In this study, *B. subtilis* restored Fe and Mn availability to near control levels and improved Ca and N accumulation, indicating coordinated recovery of nutrient homeostasis in peas. These ionomic recoveries paralleled gains in plant biomass and photosynthetic parameters, pointing to the coordinated restoration of resource supply and photochemical capacity. Such recovery aligns with established traits of *B. subtilis* relevant to alkaline soils, including siderophore and organic acid production, and antioxidant pathway activation (Kloepper et al., 2004; Egamberdieva et al., 2017). The fact that similar benefits occurred across genetically diverse pea cultivars, albeit with variable magnitudes, reflects the common but genotype-dependent nature of PGPR–legume interactions (Hasan et al., 2025; Vejan et al., 2016). Overall, these findings demonstrate that *B. subtilis* is highly effective in restoring micronutrient balance and photosynthetic performance in legumes under alkaline stress.

Systemic signaling reveals a whole-plant response

The split-root assay demonstrated that *B. subtilis* inoculation in only half of the root system conferred comparable benefits to uninoculated compartments, indicating that the effects extended beyond the local rhizosphere and involved systemic signaling. These responses point toward the generation of mobile molecular signals, potentially including phytohormones, peptides, and secondary metabolites that travel through the vascular system to modulate physiology in distant root zones and shoots (Martínez-Medina et al., 2013). The observed improvements in plant biomass and photosynthetic efficiency in both inoculated and distal root zones suggest that *B. subtilis* triggers a whole-plant acclimation program. This likely involves the integration of root-derived signals into shoot-level metabolic adjustments, ensuring resource allocation, and energy production. Such long-distance communication resembles induced systemic tolerance reported for other plant PGPR, where signal perception in one root region primes the rest of the plant for improved stress resilience (Conrath et al., 2006; Hartmann et al., 2021). Additionally, microbe-triggered systemic reprogramming often enhances antioxidant defenses, nutrient mobilization pathways, and the expression of transporter families critical for nutrient economy under abiotic stress (Verma et al., 2024; Bhattacharyya & Jha, 2012). Our transcriptomic enrichment in transporter and signaling genes aligned with this established framework, supporting the idea that *B. subtilis* activates a coordinated, plant-wide regulatory network rather than exerting effects solely on the site of inoculation. This systemic coordination also mirrors recent findings where PGPR elicit cross-compartmental benefits through volatile compounds, secreted metabolites, and immune-regulatory cues that act across the root–shoot continuum (Ryu et al., 2004; Fincheira & Quiroz, 2018). Together, these observations provide compelling evidence that *B. subtilis* initiates a whole-plant response that integrates local rhizosphere interactions with long-distance signaling, ultimately promoting pea resilience under alkaline stress.

Inorganic Fe reduces symptoms, but *B. subtilis* drives symbiotic recovery

The FeEDDHA supplementation study offered mechanistic control by directly supplying a highly stable Fe-chelate capable of maintaining Fe solubility at high pH (Rojas et al., 2008). Alkaline soils can impair root growth, thereby reducing the efficient absorption of essential minerals by plants (Saleem et al., 2023). In this study, while FeEDDHA remains stable under alkaline conditions, a portion of the Fe can still bind to soil particles, decreasing its bioavailability and overall effectiveness (Zuluaga et al., 2023). Although inorganic Fe restored chlorophyll and partially improved growth, it did not recover nodulation or siderophore-associated Fe mobilization, highlighting its narrower mode of action compared to *B. subtilis*. Once Fe limitation was chemically bypassed, the physiological necessity for *B. subtilis*-driven nutrient mobilization was removed. Although inorganic Fe supplementation provided modest improvements, it did not fully counteract the constraints imposed by high pH. Interestingly, inorganic Fe alone does not fully compensate for alkaline-induced limitations in siderophore availability and nodulation in pea plants. However, the consistently elevated siderophore production in *B. subtilis*–treated plants suggest a key role for this microbe in enhancing rhizosphere Fe-mobilization capacity under alkaline stress, outperforming inorganic Fe amendment alone.

Because alkaline stress substantially inhibits nodulation, we next examined whether *B. subtilis* promotes *Rhizobium* performance directly, using co-culture assays. The co-culture experiments

in Fe-sufficient plates revealed no significant differences in colony expansion between *B. subtilis*, *R. leguminosarum.* Under alkaline conditions, *R. leguminosarum* and *B. subtilis* respond differently to pH stress, resulting in distinct growth dynamics even within the same co-culture treatment. Such divergence did not appear under control conditions, indicating that the variation is possibly driven by species-specific sensitivity to alkaline media. However, their mixed cultures at high pH showed growth enhancement, indicating that the synergistic effects observed in planta are likely to arise from complementary interactions in alkaline media. Consistent with this, improved Fe and N status supported by increased ammonium transport in *B. subtilis*-treated roots under alkaline stress suggests that these benefits are alkaline-dependent. This further implies that the restoration of nodulation, as well as improved Fe and N status, likely stems from the positive impact of *B. subtilis* on *R. leguminosarum*, rather than from improved plant Fe availability alone. This plant-mediated facilitation aligns with previous reports in PGPR–rhizobia systems, where enhanced symbiotic performance under stress relies more on the host's metabolic and signaling reprogramming rather than on direct microbe–microbe interactions (Frey-Klett et al., 2011; Vishwakarma et al., 2020). This whole-plant response is consistent with transcriptomic enrichment in transporter and signaling genes that likely mediate these distal effects. Moreover, the complementarity and mutualistic reinforcement between *B. subtilis* and *R. leguminosarum* may facilitate higher rhizobial abundance and a more stable microbiome assembly, contributing to sustained pea performance under alkaline conditions. These findings suggest a complementary role for microbial inoculants and inorganic Fe supplementation. FeEDDHA offers rapid correction of acute Fe deficiency, while *B. subtilis* provides a sustainable, broader-spectrum improvement that also benefits Mn and Ca uptake, photosynthetic resilience, and microbiome health. In low-input or organic systems, *B. subtilis* may thus serve as a viable alternative to synthetic chelates, avoiding their higher costs and potential environmental persistence.

Transcriptomic reprogramming improves nutrient acquisition and redox homeostasis
RNA-seq analysis revealed that *B. subtilis* inoculation under alkaline stress triggered a broad transcriptional reprogramming in pea roots, with pronounced upregulation of genes associated with symbiotic association, mineral acquisition, phytohormone signaling, and redox regulation. Alkali-exposed plants inoculated with *B. subtilis* showed the upregulation of several sugar transport–related genes (*SWEET*, *GLUT*). Upregulation of multiple sugar transporters indicates enhanced carbon allocation and exchange during symbiotic association with *B. subtilis*. Such adjustments likely support microbial metabolism while maintaining the plant's energy balance (Li et al., 2025; Julius et al., 2017). Also, these transporters likely facilitate bidirectional sugar movement between the host and symbiotic partners mediated by *B. subtilis*, ensuring an adequate supply of carbon for microbial metabolism while maintaining host energy balance. Particularly, SWEET sugar transporters not only facilitate carbon allocation but also modulate nodule functioning and microbial colonization (Chen et al., 2023; Breia et al., 2021). Efficient sugar trafficking to nodules and rhizosphere microbes (Lei et al., 2025) and symbiotic associations (Pieterse et al., 2014; Zamioudis & Pieterse, 2012). Such reprogramming of carbohydrate transport pathways aligns with reports that PGPR like *B. subtilis* stimulate sugar-mediated nutrient exchange to sustain nitrogen fixation and plant resilience under high pH conditions (Berendsen et al., 2018; Kryvoruchko et al., 2018). This may be linked to the upregulated genes for ammonium transport (*PSAT0S1495G0040*), amino acid transport (*PSAT7G178640*), and Cys/Met metabolism (*PSAT6G052400*) that are associated with S and S assimilation, critical for stress adaptation (Hasan et al., 2025). Furthermore, enrichment of transporter-related transcripts,

including those for Fe, Mn, nitrate, and phosphate uptake, suggests an enhanced capacity for nutrient mobilization and assimilation under conditions where availability is otherwise restricted by high pH (Lucena, 2006; Kobayashi & Nishizawa, 2012). Ion transport–related genes, including cation/H□ exchanger (*PSAT3G088320*) and ATPase (*PSAT7G218840*), were also elevated, likely contributing to pH homeostasis and nutrient uptake under alkaline stress (Wang et al., 2014). ABC transporters (*PSAT2G058680* and *PSAT3G053400*) and oxoglutarate-related metabolism (*PSAT6G045440*) may further support metabolite exchange and carbon–nitrogen balance, as previously reported for legumes under symbiotic and abiotic stress conditions (Khan et al., 2020).

Furthermore, there was a strong induction of genes involved in redox homeostasis and osmotic adjustment, suggesting that *B. subtilis* enhances antioxidant defense and water balance under alkaline stress. Similarly, the oxidative stress regulator suggests coordinated activation of stress-responsive transcriptional programs, in agreement with other findings that PGPR treatments can activate redox signaling networks in legumes (Al-Turki et al., 2023; Ahmad et al., 2022). Upregulation of oxidoreductase genes likely bolsters the plant's oxidative defense, enabling protection of cellular structures and photosynthetic machinery from oxidative damage (Choudhury et al., 2017). Elevated ferritin-like protein transcripts (*PSAT6G040040*) indicate improved Fe storage and mitigation of Fe-induced oxidative bursts (Hasan et al., 2025). In parallel, there was a suppression of defense-associated transcripts, indicating a strategic reallocation of resources away from constitutive defense toward growth and nutrient acquisition, a hallmark of induced systemic tolerance (Conrath et al., 2006). This transcriptional profile aligns with the systemic effects observed in the split-root assay, reinforcing the idea that *B. subtilis* elicits long-distance signaling that primes the plant for improved nutrient uptake and growth under alkaline stress, while minimizing the metabolic cost of constant defense activation. Such a coordinated molecular shift is consistent with the multifaceted benefits of *B. subtilis* inoculation in garden pea, where stress tolerance and symbiotic compatibility converge to enhance overall plant performance under alkaline stress conditions.

Microbiome restructuring as an ecological layer of alkaline tolerance
The PCoA analysis of bacterial communities demonstrated that *B. subtilis* inoculation modulated rhizosphere assemblages under alkaline stress. Soil alkalinity altered bacterial richness patterns, and *B. subtilis* modestly shifted richness profiles, although diversity metrics were not fully restored. The similarity of bacterial community structure in non-stress conditions with or without *B. subtilis* suggests that *B. subtilis* does not disrupt native bacterial networks in healthy soils but rather coexists within the established community (Hartman & Tringe, 2019). In contrast, alkaline stress induced a substantial shift in community composition, likely driven by high pH-mediated alterations in nutrient availability for stress-tolerant taxa (Rousk et al., 2010). Importantly, the partial restoration of the microbial profiles under soil alkalinity toward the control cluster implies that *B. subtilis* inoculation rebalanced the microbial network under stress, potentially via root exudate modulation and production of metabolites that favor beneficial taxa (Gong et al., 2023). At the genus level, the enrichment of *Pseudomonas* and *Pseudorhizobium* influenced by *B. subtilis* under soil alkalinity aligns with their documented plant growth–promoting traits, including production of phytohormones, phosphatases, and siderophores (Compant et al., 2019; Glick, 2014). *Pseudomonas* spp. is well-known for producing siderophores and phytohormones under alkaline conditions (Lemare et al., 2022; Sainz-Mejías et al., 2020). Furthermore, studies

demonstrate that fluorescent pseudomonads secrete high-affinity siderophores (pyoverdines) that mobilize Fe and can be especially beneficial under alkaline or Fe-limiting soils (Cornelis and Matthijs, 2007; Kümmerli, 2022). Many strains solubilize inorganic phosphate via the PQQ-dependent gluconic-acid pathway and phosphatases, improving P uptake and root growth (Lugtenberg & Kamilova, 2009). *Pseudorhizobium* is a genus separated from Rhizobium by phylogenomics, with isolates from seawater, rocks, and polluted soils, adapted to harsh environments (Lassalle et al., 2021). Although its effects on crops are less documented, *Pseudorhizobium's* contribution to plant performance likely stems from metal/metalloid transformation and stress-tolerant physiology (Lassalle et al., 2021; Andres et al., 2013). The enrichment of *Klebsiella*, and *Brevundimonas* due to *B. subtilis* under non-stress conditions points toward enhanced functional redundancy in the rhizosphere, which could buffer against future environmental perturbations. Collectively, these results suggest that *B. subtilis* does not simply act as a single inoculant but rather as a microbiome engineer, reshaping other beneficial bacterial networks to support nutrient acquisition, stress resilience, and host–microbe symbiotic stability under soil alkalinity.

Similarly, ITS-based profiling showed that *B. subtilis* moderated alkaline-induced shifts in fungal assemblages, partially realigning community composition toward the control state. Alkaline stress reduced fungal richness and diversity, consistent with the suppression of pH-sensitive saprophytic and mutualistic fungi (Lauber et al., 2009). Although richness and diversity did not differ significantly, fungal community composition shifted toward that of control treatments with BS inoculation. *Pseudallescheria* (teleomorph of Scedosporium) is best supported as a saprotroph that degrades lignin and aromatics, processes that can indirectly enhance nutrient turnover in soils (Poirier et al., 2023). Also, *Pseudallescheria* detected in plant tissues may reflect transient environmental contamination rather than stable, beneficial endophyte status (Zhang et al., 2023). However, *Chaetomium* can act as a beneficial endophyte/biocontrol (Feng et al., 2023). It can also trigger induced systemic resistance since transcriptomic work in tomato shows JA/ET-linked immune priming after *C. globosum* treatment (Singh et al., 2021). Beyond defense, *C. globosum* mobilizes phosphorus via secreted phosphatases/phytases, increasing crop P uptake and yield in wheat and pearl millet (Tarafdar and Gharu, 2006). Hence, the enrichment of *Pseudallescheria* and *Chaetomium* by *B. subtilis* under alkaline soil may provide indirect benefits to pea plants by accelerating decomposition and recycling of nutrients.

CONCLUSIONS

Our findings demonstrate that *B. subtilis* inoculation effectively mitigates the adverse effects of alkaline stress in garden pea by restoring Fe, Mn, and N levels, improving photosystem efficiency, and increasing biomass across diverse cultivars (Fig. 9). Split-root assays indicated the involvement of systemic mobile signaling, while co-culture experiments revealed complementary interactions between *B. subtilis* and *Rhizobium*. Interestingly, nodulation recovery appears to result mainly from *B. subtilis*–induced enhancement of *R. leguminosarum* activity rather than from Fe availability alone. Transcriptome profiling further showed strong induction of genes associated with sugar transport, pH homeostasis, and nutrient assimilation, together with elevated expression of oxidoreductases and ferritin-like protein, suggesting improved antioxidant defense and Fe storage. In parallel, *B. subtilis* enriched beneficial taxa, such as *Pseudomonas*, *Pseudorhizobium,* and *Chaetomium*, which may function as helper

688 microbes to reinforce stress tolerance under alkaline conditions. Collectively, these findings
689 highlight how *B. subtilis* orchestrates molecular, physiological, and microbiome-level
690 reprogramming, establishing its potential as a targeted bioinoculant for sustaining legume
691 productivity in alkaline soils.
692

Data availability

Illumina sequencing data were submitted to NCBI under the following accession numbers:
PRJNA1313627 (RNA-seq), PRJNA1313634 (16S) and PRJNA1313644 (ITS).

Acknowledgments

We express our gratitude to Louisiana Biomedical Research Network for the funding. This
research was also supported by a startup grant from Lamar University.

Author contributions

AHK conceived the study, performed the molecular experiments, analyzed the data, performed
bioinformatics analysis and drafted the manuscript. AT cultivated the plants and performed split-
root assay. MRH performed microbial co-culture experiments and extracted the DNA. MGM
interpreted the results and critically revised the manuscript.

REFERENCES

Ahmad, H. M., Fiaz, S., Hafeez, S., Zahra, S., Shah, A. N., Gul, B., Aziz, O., Mahmood-Ur-
Rahman, Fakhar, A., Rafique, M., Chen, Y., Yang, S. H., & Wang, X. (2022). Plant
growth-promoting Rhizobacteria eliminate the effect of drought stress in plants: A Review.
Frontiers in plant science, 13, 875774.

Al-Turki, A., Murali, M., Omar, A. F., Rehan, M., & Sayyed, R. Z. (2023). Recent advances in
PGPR-mediated resilience toward interactive effects of drought and salt stress in plants.
Frontiers in microbiology, 14, 1214845.

Anders, S., Pyl, P.T. & Huber, W. (2015) HTSeq–a Python framework to work with high-
throughput sequencing data. Bioinformatics, 31 (2), 166–169.

Andres, J., Arsène-Ploetze, F., Barbe, V., Brochier-Armanet, C., Cleiss-Arnold, J., Coppée, J. Y.,
Dillies, M. A., Geist, L., Joublin, A., Koechler, S., Lassalle, F., Marchal, M., Médigue, C.,
Muller, D., Nesme, X., Plewniak, F., Proux, C., Ramírez-Bahena, M. H., Schenowitz, C.,
Sismeiro, O., … Bertin, P. N. (2013). Life in an arsenic-containing gold mine: genome and
physiology of the autotrophic arsenite-oxidizing bacterium *Rhizobium* sp. NT-26. Genome
Biology and Evolution, 5(5), 934–953.

Barrow NJ, Hartemink AE (2023) The effects of pH on nutrient availability depend on both soils
and plants. Plant Soil 487:21–37.

Bhattacharyya, P. N., & Jha, D. K. (2012). Plant growth-promoting rhizobacteria (PGPR):
emergence in agriculture. World journal of microbiology & biotechnology, 28(4), 1327–
1350.

Berendsen, R. L., Pieterse, C. M., & Bakker, P. A. (2012). The rhizosphere microbiome and
plant health. Trends in Plant Science, 17(8), 478–486.

Bolger, A.M., Lohse, M. & Usadel, B. (2014) Trimmomatic: a flexible trimmer for Illumina
Sequence Data. Bioinforma. btu170.

Breia, R., Conde, A., Badim, H., Fortes, A. M., Gerós, H., & Granell, A. (2021). Plant SWEETs: from sugar transport to plant-pathogen interaction and more unexpected physiological roles. Plant Physiology, 186(2), 836–852.

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. & Holmes, S. P. (2016) DADA2: High-resolution sample inference from Illumina amplicon data. Nature Methods, 13(7), 581–583.

Chandwani, S., Dewala, S., Chavan, S. M., Paul, D., Pachaiappan, R., Gopi, M., & Amaresan, N. (2023). Complete genome sequencing of *Bacillus subtilis* (CWTS 5), a siderophore-producing bacterium triggers antagonistic potential against *Ralstonia solanacearum*. Journal of Applied Microbiology, 134(4), lxad066.

Chen W, Koide RT, Adams TS, DeForest JL, Cheng L, Eissenstat DM (2016) Root morphology and mycorrhizal symbioses together shape nutrient foraging strategies of temperate trees. Proc Natl Acad Sci USA 113(31):8741–8746

Chen, L., Ganguly, D. R., Shafik, S. H., Danila, F., Grof, C. P. L., Sharwood, R. E., & Furbank, R. T. (2023). The role of SWEET4 proteins in the post-phloem sugar transport pathway of *Setaria viridis* sink tissues. Journal of Experimental Botany, 74(10), 2968–2986.

Choudhury, F. K., Rivero, R. M., Blumwald, E., & Mittler, R. (2017). Reactive oxygen species, abiotic stress and stress combination. The Plant Journal 90(5), 856–867.

Conrath, U., Beckers, G. J., Langenbach, C. J., & Jaskiewicz, M. R. (2015). Priming for enhanced defense. Annual review of phytopathology, 53, 97–119.

Cornelis, P., Matthijs, S. (2007). *Pseudomonas* siderophores and their biological significance. In: Varma, A., Chincholkar, S.B. (eds) Microbial Siderophores. Soil Biology, vol 12. Springer, Berlin, Heidelberg.

Dahl WJ, Foster LM, Tyler RT (2012) Review of the health benefits of peas (*Pisum sativum* L.). British Journal of Nutrition 108(1):3–10.

Egamberdieva, D., Wirth, S. J., Alqarawi, A. A., Abd_Allah, E. F., & Hashem, A. (2017). Phytohormones and beneficial microbes: essential components for plants to balance stress and fitness. Frontiers in Microbiology, 8, 2104.

Feng, C., Xu, F., Li, L., Zhang, J., Wang, J., Li, Y., Liu, L., Han, Z., Shi, R., Wan, X., & Song, Y. (2023). Biological control of Fusarium crown rot of wheat with *Chaetomium globosum* 12XP1-2-3 and its effects on rhizosphere microorganisms. Frontiers in Microbiology, 14, 1133025.

Fincheira, P., & Quiroz, A. (2018). Microbial volatiles as plant growth inducers. Microbiological research, 208, 63–75.

Frey-Klett, P., Burlinson, P., Deveau, A., Barret, M., Tarkka, M., & Sarniguet, A. (2011). Bacterial-fungal interactions: hyphens between agricultural, clinical, environmental, and food microbiologists. Microbiology and molecular biology reviews: MMBR, 75(4), 583–609.

Gamalero, E., Bona, E., Todeschini, V., & Lingua, G. (2020). Saline and arid soils: impact on bacteria, plants, and their interaction. Biology, 9(6), 116.

Glick B. R. (2014). Bacteria with ACC deaminase can promote plant growth and help to feed the world. Microbiological research, 169(1), 30–39.

Gong, X., Feng, Y., Dang, K., Jiang, Y., Qi, H., & Feng, B. (2023). Linkages of microbial community structure and root exudates: Evidence from microbial nitrogen limitation in soils of crop families. The Science of the Total Environment, 881, 163536.

Gong, Z., Xiong, L., Shi, H., Yang, S., Herrera-Estrella, L. R., Xu, G., Chao, D. Y., Li, J., Wang, P. Y., Qin, F., Li, J., Ding, Y., Shi, Y., Wang, Y., Yang, Y., Guo, Y., & Zhu, J. K. (2020). Plant abiotic stress response and nutrient use efficiency. Science China. Life sciences, 63(5), 635–674.

Hartman, K., & Tringe, S. G. (2019). Interactions between plants and soil shaping the root microbiome under abiotic stress. The Biochemical Journal, 476(19), 2705–2724.

Hartmann, A., Klink, S., & Rothballer, M. (2021). Plant growth promotion and induction of systemic tolerance to drought and salt stress of plants by quorum sensing auto-inducers of the N-acyl-homoserine lactone type: recent developments. Frontiers in Plant Science, 12, 683546.

Hasan, M. R., Thapa, A., & Kabir, A. H. (2025). Iron retention coupled with trade-offs in localized symbiotic effects confers tolerance to combined iron deficiency and drought in soybean. Journal of experimental botany, eraf263. Advance online publication. https://doi.org/10.1093/jxb/eraf263

Hashem, A., Tabassum, B., & Fathi Abd Allah, E. (2019). *Bacillus subtilis*: a plant-growth promoting rhizobacterium that also impacts biotic stress. Saudi Journal of Biological Sciences, 26(6), 1291–1297.

Hazarika, D. J., Bora, S. S., Naorem, R. S., Sharma, D., Boro, R. C., & Barooah, M. (2023). Genomic insights into *Bacillus subtilis* MBB3B9 mediated aluminium stress mitigation for enhanced rice growth. Scientific Reports, 13(1), 16467.

Institute of Agricultural Resources and Regional Planning, Chinese Academy of Agricultural Sciences (IARRPCAAS). (2013). Detection and identification of specific primer pairs for Bacillus subtilis by PCR (Chinese patent CN 103409549 A). China National Intellectual Property Administration. Priority date: August 27, 2013; publication date: November 27, 2013.

Jacoby, R. P., Koprivova, A., & Kopriva, S. (2021). Pinpointing secondary metabolites that shape the composition and function of the plant microbiome. Journal of experimental botany, 72(1), 57–69.

Jamil, F., Mukhtar, H., Fouillaud, M., & Dufossé, L. (2022). Rhizosphere signaling: insights into plant-rhizomicrobiome interactions for sustainable agronomy. Microorganisms, 10(5), 899.

Jeong H, Park J, Kim H. 2013. Determination of NH□□ in environmental water with interfering substances using the modified Nessler method. Journal of Chemistry 359217. https://doi.org/10.1155/2013/359217

Julius, B. T., Leach, K. A., Tran, T. M., Mertz, R. A., & Braun, D. M. (2017). Sugar Transporters in Plants: New Insights and Discoveries. Plant & cell physiology, 58(9), 1442–1460.

Kabir AH, Paltridge NG, Able AJ, Paull JG, Stangoulis JC (2012) Natural variation for Fe-efficiency is associated with upregulation of Strategy I mechanisms and enhanced citrate and ethylene synthesis in *Pisum sativum* L. Planta 235(6):1409–1419

Khan, N., You, F. M., Datla, R., Ravichandran, S., Jia, B., & Cloutier, S. (2020). Genome-wide identification of ATP binding cassette (ABC) transporter and heavy metal associated (HMA) gene families in flax (*Linum usitatissimum* L.). BMC genomics, 21(1), 722.

Kim, S. A., & Guerinot, M. L. (2007). Mining iron: iron uptake and transport in plants. FEBS letters, 581(12), 2273–2280.

Kloepper, J. W., Ryu, C.-M., & Zhang, S. (2004). Induced systemic resistance and promotion of plant growth by *Bacillus* spp. Phytopathology, 94(11), 1259–1266.

Kobayashi, T., & Nishizawa, N. K. (2012). Iron uptake, translocation, and regulation in higher plants. Annual Review of Plant Biology, 63, 131–152.

Kryvoruchko, I. S., Routray, P., Sinharoy, S., Torres-Jerez, I., Tejada-Jiménez, M., Finney, L. A., Nakashima, J., Pislariu, C. I., Benedito, V. A., González-Guerrero, M., Roberts, D. M., & Udvardi, M. K. (2018). An iron-activated citrate transporter, *MtMATE67*, is required for symbiotic nitrogen fixation. Plant Physiology, 176(3), 2315–2329.

Ku YS, Cheng SS, Ng MS, Chung G, Lam HM (2022) The tiny companion matters: the important role of protons in active transports in plants. International Journal of Molecular Sciences 23(5):2824.

Kümmerli R. (2023). Iron acquisition strategies in pseudomonads: mechanisms, ecology, and evolution. Biometals, 36(4), 777–797.

Lassalle, F., Dastgheib, S. M. M., Zhao, F. J., Zhang, J., Verbarg, S., Frühling, A., Brinkmann, H., Osborne, T. H., Sikorski, J., Balloux, F., Didelot, X., Santini, J. M., & Petersen, J. (2021). Phylogenomics reveals the basis of adaptation of *Pseudorhizobium* species to extreme environments and supports a taxonomic revision of the genus. Systematic and Applied Microbiology, 44(1), 126165.

Lauber, C. L., Hamady, M., Knight, R., & Fierer, N. (2009). Pyrosequencing-based assessment of soil pH as a predictor of soil bacterial community structure at the continental scale. Applied and Environmental Microbiology, 75(15), 5111–5120.

Lei, M., Wang, X., Chen, K., Wei, Q., Zhou, M., Chen, G., Su, S., Tai, Y., Zhuang, K., Li, D., Liu, M., Zhang, S., & Wang, Y. (2025). Sugar transporters: mediators of carbon flow between plants and microbes. Frontiers in Plant Science, 16, 1536969.

Lemare, M., Puja, H., David, S. R., Mathieu, S., Ihiawakrim, D., Geoffroy, V. A., & Rigouin, C. (2022). Engineering siderophore production in Pseudomonas to improve asbestos weathering. Microbial Biotechnology, 15(9), 2351–2363.

Li, L., Wang, X., Li, H., Ali, M. M., Hu, X., Oelmuller, R., Yousef, A. F., Alrefaei, A. F., Liu, J., & Chen, F. (2025). The SWEET14 sugar transporter mediates mycorrhizal symbiosis and carbon allocation in Dendrobium officinale. BMC plant biology, 25(1), 416.

Livak, K.J. & Schmittgen, T.D. (2001) Methods. Vol. 25. San Diego, CA: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method; pp. 402–408.

Lugtenberg, B., & Kamilova, F. (2009). Plant-growth-promoting rhizobacteria. Annual Review of Microbiology, 63, 541–556.

Mahapatra, S., Yadav, R., & Ramakrishna, W. (2022). Bacillus subtilis impact on plant growth, soil health and environment: Dr. Jekyll and Mr. Hyde. Journal of Applied Microbiology, 132(5), 3543–3562.

Martínez-Medina, A., Fernández, I., Sánchez-Guzmán, M. J., Jung, S. C., Pascual, J. A., & Pozo, M. J. (2013). Deciphering the hormonal signalling network behind the systemic resistance induced by *Trichoderma harzianum* in tomato. Frontiers in Plant Science, 4, 206.

Meisrimler CN, Wienkoop S, Lyon D, Geilfus CM, Lüthje S (2016) Long-term High pH: tracing changes in the proteome of different pea (*Pisum sativum* L.) cultivars. Journal of Proteomics 140:13–23

Ortíz-Castro, R., Contreras-Cornejo, H. A., Macías-Rodríguez, L., & López-Bucio, J. (2009). The role of microbial signals in plant growth and development. Plant Signaling & Behavior, 4(8), 701–712.

Pantigoso, H. A., Newberger, D., & Vivanco, J. M. (2022). The rhizosphere microbiome: Plant-
microbial interactions for resource acquisition. Journal of Applied Microbiology, 133(5),
2864–2876.

Pieterse, C. M., Zamioudis, C., Berendsen, R. L., Weller, D. M., Van Wees, S. C., & Bakker, P.
A. (2014). Induced systemic resistance by beneficial microbes. Annual Review of
Phytopathology, 52, 347–375.

Poirier, W., Bouchara, J. P., & Giraud, S. (2023). Lignin-modifying enzymes in *Scedosporium*
species. Journal of Fungi (Basel, Switzerland), 9(1), 105.

Rengasamy, P. (2010). Soil processes affecting crop production in salt-affected soils. Functional
Plant Biology, 37(7), 613–620.

Rojas, C. L., Romera, F. J., Alcántara, E., Pérez-Vicente, R., Sariego, C., Garcaí-Alonso, J. I.,
Boned, J., & Marti, G. (2008). Efficacy of Fe(o,o-EDDHA) and Fe(o,p-EDDHA) isomers
in supplying Fe to strategy I plants differs in nutrient solution and calcareous soil. Journal
of Agricultural and Food Chemistry, 56(22), 10774–10778.

Rousk, J., Bååth, E., Brookes, P. C., Lauber, C. L., Lozupone, C., Caporaso, J. G., Knight, R., &
Fierer, N. (2010). Soil bacterial and fungal communities across a pH gradient in an arable
soil. The ISME Journal, 4(10), 1340–1351.

Ruffel, S., Freixes, S., Balzergue, S., Tillard, P., Jeudy, C., Martin-Magniette, M. L., van der
Merwe, M. J., Kakar, K., Gouzy, J., Fernie, A. R., Udvardi, M., Salon, C., Gojon, A., &
Lepetit, M. (2008). Systemic signaling of the plant nitrogen status triggers specific
transcriptome responses depending on the nitrogen source in *Medicago truncatula*. Plant
Physiology, 146(4), 2020–2035.

Ryu, C. M., Farag, M. A., Hu, C. H., Reddy, M. S., Wei, H. X., Paré, P. W., & Kloepper, J. W.
(2003). Bacterial volatiles promote growth in Arabidopsis. Proceedings of the National
Academy of Sciences of the United States of America, 100(8), 4927–4932.

Sainz-Mejías, M., Jurado-Martín, I., & McClean, S. (2020). Understanding *Pseudomonas*
aeruginosa-Host Interactions: The Ongoing Quest for an Efficacious Vaccine. Cells, 9(12),
2617.

Saleem, A., A. Zulfiqar, M. Z. Saleem, et al., 2023. Alkaline and acidic soil constraints on iron
accumulation by rice cultivars in relation to several physio□biochemical parameters. BMC
Plant Biology 23(1): 397.

Sharma, A., Chakdar, H., Vaishnav, A., Srivastava, A. K., Khan, N., Bansal, Y. K., & Kaushik,
R. (2023). Multifarious plant growth-promoting *Rhizobacterium enterobacter* sp. CM94-
mediated systemic tolerance and growth promotion of chickpea (Cicer arietinum L.) under
salinity stress. Frontiers in Bioscience (Landmark edition), 28(10), 241.

Singh, J., Aggarwal, R., Bashyal, B. M., Darshan, K., Parmar, P., Saharan, M. S., Hussain, Z., &
Solanke, A. U. (2021). Transcriptome reprogramming of tomato orchestrate the hormone
signaling network of systemic resistance induced by *Chaetomium globosum*. Frontiers in
Plant Science, 12, 721193.

Tang, J., Li, W., Wei, T., Huang, R., & Zeng, Z. (2024). Patterns and mechanisms of legume
responses to nitrogen enrichment: A global meta-analysis. Plants (Basel, Switzerland),
13(22), 3244.

Tarafdar, J. C., & Gharu, A. (2006). Mobilization of organic and poorly soluble phosphates by
Chaetomium globosum. Applied Soil Ecology, 32(3), 273–283.

Thapa, A., Hasan, M. R., & Kabir, A. H. (2025a). *Trichoderma afroharzianum* T22 Induces
Rhizobia and Flavonoid-Driven Symbiosis to Promote Tolerance to Alkaline Stress in
Garden Pea. Plant, cell & environment, 10.1111/pce.15581.
Thapa, A., Hasan, M. R., & Kabir, A. H. (2025b). Transcriptional reprogramming and
microbiome dynamics in garden pea exposed to high pH stress during vegetative stage.
Planta, 261(4), 83.
Thilakarathna, M. S., & Cope, K. R. (2021). Split-root assays for studying legume-rhizobia
symbioses, rhizodeposition, and belowground nitrogen transfer in legumes. Journal of
experimental botany, 72(15), 5285–5299.
Toju, H., Peay, K. G., Yamamichi, M., Narisawa, K., Hiruma, K., Naito, K., Fukuda, S., Ushio,
M., Nakaoka, S., Onoda, Y., Yoshida, K., Schlaeppi, K., Bai, Y., Sugiura, R., Ichihashi, Y.,
Minamisawa, K., & Kiers, E. T. (2018). Core microbiomes for sustainable agroecosystems.
Nature Plants, 4(5), 247–257.
Vejan, P., Abdullah, R., Khadiran, T., Ismail, S., & Nasrulhaq Boyce, A. (2016). Role of Plant
Growth Promoting Rhizobacteria in Agricultural Sustainability-A Review. Molecules
(Basel, Switzerland), 21(5), 573.
Verma, K. K., Joshi, A., Song, X. P., Liang, Q., Xu, L., Huang, H. R., Wu, K. C., Seth, C. S.,
Arora, J., & Li, Y. R. (2024). Regulatory mechanisms of plant rhizobacteria on plants to
the adaptation of adverse agroclimatic variables. Frontiers in plant science, 15, 1377793.
Vishwakarma, K., Kumar, N., Shandilya, C., Mohapatra, S., Bhayana, S., & Varma, A. (2020).
Revisiting Plant-Microbe Interactions and Microbial Consortia Application for Enhancing
Sustainable Agriculture: A Review. Frontiers in Microbiology, 11, 560406.
Wang, E., Yu, N., Bano, S. A., Liu, C., Miller, A. J., Cousins, D., Zhang, X., Ratet, P., Tadege,
M., Mysore, K. S., Downie, J. A., Murray, J. D., Oldroyd, G. E., & Schultze, M. (2014). A
H^+-ATPase that energizes nutrient uptake during mycorrhizal symbioses in rice and
Medicago truncatula. The Plant Cell, 26(4), 1818–1830.
Wang, L., Ju, C., Han, C., Yu, Z., Bai, M. Y., & Wang, C. (2025). The interaction of nutrient
uptake with biotic and abiotic stresses in plants. Journal of Integrative Plant Biology, 67(3),
455–487.
Zamioudis, C., & Pieterse, C. M. (2012). Modulation of host immunity by beneficial microbes.
Molecular Plant-Microbe Interactions, 25(2), 139–150.
Zhang JC, Wang XN, Sun W, Wang XF, Tong XS, Ji XL, An JP, Zhao Q, You CX, Hao YJ
(2020) Phosphate regulates malate/citratemediated Fe uptake and transport in apple. Plant
Science 297:110526
Zhang, X. G., Guo, S. J., Wang, W. N., Wei, G. X., Ma, G. Y., & Ma, X. D. (2020). Diversity
and bioactivity of endophytes from *Angelica sinensis* in China. Frontiers in Microbiology,
11, 1489.
Zuluaga, M. Y. A., M. Cardarelli, Y. Rouphael, S. Cesco, Y. Pii, and G. Colla. 2023. Iron
nutrition in agriculture: from synthetic chelates to biochelates. Scientia Horticulturae 312:
111833.

973 **Table 1.** Genotypic screening of garden pea capable of utilizing *B. subtilis* (BS) to cope with soil
974 alkalinity. Data represent means ± SDs of three independent biological samples. Statistically
975 significant differences among different treatments are indicated by different alphabetical letters
976 at the *P* < 0.05 level.

Genotypes	Shoot height (cm)				SPAD score			
	Control	Alkaline	Alkaline+BS	BS+	Control	Alkaline	Alkaline+BS	BS+
Sugar Snap	82±2.5a	56±3.2b	78±1.5a	83±4.1a	33±0.76a	23±2.17b	32±2.47a	34±1.86a
Oregon Sugar Pod	24±1.7a	14±3.3b	22±1.8a	25±0.57a	33±1.4a	23±1.9b	34±2.3a	34±2.2a
Sugar Ann	26±2.0a	21±0.57b	24±0.57a	32±3.6ac	39±1.9a	33±2.7b	36±1.7b	36±2.4b
Little Marvel	15±0.90a	7±2.4b	12±1.3a	22±5.8c	47±2.0a	31±1.8b	36±2.2c	44±2.5a
Green Arrow	24±1.1a	19±1.9b	20±1.7b	28±3.0a	47±1.7a	39±1.9b	42±1.5b	48±1.3a
Lincoln	19±1.0a	15±0.57b	14±2.0b	19±1.5a	34±0.36a	30±1.5b	31±1.5b	37±3.8a

977
978 **Table 2.** Changes in nutrient concentrations in roots and leaves of pea plants inoculated with or
979 without *B. subtilis* (BS) under control and alkaline soil conditions. Data represent means ± SDs
980 of three independent biological samples. Statistically significant differences among different
981 treatments are indicated by different alphabetical letters at the *P* < 0.05 level.

Elements	Roots				Leaves			
	Control	Alkaline	Alkaline+BS	BS+	Control	Alkaline	Alkaline+BS	BS+
Fe (ppm)	1011±72.8a	660±96.8b	997±130.4a	999±42.1a	202±33.1a	137±22.0b	205±17.6a	196±23.2a
Mn (ppm)	70±5.5a	39±11.6b	100±25.0a	114±27.3a	49±7.1a	31±3.6b	37±5.8a	46±10.6a
S (%)	0.10±0.01a	0.09±0.01a	0.10±0.005a	0.09±0.01a	0.22±0.3a	0.13±0.01b	0.23±0.02a	0.24±0.01a
Mg (%)	0.29±0.05a	0.16±0.01b	0.36±0.07a	0.28±0.05a	0.15±0.2a	0.09±0.01a	0.14±0.02a	0.16±0.03a
Ca (%)	0.76±0.08a	0.51±0.04b	1.01±0.03a	0.99±0.09a	0.28±0.03a	0.18±0.02b	0.49±0.06c	0.53±0.11c
N (%)	3.2±0.26a	2.4±0.18b	3.2±0.22a	3.4±0.57a	7.0±1.19a	5.5±1.33b	7.1±0.68a	7.6±1.04a

Fig. 1. Effect of *Bacillus subtilis* on plant phenotypes (A), relative abundance of *B. subtilis* (BS) in the roots (B), plant fresh weight (C), plant height (D), root length (E), root fresh weight (F), root ammonium content (G), F_v/F_m: maximum efficiency of PSII (H), and PI_ABS:

photosynthetic performance index (I) in pea plants grown under four treatments: control, alkaline (pH 7.8), alkaline+BS, and BS+. Data represent means ± SD of three biological replicates ($n = 5$) on an independent cultivation. Different letters above the box plots indicate statistically significant differences among treatments ($p < 0.05$, one-way ANOVA followed by post hoc test).

Fig. 2. Growth performance and physiological responses of pea plants under different treatments (T1: control, T2: alkaline (pH 7.8), T3: alkaline + *B. subtilis* BS, T4: alkaline + inorganic Fe, T5: BS+, and T6: inorganic Fe) on phenotype of plants from each treatment (A), the relative abundance of *B. subtilis* in the roots (B), rhizosphere siderophore % (C), number of root nodules (D), plant height (E), and leaf SPAD score (F). Data represent means ± SD of three biological replicates ($n = 5$). Different letters above the box plots indicate statistically significant differences among treatments ($p < 0.05$, one-way ANOVA followed by post hoc test).

1026

1027
Fig. 3. Effects of different split-root treatments (SR1: control/control, SR2: alkaline/alkaline,
SR3: alkaline/alkaline+BS, SR4: alkaline+BS/alkaline+BS, SR5: control/control+BS, SR6:
control+BS/control+BS) on root length (A), root fresh weight (B), shoot height (C), shoot fresh
weight (D), SPAD score (E), and F_v/F_m: maximum efficiency of PSII (F) of pea plants cultivated
in two compartments (com 1 and com 2). In treatments involving bacterial inoculation,
compartment 1 received the *B. subtilis* inoculum, while compartment 2 remained uninoculated.

Data are means ± SD from three biological replicates ($n = 5$), with letters indicating statistical
differences based on one-way ANOVA followed by post hoc tests ($p < 0.05$).
1036

1037
1038 **Fig. 4.** Effect of single and co-culture growth on colony size of *R. leguminosarum* and *B. subtilis*
1039 under control (A) and alkaline conditions, pH 7.8 (B) over 3, 5, and 7 days. Data are means ± SD
1040 from three biological replicates ($n = 5$), with significance (* = $P < 0.05$, ns = not significant)
1041 determined by *t*-test comparison between single and co-culture within each time point.

1042
1043
1044
1045

A

PC2: 16.2% Variance

PC1: 29.5% Variance

Names
- Control
- Alkaline
- Alkaline+BS
- BS+

B

Up (Alkaline vs Control)   Up (Alkaline+BS vs Alkaline)

300   164   792

90   0   2

Up (BS+ vs Alkaline+BS)

C

Down (Alkaline vs Control)   Down (Alkaline+BS vs Alkaline)

248   26   1106

19   0   2

Down (BS+ vs Alkaline+BS)

D Symbiotic association

PSAT4G075040 (Sugar phosphate transporter)
PSAT1G041560 (Glucose transporter GLUT)
PSAT5G053200 (Sugar Transporter SWEET)
PSAT1G124920 (MF sugar transporter−like)
PSAT4G226520 (Sugar transport protein STP)
PSAT5G084360 (UDP−sugar pyrophosphorylase)

BS.vs.Alkaline.BS
Alkaline.vs.Control
Alkaline.BS..vs.Alkaline

E Nutrient transport and assimilation

PSAT0S1495G0040 (Ammonium transporter)
PSAT7G123960 (Nitrite and sulphite reductase)
PSAT2G157960 (Inorganic pyrophosphatase)
PSAT7G178640 (Amino acid transporter)
PSAT6G052400 (Cys/Met metabolism)
PSAT5G102080 (Iron/zinc purple acid phosphatase)
PSAT5G020560 (Malic oxidoreductase)
PSAT2G074400 (SulP transporter)
PSAT7G213080 (Zinc/iron permease)
PSAT2G173680 (MFS transporter)

BS.vs.Alkaline.BS
Alkaline.vs.Control
Alkaline.BS..vs.Alkaline

F Redox and osmotic adjustment

PSAT1G187800 (Glutaredoxin)
PSAT6G021760 (Oxidative Stress Regulator)
PSAT3G009360 (Glutathione synthase)
PSAT3G052760 (Thioredoxin)
PSAT7G068360 (Glutathione S−transferase)
PSAT6G040040 (Ferritin−like superfamily)
PSAT1G038760 (Dehydrin)
PSAT1G118120 (Aquaporin transporter)

Alkaline.BS..vs.Alkaline
Alkaline.vs.Control
BS.vs.Alkaline.BS

G Growth and stress response

PSAT2G011760 (Isoprenoid synthase)
PSAT5G063960 (Ethylene−responsive TF)
PSAT3G088320 (Cation/H+ exchanger)
PSAT2G058680 (ABC−2 type transporter)
PSAT3G053400 (ABC transporter type 1)
PSAT0S1027G0040 (Root meristem growth factor 3)
PSAT6G191920 (AP2/ERF Ethylene Responsive Factor)
PSAT6G045440 (Oxoglutarate)
PSAT3G001040 (Small auxin−up RNA)
PSAT7G218840 (ATPase)

BS.vs.Alkaline.BS
Alkaline.vs.Control
Alkaline.BS..vs.Alkaline

1046
Fig. 5. Transcriptomic changes in pea roots under control, alkaline, alkaline + *B. subtilis*
(alkaline+BS), and BS+ treatments. (A) Principal coordinate analysis (PCoA) plot showing
separation of transcriptome profiles among treatments, (B) Venn diagram of upregulated genes,
(C) Venn diagram of downregulated genes, (D–G) Heatmaps showing selected differentially
expressed genes (DEGs) grouped by functional categories. Color scale represents normalized

expression values (log$_2$ fold change), with red and blue indicating upregulation and
downregulation, respectively. Data are based on three independent biological replicates per
treatment ($n = 3$).
1055

1056
Fig. 6. Gene ontology (GO) enrichment analysis of differentially expressed genes in the
alkaline+BS *vs.* alkaline comparison using ShinyGO. (A) Top enriched GO terms for
upregulated genes and (B) Top enriched GO terms for downregulated genes associated with
biological process. Bubble size represents the number of genes in each GO term, color indicates
log10(FDR), and the x-axis shows fold enrichment. Data are based on three biological replicates
per treatment ($n = 3$).
1063

Fig. 7. Rhizosphere bacterial community structure in pea plants under different treatments: control, alkaline, alkaline+*B. subtilis* (BS), and BS+. Data represent (A) principal coordinate analysis (PCoA) plot based on Bray–Curtis dissimilarity showing separation of bacterial communities among treatments, (B) alpha diversity indices (Observed richness and Simpson index; box plots show median, interquartile range, and individual replicates, (C) relative abundance of dominant bacterial families, and (D) relative abundance of dominant bacterial genera. Asterisks (*) indicate taxa that showed statistically significant differences among treatments based on Kruskal-Wallis ($P < 0.05$). Data are from three biological replicates per treatment ($n = 3$).

1075
Fig. 8. Rhizosphere fungal community structure in pea plants under different treatments: control,
alkaline, alkaline+*B. subtilis* (BS), and BS+. Data represent (A) principal coordinate analysis
(PCoA) plot based on Bray–Curtis dissimilarity showing separation of bacterial communities
among treatments, (B) alpha diversity indices (Observed richness and Simpson index); box plots
show median, interquartile range, and individual replicates, (C) relative abundance of dominant
fungal families, and (D) relative abundance of dominant fungal genera. Asterisks (*) indicate
taxa that showed statistically significant differences among treatments based on Kruskal-Wallis
(*P* < 0.05). Data are from three biological replicates per treatment (*n* = 3).
1084
1085

1086

1087

Fig. 9. Mechanistic model of *B. subtilis*-mediated strategies enhancing pea plant tolerance to
alkaline stress. BS inoculation promotes colonization of helper microbes (*Pseudorhizobium,*
Pseudallescheria, and *Chaetomium*), which establish mutualistic interactions with the host by
exchanging sugars through SWEET and GLUT transporters. These interactions support
complementing rhizobia, enhancing nodulation through ammonium transporters and nitrogen
assimilation. In parallel, cation/H$^+$ exchangers, ATPases, and Zn/Fe permeases facilitate uptake
of essential micronutrients (Fe, Mn, Ca). Together with systemic signaling, these mechanisms
contribute to improved nutrient homeostasis and plant growth under alkaline conditions.

1096
1097