Big angiosperm genera are priorities for conservation

Matilda J.M. Brown[1]*, Tarciso C.C. Leão[1]*, Félix Forest[1], Eve Lucas[1], Barnaby E. Walker[1], Eimear Nic Lughadha[1]

[1]Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE, UK

*These two authors contributed equally to this work

Authors for correspondence:
Matilda Brown m.brown2@kew.org
Eimear Nic Lughadha e.niclughadha@kew.org

ORCID:
Matilda J.M. Brown   0000-0003-2536-8365
Tarciso C.C. Leão   0000-0002-7029-1582
Félix Forest   0000-0002-2004-433X
Eve Lucas   0000-0002-7603-435X
Barnaby E. Walker   0000-0002-3884-671X
Eimear Nic Lughadha 0000-0002-8806-4345

Abstract

Species in big angiosperm genera (≥500 species) tend to be less well-known than those in smaller genera. Consequently, species in big genera are underrepresented in conservation at all levels, from analyses to actions. However, emerging evidence indicates that species in larger genera tend to have smaller geographic ranges (a strong correlate of increased extinction risk). Here, we test the global generality of this pattern, and the relationships between plant taxon size, extinction risk and climatic zone on a global scale. Species in larger genera are more likely to have small ranges, less likely to have a global Red List assessment, but are more likely to be threatened. Persistent obstacles to improving conservation documentation of big genera include taxonomic uncertainty, data shortfalls and number of species involved. We highlight the need for closer collaboration between taxonomy and conservation to enable effective conservation of the quarter of plant species that big genera encompass.

Key words: conservation, big genera, genus size, geographic range size, extinction risk, IUCN Red List, angiosperms

Introduction

'Big' plant genera (≥500 species *per* Frodin, 2004) contain over a quarter of angiosperm diversity (Moonlight *et al.*, 2024) and include several ecologically, culturally and economically important taxa (e.g. *Euphorbia*, >2,000 spp.; *Solanum*, >1,200 spp., *Eucalyptus*, >700 spp.). Despite the familiarity of many big plant genera, there remains a

pervasive knowledge shortfall for species in big genera (Moonlight *et al.*, 2024), largely due

to the practical challenges associated with working with groups of ≥500 species. For

example, big genera are less likely to have a global taxonomic monograph and have lower

proportions of species with available DNA sequence data (Grace *et al.*, 2021). Longstanding

latitudinal biases in taxonomic effort (the 'Latitudinal Taxonomy Gradient'; (Freeman &

Pennell, 2021; Guedes *et al.*, 2025) mean that this shortfall is particularly severe in the

tropics, where the majority of big plant genera occur (Moonlight *et al.*, 2024).

Both the practical and taxonomic challenges associated with big plant genera limit our

knowledge of extinction risk in these taxa; for example, <5% of the >3,000 species of

Astragalus have global Red List assessments (far fewer than the c. 20% coverage of

angiosperms at large; IUCN, 2024). Undertaking a fully-documented species conservation

assessment (e.g. a Red List assessment; IUCN, 2012; IUCN, 2022a) is expensive in terms of

time and expertise. Thus, data-poor and taxonomically uncertain species, which are likely to

be assessed as Data Deficient (Brummitt *et al.*, 2024) or re-circumscribed (requiring re-

assessment, as assessments are specific to a particular species concept) tend to be low

priorities for assessment. Assessments are also typically undertaken as part of projects of

clearly defined geographic and/or taxonomic scope, often aiming to assess all species within

that scope (e.g. the recent, comprehensive assessment of *Dalbergia*; Cowell, 2022).

Comprehensive assessment of species in a big genus requires substantial resources, and

usually many collaborators. The successful Global Tree Assessment (>45,000 species; Rivers,

2017) demonstrates the feasibility of large assessment projects, but also the enormity of the

challenges involved (Newton *et al.*, 2015). We expect that the sheer numbers of species and

the many uncertain species circumscriptions in big genera mean that they are systematically

underrepresented on the global Red List, limiting our knowledge of extinction risk in a

substantial, non-random portion of angiosperm diversity.

There is also emerging evidence that we may be underestimating extinction risk in big plant

genera. Geographic range size is the strongest correlate of plant extinction risk (e.g. Nic

Lughadha *et al.*, 2020; Walker *et al.*, 2022), and both region-specific and flora-based studies

have found that taxa with larger numbers of species have, on average, species with smaller

geographic ranges (Schwartz & Simberloff, 2001; Lozano & Schwartz, 2005; Leão *et al.*,

2020) and higher proportions of threatened species (Fu *et al.*, 2022). Furthermore, a

disproportionate number of species described in the last two decades – the majority of

which are likely to be threatened (Brown *et al.*, 2023a) – are in the very largest genera

(>1,000 species; Moonlight *et al.*, 2024). The global generality of the relationships between

genus size, range size and extinction risk in flowering plants remain untested, particularly

across climatic zones, lifeforms and higher taxonomic ranks, but we expect that an overall

genus size - range size relationship, if supported, will also result in higher proportions of

threatened species in larger genera.

Here, we test the generality of the negative relationship between genus size and species

geographic range size (as number of botanical countries; Brummitt *et al.*, 2001) across

climate, life form and family, using data from the World Checklist of Vascular Plants

(Govaerts *et al.*, 2021). We then quantify the extent to which species in larger genera are

more or less likely to have a Red List assessment, and their probability of, when assessed,

being categorised as either Data Deficient or threatened with extinction, using data from

the IUCN Red List (version 2024-1; IUCN, 2024) and comprehensive species-level extinction

risk predictions from Bachman *et al.* (2024). Finally, we identify specific priority big genera

for conservation assessment as those for which overall extinction risk on the Red List is most

underestimated when compared to predictions.

Results

Nearly half of all big genera (and species therein) are tropical (Fig 1, Table S1). Of the 82

currently accepted big genera (from version 13 of the WCVP), 40 (49%) are entirely or

mostly tropical, containing 40,631 species (48% of species in big genera; 13% of total

angiosperm diversity). All big epiphytic genera (10 genera; 11,370 species) are

predominantly tropical; similarly, all big tree genera are either tropical (4 genera; 3,211

species) or subtropical (*Eucalyptus*, 713 species). By contrast, most of the 15 temperate big

genera are dominated by herbaceous perennials (12 genera; 16,239 species). There are no

big genera dominated by annual species.

Figure 1. Climate and lifeform distribution in big angiosperm genera. Genera coded by their

dominant (≥75% of species) climate/lifeform or as 'multiple climates/lifeforms' (see Methods). Colour

and pattern show proportion of species across covariate combinations; note that overall circle sizes

are not to scale. For full details on numbers of species and genera across covariates, see Table S1.

Covariates of genus size trends

The best-supported model formula (ranked using AIC; Table S2) across all response variables

includes both additive and interaction terms for all covariates (climate, lifeform, family).

Family-level interaction terms could not be estimated for families with very few species or

genera; the results presented below are thus based on a reduced dataset (91.8% of

angiosperm species; see Methods). Models fitted using the full dataset (excluding family

effects) are consistent with the findings reported below (Tables S3-4).

Genus size and species range size

Overall, species in big genera tend to have smaller ranges (Fig 2). The probability of being

restricted to a single botanical country is 49.7% (95% CI: 49.2-50.2%) for a species in a

monotypic genus, compared to 60.6% (CI: 60.3-60.9%) for species in the 'smallest' big

genera (i.e. genera of exactly 500 species; from marginal mean estimate; Fig 2a).

The relationship between genus size and range size is consistent across scales (i.e. when

modelling the proportion of species restricted to one, two, three, four, five and ten

botanical countries, Fig 2a), climatic zones and lifeform (Fig 2b). It is strongest for annual,

epiphytic, temperate, and subtropical genera, and comparatively weak (though still

significantly positive) for tropical and woody (tree, other woody perennial) genera (Fig 2b).

Across families included in this analysis (100 families; 91.8% of species), the genus size -

range size relationship was mixed (Fig 2c); 54 families have significant ($p<0.05$), positive

slope coefficients (i.e. species in larger genera are more likely to be restricted to a single

botanical country), five families have significant negative coefficients and 41 families had

slope coefficients not significantly different from zero.

Figure 2. Genus size-range size relationship. Panel a) shows estimated marginal mean proportions of
species under each range size threshold (as number of botanical countries) across genus size. The
median range size is one botanical country for all but the very smallest genera. Panel b) shows the
estimated marginal mean probability that species are restricted to a single botanical country across
lifeform and climate. Shaded grey area shows the 'zone of big genera'; points show the observed
proportions of single-country endemics in big genera. Panel c) shows the estimated marginal
coefficients for family (the single-country endemic threshold; only largest 50 families shown); positive
coefficients indicate that species in larger genera are more likely to be restricted to a single botanical
country. Units of coefficients on the logit scale. Error bars show 95% CIs of coefficient estimates;
colour of points denotes significance and direction; size of points denotes number of species per
family. Marginal means and trends estimated across cell weights of the non-focal terms in all panels.
For tabulated results with 95% CI values for climate, lifeform and overall marginal trends, see Table
S3.

Genus size and extinction risk

Species in big genera are less likely to have a global Red List assessment (Fig 3); for a monotypic genus, the probability of being assessed is 15.3% (95% CI: 14.4-16.2%), compared to 11.5% (CI: 11.2-11.7%) for a species in a 500-species genus (estimated marginal means).

Marginal trends across lifeform and climate are generally consistent with the overall effect, except for subtropical genera – species in these genera are more likely to be assessed if they are in a larger genus (Fig 3a). This result is likely driven by the near-comprehensive (94%) assessment of *Eucalyptus* (Myrtaceae), the only large subtropical tree genus, comprising 27.0% of species (across all lifeforms) in big subtropical genera. The impact of the Global Tree Assessment is clearly visible in our data, with trees having the highest assessment coverage of any covariate (Fig 3a).

Marginal trends across families are less consistent: 49 families have significant negative trends as expected, 19 have positive trends and 32 are not significantly different from zero (50 largest families shown in Fig 3b). Regardless of lifeform, species in tropical genera are always more likely to have been assessed than for other climates (Fig. 3a).

Figure 3. Genus size-assessment probability relationship. Panel a) shows estimated marginal means
(i.e. predictions) of assessment probability across genus size, climate and lifeform. Upper limit of
genus size values restricted to range of data for each covariate combination. Grey rectangle
highlights the 'zone of big genera'; points show the observed proportions of assessed species in big
genera. Dashed lines show n.s. trends; solid lines show trends statistically different from zero. Panel
b) shows the estimated marginal coefficients for family (the single-country endemic threshold; only
largest 50 families shown); negative coefficients indicate that species in larger genera are less likely
to have a Red List assessment. Units of coefficients on the logit scale. Error bars show 95% CIs of
coefficient estimates; colour of points denotes significance and direction; size of points denotes
number of species per family. Marginal means and trends estimated across cell weights of the non-
focal terms in all panels. For tabulated results with 95% CI values for climate, lifeform and overall
marginal trends, see Table S3.

Using Red List data, we found that species in big genera are slightly more likely to be

assessed as threatened (either Vulnerable, Endangered or Critically Endangered; Fig 4a). On

average, an assessed species in a monotypic genus has a 32.6% (CI: 31.0-34.1%) chance of

being assessed as threatened, compared to 37.5% (CI: 36.8-38.2%) if it is in a genus of 500

species (estimated marginal means). The effect is generally reversed in epiphytic and

subtropical genera (Fig 4b), though this is likely driven by a small number of assessments, as

174 species from epiphytic or subtropical genera tend to be less likely to have assessments (Fig

175 3). Across the 51 families with sufficient assessment coverage for analysis (see Methods; Fig

176 4c), 13 have a significant positive relationship as expected, 9 have a significant negative

177 relationship and 29 have relationships between genus size and assessment probability that

178 are not significantly different from zero.

Figure 4. Genus size - documented extinction risk relationship (Red List assessments only). Panel a)
shows estimated marginal means for the probability of meeting the criteria for each Red List
category (note that categories are cumulative, e.g. all species assessed as Critically Endangered also
meet the criteria for Endangered and Vulnerable. Panel b) shows estimated marginal means of being
threatened (Vulnerable, Endangered or Critically Endangered) across genus size, climate and lifeform.
Upper limit of genus size values restricted to range of data for each covariate combination. Grey
rectangle highlights the 'zone of big genera'; points show the observed proportions of species
assessed as threatened in big genera. Dashed lines show n.s. trends; solid lines show trends
statistically different from zero. Panel c) shows the estimated marginal coefficients for family (only

Species in larger genera are also more likely to be categorised as Data Deficient (Fig. 5); the

probability that a species in a monotypic genus is categorised as Data Deficient is 4.5% (95%

CI:4.1-4.9%), compared to 9.1% (CI: 8.7-9.4%) in a genus with 500 species. The effect on

Data Deficiency of being in a large genus is strongest for tropical and epiphytic species, but

reversed for annual and subtropical species (i.e. species in larger genera are less likely to be

assessed as Data Deficient), though these groups have, respectively, only 18 and 94 species

(zero and seven in big genera) that have been assessed as Data Deficient. Data were

insufficient to fit the full model including family; full model for Data Deficiency fit with

climate and lifeform covariates only.

Figure 5. Genus size – Data Deficiency relationship (Red List assessments only). Estimated marginal
means of being assessed as Data Deficient across genus size, climate and lifeform. Upper limit of
genus size values restricted to range of data for each covariate combination. Note that y-axes are

When we extended our analysis of extinction risk to include all angiosperms, by including species with extinction risk predictions (from Bachman *et al.* 2024), we observed a stronger overall relationship between genus size and extinction risk (Fig 6): a species in a monotypic genus has a 30.4% (CI: 29.8-31.1%) chance of being threatened (assessed or predicted with high confidence), compared to 45.6% (CI: 45.3-45.9%) for a genus of 500 species (estimated marginal means).

The relationship between genus size and probability of being threatened is consistent across climatic zones and lifeforms (Fig 6b). It is strongest for annual, epiphytic and genera found in multiple climates, and comparatively weak (though still significantly positive) for tropical, subtropical and woody genera (Fig 6b). Across families (Fig 6c), the genus size – extinction risk relationship was broadly consistent, though statistical power was limited; of the 100 families included in this analysis, 56 have significant ($p<0.05$), positive slope coefficients (i.e. species in larger genera are more likely to be either assessed as or predicted to be threatened), five have significant negative coefficients and 39 have slope coefficients that are not significantly different from zero. When we compared estimates of the probability that a species is threatened based on assessments versus assessments plus predictions (Extended Data Figure 1), we found strong evidence to suggest that current Red List data underestimate the proportion of species that are threatened with extinction in genera of eleven or more species, and that this underestimation is most severe for big genera.

Figure 6. Genus size - extinction risk relationship (Red List assessments and extinction risk predictions
from Bachman et al. 2024). Panel a) shows estimated marginal means across all thresholds of
estimated extinction risk. Panel b) shows estimated marginal means of being threatened (assessed or
predicted with high confidence) across genus size, climate and lifeform. Upper limit of genus size
values restricted to range of data for each covariate combination. Grey rectangle highlights the 'zone
of big genera'; points show the proportions of species in big genera that are assessed or predicted to
be threatened. Dashed lines show n.s. trends; solid lines show trends that are statistically different
from zero. Panel c) shows the estimated marginal coefficients for family (only largest 50 families
shown); positive coefficients indicate that species in larger genera are more likely threatened (either
assessed or predicted). Units of coefficients on the logit scale. Error bars show 95% CIs of coefficient
estimates; colour of points denotes significance and direction; size of points denotes number of
species per family. Marginal means and trends estimated across cell weights of the non-focal terms
in all panels. For tabulated results with 95% CI values for climate, lifeform and overall marginal
trends, see Table S3.

From the extinction risk predictions, 37,072 species in big genera are estimated (with high

confidence) to be threatened with extinction, but either lack a global Red List assessment

(36,411) or have been assessed as Data Deficient (661) – together representing 10.9% of all

angiosperm species. This number is conservative, as >13,000 species have been described or

otherwise added to the World Checklist of Vascular Plants since the release used as the

backbone for predictions. Of the species in big genera predicted to be threatened, 19,787

are in tropical genera (spanning 40 genera). We quantified the degree to which extinction

risk is likely to be underestimated in each large genus by subtracting the Red-List-derived

proportion of threatened species from that calculated using predictions (Extended Data

Table 1). Of the twenty big genera in which extinction risk is likely to be most

underestimated on the Red List, six are tropical genera in the family Orchidaceae. The most

extreme example is *Stelis* with <1% of species having a Red List assessment and none

threatened, while 79% of species are predicted to be threatened.

Discussion

Consistent with our expectations, species in big genera are more likely to have small ranges,

less likely to have Red List assessments, and if assessed, they are more likely to be assigned

to one of the threatened categories or categorised as Data Deficient. The relationships

between genus size, range size and extinction risk are broadly consistent across climatic

zones, lifeforms, and families; as over 25% of plant species are in 'big' genera (≥500 species)

and a further 35% are in moderately large genera (≥100 species), our findings are relevant

to the majority of known plant species and have significant implications for conservation

science, action and policy.

Our results support previous studies on the relationship between genus size and range size

in plants (Schwartz & Simberloff, 2001; Lozano & Schwartz, 2005; Leão *et al.*, 2020; family-

level Fu *et al.*, 2022), contrasting with earlier reports for both plants (Vamosi & Wilson,

2008) and animals (Purvis *et al.*, 2000) that smaller – especially monotypic – taxa tend to

have higher levels of extinction risk. However, the relationship between genus and range

size is not universal – in five families, species in larger genera are less likely to be restricted

to a single botanical country, contrary to expectations (Table S5). These families span seven

orders (both eudicots and monocots), vary in climate, lifeform and range sizes, and three

contain at least one big genus, so there is no clear pattern to these outliers. Similarly, we

found groups for which the general pattern of higher extinction risk in larger genera is

reversed, where species in larger genera are *less* likely to be either assessed as or predicted

to be threatened (Figs 3-4; Tables S3, S5).

Results based on published assessments were less consistent than those based on

predictions, likely due to the very low assessment coverage of many big genera. For

example, in tropical epiphytes, where seven of the ten big genera have fewer than 50

published assessments (representing <3% of species in these seven genera), assessments

show an apparent negative relationship between genus size and extinction risk (Fig 4), but

the model including predictions showed the expected positive trend (Fig 5). There are five

families with a negative relationship between genus size and extinction risk in prediction-

based models. For one of these, Melastomataceae, this may be linked to range, as it was

one of the five families where species in larger genera tended to have larger ranges (Table

S5), but for the other four the reasons for this result are unclear; differences in both range

size and extinction risk may be linked to the 'different ways to be a big genus' (an

outstanding knowledge gap noted by Moonlight *et al.*, 2024). It is also possible that there

are biases in the predictions that affect our results – this does not in any way diminish the

need for species in big genera to be prioritised for conservation assessment, whether to

formally document their predicted extinction risk, address the gaps in both knowledge and

data for big genera or to better understand the drivers of extinction risk across taxon size.

The mechanisms underpinning the link between genus and range sizes may be related to

evolutionary diversification patterns. More rapidly diversifying plant genera tend to have

smaller ranges and higher extinction risk (Davies *et al.*, 2011; Leão et al., 2020; Tanentzap et

al,. 2020), though conifers showed the opposite trend (Tanentzap et al. 2020). In animals,

genus-level diversification rate appears to be a strong correlate of both range size and

extinction risk (though there is significant variation across clades; Greenberg *et al.*, 2021),

further supporting the hypothesis that rapid speciation tends to produce range-restricted

species which are inherently more vulnerable to extinction (Davies *et al.*, 2011; Leão *et al.*,

2020).

Our results complement recent studies characterising biases in the representation of plants

on the Red List and addressing how these can be quantified and corrected (Nic Lughadha *et*

al., 2020) or circumvented (e.g. Bachman *et al.*, 2024) in future analyses to refine

understanding of global and regional patterns of plant extinction risk. For example, eight of

the 20 most under-estimated genera (when overall extinction risk is estimated using the Red

List; Table 1) are Orchidaceae (6 genera) or Asteraceae (2 genera). These two largest plant

families encompass c. 65,000 species and are recognised as being among the most under-

represented on the Red List (Nic Lughadha *et al.*, 2020).

Strikingly, the six orchid genera highlighted here for their underestimated threat levels are

all tropical, consistent with our original expectation that under-representation of big genera

in assessment processes might be greatest in the tropics. However, contrary to our

expectations, we found that species in tropical genera, across all lifeforms, were more likely

to have been assessed than species of other climates (Fig 3a). This encouraging result

suggests significant progress towards addressing the plant Red Listing deficit which was so

evident across many areas of the tropics as recently as a decade ago (Bachman *et al.*, 2019).

Much of this progress is attributable to the Global Tree Assessment, and the high

assessment coverage achieved for trees explains the relatively weak effect of genus size on

assessment in tree-dominated genera. Similarly concerted and large-scale endeavours are

urgently needed for other plants, and our findings suggest that an explicit focus on

comprehensive assessments of selected big genera could be an important and worthwhile

approach to start to redress the effects of neglect or avoidance of these groups over

decades.

Importantly, we are not suggesting that large genera be split, or that there should be an

upper limit on the size of genera. The size of genera is related to both evolutionary and

human processes (e.g. Bentham's preference for fewer, larger genera; Humphreys & Linder,

2009), and the highly skewed pattern of genus size frequency, with many small and

relatively few very large genera, is common across the tree of life (e.g. Strand & Panova,

2015; Sigwart *et al.*, 2018). Although ultimately arbitrary, there is a broad consensus that

'good' genera (monophyletic, stable and diagnosable) are useful divisions (Humphreys &

Linder, 2009; Vences *et al.*, 2013; Strand & Panova, 2015; Muñoz‑Rodríguez *et al.*, 2023),

so big plant genera (and their problems) are unlikely to be split out of existence (Moonlight

et al., 2024). However, regardless of the basis of genus size, our results show that there are

tangible implications of being a member of a big genus that can influence species'

persistence.

We strongly encourage greater focus and resourcing of the further study of big plant genera

in taxonomic, ecological and conservation contexts. Although big genera continue to

represent an immense amount of work to monograph *in toto*, molecular data can be used to

divide large taxa into more manageable subclades; as in *Eugenia* or *Myrcia* (e.g., Mazine *et*

al., 2014; Lucas *et al.*, 2018). Another approach may be the 'foundation monograph'

concept (Scotland & Wood, 2012) – a practical guide to this approach (based on the recent

monograph of *Ipomoea*; Muñoz-Rodríguez *et al.*, 2019) is detailed in Muñoz-Rodríguez *et al.*

(2024). We echo earlier recommendations for increased collaboration across conservation

and taxonomy (e.g. Nic Lughadha *et al.*, 2019; Brown *et al.*, 2023a). The success of any new

initiative focused on assessing big genera will depend on a clear-eyed, evidence-based

understanding of the challenges likely to be faced; among these is the higher proportion of

Data Deficient species in genera which lack recent monographs (Grace *et al.*, 2021) and are

likely to include unusually large proportions of species not yet described (Moonlight *et al.*,

2024). We found that species from big genera were at least twice as likely to be categorised

as Data Deficient as those in monotypic genera, a result consistent with our expectations.

However, this proportion remained relatively low overall (<10%, Fig 5) and could be further

minimised by effective collaboration between taxonomists, assessors and other local

experts to assemble the most complete information possible for species' assessments.

Where assessors identify species likely to be Data Deficient, this can feed back into

taxonomic prioritisation (e.g. targeted fieldwork or other data collection). Similarly,

taxonomists can provide input on the likelihood of species re-circumscriptions, facilitating

effective prioritisation of assessments in big genera – thus, a shared pipeline of activity

between taxonomists and assessors can allow extinction risk assessments of species in

speciose genera to be undertaken before a genus is fully monographed (Nic Lughadha *et al.*,

2019).

Species in big genera are undoubtedly conservation priorities; they are under-studied,

under-assessed, and their extinction risk has been underestimated. Over a quarter of

angiosperm diversity is in big genera, but overcoming the challenges of working with

speciose taxa will only be possible with support and further strengthening of joint efforts

across taxonomy and conservation.

Methods

Data

We calculated genus size as the number of accepted species in each genus in version 13 of the World Checklist of Vascular Plants (WCVP; Govaerts *et al.*, 2021), accessed via the 'rWCVP' and 'rWCVPdata' R packages (Brown *et al.*, 2023b). Most species lack comprehensive and/or unbiased point data from which to measure range size; instead, as a proxy for range size, we used the number of Level 3 Areas (hereafter 'botanical countries', following e.g. Bachman *et al.*, 2024) of the World Geographic Scheme for Recording Plant Distributions (Brummitt *et al.*, 2001), accessed via WCVP. Although number of botanical countries is an imperfect proxy for range size (skewed towards single-country endemics, low spatial resolution and unequally-sized botanical countries), it remains the most comprehensive and least biased available data for angiosperms and shows a robust correlation with extent of occurrence, a common measure of range size (Bureš *et al.*, 2024).

We extracted lifeform information from WCVP, then simplified the Raunkiaer lifeforms into annual, herbaceous perennial, tree, other woody perennial, and epiphyte following Humphreys et al. 2019 with modifications (2019; see Table S1). For climatic zones, we derived the broad climate types (tropical, subtropical, temperate) from the WCVP 'climate description' field or species' distributions; species spanning multiple climate zones were combined into a 'multiple climates' category. We focused on the genus-level effects of climate and lifeform, rather than species-level (e.g. species in broadly tropical genera). To assign genus-level climate and lifeform categories, we aggregated species-level climate and lifeform data from WCVP to genus-level using a 75% dominance threshold – e.g. 89.3% of species in *Tillandsia* are epiphytic and 81.4% are tropical, so we treated *Tillandsia* as a tropical, epiphytic genus. Where no single lifeform or climate dominated, genera were assigned to a single 'multiple climates/lifeforms' category. This aggregated approach had the additional benefit of allowing us to include in our analyses the c. 78,480 species lacking lifeform information in WCVP version 13. We obtained extinction risk assessments from the IUCN Red List of Threatened Species version 2024-1 (hereafter Red List; IUCN, 2022b), and extinction risk predictions from Bachman *et al.* (2024) using 'rWCVP' (Brown *et al.*, 2023b) for taxonomic reconciliation.

Analyses

We used logistic regression to model five response variables as a function of genus size: (1) range size, (2) probability of having a global Red List assessment, (3) probability of being categorised as Data Deficient on the Red List, (4) probability of being assessed as threatened on the Red List (i.e. categorised as Vulnerable, Endangered or Critically Endangered), and (5) probability of being either assessed as or predicted to be threatened. Genus size was $\log_{10}$-transformed prior to analysis; all effect sizes represent changes on this scale. To investigate the consistency of the genus size-range size relationship across angiosperm diversity, we included three covariates in our analyses: climatic zone, lifeform and family. For each of the five response variables fitted, our overall approach was the same. We first fitted a 'full model', which included all covariates and their interaction with genus size. We then performed model selection using AIC (via the 'dredge' function implemented in 'MuMIn' ; Barton, 2024) to identify the best model. To interpret the effect sizes and their associated uncertainty for models with multiple categorical interaction terms, we used estimated marginal means and estimated marginal trends for each value of covariates using the 'emmeans' and 'emtrends' functions (from the 'emmeans' R package; Lenth, 2024). To account for imbalances in these variables, we weighted the non-focal covariates by their proportional frequency in the dataset (weights = "cells"). For models that included a family-level interaction term (including those used for model selection), we removed families with fewer than five genera or fewer than 500 species. Although this retained only 25% of angiosperm families, the effect on total species and generic diversity was small; the reduced dataset of 100 families included 89.3% of angiosperm genera and 91.8% of angiosperm species. Removing Aquifoliaceae (which comprises one genus), Balsaminaceae (two genera), Begoniaceae (two genera), Dioscoreaceae (four genera) and Ebenaceae (three genera) resulted in the removal of five big genera from our initial (full) models: *Begonia* (2,092 species), *Dioscorea* (635 species), *Diospyros* (778 species), *Ilex* (564 species) and *Impatiens* (1,120 species). For models fitted to assessed species only, we removed families with fewer than 250 assessments on the Red List (49 families). To check for bias resulting from these reductions, we also fitted models without family effects to the full species dataset.

To account for the skew in range size distribution and lack of fine-scale resolution when using botanical countries, we modelled range size as a function of genus size using a series

of logistic regressions. We repeated model-fitting at multiple range-size thresholds using a

'multiple threshold' approach to test the robustness of this relationship at different scales.

We modelled the probability that species are endemic to a single botanical country,

restricted to two or fewer botanical countries, then to three, four and ten or fewer botanical

countries.

We used the same 'multiple threshold ' approach to model extinction risk as for previous

studies (e.g. Brown *et al.*, 2023a; Soto Gomez *et al.*, 2024); for the full models including

climate, lifeform and family we focused on a single threshold: the probability of being

'threatened' per the IUCN (Critically Endangered, Endangered, Vulnerable; IUCN, 2022a), or

predicted to be threatened with high confidence (Bachman *et al.*, 2024). Species assessed as

Extinct (84 species in reduced dataset) or Extinct in the Wild (31 species) could not be

modelled separately, so were included under Critically Endangered for all analyses.

To highlight the implications of under-assessment of species in big genera, we directly

compared the proportion of threatened species (using the threshold above, noting that this

differs from the IUCN calculation in the handling of DD species) across genus sizes based on

assessments only, and assessments plus predictions. From these data, we extracted the 20

big genera with the highest mismatch between their proportion threatened per the Red List,

and their estimated proportion threatened from the Bachman et al. (2024) predictions. We

present these genera as potential priorities for conservation assessment.

All models were fitted with species as the observation (i.e. data were not aggregated to

genus prior to analyses), in R version 4.4.1 (R Core Team, 2024). All data, scripts and outputs

required to reproduce our results are available at

https://github.com/matildabrown/BigAngiospermGenera [NB: currently private, materials

provided separately for review].

Extended Data

Extended Data Figure 1. Comparison of the estimated marginal means for Red List data (red) and
combined (Red List + predictions) data (grey) for the relationship between genus size and probability
of extinction risk. Dotted lines show where the lines intercept (grey; left) and where the 95% CIs
diverge (blue; right), indicating the change from over- to under-estimation of threat.

Extended Data Table 1. Numbers of species assessed as versus predicted to be threatened in big genera, arranged in descending order of underprediction (Δthreat prop.). Top 20 genera shown here, for full list of big genera see Table S4.

Genus	Climate	# spp.	prop. spp. assessed	# spp. threatened (RL)	prop. threatened (RL)	# spp. predicted threatened	prop. predicted threatened	Δthreat prop.
Stelis	tropical	1340	0.4%	0	0.0%	1060	79.1%	79.1%
Taraxacum	temperate	2490	0.1%	0	0.0%	1243	49.9%	49.9%
Lupinus	multiple climates	580	11.7%	6	8.8%	333	57.4%	48.6%
Lepanthes	tropical	1203	2.2%	11	40.7%	1058	87.9%	47.2%
Saussurea	multiple climates	516	0.6%	0	0.0%	232	45.0%	45.0%
Pedicularis	temperate	679	4.4%	3	10.0%	342	50.4%	40.4%
Ranunculus	temperate	1750	2.9%	11	21.6%	1011	57.8%	36.2%
Elatostema	multiple climates	654	0.5%	1	33.3%	435	66.5%	33.2%
Primula	temperate	545	3.5%	2	10.5%	224	41.1%	30.6%
Corydalis	temperate	538	1.3%	2	28.6%	310	57.6%	29.0%
Pleurothallis	tropical	579	0.7%	2	50.0%	439	75.8%	25.8%
Masdevallia	tropical	651	1.7%	6	54.5%	523	80.3%	25.8%
Piper	tropical	2425	6.4%	65	41.9%	1634	67.4%	25.4%
Epidendrum	tropical	1878	2.2%	17	40.5%	1210	64.4%	24.0%
Maxillaria	tropical	652	1.1%	2	28.6%	341	52.3%	23.7%
Oxytropis	temperate	608	1.5%	1	11.1%	208	34.2%	23.1%
Anthurium	tropical	1325	10.9%	46	31.9%	691	52.2%	20.2%
Ardisia	tropical	737	26.2%	68	35.2%	393	53.3%	18.1%
Berberis	multiple climates	626	9.3%	22	37.9%	350	55.9%	18.0%
Allium	temperate	1078	13.6%	31	21.1%	403	37.4%	16.3%

Acknowledgements

TCCL was funded by a Future Leader Fellowship from the Royal Botanic Gardens, Kew.

Author contributions

MJMB: Conceptualisation, Formal analysis, Visualisation, Investigation, Writing - Original Draft

TCCL: Conceptualisation, Investigation, Writing - Original Draft, Review & Editing

FF: Writing - Review & Editing

EL: Writing - Review & Editing

BEW: Writing - Review & Editing

ENL: Conceptualisation, Investigation, Writing - Review & Editing

Data availability

All source data are publicly available. Scripts and other materials required to reproduce our outputs are available at https://github.com/matildabrown/BigAngiospermGenera

Competing interests

The authors declare no competing interests.

References

Bachman SP, Brown MJM, Leao TCC, Nic Lughadha E, Walker BE. 2024. Extinction risk predictions for the world's flowering plants to support their conservation. *New Phytol* **242**(2): 797-808.

Bachman SP, Field R, Reader T, Raimondo D, Donaldson J, Schatz GE, Lughadha EN. 2019. Progress, challenges and opportunities for Red Listing. *Biological Conservation* **234**: 45-55.

Barton K 2024. MuMIn: Multi-Model Inference. R package.

Brown MJM, Bachman SP, Nic Lughadha E. 2023a. Three in four undescribed plant species are threatened with extinction. *New Phytol* **240**(4): 1340-1344.

Brown MJM, Walker BE, Black N, Govaerts RHA, Ondo I, Turner R, Nic Lughadha E. 2023b. rWCVP: a companion R package for the World Checklist of Vascular Plants. *New Phytol* **240**(4): 1355-1365.

Brummitt N, Cui Q, Bai P, Liu X, Scott B, De Palma A, Araujo AC. 2024. Reducing the number of Data Deficient plant species. *Systematics and Biodiversity* **22**(1): 2302180.

Brummitt RK, Pando F, Hollis S, Brummitt N. 2001. *World geographical scheme for recording plant distributions*. Carnegie Mellon University (Pittsburgh): Hunt Institute for Botanical Documentation.

Bureš P, Elliott TL, Veselý P, Šmarda P, Forest F, Leitch IJ, Nic Lughadha E, Soto Gomez M, Pironon S, Brown MJ. 2024. The global distribution of angiosperm genome size is shaped by climate. *New Phytologist* **242**(2): 744-759.

Cowell C. WE, Bullough L.-A., Grey J., Klitgaard B., Govaerts R., Andriambololonera S., Cervantes A., Crameri S., Lima, H.C., Lachenaud O., Li S.-J., Linares J.L., Phillipson P ., Rakotonirina N., Wilding N., van der Burgt X., Vatanparast M., Barker A., Barstow M., Beentje H., Plummer J. 2022. CITES Dalbergia Checklist. Commissioned by the CITES Secretariat. . Surrey, UK.

Davies TJ, Smith GF, Bellstedt DU, Boatwright JS, Bytebier B, Cowling RM, Forest F, Harmon LJ, Muasya AM, Schrire BD. 2011. Extinction risk and diversification are linked in a plant biodiversity hotspot. *PLoS biology* **9**(5): e1000620.

Freeman BG, Pennell MW. 2021. The latitudinal taxonomy gradient. *Trends in Ecology & Evolution* **36**(9): 778-786.

Fu Q, Huang X, Li L, Jin Y, Qian H, Kuai X, Ye Y, Wang H, Deng T, Sun H. 2022. Linking evolutionary dynamics to species extinction for flowering plants in global biodiversity hotspots. *Diversity and Distributions* **28**(12): 2871-2885.

Govaerts R, Nic Lughadha E, Black N, Turner R, Paton A. 2021. The World Checklist of Vascular Plants, a continuously updated resource for exploring global plant diversity. *Sci Data* **8**(1): 215.

Grace OM, Pérez-Escobar OA, Lucas EJ, Vorontsova MS, Lewis GP, Walker BE, Lohmann LG, Knapp S, Wilkie P, Sarkinen T. 2021. Botanical monography in the Anthropocene. *Trends in Plant Science* **26**(5): 433-441.

Greenberg DA, Pyron RA, Johnson LG, Upham NS, Jetz W, Mooers AØ. 2021. Evolutionary legacies in contemporary tetrapod imperilment. *Ecology Letters* **24**(11): 2464-2476.

Guedes JJ, Moura MR, Jardim L, Diniz-Filho JAF. 2025. Global Patterns of Taxonomic Uncertainty and its Impacts on Biodiversity Research. *Systematic biology*: syaf010.

Humphreys AM, Linder HP. 2009. Concept versus data in delimitation of plant genera. *Taxon* **58**(4): 1054-1074.

IUCN. 2012. *IUCN Red List categories and criteria: version 3.1*. Gland, Switzerland and Cambridge,
UK: Prepared by the IUCN Species Survival Commission.
IUCN 2022a. Guidelines for Using the IUCN Red List Categories and Criteria.
IUCN 2022b. The IUCN Red List of Threatened Species.
IUCN 2024. The IUCN Red List of Threatened Species.
Leão TC, Lughadha EN, Reich PB. 2020. Evolutionary patterns in the geographic range size of Atlantic
Forest plants. *Ecography* **43**(10): 1510-1520.
Lenth R 2024. emmeans: Estimated marginal means, aka least-squares means. R package version
1.10.3.
Lozano FD, Schwartz MW. 2005. Patterns of rarity and taxonomic group size in plants. *Biological
Conservation* **126**(2): 146-154.
**Lucas E, Amorim B, Lima D, Lima-Lourenço A, Nic Lughadha E, Proença C, Rosa P, Rosário A, Santos
L, Santos M. 2018.** A new infra-generic classification of the species-rich Neotropical genus
Myrcia sl. *Kew Bulletin* **73**: 1-12.
Mazine FF, Souza VC, Sobral M, Forest F, Lucas E. 2014. A preliminary phylogenetic analysis of
Eugenia (Myrtaceae: Myrteae), with a focus on Neotropical species. *Kew Bulletin* **69**: 1-14.
Moonlight PW, Baldaszti L, Cardoso D, Elliott A, Särkinen T, Knapp S. 2024. Twenty years of big
plant genera. *Proceedings of the Royal Society B* **291**(2023): 20240702.
Muñoz-Rodríguez P, Carruthers T, Wells T, Sumadijaya A, Wood JR, Scotland RW. 2024. The
research behind a taxonomic monograph: a case study from Ipomoea (Convolvulaceae). *Kew
Bulletin* **79**(4): 897-914.
**Muñoz-Rodríguez P, Carruthers T, Wood JR, Williams BR, Weitemier K, Kronmiller B, Goodwin Z,
Sumadijaya A, Anglin NL, Filer D. 2019.** A taxonomic monograph of Ipomoea integrated
across phylogenetic scales. *Nature plants* **5**(11): 1136-1144.
Muñoz‐Rodríguez P, Wood JR, Wells T, Carruthers T, Sumadijaya A, Scotland RW. 2023. The
challenges of classifying big genera such as Ipomoea. *Taxon* **72**(6): 1201-1215.
**Newton A, Oldfield S, Rivers M, Mark J, Schatz G, Garavito NT, Cantarello E, Golicher D, Cayuela L,
Miles L. 2015.** Towards a global tree assessment. *Oryx* **49**(3): 410-415.
**Nic Lughadha E, Bachman SP, Leão TCC, Forest F, Halley JM, Moat J, Acedo C, Bacon KL, Brewer
RFA, Gâteblé G, et al. 2020.** Extinction risk and threats to plants and fungi. *PLANTS, PEOPLE,
PLANET* **2**(5): 389-408.
Nic Lughadha EM, Graziele Staggemeier V, Vasconcelos TNC, Walker BE, Canteiro C, Lucas EJ. 2019.
Harnessing the potential of integrated systematics for conservation of taxonomically
complex, megadiverse plant groups. *Conserv Biol* **33**(3): 511-522.
Purvis A, Agapow P-M, Gittleman JL, Mace GM. 2000. Nonrandom extinction and the loss of
evolutionary history. *science* **288**(5464): 328-330.
R Core Team 2024. R: A language and environment for statistical computing. Vienna, Austria: R
Foundation for Statistical Computing.
Rivers M. 2017. The Global Tree Assessment—red listing the world's trees. *BGjournal* **14**(1): 16-19.
Schwartz MW, Simberloff. 2001. Taxon size predicts rates of rarity in vascular plants. *Ecology Letters*
4(5): 464-469.
Scotland RW, Wood JR. 2012. Accelerating the pace of taxonomy. *Trends in Ecology & Evolution*
27(8): 415-416.
Sigwart JD, Sutton MD, Bennett KD. 2018. How big is a genus? Towards a nomothetic systematics.
Zoological Journal of the Linnean Society **183**(2): 237-252.
**Soto Gomez M, Brown MJ, Pironon S, Bureš P, Verde Arregoitia LD, Veselý P, Elliott TL, Zedek F,
Pellicer J, Forest F. 2024.** Genome size is positively correlated with extinction risk in
herbaceous angiosperms. *New Phytologist* **243**(6): 2470-2485.
Strand M, Panova M. 2015. Size of genera—biology or taxonomy. *Zoologica Scripta* **44**(1): 106-116.

Tanentzap AJ, Igea J, Johnston MG, Larcombe MJ. 2020. Does evolutionary history correlate with
contemporary extinction risk by influencing range size dynamics? *The American Naturalist*
195(3): 569-576.
Vamosi JC, Wilson JRU. 2008. Nonrandom extinction leads to elevated loss of angiosperm
evolutionary history. *Ecology Letters* **11**(10): 1047-1053.
Vences M, Guayasamin JM, Miralles A, De La Riva I. 2013. To name or not to name: criteria to
promote economy of change in supraspecific Linnean classification schemes.
Walker BE, Tucker A, Nicolson N. 2022. Harnessing large-scale herbarium image datasets through
representation learning. *Frontiers in plant science* **12**: 806407.