Species in big angiosperm genera are priorities for conservation

Authors
Matilda J.M. Brown[1]†, Tarciso C.C. Leão[1]†, Félix Forest[1], Eve Lucas[1], Barnaby E. Walker[1], Eimear Nic Lughadha[1]

† These two authors contributed equally to this work

Corresponding author:
Matilda Brown m.brown2@kew.org
Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE, UK

Affiliations
[1]Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE, UK

Article Impact Statement
Species in big angiosperm genera are under-studied, under-assessed, and their extinction risk has been underestimated.

Keywords
Big genera, range size, extinction risk, Red List, angiosperms

Abstract

Species in big angiosperm genera (≥500 species) are often considered to be less well-known than those in smaller genera. Consequently, species in big genera are likely underrepresented in conservation at all levels, from analyses to actions. However, emerging evidence indicates that species in larger genera tend to have smaller geographic ranges (a strong correlate of increased extinction risk). Here, we test the generality of this pattern by quantifying the relationships between plant taxon size, extinction risk and climatic zone on a global scale. We find that species in larger genera are more likely to have small ranges, less likely to have a global Red List assessment, and, when assessed, are more likely to be threatened. Persistent obstacles to improving conservation documentation of big genera include taxonomic uncertainty, data shortfalls and number of species involved, requiring continued collaboration between taxonomy and conservation to reduce this bias and enable effective conservation of the quarter of plant species that big genera encompass.

Introduction

'Big' plant genera (≥500 species per Frodin, 2004) contain over a quarter of angiosperm species (Moonlight *et al.*, 2024) and include several ecologically, culturally and economically important taxa (e.g. *Euphorbia*, >2,000 spp.; *Solanum*, >1,200 spp., *Eucalyptus*, >700 spp.). Despite the familiarity of many big plant genera, there remains a pervasive knowledge shortfall for species in

big genera (Moonlight *et al.*, 2024) largely due to the practical challenges associated with

working with groups of ≥500 species – through this lens, genus size can be viewed as an

attribute that has an effect on species' individual likelihood of being studied. For example, big

genera are less likely to have a global taxonomic monograph and have lower proportions of

species with available DNA sequence data (Grace *et al.*, 2021). Longstanding latitudinal biases

in taxonomic effort (the 'Latitudinal Taxonomy Gradient'□; Freeman & Pennell, 2021; Guedes *et*

al., 2025) mean that this shortfall is particularly severe in the tropics, where the majority of big

plant genera occur (Moonlight *et al.*, 2024; though this is an effect of overall biodiversity

concentration rather than a specific concentration of big genera in the tropics; Baldaszti *et al.*,

2025).

We expect that both the practical and taxonomic challenges associated with big plant genera

limit our knowledge of extinction risk of species in these taxa; for example, <5% of the >3,000

species of *Astragalus* have global Red List assessments (far fewer than the c. 20% coverage of

angiosperms at large; IUCN, 2024), but the effect of genus size on assessment probability has

not been quantified. Undertaking a fully-documented species conservation assessment (e.g. a

Red List assessment; IUCN, 2012, 2022a) is expensive in terms of time and expertise. Thus,

species considered data-poor, which are likely to be assessed as Data Deficient (Brummitt *et al.*,

2024) tend to be low priorities for assessment. Similarly, an assessment is specific to a particular

species concept, so species that are more likely (or perceived to be more likely) to be re-

circumscribed are less likely to be prioritised for assessment. Assessments are typically

undertaken as part of projects of clearly defined geographic and/or taxonomic scope, often

aiming to assess all species within that scope (e.g. the recent assessment of Dalbergia; Cowell

et al., 2022; and 332 Mozambican endemics Darbyshire *et al.*, 2019). While the global

distribution of big genera (Baldaszti *et al.*, 2025) means that species in big genera may be

proportionally represented in geographically defined projects, they are rarely the taxonomic focus

of assessment projects. Comprehensive assessment of species in a big genus requires

substantial resources, and usually many collaborators. The successful Global Tree Assessment

(>45,000 species; Rivers, 2017) demonstrates the feasibility of large assessment projects, but

also the enormity of the challenges involved (Newton *et al.*, 2015). We expect that the sheer

numbers of species and the (real or perceived) risk of taxonomic uncertainty due to e.g. lack of

monographs for big genera mean that a) they are systematically underrepresented on the global

Red List and b) if assessed, are more likely to be assessed as Data Deficient.

There is also emerging evidence that we may be underestimating extinction risk in big plant

genera. Geographic range size is the strongest correlate of plant extinction risk (Nic Lughadha *et*

al., 2019; Walker *et al.*, 2022; Bachman *et al.*, 2024), and both region-specific and flora-based

studies have found that taxa with larger numbers of species have, on average, species with

smaller geographic ranges (Schwartz & Simberloff, 2001; Lozano & Schwartz, 2005; Leão *et al.*,

2020) and higher proportions of threatened species (Fu *et al.*, 2022). Furthermore, a

disproportionate number of species described in the last two decades – the majority of which are

likely to be threatened (Brown *et al.*, 2023a) – are in the very largest genera (>1,000 species;

Moonlight *et al.*, 2024). The global generality of the relationships between genus size, range size

and extinction risk in flowering plants remain untested, particularly across climatic zones,

lifeforms and higher taxonomic ranks, but we expect that an overall genus size - range size

relationship, if supported, will also result in higher proportions of threatened species in larger

genera.

Here, we test the generality of the negative relationship between genus size and species

geographic range size across climate, life form and family, using data from the World Checklist of

Vascular Plants ('WCVP'; Govaerts *et al.*, 2021). We then quantify the extent to which species in

larger genera are more or less likely to have a Red List assessment, and their probability of,

when assessed, being categorised as either Data Deficient or threatened with extinction, using

data from the IUCN Red List (hereafter 'Red List'; IUCN, 2024) and comprehensive species-level

extinction risk predictions from Bachman *et al.* (2024). Finally, we identify specific priority big

genera for conservation assessment as those for which overall extinction risk on the Red List is

most underestimated when compared to predictions.

Materials and Methods

Data

We calculated genus size as the number of accepted species in each genus in version 13 of the

World Checklist of Vascular Plants (Govaerts *et al.*, 2021), accessed via the 'rWCVP' and

'rWCVPdata' R packages (Brown *et al.*, 2023b). Most species lack comprehensive and/or

unbiased point data from which to measure range size; instead, as a proxy for range size, we

used the number of Level 3 Areas (hereafter 'botanical countries', following e.g. Bachman *et al.*,

2024; Svahnström *et al.*, 2025) of the World Geographic Scheme for Recording Plant

Distributions (Brummitt, 2001) accessed via WCVP. Although number of botanical countries is an

imperfect proxy for range size (skewed towards single-country endemics, low spatial resolution

and unequally-sized botanical countries), it remains the most comprehensive and least biased

available data for angiosperms. We acknowledge that these data cannot be said to be a direct

measure of range size, but instead represent an intermediate proxy, where species limited to a

single botanical country are more likely to have smaller ranges than those occurring in several

botanical countries (Svahnström *et al.*, 2025). While there are specific scenarios where the

relationship between range size and number of botanical countries breaks down (e.g. range-

restricted species occurring across country borders, species that are widespread within a large

botanical country), we assume that at a global scale, these cases are unlikely to be present with
enough frequency to significantly affect our results. Empirical support for this assumption has
been reported from a range of studies, where results are consistent across multiple proxies of
range size (e.g. extent of occurrence; Bureš *et al.*, 2024; Svahnström *et al.*, 2025).

We extracted lifeform information from WCVP, then simplified the Raunkiaer lifeforms into
annual, herbaceous perennial, tree, other woody perennial, and epiphyte following Humphreys et
al. (Humphreys *et al.*, 2019) with modifications (see Table S1). For climatic zones, we derived
the broad climate types (tropical, subtropical, temperate) from the WCVP 'climate description'
field or species' distributions; species spanning multiple climate zones were combined into a
'multiple climates' category (Supplementary text; Fig. S1). We focused on the genus-level effects
of climate and lifeform, rather than species-level (e.g. species in broadly tropical genera). To
assign genus-level climate and lifeform categories, we aggregated species-level climate and
lifeform data from WCVP to genus-level using a 75% dominance threshold – e.g., 89.3% of
species in *Tillandsia* are epiphytic and 81.4% are tropical, so we treated *Tillandsia* as a tropical,
epiphytic genus. Where no single lifeform or climate dominated, genera were assigned to a
single 'multiple climates/lifeforms' category. This aggregated approach had the additional benefit
of allowing us to include in our analyses the c. 78,480 species lacking lifeform information in
WCVP version 13. We obtained extinction risk assessments from the IUCN Red List of
Threatened Species version 2024-1 (IUCN, 2024), and extinction risk predictions from Bachman
et al. (2024) using 'rWCVP' (Brown *et al.*, 2023b) for taxonomic reconciliation, including matching
of basionyms, where the taxonomic concept matches that of the accepted species name.

Statistical analysis

We used logistic regression to model five response variables as a function of genus size: (1)
range size, (2) probability of having a global Red List assessment, (3) probability of being
categorised as Data Deficient on the Red List, (4) probability of being assessed as threatened on
the Red List (i.e. categorised as Vulnerable, Endangered or Critically Endangered), and (5)
probability of being either assessed as or predicted to be threatened. Genus size was $\log_{10}$-
transformed prior to analysis; all effect sizes represent changes on this scale. To investigate the
consistency of the genus size-range size relationship across angiosperm diversity, we included
three covariates in our analyses: climatic zone, lifeform and family. For each of the five response
variables fitted, our overall approach was the same. We first fitted a 'full model', which included
all covariates and their interaction with genus size (model formula given in Eqn 1; + denotes
additive terms; * denotes additive and interaction terms).

Equation 1.   *Response ~ GenusSize*Climate + GenusSize*Lifeform + GenusSize*Family*

We then performed model selection using AICc (via the 'dredge' function implemented in

'MuMIn'□; Barton, 2024) to identify the best model/s. To interpret the effect sizes and their

associated uncertainty for models with multiple categorical interaction terms, we used estimated

marginal means and estimated marginal trends for each value of covariates using the 'emmeans'

and 'emtrends' functions from the 'emmeans' R package (Lenth, 2025). To account for

imbalances in these variables, we weighted the non-focal covariates by their proportional

frequency in the dataset (weights = "cells"). For models that included a family-level interaction

term (including those used for model selection), we removed families with fewer than five genera

or fewer than 500 species. Although this retained only 25% of angiosperm families, the effect on

total species and generic diversity was small; the reduced dataset of 100 families included 89.3%

of angiosperm genera and 91.8% of angiosperm species. Removing Aquifoliaceae (which

comprises one genus), Balsaminaceae (two genera), Begoniaceae (two genera), Dioscoreaceae

(four genera) and Ebenaceae (three genera) resulted in the removal of five big genera from our

initial (full) models: *Begonia* (2,092 species), *Dioscorea* (635 species), *Diospyros* (778 species),

Ilex (564 species) and *Impatiens* (1,120 species). For models fitted to assessed species only, we

removed families with fewer than 250 assessments on the Red List (49 families). To check for

bias resulting from these reductions, we also fitted models without family effects to the full

species dataset.

To account for the skew in range size distribution and lack of fine-scale resolution when using

botanical countries, we modelled range size as a function of genus size using a series of logistic

regressions. We repeated model-fitting at multiple range-size thresholds using a 'multiple

threshold' approach to test the robustness of this relationship at different scales. We modelled

the probability that species are endemic to a single botanical country, restricted to two or fewer

botanical countries, then to three, four and ten or fewer botanical countries.

We used the same 'multiple threshold ' approach to model extinction risk as for previous studies

(Brown *et al.*, 2023a; Soto Gomez *et al.*, 2024); for the full models including climate, lifeform and

family we focused on a single threshold: the probability of being 'threatened' per the IUCN

(Critically Endangered, Endangered, Vulnerable; IUCN, 2022a), or predicted to be threatened

with high confidence (Bachman *et al.*, 2024). Species assessed as Extinct (84 species in

reduced dataset) or Extinct in the Wild (31 species) could not be modelled separately, so were

included under Critically Endangered for all analyses.

To highlight the implications of under-assessment of species in big genera, we directly compared

the proportion of threatened species (using the threshold above, noting that this differs from the

IUCN calculation in the handling of DD species; IUCN, 2022b) across genus sizes based on

assessments only, and assessments plus predictions. From these data, we extracted the 20 big

genera with the highest mismatch between their proportion threatened per the Red List, and their

estimated proportion threatened from the Bachman et al. (2024) predictions. We present these

genera as potential priorities for conservation assessment.

All models were fitted with species as the observation (i.e. data were not aggregated to genus

prior to analyses), in R version 4.4.1 (R Core Team, 2024). All data, scripts and outputs required

to reproduce our results are available at [link removed for double-blind review].

Results

Nearly half of all big genera (and species therein) are tropical (Fig 1, Table S2). Of the 82

currently accepted big genera (from version 13 of the WCVP), 40 (49%) are entirely or mostly

tropical, containing 40,631 species (48% of species in big genera; 13% of total angiosperm

diversity). All big epiphytic genera (10 genera; 11,370 species) are predominantly tropical;

similarly, all big tree genera are either tropical (4 genera; 3,211 species) or subtropical

(*Eucalyptus*, 713 species). By contrast, most of the 15 temperate big genera are dominated by

herbaceous perennials (12 genera; 16,239 species). There are no big genera dominated by

annual species.

Covariates of genus size trends

The best-supported model formula (ranked using AICc; Table S3) across all response variables

includes both additive and interaction terms for all covariates (climate, lifeform, family). Family-

level interaction terms could not be estimated for families with very few species or genera; the

results presented below are thus based on a reduced dataset (91.8% of angiosperm species;

see Methods). Models fitted using the full dataset (excluding family effects) are consistent with

the findings reported below (Tables S4-6).

Genus size and species range size

Overall, species in big genera tend to occur in fewer botanical countries so are, on average,

more likely to have smaller ranges (Fig 2). The probability of being restricted to a single botanical

country is 49.7% (95% CI: 49.2-50.2%) for a species in a monotypic genus, compared to 60.6%

(CI: 60.3-60.9%) for species in the 'smallest' big genera (i.e. genera of exactly 500 species; from

marginal mean estimate; Fig 2a).

The relationship between genus size and range size is consistent across scales (i.e. when

modelling the proportion of species restricted to one, two, three, four, five and ten botanical

countries, Fig 2a), climatic zones and lifeform (Fig 2b). It is strongest for annual, epiphytic,

temperate, and subtropical genera, and comparatively weak (though still significantly positive) for tropical and woody (tree, other woody perennial) genera (Fig 2b).

Across families included in this analysis (100 families; 91.8% of species), the genus size - range size relationship was mixed (Fig 2c); 54 families have significant (p<0.05), positive slope coefficients (i.e. species in larger genera are more likely to be restricted to a single botanical country), five families have significant negative coefficients and 41 families had slope coefficients not significantly different from zero.

Genus size and extinction risk

Species in big genera are less likely to have a global Red List assessment (Fig 3); for a monotypic genus, the probability of being assessed is 15.3% (95% CI: 14.4-16.2%), compared to 11.5% (CI: 11.2-11.7%) for a species in a 500-species genus (estimated marginal means).

Marginal trends across lifeform and climate are generally consistent with the overall effect, except for subtropical genera – species in subtropical genera are more likely to be assessed if they are in a larger genus (Fig 3a). This result is likely driven by the near-comprehensive (94%) assessment of *Eucalyptus* (Myrtaceae), the only large subtropical tree genus, comprising 27.0% of species (across all lifeforms) in big subtropical genera. The impact of the Global Tree Assessment is clearly visible in our data, with trees having the highest assessment coverage of any covariate (Fig 3a).

Marginal trends across families are less consistent: 49 families have significant negative trends as expected, 19 have positive trends and 32 are not significantly different from zero (50 largest families shown in Fig 3b). Regardless of lifeform, species in tropical genera are always more likely to have been assessed than those from other climates (Fig. 3a).

Using Red List data, we found that when assessed, species in big genera are slightly more likely to be assessed as threatened (either Vulnerable, Endangered or Critically Endangered; Fig 4a). On average, an assessed species in a monotypic genus has a 32.6% (CI: 31.0-34.1%) chance of being assessed as threatened, compared to 37.5% (CI: 36.8-38.2%) if it is in a genus of 500 species (estimated marginal means). The effect is generally reversed in epiphytic and subtropical genera (Fig 4b), though this is likely driven by a small number of assessments, as species from epiphytic or subtropical genera tend to be less likely to have assessments (Fig 3). Across the 51 families with sufficient assessment coverage for analysis (Fig 4c), 13 have a significant positive relationship as expected, 9 have a significant negative relationship and 29 have relationships between genus size and probability of being assessed as threatened that are not significantly different from zero.

Species in larger genera are also more likely to be categorised as Data Deficient (Fig. 5); the

probability that a species in a monotypic genus is categorised as Data Deficient is 4.5% (95%

CI:4.1-4.9%), compared to 9.1% (CI: 8.7-9.4%) in a genus with 500 species. The effect on Data

Deficiency of being in a large genus is strongest for tropical and epiphytic species, but reversed

for annual and subtropical species (i.e. species in larger genera are less likely to be assessed as

Data Deficient), though these latter groups have, respectively, only 18 and 94 species (zero and

seven in big genera) that have been assessed as Data Deficient. Data were insufficient to fit the

full model including family; full model for Data Deficiency fitted with climate and lifeform

covariates only.

When we extended our analysis of extinction risk to include all angiosperms, by including

extinction risk predictions (from Bachman *et al.*, 2024) for species lacking a Red List

assessment, we found a stronger overall relationship between genus size and extinction risk (Fig

6): a species in a monotypic genus has a 30.4% (CI: 29.8-31.1%) chance of being threatened

(assessed or predicted with high confidence), compared to 45.6% (CI: 45.3-45.9%) for a genus

of 500 species (estimated marginal means).

The relationship between genus size and predicted probability of being threatened is consistent

across climatic zones and lifeforms (Fig 6b). It is strongest for annual, epiphytic and genera

found in multiple climates, and comparatively weak (though still significantly positive) for tropical,

subtropical and woody genera (Fig 6b). Across families (Fig 6c), the genus size – extinction risk

relationship is broadly consistent, though statistical power is limited; of the 100 families included

in this analysis, 56 have significant ($p<0.05$), positive slope coefficients (i.e. species in larger

genera are more likely to be either assessed as or predicted to be threatened), five have

significant negative coefficients and 39 have slope coefficients that are not significantly different

from zero. When we compared estimates of the predicted proportions of threatened species

based on assessments versus assessments plus predictions (Fig 7), we found evidence to

suggest that current Red List data underestimate the proportion of species that are threatened

with extinction in genera of eleven or more species. This underestimation is most severe for big

genera.

From the extinction risk predictions (Bachman *et al.*, 2024), 37,072 species in big genera are

estimated (with high confidence) to be threatened with extinction, but either lack a global Red

List assessment (36,411) or have been assessed as Data Deficient (661) – together representing

10.9% of all angiosperm species. This number is conservative, as >13,000 species have been

described or otherwise added to WCVP since the release used as the backbone for predictions.

Of the species in big genera predicted to be threatened, 19,787 are in tropical genera (spanning

40 genera). We quantified the degree to which extinction risk is likely to be underestimated in

each large genus by subtracting the Red-List-derived proportion of threatened species from that

calculated using predictions (Table 1). Of the twenty big genera in which extinction risk is likely to be most underestimated on the Red List, six are tropical genera in the family Orchidaceae. The most extreme example is *Stelis* with <1% of species having a Red List assessment and none threatened, while 79% of species are predicted to be threatened.

Discussion

Consistent with our expectations, being in a larger genus is associated with having a smaller geographic range, being less likely to have a Red List assessment, and if assessed, being more likely to be either Data Deficient, or assigned to one of the threatened categories. The relationships between genus size, range size and extinction risk are broadly consistent across climatic zones, lifeforms, and families; as over 25% of plant species are in 'big' genera (≥500 species) and a further 35% are in moderately large genera (≥100 species), our findings are relevant to the majority of known plant species and have significant implications for conservation science, action and policy.

Our results support previous studies on the relationship between taxon size and range size in plants (in genera; Schwartz & Simberloff, 2001; Lozano & Schwartz, 2005; Leão *et al.*, 2020; and families; Fu *et al.*, 2022). However, the relationship between genus and range size is not universal – in five families, species in larger genera are less likely to be restricted to a single botanical country, contrary to expectations (Table S4). These families span seven orders (both eudicots and monocots), vary in climate, lifeform and range sizes, and three contain at least one big genus, so we found no clear pattern to these outliers. Similarly, we found groups (discussed below) for which the general pattern of higher extinction risk in larger genera is reversed, where species in larger genera are *less* likely to be either assessed as or predicted to be threatened (Figs 3-4; Tables S4-S6).

Results based on published assessments were less consistent than those based on predictions, likely due to the very low assessment coverage of many big genera. For example, in tropical epiphytes, where seven of the ten big genera have fewer than 50 published assessments (representing <3% of species in these seven genera), assessments show an apparent negative relationship between genus size and extinction risk (Fig 4), but the model including predictions showed the expected positive trend (Fig 5). There are five families with a negative relationship between genus size and extinction risk in prediction-based models. For one of these, Melastomataceae, this may be linked to range, as it was one of the five families where species in larger genera tended to have larger ranges (Table S4), but for the other four the reasons for this result are unclear; differences in both range size and extinction risk may be linked to the 'different ways to be a big genus' (an outstanding knowledge gap noted by Moonlight *et al.*, 2024). It is also possible that there are biases in the predictions that affect our results – for

example, the phylogenetic eigenvectors used as predictors by Bachman *et al.* (2024) make it

more likely that predictions match the Red List-derived estimates of extinction risk within a genus

(though this effect seems to be weak; Table 1). These exceptions do not in any way diminish the

need for species in big genera to be prioritised for conservation assessment, whether to formally

document their predicted extinction risk, address the gaps in both knowledge and data for big

genera or to better understand the drivers of extinction risk across taxon size.

The mechanisms underpinning the link between genus and range sizes may be related to

evolutionary diversification patterns. More rapidly diversifying plant genera tend to have smaller

ranges and higher extinction risk (Davies *et al.*, 2011; Leão *et al.*, 2020; Tanentzap *et al.*, 2020)

though conifers showed the opposite trend (Tanentzap *et al.*, 2020). In animals, genus-level

diversification rate appears to be a strong correlate of both range size and extinction risk (though

there is significant variation across clades; Greenberg *et al.*, 2021), further supporting the

hypothesis that rapid speciation tends to produce range-restricted species which are inherently

more vulnerable to extinction (Davies *et al.*, 2011; Leão *et al.*, 2020). Our findings regarding

higher extinction risk of species in larger genera are in contrast to earlier reports for both plants

(Vamosi & Wilson, 2008) and animals (Purvis *et al.*, 2000) that smaller – especially monotypic –

taxa tend to have higher levels of extinction risk.

Our results complement recent studies characterising biases in the representation of plants on

the Red List and addressing how these can be quantified and corrected (Nic Lughadha *et al.*,

2020) or circumvented (Bachman *et al.*, 2024) in future analyses to refine understanding of

global and regional patterns of plant extinction risk. For example, eight of the 20 most under-

estimated genera (when overall extinction risk is estimated using the Red List; Table 1) are

Orchidaceae (6 genera) or Asteraceae (2 genera). These two largest plant families encompass

c. 65,000 species and are recognised as being among the most under-represented on the Red

List (Nic Lughadha *et al.*, 2020).

Strikingly, the six orchid genera highlighted here for their underestimated threat levels are all

tropical, consistent with our original expectation that under-representation of big genera in

assessment processes might be greatest in the tropics. However, contrary to our expectations,

we found that species in tropical genera, across all lifeforms and independent of genus size,

were more likely to have been assessed than species of other climates (Fig 3a). This

encouraging result suggests significant progress towards addressing the plant Red Listing deficit

which was so evident across many areas of the tropics as recently as a decade ago (Bachman *et*

al., 2019). Much of this progress is attributable to the Global Tree Assessment, and the high

assessment coverage achieved for trees explains the relatively weak effect of genus size on

assessment probability in tree-dominated genera. Similarly concerted and large-scale

endeavours are urgently needed for other plants, and our findings suggest that an explicit focus

on comprehensive assessments of selected big genera could be an important and worthwhile
approach to start to redress the effects of neglect or avoidance of these groups over decades.

We are in no way suggesting that large genera be split, or that there should be an upper limit on
the size of genera. The size of genera is related to both evolutionary and human processes (e.g.
Bentham's preference for fewer, larger genera; Humphreys & Linder, 2009), and the highly
skewed pattern of genus size frequency, with many small and relatively few very large genera, is
common across the tree of life (Strand & Panova, 2015; Sigwart *et al.*, 2018). Although ultimately
arbitrary, there is a broad consensus that 'good' genera (monophyletic, stable and diagnosable)
are useful divisions (Humphreys & Linder, 2009; Vences *et al.*, 2013; Strand & Panova, 2015;
Muñoz-Rodríguez *et al.*, 2023), so big plant genera (and their problems) are unlikely to be split
out of existence (Moonlight *et al.*, 2024). However, regardless of the basis of genus size, our
results show that by treating genus size as an attribute of species, we can quantify the tangible
conservation implications of being a member of a big genus.

We strongly encourage greater focus on and resourcing of the further study of species in big
plant genera in taxonomic, ecological and conservation contexts. Although big genera continue
to represent an immense amount of work to monograph *in toto*, molecular data can be used to
divide large taxa into more manageable subclades; as in *Eugenia* or *Myrcia* (Mazine *et al.*, 2014;
Lucas *et al.*, 2018). An alternative may be the 'foundation monograph' concept (Scotland &
Wood, 2012) – a practical guide to this approach (based on the recent monograph of *Ipomoea*;
Muñoz-Rodríguez *et al.*, 2019) is detailed in Muñoz-Rodríguez *et al.* (2024). We echo earlier
recommendations for increased collaboration across conservation and taxonomy (Nic Lughadha
et al., 2019; Brown *et al.*, 2023a). The success of any new initiative focused on assessing big
genera will depend on a clear-eyed, evidence-based understanding of the challenges likely to be
faced; among these is the higher proportion of Data Deficient species in genera which lack
recent monographs (Grace *et al.*, 2021) and which are likely to include unusually large
proportions of species not yet described (Moonlight *et al.*, 2024). We found that species from big
genera were at least twice as likely to be categorised as Data Deficient as those in monotypic
genera, a result consistent with our expectations. However, this proportion remained relatively
low overall (<10%, Fig 5) and could be further minimised by effective collaboration between
taxonomists, assessors and other local experts to assemble the most complete information
possible for species' assessments. Where assessors identify species likely to be Data Deficient,
this can feed back into research prioritisation (e.g. targeted fieldwork or other data collection).
Similarly, taxonomists can provide input on the likelihood of species re-circumscriptions,
facilitating effective prioritisation of assessments in big genera – thus, a shared pipeline of
activity between taxonomists and assessors can allow extinction risk assessments of species in

speciose genera to be undertaken before a genus is fully monographed (Nic Lughadha *et al.*,

2019).

Species in big genera are undoubtedly conservation priorities; they are under-studied, under-

assessed, and their extinction risk has been underestimated. Over a quarter of angiosperm

species diversity is in big genera, but there is a systematic shortfall in extinction risk data for

these species. Moreover, we found that this knowledge gap is likely to obscure, at least in part,

the greater extinction risk faced by species in big genera than by other angiosperms. While

working with speciose taxa presents ongoing practical challenges, an urgent increase in focus on

studying, assessing and conserving species in big genera is vital to safeguard the future of

angiosperm diversity.

References

Bachman SP, Brown MJM, Leao TCC, Nic Lughadha E, Walker BE. **2024**. Extinction risk
predictions for the world's flowering plants to support their conservation. *New Phytol* **242**:
797–808.
Bachman SP, Field R, Reader T, Raimondo D, Donaldson J, Schatz GE, Lughadha EN.
2019. Progress, challenges and opportunities for Red Listing. *Biological Conservation* **234**:
45–55.
Baldaszti L, Särkinen T, Brummitt N, Cardoso D, Knapp S, Zuanny DC, Moonlight PW.
2025. On the geography of big plant genera. *Annals of Botany*: mcaf159.
Barton K. **2024**. MuMIn: Multi-Model Inference. R package.
Brown MJM, Bachman SP, Nic Lughadha E. **2023a**. Three in four undescribed plant species
are threatened with extinction. *New Phytol* **240**: 1340–1344.
Brown MJM, Walker BE, Black N, Govaerts RHA, Ondo I, Turner R, Nic Lughadha E. **2023b**.
rWCVP: a companion R package for the World Checklist of Vascular Plants. *New Phytol*
240: 1355–1365.
Brummitt RK. **2001**. *World geographical scheme for recording plant distributions*. Pittsburgh:
Hunt Inst. for Botanical Documentation.
Brummitt N, Cui Q, Bai P, Liu X, Scott B, De Palma A, Araujo AC. **2024**. Reducing the
number of Data Deficient plant species. *Systematics and Biodiversity* **22**: 2302180.
**Bureš P, Elliott TL, Veselý P, Šmarda P, Forest F, Leitch IJ, Nic Lughadha E, Soto Gomez
M, Pironon S, Brown MJ**. **2024**. The global distribution of angiosperm genome size is
shaped by climate. *New Phytologist* **242**: 744–759.
**Cowell C, Williams E, Bullough LA, Grey J, Klitgaard B, Govaerts R, Andriambololonera S,
Cervantes A, Crameri S, Lima HC**. **2022**. *CITES Dalbergia Checklist*. Commissioned by
the CITES Secretariat. Royal Botanic Gardens, Kew.
**Darbyshire I, Timberlake J, Osborne J, Rokni S, Matimele H, Langa C, Datizua C, de Sousa
C, Alves T, Massingue A, *et al.* 2019**. The endemic plants of Mozambique: diversity and
conservation status. *PhytoKeys* **136**: 45–96.
**Davies TJ, Smith GF, Bellstedt DU, Boatwright JS, Bytebier B, Cowling RM, Forest F,
Harmon LJ, Muasya AM, Schrire BD**. **2011**. Extinction risk and diversification are linked
in a plant biodiversity hotspot. *PLoS biology* **9**: e1000620.
Freeman BG, Pennell MW. **2021**. The latitudinal taxonomy gradient. *Trends in Ecology &
Evolution* **36**: 778–786.
Frodin DG. **2004**. History and concepts of big plant genera. *Taxon* **53**: 753–776.
Fu Q, Huang X, Li L, Jin Y, Qian H, Kuai X, Ye Y, Wang H, Deng T, Sun H. **2022**. Linking
evolutionary dynamics to species extinction for flowering plants in global biodiversity
hotspots. *Diversity and Distributions* **28**: 2871–2885.

Govaerts R, Nic Lughadha E, Black N, Turner R, Paton A. **2021**. The World Checklist of Vascular Plants, a continuously updated resource for exploring global plant diversity. *Sci Data* **8**: 215.

Grace OM, Pérez-Escobar OA, Lucas EJ, Vorontsova MS, Lewis GP, Walker BE, Lohmann LG, Knapp S, Wilkie P, Sarkinen T. **2021**. Botanical monography in the Anthropocene. *Trends in Plant Science* **26**: 433–441.

Greenberg DA, Pyron RA, Johnson LG, Upham NS, Jetz W, Mooers AØ. **2021**. Evolutionary legacies in contemporary tetrapod imperilment. *Ecology Letters* **24**: 2464–2476.

Guedes JJ, Moura MR, Jardim L, Diniz-Filho JAF. **2025**. Global Patterns of Taxonomic Uncertainty and its Impacts on Biodiversity Research. *Systematic biology*: syaf010.

Humphreys AM, Govaerts R, Ficinski SZ, Nic Lughadha E, Vorontsova MS. **2019**. Global dataset shows geography and life form predict modern plant extinction and rediscovery. *Nature Ecology & Evolution* **3**: 1043–1047.

Humphreys AM, Linder HP. **2009**. Concept versus data in delimitation of plant genera. *Taxon* **58**: 1054–1074.

IUCN. **2012**. *IUCN Red List categories and criteria, version 3.1, second edition*. Gland and Cambridge: IUCN.

IUCN. **2022a**. Guidelines for Using the IUCN Red List Categories and Criteria.

IUCN. **2022b**. Guidelines for Reporting on Proportion Threatened.

IUCN. **2024**. The IUCN Red List of Threatened Species.

Leão TC, Lughadha EN, Reich PB. **2020**. Evolutionary patterns in the geographic range size of Atlantic Forest plants. *Ecography* **43**: 1510–1520.

Lenth RV. **2025**. emmeans: Estimated Marginal Means, aka Least-Squares Means.

Lozano FD, Schwartz MW. **2005**. Patterns of rarity and taxonomic group size in plants. *Biological Conservation* **126**: 146–154.

Lucas EJ, Amorim BS, Lima DF, Lima-Lourenço AR, Nic Lughadha EM, Proença CEB, Rosa PO, Rosário AS, Santos LL, Santos MF. **2018**. A new infra-generic classification of the species-rich Neotropical genus Myrcia sl. *Kew Bulletin* **73**: 1–12.

Mazine FF, Souza VC, Sobral M, Forest F, Lucas E. **2014**. A preliminary phylogenetic analysis of Eugenia (Myrtaceae: Myrteae), with a focus on Neotropical species. *Kew Bulletin* **69**: 1–14.

Moonlight PW, Baldaszti L, Cardoso D, Elliott A, Särkinen T, Knapp S. **2024**. Twenty years of big plant genera. *Proceedings of the Royal Society B* **291**: 20240702.

Muñoz-Rodríguez P, Carruthers T, Wells T, Sumadijaya A, Wood JR, Scotland RW. **2024**. The research behind a taxonomic monograph: a case study from Ipomoea (Convolvulaceae). *Kew Bulletin* **79**: 897–914.

Muñoz-Rodríguez P, Carruthers T, Wood JR, Williams BR, Weitemier K, Kronmiller B, Goodwin Z, Sumadijaya A, Anglin NL, Filer D. **2019**. A taxonomic monograph of Ipomoea integrated across phylogenetic scales. *Nature plants* **5**: 1136–1144.

Muñoz-Rodríguez P, Wood JR, Wells T, Carruthers T, Sumadijaya A, Scotland RW. **2023**. The challenges of classifying big genera such as Ipomoea. *Taxon* **72**: 1201–1215.

Newton A, Oldfield S, Rivers M, Mark J, Schatz G, Garavito NT, Cantarello E, Golicher D, Cayuela L, Miles L. **2015**. Towards a global tree assessment. *Oryx* **49**: 410–415.

Nic Lughadha E, Bachman SP, Leão TCC, Forest F, Halley JM, Moat J, Acedo C, Bacon KL, Brewer RFA, Gâteblé G, *et al.* **2020**. Extinction risk and threats to plants and fungi. *PLANTS, PEOPLE, PLANET* **2**: 389–408.

Nic Lughadha EM, Graziele Staggemeier V, Vasconcelos TNC, Walker BE, Canteiro C, Lucas EJ. **2019**. Harnessing the potential of integrated systematics for conservation of taxonomically complex, megadiverse plant groups. *Conserv Biol* **33**: 511–522.

Purvis A, Agapow P-M, Gittleman JL, Mace GM. **2000**. Nonrandom extinction and the loss of evolutionary history. *science* **288**: 328–330.

R Core Team. **2024**. R: A language and environment for statistical computing.

Rivers M. **2017**. The Global Tree Assessment–red listing the world's trees. *BGjournal* **14**: 16–19.

Schwartz MW, Simberloff. **2001**. Taxon size predicts rates of rarity in vascular plants. *Ecology Letters* **4**: 464–469.

Scotland RW, Wood JR. **2012**. Accelerating the pace of taxonomy. *Trends in Ecology & Evolution* **27**: 415–416.

Sigwart JD, Sutton MD, Bennett KD. **2018**. How big is a genus? Towards a nomothetic systematics. *Zoological Journal of the Linnean Society* **183**: 237–252.

Soto Gomez M, Brown MJ, Pironon S, Bureš P, Verde Arregoitia LD, Veselý P, Elliott TL, Zedek F, Pellicer J, Forest F. **2024**. Genome size is positively correlated with extinction risk in herbaceous angiosperms. *New Phytologist* **243**: 2470–2485.

Strand M, Panova M. **2015**. Size of genera–biology or taxonomy. *Zoologica Scripta* **44**: 106–116.

Svahnström VJ, Nic Lughadha E, Forest F, Leão TCC. **2025**. Geographic range size and rarity of epiphytic flowering plants. *Nature Plants* **11**: 1380–1389.

Tanentzap AJ, Igea J, Johnston MG, Larcombe MJ. **2020**. Does evolutionary history correlate with contemporary extinction risk by influencing range size dynamics? *The American Naturalist* **195**: 569–576.

Vamosi JC, Wilson JRU. **2008**. Nonrandom extinction leads to elevated loss of angiosperm evolutionary history. *Ecology Letters* **11**: 1047–1053.

Vences M, Guayasamin JM, Miralles A, Riva IDL. **2013**. To name or not to name: Criteria to promote economy of change in Linnaean classification schemes. *Zootaxa* **3636**: 201–244.

Walker BE, Tucker A, Nicolson N. **2022**. Harnessing large-scale herbarium image datasets through representation learning. *Frontiers in plant science* **12**: 806407.

Figures and Tables

Table 1. Numbers of species assessed as versus predicted to be threatened in big genera, arranged in descending order of underprediction (Δthreat prop.).

Genus	Climate	# spp.	prop. spp. assessed	# spp. threatened (RL)	prop. threatened (RL)	# spp. predicted threatened	prop. predicted threatened	Δthreat prop.
Stelis	tropical	1340	0.4%	0	0.0%	1060	79.1%	79.1%
Taraxacum	temperate	2490	0.1%	0	0.0%	1243	49.9%	49.9%
Lupinus	multiple climates	580	11.7%	6	8.8%	333	57.4%	48.6%
Lepanthes	tropical	1203	2.2%	11	40.7%	1058	87.9%	47.2%
Saussurea	multiple climates	516	0.6%	0	0.0%	232	45.0%	45.0%
Pedicularis	temperate	679	4.4%	3	10.0%	342	50.4%	40.4%
Ranunculus	temperate	1750	2.9%	11	21.6%	1011	57.8%	36.2%
Elatostema	multiple climates	654	0.5%	1	33.3%	435	66.5%	33.2%
Primula	temperate	545	3.5%	2	10.5%	224	41.1%	30.6%
Corydalis	temperate	538	1.3%	2	28.6%	310	57.6%	29.0%
Pleurothallis	tropical	579	0.7%	2	50.0%	439	75.8%	25.8%
Masdevallia	tropical	651	1.7%	6	54.5%	523	80.3%	25.8%
Piper	tropical	2425	6.4%	65	41.9%	1634	67.4%	25.4%
Epidendrum	tropical	1878	2.2%	17	40.5%	1210	64.4%	24.0%
Maxillaria	tropical	652	1.1%	2	28.6%	341	52.3%	23.7%
Oxytropis	temperate	608	1.5%	1	11.1%	208	34.2%	23.1%

Anthurium	tropical	1325	10.9%	46	31.9%	691	52.2%	20.2%
Ardisia	tropical	737	26.2%	68	35.2%	393	53.3%	18.1%
Berberis	multiple climates	626	9.3%	22	37.9%	350	55.9%	18.0%
Allium	temperate	1078	13.6%	31	21.1%	403	37.4%	16.3%
Top 20 genera shown here, for full list of big genera see Table S7.								

Figure 1. Climate and lifeform distribution in big angiosperm genera. Genera coded by their dominant (≥75% of species) climate/lifeform or as 'multiple climates/lifeforms' (see Methods). Colour and pattern show proportion of species across covariate combinations; note that overall circle sizes are not to scale. For full details on numbers of species and genera across covariates, see Table S1.

Figure 2. Genus size-range size relationship. Panel a) shows overall estimated marginal mean proportions of species under each range size threshold (as number of botanical countries) across genus size. The median range size is one botanical country for all but the very smallest genera. Panel b) shows the estimated marginal mean probability that species are restricted to a single botanical country across lifeform and climate (note that climate and lifeform aggregated at genus level). Shaded grey area shows the 'zone of big genera'; points show the observed proportions of single-country endemics in big genera. Panel c) shows the estimated marginal slope coefficients for family (the single-country endemic threshold; only largest 50 families shown); positive coefficients indicate that species in larger genera are more likely to be restricted to a single botanical country. Units of coefficients on the logit scale. Error bars show 95% CIs of coefficient estimates; colour of points denotes significance and direction; size of points denotes number of species per family. Marginal means and trends estimated across cell weights of the non-focal terms in all panels; see Fig S1 for a graphical representation of how these are estimated from the model. For tabulated results with 95% CI values for marginal climate, lifeform, and family trends, see Table S4; for all trends (not averaged), see Table S5.

Figure 3. Genus size-assessment probability relationship. Panel a) shows estimated marginal means of assessment probability across genus size, climate and lifeform (note that climate and lifeform were aggregated at genus level). Upper limit of genus size values restricted to range of data for each covariate combination. Grey rectangle highlights the 'zone of big genera'; points show the observed proportions of assessed species in big genera. Dashed lines show n.s. trends; solid lines show trends statistically different from zero. Panel b) shows the estimated marginal slope coefficients for family (the single-country endemic threshold; only largest 50 families shown); negative coefficients indicate that species in larger genera are less likely to have a Red List assessment. Units of coefficients on the logit scale. Error bars show 95% CIs of coefficient estimates; colour of points denotes significance and direction; size of points denotes number of species per family. Marginal means and trends estimated across cell weights of the non-focal terms in all panels; see Fig S1 for a graphical representation of how these are estimated from the model. For tabulated results with 95% CI values for marginal climate, lifeform, and family trends, see Table S4; for all trends (not averaged), see Table S5.

Figure 4. Genus size - documented extinction risk relationship (Red List assessments only).
Panel a) shows estimated marginal means for the probability of meeting the criteria for each Red
List category (note that categories are cumulative, e.g. all species assessed as Critically
Endangered also meet the criteria for Endangered and Vulnerable. Panel b) shows estimated
marginal means of being threatened (Vulnerable, Endangered or Critically Endangered) across
genus size, climate and lifeform (note that climate and lifeform were aggregated at genus level).
Upper limit of genus size values restricted to range of data for each covariate combination. Grey
rectangle highlights the 'zone of big genera'; points show the observed proportions of species
assessed as threatened in big genera. Dashed lines show n.s. trends; solid lines show trends
statistically different from zero. Panel c) shows the estimated marginal slope coefficients for
family (only 34 of the largest 50 families shown; 16 of those families have <250 assessed
species so were excluded from this analysis); positive coefficients indicate that species in larger
genera are more likely to be assessed as threatened. Units of coefficients on the logit scale.
Error bars show 95% CIs of coefficient estimates; colour of points denotes significance and
direction; size of points denotes number of species per family. Marginal means and trends
estimated across cell weights of the non-focal terms in all panels; see Fig S1 for a graphical
representation of how these are estimated from the model. For tabulated results with 95% CI
values for marginal climate, lifeform, and family trends, see Table S4; for all trends (not
averaged), see Table S5.

Figure 5. Genus size – Data Deficiency relationship (Red List assessments only). Estimated marginal means of being assessed as Data Deficient across genus size, climate and lifeform (climate and lifeform were aggregated at genus level). Upper limit of genus size values restricted to range of data for each covariate combination. Shaded ribbons show 95% CIs. Dashed lines show n.s. trends; solid lines show trends that are statistically different from zero. For tabulated results with 95% CI values for climate and lifeform trends, see Table S5.

Figure 6. Genus size - extinction risk relationship (Red List assessments and extinction risk predictions). Panel a) shows estimated marginal means across all thresholds of estimated extinction risk. Panel b) shows estimated marginal means of being threatened (assessed or predicted with high confidence) across genus size, climate and lifeform (climate and lifeform were aggregated at genus level). Upper limit of genus size values restricted to range of data for each covariate combination. Grey rectangle highlights the 'zone of big genera'; points show the proportions of species in big genera that are assessed or predicted to be threatened. Dashed lines show n.s. trends; solid lines show trends that are statistically different from zero. Panel c) shows the estimated marginal slope coefficients for family (only largest 50 families shown); positive coefficients indicate that species in larger genera are more likely threatened (either assessed or predicted). Units of coefficients on the logit scale. Error bars show 95% CIs of coefficient estimates; colour of points denotes significance and direction; size of points denotes number of species per family. Marginal means and trends estimated across cell weights of the non-focal terms in all panels; see Fig S1 for a graphical representation of how these are estimated from the model. For tabulated results with 95% CI values for marginal climate, lifeform, and family trends, see Table S4; for all trends (not averaged), see Table S5.

605 **Figure 7**. Comparison of the estimated marginal means for Red List data (red) and combined
606 (Red List + predictions) data (grey) for the relationship between genus size and probability of
607 extinction risk. Dotted lines show where the lines intercept (dashed grey; left) and where the 95%
608 CIs diverge (dotted blue; right), indicating the change from over- to under-estimation of threat.
609

21