Phylogenomic challenges in polyploid-rich lineages: Insights from orthology

inference and reticulation methods using the complex genus *Packera*

(Asteraceae: Senecioneae)

Erika R. Moore-Pollard[1,2,4], Paige A. Ellestad[2], Jennifer R. Mandel[2,3]

1 *Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs,*

Connecticut 06269, USA

2 *Department of Biological Sciences, University of Memphis, Memphis, Tennessee*

38152, USA

3 *ACRE: Institute for Agricultural and Conservation Research and Education, University*

of Memphis, Memphis, Tennessee 38152, USA

4 *Address for correspondence*

Emails and ORCID IDs of all authors:

Erika R. Moore-Pollard, moore.erika.r@gmail.com, 0000-0002-1182-9274

Paige A. Ellestad, pllestad@memphis.edu, 0000-0003-3930-3239

Jennifer R. Mandel, jmandel@memphis.edu, 0000-0003-3539-2991

Abstract

Phylogenomic discordance is pervasive and cannot be fully addressed through increased data alone. Biological processes such as polyploidy, hybridization, and incomplete lineage sorting are major contributors to discordance and must be accounted for to avoid misleading evolutionary interpretations. To better understand how these processes influence phylogenetic reconstruction, we conducted a comprehensive phylogenomic study in the complex genus *Packera*. With over 90 species and varieties (40% of which exhibit polyploidy, aneuploidy, or other cytological complexities), *Packera* presents significant challenges for phylogenetic reconstruction. Given these complexities, we assessed the impact of different published orthology inference methods on the resulting evolutionary relationships and phylogenetic support of this group. We then applied three paralog-processing methods to evaluate their impact on tree topology and our understanding of *Packera*'s evolutionary history by constructing a time-calibrated phylogeny, reconstructing historical biogeography, and testing for ancient reticulation. Unsurprisingly, phylogenetic outcomes varied depending on the paralog processing method used, with no method performing the best overall over others. Our findings highlight the profound impact of orthology inference and paralog processing on phylogenomic analyses, particularly in polyploid-rich groups such as *Packera*, and we offer guidance on methodological impacts along with practical recommendations. We note that gaining a robust understanding of *Packera*'s evolutionary history requires more than computational approaches alone and while technological advancements have greatly expanded our ability to analyze genomic data, effective phylogenomic research still relies on strong taxon sampling and detailed species knowledge. Without careful attention to the biological context, such as species

45   boundaries, cytological variation, and ecological interactions, phylogenomic studies risk

46   misinterpreting evolutionary history and processes. By accounting for these factors, we

47   can begin to improve the accuracy of evolutionary reconstructions and gain deeper

48   insights into the complex history of plant diversification.

50   <u>Keywords:</u> Compositae; discordance; hybridization; paralogy; molecular systematics

Introduction

The field of phylogenetics is drastically changing with the increased use of multi-locus and genomic data. While this expansion has enabled deeper insights into evolutionary relationships across the tree of life, it has also revealed persistent and complex challenges in phylogenetic inference. A significant challenge is gene tree discordance, where trees estimated from individual loci disagree with each other and/or with the inferred species tree (Degnan and Rosenberg 2009; Mendes et al. 2016). These incongruences are not simply a consequence of limited data; rather, they often reflect differences driven by underlying biological processes such as incomplete lineage sorting (ILS), gene duplication and loss, and hybridization or introgression (Pease et al. 2018). Such processes are especially prevalent in plants, where whole-genome duplications (WGDs), ancient hybridization, and reticulate evolution are widespread and recurrent (Funk 1985; Vision et al. 2000; Wendel and Cronn 2003; Peirson et al. 2012; Panchy et al. 2016; Rose et al. 2021).

Polyploidy, or WGD, is a dominant evolutionary force in angiosperms, with estimates suggesting that up to 70% of extant flowering plants have polyploid ancestry (Grant 1971; Masterson 1994; Soltis and Soltis 1999; One Thousand Plant Transcriptomes Initiative 2019). These events gave rise to extensive paralogy in plant genomes, which poses significant challenges for orthology inference and phylogenetic analysis. Although these issues are particularly pronounced in plants, they are by no means exclusive to them. Similar complications have been documented in non-plants, such as African clawed frogs (Evans et al. 2004), desert rodents (Gallardo et al. 2004), alpine butterflies (Nice et al. 2013), and fish (Braasch and Postlethwait 2012), indicating that complex histories of gene duplication and horizontal gene transfer create

phylogenetic uncertainty across the tree of life. As the scale and scope of phylogenomic

datasets expand, so too does the need to carefully model and interpret the biological

processes underlying gene tree discordance. Fortunately, the growing availability of

phylogenomic data now makes it possible to start untangling these complex histories in

ways that were not previously feasible; however, doing so requires tools and models

that explicitly account for these processes.

In response, a number of bioinformatic tools and pipelines have been developed

to infer orthology under increasingly realistic evolutionary models. For instance,

softwares such as HybPiper (Johnson et al. 2016), Phyluce (Faircloth 2016), and

Captus (Ortiz et al. 2023) implement distinct strategies for identifying and handling

putative paralogs, with some pipelines retaining only a single gene copy (e.g., HybPiper

and Captus), while others aim to exclude all loci showing any sign of paralogy (e.g.,

Phyluce). Although Phyluce has shown greater success in some groups (e.g.,

Antennaria Gaertn.; Thapa et al. 2020), HybPiper typically outperforms across most

plant lineages because of the issues associated with polyploidy. Time will tell how

newer tools such as Captus perform compared to Phyluce or HybPiper.

Additional pipelines have been developed that implement distinct strategies for

identifying and handling putative paralogs using graph-based or tree-based methods

(e.g., paragone-nf; Yang and Smith 2014; Jackson et al. 2023), whereas others

accommodate paralogy during species tree estimation itself (e.g., ASTRAL-Pro; Zhang

et al. 2020; Zhang and Mirarab 2022a). These different approaches can yield markedly

different phylogenetic outcomes, particularly in polyploid-rich systems where paralogs

may be the rule rather than the exception. Yet, few studies have systematically

compared these methods to understand their impact on phylogenetic and downstream

evolutionary inferences.

This is especially important for groups such as *Packera* Á. Löve & D. Löve

(Asteraceae: Senecioneae), a North American genus of about 91 species known for

complex patterns of hybridization, polyploidy, and poorly resolved taxonomy. An

estimated 40% of *Packera* taxa present polyploidy, aneuploidy, and other cytological

disturbances (Barkley 1988; Trock 2006), complicating phylogenetic reconstruction of

this group (Bain and Golden 2000). Given this, *Packera* presents an ideal system to test

the effects of orthology inference differences given its extensive polyploidy, ecological

and geographic diversity, and history of reticulation—all factors that increase the

potential for gene tree discordance.

Our study provides a broadly applicable framework for evaluating orthology

inference and its downstream impacts in systems shaped by gene duplication,

reticulation, and rapid diversification. Here we present the most comprehensive

phylogenomic investigation of *Packera* to date, sampling nearly 93% of the genus and

leveraging a high-resolution Hyb-Seq dataset. We applied a robust framework for

evaluating the impact of paralog processing by comparing multiple orthology inference

methods, including pipelines that remove all paralogs, those that model duplication and

loss, and those that retain a single representative gene copy. We assessed how these

approaches influence gene tree concordance, species tree topology, and node support.

We then tested how three of these methods–a standard orthology inference method, a

method that removes all paralogous loci, and a selective orthology method (see

methods below)–alter downstream macroevolutionary analyses including divergence

dating, historical biogeography, and detection of ancient introgression. Our results

demonstrate that the choice of paralog processing approach strongly influences inferred

species relationships, gene tree concordance, node support, and evolutionary

inferences, highlighting the profound effect that orthology inference decisions can have

in phylogenomics.

Materials & Methods

Specimen collection

A nearly complete sampling of currently recognized *Packera* species and

varieties were used in this study (62 *Packera* species, 22 varieties, and one

nothospecies; totaling 85 *Packera* taxa). *Packera* species were selected based on

information published in the International Plant Names Index (IPNI; available at

http://www.ipni.org) and then curated from discussions with botanists that have

previously specialized in *Packera* (D. Trock & J. Bain, pers. comm.). Outgroup taxa from

tribes Senecioneae (n = 19) and Anthemideae (n = 4) were selected by consulting

recent studies of the group (Bain and Golden 2000; Pelser et al. 2010; Mandel et al.

2019). Additionally, three taxa from tribe Calenduleae were included for downstream

analyses that require more distantly related outgroups that do not have a known WGD

event (Barker et al. 2016; Huang et al. 2016; Zhang et al. 2021). A complete list of the

111 sampled species, herbarium vouchers, and their publication status can be found in

Table S1.

DNA extraction and sequencing

Leaf tissue was obtained from herbarium sheets or live plants that were dried in

silica beads. Dried leaf tissue (~25 mg) was ground using a Bead Mill 24 Homogenizer

(ThermoFisher Scientific, Atlanta, Georgia, USA), and total DNA was extracted using

the Omega Bio-Tek Kit (Atlanta, Georgia, USA) following manufacturer's protocols with

the addition of polyvinylpyrrolidone (PVP) and ascorbic acid to the first extraction buffer

(10 mL SQ1 buffer, 100 mg PVP, 90 mg ascorbic acid). Samples of *Packera* and

outgroup species that were obtained from other researchers followed different DNA

extraction protocols which are outlined in Table S1. All DNA samples used in this study

were quality checked using a Nanodrop 2000 (Thermo Fisher Scientific, Carlsbad,

California, USA) and quantified using a Qubit High Sensitivity assay (ThermoFisher

Scientific, Oregon, USA). If the DNA had low concentrations or possible contaminants,

the samples were cleaned following the E.Z.N.A. Cycle Pure Kit (Omega Biotek,

Georgia, USA) and were re-quantified with the Qubit HS assay. Up to 1 μg DNA per

sample was sonicated with a QSonica machine (ThermoCube, New York, USA) to

generate fragment sizes within the 400-500 base pair (bp) range. The amount of DNA

used is dependent on the initial total DNA concentration of the targeted sample. Time

allotted for sonication of each sample varied depending on approximate length and

degree of fragmentation of the DNA. Prior to sonication, total DNA was visualized on a

1% agarose gel in 1X TBE and GelRed 3x (Biotium) and the approximate length was

estimated by comparing to a 100 bp GeneRuler DNA ladder (ThermoFisher Scientific,

Vilnius, Lithuania). If the sample of DNA appeared smaller than 400 bp, the samples

were not sonicated.

Libraries for each sample were prepared using the NEBNext Ultra II DNA Library

Prep Kit and NEBNext Muliplex Oligos for Illumina Kits (New England Biolabs, Ipswich,

Massachusetts, USA) following the NEBNext Ultra II Version 5 protocol with size

selection on DNA fragments at 300-400 bp range. To make library preparation more

economic, the protocol was adjusted by halving all amounts of reagents and DNA used.

The barcoded libraries (NEBNext multiplex oligos) were quality checked with an Agilent

Bioanalyzer (Agilent Technologies, Santa Clara, California, USA) and quantified by

Qubit High Sensitivity assay. Libraries were then used for targeted sequence capture

with the custom probe set MyBaits Compositae-1061 kit (Mandel et al. 2014) from Arbor

Biosciences (Ann Arbor, Michigan, USA) following manufacturer's protocols (version

2.3.1). Approximately 500ng of total DNA library in 6 µl of library, or equal amount of

DNA per sample when pooling (maximum of 8 samples), was used in each MyBaits

reaction for hybridization with the baits for 36 hours. Hybridized targets were recovered

using Dynabeads M-280 Streptavidin (ThermoFisher Scientific, Vilnius, Lithuania)

following the manufacturer's protocol. Captured targets were amplified and quantified

using KAPA library quantification kits (Kapa Biosystems, Wilmington, Massachusetts,

USA). A 1:4 ratio of library was added to each of the captured target pools to obtain off-

target plastid sequence data. Sequence stats were obtained using the *pxlssq* function in

Phyx v. 1.3.1 (Table S2) (Brown et al. 2017).

Sequence processing

Raw FASTQ data were cleaned using the program Trimmomatic v. 0.36 (Bolger

et al. 2014) which removes Illumina adapters from low-quality bases and adapter

contamination. The Sliding Window quality filter was used in this analysis (illuminaclip

2:30:10, leading 20, trailing 20, sliding window 5:20). Cleaned reads were retained

when they had a minimum length of 36 bp and corresponding forward and reverse

pairs.

Plastid assembly

Chloroplast sequences were extracted from trimmed paired end reads using two

approaches. First, the GetOrganelle v. 1.7.7.1 toolkit (Jin et al. 2020) was used to

assemble circularized plastomes for all sequences with sufficient off-target sequencing

data. The recommended parameters for embryophyte plant plastomes were used but

the number of rounds and the disentangle time limit were increased (-R 75 and --

disentangle-time-limit 7200). All complete circular plastomes were annotated using

GeSeq (Tillich et al. 2017). From these, a *Packera* representative plastome was chosen

and used as a reference. Second, all reads were mapped to the reference plastome

using Bowtie2 (Langmead and Salzberg 2012), variants were called using Samtools

and BCFtools (Danecek et al. 2021), and consensus sequences were created using

seqtk (https://github.com/lh3/seqtk). For the aligned consensus sequences, model

selection and maximum likelihood phylogenetic analysis were then conducted using IQ-

TREE v. 2.2.2 (Minh et al. 2020) and visualized using 'ggtree' (Yu et al. 2017) in R v.

4.3.1 (R Core Team 2016; RStudio 2020).

Nuclear ortholog assembly

Cleaned FASTQ data were then assembled using the HybPiper v. 2.1.6

(Johnson et al. 2016) pipeline to match the reads to the target loci contained in the

Compositae-1061 probe set (Mandel et al. 2014). In HybPiper, the command 'assemble'

was used to map the paired-end reads to the target loci using BWA v. 0.7.17 (Li and

Durbin 2009), then assembled into contigs with specified kmer lengths (21, 33, 55, 77,

99) using SPAdes v. 3.5 (Bankevich et al. 2012). Paralog detection was carried out

using the 'paralog_retriever' command

(https://github.com/mossmatters/HybPiper/wiki/Paralogs), which flags loci that indicate

potential paralogs, then extracts the assemblies that were flagged. A multifasta was

generated for each locus containing the sequences of that specific locus for all taxa in

the run, then aligned into supercontigs using MAFFT v. 7.407 (Katoh and Standley

2013).

Phylogenetic analyses

Given the complex history of hybridization and polyploidization events in

Packera, along with the large number of paralogs flagged from HybPiper in the current

study (Table S3), we sought to compare how different published orthology inference

methods and species tree estimation tools impacted the inferred evolutionary

relationships and support of this group. Our dataset is uniquely adapted to tackle this

issue given the dense taxon sampling, high locus recovery across both orthologs and

paralogs, combined with decades of natural history and systematics work in the group,

enabling more thorough investigations into these challenges.

We generated multiple nuclear phylogenies utilizing different paralog processing

tools to infer orthology and species tree estimation: Concatenation (individual gene

histories not accounted for), HybPiper with ASTRAL-III (hereafter 'ASTRAL'; putative

single-copy orthologous loci are chosen based on coverage depth and sequence

similarity concerning the reference then inferred using coalescent methods), HybPiper

with wASTRAL (hereafter 'wASTRAL'; similar to ASTRAL but low support gene tree

branches are contracted prior to species tree estimation), ASTRAL-Pro (all single-copy

and multi-copy genes identified by HybPiper are considered, and the species tree that

optimizes a quartet similarity measure is chosen), and the 1-to-1, MI, MO, and RT

paralog processing methods by (Yang and Smith 2014) with species trees inferred

using ASTRAL-III (further details below; refer to Table S3). By performing intentional

comparisons among methods, we aim to gain a better understanding of the impact of

paralogy on phylogenomic studies, and ultimately increase phylogenomic support and

concordance across species of complex lineages.

Concatenation

244   Aligned supercontigs from MAFFT were concatenated with FASconCAT-G v.

245   1.02 (Kück and Longo 2014) then implemented in PartitionFinder v. 2.1.1 (Lanfear et al.

246   2012) to determine the most appropriate nucleotide substitution model. The overall best

247   fitting model (Table S4) was used to build a concatenated nuclear maximum likelihood

248   tree in RAxML v. 8.1.3 (Stamatakis 2014) with 1,000 bootstrap (BS) replicates.

249   ASTRAL and wASTRAL

250   For ASTRAL and wASTRAL analyses, nuclear gene alignments were separated

based on the most suitable nucleotide substitution model for each gene as determined

by PartitionFinder (Table S4), then individual gene trees were generated using RAxML.

For the ASTRAL analysis, the gene trees were implemented in ASTRAL-III v. 5.7.3

(Zhang et al. 2018) to generate the species trees. For the wASTRAL analysis, or

weighted ASTRAL, a species tree was generated using 'astral-hybrid' in the ASTER

program (https://github.com/chaoszhang/ASTER). This method has shown to be more

accurate than typical ASTRAL methods since wASTRAL incorporates gene tree

uncertainty by weighting quartets based on the confidence levels of the input gene

trees. In doing so, wASTRAL reduces the influence of erroneous gene trees, leading to

a more reliable species tree estimation (Zhang and Mirarab 2022b). Local posterior

probability (LPP) values were estimated for branch support and indicate the probability

that the branch is the true branch given the set of gene trees provided. LPP is

considered the more reliable clade support measure when running ASTRAL analyses

since LPP is computed based on a quartet score and assumes incomplete lineage

sorting (ILS) (Sayyari and Mirarab 2016).

ASTRAL-Pro

ASTRAL-Pro2 (ASTRAL for PaRalogs and Orthologs) v. 1.13.1.3 (Zhang et al.

2020; Zhang and Mirarab 2022a), hereafter referred to as ASTRAL-Pro, provides a

quartet-based species-tree inference method that allows multi-copy gene trees to be

used as input. Thus, all paralogous and orthologous sequences are used to generate

the species tree instead of selecting a single ortholog to represent an individual gene's

history (e.g., ASTRAL method). By accommodating multi-copy gene trees, ASTRAL-Pro

accounts for gene duplication and loss events, reducing biases that arise from treating

paralogs as true orthologs. This approach is supposed to improve species tree

accuracy, particularly in datasets with extensive gene family expansions, by leveraging

more information from complex gene histories. The ASTRAL-Pro pipeline is similar to

ASTRAL, as described above, but differs by having the user generate separate

alignments and conduct maximum likelihood analyses on the genes that are flagged as

paralogs and single-copy contigs, separately. Single-copy and paralogous gene

sequences were independently aligned using MAFFT, and gene trees were generated

under the best-fitting nucleotide substitution model from PartitionFinder (GTR, GTR+Γ,

GTR+I+Γ; Table S4) in RAxML with 1,000 BS replicates. The RAxML output files were

then used as input for ASTRAL-Pro2 to generate the species tree with LPP values at

each node.

Paragone-nf

Paralog processing was then carried out using the tree-based orthology

inference strategies originally proposed by (Yang and Smith 2014) via the paragone-nf

v. 1.1.0 pipeline (Jackson et al. 2023) (https://github.com/chrisjackson-

pellicle/paragone-nf). This pipeline contains a Nextflow v. 23.10.0 (Di Tommaso et al.

2017) script downloaded through a Singularity v. 3.7.1 container (Kurtzer et al. 2017,

2021) to investigate the loci flagged as paralogous from HybPiper and determines which

loci can be retained or pruned given specified conditions (Yang and Smith 2014;

Morales-Briones et al. 2021a). All orthology inference methods proposed by (Yang and

Smith 2014)–the 'one-to-one orthologs' (1-to-1), 'maximum inclusion' (MI),

'monophyletic outgroup' (MO), and 'rooted ingroup' (RT) methods–were tested. Briefly,

the 1-to-1 approach is the simplest strategy in that it only retains homologs that have no

duplications. The MI approach extracts the subtree with the highest number of taxa

without taxon duplications. The MO approach filters for homolog trees where outgroup

taxa (here tribe Calenduleae: *Calendula arvensis*, *Tripteris pinnatilobatum* (Norl.) B.

Nord., and *Chyrsanthemoides incana* (Burm.f.) Norl.) are monophyletic and non-

repeating. The RT approach searches subtrees of ingroup taxa and roots them by the

designated outgroups, then removes the ingroup taxa as rooted trees. In turn, inferred

gene duplications, or paralogs, are then pruned from root to tip in the rooted tree (Yang

and Smith 2014; Morales-Briones et al. 2021a).

To aid in consistency across analyses, the HybPiper run described above was

used as input instead of using the hybpiper-nf pipeline (Jackson et al. 2023), which is

another NextFlow script generated by the authors of the paragone-nf pipeline. The three

Calenduleae taxa were considered the internal outgroups since this tribe has not

experienced a recent whole genome duplication event, which is required for the MI and

MO analyses (Yang and Smith 2014). The resulting trimmed alignments produced via

paragone-nf were used as input for RAxML to generate gene trees, and a species tree

with LPP support values was generated using ASTRAL-III following the 'ASTRAL'

protocol described above.

The normalized quartet score was calculated for all orthology inference methods

that generated a species tree with ASTRAL-III using the '-q' option. Branch lengths were

recalculated from coalescent units to nucleotide substitutions per codon site for each

ASTRAL-III species tree using IQTree. Parsimony informative sites (PIS) and pairwise

identity (PI) were calculated for each gene matrix using the

'parsimony_informative_sites' and 'pairwise_identity' functions, respectively, in phykit v.

2.0.1 (Steenwyk et al. 2021). Nuclear and plastome trees were visualized using the

package 'phytools' (Revell 2012) in R v. 4.0.5.

Measuring discordance

Species trees from concatenation, plastid, ASTRAL, wASTRAL, and paragone-nf

(1-to-1, MI, MO, and RT) were analyzed for underlying discordance using Quartet

Sampling (Pease et al. 2018). ASTRAL-Pro was not analyzed as Quartet Sampling

does not allow multi-copy gene trees as input. Quartet Sampling assesses the

confidence, consistency, and informativeness of internal tree relationships, and the

reliability of each terminal branch by evaluating if a phylogeny has a lack of support due

to lack of information, discordance from ILS or introgression, or misplaced or erroneous

taxa. Three scores are calculated for each internal branch of the focal tree which

determines the consistency of information (Quartet Concordance [QC], the presence of

secondary evolutionary histories (Quartet Differential [QD], and the amount of

information (Quartet Informativeness [QI]). Quartet Fidelity (QF) scores are calculated at

each tip and reports how frequently a taxon is included in concordant topologies.

Quartet Sampling efficiently synthesizes phylogenetic tests and offers more

comprehensive and specific information on branch support than conventional measures

(i.e., BS or LPP; (Pease et al. 2018). Using the rooted species trees and concatenated

gene alignments from FASconCAT-G (described above), Quartet Sampling was run

using default settings on the designated analyses. The concatenation, plastid, ASTRAL,

wASTRAL, 1-to-1, MI, and MO species trees were rooted to the outgroup Calenduleae

taxon, *Tripteris pinnatilobatum* (Norl.) B. Nord., while the RT tree was rooted to

Doronicum pardalianches L. using the *pxrr* function in Phyx (Table S3).

Additionally, PhyParts v. 0.0.1 (Smith et al. 2015) was used with default settings

to quantify and visualize discordance in the ASTRAL, wASTRAL, and paragone-nf (1-to-

1, MI, MO, and RT) trees. Similar to Quartet Sampling, ASTRAL-Pro could not be

analyzed because of multi-copy gene trees. Additionally, the concatenation and plastid

trees could not be analyzed since those analyses do not use individual gene histories

as input. Resulting trees containing pie charts indicating whether gene trees were

supported, conflicted, or uninformative at each node were generated using a custom

python script

(https://github.com/mossmatters/phyloscripts/tree/master/phypartspiecharts). PhyParts

requires rooted gene trees and a rooted species tree as input. Similar to Quartet

Sampling, ASTRAL, 1-to-1, MI, and MO gene trees were rooted to *Tripteris*

pinnatilobatum and the RT tree was rooted to *Doronicum pardalianches* using the *pxrr*

function in Phyx. Rooting gene trees for PhyParts decreased the number of genes used

as input since not all gene trees contained the targeted outgroup taxon (see Table S3).

For phylogenies generated with ASTRAL technologies–ASTRAL, wASTRAL,

ASTRAL-Pro, and paragone-nf trees (Table S3)–introgression and ILS were visualized

across lineages using phytop v. 0.3.1 (Shang et al. 2025) using default settings. The

species trees were re-ran with the flag '-t 2' added to be used as input for phytop.

Tree topology comparison

To evaluate similarity among the all nuclear gene trees, pairwise adjusted

Robinson-Foulds (RF_{adj}) distances were calculated as outlined by (Mitchell et al. 2017)

between all trees using the *RF.dist()* function in package 'phangorn' (Schliep 2011) in R,

with unrooted trees as input. The Calenduleae outgroup taxa were trimmed from the

trees in R using the *drop.tip()* function in package ape (Paradis and Schliep 2019) to

make concatenation, ASTRAL, ASTRAL-Pro, and paragone-nf trees directly

comparable. The RF distance was adjusted by setting the "normalize" argument to

TRUE, which divides RF by the maximal possible distance as described in (Steel and

Penny 1993). Given that the input trees are unrooted and binary, this value is then 2n-6,

where n is the number of nodes on the tree (resulting equation: $RF_{adj} = RF/(2n-6)$). The

RF_{adj} values range from zero to one, indicating whether the tree topologies are identical

or completely dissimilar, respectively. A principal component analysis (PCA) was

generated from all RF_{adj} values in R to aid in visualization.

All nuclear and plastid phylogenies and their associated analyses described

above will be provided and compared, but for simplicity only the ASTRAL, 1-to-1, and

MO trees will be analyzed in further analyses. ASTRAL was chosen as it selects the

putative single-copy orthologs less rigorously than the remaining methods, creating a

baseline method to compare to. We then wanted to compare ASTRAL to a method that

disregards genes containing paralogs (1-to-1) and a more rigorous paralog-processing

method (MO). We chose MO over MI and RT due to its more stringent ortholog

identification, which minimizes the inclusion of paralogs and reduces phylogenetic

noise. While RT and MI retain larger gene sets by tolerating ambiguous orthologs, they

increase the risk of incorporating non-orthologous sequences, leading to potential

topological artifacts and reduced tree accuracy. Additionally, the conservative filtering of

MO enhances alignment quality and computational efficiency, providing a more reliable

foundation for downstream phylogenetic analyses (Yang and Smith 2014; Morales-

Briones et al. 2021a; Jackson et al. 2023). Unfortunately, the paralog processing

method that includes all genes, ASTRAL-Pro, could not be included since further

analyses require only single-copy orthologs and the underlying gene trees of ASTRAL-

Pro are single- and multi-copy.

Molecular clock dating

Estimates of divergence times were calculated on the ASTRAL, 1-to-1, and MO

trees using RelTime (Tamura et al. 2012, 2018), which utilizes the relative rate

framework (RRF) and has been shown to be efficient for larger datasets and produces

similar results to Bayesian approaches than other non-Bayesian dating methods such

as treePL (Smith and O'Meara 2012; Tao et al. 2020a, 2020b; Barba-Montoya 2021;

Costa et al. 2022). RelTime was implemented using the command-line version of the

MEGA software, MEGA-CC v. 11.0.11 (Kumar et al. 2012, 2016), with necessary input

files generated with MEGA-X (Kumar et al. 2018; Mello 2018). The three Calenduleae

taxa were used as outgroup taxa for this analysis. Since the branch lengths of the three

phylogenies were in coalescent units from ASTRAL-III, branch lengths of each species

tree were recalculated to the number of nucleotide substitutions per codon site prior to

running RelTime using the GTR+I+Γ model in IQTree.

There are no fossils available within *Packera* or tribe Senecioneae, so instead,

multiple dating scenarios were tested utilizing both primary and secondary age

constraints (Table S5). Briefly, a maximum (59.95 MY) and minimum (31 MY) age was

tested for the Anthemideae/Senecioneae split given those dates are the highest and

lowest values within the 95% CI from a previous diversification rate analysis of

Asteraceae (Mandel et al. 2019). Additionally, a maximum age of 34.06 MY was set at

the *Packera* crown node from the upper age limit provided by (Mandel et al. 2019).

Next, a minimum age constraint of 12 MY was tested within the outgroup-Anthemideae

clade given the presence of a fossil pollen (Wang 2004; Hobbs and Baldwin 2013;

Palazzesi et al. 2022). Finally, a maximum age constraint of 14.3 MY was tested for

Pericallis D.Don (Senecioneae), a genus endemic to the Macaronesia islands (Jones et

al. 2014), since that is the estimated age of Porto Santo, the oldest volcanic archipelago

within the Madeira Islands (McDougall and Schmincke 1976; Geldmacher et al. 2000).

Results of all tested scenarios were compared for biological relevance, though only one

scenario was chosen for the discussion. This approach allowed us to focus on the

topology that best aligned with prior systematic understanding and showed the

strongest overall support across analytical frameworks.

Biogeography

Ancestral ranges were estimated on the dated phylogenies using the package

BioGeoBEARS v. 1.1 (Matzke 2014) in R. BioGeoBEARS estimates historical,

biogeographic ranges by implementing popular models in a likelihood framework. Some

of the most common models used are dispersal-extinction-cladogenesis (DEC; (Ree

and Smith 2008), the likelihood version of DIVA (DIVALIKE; (Ronquist 1997), and the

BayArea likelihood version of the range evolution model (BAYAREALIKE; (Landis et al.

2013). In BioGeoBEARS, founder-event speciation can be added to any of these

models (+*J*) with flexibility in determining relative probability. Likelihood Ratio Test (LRT)

and Akaike Information Criterion (AIC) was used to determine which model best fit the

data.

All *Packera* taxa, along with their four most closely related outgroup genera

(*Pericallis*, *Elekmania* B.Nord., *Werneria* Kunth, and *Xenophyllum* V.A.Funk), were

coded as present or absent in ten geographic ranges outlined in Takhtajan (1986)

(Table S6): Arctic Province, Canadian Province, Appalachian Province, Atlantic and

Gulf Coast Plain Province, North American Prairies Province, Rocky Mountain and

Vancouverian Province, Madrean Region, Asia, Europe, and Central America.

Distributions were determined from previously published literature (Freeman and

Barkley 1995; Trock 1999, 2006; Gramling 2006; Mahoney and Kowal 2008; Elven et al.

2011; Yeatts et al. 2011). Only four of the 26 outgroup taxa were included to limit the

number of geographic ranges used in the analysis. The remaining outgroup taxa were

pruned from the dated tree generated with RelTime using the *pxrmt* function in Phyx.

Each taxon was limited to a max distribution of four ranges.

Reticulation

The Species Networks applying Quartets (SNaQ) function in the Julia v. 1.8.1

(Bezanson et al. 2018) package PhyloNetworks v. 0.15.2 (Solís-Lemus and Ané 2016;

Solís-Lemus et al. 2017) was used to investigate any evidence for reticulate evolution

among *Packera* taxa. This package uses unrooted gene trees to infer phylogenetic

networks in a pseudolikelihood framework with multi-locus data, while accounting for

ILS and reticulation. We tested the fit of multiple models and gradually increased the

number of reticulation events (h) from 0 to 10 to find which pseudolikelihood score

reached a plateau, given recommendations by (Solís-Lemus et al. 2017). This process

is computationally intense, so only clades containing 14 taxa or less that had counter-

support nodes in the ASTRAL, 1-to-1, and/or MO trees, as determined by a negative

Quartet Concordance (QC) score resulting from Quartet Sampling, were used as input

for PhyloNetworks, with the remaining taxa pruned from the tree using the *pxrmt*

function in Phyx. Given this, five *Packera* clades were investigated for ancient

reticulation events (Figs. 1-3). Additionally, a single species from each major clade was

then used to represent backbone relationships throughout the entire phylogeny, totaling

six separate PhyloNetwork runs that will be referenced as: Backbone: representing 11

taxa that span the entire phylogeny; Eastern: a clade of 12, 13, or 15 species (MO, 1-to-

1, and ASTRAL, respectively) either distributed along eastern North America or having

known affiliations with those species; California: eight taxa with distributions mainly

within California; Mexico: seven (MO and ASTRAL) to eight (1-to-1) taxa with

distributions mainly within Mexico; Rocky: a clade of five taxa that are distributed in the

Central US, most within the Rocky Mountains; and Arctic: nine taxa that are present in

colder climates along the Pacific Northwest and into Canada Table S7). Rocky could not

be run for 1-to-1 since less than five taxa were present in that target clade (Fig. 2).

Hybridization networks were visualized using Dendroscope v. 3.7.6 (Huson and

Scornavacca 2012).

Results

Matrix results

Illumina sequencing resulted in approximately 733.3 million reads across 111

taxa. The minimum and maximum number of reads ranged from 535,934 in *Werneria*

aretioides Wedd. and 20,685,576 in *Senecio vulgaris* L. (full list in Table S2). HybPiper

retained 1,058 of the targeted 1,061 loci across all 111 taxa with 846 of those loci

having paralog warnings (~80%). The number of loci recovered for each taxon ranged

from 566 in *Packera paupercula var. paupercula* (Michaux) Á. Löve & D. Löve to 1,054

in *Calendula arvensis* L. (Table S8). The number of gene trees varied across analyses,

with the least being 285 in the 1-to-1 analysis to 7,388 in the MI analysis (Table S3).

Parsimony informative sites (PIS) also ranged from 2.11% in the plastid (mapping) tree

to 99.64% in the MO tree (Table S3). For species trees generated with ASTRAL-III,

normalized quartet scores were used to measure gene tree discordance. In doing so,

the normalized quartet score values ranged from 0.516 with the RT tree to 0.534 with

the MO tree (Table S3), indicating that ~52-53% of the gene trees match the species

tree in those analyses.

Off-target plastid sequencing resulted in partial (i.e., mapping; n = 111) to whole

(i.e., complete; n = 18) plastomes. Complete plastomes from 7 *Packera* and one

Senecio species ranged from 150,772 bp to 151,728 bp (Fig. S1). The percentage of

missing data for each mapped consensus sequence varied drastically, with *Tephroseris*

newcombei exhibiting the most missing data (93.1%; Table S9, Fig. S2). The complete

plastome supermatrix with 18 taxa had 0.123% PIS while the mapping supermatrix with

111 taxa had 2.11% PIS (Table S3).

Raw read and assembly information for the 111 taxa can be found in Tables S2

& S8. General matrix information and summary statistics for all analyses can be found

in Table S3.

Topological comparisons and discordance

Phylogenetic topologies varied depending on sequence type (e.g., nuclear or

plastid) or orthology inference and phylogenetic tree inference method (i.e., ASTRAL,

ASTRAL-Pro, 1-to-1, etc.) (Figs. 4 & S3). With the nuclear data, *Packera* is generally

placed as the sister lineage to a clade comprising *Pericallis* and *Elekmania*, with the

exception of ASTRAL-Pro which placed *Packera* as sister to *Tephroseris* and in a larger

clade with *Pericallis* and *Elekmania* (Fig. 4). Interestingly, the plastid data showed a

similar relationship to ASTRAL-Pro; however, *Pericallis* is placed unexpectedly in a

separate clade with other Senecionineae taxa. All nuclear phylogenies recovered

monophyletic tribes and subtribes (e.g., Senecionineae and Tussilaginineae s.str.),

though the relationships within the subtribes varied. The mapped plastid tree generally

separated the subtribes, with the exception of *Roldana* (Senecionineae) in

Tussilaginineae s.str. (Figs. 4 & S3). *Tephroseris* (tribe Tephroseridineae) moved

constantly with its most consistent placement being sister to Senecioneae (Fig. 4).

Doronicum is consistently placed sister to Senecioneae, providing support for its

elevation to tribal status (Doroniceae; (Panero 2005; Fu et al. 2016).

Phylogenetic topologies remained highly incongruent independent of methods

used (Figs. 1-3, Table S10). The ASTRAL and wASTRAL tree topologies were most

similar (RF_{adj} = 0.324) (Fig. 4, Table S10), while the plastid tree was consistently the

most dissimilar to the remaining trees (RF_{adj} ranging from 0.895 to 0.914), with the

concatenated tree being the next most dissimilar (RF_{adj} ranging from 0.562 to 0.648;

Table S10). Additionally, the normalized quartet scores from ASTRAL indicate that only

half of the gene trees match the species trees in the ASTRAL and paragone-nf analyses

(Table S3). Most of the topological conflicts were among *Packera* taxa, with minor

differences among the remaining Senecioneae taxa and outgroups (Fig. S3). Support

values varied depending on which sequence type and method was used, though

generally remained low. The wASTRAL and MI trees showed the highest node support

with 58 out of 110 nodes (52.7%) having ≥0.95 LPP (Table S3 & Fig. S3). In contrast,

the plastid and concatenated trees showed much lower resolution with the fewest

number of nodes having high support (36 of 110 nodes, 32.7%, have ≥95 BS; Table S3

& Fig. S3).

Quartet Sampling indicated that underlying gene tree discordance remains high,

independent of the method used. ASTRAL and concatenated trees had the most fully

supported nodes (QS = 1/–/1) when compared to the remaining phylogenies (Fig. S4),

though the number was still low (~10%). Even so, the concatenated tree also had the

highest proportion of counter-support nodes (QC = <0; 26.4%; see Fig. S4 for all

Quartet Sampling results). PhyParts indicated that the ASTRAL and MO trees have the

most concordant gene trees. The wASTRAL analysis had the highest percentages of

gene trees supporting the top alternative bipartition. The MI tree harbored the highest

discordance and uninformative data (Fig. S5).

This underlying discordance could be caused by ILS, as it was common in all

tested ASTRAL trees (Fig. S6). Introgressive hybridization (IH) was not detected without

the presence of ILS in any of the nodes, though IH was most common in the nodes that

were shown to be discordant (Figs. S4 & S6). Interestingly, crown *Packera* consistently

had high IH values, potentially indicating introgression at that node. Backbone nodes

were generally considered well resolved with high confidence (ILS index < 50% and IH

index <10%; (Shang et al. 2025), especially in the outgroups, whereas *Packera* taxa

appeared less resolved.

Molecular clock dating

The resulting divergence rate estimates differed among ASTRAL, 1-to-1, and

MO, with estimated ages of the MO and 1-to-1 trees older and sometimes almost

double the ASTRAL ages. For ASTRAL, the age for crown *Packera* ranged from 11.4

MY in Scenario 5 and 6 to 13.5 MY in Scenario 1-4. Alternatively, the crown age of

Packera were almost doubled in the 1-to-1 and MO analyses, with ranges from ~21 MY

in Scenario 5 to 35.5 MY-35.7 MY in Scenario 4 (Table S11, Fig. S7). Scenario 2 was

chosen as the best-fitting dating model because it incorporated all available primary

calibration points and yielded divergence time estimates that align well with known

patterns of evolutionary history (Pelser et al. 2010; Mandel et al. 2019), while minimizing

reliance on secondary calibrations (Table S5) (Müller and Reisz 2005; Benton and

Donoghue 2007; Parham et al. 2012). The dated trees from all six scenarios can be

found in Fig. S7.

Biogeography

Evaluation of the six different model choices demonstrated that BAYAREALIKE

was the best fit given our data based on its lowest log likelihood (Ln L) and highest AIC

score, as well as the highest weighted AICc score for all three of our analyses (Table

S12). The BAYAREALIKE model is a simplified likelihood interpretation of the BayArea

program described by (Landis et al. 2013), which tests the importance of cladogenesis

on a particular dataset, but it does not depend on geographic distance between taxa

and allows for a larger number of areas. Generally, models including jump speciation

(+*J*) provided a better fit than those without in the DEC and DIVALIKE models, and were

the same or very similar in the BAYAREALIKE models (Table S12), indicating a

potential for jump dispersal or founder-event speciation in our data. Trees of all model

runs are in Fig. S8.

The historical biogeography analysis revealed distinct geographic patterns

across the different orthology methods (Figs. 5 & S8). ASTRAL identified the ancestral

range as concentrated in the North American Prairies Province (N), Rocky Mountain

and Vancouverian Province (R), and Mandrean (M) regions. In contrast, MO showed

the ancestral regions being only the Rocky Mountain and Vancouverian Province (R),

and Mandrean (M) regions. 1-to-1 was intermediate with crown *Packera* being ancestral

to the North American Prairies Province (N), Rocky Mountain and Vancouverian

Province (R), and Mandrean (M) regions, similar to ASTRAL; however, the remaining

backbone was Rocky Mountain and Vancouverian Province (R), and Mandrean (M)

regions, similar to MO. Notably, all analyses showed one major expansion into the

eastern United States–Appalachian Province (P) and Atlantic and Gulf Coast Plain

Province (T)–represented by the Eastern clade (Figs. 5 & S8), with a few instances of

individual introductions to the eastern United States. Many of these taxa originated

initially in the North American Prairies Province (N) and/or R ranges. ASTRAL and MO

showed two early expansions into the Arctic Province (I) and Canadian Province (C) at

the base of the phylogeny, while 1-to-1 showed at least three expansions north. 1-to-1

exhibited the most geographic inconsistency, followed by MO. Even so, geography

proved to be a reliable indicator for larger clades across all analyses, although

widespread species were scattered across the phylogeny and did not always align

consistently within geographic clades (Figs. 1-3 & S8).

Reticulation events

Five *Packera* clades were tested for ancient reticulation events given a large

number of counter-support (QC < 0) nodes from Quartet Sampling on the ASTRAL, 1-

to-1, and MO trees (Figs. 1-3). Additionally, a single species from each major clade was

then used to represent backbone relationships throughout the entire phylogeny, totaling

six separate PhyloNetwork runs named: Arctic, Backbone, California, Eastern, Mexico,

and Rocky (Table S13). Log likelihood values indicated that the most likely number of

reticulation events (h) varied depending on the run (Fig. S9). Additionally, Arctic,

Mexico, and Backbone had varying numbers of optimal reticulation events between the

paralog-processing methods (Table S7, Fig. S9). For ASTRAL, Arctic and Mexico had

the highest number of reticulation events (h = 3), followed by California and Backbone

(h = 2), then Eastern and Rocky (h = 1). For 1-to-1, Backhone had the most (h = 6),

then Mexico (h = 3), Arctic and California (h = 2), and then Eastern (h = 1). For MO,

Arctic, California, and Backbone had the highest number of reticulations (h = 2) with

Eastern, Mexico, and Rocky having the lowest (h = 1) (Figs. 6 & S9, Table S7). The

taxa also differed between analyses given different phylogenetic relationships between

clades (Table S13), impacting some of the networks identified.

Some networks showed different events depending on the phylogenetic method

and subset of data (Fig. 6); however, some consistent patterns emerged. For example,

the Eastern clade continuously showed the same reticulation event regardless of the

paralog processing method, and parentage percentages were very similar. In contrast,

Rocky did not show the same reticulation events, but the networks generally involved

Packera mancosana and *P. streptanthifolia*. The California clade showed differing

results across methods: both ASTRAL and 1-to-1 analyses identified *P. bernardina* as

the product of reticulation with the same parentage, whereas MO did not yield the same

results, and no consistent trends emerged. The Mexico clade was generally

inconsistent, but did feature reticulation events involving *P. neomexicana* and its

varieties. Additionally, *P. texensis* and *P. quercetorum* were involved in reticulation

events in two of the three paralog processing networks. Among all the clades, the Arctic

clade exhibited the most variability in the number of reticulation events, with the *P.*

bolanderi varieties consistently involved in some form of reticulation. However, the

specifics of these events varied depending on sampling and paralog processing

methods. In the Backbone subtree analysis, *Tephroseris* was frequently identified as a

product of reticulation, with 1-to-1 and MO showing the same parentage. In contrast,

ASTRAL did not show the same reticulation events but did identify a reticulation event

involving *Tephroseris* with a different parentage. Notably, the parentage of *Tephroseris*

always included a *Packera* taxon, though the specific taxon varied.

Discussion

Gene duplication was pervasive in *Packera*, with more than 80% of nuclear loci

showing evidence of paralogy (Table S3). Such widespread paralogy poses major

challenges for species tree inference, especially in Hyb-Seq datasets, where probes

may unintentionally enrich for multiple homeologs (Salmon et al. 2012; Akama et al.

2014; Crameri et al. 2022), complicating the recovery of true single-copy orthologs for

phylogenetic reconstructions. Because HybPiper's orthology inference relies largely on

single-copy filtering (Johnson et al. 2016), using its outputs alone can introduce noise

and bias into gene tree reconstruction, ultimately distorting species tree inference.

Consistent with our study, using only putative single-copy orthologs designated by

HybPiper was insufficient for accurate phylogenetic inference in *Packera*, a group

characterized by extensive polyploidy and hybridization (Trock 2006). These challenges

are not unique to *Packera*, as many plant lineages, such as Asparagaceae (Bentz and

Leebens-Mack 2024), Rosaceae (Morales-Briones et al. 2018, 2021a), and

Amaranthaceae (Morales-Briones et al. 2021b), also experience difficulties

reconstructing phylogenies due to hybridization and gene/genome duplications. Similar

issues occur in non-plant systems, including the bacterial order Enterobacteriales

(Adeolu et al. 2016), annelids (Struck 2013), and modern amphibians (Siu-Ting et al.

2019). Together, these examples highlight the complexity of orthology inference in

phylogenomics and the need for rigorous paralog processing across both plant and non-

plant systems.

The most recent phylogeny of *Packera* prior to this study was based on ITS data

and revealed extremely low sequence variation, with only 0–5% divergence and several

Packera taxa sharing identical sequences (Bain and Golden 2000). This led to a largely

unresolved tree, likely due to few phylogenetically informative sites, sparse species

sampling, and potential reticulation events (Funk 1985; Linder and Rieseberg 2004). By

using Hyb-Seq data and broader taxon sampling, we substantially improved

phylogenetic resolution across *Packera*, including within major clades. However, our

analyses also revealed that gene tree concordance and species tree resolution varied

substantially across paralog processing methods (Fig. S3), exemplifying the influence of

orthology inference strategies on downstream phylogenetic outcomes. Methods that

explicitly model gene duplication and loss, such as those implemented in paragone-nf

(Yang and Smith 2014; Jackson et al. 2023), generally yielded more concordant gene

trees and better-resolved species trees (Figs. 2-3). However, substantial conflict

persisted across the trees, indicating that even sophisticated modeling approaches

cannot fully resolve deep relationships in groups with complex histories like *Packera*.

These discrepancies may in part reflect differences in the number and identity of genes

retained after filtering (Table S3). For instance, the 1-to-1 method, which excludes all

loci with paralogs and resembles approaches used in tools like Phyluce, eliminates a

significant portion of the dataset and may oversimplify the data, especially in

duplication-rich systems such as *Packera*. While Phyluce and 1-to-1 filtering both

remove paralogs, the underlying methods differ (Faircloth 2016; Jackson et al. 2023),

causing Phyluce to outperform HybPiper in certain cases (Mandel et al. 2019; Thapa et

al. 2020). Similarly, the RT method naturally loses information since it removes

outgroups to determine orthology (Yang and Smith 2014; Morales-Briones et al. 2021b,

2021a) and potentially limits phylogenetic signals. These findings highlight that while

some approaches have potential to improve concordance, none fully resolve the

underlying discordance of *Packera*, emphasizing that the choice of method, underlying

data, and biological context must guide ortholog selection in phylogenomic analyses.

Results from a previous phylogenetic study of *Packera* showed that its

relationships tend to follow a loose geographic pattern (Bain and Golden 2000), which is

consistent with our findings (Figs. 1-3 & S8). Our nuclear phylogenies showed that

major clades tended to contain taxa within similar geographic ranges, independent of

which orthology inference method was utilized. Additionally, most taxa within these

major clades were conserved across methods, though their placement along the

backbone, and some taxa, did vary (Figs. 1-3). For example, the taxa in the Eastern

clade was fairly consistent between the orthology inference methods, though some taxa

did move in and out of the clade (Fig. S3). Interestingly, some of these taxa are

widespread species that interspecifically hybridize with other *Packera* taxa they come

into contact with. For example, *P. pseudaurea* and its varieties are typically found in

central North America, though their phylogenetic placement is generally in the Eastern

clade. These results are unsurprising as they have an extensive history of hybridization

with other *Packera* taxa in the Eastern clade, which is believed to have created much

taxonomic uncertainty (Barkley 1962; Trock 1999), potentially explaining their

inconsistent phylogenetic placements (Fig. S3).

Our results also reveal that evolutionary interpretations such as divergence

dating and biogeographic reconstruction are highly sensitive to the paralog processing

approach used to infer the underlying phylogeny. Divergence time estimates varied

across methods, with some species trees suggesting that crown *Packera*'s

diversification occurred more recently (ASTRAL: 13.5 MY), while others placed key

divergences deeper in time (1-to-1: 24.2 MY and MO: 25 MY; Fig. S7, Table S11).

These differences are not trivial as they affect our understanding of how *Packera*

responded to climatic shifts, mountain uplift, and other historical events, with broader

implications for comparative studies across Asteraceae and other radiating plant

lineages. For example, tribal- and family-wide studies predicted the crown age of

Packera to be fairly young, between 7.8 MY and 20.2 MY (Pelser et al. 2010; Mandel et

al. 2019; Zhang et al. 2021), which is more in-line with the ASTRAL analysis. While

more *Packera*-specific predictions are older, with (Barkley 1988) predicting the origin of

Packera in the later stages of the mid-Tertiary (~22-37 MY), closely aligning with our

paragone-nf analyses (Fig. S7). Similarly, biogeographic patterns shifted depending on

the inferred topology, with each analysis showing different migration patterns over time

(Figs. 5 & S8). Even so, the broad pattern is relatively unchanged. For example, all

topologies generally agreed that *Packera* originated in western North America with one

major expansion east and then later expansions north from the western clades. From

there, individual taxa expanded or became specialized to certain regions, with most of

the diversity present in western North America. The consistency across topologies lends

confidence that these patterns reflect the underlying evolutionary history within *Packera*,

rather than artifacts of methodological bias.

Despite improvements from paralogy-aware filtering, the species trees for

Packera were not fully resolved, indicating that underlying biological processes such as

ILS, hybridization, and introgression also play significant roles in shaping *Packera*'s

evolutionary history. This is supported by strong cytonuclear discordance, where plastid

phylogenetic relationships vary significantly from the nuclear trees (Figs. 4 & S3),

consistent with chloroplast capture or ancient hybridization events (Doyle 1992;

Maddison 1997; Galtier and Daubin 2008; Soltis and Soltis 2009; Smith et al. 2015; Stull

et al. 2020). Directly testing for gene flow reveals multiple introgression events,

particularly along short internal branches susceptible to ILS (Fig. S6). However, the

specific reticulation patterns recovered varied depending on the paralog processing

method used. These discrepancies highlight a critical point: gene selection—and

whether or how paralogs are retained—directly shapes reticulation inference. Each

gene has its own evolutionary trajectory, shaped by recombination and gene duplication

and loss (Betrán 2015), and filtering strategies that exclude paralogs or collapse

duplications can erase signals of past hybridization. Consequently, some reticulation

events, such as those involving Eastern *Packera* lineages, were consistently recovered

across datasets, while others appeared only in specific analyses.

Our findings emphasize that accurately inferring reticulate evolutionary histories

requires embracing the independent histories of genes and carefully considering how

orthology is defined. These differences should not be dismissed as noise. Rather, they

may reflect the underlying biological complexity of the group. In *Packera*, where

hybridization and genome duplication are prevalent, gene tree heterogeneity is

expected and informative. Our findings emphasize that accurately inferring reticulate

evolutionary histories requires embracing the independent histories of genes and

carefully considering how orthology is defined. Methods that oversimplify gene

relationships risk masking the very patterns needed to understand complex evolutionary

processes.

We note that network analyses such as PhyloNetworks, are highly sensitive to

sampling bias, so even though we sampled 85 out of a targeted 91 *Packera* taxa (~93%

coverage of the genus), missing taxa could impact the results. Including the remaining

and unsampled *Packera* taxa in future studies could help increase our understanding of

these relationships. Additionally, limiting the analyses to specified clades can ignore

other potential reticulation events occurring across groups. We recognize this limit;

however, computational requirements of current reticulation network analyses limit the

ability to sample more taxa or loci (see (Hejase and Liu 2016). With recent bioinformatic

advances and increased interest in non-bifurcating trees, we anticipate that less

computationally intense and efficient phylogenetic and network tools will soon be

developed, allowing researchers to test for reticulations across large phylogenies.

Another necessary consideration with hybridization networks is the idea of ghost

introgressions, which are ancient introgression events that leave genetic signatures of

extinct species in present-day species (Ottenburghs 2020). Accounting for ghost

introgressions has important implications for macroevolutionary studies, such as

improving the accuracy of trait evolution and biogeographic reconstructions, refining

divergence time estimates, and revealing hidden contributions of hybridization to

diversification (Ottenburghs 2020; Tricou et al. 2022). Network methods such as

PhyloNetworks can detect ghost introgressions, but these methods do not specify that

the hybrid edge is a ghost introgression, so it is up to the researcher to recognize its

presence, which can be difficult especially without background knowledge of the group

(Solís-Lemus and Ané 2016; Solís-Lemus et al. 2017).

Overall, our study highlights the necessity of employing multiple paralog

processing pipelines and integrating gene flow analyses to accurately infer species

relationships in groups with complex evolutionary histories. In such systems, pipelines

that rely solely on single-copy orthologs, like ASTRAL, can oversimplify complex

histories and mislead species tree inference. By employing methods that explicitly

model gene duplication and loss, we yielded modest improvements in resolution and

concordance, but alone were not sufficient to fully resolve relationships. By directly

testing for reticulation and ILS, we uncovered the extent to which *Packera's* evolutionary

history has been shaped by hybridization and gene flow. These findings underscore the

broader challenge of capturing evolutionary complexity in phylogenomic datasets,

particularly when orthology inference is itself a major source of uncertainty. Within this

context, the tradeoff between excluding paralogs to reduce noise and retaining them to

preserve evolutionary signal becomes especially pronounced in polyploid systems.

The results of this study exemplifies how critical tree topology is for

macroevolutionary analyses and how paralog processing shapes not only the

phylogenetic relationships but the biological inferences we make from them. These

findings accentuate a central issue in phylogenomic and systematic studies, which is

that macroevolutionary analyses are highly sensitive to orthology inference strategies

and are only as reliable as the phylogenetic framework they are built on, and that

framework is deeply shaped by the methods used to handle paralogy. Looking ahead,

improving phylogeny construction will require both technical innovation and deeper

biological insight. Targeted probe sets designed to enrich truly single-copy genes, such

as Compositae-ParaLoss-1272 (Moore-Pollard et al. 2024), offers a promising strategy

for reducing paralog-related conflict. Yet, excluding paralogs entirely may discard

informative loci, while retaining them without appropriate modeling can mislead species

tree inference. As -omics technologies and phylogenomic methods become increasingly

sophisticated, the accurate interpretation of downstream analyses, such as divergence

dating, biogeography, trait evolution, and gene flow, still hinges on robust taxonomic

and systematic knowledge. In groups like *Packera*, where orthology inference can yield

conflicting results, understanding organismal biology is essential for evaluating the

plausibility of evolutionary scenarios. Morphology, ecology, and geography provide the

critical context needed to assess whether macroevolutionary patterns reflect true

evolutionary history or methodological artifacts.

Rather than treating paralogy as a problem to be avoided, our results highlight its

potential to illuminate processes like gene duplication, polyploidy, and reticulation. While

these challenges extend across many plant and non-plant groups, our densely sampled

Packera dataset offers a uniquely powerful tool for testing and improving phylogenomic

methods. As the field pushes toward more nuanced models of genome evolution,

systems like *Packera* will be essential for developing tools that can fully capture the

intricacy of complex evolutionary histories.

Acknowledgements

The authors thank Drs. Carol Siniscalchi and Matthew Pollard for their

bioinformatic help, Drs. Randall Bayer, Duane McKenna, and Linda Watson for reading

and editing prior to submission, and Dr. John Bain, Dr. Ed Schilling, and Brandon Fuller

for donating genetic material for this study. We would also like to thank the following

herbaria for allowing us to visit and/or destructively sample leaf tissue: ALA, ARIZ, ASU,

CAS, MEM, NY, OSC, TEX, UNC, UCR, WTU. Permits to collect in the San Bernardino

National Forest and Shasta-Trinity National Forest in California were obtained prior to

collection of specimens through the USDA Forest Service. We also thank the University

of Memphis High-Performance Cluster (HPC) administrators, Eric Spangler and Kristian

Skjervold, for their assistance with the HPC and willingness to help. This work was

supported by the Center for Biodiversity (CBio) Research at the University of Memphis.

Data availability

821   Raw sequence data are available in the National Center for Biotechnology

822   Information (NCBI) Sequence Read Archive (SRA) under Bioprojects PRJNA907383,

823   PRJNA978568, PRJNA540287, and PRJNA516161.

Author contributions

ERMP designed the study, generated and analyzed data, and wrote the manuscript.

JRM provided guidance on the project. PE analyzed some of the data. JRM and PE

provided edits to the manuscript. All authors approved of the final version.

Literature Cited

Adeolu M., Alnajar S., Naushad S., S Gupta R. 2016. Genome-based phylogeny and

taxonomy of the "Enterobacteriales": proposal for Enterobacterales ord. nov.

divided into the families Enterobacteriaceae, Erwiniaceae fam. nov.,

Pectobacteriaceae fam. nov., Yersiniaceae fam. nov., Hafniaceae fam. nov.,

Morganellaceae fam. nov., and Budviciaceae fam. nov. Int. J. Syst. Evol.

Microbiol. 66:5575–5599.

Akama S., Shimizu-Inatsugi R., Shimizu K.K., Sese J. 2014. Genome-wide

quantification of homeolog expression ratio revealed nonstochastic gene

regulation in synthetic allopolyploid Arabidopsis. Nucleic Acids Res. 42:e46.

Bain J.F., Golden J.L. 2000. A phylogeny of Packera (Senecioneae; Asteraceae) based

on internal transcribed spacer region sequence data and a broad sampling of

outgroups. Mol. Phylogenet. Evol. 16:331–338.

Bankevich A., Nurk S., Antipov D., Gurevich A.A., Dvorkin M., Kulikov A.S., Lesin V.M.,

Nikolenko S.I., Pham S., Prjibelski A.D., Pyshkin A.V., Sirotkin A.V., Vyahhi N.,

Tesler G., Alekseyev M.A., Pevzner P.A. 2012. SPAdes: A new genome

assembly algorithm and its applications to single-cell sequencing. J. Comput.

Biol. 19:455–477.

Barba-Montoya J. 2021. GBE Assessing Rapid Relaxed-Clock Methods for

Phylogenomic Dating. 13:1–14.

Barker M.S., Li Z., Kidder T.I., Reardon C.R., Lai Z., Oliveira L.O., Scascitelli M.,

Rieseberg L.H. 2016. Most compositae (Asteraceae) are descendants of a

paleohexaploid and all share a paleotetraploid ancestor with the calyceraceae.

Am. J. Bot. 103:1203–1211.

Barkley T.M. 1962. A Revision of Senecio aureus. Trans. Kans. Acad. Sci. 65:318–364.

Barkley T.M. 1988. Variation among the Aureoid Senecios of North America: A

geohistorical interpretation. Bot. Rev. 54:82–106.

Benton M.J., Donoghue P.C.J. 2007. Paleontological evidence to date the tree of life.

Mol. Biol. Evol. 24:26–53.

Bentz P.C., Leebens-Mack J. 2024. Developing Asparagaceae1726: An Asparagaceae-

specific probe set targeting 1726 loci for Hyb-Seq and phylogenomics in the

family. Appl. Plant Sci. 12:e11597.

Betrán E. 2015. The "life histories" of genes. J. Mol. Evol. 80:186–188.

Bezanson J., Chen J., Chung B., Karpinski S., Shah V.B., Vitek J., Zoubritzky L. 2018.

Julia: dynamism and performance reconciled by design. Proc. ACM Program.

Lang. 2:1–23.

Bolger A.M., Lohse M., Usadel B. 2014. Trimmomatic: A flexible trimmer for Illumina

sequence data. Bioinformatics. 30:2114–2120.

Braasch I., Postlethwait J.H. 2012. Polyploidy in Fish and the Teleost Genome

Duplication. Polyploidy and Genome Evolution. Springer Berlin Heidelberg. p.

341–383.

Brown J.W., Walker J.F., Smith S.A. 2017. Phyx: Phylogenetic tools for unix.

Bioinformatics. 33:1886–1888.

Costa F.P., Schrago C.G., Mello B. 2022. Accessing the relative performance of fast

molecular dating methods for phylogenomic data. :1–14.

Crameri S., Fior S., Zoller S., Widmer A. 2022. A target capture approach for

phylogenomic analyses at multiple evolutionary timescales in rosewoods

(Dalbergia spp.) and the legume family (Fabaceae). Mol. Ecol. Resour. 22:3087–

3105.

Danecek P., Bonfield J.K., Liddle J., Marshall J., Ohan V., Pollard M.O., Whitwham A.,

Keane T., McCarthy S.A., Davies R.M., Li H. 2021. Twelve years of SAMtools

and BCFtools. Gigascience. 10.

Degnan J.H., Rosenberg N.A. 2009. Gene tree discordance, phylogenetic inference and

the multispecies coalescent. Trends Ecol. Evol. 24:332–340.

Di Tommaso P., Chatzou M., Floden E.W., Barja P.P., Palumbo E., Notredame C. 2017.

Nextflow enables reproducible computational workflows. Nat. Biotechnol.

35:316–319.

Doyle J.J. 1992. Gene trees and species trees: Molecular systematics as one-character

taxonomy. Syst. Bot. 17:144.

Elven R., Murray D.F., Volodya Yu. Razzhivin B.A.Y. 2011. Annotated Checklist of the

Panarctic Flora (PAF) Vascular plants version 1.0. Available from

http://panarcticflora.org/results?biogeographic=&bioclimatic=®ion=&name=pa

ckera+heterophylla#paf-863101.

Evans B.J., Kelley D.B., Tinsley R.C., Melnick D.J., Cannatella D.C. 2004. A

mitochondrial DNA phylogeny of African clawed frogs: Phylogeography and

implications for polyploid evolution. Mol. Phylogenet. Evol. 33:197–213.

Faircloth B.C. 2016. PHYLUCE is a software package for the analysis of conserved

genomic loci. Bioinformatics. 32:786–788.

Freeman C.C., Barkley T.M. 1995. A synopsis of the genus Packera (Asteraceae:

Senecioneae) in Mexico. SIDA Contrib. Bot. 16:699–709.

Fu Z.X., Jiao B.H., Nie B., Zhang G.J., Gao T.G., Chen Z.D., Lu A.M., Kong H.Z., Wang

X.Q., Wang Y.Z., Zhou S.L., Zhang S.Z., Wang X.M., Liu Z.J., Wang Q.F., Li

J.H., Li D.Z., Yi T.S., Hong M.A., Soltis D.E., Soltis P.S., Li J.H., Fu C.X., Liu

Q.X. 2016. A comprehensive generic-level phylogeny of the sunflower family:

Implications for the systematics of Chinese Asteraceae. J. Syst. Evol. 54:416–

437.

Funk V.A. 1985. Phylogenetic Patterns and Hybridization. Ann. Mo. Bot. Gard. 72:681.

Gallardo M.H., Kausel G., Jiménez A., Bacquet C., González C. 2004. Biological

relevance of polyploidy : ecology to genomics Whole-genome duplications in

South American desert rodents (Octodontidae). Society.:443–451.

Galtier N., Daubin V. 2008. Dealing with incongruence in phylogenomic analyses.

Philos. Trans. R. Soc. Lond. B Biol. Sci. 363:4023–4029.

Geldmacher J., Van Den Bogaard P., Hoernle K., Schmincke H.U. 2000. The 40Ar/39Ar

age dating of the Madeira Archipelago and hotspot track (eastern North Atlantic).

Geochem. Geophys. Geosyst. 1.

Gramling A. 2006. A conservation assessment of Packera millefolium, a Southern

Appalachian Endemic. .

Grant V. 1971. Plant Speciation. New York: Columbia University Press.

Hejase H.A., Liu K.J. 2016. A scalability study of phylogenetic network inference

methods using empirical datasets and simulations involving a single reticulation.

BMC Bioinformatics. 17:422.

Hobbs C.R., Baldwin B.G. 2013. Asian origin and upslope migration of Hawaiian

Artemisia (Compositae-Anthemideae). J. Biogeogr. 40:442–454.

Huang C.H., Zhang C., Liu M., Hu Y., Gao T., Qi J., Ma H. 2016. Multiple

Polyploidization Events across Asteraceae with Two Nested Events in the Early

History Revealed by Nuclear Phylogenomics. Mol. Biol. Evol. 33:2820–2835.

Huson D.H., Scornavacca C. 2012. Dendroscope 3: an interactive tool for rooted

phylogenetic trees and networks. Syst. Biol. 61:1061–1067.

Jackson C., McLay T., Schmidt-Lebuhn A.N. 2023. hybpiper-nf and paragone-nf:

Containerization and additional options for target capture assembly and paralog

resolution. APPS.

Jin J.-J., Yu W.-B., Yang J.-B., Song Y., dePamphilis C.W., Yi T.-S., Li D.-Z. 2020.

GetOrganelle: a fast and versatile toolkit for accurate de novo assembly of

organelle genomes. Genome Biol. 21:241.

Johnson M.G., Gardner E.M., Liu Y., Medina R., Goffinet B., Shaw A.J., Zerega N.J.C.,

Wickett N.J. 2016. HybPiper: Extracting coding sequence and introns for

phylogenetics from high-throughput sequencing reads using target enrichment.

Appl. Plant Sci. 4:1600016.

Jones K.E., Alfredo Reyes-Betancort J., Hiscock S.J., Carine M.A. 2014. Allopatric

diversification, multiple habitat shifts, and hybridization in the evolution of

Pericallis (Asteraceae), a Macaronesian endemic genus. Am. J. Bot. 101:637–

651.

Katoh K., Standley D.M. 2013. MAFFT multiple sequence alignment software version 7:

Improvements in performance and usability. Mol. Biol. Evol. 30:772–780.

Kück P., Longo G.C. 2014. FASconCAT-G: Extensive functions for multiple sequence

alignment preparations concerning phylogenetic studies. Front. Zool. 11:1–8.

Kumar S., Stecher G., Li M., Knyaz C., Tamura K. 2018. MEGA X: Molecular

evolutionary genetics analysis across computing platforms. Mol. Biol. Evol.

35:1547–1549.

Kumar S., Stecher G., Peterson D., Tamura K. 2012. MEGA-CC: Computing core of

950   molecular evolutionary genetics analysis program for automated and iterative

951   data analysis. Bioinformatics. 28:2685–2686.

952 Kumar S., Stecher G., Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics

953   Analysis Version 7.0 for Bigger Datasets. Mol. Biol. Evol. 33:1870–1874.

954 Kurtzer G.M., cclerget, Bauer M., Kaneshiro I., Trudgian D., Godlove D. 2021.

955   hpcng/singularity: Singularity 3.7.3. Zenodo.

956 Kurtzer G.M., Sochat V., Bauer M.W. 2017. Singularity: Scientific containers for mobility

957   of compute. PLoS One. 12:e0177459.

958 Landis M.J., Matzke N.J., Moore B.R., Huelsenbeck J.P. 2013. Bayesian analysis of

959   biogeography when the number of areas is large. Syst. Biol. 62:789–804.

960 Lanfear R., Calcott B., Ho S.Y.W., Guindon S. 2012. PartitionFinder: Combined

961   selection of partitioning schemes and substitution models for phylogenetic

962   analyses. Mol. Biol. Evol. 29:1695–1701.

963 Langmead B., Salzberg S.L. 2012. Fast gapped-read alignment with Bowtie 2. Nat.

964   Methods. 9:357–359.

965 Li H., Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler

966   transform. Bioinformatics. 25:1754–1760.

967 Linder C.R., Rieseberg L.H. 2004. Reconstructing patterns of reticulate evolution in

968   plants. Am. J. Bot. 91:1700–1708.

969 Maddison W.P. 1997. Gene trees in species trees. Systematic Bio. 46:523–536.

970 Mahoney A.M., Kowal R.R. 2008. Three new varieties of Packera paupercula

971   (Asteraceae, Senecioneae) in midwestern and southeastern North America.

972   Novon St. Louis Mo. 18:220–228.

973 Mandel J.R., Dikow R.B., Funk V.A., Masalia R.R., Staton S.E., Kozik A., Michelmore

R.W., Rieseberg L.H., Burke J.M. 2014. A Target Enrichment Method for

Gathering Phylogenetic Information from Hundreds of Loci: An Example from the

Compositae. Appl. Plant Sci. 2:1300085.

Mandel J.R., Dikow R.B., Siniscalchi C.M., Thapa R., Watson L.E., Funk V.A. 2019. A

fully resolved backbone phylogeny reveals numerous dispersals and explosive

diversifications throughout the history of Asteraceae. Proc. Natl. Acad. Sci. U. S.

A. 116:14083–14088.

Masterson J. 1994. Stomatal Size. Science. 264:421–424.

Matzke N.J. 2014. Model selection in historical biogeography reveals that founder-event

speciation is a crucial process in island clades. Syst. Biol. 63:951–970.

McDougall I., Schmincke H.U. 1976. Geochronology of Gran Canaria, Canary Islands:

Age of shield building volcanism and other magmatic phases. Bulletin

Volcanologique. 40:57–77.

Mello B. 2018. Estimating timetrees with MEGA and the timetree resource. Mol. Biol.

Evol. 35:2334–2342.

Mendes F.K., Hahn Y., Hahn M.W. 2016. Gene tree discordance can generate patterns

of diminishing convergence over time. Mol. Biol. Evol. 33:3299–3307.

Minh B.Q., Schmidt H.A., Chernomor O., Schrempf D., Woodhams M.D., Von Haeseler

A., Lanfear R., Teeling E. 2020. IQ-TREE 2: New Models and Efficient Methods

for Phylogenetic Inference in the Genomic Era. Mol. Biol. Evol. 37:1530–1534.

Mitchell N., Lewis P.O., Lemmon E.M., Lemmon A.R., Holsinger K.E. 2017. Anchored

phylogenomics improves the resolution of evolutionary relationships in the rapid

radiation of Protea L. Am. J. Bot. 104:102–115.

Moore-Pollard E.R., Jones D.S., Mandel J.R. 2024. Compositae-ParaLoss-1272: A

998  complementary sunflower-specific probe set reduces paralogs in phylogenomic

999  analyses of complex systems. Appl. Plant Sci. 12:e11568.

1000 Morales-Briones D.F., Gehrke B., Huang C.-H., Liston A., Ma H., Marx H.E., Tank D.C.,

1001  Yang Y. 2021a. Analysis of paralogs in target enrichment data pinpoints multiple

1002  ancient polyploidy events in Alchemilla s.L. (Rosaceae). Syst. Biol. 71:190–207.

1003 Morales-Briones D.F., Kadereit G., Tefarikis D.T., Moore M.J., Smith S.A., Brockington

1004  S.F., Timoneda A., Yim W.C., Cushman J.C., Yang Y. 2021b. Disentangling

1005  sources of gene tree discordance in phylogenomic data sets: Testing ancient

1006  hybridizations in Amaranthaceae s.L. Syst. Biol. 70:219–235.

1007 Morales-Briones D.F., Liston A., Tank D.C. 2018. Phylogenomic analyses reveal a deep

1008  history of hybridization and polyploidy in the Neotropical genus Lachemilla

1009  (Rosaceae). New Phytol. 218:1668–1684.

1010 Müller J., Reisz R.R. 2005. Four well-constrained calibration points from the vertebrate

1011  fossil record for molecular clock estimates. Bioessays. 27:1069–1075.

1012 Nice C.C., Gompert Z., Fordyce J.A., Forister M.L., Lucas L.K., Buerkle C.A. 2013.

1013  Hybrid speciation and independent evolution in lineages of alpine butterflies.

1014  Evolution. 67:1055–1068.

1015 One Thousand Plant Transcriptomes Initiative. 2019. One thousand plant

1016  transcriptomes and the phylogenomics of green plants. Nature. 574:679–685.

1017 Ortiz E.M., Höwener A., Shigita G., Raza M., Maurin O., Zuntini A., Forest F., Baker

1018  W.J., Schaefer H. 2023. A novel phylogenomics pipeline reveals complex pattern

1019  of reticulate evolution in Cucurbitales. bioRxiv.

1020 Ottenburghs J. 2020. Ghost introgression: Spooky gene flow in the distant past.

1021  Bioessays. 42:e2000012.

Palazzesi L., Pellicer J., Barreda V.D., Loeuille B., Mandel J.R., Pokorny L., Siniscalchi

C.M., Tellería M.C., Leitch I.J., Hidalgo O. 2022. Asteraceae as a model system

for evolutionary studies: from fossils to genomes. Bot. J. Linn. Soc. 200:143–164.

Panchy N., Lehti-Shiu M., Shiu S.H. 2016. Evolution of gene duplication in plants. Plant

Physiol. 171:2294–2316.

Panero J.L. 2005. New combinations and infrafamilial taxa in the Asteraceae.

Phytologia. 87:1–14.

Paradis E., Schliep K. 2019. Ape 5.0: An environment for modern phylogenetics and

evolutionary analyses in R. Bioinformatics. 35:526–528.

Parham J.F., Donoghue P.C.J., Bell C.J., Calway T.D., Head J.J., Holroyd P.A., Inoue

J.G., Irmis R.B., Joyce W.G., Ksepka D.T., Patané J.S.L., Smith N.D., Tarver

J.E., Van Tuinen M., Yang Z., Angielczyk K.D., Greenwood J.M., Hipsley C.A.,

Jacobs L., Makovicky P.J., Müller J., Smith K.T., Theodor J.M., Warnock R.C.M.,

Benton M.J. 2012. Best practices for justifying fossil calibrations. Syst. Biol.

61:346–359.

Pease J.B., Brown J.W., Walker J.F., Hinchliff C.E., Smith S.A. 2018. Quartet Sampling

distinguishes lack of support from conflicting support in the green plant tree of

life. Am. J. Bot. 105:385–403.

Peirson J.A., Reznicek A.A., Semple J.C. 2012. Polyploidy, infraspecific cytotype

variation, and speciation in Goldenrods: The cytogeography of Solidago subsect.

Humiles (Asteraceae) in North America. Taxon. 61:197–210.

Pelser P.B., Kennedy A.H., Tepe E.J., Shidler J.B., Nordenstam B., Kadereit J.W.,

Watson L.E. 2010. Patterns and causes of incongruence between plastid and

nuclear Senecioneae (Asteraceae) phylogenies. Am. J. Bot. 97:856–873.

R Core Team. 2016. R: A Language and Environment for Statistical Computing. Vienna:

R Foundation for Statistical Computing.

Ree R.H., Smith S.A. 2008. Maximum likelihood inference of geographic range

evolution by dispersal, local extinction, and cladogenesis. Syst. Biol. 57:4–14.

Revell L.J. 2012. phytools: An R package for phylogenetic comparative biology (and

other things). Methods Ecol. Evol. 3:217–223.

Ronquist F. 1997. Dispersal-vicariance analysis: A new approach to the quantification of

historical biogeography. Syst. Biol. 46:195–203.

Rose J.P., Toledo C.A.P., Moriarty Lemmon E., Lemmon A.R., Sytsma K.J. 2021. Out

of Sight, Out of Mind: Widespread Nuclear and Plastid-Nuclear Discordance in

the Flowering Plant Genus Polemonium (Polemoniaceae) Suggests Widespread

Historical Gene Flow Despite Limited Nuclear Signal. Syst. Biology. 70:162–180.

RStudio. 2020. RStudio: Integrated Development for R. Boston, MA: RStudio, PBC.

Salmon A., Udall J.A., Jeddeloh J.A., Wendel J. 2012. Targeted capture of

homoeologous coding and noncoding sequence in polyploid cotton. G3

(Bethesda). 2:921–930.

Sayyari E., Mirarab S. 2016. Fast Coalescent-Based Computation of Local Branch

Support from Quartet Frequencies. Mol. Biol. Evol. 33:1654–1668.

Schliep K.P. 2011. phangorn: phylogenetic analysis in R. Bioinformatics. 27:592–593.

Shang H.-Y., Jia K.-H., Li N.-W., Zhou M.-J., Yang H., Tian X.-L., Ma Y.-P., Zhang R.-G.

2025. Phytop: a tool for visualizing and recognizing signals of incomplete lineage

sorting and hybridization using species trees output from ASTRAL. Hortic. Res.

12:uhae330.

Siu-Ting K., Torres-Sánchez M., San Mauro D., Wilcockson D., Wilkinson M., Pisani D.,

O'Connell M.J., Creevey C.J. 2019. Inadvertent paralog inclusion drives

artifactual topologies and timetree estimates in phylogenomics. Mol. Biol. Evol.

36:1344–1356.

Smith S.A., Moore M.J., Brown J.W., Yang Y. 2015. Analysis of phylogenomic datasets

reveals conflict, concordance, and gene duplications with examples from animals

and plants. BMC Evol. Biol. 15:150.

Smith S.A., O'Meara B.C. 2012. TreePL: Divergence time estimation using penalized

likelihood for large phylogenies. Bioinformatics. 28:2689–2690.

Solís-Lemus C., Ané C. 2016. Inferring Phylogenetic Networks with Maximum

Pseudolikelihood under Incomplete Lineage Sorting. PLoS Genet. 12:1–21.

Solís-Lemus C., Bastide P., Ané C. 2017. PhyloNetworks: A package for phylogenetic

networks. Mol. Biol. Evol. 34:3292–3298.

Soltis D.E., Soltis P.S. 1999. Polyploidy: Recurrent formation and genome evolution.

Trends Ecol. Evol. 14:348–352.

Soltis P.S., Soltis D.E. 2009. The role of hybridization in plant speciation. Annu. Rev.

Plant Biol. 60:561–588.

Stamatakis A. 2014. RAxML version 8: A tool for phylogenetic analysis and post-

analysis of large phylogenies. Bioinformatics. 30:1312–1313.

Steel M.A., Penny D. 1993. Distributions of tree comparison metrics--some new results.

Syst. Biol. 42:126–141.

Steenwyk J.L., Buida T.J., Labella A.L., Li Y., Shen X.-X., Rokas A. 2021. PhyKIT: a

broadly applicable UNIX shell toolkit for processing and analyzing phylogenomic

data. Bioinformatics. 37:2325–2331.

Struck T.H. 2013. The impact of paralogy on phylogenomic studies - a case study on

annelid relationships. PLoS One. 8:e62892.

Stull G.W., Soltis P.S., Soltis D.E., Gitzendanner M.A., Smith S.A. 2020. Nuclear

phylogenomic analyses of asterids conflict with plastome trees and support novel

relationships among major lineages. Am. J. Bot. 107:790–805.

Takhtajan A. 1986. Floristic Regions of the World. Berkley and Los Angeles, California:

University of California Press.

Tamura K., Battistuzzi F.U., Billing-Ross P., Murillo O., Filipski A., Kumar S. 2012.

Estimating divergence times in large molecular phylogenies. Proc. Natl. Acad.

Sci. U. S. A. 109:19333–19338.

Tamura K., Tao Q., Kumar S. 2018. Theoretical foundation of the RelTime method for

estimating divergence times from variable evolutionary rates. Mol. Biol. Evol.

35:1770–1782.

Tao Q., Tamura K., Kumar S. 2020a. The Molecular Evolutionary Clock. The Molecular

Evolutionary Clock.

Tao Q., Tamura K., Mello B., Kumar S. 2020b. Reliable Confidence Intervals for

RelTime Estimates of Evolutionary Divergence Times. Mol. Biol. Evol. 37:280–

290.

Thapa R., Bayer R.J., Mandel J.R. 2020. Phylogenomics Resolves the Relationships

within Antennaria (Asteraceae, Gnaphalieae) and Yields New Insights into its

Morphological Character Evolution and Biogeography. Syst. Bot. 45:387–402.

Tillich M., Lehwark P., Pellizzer T., Ulbricht-Jones E.S., Fischer A., Bock R., Greiner S.

2017. GeSeq - Versatile and accurate annotation of organelle genomes. Nucleic

Acids Res. 45:W6–W11.

Tricou T., Tannier E., de Vienne D.M. 2022. Ghost lineages highly influence the

interpretation of introgression tests. Syst. Biol. 71:1147–1158.

Trock D.K. 1999. A revisionary synthesis of the genus Packera (Asteraceae:

Senecioneae). .

Trock D.K. 2006. Packera Á. Löve & D. Löve. Flora of North America Editorial

Committee, eds. 1993+. New York and Oxford: p. 570–602.

Vision T.J., Brown D.G., Tanksley S.D. 2000. The origins of genomic duplications in

Arabidopsis. Science. 290:2114–2117.

Wang W.M. 2004. On the origin and development of Artemisia (Asteraceae) in the

geological past. Bot. J. Linn. Soc. 145:331–336.

Wendel J.F., Cronn R.C. 2003. Polyploidy and the Evolutionary History of Cotton.

Botany Publication and Papers. 23:139–186.

Yang Y., Smith S.A. 2014. Orthology inference in nonmodel organisms using

transcriptomes and low-coverage genomes: Improving accuracy and matrix

occupancy for phylogenomics. Mol. Biol. Evol. 31:3081–3092.

Yeatts L., Schneider A., Schneider B. 2011. Packera mancosana (Asteraceae:

Senecioneae), a new species and shale barren endemic of southwestern

colorado. Phytoneuron.:1–8.

Yu G., Smith D.K., Zhu H., Guan Y., Lam T.T.-Y. 2017. Ggtree: An r package for

visualization and annotation of phylogenetic trees with their covariates and other

associated data. Methods Ecol. Evol. 8:28–36.

Zhang C., Huang C.-H., Liu M., Hu Y., Panero J.L., Luebert F., Gao T., Ma H. 2021.

Phylotranscriptomic insight into Asteraceae diversity, polyploidy, and

morphological innovation. Journal of Integrative Plant Biolog. 63:1273–1293.

Zhang C., Mirarab S. 2022a. ASTRAL-Pro 2: ultrafast species tree reconstruction from

multi-copy gene family trees. Bioinformatics. 38:4949–4950.

Zhang C., Mirarab S. 2022b. Weighting by gene tree uncertainty improves accuracy of

quartet-based species trees. Mol. Biol. Evol. 39.

Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018. ASTRAL-III: Polynomial time species

tree reconstruction from partially resolved gene trees. BMC Bioinformatics.

19:15–30.

Zhang C., Scornavacca C., Molloy E.K., Mirarab S. 2020. ASTRAL-pro: Quartet-based

species-tree inference despite paralogy. Mol. Biol. Evol. 37:3292–3307.

1150

ASTRAL

Quartet Concordance (QC)
- 🔵 QC > 0.2
- 🔵 0.2 ≥ QC > 0
- 🟠 0 ≥ QC ≥ -0.05
- 🔴 QC < -0.05

Tribes/Subtribes
- 🟨 Senecioneae/Senecioninae
- 🟩 Senecioneae/Tussilagininae s.str.
- 🟪 Tephroseridineae
- 🟪 Anthemideae
- 🟧 Doroniceae
- 🟩 Calenduleae

Reticulation clades
- ▢ Mexico
- ▢ Rocky
- ▢ California
- ▢ Eastern
- ▢ Arctic
- ⭐ Backbone

1151

Figure 1. Full phylogeny of the ASTRAL analysis. Tribal/subtribal status is indicated on

the right of the phylogeny and follows the color code as indicated in the key. Numbers

next to tip labels correspond to the Quartet Fidelity (QF) score from Quartet Sampling,

which ranges from 0 to 1, with values closer to 0 indicating erroneous taxa. Circles at

nodes show the Quartet Concordance (QC) factors from Quartet Sampling, where red

and orange indicate counter-support nodes. Asterisks next to QC circles indicate high

support (≥95 LPP) at that node. Highlighted clades, and teal stars next to some tip

labels, indicate clades where reticulation events were tested, following the color key on

the left.

1161

1-to-1

Figure 2. Full phylogeny of the paragone-nf: 1-to-1 analysis. Tribal/subtribal status is

indicated on the right of the phylogeny and follows the color code as indicated in the

key. Numbers next to tip labels correspond to the Quartet Fidelity (QF) score from

Quartet Sampling, which ranges from 0 to 1, with values closer to 0 indicating erroneous

taxa. Circles at nodes show the Quartet Concordance (QC) factors from Quartet

Sampling, where red and orange indicate counter-support nodes. Asterisks next to QC

circles indicate high support (≥95 LPP) at that node. Highlighted clades, and teal stars

next to some tip labels, indicate clades where reticulation events were tested, following

the color key on the left.

1172

MO

Quartet Concordance (QC)
- QC > 0.2
- 0.2 ≥ QC > 0
- 0 ≥ QC ≥ -0.05
- QC < -0.05

Tribes/Subtribes
- Senecioneae/Senecionineae
- Senecioneae/Tussilagininae s.str.
- Tephroseridineae
- Anthemideae
- Doroniceae
- Calenduleae

Reticulation clades
- Mexico
- Rocky
- California
- Eastern
- Arctic
- Backbone

Packera neomexicana var. *toumeyi* (0.432)
Packera neomexicana var. *mutabilis* (0.517)
Packera quercetorum (0.509)
Packera neomexicana var. *neomexicana* (0.481)
Packera thurberi (0.446)
Packera texensis (0.454)
Packera multilobata (0.47)
Packera hintoniorum (0.585)
Packera scalaris var. *parrasiana* (0.628)
Packera eurycephala var. *eurycephala* (0.578)
Packera layneae (0.534)
Packera eurycephala var. *lewisrosei* (0.608)
Packera ionophylla (0.644)
Packera bernardina (0.56)
Packera clevelandii (0.58)
Packera hesperia (0.607)
Packera macounii (0.557)
Packera fendleri (0.423)
Packera cynthioides (0.396)
Packera streptanthifolia var. *rubicaulis* (0.501)
Packera malmstenii (0.594)
Packera werneriifolia (0.542)
Packera franciscana (0.489)
Packera mancosana (0.529)
Packera cana (0.445)
Packera candidissima (0.599)
Packera bellidifolia (0.712)
Packera streptanthifolia var. *borealis* (0.615)
Packera neomexicana var. *metcalfei* (0.656)
Packera millelobata (0.5)
Packera dimorphophylla var. *dimorphophylla* (0.695)
Packera crocata (0.699)
Packera sanguisorboides (0.668)
Packera quebradensis (0.687)
Packera cardamine (0.656)
Packera musiniensis (0.708)
Packera castoreus (0.536)
Packera paupercula var. *paupercula* (0.426)
Packera anonyma (0.559)
Packera schweinitziana (0.605)
Packera xmemmingeri (0.624)
Packera millefolium (0.613)
Packera paupercula var. *appalachiana* (0.501)
Packera plattensis (0.522)
Packera obovata (0.613)
Packera aurea (0.643)
Packera spellenbergii (0.644)
Packera pseudaurea var. *flavula* (0.688)
Packera pseudaurea var. *semicordata* (0.683)
Packera crawfordii (0.549)
Packera streptanthifolia var. *streptanthifolia* (0.671)
Packera dimorphophylla var. *paysonii* (0.652)
Packera paupercula var. *savannarum* (0.434)
Packera dubia (0.479)
Packera pseudaurea var. *pseudaurea* (0.546)
Packera subnuda var. *moresbiensis* (0.527)
Packera indecora (0.658)
Packera pauciflora (0.647)
Packera subnuda var. *subnuda* (0.545)
Packera rosei (0.645)
Packera antennariifolia (0.634)
Packera bolanderi var. *bolanderi* (0.649)
Packera bolanderi var. *harfordii* (0.624)
Packera flettii (0.673)
Packera ganderi (0.677)
Packera brewerii (0.685)
Packera porteri (0.592)
Packera cymbalaria (0.632)
Packera heterophylla (0.566)
Packera debilis (0.559)
Packera contermina (0.614)
Packera greenei (0.571)
Packera hyperborealis (0.542)
Packera ogotorukensis (0.602)
Packera glabella (0.768)
Packera tampicana (0.745)
Packera coahuilensis (0.749)
Packera sanguisorbae (0.718)
Packera hartiana (0.712)
Pericallis echinata (0.791)
Pericallis murrayi (0.788)
Pericallis cruenta (0.782)
Elekmania barahonensis (0.727)
Xenophyllum lycopodioides (0.74)
Werneria aretioides (0.696)
Packera moranii (0.484)
Packera loratifolia (0.607)
Packera zimapanica (0.525)
Packera actinella (0.731)
Packera toluccana (0.672)
Packera umbraculifera (0.736)
Senecio vulgaris (0.827)
Senecio flavus (0.755)
Jacobaea vulgaris (0.56)
Erechtites hieraciifolius (0.615)
Emilia fosbergii (0.874)
Sinacalia tangutica (0.616)
Tussilago farfara (0.643)
Telanthophora andrieuxii (0.728)
Roldana gilgii (0.693)
Cacaliopsis nardosmia (0.777)
Brachyglottis repanda (0.732)
Tephroseris newcombei (0.657)
Achillea nobilis (0.829)
Anthemis arvensis (0.874)
Artemisia vulgaris (0.84)
Cotula turbinata (0.863)
Doronicum pardalianches (0.852)
Tripteris pinnatilobatum (0.833)
Chyrsanthemoides incana (0.85)
Calendula arvensis (0.863)

PACKERA

Figure 3. Full phylogeny of the paragone-nf: MO analysis. Tribal/subtribal status is

indicated on the right of the phylogeny and follows the color code as indicated in the

key. Numbers next to tip labels correspond to the Quartet Fidelity (QF) score from

Quartet Sampling, which ranges from 0 to 1, with values closer to 0 indicating erroneous

taxa. Circles at nodes show the Quartet Concordance (QC) factors from Quartet

Sampling, where red and orange indicate counter-support nodes. Asterisks next to QC

circles indicate high support (≥95 LPP) at that node. Highlighted clades, and teal stars

next to some tip labels, indicate clades where reticulation events were tested, following

the color key on the left.

1183

1184

Figure 4. Genus-level phylogenies of all trees generated in this study. Genera are

colored according to their tribal/subtribal classifications following the key on the top of

the figure. Packera is marked with a black asterisk next to its placement in the trees.

Navy blue text at node represents support values (LPP or BS). Bottom left is a matrix of

1189 RF_{adj} values between the nine different trees, with most similar topologies as dark blue

1190 squares (0) and least similar as yellow squares (1). Bottom right are the RF_{adj} values

1191 plotted as a principal components analysis. Points closer together indicate higher

1192 similarity, while dots further away indicate higher dissimilarity.

1193

Figure 5. Predicted biogeographic region summaries generated from BioGeoBEARS of

the HybPiper, paragone-nf: 1-to-1, and paragone-nf: MO analyses. The maps of North

America are colored according to the ten geographic regions outlined by Takhtajan

(1986): Arctic Province (I), Canadian Province (C), Appalachian Province (P), Atlantic

and Gulf Coast Plain Province (T), North American Prairies Province (N), Rocky

Mountain and Vancouvarian Province (R), Madrean Region (M), Asia (A), Europe (E),

and Central America (S). Stars indicate the predicted ancestral range of crown *Packera*.

Arrows and numbers correspond to their predicted range expansions in chronological

order, 5 being the most recent major expansion. All BioGeoBears results can be found

in Supplemental Figure I. Images of *Packera* species highlighting some of the

morphological diversity seen in the group. Colored bars on the bottom of the image

correspond to their current distributions, following the color scheme of the map and key.

From left to right: Flame ragwort, *P. greenei* (A.Gray) W.A.Weber & Á.Löve; Tehachapi

ragwort, *P. ionophylla* (Greene) W.A.Weber & Á.Löve; Fendler's ragwort, *Packera*

fendleri (A.Gray) W.A.Weber & Á.Löve; Weak groundsel, *P. debilis* (Nutt.) W.A.Weber &

1210   Á.Löve; and Rayless mountain ragwort, *P. indecora* (Greene) Á.Löve & D.Löve. Image

1211   credits: Erika R. Moore-Pollard and Robert Lagier.

1212

1213

Figure 6. PhyloNetworks figures of the three tested analyses: ASTRAL, 1-to-1, and MO.

The various areas of the phylogeny that were tested is shown on the left and can be

read as a row. Dark green edges indicate the major hybrid edge, while the light blue

shows the minor hybrid edge. The associated numbers next to each branch

corresponds to the percentage of parentage given by that branch. Number of

hybridizations correspond with the best log likelihood value. The 1-to-1 analysis does

not have a Rocky analysis since the clade in that tree only had four taxa.

Table S1. Voucher specimens of the 111 taxa used in this study. Publication status and

authorities assigned by IPNI. FNA = Flora of North America (2006).

Table S2. Raw sequence statistics using the pxlssq function in *Phyx* for each of the 111

samples in this study.

Table S3. Matrix and tree statistics of the nine trees generated with 111 taxa.

Table S4. Best fitting models for each gene from PartitionFinder given the overall run

(General) and the ASTRAL-Pro run where the paralogs and single-copy orthologs were

run separately. The three models tested were GTR, GTR+G, and GTR+I+G. Numbers

in parenthesis next to the model type corresponds to the number of genes for that

model.

Table S5. Dating scenarios with various age constraints (A-E) implemented in RelTime.

y = age constraint was used in the scenario; n = age constraint was not used in the

scenario.

Table S6. Presence/Absence values of each species given the ten geographic regions

outlined by Takhtajan (1986). A value of "0" indicates absence, "1" indicates presence.

Letters to the right of each column heading relates to Fig. 5. The four column headings

above the geographic regions refer to the four groupings of *Packera* used in the

discussion, no groupings were designated for Europe or Central America since they

represent outgroup distributions.

Table S7. Log likelihood values and taxon lists on the five PhyloNetworks runs–

Backbone, Arctic/Alpine, California/Mexico, Eastern, and Rocky Mountains–across the

ASTRAL, 1-to-1, and MO trees. The number of reticulations with the lowest log

likelihood value, indicated by blue font, is considered the best fit.

Table S8. Assembly statistics generated from the HybPiper pipeline.

Table S9. Percent missing data of each plastid assembly generated.

Table S10. Pairwise adjusted Robinson-Foulds (RFadj) distances between the nine

main trees compared in this study. The values range from zero to one, indicating

whether the tree topologies are identical or completely dissimilar, respectively.

Table S11. Resulting divergence rates of crown *Packera*, Senecioneae, and

Anthemideae from ASTRAL, MO, 1-to-1 using RelTime.

Table S12. Biogeographical model selection for all *Packera* taxa and four outgroup

genera (*Pericallis, Elekmania, Werneria, Xenophyllum*). Ln L = log likelihood values;

Num params = number of free parameters; d = "dispersal" parameter; e = "extinction"

parameter; j = "jump" parameter which specifies the weight of each jump dispersal event

in the cladogenesis matrix; AIC = Akaike information criterion; AICc = Akaike

information criterion with corrections for small sample sizes. "wt" represents the

resulting model weights with AIC and AICc. Blue font indicates the chosen BioGeoBears

analysis.

Table S13. List of species in each of the five tested networks for the ASTRAL, 1-to-1,

and MO trees. Colored cells indicate that the species was present in the network, no

color indicates absence. Colors are according to the network shading from Figs. 1, 2,

and 3.

Figure S1. Annotated plastids of 18 taxa with complete plastomes.

Figure S2. Full chloroplast tree of 111 mapped plastid data. Colors at the end of tips

corresponds to the percentage of missing data for each taxon.

Figure S3. All tested trees containing 111 taxa (excluding the complete plastid tree).

Scale bar under the tree on the left represents the mean number of nucleotide

substitutions per site. Support values–bootstraps (BS, ranging from 0-100) or local

posterior probabilities (LPP, ranging from 0-1)–are indicated next to the node. Species

tree inference methods are detailed in Table 1. Phylogenies generated using phytools in

R.

Figure S4. Results of Quartet Sampling on the different, tested trees. Values at each

node correspond to the Quartet Concordance (QC), Quartet Differential (QD), and

Quartet Informativeness (QI) values (QC/QD/QI). Colored circles at tips correspond to

the Quartet Fidelity (QF) score.

Figure S5. PhyParts results of the different phylogenies. Numbers above the branch

indicate the number of concordant gene trees, while the numbers below the branch

indicate the number of conflicting gene trees. Blue: concordant gene trees, green: top

alternative bipartition, red: all other alternative bipartitions; gray: uninformative or

missing data.

Figure S6. PhyTop results.

Figure S7. Divergence rate estimates of Scenarios 1-6 using RelTime for the ASTRAL,

1-to-1, and MO trees in millions of years. Left tree contains the mean age of that

designated node, the right tree shows the 95% age range for the designated node.

Figure S8. Predicted biogeographic regions generated from BioGeoBEARS on all

Packera taxa and four outgroup genera (*Pericallis*, *Elekmania*, *Werneria*, and

Xenophyllum) using all tested models for the ASTRAL, 1-to-1, and MO trees. Boxes and

circles at nodes represent the geographic regions provided in Figure 5.

Figure S9. Plots of loglikelihood values from Table S7 of each hybridization run (along

the left y-axis) for each tree tested (along the top x-axis). The bottom x-axis indicates

the 11 tested hybridization numbers (n=0-10). Colored points indicate the best fitting

hybridization for that analysis.