Phylogenomic challenges in polyploid-rich lineages: Insights from paralog

processing and reticulation methods using the complex genus *Packera*

(Asteraceae: Senecioneae)

Erika R. Moore-Pollard[1,2], Paige A. Ellestad[2], Jennifer R. Mandel[2,3]

[1]*Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs,*

Connecticut 06269, USA

[2]*Department of Biological Sciences, University of Memphis, Memphis, Tennessee*

38152, USA

[3]*ACRE: Institute for Agricultural and Conservation Research and Education, University*

of Memphis, Memphis, Tennessee 38152, USA

ABSTRACT

Phylogenomic discordance is pervasive and cannot always be resolved by increasing

the amount of sequencing data alone. Biological processes such as polyploidy,

hybridization, and incomplete lineage sorting are major contributors to discordance and

must be accounted for to avoid misleading evolutionary interpretations. To better

understand how these processes influence phylogenetic reconstruction, we conducted a

comprehensive phylogenomic study in the complex genus *Packera*. With over 90

species and varieties, 40% of which exhibit polyploidy, aneuploidy, or other cytological

complexities, *Packera* presents significant challenges for phylogenetic reconstruction.

Given these complexities, we assessed different published paralog processing methods

on the resulting evolutionary relationships and phylogenetic support of this group. We

then applied three of these methods to evaluate their impact on tree topology and our

understanding of *Packera*'s evolutionary history by constructing a time-calibrated

phylogeny, reconstructing historical biogeography, and testing for ancient reticulation.

Phylogenetic outcomes varied based on the paralog processing method used, with no

method performing the best over others. Our findings highlight the large impact of

orthology inference and paralog processing on phylogenomic analyses, particularly in

polyploid-rich groups such as *Packera*, and we offer guidance on methodological

impacts along with practical recommendations. We note that gaining a robust

understanding of *Packera*'s evolutionary history requires more than computational

approaches alone. While technological advancements have greatly expanded our ability

to analyze genomic data, effective phylogenomic research still relies on strong taxon

sampling and detailed species knowledge. Without careful attention to the biological

context, such as reproductive boundaries, cytological variation, ecological interactions,

and historical biogeographic processes, phylogenomic studies risk misinterpreting

evolutionary history and processes. By accounting for these factors, we can begin to

improve the accuracy of evolutionary reconstructions and gain deeper insights into the

complex history of plant diversification.

KEYWORDS

Compositae, discordance, hybridization, molecular systematics, orthology inference

INTRODUCTION

The field of phylogenetics is drastically changing and now routinely incorporates

multi-locus and genomic data. While this expansion has enabled deeper insights into

evolutionary relationships across the tree of life, it has also revealed persistent and

complex challenges in phylogenetic inference. One significant challenge is gene tree

discordance, where phylogenetic trees estimated from individual loci disagree with each

other and/or with the inferred species tree (Degnan and Rosenberg 2009; Mendes et al.

2016). These incongruences are not simply a consequence of limited data; rather, they

often reflect differences driven by underlying biological processes such as incomplete

lineage sorting (ILS), gene duplication and loss, and hybridization or introgression

(Pease et al. 2018). Such processes are especially prevalent in plants, where whole-

genome duplications (WGDs), ancient hybridization, and reticulate evolution are

widespread and recurrent (Funk 1985; Vision et al. 2000; Wendel and Cronn 2003;

Peirson et al. 2012; Panchy et al. 2016; Rose et al. 2021).

Polyploidy, or WGD, is a dominant evolutionary force in angiosperms, with

estimates suggesting that up to 70% of extant flowering plants have polyploid ancestry

(Grant 1971; Masterson 1994; Soltis and Soltis 1999; One Thousand Plant

Transcriptomes Initiative 2019). These events gave rise to extensive paralogy in plant

genomes, which poses significant challenges for orthology inference and phylogenetic

analysis. Although these issues are particularly pronounced in plants (e.g.,

Asparagaceae, Bentz and Leebens-Mack 2024; Rosaceae, Morales-Briones et al. 2018,

2021a; and Amaranthaceae, Morales-Briones et al. 2021b), they are by no means

exclusive to them. Similar complications have been documented in non-plants, such as

African clawed frogs (Evans et al. 2004), desert rodents (Gallardo et al. 2004), alpine

butterflies (Nice et al. 2013), and fish (Braasch and Postlethwait 2012), indicating that

complex histories of gene duplication and horizontal gene transfer create phylogenetic

uncertainty across the tree of life. As the scale and scope of phylogenomic datasets

expand, so too does the need to carefully model and interpret the biological processes

underlying gene tree discordance. Fortunately, the growing availability of phylogenomic

data now makes it possible to start untangling these complex histories in ways that were

not previously feasible; however, doing so requires tools and models that explicitly

account for these processes.

In response, several bioinformatic tools and pipelines have been developed to

infer orthology under increasingly realistic evolutionary models. For instance, softwares

such as HybPiper (Johnson et al. 2016) and Phyluce (Faircloth 2016) implement distinct

strategies for identifying and handling putative paralogs, with some pipelines retaining

only a single gene copy (e.g., HybPiper), while others exclude all loci which exhibit

signs of paralogy (e.g., Phyluce). Although Phyluce has shown greater success in some

groups (e.g., *Antennaria* Gaertn.; Thapa et al. 2020), HybPiper typically outperforms

Phyluce across most plant lineages because of its ability to better resolve issues

associated with polyploidy.

Additional pipelines have been developed that implement distinct strategies for

identifying and handling putative paralogs using graph- or tree-based methods (e.g.,

paragone-nf; Yang and Smith 2014; Jackson et al. 2023), whereas others consider

paralogy during species tree estimation itself (e.g., ASTRAL-Pro; Zhang et al. 2020;

Zhang and Mirarab 2022a). These different approaches can yield markedly different

phylogenetic outcomes, particularly in polyploid-rich systems where paralogs may be

the rule rather than the exception. Yet, few studies have systematically compared these

methods to understand their impact on phylogenetic and downstream evolutionary

inferences.

This is especially relevant for groups such as *Packera* Á. Löve & D. Löve

(Asteraceae: Senecioneae), a North American genus of about 91 species known for

polyploidy, complex patterns of hybridization, and poorly resolved taxonomy. An

estimated 40% of *Packera* taxa present polyploidy, aneuploidy, and/or other cytological

disturbances (Barkley 1988; Trock 2006), complicating phylogenetic reconstruction of

this group (Bain and Golden 2000). Given this, *Packera* provides an ideal system for

testing how alternative strategies for handling paralogs influence phylogenomic

inference, owing to its extensive polyploidy, ecological and geographic diversity, and

history of reticulation; all factors that increase the potential for gene tree discordance.

Here we present the most comprehensive phylogenomic investigation of *Packera*

to date, sampling nearly 93% of the genus and leveraging a high-resolution Hyb-Seq

dataset. We applied a robust framework for evaluating the impact of gene duplication by

comparing multiple paralog processing methods, including pipelines that remove all

paralogs, those that model duplication and loss, and those that retain a single

representative gene copy. We assessed how these approaches influence gene tree

concordance, species tree topology, and node support. We then tested how three of

these methods (a standard paralog processing method, a method that removes all

paralogs, and a selective paralog-aware method; see methods below) alter the

interpretations of downstream macroevolutionary analyses including divergence dating,

historical biogeography, and detection of ancient introgression. Our results demonstrate

that the choice of paralog processing approach strongly influences inferred species

relationships, gene tree concordance, node support, and evolutionary inferences,

highlighting the profound effect that paralog-aware decisions can have in

phylogenomics. Additionally, this study provides a broadly applicable framework for

assessing paralogs and their downstream impacts on phylogenetic inference in systems

shaped by gene duplication, reticulation, and rapid diversification.

MATERIALS & METHODS

Specimen Collection

A nearly complete sampling of currently recognized *Packera* species and

varieties were used in this study (62 *Packera* species, 22 varieties, and one

nothospecies; totaling 85 *Packera* taxa). *Packera* species were selected based on

information published in the International Plant Names Index (IPNI; available at

http://www.ipni.org) and then curated from discussions with botanists that have

previously specialized in *Packera* (D. Trock & J. Bain, pers. comm.). Outgroup taxa from

tribes Senecioneae (n = 19) and Anthemideae (n = 4) were selected by consulting

recent studies of the group (Bain and Golden 2000; Pelser et al. 2010; Mandel et al.

2019). Additionally, three taxa from tribe Calenduleae were included for downstream

analyses that require more distantly related outgroups that do not have a known WGD

event (Barker et al. 2016; Huang et al. 2016; Zhang et al. 2021). A complete list of the

111 sampled species, herbarium vouchers, and their publication status can be found in

Supplementary Table 1.

DNA Extraction and Sequencing

Leaf tissues were obtained from herbarium specimens or silica-dried field

collections. Dried leaf tissue (~25 mg) was ground using a Bead Mill 24 Homogenizer

(ThermoFisher Scientific, Atlanta, Georgia, USA), and total DNA was extracted using

the Omega Bio-Tek Kit (Atlanta, Georgia, USA) following manufacturer's protocols with

the addition of polyvinylpyrrolidone (PVP) and ascorbic acid to the first extraction buffer

(10 mL SQ1 buffer, 100 mg PVP, 90 mg ascorbic acid per reaction). Samples of

Packera and outgroup species that were obtained from other researchers followed

different DNA extraction protocols which are described in Supplementary Table 1. All

DNA samples used in this study were quality checked using a Nanodrop 2000 (Thermo

Fisher Scientific, Carlsbad, California, USA) and quantified using a Qubit High

Sensitivity assay (ThermoFisher Scientific, Oregon, USA). If the DNA had low

concentrations or possible contaminants, the samples were cleaned following the

E.Z.N.A. Cycle Pure Kit (Omega Biotek, Georgia, USA) and re-quantified with the Qubit

High Sensitivity (HS) assay. Up to 1 μg DNA per sample was sonicated with a QSonica

machine (ThermoCube, New York, USA) to generate fragment sizes within the 400-500

base pair (bp) range. The amount of DNA used was dependent on the initial total DNA

concentration of the targeted sample. Time allotted for sonication of each sample varied

depending on approximate length and degree of fragmentation of the DNA. Prior to

sonication, total DNA was visualized on a 1% agarose gel in 1X TBE and GelRed 3x

(Biotium) and the approximate length was estimated by comparing to a 100 bp

GeneRuler DNA ladder (ThermoFisher Scientific, Vilnius, Lithuania). If the DNA sample

appeared smaller than 400 bp, it was not sonicated.

Libraries for each sample were prepared using the NEBNext Ultra II DNA Library

Prep and NEBNext Multiplex Oligos for Illumina Kits (New England Biolabs, Ipswich,

Massachusetts, USA) following the NEBNext Ultra II Version 5 protocol with size

selection on DNA fragments at 300-400 bp range. The protocol was adjusted by halving

all amounts of reagents and DNA used. The barcoded libraries (NEBNext multiplex

oligos) were quality checked with an Agilent Bioanalyzer (Agilent Technologies, Santa

Clara, California, USA) and quantified by Qubit HS assay. Libraries were then used for

targeted sequence capture with the custom MyBaits probe set, Compositae-1061

(Mandel et al. 2014), from Arbor Biosciences (Ann Arbor, Michigan, USA) following the

manufacturer's protocol (v. 2.3.1). Approximately 500 ng of total DNA library in 6 µl of

library, or equal amount of DNA per sample when pooling (maximum of 8 samples), was

used in each MyBaits reaction for hybridization with the baits for 36 hours. Hybridized

targets were recovered using Dynabeads M-280 Streptavidin (ThermoFisher Scientific,

Vilnius, Lithuania) following the manufacturer's protocol. Captured targets were

amplified and quantified using KAPA library quantification kits (Kapa Biosystems,

Wilmington, Massachusetts, USA). A 1:4 ratio of library was added to each of the

captured target pools to obtain off-target plastid sequence data. Libraries were

sequenced on Illumina MiSeq (300 cycles) and NovaSeq 6000 (300 cycles) instruments

at Psomagen (Rockville, Maryland, USA) across several runs. Sequence stats were

obtained using the pxlssq function in Phyx v. 1.3.1 (Supplementary Table 2) (Brown et

al. 2017).

Sequence Processing

Raw FASTQ data were trimmed of low quality bases and adapter contamination

using Trimmomatic v. 0.36 (Bolger et al. 2014). The Sliding Window quality filter was

used in this analysis (illuminaclip 2:30:10, leading 20, trailing 20, sliding window 5:20).

Cleaned reads with a minimum length of 36 bp and corresponding forward and reverse

pairs were retained. FastQC v. 0.11.9 (Andrews 2010) was used to ensure good

sequence quality.

Plastid Assembly

Chloroplast sequences were extracted from trimmed reads using two

approaches. First, the GetOrganelle v. 1.7.7.1 toolkit (Jin et al. 2020) was used to

assemble circularized plastomes for all sequences with sufficient off-target sequencing

data. The recommended parameters for embryophyte plant plastomes were used,

except that the number of rounds and the disentangle time limit were increased (-R 75

and --disentangle-time-limit 7200). All complete circular plastomes were annotated

using GeSeq (Tillich et al. 2017). From these, a complete *Packera* representative

plastome was chosen and used as a reference. Alternatively, all reads were mapped to

the reference plastome using Bowtie2 (Langmead and Salzberg 2012), variants were

called using SAMtools v. 1.18 and BCFtools v. 1.18 (Danecek et al. 2021), and

consensus sequences were created using seqtk v. 1.4 (https://github.com/lh3/seqtk).

For the aligned consensus sequences, model selection and maximum likelihood

phylogenetic analysis were then conducted using IQ-TREE v. 2.2.2 (Minh et al. 2020)

and visualized using 'ggtree' (Yu et al. 2017) in R v. 4.3.1 (R Core Team 2016; RStudio

2020).

Nuclear Ortholog Assembly

Cleaned FASTQ data were assembled using the HybPiper v. 2.1.6 (Johnson et

al. 2016) pipeline to match the reads to the target loci contained in the Compositae-

1061 probe set (Mandel et al. 2014). In HybPiper, the command 'assemble' was used to

map the paired-end reads to the target loci using BWA v. 0.7.17 (Li and Durbin 2009.

These were then assembled into contigs with specified kmer lengths (21, 33, 55, 77, 99)

using SPAdes v. 3.5 (Bankevich et al. 2012). Paralog detection was carried out using

the 'paralog_retriever' command

(https://github.com/mossmatters/HybPiper/wiki/Paralogs), which flags loci that indicate

potential paralogs, then extracts the assemblies that were flagged. A supercontig

multifasta for each locus was generated and aligned in MAFFT v. 7.407 (Katoh and

Standley 2013) using default parameters.

Phylogenetic Analyses

Given the complex history of hybridization and polyploidization events in

Packera, along with the large number of paralogs flagged from HybPiper in the current

study (Supplementary Table 3), we sought to compare how different published

orthology inference methods and species tree estimation tools impacted the inferred

evolutionary relationships and support of this group. Our dataset is uniquely adapted to

tackle this issue given dense taxon sampling and high locus recovery across both

orthologs and paralogs, combined with decades of natural history and systematics work

in the group, enabling more thorough investigations into these challenges.

We generated multiple nuclear phylogenies utilizing different paralog processing

tools to infer orthology and species tree estimation: Concatenation (individual gene

histories not accounted for), HybPiper with ASTRAL-III (hereafter 'ASTRAL'; putative

single-copy orthologous loci are chosen based on coverage depth and sequence

similarity concerning the gene references from Compositae-1061, then inferred using

coalescent methods), HybPiper with wASTRAL (hereafter 'wASTRAL'; similar to

ASTRAL, but low support gene tree branches are contracted prior to species tree

estimation), HybPiper with ASTRAL-Pro (hereafter 'ASTRAL-Pro'; all single-copy and

multi-copy genes identified by HybPiper are considered, and the species tree that

optimizes a quartet similarity measure is chosen), and the 'one-to-one orthologs' (1-to-

1), 'maximum inclusion' (MI), 'monophyletic outgroup' (MO), and 'rooted ingroup' (RT)

paralog processing methods by Yang and Smith (2014) with species trees inferred

using ASTRAL-III (further details below; refer to Supplementary Table 3, Supplementary

Fig. 1). By performing comparisons among methods, we aimed to gain a better

understanding of the impact of paralogy on phylogenomic studies, and ultimately

increase phylogenomic support and concordance across species of complex lineages.

Concatenation.—Aligned supercontigs from MAFFT were concatenated with

FASconCAT-G v. 1.02 (Kück and Longo 2014), then implemented in PartitionFinder v.

2.1.1 (Lanfear et al. 2012) to determine the most appropriate nucleotide substitution

model (GTR, GTR+Γ, GTR+I+Γ). The overall best fitting model (Supplementary Table 4)

was used to build a concatenated nuclear maximum likelihood tree in RAxML v. 8.1.3

(Stamatakis 2014) with 1,000 bootstrap (BS) replicates.

ASTRAL and wASTRAL.—For the ASTRAL and wASTRAL analyses, nuclear

gene alignments were separated based on the most suitable nucleotide substitution

model for each gene as determined by PartitionFinder (Supplementary Table 4), then

individual gene trees were generated using RAxML. For the ASTRAL analysis, the gene

trees were used in ASTRAL-III v. 5.7.3 (Zhang et al. 2018) to generate the species

trees. For the wASTRAL analysis, or weighted ASTRAL, a species tree was generated

using 'astral-hybrid' in the ASTER program (https://github.com/chaoszhang/ASTER).

This method has shown to be more accurate than typical ASTRAL methods since

wASTRAL incorporates gene tree uncertainty by weighting quartets based on the

confidence levels of the input gene trees. In doing so, wASTRAL reduces the influence

of erroneous gene trees, leading to a more reliable species tree estimation (Zhang and

Mirarab 2022b). Local posterior probability (LPP) values were estimated for branch

support and indicate the probability that the branch is the true branch given the set of

gene trees provided (Sayyari and Mirarab 2016).

ASTRAL-Pro.—ASTRAL-Pro2 (ASTRAL for PaRalogs and Orthologs) v. 1.13.1.3

(Zhang et al. 2020; Zhang and Mirarab 2022a), hereafter referred to as ASTRAL-Pro,

provides a quartet-based species-tree inference method that allows multi-copy gene

trees to be used as input. Thus, all paralogous and orthologous sequences as assigned

by HybPiper were used to generate the species tree instead of selecting a single

ortholog to represent an individual gene's history (e.g., ASTRAL method). By

accommodating multi-copy gene trees, ASTRAL-Pro accounts for gene duplication and

loss events, reducing biases that arise from treating paralogs as true orthologs. This

approach is intended to improve species tree accuracy, particularly in datasets with

extensive gene family expansions, by leveraging more information from complex gene

histories. The ASTRAL-Pro pipeline is similar to ASTRAL, as described above, but

differs by having the user generate separate alignments and conduct maximum

likelihood analyses on the genes that are flagged as paralogs and single-copy contigs.

Single-copy and paralogous gene sequences were independently aligned using

MAFFT, and gene trees were generated under the best-fitting nucleotide substitution

model from PartitionFinder (Supplementary Table 4) in RAxML with 1,000 BS replicates.

The RAxML output files were then used as input for ASTRAL-Pro2 to generate the

species tree with LPP values at each node.

1-to-1, MI, MO, and RT methods.—Paralog processing was then carried out

using the tree-based orthology inference strategies originally proposed by Yang and

Smith (2014) via the paragone-nf v. 1.1.0 pipeline (Jackson et al. 2023;

https://github.com/chrisjackson-pellicle/paragone-nf). This pipeline investigates the loci

flagged as paralogous from HybPiper and determines which loci can be retained or

pruned given specified conditions (Yang and Smith 2014; Morales-Briones et al. 2021a).

All orthology inference methods proposed by Yang and Smith (2014)–the 'one-to-one

orthologs' (1-to-1), 'maximum inclusion' (MI), 'monophyletic outgroup' (MO), and 'rooted

ingroup' (RT) methods–were tested. Briefly, the 1-to-1 approach is the simplest strategy

in that it only retains homologs that have no duplications. The MI approach extracts the

subtree with the highest number of taxa without taxon duplications. The MO approach

filters for homolog trees where outgroup taxa (here tribe Calenduleae: *Calendula*

arvensis, *Tripteris pinnatilobatum* (Norl.) B.Nord., and *Chyrsanthemoides incana*

(Burm.f.) Norl.) are monophyletic and non-repeating. The RT approach searches

subtrees of ingroup taxa and roots them by the designated outgroups, then removes the

ingroup taxa as rooted trees. In turn, inferred gene duplications, or paralogs, are then

pruned from root to tip in the rooted tree (Yang and Smith 2014; Morales-Briones et al.

2021a).

To aid in consistency across analyses, the HybPiper run described above was

used as input instead of using the hybpiper-nf pipeline (Jackson et al. 2023). The three

Calenduleae taxa were considered the internal outgroups since a recent whole genome

duplication event has not been reported for this tribe, as is required for the MI and MO

analyses (Yang and Smith 2014). The resulting trimmed alignments produced via

paragone-nf were used as input for RAxML to generate gene trees, and a species tree

with LPP support values was generated using ASTRAL-III following the 'ASTRAL'

protocol described above.

Measuring phylogenetic tree support

The normalized quartet score was calculated for all orthology inference methods

that generated a species tree with ASTRAL-III using the '-q' option. Branch lengths were

recalculated from coalescent units to nucleotide substitutions per codon site for each

ASTRAL-III species tree using IQ-TREE. Parsimony informative sites (PIS) and pairwise

identity (PI) were calculated for each gene matrix using the

'parsimony_informative_sites' and 'pairwise_identity' functions, respectively, in phykit v.

2.0.1 (Steenwyk et al. 2021). Nuclear and plastome trees were visualized using the

package 'phytools' (Revell 2012) in R v. 4.0.5.

Measuring Discordance

Species trees from concatenation, plastid, ASTRAL, wASTRAL, and paragone-nf

(1-to-1, MI, MO, and RT) were analyzed for underlying discordance using Quartet

Sampling (Pease et al. 2018). ASTRAL-Pro was not analyzed as Quartet Sampling

does not allow multi-copy gene trees as input. Quartet Sampling assesses the

confidence, consistency, and informativeness of internal tree relationships, and the

reliability of each terminal branch by evaluating if a phylogeny has a lack of support due

to lack of information, discordance from ILS or introgression, or misplaced or erroneous

taxa. Three scores were calculated for each internal branch of the focal tree which

determines the consistency of information (Quartet Concordance [QC]), the presence of

secondary evolutionary histories (Quartet Differential [QD]), and the amount of

information (Quartet Informativeness [QI]). Quartet Fidelity (QF) scores were calculated

at each tip to report how frequently a taxon is included in concordant topologies. Using

the rooted species trees and concatenated gene alignments from FASconCAT-G

(described above), Quartet Sampling was run using default settings on the designated

analyses. The concatenation, plastid, ASTRAL, wASTRAL, 1-to-1, MI, and MO species

trees were rooted to the outgroup Calenduleae taxon, *Tripteris pinnatilobatum*, while the

RT tree was rooted to *Doronicum pardalianches* L. using the *pxrr* function in Phyx

(Supplementary Table 3).

Additionally, PhyParts v. 0.0.1 (Smith et al. 2015) was used with default settings

to quantify and visualize discordance in the ASTRAL, wASTRAL, and paragone-nf (1-to-

1, MI, MO, and RT) trees. Similar to Quartet Sampling, ASTRAL-Pro could not be

analyzed because of multi-copy gene trees. Additionally, the concatenation and plastid

trees could not be analyzed since those analyses do not use individual gene histories

as input. Resulting trees containing pie charts indicating whether gene trees were

supported, conflicted, or uninformative at each node were generated using a custom

python script

(https://github.com/mossmatters/phyloscripts/tree/master/phypartspiecharts). PhyParts

requires rooted gene trees and a rooted species tree as input. Similar to Quartet

Sampling, ASTRAL, 1-to-1, MI, and MO gene trees were rooted to *Tripteris*

pinnatilobatum and the RT tree was rooted to *Doronicum pardalianches* using the *pxrr*

function in Phyx. Rooting gene trees for PhyParts decreased the number of genes used

as input since not all gene trees contained the targeted outgroup taxon (see

Supplementary Table 3). For phylogenies generated with ASTRAL technologies–

ASTRAL, wASTRAL, ASTRAL-Pro, and paragone-nf trees (Supplementary Table 3)–

introgression and ILS were visualized across lineages using phytop v. 0.3.1 (Shang et

al. 2025) with default settings. The species trees were re-ran with the flag '-t 2' added to

be used as input for phytop.

Tree Topology Comparison

To evaluate similarity among the nuclear gene trees, pairwise adjusted

Robinson-Foulds (RF_{adj}) distances were calculated as outlined by (Mitchell et al. 2017)

between all trees using the *RF.dist()* function in package 'phangorn' (Schliep 2011) in R,

with unrooted trees as input and setting the "normalize" argument to TRUE (Steel and

Penny 1993). The Calenduleae outgroup taxa were trimmed from the trees in R using

the *drop.tip()* function in package 'ape' (Paradis and Schliep 2019) to make

concatenation, ASTRAL, ASTRAL-Pro, and paragone-nf trees directly comparable. The

RF_{adj} values range from zero to one, indicating whether the tree topologies are identical

or completely dissimilar, respectively. A principal component analysis (PCA) was

generated from all RF_{adj} values in R to aid in visualization.

All nuclear and plastid phylogenies and their associated analyses described

above were compared, but for simplicity only the ASTRAL, 1-to-1, and MO trees were

analyzed in further analyses. ASTRAL was chosen as it selects the putative single-copy

orthologs less rigorously than the remaining methods, creating a baseline method for

comparison. We then compared ASTRAL to a method that disregards all genes

containing paralogs (1-to-1) and a more rigorous paralog-processing method (MO). We

chose MO over MI and RT due to its more stringent ortholog identification, which

minimizes the inclusion of paralogs and reduces phylogenetic noise. Additionally, the

conservative filtering of MO enhances alignment quality and computational efficiency,

providing a more reliable foundation for downstream phylogenetic analyses (Yang and

Smith 2014; Morales-Briones et al. 2021a; Jackson et al. 2023).

Molecular Clock Dating

Estimates of divergence times were calculated on the ASTRAL, 1-to-1, and MO

trees with RelTime (Tamura et al. 2012, 2018) using the command-line version of the

MEGA software, MEGA-CC v. 11.0.11 (Kumar et al. 2012, 2016), with necessary input

files generated with MEGA-X (Kumar et al. 2018; Mello 2018). The three Calenduleae

taxa were used as outgroup taxa for this analysis.

There are no fossils available within *Packera* or tribe Senecioneae, so instead,

multiple dating scenarios were tested utilizing both primary and secondary age

constraints (Supplementary Table 5). Briefly, a maximum (59.95 MY) and minimum (31

MY) age was tested for the Anthemideae/Senecioneae split given those dates are the

highest and lowest values within the 95% CI from a previous diversification rate analysis

of Asteraceae (Mandel et al. 2019). Next, a maximum age of 34.06 MY was set at the

Packera crown node from the upper age limit provided by (Mandel et al. 2019).

Additionally, a minimum age constraint of 12 MY was tested within the outgroup (the

Anthemideae clade) given the presence of a fossil pollen (Wang 2004; Hobbs and

Baldwin 2013; Palazzesi et al. 2022). Finally, a maximum age constraint of 14.3 MY

was tested for *Pericallis* D.Don (Senecioneae), a genus endemic to the Macaronesia

islands (Jones et al. 2014), since that is the estimated age of Porto Santo, the oldest

volcanic archipelago within the Madeira Islands (McDougall and Schmincke 1976;

Geldmacher et al. 2000). Results of all tested scenarios were compared for biological

relevance, though only one scenario was chosen for the discussion. This approach

allowed us to focus on the topology that best aligned with prior systematic

understanding and showed the strongest overall support across analytical frameworks.

Biogeography

Ancestral ranges were estimated on the dated phylogenies using the package

BioGeoBEARS v. 1.1 (Matzke 2014) in R. BioGeoBEARS estimates historical,

biogeographic ranges by implementing popular models in a likelihood framework. Some

of the most common models used are dispersal-extinction-cladogenesis (DEC; Ree and

Smith 2008), the likelihood version of DIVA (DIVALIKE; Ronquist 1997), and the

BayArea likelihood version of the range evolution model (BAYAREALIKE; Landis et al.

2013). In BioGeoBEARS, founder-event speciation can be added to any of these

models (+*J*) with flexibility in determining relative probability. Likelihood Ratio Test (LRT)

and Akaike Information Criterion (AIC) were used to determine which model best fits the

data.

All *Packera* taxa, along with their four most closely related outgroup genera

(*Pericallis*, *Elekmania* B.Nord., *Werneria* Kunth, and *Xenophyllum* V.A.Funk), were

coded as present or absent in ten geographic ranges outlined in Takhtajan (1986)

(Supplementary Table 6): Arctic Province, Canadian Province, Appalachian Province,

Atlantic and Gulf Coast Plain Province, North American Prairies Province, Rocky

Mountain and Vancouverian Province, Madrean Region, Asia, Europe, and Central

America. Distributions were determined from previously published literature (Freeman

and Barkley 1995; Trock 1999, 2006; Gramling 2006; Mahoney and Kowal 2008; Elven

et al. 2011; Yeatts et al. 2011). Only four of the 26 outgroup taxa were included to limit

the number of geographic ranges used in the analysis. The remaining outgroup taxa

were pruned from the dated tree generated with RelTime using the *pxrmt* function in

Phyx. Each taxon was limited to a maximum distribution of four ranges.

Reticulation

The Species Networks applying Quartets (SNaQ) function in the Julia v. 1.8.1

(Bezanson et al. 2018) package PhyloNetworks v. 0.15.2 (Solís-Lemus and Ané 2016;

Solís-Lemus et al. 2017) was used to investigate reticulate evolution among *Packera*

taxa. This package uses unrooted gene trees to infer phylogenetic networks in a

pseudolikelihood framework with multi-locus data, while accounting for ILS and

reticulation. We tested the fit of multiple models and gradually increased the number of

reticulation events (h) from 0 to 10 to find which pseudolikelihood score reached a

plateau, given recommendations by (Solís-Lemus et al. 2017). This process is

computationally intense, so only clades containing 14 taxa or less that had counter-

support nodes in the ASTRAL, 1-to-1, and/or MO trees, as determined by a negative

QC score resulting from Quartet Sampling, were used as input for PhyloNetworks, with

the remaining taxa pruned from the tree using the *pxrmt* function in Phyx. Given this,

five Packera clades were investigated for ancient reticulation events (Figs. 2-4).

Additionally, a single species from each major clade was then used to represent

backbone relationships throughout the entire phylogeny, totaling six separate

PhyloNetwork runs that will be referenced as follows: Backbone, representing 11 taxa

that span the entire phylogeny; Eastern, a clade of 12, 13, or 15 species (MO, 1-to-1,

and ASTRAL, respectively) either distributed along eastern North America or having

known affiliations with those species; California, eight taxa with distributions mainly

within California; Mexico, seven (MO and ASTRAL) to eight (1-to-1) taxa with

distributions mainly within Mexico; Rocky, a clade of five taxa that are distributed in the

Central US, most within the Rocky Mountains; and Arctic, nine taxa that are present in

colder climates along the Pacific Northwest and into Canada (Supplementary Table 7).

Rocky could not be run for 1-to-1 since fewer than five taxa were present in that target

clade.

RESULTS

Sequencing and Matrix Results

Illumina sequencing resulted in approximately 733.3 million reads across 111

taxa. Read numbers ranged from 535,934 in *Werneria aretioides* Wedd. to 20,685,576

in *Senecio vulgaris* L. (full list in Supplementary Table 2). HybPiper retained 1,058 of

the targeted 1,061 loci across all 111 taxa with 846 of those loci having paralog

warnings (~80%). The number of loci recovered for each taxon ranged from 566 in

Packera paupercula var. paupercula (Michx.) Á.Löve & D.Löve to 1,054 in *Calendula*

arvensis L. (Supplementary Table 8). The number of gene trees varied across analyses,

from 285 in the 1-to-1 analysis to 7,388 in the MI analysis (Supplementary Table 3).

Parsimony informative sites (PIS) also ranged from 2.11% in the plastid (mapping) tree

to 99.64% in the MO tree (Supplementary Table 3). For species trees generated with

ASTRAL-III, normalized quartet scores were used to measure gene tree discordance,

which ranged from 0.516 for the RT tree to 0.534 for the MO tree (Supplementary Table

3), indicating that ~52-53% of the gene trees matched the species tree in those

analyses.

Off-target plastid sequencing resulted in partial (i.e., mapping; n = 111) to whole

(i.e., complete; n = 18) plastomes. Complete plastomes from 17 *Packera* and one

Senecio species ranged from 150,772 bp to 151,728 bp in size (Supplementary Fig. 2).

For each consensus sequence mapped to the reference plastid assembly, *Packera*

cymbalaria, the percentage of missing data varied drastically, with *Tephroseris*

newcombei exhibiting the most missing data (93.1%; Supplementary Table 9,

Supplementary Fig. 3). The mapped plastid supermatrix with 111 taxa had 2.11% PIS

(Supplementary Table 3).

Topological Comparisons and Discordance

Phylogenetic topologies varied depending on sequence type (e.g., nuclear or

plastid) as well as paralog processing and phylogenetic tree inference method (i.e.,

ASTRAL, ASTRAL-Pro, 1-to-1, etc.) (Fig. 1, Supplementary Fig. 4). When building

phylogenies with nuclear data, *Packera* was generally placed as the sister lineage to a

clade comprising *Pericallis* and *Elekmania*, with the exception of ASTRAL-Pro which

Figure 1. Genus-level phylogenies of all trees generated in this study. **A**) Genera are colored according to their tribal/subtribal classifications following the key on the top of the figure. *Packera* is marked with a black asterisk next to its placement in the trees. Navy blue text at node represents support values (LPP or BS). **B**) Bottom left is a matrix of RF$_{adj}$ values between the nine different trees, with most similar topologies as dark blue squares (0) and least similar as yellow squares (1). The bottom right shows the RF$_{adj}$ values plotted as a principal components analysis (PCA). Points closer together indicate higher similarity, while dots further away indicate higher dissimilarity.

placed *Packera* as sister to *Tephroseris* and in a larger clade with *Pericallis* and

Elekmania (Fig. 1). Interestingly, the plastid data showed a similar relationship to

ASTRAL-Pro; however, *Pericallis* was placed unexpectedly in a separate clade with

other Senecionineae taxa. All nuclear phylogenies recovered monophyletic tribes and

subtribes (e.g., Senecionineae and Tussilaginineae s.str.), though the relationships

within the subtribes varied. The mapped plastid tree generally separated the subtribes,

apart from *Roldana* (Senecionineae) in Tussilaginineae s. str. (Fig. 1, Supplementary

Fig. 4). The placement of *Tephroseris* (tribe Tephroseridineae) varied across

phylogenies, with its most consistent placement as sister to Senecioneae (Fig. 1).

Doronicum was consistently placed sister to Senecioneae, providing support for its

elevation to tribal status (Doroniceae; Panero 2005; Fu et al. 2016).

Phylogenetic topologies remained highly incongruent independent of methods

used (Figs. 2-4, Supplementary Table 10). The ASTRAL and wASTRAL tree topologies

were most similar (RF_{adj} = 0. 324) (Fig. 1, Supplementary Table 10), while the plastid

tree was consistently the most dissimilar to the remaining trees (RF_{adj} ranging from

0.895 to 0.914), with the concatenated tree being the next most dissimilar (RF_{adj} ranging

from 0.562 to 0. 648; Supplementary Table 10). Additionally, the normalized quartet

scores from ASTRAL indicate that only about half of the gene trees match the species

trees in the ASTRAL and paragone-nf analyses (Supplementary Table 3). Most of the

topological conflict was among *Packera* taxa, with minor differences among the

remaining Senecioneae taxa and outgroups (Supplementary Fig. 4). Support values

varied depending on which sequence type and method was used, though generally

remained low. The wASTRAL and MI trees showed the highest node support with 58

ASTRAL

Figure 2. Full phylogeny of the ASTRAL analysis. Tribal/subtribal status is indicated on the right of the phylogeny and follows the color code as indicated in the key. Numbers next to tip labels correspond to the Quartet Fidelity (QF) score from Quartet Sampling, which ranges from 0 to 1, with values closer to 0 indicating erroneous taxa. Circles at nodes show the Quartet Concordance (QC) factors from Quartet Sampling, where red and orange indicate counter-support nodes. Asterisks next to QC circles indicate high support (≥ 95 LPP) at that node. Highlighted clades and teal stars next to some tip labels, indicate clades where reticulation events were tested, following the color key on the left.

Figure 3. Full phylogeny of the 'one-to-one ortholog' (1-to-1) analysis. Tribal/subtribal status is indicated on the right of the phylogeny and follows the color code as indicated in the key. Numbers next to tip labels correspond to the Quartet Fidelity (QF) score from Quartet Sampling, which ranges from 0 to 1, with values closer to 0 indicating erroneous taxa. Circles at nodes show the Quartet Concordance (QC) factors from Quartet Sampling, where red and orange indicate counter-support nodes. Asterisks next to QC circles indicate high support (≥ 95 LPP) at that node. Highlighted clades and teal stars next to some tip labels, indicate clades where reticulation events were tested, following the color key on the left.

MO

Figure 4. Full phylogeny of the 'monophyletic outgroup' (MO) analysis. Tribal/subtribal status is indicated on the right of the phylogeny and follows the color code as indicated in the key. Numbers next to tip labels correspond to the Quartet Fidelity (QF) score from Quartet Sampling, which ranges from 0 to 1, with values closer to 0 indicating erroneous taxa. Circles at nodes show the Quartet Concordance (QC) factors from Quartet Sampling, where red and orange indicate counter-support nodes. Asterisks next to QC circles indicate high support (≥ 95 LPP) at that node. Highlighted clades and teal stars next to some tip labels, indicate clades where reticulation events were tested, following the color key on the left.

out of 110 nodes (52.7%) having ≥ 0.95 LPP (Supplementary Table 3,

Supplementary Fig. 4). In contrast, the plastid and concatenated trees showed much

lower resolution with the fewest number of nodes having high support (36 of 110 nodes,

32.7%, have ≥ 95 BS; Supplementary Table 3, Supplementary Fig. 4).

Quartet Sampling indicated that underlying gene tree discordance remained high,

independent of the method used. ASTRAL and concatenated trees had the most fully

supported nodes (QS = 1/–/1) when compared to the remaining phylogenies

(Supplementary Fig. 5), though the number was still notably low (~10%). Even so, the

concatenated tree also had the highest proportion of counter-support nodes (QC < 0;

26.4%; see Supplementary Fig. 5 for all Quartet Sampling results). PhyParts indicated

that the ASTRAL and MO trees had the most concordant gene trees. The wASTRAL

analysis had the highest percentages of gene trees supporting the top alternative

bipartition. The MI tree harbored the highest discordance and uninformative data

(Supplementary Fig. 6).

This underlying discordance could be caused by ILS, which was common in all

tested ASTRAL trees (Supplementary Fig. 7). Introgressive hybridization (IH) was not

detected without the presence of ILS in any of the nodes, though IH was most common

in the nodes that were shown to be discordant (Supplemental Figs. 5, 7). Interestingly,

crown *Packera* consistently had high IH values, potentially indicating introgression at

that node. Backbone nodes were generally considered well resolved with high

confidence (ILS index < 50% and IH index < 10%; Shang et al. 2025), especially in the

outgroups, whereas *Packera* taxa appeared less resolved.

Molecular Clock Dating

The resulting divergence rate estimates differed among ASTRAL, 1-to-1, and

MO, with estimated ages of the MO and 1-to-1 trees older and sometimes almost

double the ASTRAL ages. For ASTRAL, the mean crown age for *Packera* ranged from

11.4 MY in Scenario 5 and 6 to 13.5 MY in Scenario 1-4. Alternatively, the mean crown

age of *Packera* were almost doubled in the 1-to-1 and MO analyses, with ranges from

~21 MY in Scenario 5 to 35.5–35.7 MY in Scenario 4 (Supplementary Table 11,

Supplementary Fig. 8). Scenario 2 was chosen as the best-fitting dating model because

it incorporated all available primary calibration points and yielded divergence time

estimates that align well with known patterns of evolutionary history (Pelser et al. 2010;

Mandel et al. 2019), while minimizing reliance on secondary calibrations

(Supplementary Table 5) (Müller and Reisz 2005; Benton and Donoghue 2007; Parham

et al. 2012). The dated trees from all six scenarios can be found in Supplementary

Figure 8.

Biogeography

Evaluation of the six different model choices demonstrated that BAYAREALIKE

was the best fit given our data based on its lowest log likelihood (Ln L) and highest AIC

score, as well as the highest weighted AICc score for all three of our analyses

(Supplementary Table 12). Generally, models including jump speciation (+J) provided a

better fit than those without in the DEC and DIVALIKE models, and were the same or

very similar in the BAYAREALIKE models (Supplementary Table 12), indicating a

potential for jump dispersal or founder-event speciation in our data. Trees of all model

runs are in Supplementary Figure 9.

The historical biogeography analysis revealed distinct geographic patterns

across the different orthology methods (Fig. 5, Supplementary Fig. 9). ASTRAL

identified the ancestral range as concentrated in the North American Prairies Province

(N), Rocky Mountain and Vancouverian Province (R), and Madrean (M) regions. In

contrast, MO showed the ancestral regions being only the Rocky Mountain and

Vancouverian Province (R), and Madrean (M) regions. 1-to-1 was intermediate with

crown *Packera* being ancestral to the North American Prairies Province (N), Rocky

Mountain and Vancouverian Province (R), and Madrean (M) regions, similar to

ASTRAL; however, the remaining backbone was Rocky Mountain and Vancouverian

Province (R), and Madrean (M) regions, similar to MO. Notably, all analyses showed

one major expansion into the eastern United States–Appalachian Province (P) and

Atlantic and Gulf Coast Plain Province (T)–represented by the Eastern clade (Fig. 5,

Supplementary Fig. 9), with a few instances of individual introductions to the eastern

United States. Many of these taxa originated initially in the North American Prairies

Province (N) and/or Rocky Mountain and Vancouverian Province (R) ranges. ASTRAL

and MO showed two early expansions into the Arctic Province (I) and Canadian

Province (C) at the base of the phylogeny, while 1-to-1 showed at least three

expansions north. 1-to-1 exhibited the most geographic inconsistency, followed by MO.

Even so, geography proved to be a reliable indicator for larger clades across all

analyses, although widespread species were scattered across the phylogeny and did

not always align consistently within geographic clades (Figs. 2-4, Supplementary Fig.

9).

Figure 5. Predicted biogeographic region summaries generated from BioGeoBEARS of the ASTRAL, 1-to-1, and MO analyses. The maps of North America are colored according to the ten geographic regions outlined by Takhtajan (1986): Arctic Province (I), Canadian Province (C), Appalachian Province (P), Atlantic and Gulf Coast Plain Province (T), North American Prairies Province (N), Rocky Mountain and Vancouvarian Province (R), Madrean Region (M), Asia (A), Europe (E), and Central America (S). Stars indicate the predicted ancestral range of crown *Packera*. Arrows and numbers correspond to their predicted range expansions in chronological order, with 5 being the most recent major expansion. Numbers also correspond to the dated tree to the right of the phylogeny, which indicates the nodes the expansions occurred in relation to the major geographic clades designated in Figures 2-4. All BioGeoBears results can be found in Supplementary Figure 9. Images of *Packera* species highlighting some of the morphological diversity seen in the group. Colored bars on the bottom of the image correspond to their current distributions, following the color scheme of the map and key. From left to right: Flame ragwort, *P. greenei* (A.Gray) W.A.Weber & Á.Löve; Tehachapi ragwort, *P. ionophylla* (Greene) W.A.Weber & Á.Löve; Fendler's ragwort, *Packera fendleri* (A.Gray) W.A.Weber & Á.Löve; Weak groundsel, *P. debilis* (Nutt.) W.A.Weber & Á.Löve; and Rayless mountain ragwort, *P. indecora* (Greene) Á.Löve & D.Löve. Image credits: Erika R. Moore-Pollard and Robert Lagier.

Reticulation events

Five *Packera* clades were tested for ancient reticulation events given a large

number of counter-support (QC < 0) nodes from Quartet Sampling on the ASTRAL, 1-

to-1, and MO trees (Figs. 2-4). Additionally, a single species from each major clade was

then used to represent backbone relationships throughout the entire phylogeny, totaling

six separate PhyloNetwork runs named: Arctic, Backbone, California, Eastern, Mexico,

and Rocky (Supplementary Table 13). Log-likelihood values indicated that the most

likely number of reticulation events (h) varied depending on the run (Supplementary Fig.

10). Additionally, Arctic, Mexico, and Backbone had varying numbers of optimal

reticulation events between the paralog-processing methods (Supplementary Table 7,

Supplementary Fig. 10). For ASTRAL, Arctic and Mexico had the highest number of

reticulation events (h = 3), followed by California and Backbone (h = 2), then Eastern

and Rocky (h = 1). For 1-to-1, Backhone had the most (h = 6), then Mexico (h = 3),

Arctic and California (h = 2), and then Eastern (h = 1). For MO, Arctic, California, and

Backbone had the highest number of reticulations (h = 2) with Eastern, Mexico, and

Rocky having the lowest (h = 1) (Fig. 6, Supplementary Fig. 10, Supplementary Table

7). The taxa also differed between analyses given different phylogenetic relationships

between clades (Supplementary Table 13), impacting some of the networks identified.

Some networks showed different events depending on the phylogenetic method

and subset of data (Fig. 6); however, some consistent patterns emerged. For example,

the Eastern clade continuously showed the same reticulation event regardless of the

paralog processing method, and parentage percentages were very similar. In contrast,

Rocky did not show the same reticulation events, but the networks generally involved

Packera mancosana and *P. streptanthifolia*. The California clade showed differing

results across methods: both ASTRAL and 1-to-1 analyses identified *P. bernardina* as

the product of reticulation with the same parentage, whereas MO did not yield the same

results, and no consistent trends emerged. The Mexico clade was generally

inconsistent, but featured reticulation events involving *P. neomexicana* and its varieties.

Additionally, *P. texensis* and *P. quercetorum* were involved in reticulation events in two

of the three paralog processing networks. Among all the clades, the Arctic clade

Figure 6. PhyloNetworks figures of the three tested analyses: ASTRAL, 1-to-1, and MO. The various areas of the phylogeny that were tested are shown on the left and can be read as a row. Dark green edges indicate the major hybrid edge, while the light blue shows the minor hybrid edge. The associated numbers next to each branch correspond to the percentage of parentage given by that branch. Number of hybridizations correspond with the best log likelihood value. The 1-to-1 analysis does not have a Rocky analysis since the clade in that tree only had four taxa.

exhibited the most variability in the number of reticulation events, with the *P. bolanderi*

varieties consistently involved in some form of reticulation. However, the specifics of

these events varied depending on sampling and paralog processing methods. In the

Backbone subtree analysis, *Tephroseris* was frequently identified as a product of

reticulation, with 1-to-1 and MO showing the same parentage. In contrast, ASTRAL did

not show the same reticulation events but did identify a reticulation event involving

Tephroseris with a different parentage. Notably, the parentage of *Tephroseris*

always included a *Packera* taxon, though the specific taxon varied.

DISCUSSION

Gene duplication was pervasive in *Packera*, with more than 80% of nuclear loci

showing evidence of paralogy (Supplementary Table 3). Such widespread paralogy

poses major challenges for species tree inference, especially in Hyb-Seq datasets,

where probes may unintentionally enrich for multiple homeologs (Salmon et al. 2012;

Akama et al. 2014; Crameri et al. 2022), complicating the recovery of true single-copy

orthologs for phylogenetic reconstructions. Although some paralogy is expected in

target enrichment studies (Siniscalchi et al. 2021), the extent observed in *Packera* is

unusually high and reflects lineage-specific genome dynamics, rather than a general

feature of all species. This has important methodological implications, as HybPiper's

orthology inference relies largely on single-copy filtering (Johnson et al. 2016), and

using its outputs alone can introduce noise and bias into gene tree reconstruction,

ultimately distorting species tree inference. Consistent with our study, using only

putative single-copy orthologs designated by HybPiper was insufficient for accurate

phylogenetic inference in *Packera*, a group characterized by extensive polyploidy and

hybridization (Fig. 2) (Trock 2006). These challenges are not unique to *Packera*, as

many plant and non-plant lineages also experience difficulties reconstructing

phylogenies due to hybridization and gene/genome duplications. Together, these

examples highlight the complexity of orthology inference in phylogenomics and the need

for rigorous paralog processing across both plant and non-plant systems.

The most recent phylogeny of *Packera* prior to this study was based on ITS data

and revealed extremely low sequence variation, with only 0–5% divergence and several

Packera taxa sharing identical sequences (Bain and Golden 2000). This led to a largely

unresolved tree, likely due to few phylogenetically informative sites, sparse species

sampling, and potential reticulation events (Funk 1985; Linder and Rieseberg 2004). By

using Hyb-Seq data and broader taxon sampling, we substantially improved

phylogenetic resolution across *Packera*, including within major clades. However, our

analyses also revealed that gene tree concordance and species tree resolution varied

substantially across methods (Supplementary Fig. 4), exemplifying the influence of

paralog processing strategies on downstream phylogenetic outcomes. Methods that

explicitly model gene duplication and loss, such as those implemented in paragone-nf

(Yang and Smith 2014; Jackson et al. 2023), generally yielded more concordant gene

trees and better-resolved species trees (Figs. 3, 4). However, substantial conflict

persisted across the trees, indicating that even sophisticated modeling approaches

cannot fully resolve deep relationships in groups with complex histories like *Packera*.

These discrepancies may in part reflect differences in the number and identity of genes

retained after filtering (Supplementary Table 3). For instance, the 1-to-1 method, which

excludes all loci with paralogs and resembles approaches used in tools like Phyluce,

eliminates a significant portion of the dataset and may oversimplify the data, especially

in duplication-rich systems such as *Packera*. While Phyluce and 1-to-1 filtering both

remove paralogs, the underlying methods differ (Faircloth 2016; Jackson et al. 2023),

causing Phyluce to outperform HybPiper in certain cases (Mandel et al. 2019; Thapa et

al. 2020). Similarly, the RT method naturally loses information since it removes

outgroups to determine orthology (Yang and Smith 2014; Morales-Briones et al. 2021b,

2021a) and potentially limits phylogenetic signals. These findings highlight that while

some approaches have potential to improve concordance, none fully resolved the

underlying discordance of *Packera*, emphasizing that the choice of method, underlying

data, and biological context must guide ortholog selection in phylogenomic analyses.

Results from a previous phylogenetic study of *Packera* showed that its

evolutionary relationships tend to loosely follow a geographic pattern (Bain and Golden

2000), which is consistent with our findings (Figs. 2-4, Supplementary Fig. 9). Our

nuclear phylogenies showed that major clades tended to contain taxa within similar

geographic ranges, independent of which paralog processing method was utilized.

Additionally, most taxa within these major clades were conserved across methods,

though their placement along the backbone, and some taxa, did vary (Figs. 2-4). For

example, the taxa in the Eastern clade was fairly consistent between the orthology

inference methods, although some taxa did move in and out of the clade

(Supplementary Fig. 4). Interestingly, several of these taxa are widespread and engage

in interspecific hybridization with other *Packera* species where their ranges overlap. For

example, *P. pseudaurea* and its varieties are typically found in central North America,

though their phylogenetic placement is generally in the Eastern clade. These results are

unsurprising as they have an extensive history of hybridization with other *Packera* taxa

in the Eastern clade, which is believed to have created much taxonomic uncertainty

(Barkley 1962; Trock 1999), potentially explaining their inconsistent phylogenetic

placements (Supplementary Fig. 4). However, conflicting signals may also reflect

additional evolutionary processes beyond hybridization, such as cryptic speciation

within widespread taxa or the influence of unsampled and/or extinct lineages.

Our results also reveal that evolutionary interpretations such as divergence

dating and biogeographic reconstruction are highly sensitive to the paralog processing

approach used to infer the underlying phylogeny. Divergence time estimates varied

across methods, with some species trees suggesting that crown *Packera*'s

diversification occurred more recently (ASTRAL: 13.5 MY), while others placed key

divergences deeper in time (1-to-1: 24.2 MY and MO: 25 MY; Supplementary Fig. 8,

Supplementary Table 11). These differences in divergence times are not trivial, as they

affect our understanding of how *Packera* taxa respond to climatic shifts, mountain uplift,

and other historical events, and have broader implications for comparative studies

across Asteraceae and other radiating plant lineages. For example, tribal- and family-

wide studies predicted the crown age of *Packera* to be fairly young, between 7.8 MY

and 20.2 MY (Pelser et al. 2010; Mandel et al. 2019; Zhang et al. 2021), which is more

in-line with the ASTRAL analysis. While more *Packera*-specific predictions are older,

with (Barkley 1988) predicting the origin of *Packera* in the later stages of the mid-

Tertiary (~22-37 MY), closely aligning with the paragone-nf analyses (Supplementary

Fig. 8). Similarly, biogeographic patterns shifted depending on the inferred topology,

with each analysis showing different migration patterns over time (Fig. 5,

Supplementary Fig. 9). Even so, the broad pattern is relatively unchanged. For

example, all topologies generally agreed that *Packera* originated in western North

America with one major expansion east and then later expansions north from the

western clades. From there, individual taxa expanded or became specialized to certain

regions, with most of the diversity present in western North America. Because the

topologies are consistent across methods, we can be more confident that they likely

reflect the true evolutionary history of *Packera* instead of methodological artifacts.

701   Despite improvements from paralogy-aware filtering, the species trees for

Packera were not fully resolved, indicating that underlying biological processes such as

ILS, hybridization, and introgression also play significant roles in shaping *Packera*'s

evolutionary history. This is supported by strong cytonuclear discordance, where plastid

phylogenetic relationships vary significantly from the nuclear trees (Fig. 1,

Supplementary Fig. 4), consistent with chloroplast capture or ancient hybridization

events (Doyle 1992; Maddison 1997; Galtier and Daubin 2008; Soltis and Soltis 2009;

Smith et al. 2015; Stull et al. 2020). Directly testing for gene flow revealed multiple

introgression events, particularly along short internal branches susceptible to ILS

(Supplementary Fig. 7). However, the specific reticulation patterns recovered varied

depending on the paralog processing method used. These discrepancies highlight a

critical point: gene selection and paralog processing directly shapes reticulation

inference. Each gene has its own evolutionary trajectory, shaped by recombination and

gene duplication and loss (Betrán 2015), and filtering strategies that exclude paralogs or

collapse duplications can erase signals of past hybridization. As a consequence of

different orthology and paralog processing methods, some reticulation events, such as

those involving Eastern *Packera* lineages, were consistently recovered across datasets,

while others appeared only in specific analyses.

Network analyses, such as PhyloNetworks, are inherently sensitive to sampling

bias (Solís-Lemus and Ané 2016). Even though we sampled 85 out of a targeted 91

Packera taxa (~93% coverage of the genus), missing taxa may have impacted our

results. Including the remaining and unsampled *Packera* taxa in future studies could

help increase our understanding of these relationships. Additionally, limiting the

analyses to specified clades can ignore other potential reticulation events occurring

across groups. We recognize this limitation; however, expanding sampling to more taxa

or loci is currently restricted by the intensive and often unfeasible computational

requirements of reticulation network analyses (see Hejase and Liu 2016). With recent

bioinformatic advances and increased interest in non-bifurcating trees, we anticipate

that less computationally intense and more efficient phylogenetic and network tools will

soon be developed, allowing researchers to test for reticulations across large

phylogenies.

Ghost introgressions, which are ancient introgression events that leave genetic

signatures of extinct species in present-day species (Ottenburghs 2020), also pose

challenges to inferences made by hybridization networks. Accounting for ghost

introgressions in macroevolutionary studies may help improve the accuracy of trait

evolution and biogeographic reconstructions, refine divergence time estimates, and

reveal hidden contributions of hybridization to diversification (Ottenburghs 2020; Tricou

et al. 2022). Network methods such as PhyloNetworks can detect ghost introgressions,

but these methods do not specify that the hybrid edge is a ghost introgression, so it is

up to the researcher to recognize its presence, which can be difficult especially without

background knowledge of the group (Solís-Lemus and Ané 2016; Solís-Lemus et al.

2017).

Our findings emphasize that accurately reconstructing reticulate evolutionary

histories requires embracing the independent histories of genes and carefully

considering how orthology is defined. In *Packera*, gene tree heterogeneity is expected

and evolutionarily informative rather than mere noise. Methods that oversimplify gene

relationships, such as pipelines restricted to single-copy orthologs, risk masking the

very patterns needed to understand complex evolutionary processes. By comparing

alternative strategies for handling paralogs and integrating gene flow analyses, we show

that approaches which explicitly model gene duplication and loss provide modest

improvements in resolution and concordance but are insufficient on their own to fully

resolve relationships. Direct tests for reticulation and ILS revealed the extent to which

Packera's evolutionary history has been shaped by hybridization and gene flow,

underscoring the broader challenge of capturing complexity in phylogenomic datasets.

Within polyploid systems, the tradeoff between excluding paralogs to reduce conflict

and retaining them to preserve evolutionary signals becomes especially pronounced.

These results also highlight how paralog handling influences not only

phylogenetic reconstruction but also the biological inferences drawn from the inferred

relationships. Macroevolutionary analyses such as divergence dating, trait evolution,

and biogeography are highly sensitive to orthology inference strategies and are only as

reliable as the phylogenetic framework on which they are built (e.g., Magalhaes et al.

2021). Improving that framework will require both methodological innovation and deeper

biological insight. For example, targeted probe sets designed to enrich truly single-copy

loci (e.g., Compositae-ParaLoss-1272; Moore-Pollard et al. 2024) offer one path

forward. Yet entirely excluding paralogs may discard informative loci, while retaining

them without appropriate modeling risks misleading inference. Integrating

morphological, ecological, and geographic evidence is therefore essential for assessing

whether patterns inferred from large-scale datasets reflect genuine evolutionary history

or methodological artifacts.

Rather than treating paralogy as a technical obstacle to be circumvented in

phylogenetic studies, our findings underscore its potential to illuminate fundamental

evolutionary processes, including gene duplication, polyploidy, and reticulation, that

have shaped the diversification of lineages across the tree of life. While these

challenges are common to many plant and non-plant groups, our densely sampled

Packera dataset provides a uniquely powerful system for testing and improving

phylogenomic methods. As the field advances toward increasingly nuanced and

biologically realistic models of genome evolution, systems like *Packera* will be essential

for developing tools that can fully capture the intricacy of complex evolutionary histories.

ACKNOWLEDGMENTS

The authors thank Drs. Carol Siniscalchi and Matthew Pollard for their

bioinformatic help, Drs. Randall Bayer, L. Ellie Buckland, Duane McKenna, Anna Pardo,

and Linda Watson for reading and editing prior to submission, and Dr. John Bain, Dr. Ed

Schilling, and Brandon Fuller for donating genetic material for this study. We would also

like to thank the following herbaria for allowing us to visit and/or destructively sample

leaf tissue: ALA, ARIZ, ASU, CAS, MEM, NY, OSC, TEX, UNC, UCR, WTU. Permits to

collect in the San Bernardino National Forest and Shasta-Trinity National Forest in

California were obtained prior to collection of specimens through the USDA Forest

Service. We also thank the University of Memphis High-Performance Cluster (HPC)

administrators, Eric Spangler and Kristian Skjervold, for their assistance with the HPC

and willingness to help. This work was supported by the Center for Biodiversity

Research at the University of Memphis.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

Raw sequence data are available in the National Center for Biotechnology

Information (NCBI) Sequence Read Archive (SRA) under BioProjects PRJNA907383,

PRJNA978568, PRJNA540287, and PRJNA516161.

REFERENCES

Akama S., Shimizu-Inatsugi R., Shimizu K.K., Sese J. 2014. Genome-wide
quantification of homeolog expression ratio revealed nonstochastic gene regulation in
synthetic allopolyploid *Arabidopsis*. *Nucleic Acids Res.* 42:e46.

Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data.
Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

Bain J.F., Golden J.L. 2000. A phylogeny of *Packera* (Senecioneae; Asteraceae) based
on internal transcribed spacer region sequence data and a broad sampling of
outgroups. *Mol. Phylogenet. Evol*. 16:331–338.

Bankevich A., Nurk S., Antipov D., Gurevich A.A., Dvorkin M., Kulikov A.S., Lesin V.M.,
Nikolenko S.I., Pham S., Prjibelski A.D., Pyshkin A.V., Sirotkin A.V., Vyahhi N., Tesler
G., Alekseyev M.A., Pevzner P.A. 2012. SPAdes: A new genome assembly algorithm
and its applications to single-cell sequencing. *J. Comput. Biol*. 19:455–477.

Barker M.S., Li Z., Kidder T.I., Reardon C.R., Lai Z., Oliveira L.O., Scascitelli M.,
Rieseberg L.H. 2016. Most Compositae (Asteraceae) are descendants of a
paleohexaploid and all share a paleotetraploid ancestor with the Calyceraceae. *Am. J.*
Bot. 103:1203–1211.

Barkley T.M. 1962. A Revision of *Senecio aureus*. *Trans. Kans. Acad. Sci*. 65:318–364.

Barkley T.M. 1988. Variation among the Aureoid Senecios of North America: A
geohistorical interpretation. *Bot. Rev*. 54:82–106.

Benton M.J., Donoghue P.C.J. 2007. Paleontological evidence to date the tree of life.
Mol. Biol. Evol. 24:26–53.

Bentz P.C., Leebens-Mack J. 2024. Developing Asparagaceae1726: An Asparagaceae-
specific probe set targeting 1726 loci for Hyb-Seq and phylogenomics in the family.
Appl. Plant Sci. 12:e11597.

Betrán E. 2015. The "life histories" of genes. *J. Mol. Evol*. 80:186–188.

Bezanson J., Chen J., Chung B., Karpinski S., Shah V.B., Vitek J., Zoubritzky L. 2018.
Julia: dynamism and performance reconciled by design. *Proc. ACM Program. Lang*.
2:1–23.

Bolger A.M., Lohse M., Usadel B. 2014. Trimmomatic: A flexible trimmer for Illumina
sequence data. *Bioinformatics*. 30:2114–2120.

Braasch I., Postlethwait J.H. 2012. Polyploidy in Fish and the Teleost Genome
Duplication. In *Polyploidy and Genome Evolution*. Springer Berlin Heidelberg. p. 341–
383.

Brown J.W., Walker J.F., Smith S.A. 2017. Phyx: Phylogenetic tools for unix.
Bioinformatics. 33:1886–1888.

Crameri S., Fior S., Zoller S., Widmer A. 2022. A target capture approach for
phylogenomic analyses at multiple evolutionary timescales in rosewoods (*Dalbergia*
spp.) and the legume family (Fabaceae). *Mol. Ecol. Resour*. 22:3087–3105.

Danecek P., Bonfield J.K., Liddle J., Marshall J., Ohan V., Pollard M.O., Whitwham A.,
Keane T., McCarthy S.A., Davies R.M., Li H. 2021. Twelve years of SAMtools and
BCFtools. *Gigascience*. 10:giab008.

Degnan J.H., Rosenberg N.A. 2009. Gene tree discordance, phylogenetic inference and
the multispecies coalescent. *Trends Ecol. Evol*. 24:332–340.

Doyle J.J. 1992. Gene trees and species trees: Molecular systematics as one-character
taxonomy. *Syst. Bot*. 17:144.

Elven R., Murray D.F., Volodya Yu. Razzhivin B.A.Y. 2011. Annotated Checklist of the
Panarctic Flora (PAF) Vascular plants version 1.0. Available from
http://panarcticflora.org/results?biogeographic=&bioclimatic=®ion=&name=packera+
heterophylla#paf-863101 .

Evans B.J., Kelley D.B., Tinsley R.C., Melnick D.J., Cannatella D.C. 2004. A
mitochondrial DNA phylogeny of African clawed frogs: Phylogeography and implications
for polyploid evolution. *Mol. Phylogenet. Evol*. 33:197–213.

Faircloth B.C. 2016. PHYLUCE is a software package for the analysis of conserved
genomic loci. *Bioinformatics*. 32:786–788.

Freeman C.C., Barkley T.M. 1995. A synopsis of the genus *Packera* (Asteraceae:
Senecioneae) in Mexico. *SIDA Contrib. Bot*. 16:699–709.

Fu Z.X., Jiao B.H., Nie B., Zhang G.J., Gao T.G., Chen Z.D., Lu A.M., Kong H.Z., Wang
X.Q., Wang Y.Z., Zhou S.L., Zhang S.Z., Wang X.M., Liu Z.J., Wang Q.F., Li J.H., Li
D.Z., Yi T.S., Hong M.A., Soltis D.E., Soltis P.S., Li J.H., Fu C.X., Liu Q.X. 2016. A
comprehensive generic-level phylogeny of the sunflower family: Implications for the
systematics of Chinese Asteraceae. *J. Syst. Evol*. 54:416–437.

Funk V.A. 1985. Phylogenetic Patterns and Hybridization. *Ann. Mo. Bot. Gard*. 72:681.

Gallardo M.H., Kausel G., Jiménez A., Bacquet C., González C., Figueroa J., Köhler N.,
Ojeda, R. 2004. Whole-genome duplications in South American desert rodents
(Octodontidae). *Biol. J. Linn. Soc*. 82:443–451.

Galtier N., Daubin V. 2008. Dealing with incongruence in phylogenomic analyses.
Philos. Trans. R. Soc. Lond. B Biol. Sci. 363:4023–4029.

Geldmacher J., Van Den Bogaard P., Hoernle K., Schmincke H.U. 2000. The 40Ar/39Ar
age dating of the Madeira Archipelago and hotspot track (eastern North Atlantic).
Geochem. Geophys. Geosyst. 1.

Gramling A. 2006. A conservation assessment of *Packera millefolium*, a Southern
Appalachian Endemic [Dissertation]. The University of North Carolina at Chapel Hill.

https://www.proquest.com/dissertations-theses/conservation-assessment-i-packera-
millefolium/docview/305279534/se-2?accountid=30699.

Grant V. 1971. Plant Speciation. New York: Columbia University Press.

Hejase H.A., Liu K.J. 2016. A scalability study of phylogenetic network inference
methods using empirical datasets and simulations involving a single reticulation. *BMC*
Bioinf. 17:422.

Hobbs C.R., Baldwin B.G. 2013. Asian origin and upslope migration of Hawaiian
Artemisia (Compositae-Anthemideae). *J. Biogeogr*. 40:442–454.

Huang C.H., Zhang C., Liu M., Hu Y., Gao T., Qi J., Ma H. 2016. Multiple
Polyploidization Events across Asteraceae with Two Nested Events in the Early History
Revealed by Nuclear Phylogenomics. *Mol. Biol. Evol*. 33:2820–2835.

Jackson C., McLay T., Schmidt-Lebuhn A.N. 2023. hybpiper-nf and paragone-nf:
Containerization and additional options for target capture assembly and paralog
resolution. *Appl. Plant Sci*. 11:e11532.

Jin J.-J., Yu W.-B., Yang J.-B., Song Y., dePamphilis C.W., Yi T.-S., Li D.-Z. 2020.
GetOrganelle: a fast and versatile toolkit for accurate de novo assembly of organelle
genomes. *Genome Biol*. 21:241.

Johnson M.G., Gardner E.M., Liu Y., Medina R., Goffinet B., Shaw A.J., Zerega N.J.C.,
Wickett N.J. 2016. HybPiper: Extracting coding sequence and introns for phylogenetics
from high-throughput sequencing reads using target enrichment. *Appl. Plant Sci*.
4:1600016.

Jones K.E., Alfredo Reyes-Betancort J., Hiscock S.J., Carine M.A. 2014. Allopatric
diversification, multiple habitat shifts, and hybridization in the evolution of *Pericallis*
(Asteraceae), a Macaronesian endemic genus. *Am. J. Bot*. 101:637–651.

Katoh K., Standley D.M. 2013. MAFFT multiple sequence alignment software version 7:
Improvements in performance and usability. *Mol. Biol. Evol*. 30:772–780.

Kück P., Longo G.C. 2014. FASconCAT-G: Extensive functions for multiple sequence
alignment preparations concerning phylogenetic studies. *Front. Zool*. 11:1–8.

Kumar S., Stecher G., Li M., Knyaz C., Tamura K. 2018. MEGA X: Molecular
evolutionary genetics analysis across computing platforms. *Mol. Biol. Evol*. 35:1547–
1549.

Kumar S., Stecher G., Peterson D., Tamura K. 2012. MEGA-CC: Computing core of
molecular evolutionary genetics analysis program for automated and iterative data
analysis. *Bioinformatics*. 28:2685–2686.

Kumar S., Stecher G., Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics
Analysis Version 7.0 for Bigger Datasets. *Mol. Biol. Evol*. 33:1870–1874.

Landis M.J., Matzke N.J., Moore B.R., Huelsenbeck J.P. 2013. Bayesian analysis of
biogeography when the number of areas is large. *Syst. Biol*. 62:789–804.

Lanfear R., Calcott B., Ho S.Y.W., Guindon S. 2012. PartitionFinder: Combined
selection of partitioning schemes and substitution models for phylogenetic analyses.
Mol. Biol. Evol. 29:1695–1701.

Langmead B., Salzberg S.L. 2012. Fast gapped-read alignment with Bowtie 2. *Nat.*
Methods. 9:357–359.

Li H., Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics*. 25:1754–1760.

Linder C.R., Rieseberg L.H. 2004. Reconstructing patterns of reticulate evolution in
plants. *Am. J. Bot*. 91:1700–1708.

Maddison W.P. 1997. Gene trees in species trees. *Syst. Biol*. 46:523–536.

Magalhaes I.L.F., Santos A.J., Ramírez M.J. 2021. Incorporating Topological and Age
Uncertainty into Event-Based Biogeography of Sand Spiders Supports Paleo-Islands in
Galapagos and Ancient Connections among Neotropical Dry Forests. *Diversity*. 13:418.

Mahoney A.M., Kowal R.R. 2008. Three new varieties of *Packera paupercula*
(Asteraceae, Senecioneae) in midwestern and southeastern North America. *Novon: J.*
Bot. Nomencl. 18:220–228.

Mandel J.R., Dikow R.B., Funk V.A., Masalia R.R., Staton S.E., Kozik A., Michelmore
R.W., Rieseberg L.H., Burke J.M. 2014. A Target Enrichment Method for Gathering
Phylogenetic Information from Hundreds of Loci: An Example from the Compositae.
Appl. Plant Sci. 2:1300085.

Mandel J.R., Dikow R.B., Siniscalchi C.M., Thapa R., Watson L.E., Funk V.A. 2019. A
fully resolved backbone phylogeny reveals numerous dispersals and explosive
diversifications throughout the history of Asteraceae. *Proc. Natl. Acad. Sci. U.S.A*.
116:14083–14088.

Masterson J. 1994. Stomatal size in fossil plants: evidence for polyploidy in majority of
angiosperms. *Science*. 264:421–424.

Matzke N.J. 2014. Model selection in historical biogeography reveals that founder-event
speciation is a crucial process in island clades. *Syst. Biol*. 63:951–970.

McDougall I., Schmincke H.U. 1976. Geochronology of Gran Canaria, Canary Islands:
Age of shield building volcanism and other magmatic phases. *Bulletin Volcanologique*.
40:57–77.

Mello B. 2018. Estimating timetrees with MEGA and the timetree resource. *Mol. Biol.*
Evol. 35:2334–2342.

Mendes F.K., Hahn Y., Hahn M.W. 2016. Gene tree discordance can generate patterns

of diminishing convergence over time. *Mol. Biol. Evol*. 33:3299–3307.

Minh B.Q., Schmidt H.A., Chernomor O., Schrempf D., Woodhams M.D., Von Haeseler
A., Lanfear R., Teeling E. 2020. IQ-TREE 2: New Models and Efficient Methods for
Phylogenetic Inference in the Genomic Era. *Mol. Biol. Evol*. 37:1530–1534.

Mitchell N., Lewis P.O., Lemmon E.M., Lemmon A.R., Holsinger K.E. 2017. Anchored
phylogenomics improves the resolution of evolutionary relationships in the rapid
radiation of *Protea* L. *Am. J. Bot*. 104:102–115.

Moore-Pollard E.R., Jones D.S., Mandel J.R. 2024. Compositae-ParaLoss-1272: A
complementary sunflower-specific probe set reduces paralogs in phylogenomic
analyses of complex systems. *Appl. Plant Sci*. 12:e11568.

Morales-Briones D.F., Gehrke B., Huang C.-H., Liston A., Ma H., Marx H.E., Tank D.C.,
Yang Y. 2021a. Analysis of paralogs in target enrichment data pinpoints multiple ancient
polyploidy events in *Alchemilla* s.L. (Rosaceae). *Syst. Biol*. 71:190–207.

Morales-Briones D.F., Kadereit G., Tefarikis D.T., Moore M.J., Smith S.A., Brockington
S.F., Timoneda A., Yim W.C., Cushman J.C., Yang Y. 2021b. Disentangling sources of
gene tree discordance in phylogenomic data sets: Testing ancient hybridizations in
Amaranthaceae s.l. *Syst. Biol*. 70:219–235.

Morales-Briones D.F., Liston A., Tank D.C. 2018. Phylogenomic analyses reveal a deep
history of hybridization and polyploidy in the Neotropical genus *Lachemilla* (Rosaceae).
New Phytol. 218:1668–1684.

Müller J., Reisz R.R. 2005. Four well-constrained calibration points from the vertebrate
fossil record for molecular clock estimates. *BioEssays*. 27:1069–1075.

Nice C.C., Gompert Z., Fordyce J.A., Forister M.L., Lucas L.K., Buerkle C.A. 2013.
Hybrid speciation and independent evolution in lineages of alpine butterflies. *Evolution*.
67:1055–1068.

One Thousand Plant Transcriptomes Initiative. 2019. One thousand plant
transcriptomes and the phylogenomics of green plants. *Nature*. 574:679–685.

Ottenburghs J. 2020. Ghost introgression: Spooky gene flow in the distant past.
BioEssays. 42:e2000012.

Palazzesi L., Pellicer J., Barreda V.D., Loeuille B., Mandel J.R., Pokorny L., Siniscalchi
C.M., Tellería M.C., Leitch I.J., Hidalgo O. 2022. Asteraceae as a model system for
evolutionary studies: from fossils to genomes. *Bot. J. Linn. Soc*. 200:143–164.

Panchy N., Lehti-Shiu M., Shiu S.H. 2016. Evolution of gene duplication in plants. *Plant
Physiol*. 171:2294–2316.

Panero J.L. 2005. New combinations and infrafamilial taxa in the Asteraceae.
Phytologia. 87:1–14.

Paradis E., Schliep K. 2019. Ape 5.0: An environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics*. 35:526–528.

Parham J.F., Donoghue P.C.J., Bell C.J., Calway T.D., Head J.J., Holroyd P.A., Inoue
J.G., Irmis R.B., Joyce W.G., Ksepka D.T., Patané J.S.L., Smith N.D., Tarver J.E., Van
Tuinen M., Yang Z., Angielczyk K.D., Greenwood J.M., Hipsley C.A., Jacobs L.,
Makovicky P.J., Müller J., Smith K.T., Theodor J.M., Warnock R.C.M., Benton M.J.
2012. Best practices for justifying fossil calibrations. *Syst. Biol*. 61:346–359.

Pease J.B., Brown J.W., Walker J.F., Hinchliff C.E., Smith S.A. 2018. Quartet Sampling
distinguishes lack of support from conflicting support in the green plant tree of life. *Am.*
J. Bot. 105:385–403.

Peirson J.A., Reznicek A.A., Semple J.C. 2012. Polyploidy, infraspecific cytotype
variation, and speciation in Goldenrods: The cytogeography of *Solidago* subsect.
Humiles (Asteraceae) in North America. *Taxon*. 61:197–210.

Pelser P.B., Kennedy A.H., Tepe E.J., Shidler J.B., Nordenstam B., Kadereit J.W.,
Watson L.E. 2010. Patterns and causes of incongruence between plastid and nuclear
Senecioneae (Asteraceae) phylogenies. *Am. J. Bot*. 97:856–873.

R Core Team. 2016. R: A Language and Environment for Statistical Computing. Vienna:
R Foundation for Statistical Computing.

Ree R.H., Smith S.A. 2008. Maximum likelihood inference of geographic range
evolution by dispersal, local extinction, and cladogenesis. *Syst. Biol*. 57:4–14.

Revell L.J. 2012. phytools: An R package for phylogenetic comparative biology (and
other things). *Methods Ecol. Evol*. 3:217–223.

Ronquist F. 1997. Dispersal-vicariance analysis: A new approach to the quantification of
historical biogeography. *Syst. Biol*. 46:195–203.

Rose J.P., Toledo C.A.P., Moriarty Lemmon E., Lemmon A.R., Sytsma K.J. 2021. Out
of Sight, Out of Mind: Widespread Nuclear and Plastid-Nuclear Discordance in the
Flowering Plant Genus *Polemonium* (Polemoniaceae) Suggests Widespread Historical
Gene Flow Despite Limited Nuclear Signal. *Syst. Biol*. 70:162–180.

RStudio. 2020. RStudio: Integrated Development for R. Boston, MA: RStudio, PBC.

Salmon A., Udall J.A., Jeddeloh J.A., Wendel J. 2012. Targeted capture of
homoeologous coding and noncoding sequence in polyploid cotton. *G3*. 2:921–930.

Sayyari E., Mirarab S. 2016. Fast Coalescent-Based Computation of Local Branch
Support from Quartet Frequencies. *Mol. Biol. Evol*. 33:1654–1668.

Schliep K.P. 2011. phangorn: phylogenetic analysis in R. *Bioinformatics*. 27:592–593.

Shang H.-Y., Jia K.-H., Li N.-W., Zhou M.-J., Yang H., Tian X.-L., Ma Y.-P., Zhang R.-G.
2025. Phytop: a tool for visualizing and recognizing signals of incomplete lineage sorting

and hybridization using species trees output from ASTRAL. *Hortic. Res*. 12:uhae330.

Siniscalchi C.M., Hidalgo O., Palazzesi L., Pellicer J., Pokorny L., Maurin O., Leitch I.J.,
Forest, F., Baker W.J. Mandel J.R. 2021. Lineage-specific vs. universal: A comparison
of the Compositae1061 and Angiosperms353 enrichment panels in the sunflower family.
Appl. Plant Sci. 9.

Smith S.A., Moore M.J., Brown J.W., Yang Y. 2015. Analysis of phylogenomic datasets
reveals conflict, concordance, and gene duplications with examples from animals and
plants. *BMC Evol. Biol*. 15:150.

Solís-Lemus C., Ané C. 2016. Inferring Phylogenetic Networks with Maximum
Pseudolikelihood under Incomplete Lineage Sorting. *PLoS Genet*. 12:1–21.

Solís-Lemus C., Bastide P., Ané C. 2017. PhyloNetworks: A package for phylogenetic
networks. *Mol. Biol. Evol*. 34:3292–3298.

Soltis D.E., Soltis P.S. 1999. Polyploidy: Recurrent formation and genome evolution.
Trends Ecol. Evol. 14:348–352.

Soltis P.S., Soltis D.E. 2009. The role of hybridization in plant speciation. *Annu. Rev.*
Plant Biol. 60:561–588.

Stamatakis A. 2014. RAxML version 8: A tool for phylogenetic analysis and post-
analysis of large phylogenies. *Bioinformatics*. 30:1312–1313.

Steel M.A., Penny D. 1993. Distributions of tree comparison metrics—some new results.
Syst. Biol. 42:126–141.

Steenwyk J.L., Buida T.J., Labella A.L., Li Y., Shen X.-X., Rokas A. 2021. PhyKIT: a
broadly applicable UNIX shell toolkit for processing and analyzing phylogenomic data.
Bioinformatics. 37:2325–2331.

Stull G.W., Soltis P.S., Soltis D.E., Gitzendanner M.A., Smith S.A. 2020. Nuclear
phylogenomic analyses of asterids conflict with plastome trees and support novel
relationships among major lineages. *Am. J. Bot*. 107:790–805.

Takhtajan A. 1986. Floristic Regions of the World. Berkley and Los Angeles, California:
University of California Press.

Tamura K., Battistuzzi F.U., Billing-Ross P., Murillo O., Filipski A., Kumar S. 2012.
Estimating divergence times in large molecular phylogenies. *Proc. Natl. Acad. Sci.*
U.S.A. 109:19333–19338.

Tamura K., Tao Q., Kumar S. 2018. Theoretical foundation of the RelTime method for
estimating divergence times from variable evolutionary rates. *Mol. Biol. Evol*. 35:1770–
1782.

Thapa R., Bayer R.J., Mandel J.R. 2020. Phylogenomics Resolves the Relationships
within *Antennaria* (Asteraceae, Gnaphalieae) and Yields New Insights into its

Morphological Character Evolution and Biogeography. *Syst. Bot*. 45:387–402.

Tillich M., Lehwark P., Pellizzer T., Ulbricht-Jones E.S., Fischer A., Bock R., Greiner S.
2017. GeSeq–versatile and accurate annotation of organelle genomes. *Nucleic Acids*
Res. 45:W6–W11.

Tricou T., Tannier E., de Vienne D.M. 2022. Ghost lineages highly influence the
interpretation of introgression tests. *Syst. Biol*. 71:1147–1158.

Trock D.K. 1999. A revisionary synthesis of the genus *Packera* (Asteraceae:
Senecioneae) [Dissertation]. Kansas State University.
https://www.proquest.com/dissertations-theses/revisionary-synthesis-genus-packera-
asteraceae/docview/304521955/se-2?accountid=30699.

Trock D.K. 2006. *Packera* Á. Löve & D. Löve. Flora of North America Editorial
Committee, eds. 1993+. New York and Oxford: p. 570–602.

Vision T.J., Brown D.G., Tanksley S.D. 2000. The origins of genomic duplications in
Arabidopsis. *Science*. 290:2114–2117.

Wang W.M. 2004. On the origin and development of *Artemisia* (Asteraceae) in the
geological past. *Bot. J. Linn. Soc*. 145:331–336.

Wendel J.F., Cronn R.C. 2003. Polyploidy and the evolutionary history of cotton. *Adv.*
Agron. 23:139–186.

Yang Y., Smith S.A. 2014. Orthology inference in nonmodel organisms using
transcriptomes and low-coverage genomes: Improving accuracy and matrix occupancy
for phylogenomics. *Mol. Biol. Evol*. 31:3081–3092.

Yeatts L., Schneider A., Schneider B. 2011. *Packera mancosana* (Asteraceae:
Senecioneae), a new species and shale barren endemic of southwestern Colorado.
Phytoneuron. 26:1–8.

Yu G., Smith D.K., Zhu H., Guan Y., Lam T.T.-Y. 2017. GGTREE: An R package for
visualization and annotation of phylogenetic trees with their covariates and other
associated data. *Methods Ecol. Evol*. 8:28–36.

Zhang C., Huang C.-H., Liu M., Hu Y., Panero J.L., Luebert F., Gao T., Ma H. 2021.
Phylotranscriptomic insight into Asteraceae diversity, polyploidy, and morphological
innovation. *J. Integr. Plant Biol*. 63:1273–1293.

Zhang C., Mirarab S. 2022a. ASTRAL-Pro 2: ultrafast species tree reconstruction from
multi-copy gene family trees. *Bioinformatics*. 38:4949–4950.

Zhang C., Mirarab S. 2022b. Weighting by gene tree uncertainty improves accuracy of
quartet-based species trees. *Mol. Biol. Evol*. 39:msac215.

Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018. ASTRAL-III: Polynomial time species
tree reconstruction from partially resolved gene trees. *BMC Bioinformatics*. 19:15–30.

Zhang C., Scornavacca C., Molloy E.K., Mirarab S. 2020. ASTRAL-Pro: Quartet-based
species-tree inference despite paralogy. *Mol. Biol. Evol*. 37:3292–3307.