Mathematical modeling for a primitive form of habituation in an amoeba

Kota Nishi[1], Atsushi Tero[2*], Yukinori Nishigami[3],
Toshiyuki Nakagaki[3]

[1]Joint Graduate School of Mathematics for Innovation, Kyushu University, Fukuoka, Japan.
[2]Institute of Mathematics for Industry, Kyushu University, Fukuoka, Japan.
[3]Research Institute for Electronic Science, Hokkaido University, Hokkaido, Japan.

*Corresponding author. E-mail: tero@imi.kyushu-u.ac.jp

Abstract

Learning abilities, once thought to be unique to higher animals, have been reported to exist in their primitive form in single-celled organisms. This has triggered a growing interest in carefully examining the nature and mechanisms of the primitive versions of learning abilities, which would provide important clues for understanding the evolution of behavioral capabilities in organisms. In this study, we focused on previous experimental studies showing that the slime mold *Physarum polycephalum*, a model organism for studying protist behavior, exhibits the ability to adapt to chemical environments. We propose a possible dynamic mechanism underlying this habituation, reproducing reported experimental observations with accuracy. By refining a mathematical model that was as simple as possible and based on non-specific biochemical processes within cells, we clarified a plausible mechanistic framework. Based on these results, we examined the similarities and differences between this framework and previously proposed habituation models of single-cell movement and animal neural-circuit regulation. These findings are significant because they open new avenues for research into the generality and evolutionary origins of acclimation learning.

Keywords: True slime mold, Habituation, Ordinary differential equation, Reaction–diffusion process

1 Introduction

Studies on information processing in simpler organisms, such as single-celled protists that can evoke an adaptive response to complex environmental situations, have deepened our understanding of cognitive behavior in neural organisms (Lyon et al. 2021). Several types of adaptive behavior have been reported in the amoeboid organism *Physarum polycephalum*, making it an important model organism for elucidating a primitive form of cognitive function in higher animals (Reid 2023).

Plasmodium of *P. polycephalum* is an amoeba-like multinucleated unicellular organism with a large sheet-like body up to several m^2. Its protoplasm exists in two phases; protoplasmic gel (ectoplasm), which exhibits passive elasticity and active rhythmic contractility, and protoplasmic sol (endoplasm), which is a viscous liquid. Inside the sheet-like body of plasmodium, an intricate network of veins (tubular channels for protoplasmic streaming) develops toward the rear from the extending front.

The ectoplasmic gel periodically contracts based on the biochemical regulation of the actomyosin system, and these self-sustained oscillations of contraction show some phase and force differences within the body. As this contraction generates pressure on the sol, these differences cause it to flow within the body through the tube network. This sol flow causes amoeboid migration of plasmodium while inducing major morphological changes in the tube network.

Plasmodium can solve difficult foraging problems by reforming its efficient tube-network (Nakagaki et al. 2000; Tero et al. 2010), and can memorize and anticipate the environmental periodic changes as well (Saigusa et al. 2008). These adaptive behaviors and their underlying mechanisms have been analyzed using mathematical modeling (Nakagaki et al. 2007; Saigusa et al. 2008; Tero et al. 2007, 2010).

As this mathematical model elucidated the key mechanism underlying adaptive behavior, it was applicable to similar types of behavior in higher animals, such as the social design of public transportation networks in human society (Tero et al. 2010) and ant trail formation in complicated landscapes (Ma et al. 2013).

As discussed above, studies on *P. polycephalum* have provided a new perspective on the evolution of behavioral intelligence (right from cells to animal societies). Along this line of study, we aimed to investigate habituation in *P. polycephalum* and analyze the underlying mechanism by mathematical modeling.

Habituation, a type of learning, refers to the decrease in behavioral responses when stimuli are repeatedly presented (Rankin et al. 2009). Habituation is distinct from sensory adaptation, sensory fatigue, and motor fatigue (Rankin et al. 2009). Several studies have attempted to elucidate the mechanisms underlying habituation. Most habituation studies have focused on multicellular organisms that possess a nervous system (Castellucci et al. 1970; Johnson and Wuensch 1994).

However, some single-celled microorganisms, such as ciliates (Wood 1988a,b; Rajan et al. 2023) and amoebae (Boisseau et al. 2016), also exhibit habituation. Habituation to mechanical stimuli has been extensively demonstrated in ciliates (Dussutour 2021). For example, in the ciliate *Stentor*, the contraction response to mechanical stimulation is an example of habituation (Wood 1988a). In this habituation, changes in intracellular Ca^{2+} concentration are crucial (Dussutour 2021).

Fig. 1 Illustration depicting the workflow of the bridge crossing experiment conducted by Boisseau et al. (2016). The b-Q represents the agar bridge containing quinine, and the b-N represents the normal agar bridge

Fig. 2 Time taken by plasmodium to cross the bridge (from day 1 to 9) (Boisseau et al. 2016). Black point represents the mean time to cross the bridge, and the error bar represents the 95 percent confidence interval: the gray background represents the day when plasmodium crossed the b-Q, and the white background represents the day when plasmodium crossed the b-N. Data adapted from Boisseau et al. (2016)

In amoebae, habituation to chemical stimulation has been observed in plasmodium of *P. polycephalum*, as demonstrated by the hallmarks of habituation (Boisseau et al. 2016). However, the underlying mechanisms have not been adequately explored. Therefore, in this study, we propose a mathematical model to reproduce habituation in *P. polycephalum*. We compared this model with previous models and discussed the commonalities in the habituation mechanism.

2 Nature of Habituation in Plasmodium and Aim of Mathematical Modeling

Plasmodium usually locomotes when searching for food products. It decreases its speed when encountering quinine, a chemical repellent (Takagi et al. 2007). Boisseau et al. (2016) demonstrated the hallmarks of habituation in plasmodium by leveraging this property. They conducted bridge-crossing experiments on plasmodium (Fig. 1). In their experiment, a plasmodium was placed on a food patch, with a new food patch connected by one of two agar bridges: an agar bridge containing quinine (hereinafter, b-Q) or a normal agar bridge without quinine (hereinafter, b-N). The bridge was set between plasmodium and the new food patch, and plasmodium was then made to cross either the b-Q or the b-N once per day (Fig. 1). They recorded the time from when the tip of plasmodium entered the bridge to when it reached a new food patch (hereinafter

Table 1 The hallmarks of habituation (excerpt)

Name	Feature
habituation	Repeated application of a stimulus results in a progressive decrease in some parameter of a response to an asymptotic level.
spontaneous recovery	If the stimulus is withheld after response decrement, the response recovers at least partially over the observation time.

This table adapted from Rankin et al. (2009).

called the time to cross the bridge). From their experiments, the following observations were made: (i), the time taken to cross the b-Q decreased when plasmodium repeatedly crossed the b-Q (Fig. 2(a)), (ii), the time to cross the b-Q recovered after plasmodium crossed the b-N for 2 consecutive days (Fig. 2(a)), (iii), the time required to cross the b-N does not depend on the number of previous passes through the b-Q (Fig. 2 at day 7). In particular, features (i) and (ii) correspond to habituation and spontaneous recovery, the hallmarks of habituation (Table 1), respectively. Although feature (iii) is not directly related to habituation, feature (iii) suggests that plasmodium reflects the stimulus history of its motility only under necessary circumstances.

In this study, we developed a mathematical model to account for the three behavioral features (i) $\sim$ (iii). We formulated a locomotion model of plasmodium driven by a chemical reaction–diffusion process. In the next section, we describe how the physiological and behavioral properties of plasmodium are represented by a mathematical model.

3 Formulation of the Mathematical Model

3.1 Overview of Our Model

First, we present an overview of the formulated model. We developed the following model, which comprises the equation describing plasmodium locomotion (Eq. (1)) and the reaction–diffusion system regulating locomotion (Eq. (2) and (3)).

$$\frac{d\ell}{dt}(t) = \gamma(p_{\mathrm{r}}(t) - p_{\mathrm{f}}(t)), \tag{1}$$

$$\frac{dp_{\mathrm{f}}}{dt}(t) = d(p_{\mathrm{r}} - p_{\mathrm{f}}) + F_{\mathrm{f}}(p_{\mathrm{f}}), \tag{2}$$

$$\frac{dp_{\mathrm{r}}}{dt}(t) = r_{\mathrm{v}}d(p_{\mathrm{f}} - p_{\mathrm{r}}) + F_{\mathrm{r}}(p_{\mathrm{r}}). \tag{3}$$

where

$$F_{\mathrm{f}}(p_{\mathrm{f}}) = (1 - h(t))(-a_{\mathrm{de}}p_{\mathrm{f}} + g_{\mathrm{f}}) + h(t)(-a_{\mathrm{de}}^{\mathrm{s}}p_{\mathrm{f}} + g_{\mathrm{f}}^{\mathrm{s}}),$$
$$F_{\mathrm{r}}(p_{\mathrm{r}}) = -a_{\mathrm{de}}p_{\mathrm{r}} + g_{\mathrm{r}},$$

ℓ represents the position of plasmodium tip and p_f and p_r represent the sol pressures at the front and rear parts, respectively (Fig. 3). Function $h(t)$ equals 1 in the presence of stimulation and 0 in its absence. Plasmodium expands its tip through sol flow in the tube network. The sol flow is driven by the gradient of the sol pressure. Therefore, we assumed that the tip expansion velocity is proportional to the pressure gradient (Eq. (1)). In this study, plasmodium locomotion was restricted to a straight line. In addition, we regarded plasmodium as consisting of two parts: the front part, which is the extendable tip for locomotion, and the rear part, which is the section opposite to the tip (Fig. 3). Accordingly, we introduced the sol pressure variable at each part, p_f and p_r. Thus, $p_r - p_f$ represents the pressure gradient (Eq. (1)). The locomotion model (Eq. (1)) indicated that plasmodium migration is driven by the difference between the pushing force from the rear, which promotes forward movement, and the pushing-back force from the front, which inhibits it. By identifying the sol pressure with the concentration of the chemical involved, we formulated the dynamics of the sol pressure using a simplified reaction–diffusion system (Eq. (2) and (3)). The first term on the right-hand side is the diffusion term and the second term is the reaction term (Eq. (2) and (3)). Here, the parameters of the reaction term at the front switch in the presence or absence of stimulation because plasmodium detects the repellent at the front.

We developed a plasmodium locomotion model driven by a special difference in the chemical concentration that regulates the sol pressure. The stimulation directly influenced the chemical reaction at the front, whereas the reaction at the rear was indirectly affected by the diffusion between the front and rear. In the following section, we describe the model formation in more detail.

3.2 Detail of Formulation

First, we derived a locomotion model in plasmodium. We assumed that the tip expansion velocity was proportional to the volumetric flow rate of the sol toward its tip because plasmodium expanded its tip due to the sol flow in the tube network. Thus, we formulated the tip expansion velocity (hereafter called the locomotion velocity) as follows:

$$\frac{d\ell}{dt}(t) = k_\ell Q(t),$$

where $Q(t)$ denotes the volumetric flow rate of the sol. The sol flow inside the tube can be approximated using a Hagen-Poiseuille flow: The volumetric flow rate is expressed as follows:

$$Q = \frac{\pi R^4}{8\mu L}(p_1 - p_2),$$

where Q is the volumetric flow rate, R is the tube radius. L is the tube length, μ is the viscosity coefficient, and Here, p_1 and p_2 denote the pressures at the tube edges. In this study, we considered plasmodium to be composed of two parts: front and rear (Fig. 3). We assumed that n tubes cross between the front and rear parts and that

Fig. 3 Schematic illustration of a cross-section of plasmodium. Each of c_f and c_r represents a chemical that regulates the gel contraction. p_f and p_r represent the sol pressure. ℓ represents the coordinate of the tip position. A line with a circle represents an activation effect, a line with a T-shaped end represents an inhibition effect, an arrow represents the generation effect of a chemical, and a double-headed arrow represents the diffusion effect

they are identical. Therefore, the volumetric flow rate of the sol toward the front part is as follows:

$$Q(t) = n\frac{\pi R^4}{8\mu L}(p_r(t) - p_f(t)).$$

Consequently, the locomotion velocity of plasmodium is represented as

$$\frac{d\ell}{dt}(t) = \gamma(p_r(t) - p_f(t)),$$

where

$$\gamma = k_\ell n\frac{\pi R^4}{8\mu L}.$$

Next, we formulated a reaction–diffusion model for chemically involved gel contraction with simplified diffusion terms. We considered the chemical reactions, generation and decomposition, and the chemical diffusion between the front and rear parts. Then, the reaction–diffusion model was formulated as follows:

$$\frac{dc_f}{dt}(t) = d(c_r - c_f) + F_f(c_f), \tag{4}$$

$$\frac{dc_r}{dt}(t) = r_v d(c_f - c_r) + F_r(c_r), \tag{5}$$

where

$$F_f(c_f) = (1 - h(t))(-a_{de}c_f + g_f) + h(t)(-a_{de}^s c_f + g_f^s),$$
$$F_r(c_r) = -a_{de}c_r + g_r.$$

The variables c_f and c_r represent the concentrations of a chemical involved in gel contraction in the front and rear parts, respectively. The terms $d(c_r - c_f)$ and $r_v d(c_f - c_r)$ represent the chemical diffusion between the front and rear parts, respectively, where r_v is their volume ratio. Note that we assume diffusion between solutions with different

Fig. 4 Time course of our model (Eq. (1), (2), and (3)) where the bridge condition at day 1 is the b-Q, and that at day 2 is the b-N. (a) time course of sol pressure. The red line represents p_f, and the blue line represents p_r. (b) time course of the tip position of plasmodium. The value of ℓ resets to 0 at the start of the day. L_r represents the bridge length. TCB means the time to cross the bridge. Gray background represents the term under stimulation, $h(t) = 1$, and white background represents the term in the absence of stimulation, $h(t) = 0$

volumes, and therefore diffusion terms describe above (Appendix A). The terms $F_f(c_f)$ and $F_r(c_r)$ represent chemical reactions. $-a_{de}c_f$, $-a^s_{de}c_f$ and $-a_{de}c_r$ represent the decomposition of chemicals and g_f, g^s_f and g_r represent the generation of the chemicals. Function $h(t)$ equals 1 in the presence of stimulation and 0 in its absence. Thus, the chemical reaction at the front switches depending on its presence or absence.

Finally, we derived the temporal dynamics of the sol pressure. Sol pressure was generated by gel contraction. In this study, we assumed a linear relationship between sol pressure and the concentration of a chemical involved in gel contraction, although a more complex relationship may exist. From this assumption, the following equations were obtained:

$$p_f(t) = kc_f(t), p_r(t) = kc_r(t), \qquad (6)$$

where k denotes a positive constant. From Eq. (4), (5), and (6), the sol-pressure dynamics were derived as follows:

$$\frac{dp_f}{dt}(t) = d(p_r - p_f) + F_f(p_f),$$
$$\frac{dp_r}{dt}(t) = r_v d(p_f - p_r) + F_r(p_r),$$

where

$$F_f(p_f) = (1 - h(t))(-a_{de}p_f + g_f) + h(t)(-a^s_{de}p_f + g^s_f),$$
$$F_r(p_r) = -a_{de}p_r + g_r.$$

Here, kg_f, kg^s_f and kg_r replace g_f, g^s_f and g_r, respectively.

In conclusion, the model presented in Sec. 5.1. In the next section, we describe simulations performed using this model.

4 Simulation

First, we explain the simulation settings using a simple case study (Fig. 4). The purpose of our simulation was to compute the time required to cross the bridge daily, as in Boisseau et al. (2016). First, given the initial value of $(p_f(t), p_r(t))$ for each day, we compute the crossing time using Eq. (1), (2), and (3) (Fig. 4). We applied stimulation in the following pattern, depending on the bridge type. If the bridge type is the b-Q on day n $(t_{n-1} \leq t < t_n)$,

$$h(t) = \begin{cases} 1, & (t_{n-1} \leq t < t_{n-1} + T_s), \\ 0, & (t_{n-1} + T_s \leq t < t_n), \end{cases}$$

where $t_n = 24n$, $0 < T_s < 24$. If the bridge is the b-N on day n,

$$h(t) = 0, \ (t_{n-1} \leq t < t_n).$$

The initial value of $(p_f(t), p_r(t))$ on Day 1 is (P_f^u, P_r^u), which is the equilibrium state in the absence of stimulation. After Day 2, the final states of $(p_f(t), p_r(t))$ on the previous day were used as the initial conditions (Fig. 4(a)). Second, we computed the time taken to cross the bridge as the time required for $\ell(t)$, the position of plasmodium tips, to reach L_r, the length of the bridge, on each day (Fig. 4(b)).

Our simulation results reproduced three features (i) $\sim$ (iii) observed in the experimental results (Fig. 5). We focused on the temporal dynamics of the tip expansion velocity, corresponding to Fig. 5(a) (Fig. 6). Mitigation of deceleration due to repeated stimulation, followed by recovery of deceleration after a temporary interruption in stimulation, was observed (Fig. 6). In addition, the velocity immediately returned to $\gamma(P_r^u - P_f^u)$ when the stimulation was withheld (Fig. 6). The mitigation of deceleration under stimulation indicated a decline in the time to cross the b-Q, and its recovery indicated an increase in the time to cross the b-Q (Fig. 5(a)). In addition, the immediate return of velocity to $\gamma(P_r^u - P_f^u)$ corresponds to the consistency of the time required to cross the b-N (Fig. 5). In other words, these temporal behaviors enable the reproduction of three features (i) $\sim$ (iii). Therefore, in the subsequent section, we discuss the relationship between the sol pressures (p_f, p_r) and the velocity dynamics.

5 Mathematical Analysis

We analyzed the behavior of the locomotion velocity from the behavior of the sol pressure on the (p_f, p_r) plane. At this point, the dynamics of the sol pressure in the presence and absence of stimulation on the (p_f, p_r) plane and the relationship between the pressure and locomotion output are required. In Sec. 5.1, we analyze the dynamics of the pressure on the (p_f, p_r) plane and the relationship between the pressure and locomotion output. In Sec. 5.2 and 5.3, we discuss the behavior of the locomotion velocity based on the analysis in Sec. 5.1.

Fig. 5 Our simulation results of time to cross the bridge. The Red cross mark represents our simulation results, and the circle represents the mean time to cross the bridge in the experiment (Boisseau et al. 2016). Gray background represents the day when the bridge condition is quinine, and white background represents the day when the bridge condition is normal. (Parameters: $0 \leq t \leq 216$, $\Delta t = 1.0 \times 10^{-5}$, $L_\mathrm{r} = 1.3$, $T_\mathrm{s} = 8.0$, $\gamma = 2.3$, $a_\mathrm{de} = 3.33 \times 10^{-2}$, $g_\mathrm{f} = 1.17 \times 10^{-1}$, $g_\mathrm{r} = 4.08 \times 10^{-1}$, $a_\mathrm{de}^\mathrm{s} = 9.82 \times 10^{-1}$, $g_\mathrm{f}^\mathrm{s} = 11.2$, $d = 5.00 \times 10^{-1}$, $r_\mathrm{v} = 1.00 \times 10^{-1}$)

Fig. 6 The temporal dynamics of the tip expansion velocity corresponding to Fig. 5(a). Gray background represents the term under stimulation, $h(t) = 1$, white background represents the term in the absence of stimulation, $h(t) = 0$, and green broken line represents that the velocity is equal to $\gamma(P_\mathrm{r}^\mathrm{u} - P_\mathrm{f}^\mathrm{u})$

5.1 Mathematical Preparation

First, we analyzed the dynamics of the sol pressure. In the presence of a stimulation ($h(t) = 1$), Eq. (2) and (3) can be transformed as follows.

$$\frac{d}{dt}\begin{pmatrix} p_\mathrm{f} \\ p_\mathrm{r} \end{pmatrix} = \begin{pmatrix} -(a_\mathrm{de}^\mathrm{s} + d) & d \\ r_\mathrm{v}d & -(a_\mathrm{de} + r_\mathrm{v}d) \end{pmatrix}\begin{pmatrix} p_\mathrm{f} - P_\mathrm{f}^\mathrm{s} \\ p_\mathrm{r} - P_\mathrm{r}^\mathrm{s} \end{pmatrix}, \qquad (7)$$

where $(P_\mathrm{f}^\mathrm{s}, P_\mathrm{r}^\mathrm{s})$ denotes the equilibrium point.

$$P_\mathrm{f}^\mathrm{s} = \frac{(a_\mathrm{de} + r_\mathrm{v}d)g_\mathrm{f}^\mathrm{s} + dg_\mathrm{r}}{a_\mathrm{de}^\mathrm{s}(a_\mathrm{de} + r_\mathrm{v}d) + a_\mathrm{de}d}, P_\mathrm{r}^\mathrm{s} = \frac{r_\mathrm{v}dg_\mathrm{f}^\mathrm{s} + (a_\mathrm{de}^\mathrm{s} + d)g_\mathrm{r}}{a_\mathrm{de}^\mathrm{s}r_\mathrm{v}d + a_\mathrm{de}(a_\mathrm{de}^\mathrm{s} + d)}.$$

Fig. 7 (a) The solution trajectory of Eq. (7). The arrows at top-left represent the directions of eigenvector $\boldsymbol{v}_1$ and $\boldsymbol{v}_2$ (Eq. (8) and (9)). (b) The solution trajectory of Eq. (10). The arrows at top-left represent the directions of eigenvector $\boldsymbol{w}_1$ and $\boldsymbol{w}_2$ (Eq. (11) and (12)). The red broken line represents the straight line $p_r = p_f + P_r^u - P_f^u$. The white circle represents the equilibrium point in the absence of stimulation, (P_f^u, P_r^u), and the black circle represents the equilibrium point under stimulation, (P_f^s, P_r^s). The parameters are same at Fig. 5

The eigenvalues and eigenvectors in Eq. (7) are

$$\lambda_1 = \frac{-B - \sqrt{D}}{2}, \ \boldsymbol{v}_1 = \begin{pmatrix} \lambda_1 + a_{de} + r_v d \\ r_v d \end{pmatrix}, \tag{8}$$

$$\lambda_2 = \frac{-B + \sqrt{D}}{2}, \ \boldsymbol{v}_2 = \begin{pmatrix} d \\ \lambda_2 + a_{de}^s + d \end{pmatrix}, \tag{9}$$

where λ_i represents the eigenvalue, $\boldsymbol{v}_i$ represents the eigenvector corresponding to eigenvalue λ_i, and

$$B = a_{de}^s + a_{de} + d + r_v d,$$
$$D = (a_{de}^s - a_{de} + d - r_v d)^2 + 4 r_v d^2.$$

The dynamics of Eq. (7) depict in Fig. 7(a). In the absence of a stimulation ($h(t) = 0$), Eq. (2) and (3) can be modified as follows:

$$\frac{d}{dt} \begin{pmatrix} p_f \\ p_r \end{pmatrix} = \begin{pmatrix} -(a_{de} + d) & d \\ r_v d & -(a_{de} + r_v d) \end{pmatrix} \begin{pmatrix} p_f - P_f^u \\ p_r - P_r^u \end{pmatrix}, \tag{10}$$

where (P_f^u, P_r^u) denotes the equilibrium point.

$$P_f^u = \frac{(a_{de} + r_v d) g_f + d g_r}{a_{de}(a_{de} + r_v d + d)}, P_r^u = \frac{r_v d g_f + (a_{de} + d) g_r}{a_{de}(r_v d + a_{de} + d)}.$$

The eigenvalues and eigenvectors in Eq. (10) are

$$\mu_1 = -(a_{de} + d + r_v d), \ \boldsymbol{w}_1 = \begin{pmatrix} 1 \\ -r_v \end{pmatrix}, \tag{11}$$

Fig. 8 (a) The plot of $V(p_\mathrm{f}, p_\mathrm{r})$ on the $(p_\mathrm{f}, p_\mathrm{r})$ plane. Red broken lines represent the iso-velocity lines. The value on each line represents the velocity corresponding to the sol pressure $(p_\mathrm{f}, p_\mathrm{r})$ on the line. (b) The plot of $A_\mathrm{s}(p_\mathrm{f}, p_\mathrm{r})$ on the $(p_\mathrm{f}, p_\mathrm{r})$ plane. Blue dotted lines represent the iso-acceleration lines. The value on each line represents the acceleration under stimulation corresponding to the sol pressure $(p_\mathrm{f}, p_\mathrm{r})$ on the line. The white circle represents $(P_\mathrm{f}^\mathrm{u}, P_\mathrm{r}^\mathrm{u})$, and the black circle represents $(P_\mathrm{f}^\mathrm{s}, P_\mathrm{r}^\mathrm{s})$. The parameters are same at Fig. 5

$$\mu_2 = -a_\mathrm{de}, \;\; \boldsymbol{w}_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \tag{12}$$

where μ_i represents the eigenvalue, and $\boldsymbol{w}_i$ represents the eigenvector corresponding to the eigenvalue μ_i. The dynamics of Eq. (10) depict in Fig. 7(b).

Second, we analyzed the relationship between the sol pressure $(p_\mathrm{f}, p_\mathrm{r})$ and behavioral output on the $(p_\mathrm{f}, p_\mathrm{r})$ plane. From Eq. (1), we define the function $V(p_\mathrm{f}, p_\mathrm{r})$ that returns the locomotion velocity corresponding to the sol pressure $(p_\mathrm{f}, p_\mathrm{r})$.

$$V(p_\mathrm{f}, p_\mathrm{r}) = \gamma(p_\mathrm{r} - p_\mathrm{f}). \tag{13}$$

From Eq. (13), $V(p_\mathrm{f}, p_\mathrm{r})$ has a constant value on the line

$$p_\mathrm{r} = p_\mathrm{f} + (\mathrm{const}). \tag{14}$$

For example, $V(p_\mathrm{f}, p_\mathrm{r}) = 1$ for all $(p_\mathrm{f}, p_\mathrm{r})$ satisfying $p_\mathrm{r} = p_\mathrm{f} + 1/\gamma$. The lines in Eq. (14) is referred to as the iso-velocity line. The iso-velocity lines are shown in Fig. 8(a). Moreover, by differentiating both sides of Eq. (1), and substituting Eq. (7), we obtain the acceleration under stimulation as follows:

$$\frac{d^2\ell}{dt^2}(t) = -\gamma(a_\mathrm{de} + d + r_\mathrm{v}d)\left(p_\mathrm{r}(t) - \frac{a_\mathrm{de}^\mathrm{s} + d + r_\mathrm{v}d}{a_\mathrm{de} + d + r_\mathrm{v}d}p_\mathrm{f}(t)\right) - \gamma(g_\mathrm{f}^\mathrm{s} - g_\mathrm{r}).$$

We define the function $A_\mathrm{s}(p_\mathrm{f}, p_\mathrm{r})$, which returns the acceleration under stimulation for sol pressure $(p_\mathrm{f}, p_\mathrm{r})$.

$$A_\mathrm{s}(p_\mathrm{f}, p_\mathrm{r}) = -\gamma(a_\mathrm{de} + d + r_\mathrm{v}d)\left(p_\mathrm{r} - \frac{a_\mathrm{de}^\mathrm{s} + d + r_\mathrm{v}d}{a_\mathrm{de} + d + r_\mathrm{v}d}p_\mathrm{f}\right) - \gamma(g_\mathrm{f}^\mathrm{s} - g_\mathrm{r}). \tag{15}$$

Fig. 9 The sol pressure dynamics on the $(p_{\mathrm{f}}, p_{\mathrm{r}})$ plane corresponding to Fig. 5(a). (a) The trajectory of day 1∼2: the gray solid line represents the trajectory of day 1 (b-Q), and the black solid line represents the trajectory of day 2 (b-Q). (b) The trajectory of day 1∼8: the gray solid line represents the trajectory of day 1∼6 (b-Q), and the black solid line represents the trajectory of day 7∼8 (b-N). The white circle represents $(P_{\mathrm{f}}^{\mathrm{u}}, P_{\mathrm{r}}^{\mathrm{u}})$, the black circle represents $(P_{\mathrm{f}}^{\mathrm{s}}, P_{\mathrm{r}}^{\mathrm{s}})$, red broken lines represent the iso-velocity lines, blue dotted lines represent the iso-acceleration lines, and the red triangle with integer n points to the initial state at day n

From Eq. (15), $A_{\mathrm{s}}(p_{\mathrm{f}}, p_{\mathrm{r}})$ has a constant value on the line

$$p_{\mathrm{r}} = \frac{a_{\mathrm{de}}^{\mathrm{s}} + d + r_{\mathrm{v}}d}{a_{\mathrm{de}} + d + r_{\mathrm{v}}d} p_{\mathrm{f}} + (\text{const}). \tag{16}$$

The lines in Eq. (16) is referred to as the iso-acceleration line. The iso-acceleration lines are shown in Fig. 8(b). Because $a_{\mathrm{de}}^{\mathrm{s}} > a_{\mathrm{de}} > 0$ holds for our simulation parameters, the slope of the iso-acceleration line is greater than 1 (Eq. (16)). Hence, the iso-acceleration and iso-velocity lines intersect, as shown in Fig. 9.

5.2 Mitigation and Recovery of Deceleration

In this section, we discuss the mitigation and recovery of the deceleration under stimulation. The shift in the initial pressure state each day was crucial for the mitigation and recovery of deceleration under stimulation. By repeatedly crossing the b-Q, the initial state of each day shifts to the upper right (Fig. 9(a)). Focusing on the iso-acceleration lines, the initial state transitions in the direction in which the acceleration increases (Fig. 9(a)). It follows that the deceleration is mitigated (Fig. 6). However, the initial state shifted to the lower left each day when crossing the b-N (Fig. 9(b)). In other words, the initial state transitions in the opposite direction when repeatedly crossing the b-Q. Thus, deceleration under stimulation recovered when crossing the b-N. In summary, the mitigation of deceleration when repeatedly crossing the b-Q corresponded to an upper right shift of the initial state (Fig. 9(a)), and its recovery when crossing the b-N corresponded to a lower left shift (Fig. 9(b)).

Next, we discuss why the initial state shift occurs when the crossing the b-Q is repeated. To focus on the displacement in the direction along the iso-velocity line, namely, $(1,1)^t$, we introduce the following variable:

$$m(t) = p_{\mathrm{r}}(t) + p_{\mathrm{f}}(t).$$

The variable $m(t)$ represents the location in the direction of $(1,1)^t$ on the (p_f, p_r) plane. From the temporal change in $m(t)$, we can observe the temporal change in the coordinates on the axis parallel to the iso-velocity lines.

We focused on the pressure dynamics on day 1 when crossing the b-Q. The initial period of the day was during stimulation, followed by a period without stimulation. It follows that the pressure transitions toward the black point, (P_f^s, P_r^s), firstly, and then toward the white point, (P_f^u, P_r^u) (Fig. 10(a)). Considering the behavior of $m(t)$, the increase in $m(t)$ under stimulation exceeds its decrease in the absence of stimulation when $0 \le t < 24$ (Fig. 10(b)). Consequently, the initial state on Day 2 was positioned to the upper right of that on Day 1 (Fig. 10(a)). Next, we focus on the case where the crossing the b-Q is repeated. The increase in $m(t)$ under stimulation gradually balanced its decrease in the absence of stimulation by repeatedly crossing the b-Q (Fig. 10(b)). Consequently, the displacement of the initial state to the upper right decreased and the deceleration converged (Fig. 6(b)).

In summary, a shift in the initial state of each day is important for mitigation and recovery from deceleration. Additionally, we found that its mitigation caused a difference between the displacement to the axis parallel to the iso-velocity lines under stimulation and in its absence.

5.3 Velocity Consistency on the b-N

Here, we discuss the reason for the constant locomotion velocity of the b-N. The initial locomotion velocity on each day was almost the same as $\gamma(P_r^u - P_f^u)$ regardless of whether the bridge condition on the previous day was quinine or normal (Fig. 6(b)). In short, the initial states were sufficiently close to the straight line $p_r = p_f + P_r^u - P_f^u$, where the velocity was $\gamma(P_r^u - P_f^u)$ (Fig. 9). We now consider why the initial states are distributed in this manner.

The sol pressure $(p_f(t), p_r(t))$ first approaches a straight line $p_r = p_f + P_r^u - P_f^u$ in the absence of stimulation (Fig. 7(b)). The eigenvalues of Eq. (10) are negative, and the absolute value of the eigenvalue corresponding to the eigenvector $(1, -r_v)^t$ is larger than that corresponding to the eigenvector $(1, 1)^t$ due to the diffusion effect (Eq. (11) and (12)). Thus, the sol pressure decays faster toward $(1, -r_v)^t$ on the (p_f, p_r) plane. In addition, the sol pressure transitions along the straight line $p_r = p_f + P_r^u - P_f^u$ after approaching this line, because Eq. (10) has the eigenvector $(1, 1)^t$ (Fig. 7(b)).

When crossing the b-Q, stimulation was absent during the last period of the day and occurred before this period. Therefore, the sol pressure returns to a straight line $p_r = p_f + P_r^u - P_f^u$ although it leaves the line for some time when crossing the b-Q (Fig. 9(a) and 10(a)). When crossing the b-N, the sol pressure transitions along the straight line $p_r = p_f + P_r^u - P_f^u$ (Fig. 9(b)).

Consequently, the initial states are distributed along a straight line $p_r = p_f + P_r^u - P_f^u$ (Fig. 9). From this distribution, the sol pressure transitions along a straight line when crossing the b-N, independent of the bridge conditions of the previous day (Fig. 9(b)). Thus, the locomotion velocity remained constant (Fig. 6(b)).

Fig. 10 (a) The trajectory of day 1 on the $(p_\mathrm{f}, p_\mathrm{r})$ plane. The white circle represents $(P_\mathrm{f}^\mathrm{u}, P_\mathrm{r}^\mathrm{u})$, and the black circle represents $(P_\mathrm{f}^\mathrm{s}, P_\mathrm{r}^\mathrm{s})$. The red triangle with integer n points to the initial value at day n. (b) The temporal dynamics of $m(t)$. Gray background represents the period under stimulation, $h(t) = 1$, and white background represents the period without stimulation, $h(t) = 0$

6 Discussion

In the mathematical model proposed in this study, crawling migration (Eq. (1)) is regulated by the spatial difference in the protoplasmic pressure between the frontal and rear parts (Eq. (2) and (3)), and exhibits the two hallmarks defined in Table 1: habituation and spontaneous recovery.

In this section, we discuss the proposed model from a biological perspective. Deceleration mitigation as an adaptive behavior is realized by an initial state shift along the straight line $p_\mathrm{r} = p_\mathrm{f} + P_\mathrm{r}^\mathrm{u} - P_\mathrm{f}^\mathrm{u}$. Chemical diffusion and spatial differences in chemical decomposition rates are important for adaptive behavior.

The diffusion effect contributed to the transition of the initial state. Owing to diffusion effects, p_r increases with increasing p_f under stimulation. This, in turn, causes the displacement of the sol pressure $(p_\mathrm{f}, p_\mathrm{r})$ parallel to the iso-velocity lines to increase (Fig. 7(a) and 9(a)). In the absence of stimulation, the sol pressure $(p_\mathrm{f}, p_\mathrm{r})$ returns to a straight line $p_\mathrm{r} = p_\mathrm{f} + P_\mathrm{r}^\mathrm{u} - P_\mathrm{f}^\mathrm{u}$ (Sec. 5.3). Based on these factors, an initial state shift along the straight line $p_\mathrm{r} = p_\mathrm{f} + P_\mathrm{r}^\mathrm{u} - P_\mathrm{f}^\mathrm{u}$ is realized.

The spatial difference in the chemical decomposition rate contributes to the connection between the initial state shift and mitigation of deceleration. The relationship between the decomposition rates under stimulation in the front and rear parts determines the slope of the iso-acceleration line Eq. (16). In our simulation, $a_\mathrm{de}^\mathrm{s} > a_\mathrm{de}$ is true. Thus, the slope is greater than 1. However, the direction of the initial state shift is parallel to the iso-velocity line, whose slope is equal to 1 (Eq. (14)). Therefore, the direction of the initial state shift and the iso-velocity line intersect because $a_\mathrm{de}^\mathrm{s} > a_\mathrm{de}$ (Fig. 9(a)). Consequently, the deceleration was mitigated.

In summary, the diffusion effect between the front and rear parts is crucial for the transition of the initial state, and the difference in the chemical decomposition rate between the front and rear parts is important for reflecting the transition in the locomotion velocity. In short, the spatial scale of plasmodium played a key role in adaptive behavior in our model.

Some models exhibiting these two hallmarks (Table 1) have been proposed (Eckert et al. 2024; Stanley 1976). In a model based on the biochemical kinetics of cellular

Fig. 11 (a) The schematic diagram of IFF model in Eckert et al. (2024). (b) The schematic diagram of our model. r is equal to $-\dot{\ell}$. Line with a circle represents activation effect, line with a T-shaped end represents inhibition effect, arrow represents the generation effect of chemical and double-headed arrow represents diffusion effect

movement, two distinct chemical substances—activators and inhibitors—control the output (Eckert et al. 2024). In a model based on signal processing in neural circuits, the output is activated by the firing of other neurons and inhibited by a decline in the synaptic connection strength (Stanley 1976). In both models, the core mechanism controls the balance between two antagonistic factors: activation and inhibition of the response.

Although the biological realities represented by these models differ from those of our model, we considered whether there is any similarity in the core mechanism between our model and previously proposed models. In previous models, a common dynamic, which controls the balance between the inhibitory and activating effects on the response, was found.

In our model, the core mechanism lies in the relationship between the pressure difference of the protoplasm and the locomotion speed. In our simulation, the deceleration of locomotion speed under stimulation was decreased by repeated stimulation (Fig. 6(b)). Thus, we regarded deceleration during stimulation as a response to the stimulation. The variable p_{f} acts as an activator for deceleration under stimulation because a higher p_{f} decreases locomotion speed (Eq. (1)). In contrast, the variable p_{r} acts as an inhibitor of deceleration under stimulation because a higher p_{r} increases locomotion speed (Eq. (1)). In this sense of a dynamical system, our model has a mechanism similar to that of previous models.

However, the novel point worth noting in our model is that the two variables of the activator and inhibitor are not different chemical components, but the same component in different places in plasmodium (Fig. 3). Spatial differences in identical biochemical processes can exhibit two hallmarks of habituation. In other words, our model suggested that the spatial inhomogeneity of a chemical plays a crucial role in the habituation of spatially extending cells, such as *P. polycephalum*.

We compared our model with that of Eckert et al. (2024) based on an incoherent feed-forward (IFF). They reproduced the two features of habituation that our model does (Table 1) using the IFF model. In their model, two factors, I and M, act on the response R (Fig. 11(a)). The activation effect under stimulation from I to R was constant for each stimulation (top of Figure 2C and 3B in Eckert et al. (2024)). However, the inhibitory effect under stimulation from M to R depends on the series of stimulations. For example, the inhibitory effect gradually increases with each stimulation

(top of Figure 2C and 3B in Eckert et al. (2024)). This is because M increases with repeated stimulations. In contrast, the inhibition effect decreases when stimulation is withheld, owing to the decrease in M. In brief, the active effect on R was constant, but the inhibitory effect depended on the stimulation history. The IFF model realized two features of habituation that our model reproduced from the above properties.

In our model, p_f and p_r affect the locomotion velocity $\dot{\ell}$. As previously mentioned, we defined deceleration under stimulation as the response to stimulation. To make it easier to compare the IFF model with ours, we regard an increase in $r(t) = -\dot{\ell}(t)$ as the response. p_f is an activator of r and p_r is its inhibitor (Fig. 11(b)). p_f increased under stimulation. In addition, p_r increased under stimulation because of the diffusion effect (Fig. 7(a), 9(a), and 10(a)). For a more complex situation in which both the activator and inhibitor change, we analyzed the behavior of $(p_f(t), p_r(t))$ in the phase plane. The initial state shift was important for the decline and recovery of the response (Sec. 5.2 and 5.3). In our model, stimulation history was reflected in the initial state shift. To summarize, a common feature of the IFF model and our model is that they include a structure that stores the stimulation history. In the IFF model, variable $M(t)$ plays the role of memory. In our model, this is the initial state transition in the phase plane.

In the behavioral definition of habituation, there are ten hallmarks (Rankin et al. 2009). Some models have already been proposed to reproduce three or more hallmarks, but these models are more complicated than our model, as they involve more variables and incorporate nonlinearity in ODEs (Eckert et al. 2024; Smart et al. 2024), whereas our model essentially comprises two linear ODEs with switching parameters (Eq. (2) and (3)). For example, Eckert et al. (2024) reproduced other hallmarks by concatenating two IFF models. Our model was constructed in a bottom-up approach and it has the potential to be improved by adding new variables and/or nonlinearities, thereby reproducing three or more hallmarks.

In this study, we analyzed the behavior of $(p_f(t), p_r(t))$ on the phase plane to elucidate the fundamental structure of the two hallmarks: habituation and spontaneous recovery (Table 1). The analysis revealed that the initial state transition on an iso-velocity line is important for two hallmarks (Sec. 5.2 and 5.3). Our model provides an understanding of these two hallmarks from the perspective of a dynamic system. Our future research direction will be to analyze other hallmarks of habituation from the same perspective. This approach will bring new progress toward a comprehensive understanding of habituation mechanisms among a wide range of species.

A Derivation of the Diffusion Term

We derived the diffusion term for different solution volumes. We consider adjacent solutions 1 and 2 (Fig. 12). The volume of solution i was defined as V_i [m^3]. The chemical concentration of solution i is defined as c_i [mol/m^3]. J_i represents the flux [mol/m^2·s]. We derived the diffusion term when $V_1 \neq V_2$. In this study, we assumed that the flux from both ends is zero; that is, $J_0 = J_2 = 0$. From Fick's first law, J_1 can be expressed as follows:

$$J_1 = -D\frac{c_2 - c_1}{\Delta x},$$

Fig. 12 The problem setting of the diffusion between the different volumes of solutions. Arrow represents the flux

where D represents the diffusion coefficient $[\mathrm{m^2/s}]$, and Δx is the distance between solutions 1 and 2 $[\mathrm{m}]$. Note that the flux J_i is positive to the right. The inflow and outflow of solute particles per unit time in solution 1 are $S(J_0 - J_1)$ and where S denotes the particle passage. Thus, the following equation is obtained:

$$\frac{d}{dt}(V_1 c_1) = S(J_0 - J_1)$$
$$= \frac{SD}{\Delta x}(c_2 - c_1). \tag{17}$$

From Eq. (17), the differential equation for c_1 is derived as

$$\frac{d}{dt}c_1(t) = \frac{SD}{V_1 \Delta x}(c_2 - c_1).$$

Similarly, the differential equation for c_2 is derived as follows:

$$\frac{d}{dt}c_2(t) = \frac{SD}{V_2 \Delta x}(c_1 - c_2).$$

We now define d and r_v as follows:

$$d = \frac{SD}{V_2 \Delta x}, r_\mathrm{v} = \frac{V_2}{V_1}.$$

Ultimately, we obtain the differential equation with the diffusion term between different volumes of the solutions as follows:

$$\frac{d}{dt}c_1(t) = r_\mathrm{v} d(c_2 - c_1),$$
$$\frac{d}{dt}c_2(t) = d(c_1 - c_2).$$

Acknowledgements

K. N. was supported by the Doctoral Program for World-Leading Innovative & Smart Education (WISE program) from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT) at Kyushu University. This work was supported by the Cooperative Research Program of the "Network Joint Research Center for Materials and Devices" of MEXT, Japan.

Statements and Declarations

Funding

K. N. was supported by the WISE Program of MEXT at Kyushu University. This work was supported by the Cooperative Research Program of the "Network Joint Research Center for Materials and Devices" of MEXT, Japan.

Competing Interests

The authors declare no competing interests relevant to the contents of this article.

Authors' Contribution

A.T. proposed the fundamental ideas for the mathematical model, while K.N. performed the formulation, simulation, and analysis under A.T.'s supervision. T.N. and Y.N. examined the biological validity of the model. K.N. drafted the manuscript, which was revised by T.N. and Y.N. Throughout the entire research process, all authors collaborated and advanced the work through discussions under A.T.'s leadership.

Data Availability

The experimental data used in this study were originally generated by Boisseau et al. (2016) and are publicly available at https://doi.org/10.5061/dryad.51j89.

References

Boisseau RP, Vogel D, Dussutour A (2016) Habituation in non-neural organisms: evidence from slime moulds. Proc Biol Sci 283(1829):20160446. https://doi.org/10.1098/rspb.2016.0446

Castellucci V, Pinsker H, Kupfermann I, Kandel ER (1970) Neuronal mechanisms of habituation and dishabituation of the gill-withdrawal reflex in *Aplysia*. Science 167(3926):1745–1748. https://doi.org/10.1126/science.167.3926.1745

Dussutour A (2021) Learning in single cell organisms. Biochem Biophys Res Commun 564:92–102. https://doi.org/10.1016/j.bbrc.2021.02.018

Eckert L, Vidal-Saez MS, Zhao Z, Garcia-Ojalvo J, Martinez-Corral R, Gunawardena J (2024) Biochemically plausible models of habituation for single-cell learning. Curr Biol 34(24):5646–5658.e3. https://doi.org/10.1016/j.cub.2024.10.041

Johnson MC, Wuensch KL (1994) An investigation of habituation in the jellyfish *Aurelia aurita*. Behav Neural Biol 61(1):54–59. https://doi.org/10.1016/S0163-1047(05)80044-5

Lyon P, Keijzer F, Arendt D, Levin M (2021) Reframing cognition: getting down to biological basics. Philos Trans R Soc Lond B Biol Sci 376(1820):20190750. https://doi.org/10.1098/rstb.2019.0750

Ma Q, Johansson A, Tero A, Nakagaki T, Sumpter DJT (2013) Current-reinforced random walks for constructing transport networks. J R Soc Interface 10(80):20120864. https://doi.org/10.1098/rsif.2012.0864

Nakagaki T, Yamada H, Tóth A (2000) Maze-solving by an amoeboid organism. Nature 407(6803):470–470. https://doi.org/10.1038/35035159

Nakagaki T, Iima M, Ueda T, Nishiura Y, Saigusa T, Tero A, Kobayashi R, Showalter K (2007) Minimum-risk path finding by an adaptive amoebal network. Phys Rev Lett 99(6). https://doi.org/10.1103/PhysRevLett.99.068104

Rajan D, Makushok T, Kalish A, Acuna L, Bonville A, Correa Almanza KC, Garibay B, Tang E, Voss M, Lin A, Barlow K, Harrigan P, Slabodnick MM, Marshall WF (2023) Single-cell analysis of habituation in *Stentor coeruleus*. Curr Biol 33(2):241–251.e4. https://doi.org/10.1016/j.cub.2022.11.010

Rankin CH, Abrams T, Barry RJ, Bhatnagar S, Clayton DF, Colombo J, Coppola G, Geyer MA, Glanzman DL, Marsland S, McSweeney FK, Wilson DA, Wu CF, Thompson RF (2009) Habituation revisited: an updated and revised description of the behavioral characteristics of habituation. Neurobiol Learn Mem 92(2):135–138. https://doi.org/10.1016/j.nlm.2008.09.012

Reid CR (2023) Thoughts from the forest floor: a review of cognition in the slime mould physarum polycephalum. Anim Cogn 26(6):1783–1797. https://doi.org/10.1007/s10071-023-01782-1

Saigusa T, Tero A, Nakagaki T, Kuramoto Y (2008) Amoebae anticipate periodic events. Phys Rev Lett 100(1):018101. https://doi.org/10.1103/PhysRevLett.100.018101

Smart M, Shvartsman SY, Mönnigmann M (2024) Minimal motifs for habituating systems. Proc Natl Acad Sci U S A 121(41):e2409330121. https://doi.org/10.1073/pnas.2409330121

Stanley JC (1976) Computer simulation of a model of habituation. Nature 261(5556):146–148. https://doi.org/10.1038/261146a0

Takagi S, Nishiura Y, Nakagaki T, Ueda T, Ueda KI (2007) Indecisive behavior of amoeba crossing an environmental barrier. In: Topological aspects of critical systems and networks. World Scientific, pp 86–93, https://doi.org/10.1142/9789812708687_0011

Tero A, Kobayashi R, Nakagaki T (2007) A mathematical model for adaptive transport network in path finding by true slime mold. J Theor Biol 244(4):553–564. https://doi.org/10.1016/j.jtbi.2006.07.015

Tero A, Takagi S, Saigusa T, Ito K, Bebber DP, Fricker MD, Yumiki K, Kobayashi R, Nakagaki T (2010) Rules for biologically inspired adaptive network design. Science 327(5964):439–442. https://doi.org/10.1126/science.1177894

Wood DC (1988a) Habituation in *Stentor*: a response-dependent process. J Neurosci 8(7):2248–2253. https://doi.org/10.1523/JNEUROSCI.08-07-02248.1988

Wood DC (1988b) Habituation in *Stentor*: produced by mechanoreceptor channel modification. J Neurosci 8(7):2254–2258. https://doi.org/10.1523/JNEUROSCI.08-07-02254.1988