Body temperature drives azole tolerance in *Candida albicans* by hindering the autophagic degradation of Erg11

Yanru Feng[1], Cheng Zhen[1], Wanqian Li[1], Malcolm Whiteway[2], Xinyu Fang[1], Xuqing Shen[1], Hui Lu[1*], Yuanying Jiang[1*]

1. Department of Pharmacy, Shanghai Tenth People's Hospital, School of Medicine, Tongji University, Shanghai, China.

2. Department of Biology, Concordia University, Montreal, QC, Canada

*To whom correspondence should be addressed:

Hui Lu

Associate Senior Research Fellow in the Department of Pharmacology, Shanghai Tenth People's Hospital, School of Medicine, Tongji University

No.1239 Siping Road, Shanghai, 200092, China.

E-mail: luhui2019@tongji.edu.cn (HL)

Yuanying Jiang

Professor in the Department of Pharmacology, Shanghai Tenth People's Hospital, School of Medicine, Tongji University

No.1239 Siping Road, Shanghai, 200092, China.

E-mail: jiangyy@tongji.edu.cn (YYJ)

Abstract

Human body temperature has been shown to limit the antifungal efficacy of azoles; however, the precise mechanisms underlying this limitation remain insufficiently understood. In this study, we observed that brief exposure (24 h) to human body temperature (37 °C), in contrast to 30 °C, significantly enhances the tolerance of *Candida albicans* to azoles without inducing resistance. Our findings suggest that this increased tolerance is due to a reduction in the degradation of *C. albicans* Erg11 at 37 °C compared to 30 °C, thereby promoting azole tolerance. This phenomenon occurs because Erg11 degradation is mediated by autophagy, which is inhibited in *C. albicans* at 37 °C. The suppression of autophagy at this temperature may be associated with elevated mitochondrial production of reactive oxygen species, leading to mitochondrial dysfunction. In response to heat stress-induced mitochondrial dysfunction, autophagy-related proteins such as Atg8 accumulate around the mitochondria, while their levels in the endoplasmic reticulum decrease, consequently inhibiting autophagy in *C. albicans*. These findings suggest that promoting the degradation of Erg11 in *C. albicans* may serve as a viable strategy to augment the fungicidal efficacy of azoles in the human host.

Keywords: Erg11 degradation, azole tolerance, autophagy, heat stress, *Candida albicans*.

Introduction

In recent years, the morbidity and mortality rates associated with candidiasis have remained consistently high, presenting significant threats to public health and imposing a substantial medical burden[1]. Recent epidemiological data suggest that approximately 1.56 million individuals are affected by candidiasis annually, resulting in approximately 100,000 fatalities[2]. Azoles have emerged as the most frequently utilized antifungal agents for the clinical prevention and treatment of fungal infections, owing to their broad antimicrobial spectrum, favorable safety profile, and diverse routes of administration[3]. However, a primary challenge associated with azole therapy is their fungistatic nature, which allows *Candida albicans* to develop tolerance to azoles, ultimately leading to resistance[4]. The azole tolerance exhibited by *C. albicans* is evidenced by its ability to proliferate at azole concentrations surpassing the minimum inhibitory concentration (MIC). Although the prevalence of fluconazole (FLC) resistance in *C. albicans* is generally low, reported at less than 1%[5], FLC often proves ineffective in treating candidiasis caused by susceptible *C. albicans* isolates. The observed discrepancy between overall treatment outcomes and minimal levels of clinical resistance is likely attributable to multiple factors[6]. Nevertheless, FLC tolerance in *C. albicans* is a critical consideration, as both in vitro and in vivo antifungal efficacy of FLC is substantially reduced against *C. albicans* strains exhibiting high azole tolerance[7]. Furthermore, elevated levels of FLC tolerance are associated with persistent candidemia[8,9], which contributes to approximately 50% of the mortality associated with neonatal candidiasis[10]. However, the specific factors and mechanisms regulating the tolerance of *C. albicans* to azoles within the human host environment remain unclear.

It has been reported that the human body temperatures play a critical role in driving the development of drug resistance in pathogenic fungi. Studies have shown that, compared to a culture temperature of 30 °C, a temperature of 37 °C can decrease the susceptibility of *C. albicans* to azoles[11,12]. Research on the fungal pathogen *Rhodosporidiobolus fluvialis* suggests that exposure to mammalian body temperature can induce mutagenesis, leading to the emergence of hypervirulent and pan-drug-resistant strains[13]. Moreover, it has been demonstrated that anthropogenic global warming significantly

affects azole resistance[14,15], with increased occurrences of azole-resistant *Aspergillus*

fumigatus and *Candida parapsilosis* reported in warmer environments, such as

greenhouses[16] and tropical regions[17,18]. Additionally, the rapid global spread of

multidrug-resistant *Candida auris* in humans has been linked to temperature adaptation

driven by climate warming[19,20]. Body temperature may influence fungal drug

susceptibility through non-genetic mechanisms within the human host[21]. However, the

exact mechanisms by which short-term exposure to body temperature reduces fungal

susceptibility to azoles have yet to be elucidated. This gap in understanding poses a

challenge for developing targeted antifungal strategies aimed at enhancing the efficacy

of azoles in human hosts.

In this study, we observed that a short-term exposure (24 h) to the human body

temperature of 37 °C increases azole tolerance, but not azole resistance, in *C. albicans.*

Our investigations further revealed that Erg11 is subject to degradation via the

autophagy pathway. However, exposure to a body temperature of 37 °C, as opposed to

30 °C, inhibits autophagy, thereby decelerating the degradation rate of Erg11. This

process ultimately enhances the tolerance of *C. albicans* to azoles. These findings

expand the conceptual framework of azole tolerance mechanisms in *C. albicans*,

demonstrating that the human host body temperature can modulate the degradation rate

of the azole drug target Erg11 through epigenetic regulatory pathways.

Results

Short-term exposure to body temperature enhances the tolerance of *C. albicans* to

azoles

At a physiological body temperature of 37 °C, compared to 30 °C, we observed a

decrease in the susceptibility of the *C. albicans* wild-type strain SN152 to FLC (Figure

1a). However, it remains uncertain whether this decreased drug susceptibility is due to

an increase in the resistance level or tolerance level of *C. albicans* to FLC. In this study,

we characterized the enhancement of FLC resistance in *C. albicans* by assessing the

elevation of the minimum inhibitory concentration (MIC) value of FLC, while the

increase in the super-MIC growth (SMG) value was used to represent the elevation of

FLC tolerance in *C. albicans*[8,22](Figure 1b). Our findings indicate that, compared to

30 °C, the temperature of 37 °C significantly increased the SMG value of FLC against

the strain SN152 from 0.070 to 0.440, whereas the MIC value of FLC against the strain

SN152 remained constant at 0.5 μg/ml (Figure 1c). Cyclosporin A, a known calcineurin

inhibitor, was observed to abolish the tolerance of *C. albicans* to FLC by reducing the

SMG value of FLC, but not affect the resistance of *C. albicans* to FLC, as indicated by

the unchanged MIC value[8,22]. Our study demonstrated that the decreased susceptibility

of *C. albicans* to FLC after incubation at 37 °C for 24 h was reversed by cyclosporin A

(Extended Data Figure 1a). Furthermore, the deletion of the *CDR1* gene, which encodes

the efflux pump Cdr1, resulted in a reduced MIC value of FLC against *C. albicans*, yet

it did not eliminate FLC tolerance in the organism[8,23]. Our findings indicated that, in

comparison to 30 °C, culturing at 37 °C further diminished the susceptibility of the *C.*

albicans cdr1Δ/Δ null mutant to FLC (Extended Data Figure 1b). This observation is

attributed to the physiological temperature, which increased the SMG value of FLC

against the *cdr1Δ/Δ* null mutant, although it did not elevate the MIC value for this

mutant (Figure 1c). These results suggest that the reduced susceptibility of *C. albicans*

to FLC following short-term exposure (24 h) to physiological temperature is primarily

due to an enhanced tolerance, rather than the development of resistance of *C. albicans*

to azoles.

Our investigation revealed that elevating the culture temperature from 30 °C to 37 °C

significantly increased the SMG values of FLC against clinical isolates of *C. albicans*

($n = 104$), *Candida glabrata* ($n = 10$), *Candida parapsilosis* ($n = 10$), *Candida krusei*

($n = 10$), and *Candida guilliermondii* ($n = 10$) (Figure 1d). This observation suggests

that physiological body temperature generally enhances the tolerance of *Candida*

species isolates to FLC. Additionally, the SMG values for itraconazole, miconazole,

ketoconazole and voriconazole against *C. albicans* were significantly elevated when

cells were cultured at 37 °C (Figure 1e). Moreover, the antifungal efficacy of other

ergosterol biosynthesis inhibitors, such as terbinafine (targeting Erg1) and fluvastatin

(targeting Hmg1), were reduced at 37 °C (Figure 1f). However, the culture temperature

did not affect the antifungal activity of amphotericin B (binding and destroying plasma

ergosterol), caspofungin (targeting Fks1 to inhibit the synthesis of 1,3-β-D-glucan),

tunicamycin (inducing ER stress), myriocin (blocking serine-palmitoyltransferase in sphingolipid biosynthesis) and staurosporine (inhibiting protein kinase C) against *C. albican*s (Extended Data Figure 1c). Overall, these findings indicate that brief exposure to physiological body temperature substantially increases the tolerance of *C. albicans* to azoles.

The augmented tolerance of *C. albicans* to azoles at body temperature is dependent on Erg11

Upon brief exposure to body temperature, *C. albicans* exhibited reduced susceptibility to inhibitors targeting ergosterol biosynthesis, including azoles, terbinafine and fluvastatin (Figures 1e and 1f). Conversely, this exposure did not significantly impact the efficacy of inhibitors targeting non-ergosterol biosynthesis pathways, such as caspofungin, tunicamycin, myriocin, and staurosporine (Extended Data Figure 1c). Furthermore, our study demonstrated that both fluvastatin (1 µg/ml) and terbinafine (1 µg/ml) effectively decreased the SMG values of FLC against *C. albicans* at 37 °C (Figure 2a). These findings suggest that exposure to body temperature may influence the ergosterol biosynthesis pathway in *C. albicans*, thereby enhancing its tolerance to azoles.

We proceeded to delete the *EFG1*, *NDT80*, and *UPC2* genes in *C. albicans*, which encode transcriptional regulators involved in ergosterol biosynthesis[24-27]. Our findings demonstrated that at 37 °C, there was an increase in the SMG value of FLC against the *efg1*Δ/Δ and *ndt80*Δ/Δ null mutants (Extended Data Figure 2a). In contrast, the *upc2*Δ/Δ null mutant consistently exhibited low MIC and SMG values for FLC at both 30 °C and 37 °C (Figure 2b). This observation was further corroborated by spot assays, which confirmed that 37 °C did not reduce the susceptibility of the *upc2*Δ/Δ null mutant to FLC (Extended Data Figure 2b). These results suggest that Upc2 may contribute to the tolerance of *C. albicans* to azoles at physiological body temperature.

Body temperature may activate Upc2 transcription, thereby enhancing azole tolerance in *C. albicans*. We tagged the C-terminal of Upc2 with GFP, and subsequent laser confocal microscopy demonstrated that brief exposure to body temperature did not influence the nuclear import of Upc2 in *C. albicans* in the presence of FLC (16 µg/ml)

(Extended Data Figures 2c and 2d). Furthermore, RNA-seq analysis indicated that short-term exposure to 37 °C (for 4 h, 8 h, and 16 h) did not significantly alter the transcription levels of most ergosterol synthesis genes (Extended Data Figure 2e). Additionally, we tagged the C-terminal regions of key enzymes involved in ergosterol biosynthesis with GFP, including Erg6, Erg9, Erg10, Erg11, Erg12, Erg13, Erg20, Erg26, and Erg251. Consistent with the RNA-seq findings, no upregulation of these enzymes was observed at 37 °C compared to 30 °C. In fact, western blot analysis suggested a slight decrease in expression levels at 37 °C (Figures 2c and 2d). Furthermore, the analysis conducted using gas chromatography-mass spectrometry (GC-MS) indicated that there were no notable differences in intracellular ergosterol concentrations between the two temperature conditions at the 4-hour, 8-hour, and 16-hour incubation intervals (Extended Data Figure 2f). These results collectively imply that the enhanced tolerance of *C. albicans* to azoles at 37 °C is not mediated by the activation of Upc2 transcriptional activity.

In *C. albicans*, Erg7 (lanosterol synthase) catalyzes the conversion of squalene epoxide to lanosterol. This lanosterol is subsequently transformed into eburicol through the activity of Erg6 (Δ-(24)-sterol C-methyltransferase). Following this conversion, Erg11 mediates the oxidative demethylation of the 14-α-methyl group from eburicol, producing 4,4-dimethylcholesta-8,14,24-trienol[28]. Inhibition of Erg11 by azoles induces an alternative biosynthetic pathway involving Erg251 (C-4 sterol methyl oxidase), Erg26 (C-3 sterol dehydrogenase), Erg27 (3-keto sterol reductase), and Erg3 (C-5 sterol desaturase), which results in the synthesis of compensatory sterols that retain 14α-methyl groups[22] (Figure 2e). GC-MS analysis demonstrated a significant reduction in lanosterol levels, with no statistically significant changes in obtusifoliol concentrations, when cells were exposed to 16 μg/ml FLC at 37 °C compared to 30 °C (Figure 2f). Furthermore, utilizing the *ADH1* promoter to enhance the expression of ergosterol biosynthesis genes in the *upc2Δ/Δ* null mutant, specifically *ERG2*, *ERG4*, *ERG6, ERG9, ERG11, ERG24, ERG26*, and *ERG27*, showed that only the upregulation of *ERG11* significantly improved the relative growth of *C. albicans* under FLC exposure (0.25 μg/ml) at 37 °C (Figure 2g). These results are corroborated by chemical

genomics experiments, which demonstrated that fluvastatin, targeting Hmg1, and terbinafine, targeting Erg1 — both acting upstream of Erg11 — can inhibit azole tolerance induced by heat stress (Figure 2a). Conversely, amorolfine, which targets Erg24 and is positioned downstream of Erg11[29], did not show a similar inhibitory effect (Extended Data Figure 2g). Additionally, a notable increase in FLC tolerance was observed in the *erg24Δ/Δ* null mutant at 37 °C (Extended Data Figure 2h). The deletion of either the *ERG251* or *ERG3* gene[22], which are components of an alternative pathway, did not abolish the heat stress-induced tolerance to azoles (Extended Data Figure 2i). These findings suggest that body temperature augments the tolerance of *C. albicans* to azoles via a mechanism reliant on Erg11.

Body temperature inhibits the degradation of Erg11

We further modulated the expression of the *ERG11* and *ENO1* genes in the *ERG11/erg11Δ* and *ENO1/eno1Δ* heterozygous deletion mutants using the doxycycline (DOX)-inducible Tet-off promoter system (Extended Data Figure 3a). This system facilitates the regulation of the transcription levels of the *ERG11* and *ENO1* genes in response to varying DOX concentrations (Extended Data Figure 3b). Notably, the MIC values of DOX against the $P_{tet-off}$-*ENO1/eno1Δ* mutant remained consistent at both 30 °C and 37 °C. Conversely, the MIC value of DOX against the $P_{tet-off}$-*ERG11/erg11Δ* mutant at 37 °C was observed to be twice as high as at 30 °C (Figure 3a). These findings indicate that the increased tolerance of *C. albicans* to FLC at 37 °C cannot be attributed to the expression level of Erg11 at this temperature.

Moreover, the induction of azole tolerance by body temperature was determined to be contingent upon the culture medium. Specifically, an increased azole tolerance phenotype at 37 °C was observed in the nutrient-rich YPD medium, whereas this was not the case in the nutrient-poor SD medium (Figure 3b). These results imply that the heightened tolerance of *C. albicans* to FLC at 37 °C cannot be ascribed to an augmented catalytic activity of Erg11 at this temperature.

Cycloheximide (CHX) chase analysis was employed to assess the degradation rates of Erg11 at 30 °C and 37 °C. The analysis revealed that, unlike at 30 °C, the degradation of Erg11 was markedly inhibited at 37 °C (Figures 3c and 3d). At 37 °C, the inhibition

of Erg11 degradation leads to the sustained preservation of its catalytic activity. As a

result, *C. albicans* requires a reduced amount of Erg11 to produce an equivalent amount

of ergosterol at 37 °C compared to 30 °C. This induces a sterol-mediated negative

feedback mechanism that decreases Erg11 levels in *C. albicans* at 37 °C in the absence

of FLC (Figures 3e and 3f). However, when FLC inhibits Erg11 activity, the resulting

reduction in ergosterol content triggers an upregulation of Erg11 expression. Given that

37 °C suppresses Erg11 degradation, the exposure of azoles leads to elevated levels of

Erg11 protein in *C. albicans* at 37 °C relative to 30 °C (Figures 3e and 3f), thereby

enhancing the azole tolerance of *C. albicans*.

Erg11 undergoes degradation through autophagy

In *S. cerevisiae,* it has been documented that Erg11 is subject to degradation via the

Asi-mediated Endoplasmic Reticulum-Associated Degradation (ERAD) pathway

following ubiquitination[30-32]. However, the precise mechanism by which ubiquitinated

Erg11 is degraded in *C. albicans*—whether through the 26S proteasome or autophagy

in *C. albicans*—remains unclear. To investigate the degradation pathway of Erg11 in

C. albicans, we employed wortmannin, an autophagy inhibitor[33], and MG132, a

proteasome inhibitor[34], to ascertain their effects on Erg11 degradation. Our results

demonstrated that wortmannin significantly enhanced the FLC tolerance of *C. albicans*

at 30 °C and further augmented FLC tolerance at 37 °C. Conversely, MG132

diminished FLC resistance at both 30 °C and 37 °C, and reduced FLC tolerance at 37 °C

(Figure 4a). Moreover, CHX chase analysis revealed that wortmannin, but not MG132,

inhibited the degradation of Erg11 at 30 °C (Figures 4b and 4c). We employed the

CellTracker Blue CMAC dye to stain vacuoles and tagged Erg11 with GFP. Laser

confocal microscopy experiments revealed that Erg11 was localized within vacuoles,

as indicated by the co-localization of green and red fluorescence, resulting in a yellow

fluorescence signal (Figures 4d and 4e). Furthermore, our findings revealed that

incubation at 37 °C, as opposed to 30 °C, significantly hindered the entry of Erg11 into

vacuoles (Figures 4d and 4e). These results collectively suggest that Erg11 undergoes

degradation via autophagy, and that this process may be inhibited at 37 °C, thereby

suppressing the degradation of Erg11.

Body temperature inhibits autophagy in *C. albicans*

Considering that body temperature inhibits the degradation of Erg11, through the autophagy pathway, it can be posited that body temperature may suppress the autophagic process in *C. albicans*. To investigate this hypothesis, we employed a *C. albicans* GFP-Atg8 mutant, in which GFP was fused to the N-terminus of Atg8[35]. During the activation of autophagy, GFP-Atg8 localizes to autophagosomes and is subsequently released into vacuoles. Notably, GFP exhibits greater resistance to vacuolar hydrolases, leading to the degradation of Atg8 while GFP remains intact. Thus, the conversion of GFP-Atg8 to GFP serves as an indicator of autophagic activity. Our findings indicated that at 37 °C, the ratio of GFP to GFP-Atg8 was significantly lower compared to that observed at 30 °C (Figures 5a and 5b). Additionally, laser confocal microscopy demonstrated that at 30 °C, *C. albicans* cells exhibited a higher number of cells with green fluorescent dots, which are indicative of autophagosomes containing GFP-Atg8; conversely, at 37 °C, the number of such cells was markedly diminished (Figures 5c and 5d). In addition, we utilized MitoTracker Deep Red FM dye for mitochondrial labeling, ER-Tracker Red dye for ER labeling, and GFP for Atg8 labeling, employing confocal laser scanning microscopy to examine the effects of elevated environmental temperature on the intracellular localization of Atg8. Our findings indicate that, relative to the condition at 30 °C, the proximity of Atg8 to mitochondria increased at 37 °C (Figures 5e and 5f), whereas its colocalization with the ER diminished (Figures 5g and 5h), consequently resulting in the inhibition of autophagy.

Body temperature-induced mitochondrial oxidative damage that inhibits autophagy

Research has demonstrated that heat stress induces mitochondrial dysfunction via oxidative stress mechanisms[36]. In this study, we cultured *C. albicans* at 30 °C and 37 °C for 4 h, followed by RNA sequencing analysis. The results revealed that genes differentially expressed and associated with mitochondrial function were the most enriched, suggesting that exposure to 37 °C may impair mitochondrial function (Figure 6a). To further investigate, we employed 2′,7′-dichlorofluorescin diacetate (DCFH-DA)

to quantify reactive oxygen species (ROS) levels and observed that the 37 °C condition,

compared to 30 °C, resulted in elevated ROS levels in *C. albicans* (Figure 6b).

Furthermore, the antioxidant N-acetyl cysteine (NAC) was shown to effectively reduce

ROS levels and synergize with FLC to mitigate the tolerance of *C. albicans* at 37 °C

(Figure 6c). These findings collectively suggest that mitochondrial oxidative damage

in *C. albicans* induced by body temperature may contribute to the increased tolerance

of *C. albicans* to azoles.

Under conditions of heat stress, Atg8 in plants has been documented to engage

biological processes beyond autophagy, including its relocation to the Golgi apparatus

to facilitate Golgi reorganization[37]. Consequently, we hypothesized that the

redistribution of Atg8 may provide protection to mitochondria against oxidative

damage induced by body temperatures. To test this hypothesis, we generated an

atg8Δ/Δ null mutant in *C. albicans* and assessed the growth rates of both the *atg8Δ/Δ*

null mutant and its parental strain SN152 at 30 °C and 37 °C, respectively. Our findings

indicated that the *atg8Δ/Δ* null mutant exhibited growth rates comparable to those of

the SN152 strain at 30 °C. However, at 37 °C, the *atg8Δ/Δ* null mutant demonstrated

reduced growth relative to the SN152 strain (Figure 6d), suggesting that Atg8 plays a

critical role in the response to oxidative stress at 37 °C. Additionally, we employed JC-

1 to evaluate the impact of the *ATG8* gene on mitochondrial membrane potential

(MMP). The lipophilic JC-1 dye, with a delocalized positive charge, accumulates in

functional mitochondria, emitting red fluorescence. In dysfunctional mitochondria, it

remains dispersed in the cytosol, emitting green fluorescence. A decrease in red and

increase in green fluorescence signals mitochondrial dysfunction. Mitochondria with a

red-to-green ratio over 1 are considered normal (P1 in Figure 6e), while those below 1

are dysfunctional (P2 in Figure 6e). Our findings indicated that at 30 °C, the SN152

strain (0.290% ± 0.096%) and the *atg8Δ/Δ* null mutant (0.730% ± 0.131%) exhibit

comparable proportions of cells in the P2 region ($P > 0.999$). In contrast, at 37 °C, the

atg8Δ/Δ null mutant (8.42% ± 0.800%) shows a significantly higher proportion of cells

in the P2 region compared to the SN152 strain (5.33% ± 1.19%; $P = 1.07 \times 10^{-2}$)

(Figures 6e and 6f). These results suggest that mitochondrial dysfunction in *C. albicans*,

triggered by elevated temperatures, may lead to a reduced presence of autophagy-

related proteins, such as Atg8, within the ER, thereby inhibiting autophagy.

Discussion

In this study, we observed that short-term exposure of *C. albicans* to the human body

temperature of 37 °C, as opposed to 30 °C, results in increased azole tolerance, but not

resistance. This phenomenon is primarily attributed to the reduced degradation rate of

Erg11 at 37 °C compared to 30 °C. The slower degradation of Erg11 at 37 °C is due to

its degradation via autophagy, which is inhibited under this condition in *C. albicans*.

The inhibition of autophagy at 37 °C may be mediated by elevated mitochondrial

production of ROS, leading to mitochondrial dysfunction. In response to heat stress-

induced mitochondrial dysfunction, autophagy-related proteins such as Atg8

accumulate around the mitochondria, while their presence in the ER diminishes, thereby

suppressing autophagy in *C. albicans* (Figure 7).

Our investigation revealed that brief exposure of *C. albicans* to human body

temperature markedly increases the fungus's tolerance to azoles. This phenomenon may

significantly contribute to the ineffectiveness of azole treatments in infections caused

by non-resistant *C. albicans*. While the development of azole resistance in *C. albicans*

is a well-established cause of therapeutic failure, azole therapies may also prove

ineffective against infections involving non-resistant strains[4,38]. This ineffectiveness is

largely due to the intrinsic tolerance of *C. albicans* to azoles[4,12,38]. Previous studies have

confirmed that both in vitro and in vivo antifungal efficacy of FLC is substantially

reduced against *C. albicans* strains exhibiting high azole tolerance[7]. Research has

further shown that this tolerance is a pivotal factor in the persistence of candidemia[8,9].

Importantly, among patients who received their initial dose of FLC within 24 h of

candidemia onset, the probability of treatment failure was significantly higher when the

infecting strain demonstrated elevated tolerance to FLC[9]. This study illustrates that

even brief exposure to body temperature conditions can markedly increase the azole

tolerance of non-resistant *C. albicans*, potentially resulting in the ineffectiveness of

azole therapy for candidemia. These findings suggest that in the development of azole

adjuvants to augment antifungal efficacy against *C. albicans*, research should prioritize

evaluating whether these compounds facilitate the fungicidal action of azoles against

C. albicans[4,39,40].

Erg11 is integral to the biosynthesis of ergosterol and serves as a primary target for

azoles[41,42]. While the implications of Erg11 mutations and overexpression have been

extensively explored, its stability is equally crucial for maintaining fungal susceptibility

to azoles[43-45]. Current evidence suggests that Erg11 is processed via the Asi-mediated

Endoplasmic Reticulum-Associated Degradation (ERAD) pathway, which involves

ubiquitin-conjugating enzymes Ubc4 and Ubc7, the Asi1/Asi2/Asi3 ubiquitin ligase

complexes, and Cdc48-facilitated extraction, which culminates in Erg11 ubiquitination

and subsequent degradation in *S. cerevisiae*[30-32]. However, it remains uncertain whether

ubiquitinated Erg11 is degraded by the 26S proteasome—a pathway commonly

associated with many ERAD substrates[46]—or through autophagy, which is typical for

membrane proteins[47]. In this study, we demonstrated that the degradation of Erg11 is

inhibited by the autophagy inhibitor wortmannin, whereas the 26S proteasome inhibitor

MG132 did not significantly affect Erg11 degradation. Additionally, we observed that

Erg11 can translocate to the vacuole, suggesting that Erg11 in *C. albicans* is degraded

via autophagy. Our investigation further revealed that brief exposure to a body

temperature environment decelerates the degradation rate of Erg11 in *C. albicans* by

inhibiting the autophagy process, thereby affecting azole drug tolerance. This finding

expands the cognitive framework of azole tolerance mechanisms in *C. albicans*,

highlighting that host temperature can dynamically regulate the abundance of the target

enzyme Erg11 through epigenetic regulatory pathways.

In conclusion, this study demonstrates that under physiological temperature conditions,

the autophagy process in *C. albicans* is inhibited, leading to a reduced degradation rate

of Erg11. This inhibition consequently enhances the tolerance of *C. albicans* to azoles.

These findings suggest that promoting the degradation of Erg11 in *C. albicans* may

serve as a viable strategy to augment the fungicidal efficacy of azoles in the human

host.

Methods

Plasmids, strains, primers, agents, and cultural conditions

All plasmids, strains, and primers utilized in this study are detailed in Supplementary Tables 1, 2, and 3, respectively. The *C. albicans* strains were cultured in YPD medium, comprising 1% (w/v) yeast extract (Oxoid Ltd, Basingstoke, UK), 2% (w/v) peptone (Oxoid Ltd, Basingstoke, UK), and 2% (w/v) glucose (Sangon Biotech, Shanghai, China), at temperatures of either 30 °C or 37 °C. For the spotting assay, YPD medium plates were supplemented with 2% (w/v) agar (Sangon Biotech, Shanghai, China) and various concentrations of compounds. To construct mutant *C. albicans* strains, SD selective medium was employed, consisting of 0.67% (w/v) yeast nitrogen base without amino acids (Sangon Biotech, Shanghai, China), 2% (w/v) glucose, 2% (w/v) agar, and an appropriate amino acid mix (Sangon Biotech, Shanghai, China). FLC, itraconazole, miconazole, ketoconazole, voriconazole, fluvastatin, terbinafine, caspofungin, and amphotericin B were procured from Aladdin, Shanghai, China. Myriocin, staurosporine, brefeldin A, and amorolfine were obtained from MedChemExpress, Shanghai, China. Tunicamycin was sourced from TargetMol, Boston, USA. All these compounds were dissolved in dimethyl sulfoxide (DMSO) supplied by Sangon Biotech, Shanghai, China, and stored at -20 °C.

Spotting assays

The spotting assay was performed as described previously[7]. FLC was dissolved in DMSO, added to YPD medium, and used to prepare solid plates. Cultures of *C. albicans* grown overnight were standardized to a concentration of 1×10^7 cells/ml in YPD. These cultures were subsequently subjected to a series of 1:10 serial dilutions in YPD to five concentration gradients. The plates were incubated at either 30 °C or 37 °C for a duration of 48 h. Post-incubation, images of the plates were captured using the Tanon 5200 Imaging System (Tanon, Shanghai, China) in conjunction with AllDoc_X software.

In vitro antifungal susceptibility testing

To examine the in vitro susceptibility to five azoles (FLC, itraconazole, miconazole, ketoconazole, and voriconazole), other ergosterol biosynthesis inhibitors (terbinafine

and fluvastatin) and non-ergosterol biosynthesis antifungals (amphotericin B, caspofungin, tunicamycin, myriocin, and staurosporine), the broth microdilution assay was conducted according to previously reported protocol with slight modifications[48]. Briefly, 100 μl of drugs at 2-fold the final concentration was serially diluted in flat-bottom 96-well plates and combined with 100 μl of overnight *C. albicans* cultures adjusted to 1×10^3 cells/ml. The plates were incubated separately at 30 °C and 37 °C without shaking, and the optical density at 600 nm (OD_{600}) was measured with a spectrophotometer, Multiskan SkyHigh (Thermo Scientific, Singapore) after 24 h. The MIC value was determined as the drug concentration at which 50% of the growth was inhibited in terms of OD_{600} values compared to the no-drug wells.

To compare the values of SMG of FLC against different *Candida* isolates (*C. albicans*, *C. parapsilosis*, *C. glabrata*, *C. krusei*, and *C. guilliermondii*), the *C. albicans* wild-type strain SN152 and different mutants (*cdr1Δ/Δ*, *efg1Δ/Δ*, *ndt80Δ/Δ*, *upc2Δ/Δ*, *erg24Δ/Δ*, and *erg251Δ/Δ*) between 30 °C and 37 °C, based on the broth microdilution assay, the SMG value at 30 °C was calculated as the average growth per well above the MIC divided by the level of growth without the drug, and the SMG at 37 °C was calculated using the same MIC at 30 °C. The specific formula for calculating the value of the SMG is as follows:

$$SMG = \frac{\text{Average 24 h growth above MIC}}{\text{Growth without FLC}}$$

To examine whether one drug (cyclosporin A, terbinafine, fluvastatin, amorolfine, wortmannin, MG132, and NAC) combined with FLC can clear trailing in *C. albicans* at 30 °C and 37 °C, the checkerboard assays were used as described previously[49]. 50 μl of 4-fold the final drug concentration of drug A was dispensed in 2-fold serial dilution steps across plate columns, and then, 50 μl of 4-fold the final drug concentration of drug B was dispensed in 2-fold serial dilution steps down rows of the plate. 100 μl of overnight *C. albicans* cultures adjusted to 1×10^3 cells/ml was dispensed in all drug-containing wells plus one control well containing no drugs. Cell concentrations were recorded as the absorbance at OD_{600} after incubation at 30 °C or 37 °C for 24 h.

The homozygous null mutants constructions

To construct homozygous null mutants (*ndt80Δ/Δ* and *efg1Δ/Δ*) in this study, *C.*

albicans wild-type strain SN152 was used to permit disruption of both alleles of a given

target gene with two auxotrophic markers; In this experiment, the first round of gene

disruption was carried out with one of the three selectable markers, usually *Candida*

dubliniensis (*C. dubliniensis*) *HIS1*, and the second round was carried out with one of

the two remaining markers, usually *C. dubliniensis ARG4*. After selection of

transformants on the SD medium without histidine (SD/-His) or without histidine and

arginine (SD/-His/-Arg), gene disruption candidates were screened by PCR for

expected 5' and 3' junctions as well as the size of the disrupted gene[50].

The doxycycline-inducible Tet-off promoter of genes mutants constructions

To construct the doxycycline-inducible Tet-off promoter of *ERG11* or *ENO1* mutants,

first the *ERG11* and *ENO1* gene on one allele were deleted separately like what we

described above to generate *ERG11/erg11Δ* and *ENO1/eno1Δ*. Then the Tet-off vectors,

pCPC44 contained an ADH1 promoter, an ACT1 terminator and a tetracycline

repressor (TetR) element between them. TetR element was codon-optimized by

chemical synthesis and fused to GAL4AD to generate the fusion gene, *cat*TA. A

tetracycline-responsive elements (P_{Tet}) which was introduced into the downstream of

selection markers (*C. dubliniensis ARG4*). Through two rounds of PCR amplification

of pCPC44, the DNA cassettes with 78 bp homology regions to the target gene ($P_{tet-off}$-

ERG11 and $P_{tet-off}$-*ENO1*) could be transformed into *ERG11/erg11Δ* and *ENO1/eno1Δ*

separately to generate the strain with the Tet-off promoter-controlled target gene. After

selection of transformants on the SD medium without arginine (SD/-Arg), verification

primers were used to verify the successful construction of the strain with the Tet-off

promoter-controlled *ERG11* or *ENO1* ($P_{tet-off}$-*ERG11/erg11Δ* and $P_{tet-off}$-

ENO1/eno1Δ)[51].

The ectopic overexpression gene mutants constructions

To construct the ectopic overexpression gene mutants, the ectopic overexpression

vectors, plasmids pCPC18-*ERG2*, pCPC18-*ERG4*, pCPC18-*ERG6*, pCPC18-*ERG9*,

pCPC18-*ERG11*, pCPC18-*ERG24*, pCPC18-*ERG26* and pCPC18-*ERG27* were

generated. In the plasmid pCPC18, the ADH1 promoter and the ACT1 terminator were

sequentially inserted into the coding region of the ADE2 gene and loxP-flanked

selection marker, *C. maltosa LEU2*, was placed downstream of the ACT1 terminator.

ERG2, *ERG4*, *ERG6*, *ERG9*, *ERG11*, *ERG24*, *ERG26* and *ERG27* open reading frame

(ORF) fragments were PCR amplified from *C. albicans* SN152 genomic DNA and

inserted into the plasmid pCPC18 backbone in one-step assembly reaction via Exo III-

mediated ligation-independent cloning (LIC)[51].

The plasmids, pCPC18-*ERG2*, pCPC18-*ERG4*, pCPC18-*ERG6*, pCPC18-*ERG9*,

pCPC18-*ERG11*, pCPC18-*ERG24*, pCPC18-*ERG26*, and pCPC18-*ERG27*, which we

described above were used to construct the ectopic overexpression mutants of *ERG2*,

ERG4, *ERG6*, *ERG9*, *ERG11*, *ERG24*, *ERG26* and *ERG27* in *upc2*Δ/Δ null mutant.

Through two rounds of PCR amplification, the ectopic expression cassettes were

transformed and integrated into the *ADE2* locus and were controlled by the constitutive

ADH1 promoter. After selection of transformants on the SD medium without leucine

(SD/-Leu), verification primers were used to verify the successful construction of the

strain with the ectopic overexpression gene[51].

RNA isolation and qRT–PCR

The overnight cultured $P_{tet-off}$-*ERG11*/*erg11*Δ mutant and $P_{tet-off}$-*ENO1*/*eno1*Δ mutant

with the doxycycline (DOX)-inducible Tet-off promoter of *ERG11* or *ENO1* were

1:100 diluted to fresh YPD medium with different concentrations of DOX (0 μg/ml, 8

μg/ml, 16 μg/ml and 32 μg/ml; Aladdin, Shanghai, China) and cultured at 30 °C and

37 °C for 4 h to10^7 cells/ml and collected by centrifugation. RNA was isolated using

the YeaStar RNA Kit (Zymo reserch, USA). The quality of RNA preparations was

assessed by NanoDrop One (Thermo scientific, shanghai, China). One microgram of

RNA was used for complementary DNA (cDNA) synthesis using 5 × PrimeScript RT

Master Mix (Perfect Real Time) (Takara, Beijing, China). Quantitative reverse

transcriptasePCR (qRT–PCR) reactions were prepared using TB Green® Premix Ex

Taq™ II (Tli RNaseH Plus) (Takara, Beijing, China). cDNA mixtures were diluted 1:20

and 2 μl were used in a reaction volume of 20 μl with the primer pairs in Supplementary

Table 3 with the following cycle conditions: preincubation step was 95 °C for 30s; the

amplification step was 95 °C for 5s and 50 °C for 20s, repeat for 40 cycles in Bio-Rad

CFX Connect Real-Time System RT-PCR 96-well (Bio-Rad, Singapore). The levels of

gene expression with three replicates were determined in four separate amplifications

and analyzed using the 2^-ΔΔCt method. Relative levels of *ERG11* and *ENO1*

messenger RNA were normalized with respect to the levels of the invariant reference

gene *PGK1*; the levels of *ERG11* and *ENO1* in mutants treated with different

concentrations of DOX were subsequently averaged and normalized to the levels of

expression in the corresponding mutants without DOX.

RNA-seq and data analysis

C. albicans SN152 were overnight cultured at 30 °C and then 1:100 diluted to fresh

YPD medium cultured at 30 °C and 37 °C and collected at 4-hour, 8-hour and 16-hour

time points. Samples in biological triplicates were submitted to the BGISEQ platform

(BGI Genomics, Wuhan, China) for RNA sequencing analyzes. Transcriptome libraries

were generated using the Standard Sensitivity RNA Analysis Kit (15 nt) (DNF-471)

following the manufacturer's protocols performed by Fragment Analyzer. The raw

RNA-seq reads were quality controlled with SOAPnuke (v1.5.6)[52]. The trimmed reads

were mapped to the final assembly genomes *C. albicans* SC5314 (reference

candidagenome: Assembly22) using HISAT2 (v2.1.0)[53]. Bowtie2 (v2.3.4.3)[54] was used

to compare clean reads to reference gene sequences, and then RSEM (v1.3.1)[55] was

used to calculate gene expression levels for individual samples. DESeq2 (v1.4.5)[56]

used to perform differential gene testing under the condition of Q value ≤ 0.05 or FDR

≤ 0.001. GO enrichment analysis further explored the gene functions associated with

phenotypic changes. Based on hypergeometric testing, we used Phyper to perform GO

on differentially expressed genes enrichment analysis. The gene with Qvalue ≤ 0.05

was defined as significant enrichment in candidate genes that meet this condition. The

value of log2 Fold change (37 °C vs 30 °C) was not directly calculated by taking the

mean expression level (FPKM) and then calculating the log. Instead, it was

standardized using the local standard deviation of the FPKM (counts) based on a linear

regression of all gene expression levels, taking a small interval of the corresponding

gene, and then calculating the corresponding standard deviation. When correcting for

each mRNA expression level, it was related to the distribution of all mRNA expression

levels in each sample.

Protein extractions and western blotting.

To contrast the expression of Erg proteins (Erg6, Erg9, Erg10, Erg11, Erg12, Erg13,

Erg20, Erg26, and Erg251) between 30 °C and 37 °C, the *C. albicans* tagged ERG fused

to a C-terminal GFP were cultured overnight in YPD at 30 °C or 37 °C with shaking.

In order to examine the expression of Erg11 in the presence of FLC, the Erg11-GFP

mutants were cultured overnight in YPD at 30 °C and 1:10 diluted to fresh YPD

medium with or without 4 μg/ml FLC cultured at 30 °C or 37 °C for 4 h. All strains

were harvested at 4 °C, resuspended in 500 μl of protein lysis buffer which consists of

phosphate-buffered saline (PBS; Sangon biotech, Shanghai, China), 5 mM EDTA (pH

8.0; Sangon biotech, Shanghai, China), 1mM PMSF (Sangon biotech, Shanghai, China),

and 1.0% Protease Inhibitor cocktail (TargetMol, Shanghai, China), transferred into

screw-cap tubes with the addition of 500 μl of glass beads, and homogenized by a bead-

beater for 4 cycles of 1min, with 1 min cooling on ice in each cycle. Then the tubes

were centrifuged at 10,000 × g for 10 min at 4 °C and supernatant was transferred to a

new tube. The protein concentration was measured with Bradford Protein Assay Kit

(Epizyme, Shanghai, China). Equal amounts of protein were separated by 8-20% SDS-

PAGE (Byotime, Shanghai, China) and then transferred onto PVDF membranes (Merck

Millipore Ltd., Carrigtwohill, Ireland). After blocking with 1 × Protein Free Rapid

Blocking Buffer (Epizyme, Shanghai, China) for 10 min at room temperature, the

membranes were incubated with primary antibodies against GFP (Anti-GFP [B-2])

mouse monoclonal antibody, 1:1000 dilution; SANTA CRUZ BIOTECHNOLOGY,

USA) or α Tubulin (Anti-α Tubulin [YL1/2] rat monoclonal antibody, 1:1000 dilution;

SANTA CRUZ BIOTECHNOLOGY, USA) or Pgk1 (Anti-Pgk1 [14] mouse

monoclonal antibody, 1:1000 dilution; SANTA CRUZ BIOTECHNOLOGY, USA)

overnight at 4 °C. Then, the membranes were incubated with secondary antibodies

(Peroxidase-conjugated Anti-Mouse IgG$_k$, 1:10000 dilution; SANTA CRUZ

BIOTECHNOLOGY, USA) for 1 h at room temperature. The probed protein was

visualized using an ECL western blotting kit (Epizyme, Shanghai, China) and

photographed by the Tanon 5200 Chemiluminescence Imaging System (Tanon,

Shanghai, China) with AllDoc_X software. The α Tubulin and Pgk1 were used as the

loading control.

Laser scanning confocal microscopy

For the colocalization of Upc2 and nucleus, the Upc2-GFP mutants cultured overnight

at 30 °C were 1:10 diluted to YPD medium containing 16 μg/ml FLC and cultured at

30 °C or 37 °C for 4 h. After washing twice with PBS, cells were collected and

incubated with 10 μg/ml the nuclear dye, Hoechst 33342 (MedChemExpress, Shanghai,

China) for 10 min at room temperature in the dark. Finally, cells were washed with PBS

twice and photographed. The fluorescence distribution curve of the GFP green

fluorescence and nuclear dye (Hoechst) with a red fluorescent pseudocolor in the region

of interest in single cell was used to show the colocalization of Upc2 and nucleus at

30 °C and 37 °C by Image J (1.54g).

For the colocalization of Erg11 and vacuole, the Erg11-GFP mutant grown overnight

were 1:10 diluted to YPD medium and cultured at 30 °C or 37 °C for 1 h. After washing

twice with PBS, cells were collected and incubated with 100 μM vacuole dye,

CellTracker Blue CMAC (MedChemExpress, Shanghai, China) for 30 min at 30 °C or

37 °C in the dark. After washing with PBS twice, the cells were photographed. The

colocalization rate was calculated by dividing the number of cells colocalized showing

yellow fluorescence signal by the total number of cells.

For comparing the autophagy activity between 30 °C and 37 °C, the GFP-Atg8 mutants

grown overnight were 1:10 diluted to YPD medium and cultured at 30 °C or 37 °C for

1 h. After washing twice with PBS the cells were photographed. The autophagy activity

is reflected as the ratio of the number of cells with GFP-ATG8 dots to the total number

of cells at 30 °C and 37 °C.

For the colocalization of Atg8 and mitochondria or ER, the GFP-Atg8 mutants grown

overnight were 1:10 diluted to YPD mediut m and cultured at 30 °C or 37 °C for 30

min. After washing twice with PBS, cells were collected and incubated with 0.3 μM

mitochondria dye, MitoTracker Deep Red FM (MedChemExpress, Shanghai, China) or 1 μM ER dye, ER-tracker Red (MedChemExpress, Shanghai, China) for 30 min at 30 °C or 37 °C in the dark. After washing with PBS twice, the cells were photographed. The colocalization ratio was calculated by dividing the number of GFP-Atg8 dots colocalized with mitochondria or ER by the total number of cells containing GFP-Atg8 dots.

All the images were acquired on an Andor Dragonfly Confocal Imaging System (Andor, Oxford instrument, UK), ImarisViewer (10.1.0) equipped with a white light laser. A 100 × oil-immersion objective was employed. The excitation (Ex)/emission (Em) wavelength for GFP was 488 nm/521 nm. The Ex/Em wavelength of Hoechst 33342 for nucleus was 405 nm/445 nm. The Ex/Em wavelength of CellTracker Blue CMAC for vacuole was 405 nm/445 nm. The Ex/Em of MitoTracker Deep Red FM for mitochondria was 637 nm/660 nm. The Ex/Em of ER-tracker Red for ER was 572 nm/582 nm - 750 nm.

Quantitation of the content of sterols

The content of sterols in *C. albicans* was determined by a method described previously with slight modifications[57]. *C. albicans* SN152 were overnight cultured at 30 °C and then 1:100 diluted to fresh YPD medium cultured at 30 °C or 37 °C and collected at 4-hour, 8-hour and 16-hour time points. For the group with FLC, *C. albicans* SN152 were overnight cultured at 30 °C and then 1:100 diluted to fresh YPD medium with 16 μg/ml FLC, cultured at 30 °C and 37 °C for 16 h. Cells were collected and washed three times with double-distilled water (ddH2O), and the weight of the wet cell pellet was adjusted to approximately 0.5 g. Furthermore, the 15% (m/v) NaOH (Sangon biotech, Shanghai, China) dissolved in 90% (v/v) ethanol (Sangon biotech, Shanghai, China) were added to each pellet and mixed thoroughly. The suspension was incubated at 80 °C for 1 h in a water bath. Sterols were extracted by adding 18 ml of petroleum ether (boiling range: 30 °C-60 °C; SCR, Shanghai, China) in total three times, and each time the mixture was vortexed vigorously for 2 min and allowed to stand for 3 min. The petroleum ether layer was transferred to the same clear glass tubes and washed twice with ddH2O. The collected petroleum ether was volatilized at 65 °C for 20 min in the water bath. Finally,

the sterols were extracted by the addition of 600 μl of n-hexane (Sangon biotech,

Shanghai, China), and a 200-μl aliquot was injected for gas chromatography-mass

spectrometry (Finnigan Voyager, USA) with HP-50 columns (50% phenyl–50%

methylpolysiloxane, 30 m by 0.25 mm by 0.25 μm). Before injection, 20 mg of

cholesterol (Sangon biotech, Shanghai, China) was added to each sample as the internal

reference for quantifying other sterols. Sterols of interest were identified by their

relative retention times, and mass spectra were compared with the sterol profiles of

NIST. The contents of sterol were calculated as the sterol mass corresponding to the

unit mass of cells (μg/mg). Assays were performed in duplicate three times.

Cycloheximide chase analysis

Cycloheximide (CHX) chase analysis was essentially performed by a method with

some modifications[58]. Briefly, The Erg11-GFP mutants were inoculated overnight in

YPD at 30 °C with shaking. The overnighted cultures were diluted to an OD_{600} value

of 0.2 in fresh YPD medium and incubated at 30 °C with shaking for about 4 h, until

the cells reach mid-logarithmic growth phase (OD_{600}=1.0). Then 25 OD_{600} units of each

culture per time point were collected. Then each cell pellet was resuspended in 10 ml

of 30 °C or 37 °C pre-warmed fresh YPD medium per 25 OD_{600} units of cells. After the

equilibrium of cell suspensions by incubation for 5 min in the 30 °C or 37 °C heat block,

cycloheximide (MedChemExpress, Shanghai, China) was added to a final

concentration of 1 mg/ml to the cell suspension and the timer started counting from

0:00. Immediately after adding cycloheximide and vertexing, 10 ml (~25 OD_{600} units)

of the cell suspension with added cycloheximide was transferred to a pre-cooling

microcentrifuge tube to centrifuge at 4,000 × g at 4 °C for 5 min. After that, the cell

suspension was removed the supernatant and washed with sterile water twice at 4,000

× g at 4 °C for 5 min and stored at -20 °C. When all samples at different time points

have been collected, the cells are ready to lysis for western blotting like we described

above. To investigate the effects of Wortmannin and MG132 on the degradation of

Erg11, Wortmannin (1 μM) and MG132 (100 μM) were added after adding

cycloheximide to inhibit the protein synthesis. Pgk1 was used as the loading control.

Assays were performed in duplicate three times.

GFP−ATG8 cleavage assay

In principle, GFP-Atg8 translocates along autophagosomes and is subsequently released into vacuoles. GFP is more resistant to vacuolar hydrolases, leading to the degradation of Atg8 while retaining GFP. Therefore, the ratio of free GFP to GFP−ATG8 can reflect the activity of autophagy. To investigate the effect of different temperatures on the activity of autophagy, the GFP-Atg8 mutants were overnight cultured and then 1:10 diluted in YPD medium at 30 °C or 37 °C for 4 h. Total proteins were extracted, measured with Bradford Protein Assay Kit and immunoblotted with primary antibodies against GFP like what we describe in the western blotting assay. Pgk1 was used as the loading control. The GFP cleavage activity was quantified by determining the ratio of free GFP to GFP−ATG8.

Measurement of intracellular reactive oxygen species

Intracellular ROS was measured using 2',7'-dichlorodihydrofluoroscein diacetate (DCFH-DA) with some modifications[59]. In principle, DCFH-DA entering the cell can be hydrolyzed to form 2',7'-dichlorodihydrofluorescein (DCFH) which cannot permeate cell membranes and then is oxidized by ROS to form fluorescent 2',7'-Dichlorofluorescein (DCF). ROS level can be reflected by the measurement of fluorescent DCF and calculation of specific fluorescence intensity (SFI). Specifically, the *C. albicans* SN152 were overnight cultured at 30 °C and then 1:5 diluted and treated with 10 μM DCFH-DA (MedChemExpress, Shanghai, China) and incubated at 30 °C or 37 °C with shaking for 30 min. The cells were harvested at 4 °C and washed twice with PBS. Then cells were suspended in 500 μl PBS and transferred into screw-cap tubes with the addition of glass beads equal in volume to the cell pellet, and homogenized by a bead-beater for 4 cycles of 1 min, with 1 min cooling on ice in each cycle. Then the tubes were centrifuged at $10,000 \times$ g for 10 min at 4 °C and supernatant was transferred to a new tube. The fluorescence was recorded, at 25 °C, at an excitation wavelength (Ex) of 485 nm and an emission (Em) wavelength of 535 nm, using the microplate reader, Infinite F Plex (Tecan, Groedig, Austria). Em measurements were

normalized by estimation of the protein content of the supernatant using a Bradford

protein assay kit. The SFI was expressed as fluorescence [(Em = 535 nm/Ex = 485

nm)/mg protein]. As a control, background fluorescence of cells not treated with

DCFH-DA, was also recorded. In order to ensure that the fluorescence readings were

in the linear dynamic range of DCF concentrations (1 nM – 400 nM), a calibration curve

was constructed of fluorescence (Em = 535 nm, Ex = 485 nm) vs. DCF concentration.

Growth curve assays

Growth curve assay was performed as described previously[60]. *C. albicans* SN152 and

atg8Δ/Δ null mutant were overnight cultured at 30 °C. Then cells were adjusted to the

concentration of 10^3 cells/ml by the hemacytometer. Then cells were cultured at 30 °C

or 37 °C with shaking (200 rpm). The optical density at 600 nm (OD_{600}) was measured

every hour until the stationary phase of the growth curve was reached by using the

microplate reader, Infinite M Nano (Tecan, Groedig, Austria).

Mitochondrial Membrane Potential (MMP)

Changes in MMP were determined by the MMP probe JC-1 as previously described

with some modification. *C. albicans* SN152 and *atg8Δ/Δ* null mutant were overnight

cultured at 30 °C and then 1:10 diluted to fresh YPD medium cultured at 30 °C or 37 °C

for 4 h. Then cells were washed and resuspended in PBS with 2.5 μg/ml JC-1 at 37 °C

for 20 min in the dark, then washed twice. All samples were estimated via the BD

LSRFortessa Cell Analyzer (LSRFortessa; BD Bioscience, USA). Gating of single cells

was achieved as follows. The green fluorescence signal of the JC-1 monomer was

detected using the FITC channel equipped with a 530 nm band-pass filter (bandwidth

30 nm) while the red fluorescence signal of JC-1 aggregates was detected using the PE

channel equipped with a 585 nm band-pass filter (bandwidth 15 nm). First, the cells

populations were separated from cell debris and selected via SSC-A (side scatter; U =

302 V) and FSC-A (forward scatter; U = 640 V) signals. Second, duplet cells were

excluded via FSC-W and FSC-H signals. A total of 50,000 events was analyzed per

sample. Flow cytometry data were collected using FACS Diva Software and analysed

using FlowJo (10.8.1). SN152 cells cultured at 30 °C for 4 h with healthy mitochondria

were considered as negative control sample. Here, according to the negative control

sample, P1 was set as a region in which > 95% of the dual fluorescent population falls

within it and the JC-1 red/green fluorescence ratio > 1[61-63]. The loss of MMP was

reflected by the proportion of cells in P2 with less aggregates (red fluorescence) and

more monomer (green fluorescence) among different groups. The ordinary one-way

ANOVA, Bonferroni's multiple comparisons test was used to evaluate multiple

comparisons of the proportion of cells with damaged mitochondria.

Statistical analysis

Statistical analysis was performed using Graphpad Prism 9.0 software. The two-tailed

paired t-test was used to compare the values of SMG of FLC against clinical *Candida*

isolates (*C. glabrata*, *C. krusei* and *C. guilliermondii*) between 30 °C and 37 °C. The

Wilcoxon matched-pairs signed rank test was used to compare the values of SMG of

FLC against *C. albicans* and *C. parapsilosis* between 30 °C and 37 °C. The two-tailed

unpaired t-test was used to compare the expression of proteins in WB, sterol content,

the degradation of Erg11-GFP, the colocalization rate, the autophagic activity, Atg8

dots and ROS level between two groups. The ordinary one-way ANOVA, Bonferroni's

multiple comparisons test was used to evaluate multiple comparisons of the degradation

of Erg11-GFP in the presence of different compounds and the proportion of cells with

damaged mitochondria. For all analyses, $P < 0.05$ was considered significant. Data are

shown as means ± standard deviation (s.d.) from three or more replicates.

Data availability

713   The RNA sequencing data of *C. albicans* SN152 cultured at 30 °C and 37 °C for 4 h, 8

714   h and 16 h are publicly available at the NCBI Sequence Read Archive (SRA) and can

715   be accessed under BioProject PRJNA994182.

Acknowledgments

717   This study received financial support from the National Natural Science Foundation of

718   China (No. 82020108032), the Innovation Program of Shanghai Municipal Education

719   Commission (202101070007-E00094), and the National Key Research and

720   Development Program of China (No.2022YFC2303004).

Author Contributions

YR.F. conducted most of the experiments and performed data analysis. **YR.F., H. L., M.W.,** and **YY.J.** wrote the manuscript draft and revised the manuscript. **YR.F., W.L., C.Z., XY.F,** and **XQ.S.** constructed mutant strains. **YR.F., W.L., C.Z., XY.F,** and **XQ.S.** evaluated the antifungal activities and data collection. **YR.F., H. L., M. W.,** and **YY.J.** discussed and analyzed the data. **H.L.** and **YY.J.** conceived the idea. **H.L.** and **YY.J.** directed the experiments.

Competing Interests statement

The authors declare no conflicts of interest.

References

Lu, H., Hong, T., Jiang, Y., Whiteway, M. & Zhang, S. Candidiasis: From cutaneous to systemic, new perspectives of potential targets and therapeutic strategies. *Adv Drug Deliv Rev* **199**, 114960, doi:10.1016/j.addr.2023.114960 (2023).

Denning, D. W. Global incidence and mortality of severe fungal disease. *Lancet Infect Dis* **24**, e428-e438, doi:10.1016/s1473-3099(23)00692-8 (2024).

Shafiei, M., Peyton, L., Hashemzadeh, M. & Foroumadi, A. History of the development of antifungal azoles: A review on structures, SAR, and mechanism of action. *Bioorg Chem* **104**, 104240, doi:10.1016/j.bioorg.2020.104240 (2020).

Berman, J. & Krysan, D. J. Drug resistance and tolerance in fungi. *Nat Rev Microbiol* **18**, 319-331, doi:10.1038/s41579-019-0322-2 (2020).

Pfaller, M. A., Diekema, D. J., Turnidge, J. D., Castanheira, M. & Jones, R. N. Twenty Years of the SENTRY Antifungal Surveillance Program: Results for Candida Species From 1997-2016. *Open Forum Infect Dis* **6**, S79-S94, doi:10.1093/ofid/ofy358 (2019).

Lepak, A. J. & Andes, D. R. Antifungal pharmacokinetics and pharmacodynamics. *Cold

Spring Harb Perspect Med **5**, a019653, doi:10.1101/cshperspect.a019653 (2014).

7 Li, W. *et al.* Pitavastatin Calcium Confers Fungicidal Properties to Fluconazole by

Inhibiting Ubiquinone Biosynthesis and Generating Reactive Oxygen Species.

Antioxidants (Basel) **13**, doi:10.3390/antiox13060667 (2024).

8 Rosenberg, A. *et al.* Antifungal tolerance is a subpopulation effect distinct from

resistance and is associated with persistent candidemia. *Nat Commun* **9**, 2470,

doi:10.1038/s41467-018-04926-x (2018).

9 Levinson, T. *et al.* Impact of tolerance to fluconazole on treatment response in Candida

albicans bloodstream infection. *Mycoses* **64**, 78-85, doi:10.1111/myc.13191 (2021).

10 Hammoud, M. S., Al-Taiar, A., Fouad, M., Raina, A. & Khan, Z. Persistent candidemia

in neonatal care units: risk factors and clinical significance. *Int J Infect Dis* **17**, e624-

628, doi:10.1016/j.ijid.2012.11.020 (2013).

11 Xu, Y. *et al.* Multifactorial Mechanisms of Tolerance to Ketoconazole in Candida

albicans. *Microbiol Spectr* **9**, e0032121, doi:10.1128/Spectrum.00321-21 (2021).

12 Yang, F. *et al.* Antifungal Tolerance and Resistance Emerge at Distinct Drug

Concentrations and Rely upon Different Aneuploid Chromosomes. *mBio* **14**, e0022723,

doi:10.1128/mbio.00227-23 (2023).

13 Huang, J. *et al.* Pan-drug resistance and hypervirulence in a human fungal pathogen

are enabled by mutagenesis induced by mammalian body temperature. *Nat Microbiol*

9, 1686-1699, doi:10.1038/s41564-024-01720-y (2024).

14 Fisher, M. C. *et al.* Tackling the emerging threat of antifungal resistance to human

health. *Nat Rev Microbiol* **20**, 557-571, doi:10.1038/s41579-022-00720-1 (2022).

15   Fisher, M. C., Hawkins, N. J., Sanglard, D. & Gurr, S. J. Worldwide emergence of

resistance to antifungal drugs challenges human health and food security. *Science* **360**,

739-742, doi:10.1126/science.aap7999 (2018).

16   Zhou, D. *et al.* Extensive Genetic Diversity and Widespread Azole Resistance in

Greenhouse Populations of Aspergillus fumigatus in Yunnan, China. *mSphere* **6**,

doi:10.1128/mSphere.00066-21 (2021).

17   Duong, T. N. *et al.* Azole-resistant Aspergillus fumigatus is highly prevalent in the

environment of Vietnam, with marked variability by land use type. *Environ Microbiol* **23**,

7632-7642, doi:10.1111/1462-2920.15660 (2021).

18   Daneshnia, F. *et al.* Worldwide emergence of fluconazole-resistant Candida

parapsilosis: current framework and future research roadmap. *Lancet Microbe* **4**, e470-

e480, doi:10.1016/S2666-5247(23)00067-8 (2023).

19   Casadevall, A., Kontoyiannis, D. P. & Robert, V. On the Emergence of Candida auris:

Climate Change, Azoles, Swamps, and Birds. *mBio* **10**, doi:10.1128/mBio.01397-19

(2019).

20   Akinbobola, A. B., Kean, R., Hanifi, S. M. A. & Quilliam, R. S. Environmental reservoirs

of the drug-resistant pathogenic yeast Candida auris. *PLoS Pathog* **19**, e1011268,

doi:10.1371/journal.ppat.1011268 (2023).

21   Bottery, M. J. & Denning, D. W. Body heat drives antifungal resistance. *Nat Microbiol*

9, 1638-1639, doi:10.1038/s41564-024-01738-2 (2024).

22   Lu, H. *et al.* A Small Molecule Inhibitor of Erg251 Makes Fluconazole Fungicidal by

Inhibiting the Synthesis of the 14alpha-Methylsterols. *mBio* **14**, e0263922,

790   doi:10.1128/mbio.02639-22 (2023).

791   23   Wang, H. *et al.* Myriocin enhances the antifungal activity of fluconazole by blocking the

792   membrane localization of the efflux pump Cdr1. *Front Pharmacol* **13**, 1101553,

793   doi:10.3389/fphar.2022.1101553 (2022).

794   24   Vasicek, E. M., Berkow, E. L., Flowers, S. A., Barker, K. S. & Rogers, P. D. UPC2 is

795   universally essential for azole antifungal resistance in Candida albicans. *Eukaryot Cell*

796   **13**, 933-946, doi:10.1128/ec.00221-13 (2014).

797   25   Lo, H. J. *et al.* Efg1 involved in drug resistance by regulating the expression of ERG3

798   in Candida albicans. *Antimicrob Agents Chemother* **49**, 1213-1215,

799   doi:10.1128/AAC.49.3.1213-1215.2005 (2005).

800   26   Sellam, A., Tebbji, F. & Nantel, A. Role of Ndt80p in sterol metabolism regulation and

801   azole resistance in Candida albicans. *Eukaryot Cell* **8**, 1174-1183,

802   doi:10.1128/EC.00074-09 (2009).

803   27   Yang, H. *et al.* Structural mechanism of ergosterol regulation by fungal sterol

804   transcription factor Upc2. *Nat Commun* **6**, 6129, doi:10.1038/ncomms7129 (2015).

805   28   Martel, C. M. *et al.* Identification and characterization of four azole-resistant erg3

806   mutants of Candida albicans. *Antimicrob Agents Chemother* **54**, 4527-4533,

807   doi:10.1128/AAC.00348-10 (2010).

808   29   Polak-Wyss, A., Lengsfeld, H. & Oesterhelt, G. Effect of oxiconazole and Ro 14-

809   4767/002 on sterol pattern in Candida albicans. *Sabouraudia* **23**, 433-441,

810   doi:10.1080/00362178585380631 (1985).

30   Foresti, O., Rodriguez-Vaello, V., Funaya, C. & Carvalho, P. Quality control of inner

nuclear membrane proteins by the Asi complex. *Science* **346**, 751-755,

doi:10.1126/science.1255638 (2014).

31   Natarajan, N., Foresti, O., Wendrich, K., Stein, A. & Carvalho, P. Quality Control of

Protein Complex Assembly by a Transmembrane Recognition Factor. *Mol Cell* **77**, 108-

119 e109, doi:10.1016/j.molcel.2019.10.003 (2020).

32   Khmelinskii, A. *et al.* Protein quality control at the inner nuclear membrane. *Nature* **516**,

410-413, doi:10.1038/nature14096 (2014).

33   Delorme-Axford, E., Guimaraes, R. S., Reggiori, F. & Klionsky, D. J. The yeast

Saccharomyces cerevisiae: an overview of methods to study autophagy progression.

Methods **75**, 3-12, doi:10.1016/j.ymeth.2014.12.008 (2015).

34   Marshall, R. S. & Vierstra, R. D. A trio of ubiquitin ligases sequentially drives

ubiquitylation and autophagic degradation of dysfunctional yeast proteasomes. *Cell*

Rep **38**, 110535, doi:10.1016/j.celrep.2022.110535 (2022).

35   Zhen, C. *et al.* Otilonium Bromide Exhibits Potent Antifungal Effects by Blocking

Ergosterol Plasma Membrane Localization and Triggering Cytotoxic Autophagy in

Candida Albicans. *Adv Sci (Weinh)*, e2406473, doi:10.1002/advs.202406473 (2024).

36   Chang-Chien, J. *et al.* Vitamin D ameliorates particulate matter induced mitochondrial

damages and calcium dyshomeostasis in BEAS-2B human bronchial epithelial cells.

Respir Res **25**, 321, doi:10.1186/s12931-024-02951-7 (2024).

37   Zhou, J. *et al.* A non-canonical role of ATG8 in Golgi recovery from heat stress in plants.

Nat Plants **9**, 749-765, doi:10.1038/s41477-023-01398-w (2023).

38   Feng, Y., Lu, H., Whiteway, M. & Jiang, Y. Understanding fluconazole tolerance in

Candida albicans: implications for effective treatment of candidiasis and combating

invasive fungal infections. *J Glob Antimicrob Resist* **35**, 314-321,

doi:10.1016/j.jgar.2023.10.019 (2023).

39   Xiong, J., Lu, H. & Jiang, Y. Mechanisms of Azole Potentiation: Insights from Drug

Repurposing Approaches. *ACS Infect Dis* **11**, 305-322,

doi:10.1021/acsinfecdis.4c00657 (2025).

40   Lu, H., Shrivastava, M., Whiteway, M. & Jiang, Y. Candida albicans targets that

potentially synergize with fluconazole. *Crit Rev Microbiol* **47**, 323-337,

doi:10.1080/1040841x.2021.1884641 (2021).

41   Jordá, T. & Puig, S. Regulation of Ergosterol Biosynthesis in Saccharomyces

cerevisiae. *Genes (Basel)* **11**, doi:10.3390/genes11070795 (2020).

42   Van Rhijn, N., Bromley, M., Richardson, M. & Bowyer, P. CYP51 Paralogue Structure

Is Associated with Intrinsic Azole Resistance in Fungi. *mBio* **12**, e0194521,

doi:10.1128/mBio.01945-21 (2021).

43   Hu, T. *et al.* Hotspot mutations and genomic expansion of ERG11 are major

mechanisms of azole resistance in environmental and human commensal isolates of

Candida tropicalis. *Int J Antimicrob Agents* **62**, 107010,

doi:10.1016/j.ijantimicag.2023.107010 (2023).

44   Handelman, M. *et al.* Point Mutation or Overexpression of Aspergillus fumigatus

cyp51B, Encoding Lanosterol 14alpha-Sterol Demethylase, Leads to Triazole

Resistance. *Antimicrob Agents Chemother* **65**, e0125221, doi:10.1128/AAC.01252-21

(2021).

45   Villasmil, M. L., Barbosa, A. D., Cunningham, J. L., Siniossoglou, S. & Nickels, J. T.,

Jr. An Erg11 lanosterol 14-alpha-demethylase-Arv1 complex is required for Candida

albicans virulence. *PLoS One* **15**, e0235746, doi:10.1371/journal.pone.0235746

(2020).

46   Krshnan, L., van de Weijer, M. L. & Carvalho, P. Endoplasmic Reticulum-Associated

Protein Degradation. *Cold Spring Harb Perspect Biol* **14**,

doi:10.1101/cshperspect.a041247 (2022).

47   Li, Y., Li, S. & Wu, H. Ubiquitination-Proteasome System (UPS) and Autophagy Two

Main Protein Degradation Machineries in Response to Cell Stress. *Cells* **11**,

doi:10.3390/cells11050851 (2022).

48   Lu, H. *et al.* A Small Molecule Inhibitor of Erg251 Makes Fluconazole Fungicidal by

Inhibiting the Synthesis of the 14α-Methylsterols. *mBio* **14**, e0263922,

doi:10.1128/mbio.02639-22 (2023).

49   Epp, E. *et al.* Reverse genetics in Candida albicans predicts ARF cycling is essential

for drug resistance and virulence. *PLoS Pathog* **6**, e1000753,

doi:10.1371/journal.ppat.1000753 (2010).

50   Noble, S. M. & Johnson, A. D. Strains and strategies for large-scale gene deletion

studies of the diploid human fungal pathogen Candida albicans. *Eukaryot Cell* **4**, 298-

309, doi:10.1128/ec.4.2.298-309.2005 (2005).

51   Chang, P., Wang, W., Igarashi, Y., Luo, F. & Chen, J. Efficient vector systems for

economical and rapid epitope-tagging and overexpression in Candida albicans. *J*

Microbiol Methods **149**, 14-19, doi:10.1016/j.mimet.2018.04.016 (2018).

52   Li, R., Li, Y., Kristiansen, K. & Wang, J. SOAP: short oligonucleotide alignment program.

Bioinformatics **24**, 713-714, doi:10.1093/bioinformatics/btn025 (2008).

53   Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low memory

requirements. *Nat Methods* **12**, 357-360, doi:10.1038/nmeth.3317 (2015).

54   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat*

Methods **9**, 357-359, doi:10.1038/nmeth.1923 (2012).

55   Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with

or without a reference genome. *BMC Bioinformatics* **12**, 323, doi:10.1186/1471-2105-

12-323 (2011).

56   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol* **15**, 550, doi:10.1186/s13059-014-0550-

8 (2014).

57   Lv, Q. Z. *et al.* NSG2 (ORF19.273) Encoding Protein Controls Sensitivity of Candida

albicans to Azoles through Regulating the Synthesis of C14-Methylated Sterols. *Front*

Microbiol **9**, 218, doi:10.3389/fmicb.2018.00218 (2018).

58   Buchanan, B. W., Lloyd, M. E., Engle, S. M. & Rubenstein, E. M. Cycloheximide Chase

Analysis of Protein Degradation in Saccharomyces cerevisiae. *J Vis Exp*,

doi:10.3791/53975 (2016).

59   Moraitis, C. & Curran, B. P. Reactive oxygen species may influence the heat shock

response and stress tolerance in the yeast Saccharomyces cerevisiae. *Yeast* **21**, 313-

323, doi:10.1002/yea.1078 (2004).

60   Lu, H. *et al.* Loss of RPS41 but not its paralog RPS42 results in altered growth,

filamentation and transcriptome changes in Candida albicans. *Fungal Genet Biol* **80**,

31-42, doi:10.1016/j.fgb.2015.03.012 (2015).

61   Guthrie, H. D. & Welch, G. R. Determination of intracellular reactive oxygen species

and high mitochondrial membrane potential in Percoll-treated viable boar sperm using

fluorescence-activated flow cytometry. *J Anim Sci* **84**, 2089-2100,

doi:10.2527/jas.2005-766 (2006).

62   Yamakawa, K. *et al.* Platelet mitochondrial membrane potential correlates with severity

in patients with systemic inflammatory response syndrome. *J Trauma Acute Care Surg*

74, 411-417; discussion 418, doi:10.1097/TA.0b013e31827a34cf (2013).

63   Elefantova, K., Lakatos, B., Kubickova, J., Sulova, Z. & Breier, A. Detection of the

Mitochondrial Membrane Potential by the Cationic Dye JC-1 in L1210 Cells with

Massive Overexpression of the Plasma Membrane ABCB1 Drug Transporter. *Int J Mol*

Sci **19**, doi:10.3390/ijms19071985 (2018).

Figures

Figure 1. Short-term exposure to body temperature enhances the tolerance of *C. albicans* to azoles. **(a)** Spotting assays of *C. albicans* SN152. Cells were spotted onto YPD agar containing different concentrations (0 μg/ml, 0.5 μg/ml, 1 μg/ml, 2 μg/ml, 4 μg/ml, 8 μg/ml and 16 μg/ml) of fluconazole (FLC) and cultured at 30 °C or 37 °C for 2 days. The different colors of right triangle indicate cells were 1:10 serially diluted and cultured at 30 °C (blue one) or 37 °C (red one). **(b)** Illustration of minimum inhibitory concentration (MIC) and supra-MIC growth (SMG) calculations. MIC was calculated at 24 h as the FLC concentration at which 50% of the growth was inhibited at 30 °C, relative to growth without FLC. SMG was calculated as the average growth per well above the MIC divided by the level of growth without FLC. The darker the yellow color, the more vigorous the cell growth. **c**, Broth microdilution assays showing MIC and

SMG levels at 24 h for *C. albicans* wild-type SN152 and *cdr1Δ/Δ* null mutant cultured

at 30 °C or 37 °C in the presence of FLC. **d**, The values of SMG of FLC against clinical

Candida isolates which were cultured at 30 °C or 37 °C. Data represent mean ± standard

deviation (s.d). The *P* values were determined by the Wilcoxon matched-pairs signed

rank test for *C. albicans* ($n = 104$) and *C. parapsilosis* ($n = 10$) and the two-tailed paired

t test for *C. glabrata* ($n = 10$), *C. krusei* ($n = 10$) and *C. guilliermondii* ($n = 10$). **e**, Broth

microdilution assays showing SMG levels at 24 h for *C. albicans* SN152 cultured at

30 °C or 37 °C in the presence of azoles. Technical duplicates were averaged. **f**, Broth

microdilution assays showing MIC levels at 24 h for *C. albicans* SN152 cultured at

30 °C or 37 °C in the presence of terbinafine and fluvastatin. Technical duplicates were

averaged.

Figure 2. Body temperature enhances *C. albicans*' tolerance to azoles through Erg11.

a, Broth microdilution assays showing SMG levels at 24 h for *C. albicans* SN152

cultured at 30 °C or 37 °C in the presence of FLC combined with fluvastatin (1 μg/ml)

or terbinafine (1 μg/ml). The control group represents cells were in the presence of FLC

without combined compounds. **b**, Broth microdilution assays showing MIC and SMG

levels at 24 h for *upc2Δ/Δ* null mutant cultured at 30 °C or 37 °C in the presence of

FLC. Technical duplicates were averaged. **c**, The expression of proteins involved in

ergosterol synthesis (Erg6, Erg9, Erg10, Erg11, Erg12, Erg 13, Erg20, Erg26 and

Erg251) examined by western blotting. Cells were precultured at 30 °C or 37 °C for 16

h. The representative images shown are from three replicates. **d**, Quantification of the

signal intensity ratio of Erg-GFP to Tubulin by Image J. Data represent means ± s.d. of

three replicates, two-tailed, unpaired t test. **e**, Illustration of ergosterol synthesis

pathway and ergosterol synthesis inhibitors. **f**, The determination of sterol content in

the presence of 16 μg/ml FLC based on GC-MS at the 16-hour time point. Data

represent means ± s.d. of three replicates, two-tailed, unpaired t test. **g**, The relative

growth at optical density at 600 nm (OD_{600}) of *upc2Δ/Δ* null mutants in which the

expression of ergosterol biosynthesis genes (*ERG2*, *ERG4*, *ERG6, ERG9, ERG11*,

ERG24, ERG26, and *ERG27*) were increased by ADH1 promoter in the presence of

0.25 μg/ml FLC at 30 °C or 37 °C for 24 h. Technical duplicates were averaged.

Figure 3. Body temperature inhibits the degradation of Erg11. **a**, Broth microdilution assays showing MIC levels at 24 h for P$_{\text{tet-off}}$-*ERG11/erg11*Δ mutant and P$_{\text{tet-off}}$-*ENO1/eno1*Δ mutant cultured at 30 °C or 37 °C in the presence of DOX. Technical duplicates were averaged. **b**, Broth microdilution assays showing SMG levels at 24 h for *C. albicans* SN152 cultured in different culture medium (YPD or SD) at 30 °C or 37 °C in the presence of FLC. Technical duplicates were averaged. **c**, Degradation of Erg11-GFP in *C. albicans* SN152 after inhibition of protein synthesis by cycloheximide upon a 3-hour shift to 30 °C or 37 °C. The representative images shown are from three replicates. **d**, Quantification of the degradation of Erg11-GFP calculated as the ratio of Erg11-GFP to Pgk1 by Image J. Set the protein level at the initial time point to 100 %. Data represent means ± s.d. of three replicates, two-tailed, unpaired t test. **e**, The expression of Erg11 protein examined by western blotting. Cells were precultured at 30 °C and 37 °C for 4 h with or without 4 μg/ml FLC. The representative images shown are from three replicates. **f**, Quantification of the signal intensity ratio of Erg11-GFP to Pgk1 by Image J. Data represent means ± s.d. of three replicates, two-tailed, unpaired t test.

Figure 4. Erg11 undergoes degradation through autophagy. **a**, Broth microdilution
assays showing SMG levels at 24 h for *C. albicans* SN152 cultured at 30 °C or 37 °C
in the presence of FLC combined with wortmannin (1 μM) or MG132 (6.25 μM).
Technical duplicates were averaged. **b**, Degradation of Erg11-GFP in *C. albicans*
SN152 in the presence of Wortmannin (1 μM) and MG132 (100 μM) after inhibition of
protein synthesis by cycloheximide upon a 2-hour shift to 30 °C. **c**, Quantification of
the degradation of Erg11-GFP calculated as the ratio of Erg11-GFP to Pgk1 by Image
J. Data represent means ± s.d. of three replicates, ordinary one-way ANOVA,
Bonferroni's multiple comparisons test. **d**, Colocalization between Erg11-GFP and
vacuole in cells precultured at 30 °C or 37 °C for 1 h. Pseudo-color was used to display
the blue fluorescence of the vacuole dye (CellTracker Blue CMAC) as red. The white
arrow indicated vacuoles with Erg11-GFP. The representative images shown are from
three replicates. Scale bar, 5 μm. **e**, Quantification of the colocalization ratio of the
number of cells colocalized with Erg11 and vacuoles to the total number of cells. The
data represent means ± s.d of three replicates, two-tailed, unpaired t test.

Figure 5. Body temperature inhibits autophagy in *C. albicans*. **a**, GFP-ATG8 cleavage

assay showing the autophagic activity after pre-cultivation at 30 °C or 37 °C for 4 h. **b**,

Quantification of the autophagic activity displaying as the ratio of free GFP to GFP-

ATG8 by Image J. Data represent means ± s.d. of three replicates, two-tailed, unpaired

t test. **c**, Confocal images of GFP-ATG8 dots in cells precultured at 30 °C or 37 °C for

1 h. BF represents cells in bright filed. Scale bar, 2 μm. **d**, The ratio of the number of

cells with GFP-ATG8 dots to the total number of cells. Data represent means ± s.d.

form three replicates, two-tailed, unpaired t test. **e**, Colocalization between GFP-Atg8

dots and mitochondria in cells precultured at 30 °C or 37 °C for 30 min. Scale bar, 2 μm.

f, Quantification of the colocalization ratio of the number of GFP-Atg8 dots colocalized

with mitochondria to the number of cells containing GFP-Atg8 dots. The data represent

means ± s.d. n = 6 confocal microscopic images form three replicates, two-tailed,

1010   unpaired t test. **g**, Colocalization between GFP-Atg8 dots and ER in cells precultured

1011   at 30 °C or 37 °C for 30 min. Scale bar, 2 μm. **h**, Quantification of the colocalization

1012   ratio of the number of GFP-Atg8 dots colocalized with ER to the number of cells

1013   containing GFP-Atg8 dots. The data represent means ± s.d. n = 6 confocal microscopic

1014   images form three replicates, two-tailed, unpaired t test.

1015

1016

Figure 6. Atg8 relocalizes to mitochondria to safeguard them against oxidative damage

induced by Body temperature. **a**, Gene Ontology categories according to cellular

components (GO-CC) enrichment based on the 1733 genes with significant differences

of expression level (Qvalue ≤ 0.05) in *C. albicans* SN152 cultured at 30 °C and 37 °C

for 4 h. **b**, The intracellular ROS levels measured by specific fluorescence intensity

(SFI) of the oxidant-sensitive probe 2′,7′-dichlorofluorescin diacetate (DCFH-DA) in

cells precultured at 30 °C or 37 °C for 30 min. SFI was calculated as fluorescence [(Em

= 535 nm/Ex = 485 nm)/mg protein]. Data represent means ± s.d. of three replicates,

two-tailed, unpaired t test. **c**, Broth microdilution assays showing SMG levels at 24 h

for *C. albicans* SN152 cultured at 37 °C in the presence of FLC combined with NAC

(0 mM and 50mM). Technical duplicates were averaged. **d**, Growth curve of *C.*

albicans SN125 (blue) and *atg8Δ/Δ* null mutant (red) at 30 °C (left) and 37 °C (right)

for 24 h. The optical density at 600 nm (OD_{600}) was measured every hour. The colored

bands show the means ± s.d. of six replicates. **e**, Determination of mitochondrial

membrane potential (MMP) by JC-1 for *C. albicans* SN125 and the *atg8Δ/Δ* null

mutant at 30 °C and 37 °C for 4 h through flow cytometry. JC-1 aggregates show red

fluorescence (FL-1 channel, y-axis) and JC-1monomer shows green fluorescence (FL-

1034   2 channel, x-axis).   Healthy cells were classified in P1, while cells with damaged

1035   mitochondria were classified in P2. The representative images shown are from three

1036   replicates. **f**, Quantification of the proportion of cells in P2 with damaged mitochondria

1037   by FlowJo. Data represent means ± s.d. of three replicates, ordinary one-way ANOVA,

1038   Bonferroni's multiple comparisons test.

1040

Fig.7 Schematic diagram showing a model of body temperature-induced azoles tolerance. Compared with 30 °C, body temperature (37 °C) results in an elevated mitochondrial production of ROS and subsequently leads to mitochondrial dysfunction which causes the relocation of Atg8 to mitochondria and the inhibition of autophagy. As the result, the slower degradation of Erg11 results in azole tolerance in *C. albicans* at 37 °C.

1047