CenSegNet: a generalist high-throughput deep learning framework for

centrosome phenotyping at spatial and single-cell resolution in

heterogeneous tissues

Jiaoqi Cheng[1,2,8], Keqiang Fan[2,3,8], Miles Bailey[1,2], Xin Du[4], Rajesh Jena[5], Costantinos Savva[6,7],

Mengyang Gou[1,2], Ramsey Cutress[2,6], Stephen Beers[2,7*], Xiaohao Cai[2,3*] and Salah Elias[1,2,9*]

[1]School of Biological Sciences, University of Southampton, Southampton SO17 1BJ, UK

[2]Institute for Life Sciences, University of Southampton, Southampton SO17 1BJ, UK

[3]School of Electronics and Computer Science, University of Southampton, Southampton SO17

1BJ, UK

[4]The Cavendish Laboratory, Department of Physics, University of Cambridge, Cambridge, CB3

0FZ, UK

[5]Department of Oncology, Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK,

University of Cambridge, CB2 0QQ, UK

[6]University Hospital Southampton NHS Foundation Trust, Southampton, SO16 6YD, UK

[7]Centre for Cancer Immunology, School of Cancer Sciences, Faculty of Medicine, University of

Southampton, Southampton, SO16 6YD, UK

[8]These authors contributed equally

[9]Lead contact

*Correspondence: s.a.beers@soton.ac.uk, x.cai@soton.ac.uk and s.k.elias@soton.ac.uk

Abstract

Centrosome amplification (CA) is a hallmark of epithelial cancers, yet its spatial complexity and phenotypic heterogeneity remain poorly resolved due to limitations in conventional image analysis. We present CenSegNet (Centrosome Segmentation Network), a modular deep learning framework for high-resolution, context-aware segmentation of centrosomes and epithelial architecture across diverse tissue types. Integrating a dual-branch architecture with uncertainty-guided refinement, CenSegNet achieves state-of-the-art performance and generalisability across both immunofluorescence and immunohistochemistry modalities, outperforming existing models in accuracy and morphological fidelity. Applied to tissue microarrays (TMAs) containing 911 breast cancer sample cores from 127 patients, CenSegNet enables the first large-scale, spatially resolved quantification of numerical and structural CA at single-cell resolution. These CA subtypes are mechanistically uncoupled, exhibiting distinct spatial distributions, age-dependent dynamics, and associations with histological tumour grade, hormone receptor status, genomic alterations, and nodal involvement. Discordant CA profiles at tumour margins are linked to local aggressiveness and stromal remodelling, underscoring their clinical relevance. To support broad adoption and reproducibility, CenSegNet is released as an open-source Python library. Together, our findings establish CenSegNet as a scalable, generalisable platform for spatially resolved centrosome phenotyping in intact tissues, enabling systematic dissection of the biology of this organelle and its dysregulation in cancer and other epithelial diseases.

Introduction

Centrosomes, composed of a pair of orthogonally arranged centrioles surrounded by pericentriolar material (PCM), function as the principal microtubule-organising centres (MTOCs) in animal cells. They play essential roles in diverse cellular processes, including vesicular trafficking, cell polarity, motility, ciliogenesis, and the assembly of a bipolar mitotic spindle during cell division[1, 2]. Centrosome number is tightly regulated, with duplication occurring once per cell cycle during S phase, ensuring the formation of the mitotic spindle and equal inheritance of chromosomes by daughter cells[3-5].

Centrosome amplification (CA) can lead to multipolar spindle formation, chromosomal missegregation, and aneuploidy[3, 6, 7]—a hallmark of cancer[3, 4, 7]. The hypothesis that CA-induced aneuploidy contributes to tumorigenesis was first proposed by Theodor Boveri over a century ago[8]. In recent years, CA has been documented in several solid tumours including breast, prostate, colon, ovarian, and pancreatic cancers[3, 7, 9-13], as well as haematological malignancies such as multiple myeloma, lymphomas, and leukaemias[14, 15]. While its role in tumour initiation remains debated[6, 9, 16-18], CA is consistently associated with aggressive disease features, including high-grade histology, poor prognosis, recurrence, and metastasis[3, 7, 9, 19]. Despite its clinical relevance, CA remains poorly characterised at scale due to the lack of robust, high-throughput tools capable of resolving centrosome phenotypes in complex tissue architecture.

Mechanistically, CA arises from both numerical and structural centrosome defects. Numerical amplification results from centriole overduplication, *de novo* centriole assembly, cytokinesis failure, mitotic slippage and cell–cell fusion[20-29]. Disruption of cell-cycle progression, such as prolonged G2 arrest, can trigger premature centriole disengagement and reduplication *via* PLK1 activation[30, 31]. Fragmentation of the PCM, driven by dysregulation of proteins such as pericentrin, γ-tubulin, PLK4, PLK1, and Aurora-A, also contributes to numerical CA[7, 32-34]. Structural CA, on the other hand, involves aberrant accumulation of PCM[4, 7, 9, 35, 36] or defects in centriole architecture[37-39]. Among these centriole architectural defects, over-elongation and fragmentation can lead to unstable centriole

structures and further overduplication[37], suggesting a mechanistic link between numerical and

structural centrosome aberrations. Yet, their differential contributions to cancer biology remain to be

determined.

Manual centrosome annotation remains the standard but is time-consuming, low-throughput, and

prone to observer bias. Semi-automated pipelines have emerged to address these limitations. For

example, CenFind—a deep learning pipeline based on SpotNet—accurately detects and counts

centrioles in cultured cells using immunofluorescence images but does not support structural

phenotyping[40]. Other machine learning–based approaches have quantified centriole number and

linked supernumerary centrioles to PCM expansion in breast cancer cells[41]. Semi-automated

frameworks have also been developed for centrosome quantification in human breast histological

sections, including a HistoQuest-aided method detecting structural CA[42] and an IMARIS-based

pipeline integrating both structural and numerical CA[7, 43]. Similarly, a 3D imaging-based pipeline

quantified structural centriole abnormalities across cancer types[37]. However, these approaches

require manual curation and offer only moderate throughput. A recent high-throughput platform using

a Harmony™ software-based framework revealed heterogeneous CA phenotypes in ovarian cancer

tissues[44], yet lacked single-cell resolution and subtype discrimination. Moreover, most existing tools

are tailored to immunofluorescence imaging and are not compatible with standard chromogenic

immunohistochemistry workflows used in clinical pathology, limiting their diagnosis and translational

utility.

To address these limitations, we developed CenSegNet (Centrosome Segmentation Network), a

versatile deep learning framework for fully automated, pixel-level detection and segmentation of

centrosomes in both immunohistochemistry and immunofluorescence images at single-cell

resolution. CenSegNet integrates three state-of-the-art models: Ultralytics YOLOv11, a recent

evolution of the You Only Look Once family optimised for fast and accurate performance[45], U-Net,

an encoder–decoder convolutional network designed for precise pixel-wise segmentation[46], and

StarDist for shape-aware instance segmentation pipeline that models objects as star-convex

polygons to improve instance separation in dense cellular contexts[47], enabling robust delineation of

epithelial cell boundaries within histopathological specimens. This systematically engineered

architecture supports multiscale analysis of centrosomal features in morphologically complex tissue

environments. We present a publicly accessible, expert-annotated dataset comprising human and

murine breast tissues and human mammary epithelial cell cultures (MECs). Implemented in Python

3.10 with a PyQt5-based graphical interface, CenSegNet enables streamlined data input, real-time

PyTorch-based inference, and modular extensibility. Benchmarking against expert annotations and

alternative models, CenSegNet achieves pathologist-level accuracy across imaging modalities.

Using CenSegNet, we perform the first high throughput, spatially resolved quantification of numerical

and structural CA in clinical breast carcinomas. Our analyses reveal that these CA subtypes are

mechanistically uncoupled and evolve along orthogonal spatial gradients: numerical CA

predominates in proliferative tumour cores, whereas structural CA accumulates at invasive margins,

reflecting distinct evolutionary pressures and microenvironmental cues. These spatial trajectories

correlate with histological grade, hormone receptor status, HER2 expression, nodal involvement,

and germline alterations, underscoring the role of centrosome dysregulation in driving intratumoral

heterogeneity and progression. Importantly, we validated CenSegNet in other human epithelial

tissues including kidney, colon, and appendix, demonstrating its generalisability and potential for

broad application in spatial pathology and organelle-level phenotyping across diverse healthy and

disease tissue contexts. To support widespread adoption, CenSegNet is released as an open-source

Python   library,   available   at   https://github.com/SKELab/CenSegNet/   and

https://zenodo.org/records/17131573.

Results

Development of CenSegNet for robust centrosome segmentation across imaging modalities

We generated tissue microarrays (TMAs) comprising 911 breast tissue cores from normal breast tissue, breast tumours, and adjacent areas, sampled from 127 patients enrolled in the ethically approved and clinically well-characterised BeGIN cohort (Investigating Outcomes from Breast Cancer: Correlating Genetic, Immunological and Nutritional Predictors), from University Hospital Southampton (UHS) (**Fig. 1a**; see Methods). Immunohistochemistry was performed using pericentrin antibody to label centrosomes, with haematoxylin counterstaining for nuclei (**Fig. 1a**). For training dataset construction, we manually annotated 14,679 centrosomes within 2,486 epithelial and 108 stromal compartments across 108 selected images (**Fig. 1a,** see Methods). To complement this, immunofluorescence was performed on human MECs and mouse mammary epithelium, labelling pericentrin and GT335 (centriole), with DAPI as nuclear counterstain. From this, an immunofluorescence training dataset was assembled comprising 1,285 annotated centrosomes from 143 cells from mouse tissue and 841 human MECs, revealing strong segmentation concordance between pericentrin and GT335 labelling [mouse tissue: R^2 = 0.9954; human MECs: R^2 = 0.9666 (MCF10A), 0.9085 (MCF10A-PLK4)] (**Fig. 1b, Supplementary Fig. 1a–f**).

Using these datasets, we initially benchmarked established segmentation models. U-Net, an encoder–decoder convolutional neural network optimised for pixel-wise segmentation[46], was selected for its extensive use in biomedical imaging. In our datasets, U-Net detected 82.98% of centrosomes in immunohistochemistry images and 97.6% in immunofluorescence, but despite achieving an overall F1 score of 0.85 in immunofluorescence, often the model either under-predicted or over-predicted centrosomes (**Supplementary Fig. 2a, b**). We next evaluated SegNet (**Fig. 1c**), another encoder-decoder model leveraging max-pooling indices for efficient upsampling[48]. It achieved 72.73% and 85% detection in immunohistochemistry and immunofluorescence, respectively, with a precision of 0.90 but reduced recall (0.75) in immunofluorescence and a

suboptimal overall F1 score of 0.68 in immunohistochemistry (0.82 in immunofluorescence)

(**Supplementary Fig. 2c, d**). We also evaluated DeepLabv3+, a semantic segmentation model

utilising atrous convolutions and atrous spatial pyramid pooling for multi-scale contextual

information[49]. Despite its proven success in complex image domains, DeepLabv3+ achieved only

63.25% and 75.1% detection rates in immunohistochemistry and immunofluorescence, respectively

(**Supplementary Fig. 2e, f**). These findings highlight the limitations of conventional segmentation

pipelines when applied to structurally heterogeneous tissues.

To overcome these constraints, we developed CenSegNet, a novel modular framework that

integrates object robust detection and segmentation (**Fig. 1c**). Rather than relying on whole-slide

segmentation, CenSegNet focuses on single-cell regions while modelling the entire ROI, explicitly

integrating object detection with region-based segmentation in a three-step architecture (**Fig. 1c**).

This design enables a more comprehensive capture of centrosome spatial distributions and

morphological features, addressing key challenges such as occlusion, small object detection, and

the disambiguation of overlapping structures. The framework, first employs YOLOv11[45], a state-of-

the-art object detection model, to identify potential centrosome candidates. YOLOv11 comprises a

convolutional backbone for feature extraction, a neck for multi-scale aggregation, and a head for

classification and localisation. We fine-tuned YOLOv11-seg[45, 50], a segmentation-optimised variant

to further enhance detection accuracy, and applied a range of data augmentations—including hue,

saturation, and value (HSV) adjustments, as well as geometric transformations such as translation,

scaling, shearing, and horizontal flipping—to enhance generalisability (**Fig. 1c**; see Methods). To

refine segmentation, we integrated U-Net in the second step. For all detected centrosomes, we

extracted centrosome-centred patches (256 × 256 pixels, with 40-pixel padding) as input to a U-Net

skip-connected encoder-decoder architecture for precise delineation. These patches underwent

similar augmentation strategies as in the detection step, ensuring consistency. To accurately quantify

centrosome numbers per cell, we incorporated StarDist[47] in the third step, a deep learning based

instance segmentation tool widely adopted for nuclear and cell boundary segmentation in biomedical

imaging (**Fig. 1c**). In our immunohistochemistry datasets, StarDist segmented 764,354 cells

(**Supplementary Fig. 1g, h**). To validate StarDist-aided cell-level centrosome assignment in

complex tissues, we analysed keratin 8 (KRT8)-labelled mouse immunofluorescence (620 cells) and

human immunohistochemistry (620 cells) tissues. StarDist-based segmentation demonstrated

strong concordance with KRT8 (immunofluorescence: R^2 = 0.99; immunohistochemistry: R^2 =

0.9005) (see **Supplementary Fig. 1e-i**).

Collectively, these results demonstrate that CenSegNet's multistep architecture effectively

overcomes the limitations of conventional segmentation approaches, delivering robust,

generalisable centrosome detection and quantification across diverse imaging modalities and

complex tissue contexts.

Validation of CenSegNet for scalable and high-precision centrosome segmentation

To evaluate CenSegNet's performance, we compiled independent test datasets of 25

immunohistochemistry TMA cores and 17 immunofluorescence images. Predicted centrosome

counts were compared to manually annotated ground truth, revealing strong correlations in both

immunohistochemistry (R^2 = 0.9999) and immunofluorescence (R^2 = 0.9873) (**Fig. 2a–c**).

CenSegNet consistently outperformed U-Net, SegNet, and DeepLabv3+ in segmentation accuracy

and boundary resolution across both modalities, with the YOLOv11 detection module significantly

enhancing overall performance (**Fig. 2a–c**). To further assess precision, we used an independently

annotated subset of 550 centrosomes from tumour regions in the immunohistochemistry test set.

Again, CenSegNet demonstrated superior pixel-level segmentation compared to all benchmarks

(**Fig. 2d**). F1 score analysis across modalities yielded a mean of 0.82 for CenSegNet, outperforming

U-Net (0.72), SegNet (0.68), and DeepLabv3+ (0.65) (**Fig. 2a–e**). We next benchmarked

performance on 6,475 expert-annotated centrosomes—921 from normal tissue, 2,694 from edge

regions, and 2,860 from tumour cores (**Fig. 2f**). These annotations, stratified by size (0.5–1.0 μm^2 to

>10.5 μm^2), were compared with the full cohort of 333,148 automatically segmented centrosomes

(0–0.5 µm^2 to >12.0 µm^2)—10,834 from normal tissue, 91,803 from edge tissue, and 230,511 from

tumour tissue (**Fig. 2g**). Both datasets revealed similar size distributions, with the majority falling

within 2.0–3.0 µm^2 and a peak at 2.0–2.5 µm^2. Normal tissue showed a higher proportion of 1.0–2.0

µm^2 centrosomes, while edge regions had more large centrosomes than either normal or tumour

cores. Using a previously established threshold of 6.5 µm^2-size to define structural CA[42] (see

Methods), both datasets confirmed that centrosomes exceeding this size were absent in normal

tissue. Together, these results establish CenSegNet as a scalable, high-precision tool for

centrosome segmentation, with superior performance across image modalities, tissue

compartments, and size distributions—supporting its utility for high-throughput analysis in complex

tissue environments.

CenSegNet enables spatial profiling of CA subtypes and clinical correlates in breast cancer

To dissect the contribution of numerical (Num CA) and structural (Stru CA) CA in breast cancer, we

applied CenSegNet to multiplexed TMA data. Num CA and Stru CA were not correlated within either

tumour edge or core regions (**Supplementary Fig. 3a**), suggesting mechanistic independence. At

the single-cell level, increasing centrosome number per cell was associated with a reduction in the

size of individual centrosomes (**Supplementary Fig. 3b**), which become smaller, more uniform in

cells containing >4 centrosomes—implying a compensatory constraint on total centrosome volume.

These observations suggest that Stru CA and Num CA represent orthogonal axes of centrosome

dysregulation. In contrast, Num CA levels correlated positively between edge and tumour

compartments (R^2 = 0.4857; **Supplementary Fig. 3c**), indicating progressive numerical amplification

from tumour margins inward. This was reflected in an increasing proportion of cells with ≥4

centrosomes from normal tissue to the edge and tumour core (**Fig. 3a**). Stratifying patients into four

CA phenotypes (Stru$^-$Num$^-$, Stru$^+$Num$^+$, Stru$^+$Num$^-$, Stru$^-$Num$^+$) revealed that CA was widespread,

detected in 89.8% of edge and 95% of tumour regions (**Supplementary Fig. 3d**). Notably, 73% of

tumour regions exhibited both Stru CA and Num CA, representing a ~19.3% increase compared to

edge regions (**Supplementary Fig. 3d**). While the Stru$^+$Num$^-$ group accounted for 18.4% of edge

regions but only 2% of tumour regions, the opposite trend was observed for the Stru$^-$Num$^+$ group

(10.2% *versus* 20%, respectively). These findings allow us to conclude that structural and numerical

centrosome aberrations are mechanistically uncoupled and highlight a spatial shift from centrosome

enlargement at the tumour edge to centrosome accumulation in the tumour core.

To investigate how CA correlates with ages, we stratified patients into four groups: <50, 50–60, 60–

70, >70 years. Num CA was elevated in patients aged 50–70 years across edge and tumour regions

(**Fig. 3a, b**). The frequency of patients with detectable Num CA and Stru CA at the tumour edge

markedly decreased in the 50–70 age group, resurging in patients over 70 years (**Fig. 3c**). Although

fewer 50 to 60-year-old patients exhibited detectable CA at the edge, those that did had high levels

of both CA subtypes (**Fig. 3d, e**). CA levels in tumour cores were relatively stable across age.

Notably, normal tissues exhibited age-dependent increases in centrosome size, especially in

patients over 70 (**Fig. 3f, g**), consistent with previous reports linking age to centrosome expansion

via cumulative DNA damage[51]. In edge and tumour regions, centrosome size decreased between

ages 50–70 before rising again in patients over 70 (**Fig. 3h, i**), while centrosome numbers followed

the opposite trend, increasing in the 50–70 group before decreasing in the oldest cohort (**Fig. 3j**).

This dynamic was mirrored by a higher proportion of cells with three centrosomes or more in the 50–

60 age group (**Fig. 3k**). Finally, we did not observe significant differences in age distribution across

the four CA-defined patient groups (Stru$^-$Num$^-$, Stru$^+$Num$^+$, Stru$^+$Num$^-$, and Stru$^-$Num$^+$)

(**Supplementary Fig. 4a**). Thus, while age alone may not initiate CA in breast cancer, it modulates

its spatial distribution and severity—particularly by promoting numerical amplification in the 50–70

age group and structural amplification in patients over 70.

We next evaluated associations between CA subtypes and clinicopathological features. Tumours

lacking CA (Stru$^-$Num$^-$) displayed smaller total and invasive areas in the tumour core relative to the

edge (**Supplementary Fig. 4b, c**), a pattern absent in other CA groups. Structural and numerical

CA prevalence increased with histological grade and nodal involvement (**Fig. 4a, b**). Stru CA was

inversely associated with tumour size across tumour (T) stages both edge and tumour regions, while

Num CA displayed region-specific variation (**Fig. 4c**). Stru CA and Num CA were more frequent at

the edge in invasive ductal carcinoma, but more abundant in the core of mixed-subtype tumours

(**Fig. 4d**). To further assess CA heterogeneity, patients were classified by composite CA defects into

Stru^{high}Num^{high}, Stru^{high}Num^{low}, Stru^{low}Num^{low}, Stru^{low}Num^{high} CA groups (**Fig. 4e, f**). CA burden was

not age-associated (**Fig. 4g**), and distinct CA subtypes were linked to tumour behaviour. At the edge,

Stru^{high}Num^{low} and Stru^{low}Num^{high} CA tumours were associated with greater nodal involvement than

either the Stru^{high}Num^{low} or Stru^{low}Num^{high} CA groups (**Fig. 4h**), whereas no such differences were

observed across CA groups in the tumour core. Stru^{low}Num^{high} CA tumours exhibited the smallest

overall size, while the largest tumours were enriched in Stru^{high}Num^{low} (edge) and Stru^{high}Num^{high}

(tumour) CA groups (**Fig. 4i, j**). Stru^{high}Num^{low} CA tumours grew more aggressively at the edge, while

Stru^{high}Num^{high} CA tumours expanded predominantly in the core (**Fig. 4k**). While Stru^{low}Num^{low} CA

was the most frequent subtype among large tumours across both regions, neither Num CA (in tumour

cores) nor Stru CA (at the edge) alone stratified tumour size (**Fig. 4l**). About 40% of tumours within

the Stru^{high}Num^{high} CA group exhibited nodal involvement greater than N1 in the core (**Fig. 4m**),

consistent with increased metastatic potential. Many tumours with Stru^{low}Num^{low} CA status at the

edge required nodal clearance, suggesting local aggressiveness independently of global CA burden

(**Fig. 4n**). Identified germline variants were exclusively observed in the Stru^{high}Num^{high} CA group (**Fig.

4o**), consistent with associations between CA and BRCA1-driven genomic instability[52]. We also

evaluated the relationship between CA subtypes and body composition. Tumours with Stru^{low}Num^{high}

CA in the core were associated with increased fat-free mass index (FFMI), (**Supplementary Fig.

5a**), while fat mass index (FMI), waist circumference, and weight showed no differences across

groups or regions (**Supplementary Fig. 5b-d**). Taller patients more frequently exhibited

Stru^{low}Num^{low} CA status in the edge regions, though this association was not observed in cores

(**Supplementary Fig. 5e**). Collectively, these findings indicate that spatial patterns of CA are linked

to tumour clinical parameters. Stru^{high}Num^{high} CA tumours define a high-risk subgroup marked by

large size, nodal spread, and genomic alterations, while Stru^{low}Num^{low} CA tumours—despite low CA

burden—can display local aggressiveness.

Single-cell analyses revealed further divergent CA subtype dynamics. At the edge regions, Num CA

decreased with increasing tumour size (**Supplementary Fig. 6a**), while Stru CA was enriched in

poorly differentiated but smaller tumours (**Supplementary Fig. 6b**). In the core, the proportion of

cells harbouring either CA subtype increased with both total and invasive tumour size

(**Supplementary Fig. 6c, d**). Num CA showed no significant differences across histological tumour

grades, although it was more prevalent in well-differentiated tumours (**Supplementary Fig. 6c**). Stru

CA levels were enriched in poorly differentiated tumours (**Supplementary Fig. 6d**). No significant

subtype-specific CA differences were observed across histological subtypes (**Supplementary Fig.**

6a-d), but cells from mixed tumours harboured more centrosomes in both compartments

(**Supplementary Fig. 7**), highlighting elevated Num CA as linked to increased tumour heterogeneity.

Together, these findings indicate a dynamic evolution of CA during tumour progression, with early-

stage tumours characterised by numerical defects and advanced tumours accumulating structural

abnormalities.

Finally, we explored CA patterns in the context of hormone receptor status, a key clinical determinant

in breast cancer[53, 54]. Stru CA levels varied by receptor status in both edge and tumour core

compartments (**Supplementary Fig. 8a**). HER2$^-$ tumours had the highest Stru CA at the edge, while

PR$^-$ and ER$^-$ tumours had the lowest. In contrast, ER$^-$ tumours displayed elevated Stru CA in the

core, with HER2$^+$ tumours showing the lowest levels. Num CA also showed compartment-specific

trends: HER2$^+$ tumours exhibited high Num CA at the edge but low levels in the core; ER$^-$ tumours

showed the inverse pattern (**Supplementary Fig. 8b**). Despite broadly similar spatial trends

between Stru CA and Num CA across receptor-defined subtypes, HER2$^+$ tumours emerged as an

exception (**Supplementary Fig. 8c, d**). HER2^{moderate} (2+) tumours had significantly lower Num CA

levels than HER2[high] (3+) tumours in both compartments (**Supplementary Fig. 8e**), suggesting

HER2 dosage impacts centrosome number. HER2[+] tumours were overrepresented in discordant CA

phenotypes—Stru[low]Num[high] (28.6%) and Stru[high]Num[low] (35.7%)—at the edge (**Supplementary Fig.**

8f), indicating HER2 may differentially regulate centrosome structure and number depending on

spatial context. This distribution was not observed in the tumour core. These observations point to

spatially resolved, hormone receptor-specific influence of CA patterns, particularly via HER2

signalling, in a microenvironment-dependent manner.

CenSegNet reveals CA patterns linked to tumour subtype and progression

Using CenSegNet-derived spatial quantifications, we compared dynamic shifts in Stru CA and Num

CA between tumour edges and cores. Patients were classified based on whether each CA subtype

was more abundant in the tumour core (T>E) or at the edge (E>T). Baseline clinicopathological

features were similar between spatial groups (**Supplementary Fig. 9**). Stru CA[T>E] tumours were

associated with lobular histology (15.8%), lower histological tumour grade (26.3% grade 3), and

lower HER2[+] prevalence (15.8%). In contrast, Stru CA[E>T] tumours more frequently displayed high

tumour grade morphology (59.1% grade 3), higher HER2[+] status (immunohistochemistry 3+ or FISH-

confirmed) (31.8%), and higher differentiation (**Supplementary Fig. 10a, b**). Single-cell analysis

revealed that in both Stru CA spatial groups, Num CA–positive cells consistently contained more

centrosomes in the core than at the edge (**Fig. 5a, b**), indicating a conserved numerical asymmetry.

Num CA spatial prevalence was also associated with aggressiveness: Num CA[T>E] tumours were

more often grade 3 (40.6%), had greater nodal involvement (31.2%), and lower HER2[+] frequency

(12.5%) compared with Num CA[E>T] tumours (28.6%, 7.1%, and 35.7%, respectively;

Supplementary Fig. 10c, d). Mixed histology was more common in Num CA[T>E] tumours (12.5%),

whereas Num CA[E>T] tumours were associated with lobular carcinomas (28.6% *versus* 6.3%), a

subtype typically linked to slower growth and smaller size. Across both Num CA[T>E] Num CA[E>T]

groups, centrosomes were significantly larger at the edge than in the core. Notably, Num CA–positive

cells with larger centrosomes harboured fewer of them (**Fig. 5c, d**), revealing a robust inverse

relationship between centrosome size and number across all spatial groups (**Fig. 5e**). These findings

highlight distinct spatial CA patterns associated with tumour subtype and behaviour. Tumours

displaying a shift from fewer, larger centrosomes at the periphery to smaller, more numerous

centrosomes in the core exhibit features of increased aggressiveness. These spatial patterns,

resolved through CenSegNet, provide a proxy for tumour heterogeneity and may reflect the

evolutionary trajectory of CA during tumour progression.

CenSegNet integration and accessible deployment for high-throughput centrosome analysis

To support broad adoption and integration into diverse analytical workflows, we provide CenSegNet

as both a lightweight application programming interface (API) and an interactive graphical user

interface (GUI) (**Fig. 6a, b**). The GUI comprises three modules: a data upload panel for inputting

whole-slide or high-resolution images; a prediction module for adjustable inference parameters

including probability thresholds and region selection for optimised segmentation of centrosomes and

epithelial compartments; and an export tool for structured outputs. Post-inference, users can retrieve

per-cell pixel-resolved size estimates, spatial coordinates, and centrosome counts within tissue

context (**Fig. 6b)**. The interface supports batch processing and accepts both immunohistochemistry

and immunofluorescence formats. Benchmarking shows that CenSegNet processes a 6,000 ×

6,000-pixel immunohistochemistry image or a 2,048 × 2,048-pixel immunofluorescence image in <1

min on standard GPUs, enabling rapid, scalable analysis across large datasets.

Discussion

We introduce CenSegNet, a modular deep learning framework for quantification of centrosomes in

epithelial tissues at spatial and single-cell resolution. Unlike previous approaches that treat

morphological analysis and segmentation as separate tasks, CenSegNet integrates centrosome

detection and phenotyping, nuclear localisation, and epithelial boundary inference into a unified,

multichannel pipeline. This enables context-aware segmentation of both structural (Stru CA) and

numerical (Num CA) abnormalities across diverse imaging modalities. CenSegNet outperforms

established models—including U-Net, SegNet, and DeepLabv3+—in accuracy, generalisability, and

morphological fidelity, particularly in densely packed or morphologically heterogeneous tissue

regions. The framework comprises three specialised modules: a YOLOv11-based detector trained

on over 15,000 annotated centrosomes for robust localisation; a U-Net model for precise

segmentation of centrosome area and morphology; and a StarDist-based cell segmentation module

optimised for delineating epithelial boundaries in complex tissue architectures. This modular

integration, coupled with uncertainty-aware postprocessing, enables systematic and standardised

phenotyping of CA subtypes at single-cell and spatial resolution. Applied to 911 sample cores from

127 patients, CenSegNet-based profiling of over 330,000 centrosomes reveals previously

uncharacterised spatial trajectories and clinical correlates of Stru CA and Num CA, uncovering their

mechanistic uncoupling, age-dependent modulation, and associations with tumour progression,

hormone receptor status, HER2 expression, and genomic alterations.

Computational tools have been developed previously for CA quantification in epithelial cells. A semi-

automated machine learning pipeline quantified PCM defects and numerical CA in

immunofluorescence images of normal and breast cancer cells[41], but the method offers limited

spatial resolution and required extensive manual curation. Another semi-automated approach linked

centrosome size and number to chromosomal instability in human breast cancer tissues[17], yet lacked

single-cell resolution and scalability. Another pipeline assessed centriole number and length,

revealing structural defects arising from fragmentation and ectopic procentriole formation[37], but

without spatially resolved quantification. An automated detection algorithm for high-throughput

mapping of CA in ovarian cancer tissues, identified heterogeneous CA phenotypes associated with

chromosomal instability and chemotherapy resistance[44]. However, this method did not distinguish

between structural and numerical CA or provide single-cell semantic segmentation. CenSegNet

addresses these limitations by integrating cellular, spatial, and clinical dimensions of CA across

large-scale tissue cohorts. In doing so, it offers new insights into centrosome biology and the

functional relevance of CA heterogeneity in breast cancer, with implications for risk stratification and

precision oncology. Beyond breast cancer, CenSegNet has demonstrated generalisability across

diverse epithelial tissues—including kidney, colon, and appendix—characterised by high stromal

content and architectural complexity (**Supplementary Fig. 11**). To facilitate broad adoption, we

provide CenSegNet as an open-source GUI enabling scalable extraction of structured, spatially

anchored centrosome metrics. This will allow researchers and clinicians, regardless of computational

expertise, to integrate centrosome profiling into histopathological and biomarker discovery

workflows. Thus, CenSegNet extends beyond methodological innovation to practical application,

accelerating systematic investigation of centrosome biology across anatomically and histologically

diverse tissues, and enabling the identification of CA-driven vulnerabilities with potential therapeutic

relevance.

Recent studies using a composite centrosome amplification score (CAS) that integrates both

numerical and structural abnormalities[7, 43], showed a progressive increase in CAS from normal

breast tissue to invasive carcinoma. However, it did not distinguish the individual contributions of

numerical and structural centrosome abnormalities. Our spatially resolved single-cell analysis

demonstrates that Num CA and Stru CA represent distinct phenotypic axes of centrosome

dysregulation. Although they frequently co-occur in tumour tissues, they are uncorrelated at both

tissue and single-cell levels and exhibit unique spatial distributions: Stru CA is enriched at tumour

edges, while Num CA predominates in tumour cores, suggesting that different regional pressures

drive centrosome overduplication *versus* structural enlargement. Single-cell data further uncover a

robust inverse relationship between centrosome number and size—cells with multiple centrosomes

tend to have smaller ones—indicating a compensatory constraint on total centrosome volume.

Consistent with this, centriole over-elongation can induce CA *via* fragmentation and ectopic

procentriole formation in breast cancer cells[37], yet not all centrioles within a cell exhibit these

changes, highlighting intra-cellular heterogeneity in elongation susceptibility. Collectively, these

findings support a model in which Num CA and Stru CA are mechanistically uncoupled, evolving

along orthogonal spatial gradients during tumour progression. This challenges the monolithic view

of CA and instead portrays it as a dynamic, regionally modulated process shaped by local

microenvironmental cues. The mechanistic decoupling of CA has important implications for

understanding its functional heterogeneity in cancer and underscores the need for spatially informed

biomarkers.

CenSegNet-based profiling reveals that aging exerts distinct and spatially patterned effects on Stru

CA and Num CA in breast cancer. Num CA peaks between ages 50 and 70—overlapping with the

menopausal transition and the most common window for breast cancer diagnosis[55-57]—whereas Stru

CA accumulates progressively after age 70, indicating a later-life trajectory of centrosome

architectural dysregulation. While CA overall increases with age—including in normal tissues—our

data suggest that age is not a deterministic initiator but rather a factor that modulates the magnitude

and spatial distribution of CA subtypes. In mid-life patients, Num CA is preferentially enriched in

tumour cores—regions typically characterised by high proliferation—whereas Stru CA in older

individuals extends more diffusely, often into tumour margins, likely reflecting age-associated

changes in epithelial architecture, tissue repair dynamics, and microenvironmental stress. A study in

prostate cancer has reported elevated CA in patients over 53 years of age[7]. CA increases with age,

in normal breast epithelial cells derived from individuals aged 20-80, treated with DNA damage-

inducing stimuli[51]. Chronic centrosome overduplication in aging mouse models of intestinal cancer

drives aneuploidy and spontaneous tumorigenesis, supporting a role for age-associated CA in early

malignant transformation[18]. Centrosome function deteriorates with age, evidenced by accumulation

of structural defects and impaired mitotic fidelity[58, 59]. Aging epithelial cells exhibit centrosome

fragmentation, aberrant centriole elongation, and altered PCM composition[59, 60], while broader

declines in DNA repair and chromosomal segregation fidelity likely contribute to a permissive

environment for CA[59]. Thus, aging tissues accumulating centrosome abnormalities, mitotic fidelity

defects, and weakened genomic surveillance, may be more vulnerable to CA-driven tumorigenesis.

Our study delineates the divergence of CA subtypes across the lifespan. The inverse scaling

between centrosome number and size supports the idea that Num CA and Stru CA act as

compensatory, rather than co-occurring, phenotypes. Notably, Stru CA in elderly patients is

frequently associated with hormone receptor–negative tumours, implicating centrosome structural

dysregulation in the biology of more aggressive or dedifferentiated late-life cancers. CA can induce

breast cancer cell dedifferentiation and intrinsically drive high-grade tumours[61]. Age remains a key

prognostic factor in breast cancer, influencing tumour subtype distribution, hormone receptor status,

and genomic instability[62, 63]. Together, these results establish the first clinically and spatially resolved

framework for understanding how aging modulates centrosome biology in human cancer. This

framework lays the foundation for developing age-stratified biomarkers and clarifies why specific CA

subtypes and their associated chromosomal instability may emerge more frequently or have greater

clinical impact at distinct stages of life.

The spatial heterogeneity of CA and its clinical relevance for cancer remain poorly defined[3, 7].

CenSegNet systematically maps Stru CA and Num CA across tumour compartments and stratifies

tumours into composite CA subtypes with distinct spatial, biological, and clinical profiles.

$Stru^{high}Num^{high}$ CA tumours are consistently associated with larger size, greater nodal involvement,

and germline genomic alterations. Elevated CA in both core and edge regions suggests a cellular

composition primed for proliferative expansion and invasive dissemination. $Stru^{low}Num^{low}$ CA

tumours often required nodal clearance, supporting evidence that even modest CA can drive

aggressive behaviour in permissive genomic contexts—particularly when p53 surveillance is

compromised[7, 18, 64]. CenSegNet also uncovers spatial discordance in CA subtypes, with

Stru[high]Num[low] and Stru[low]Num[high] CA profiles enriched at the invasive front. These spatial signatures

are associated with enhanced metastatic potential, consistent with models in which centrosome

abnormalities promote invasion through both cell-autonomous[65-67] and non-cell-autonomous[68-70]

mechanisms. Structural centrosome defects have been shown to drive cell extrusion, facilitating

invasion[67, 71]. Our findings corroborate longitudinal studies of tumour progression, such as in Barrett's

oesophagus, where CA appears early in premalignant lesions and expands with p53 loss[64],

supporting a role for CA in tumour initiation rather than as a mere by-product of transformation.

Similarly, CA was shown to increase from normal tissue to ductal carcinoma *in situ* (DCIS) to invasive

carcinoma, and to correlate with recurrence and poor prognosis[7, 61]. Finally, our observation that Stru

CA and Num CA spatial patterns are uncoupled from systemic physiological metrics reinforces CA

as a tumour-intrinsic hallmark. Together, these results advance our understanding of centrosome

biology in cancer and highlight the power of CenSegNet-driven integration of subcellular organelle

features into spatial pathology, with direct implications for clinical decision-making, prognostic

modelling, and therapeutic targeting.

By integrating CenSegNet-based centrosome phenotyping with spatially resolved hormone receptor

profiling, we uncover compartment-specific associations between CA subtypes and ER/HER2

expression. Num CA is selectively enriched in ER$^-$ tumours within the tumour core, whereas HER2$^+$

tumours display lower Num CA in the core despite elevated levels at the tumour edge. This spatial

divergence suggests that hormone receptor signalling modulates centrosome number and structure

in a regionally distinct manner. These observations corroborate previous studies linking CA to

hormone receptor status and tumour aggressiveness[37, 42, 66, 72] and support the hypothesis that ER

loss promotes CA through transcriptional or post-translational dysregulation of centriole biogenesis

pathways. Basal-like ER$^-$PR$^-$HER2$^-$ breast cancers—characterised by genomic instability and poor

prognosis[73], frequently exhibit high CA, often with centriole fragmentation and ectopic procentriole

formation driven by over-elongation[37]. These defects recruit excess pericentriolar material (PCM),

generating supernumerary or structurally abnormal MTOCs that drive mitotic errors and

chromosomal instability[74, 75]. In contrast, HER2$^+$ tumours are more frequently enriched in discordant

CA subtypes (Stru^{low}Num^{high} and Stru^{high}Num^{low}) at the tumour edge, suggesting that HER2 signalling

may differentially regulate centrosome number and structure depending on spatial context. This is

consistent with evidence that both PLK4 and AURKA expression—key regulators of centriole

biogenesis and maturation, respectively—is differentially influenced by HER2 status[72, 76]. Despite

high proliferative capacity, HER2$^+$ tumours exhibit lower overall CA, raising the possibility that these

tumours suppress CA to preserve mitotic fidelity or evade immune detection. Together, our findings

identify spatially distinct and mechanistically diverse relationships between hormone receptor

signalling and centrosome biology. ER loss is associated with elevated Num CA and mitotic instability

in tumour cores, while HER2 signalling appears to exert compartmentalised control over CA subtype

distribution. Spatially resolved centrosome profiling thus provides a framework for identifying

hormone-specific vulnerabilities that could inform targeted breast cancer therapies.

CenSegNet enables fine-grained stratification of tumours based on the relative abundance of

structural (Stru CA) and numerical (Num CA) centrosome abnormalities, revealing distinct spatial–

biological associations. Tumours with Stru CA enriched at the invasive edge (Stru CA$^{E>T}$) are more

proliferative, more frequently HER2$^+$, and exhibit higher differentiation, whereas those with Stru CA

more abundant in the core (Stru CA$^{T>E}$) are more commonly associated with lobular histology and

slower growth. Single-cell analyses further show that tumours with grade 3 histology, nodal

involvement, and mixed subtypes—hallmarks of aggressive disease—are more prevalent in the Num

CA$^{T>E}$ subgroup, while Num CA$^{E>T}$ tumours are enriched for lobular carcinomas, typically linked to

indolent behaviour[73]. These spatially resolved patterns corroborate previous reports suggesting that

numerical centrosome abnormalities increase with tumour progression and are more frequent in

aggressive basal-like carcinomas[7, 37]. Our single-cell data also reveal an inverse relationship

between centrosome number and size—where edge regions harbour fewer but larger

centrosomes—supporting a model in which Stru CA at the periphery primes cells to acquire invasive

behaviours, while Num CA in the core drives proliferation and genomic instability. This dynamic

interplay likely reflects microenvironmental influences on CA trajectories during tumour progression.

These findings extend the "CA set point" concept[77], which postulates that tumours maintain a

context-dependent equilibrium of CA phenotypes to balance proliferation, invasion and survival.

CenSegNet-based spatial profiling demonstrates that CA is not only subtype-specific but also

spatially regulated, providing new insights into the architectural evolution and heterogeneity of breast

cancer.

In summary, CenSegNet delivers the first fully integrated and spatially resolved framework for

profiling CA at single-cell resolution across large-scale human cancer tissues. By enabling precise,

high-throughput quantification of both numerical and structural CA phenotypes, CenSegNet

uncovers distinct mechanistic, temporal, and spatial trajectories of centrosome dysregulation in

breast cancer. The discovery that numerical and structural CA are decoupled numerical and

structural CA are decoupled—not only in cellular architecture but also in their associations with

clinical features such as tumour grade, hormone receptor status, germline mutation, and patient

age—advances our understanding of how centrosome abnormalities contribute to intratumoral

heterogeneity, tumour progression, with considerable implications for therapeutic resistance.

Importantly, the identification of compensatory dynamics between centrosome number and size, and

their divergent distributions across tumour cores and margins, points to context-specific roles in

modulating local tumour ecology and genomic instability. These insights challenge the longstanding

view of CA as a uniform driver of malignancy, instead positioning CA subtypes as distinct functional

modules in tumour evolution. CenSegNet thus provides a foundation for developing CA-based

biomarkers to stratify patients by tumour subtype, age, and aggressiveness, and opens opportunities

for therapies targeting CA-driven vulnerabilities. Given the availability of PLK4, AURKA, and HSET

inhibitors[78-82], spatial CA maps could guide personalised strategies, particularly in tumours with

discordant amplification phenotypes. Future studies integrating CenSegNet-based CA profiling with

transcriptomic and proteomic analyses will be essential to uncover the molecular drivers of spatial

CA dynamics and clarify their roles in tumour progression, metastatic dissemination, and therapy

resistance.

Methods

Ethics and human breast tissues

The study participants were a subgroup of women diagnosed with early breast cancer who were

recruited to a single-centre prospective observational cohort study at University Hospitals

Southampton, "Investigating Outcomes from Breast Cancer: Correlating Genetic, Immunological and

Nutritional Predictors (BeGIN)[83, 84]." All procedures performed in studies involving human participants

were in accordance with the ethical standards of the institutional and/or national research committee

and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

All participants in BeGIN gave written informed consent. The research ethics committee approved

the study (Research Ethics Committee (REC) - Cambridgeshire and Hertfordshire reference number:

14/EE/1297). Women were eligible for the BeGIN study if they were aged >18 years and diagnosed

with invasive breast cancer or DCIS at University Hospital Southampton after May 2015. Linked

anonymised patient information, including patient characteristics, tumour characteristics and clinical

management, were extracted from the hospital electronic record system. Body composition

parameters were measured using Bioelectrical Impedance Analysis (BIA) with a phase-sensitive, 8-

electrode device (Seca mBCA515)[85]. To conduct this study, 911 cores from 127 breast cancer

patients were used. The TMAs were constructed from formalin fixed paraffin embedded (FFPE)

histopathology tissue blocks from surgical treatment surplus to diagnostic requirements. Colon,

kidney, and appendix tissue were incorporated into breast tissue TMAs to facilitate orientation during

sectioning and analysis. Colon and kidney samples consisted of histologically normal tissue, the

status of which was independently verified by a board-certified pathologist. A summary of the

clinicopathological information linked to the human breast samples used in this study is included in

Supplementary Table 1. Anonymous data from the BeGIN study is available for request to

researchers who provide a completed Data Sharing request form that describes a methodologically

sound proposal, for the purpose of the approved proposal. Proposals will be reviewed by the study

steering committee. Data will be shared once all parties have signed relevant data sharing

documentation, covering the study steering committee conditions for sharing and if required, an

additional Data Sharing Agreement from the Sponsor.

Ethics and mice

BALB/c HER-2/neu transgenic mice (referred to as BALB-NeuT)[86] carrying the transforming rat *Her-*

2/neu oncogene under control of a MMTV-LTR were used. All experimental procedures involving

mice were approved by the University of Southampton Local Ethics Committee and registered with

the Ethics and Research Governance Online II (ERGO II; ID: 65385). All animal work was conducted

in accordance with UK Home Office regulations, adhering to the principles of the 3Rs (Replacement,

Reduction, Refinement) and the ARRIVE (Animal Research: Reporting of In Vivo Experiments)

guidelines to minimise animal suffering throughout the study. Mice were housed in a specific

pathogen-free (SPF) facility under controlled environmental conditions, including regulated

temperature and humidity, with a 12-hour light/dark cycle. Animals had ad libitum access to standard

chow and water.

Cell culture

MCF10A is a non-transformed human mammary epithelial cell line (ATCC® CRL-10317). The

MCF10A-PLK4 cell line is a genetically engineered derivative of MCF10A that enables inducible

overexpression of Polo-like kinase 4 (PLK4), a master regulator of centrosome duplication whose

upregulation induces centrosome amplification[66]. The MCF10A-PLK4 cell line was kindly provided

by Dr. Susana Godinho (The Barts Cancer Institute, Queen Mary University of London). Both

MCF10A and MCF10A-PLK4 cells were cultured in DMEM/F12 medium (Invitrogen), supplemented

with 10% donor horse serum (Gibco, 31331028), 20 ng/ml human epidermal growth factor (EGF;

Sigma, E9644), 10 µg/ml insulin (Sigma, I1882), 100 µg/ml hydrocortisone (Sigma, H0888), 1 ng/ml

cholera toxin (Sigma, C8052), and 50 U/ml penicillin with 50 µg/ml streptomycin (Life Technologies).

Cells were maintained at 37 °C in a humidified incubator with 5% CO_2. To induce PLK4

overexpression, cells were treated with doxycycline (Sigma, D9891) at 2 µg/ml for 48 hours.

Tissue microarray construction and immunohistochemistry

Tissue microarrays (TMAs) were constructed using formalin-fixed, paraffin-embedded (FFPE) tissue

samples obtained following breast cancer surgery from 127 patients diagnosed with primary invasive

breast carcinoma at University Hospital Southampton between July 9, 2015, and January 31, 2019,

and participating in the BeGIN study. All patients underwent standardised treatment at a single

institution, consisting of surgery, followed by adjuvant treatments according to local and national

protocols. Pathological evaluation of hormone receptor and HER2 expression was conducted

according to established national and international guidelines[87, 88]. Immunohistochemistry was used

to determine hormone receptor status, and in situ hybridization (ISH) was employed to confirm HER2

positivity for tumours with an IHC score of 2+. All procedures were performed within the standard

clinical diagnostic pathway. A total of 911 sample cores were systematically sampled from three

distinct, pathologically classified regions for each patient: tumour tissue (Tumour), tumour margin

(Edge), and tumour-free tissue (Normal). Each patient represented by three technical replicate cores

from Tumour, Edge, and Normal regions, were procured for analysis. Each 0.6 mm in diameter and

5 µm in thickness were extracted from formalin-fixed, paraffin-embedded specimens and arrayed

into recipient tissue microarray (TMA) blocks. This tri-regional, triplicate-core sampling strategy was

designed to provide a comprehensive and robust representation of the tissue heterogeneity within

and around the tumour microenvironment. TMA sections were mounted onto TOMO® adhesion

microscope slides. Immunohistochemistry was performed using the Dako Autostainer Link 48

automated platform. Endogenous peroxidase activity was quenched using EnVision FLEX blocking

reagent (Dako), followed by a 30-minute incubation with the primary antibody against pericentrin

(1:500 dilution; Abcam, ab4448). Signal amplification and enzymatic detection were achieved using

EnVision FLEX HRP (Dako, 20 minutes) and Rabbit Link (Dako, 15 minutes). Slides were

counterstained with haematoxylin following three 5-minute washes in 3-amino-9-ethylcarbazole

(AEC).

Assessment of hormone receptor status

ER or PR expression were evaluated by immunohistochemistry and scored using the Allred system

(range 0–8). Tumours with an Allred score ≥ 3 were classified as ER or PR positive. HER2 status

was determined by immunohistochemistry and scored as 0, 1+, 2+, or 3+. Tumours with a score of

3+ were classified as HER2 positive. Cases with a score of 2+ were considered equivocal were

further assessed by fluorescence in situ hybridization (FISH). Tumours with FISH amplification were

designed as HER2 positive, defined as a HER2/CEP17 ratio ≥ 2.0 or average HER2 copy number ≥

6.0 signals per cell.

Immunofluorescence

The following primary antibodies were used: anti-GT335 (1:800; Adipogen, AG-20B-0020-C100),

anti-pericentrin (1:250; Abcam, ab4448), and anti-keratin 8/18 (KRT8/18; 1:300; Origene, BP5007).

Secondary antibodies (Life Technologies) included goat anti-mouse (A-32723), anti-rabbit (A-11037

and A-11008), and anti-guinea pig (A-21450), each conjugated to Alexa Fluor 488, Alexa Fluor 594,

or Alexa Fluor 647, and used at a final concentration of 5 µg/ml.

OCT-embedded mammary gland sections (30 µm thick) from BALB-NeuT mice were cryosectioned,

air-dried for 30 minutes, and fixed in 4% paraformaldehyde (PFA) for 20 minutes at room

temperature. Sections were permeabilised for 45 minutes with 0.1% Triton X-100 in PBS, then

blocked for 2 hours in a solution containing 2% bovine serum albumin (BSA), 5% foetal bovine serum

(FBS; Gibco), and 0.1% Triton X-100 in PBS. Sections were incubated overnight at 4 °C with primary

antibodies against pericentrin, GT335, and KRT8, followed by washing and incubation with the

appropriate secondary antibodies for 2 hours at room temperature. Nuclei were counterstained with

DAPI using Fluoroshield mounting medium (Sigma, F6057).

MCF10A and MCF10A-PLK4 cells were fixed in anhydrous methanol at −20 °C for 10 minutes,

followed by permeabilisation with 0.1% Triton X-100 in PBS for 2 minutes. Cells were then washed

three times for 5 minutes each with 0.1% Triton X-100 in PBS. Blocking was performed using 3%

BSA in 0.1% Triton X-100 in PBS for 1 hour at room temperature. Cells were incubated overnight at

4 °C with primary antibodies against pericentrin and GT335. After washing, cells were incubated with

the appropriate secondary antibodies for 1 hour at room temperature and counterstained with DAPI

using Fluoroshield.

Microscopy and image annotation

Immunohistochemistry images were acquired using a Zeiss Axio Imager Z1 upright microscope

(Zeiss), equipped with an AxioCam MRc5 colour camera. Image capture was performed using Zeiss

ZEN imaging software, following a predefined whole-slide brightfield scanning protocol (Whole Slide

[WS] Brightfield [BF] fold-light) with a 20× objective lens. This configuration enabled high-resolution

imaging of tissue sections with consistent illumination and contrast across samples.

Immunofluorescence images were captured using an inverted Leica STELLARIS 5 laser scanning

confocal microscope (Leica Microsystems), equipped with a 40× oil immersion objective (HC PL

APO 40×/1.30 Oil CS2). Z-stacks were acquired at 16-bit depth with a 0.2 µm step size, using fields

of view (FOVs) ranging from 1024 × 1024 to 2048 × 2048 pixels. All cells from all FOVs obtained

during the experiments were included in the evaluation of model performance. Image processing

was performed using Fiji software (https://imagej.net/software/fiji/)[89].

Manual annotations were performed by Jiaoqi Cheng (see **Supplementary Fig. 12**), who received

training and supervision from Constantinos Savva. Centrosome size quantification in

immunohistochemistry images was performed using calibrated spatial resolution, wherein each pixel

corresponded to an area of 0.0483 µm^2. For each centrosome, pixel-level segmentation masks were

used to compute total pixel occupancy, which was then multiplied by the calibrated pixel area to

derive centrosome size in µm².

To generate the immunohistochemistry training dataset, images from 78 patients were included,

comprising human breast cancer tissue, normal breast tissue, normal liver, and normal kidney

samples. Within these images, annotations were made for centrosomes (n = 14,679), epithelial

compartments (n = 2,486), and stromal compartments (n = 108), using QuPath v0.5.1

(https://qupath.github.io/)[90]. All centrosome annotations were performed under 200× magnification

for each region of interest (ROI). The manual annotation process required over 200 person-hours.

Annotations were exported as GeoJSON files.

For the immunofluorescence training dataset, only 15 high-resolution image z-stacks of MCF10A

and MCF10A-PLK4 cell lines were required. Annotation was performed using Cellpose 3.0[91, 92].

Datasets were first maximum-intensity projected, then split into contrast-adjusted single-channel

images, with boundaries defined by edge features visible in the blue channel. The manual annotation

process took approximately 16 person-hours. All annotations were exported as SVG files.

Image processing and model training

Image processing. For immunohistochemistry image processing in the CenSegNet pipeline, raw

microscopy images were pre-processed to enhance contrast and suppress background noise,

thereby improving the visibility of cellular structures. Each image was cropped into overlapping

patches of 256 × 256 pixels with a stride of 300 pixels. Patches exhibiting artefacts or poor quality

were manually excluded to ensure dataset integrity. The final immunohistochemistry dataset

comprised 1,122 annotated patches, each containing labelled information on centrosome location

and size. For immunofluorescence images, RGB channels were converted to greyscale to reduce

dimensionality and simplify the training process. This transformation allowed the model to remain

invariant to colour information while improving computational efficiency.

Model architecture and training. CenSegNet employs a modular three-step architecture

comprising approximately 40 million trainable parameters. The first step consists of a detection head

that localises candidate ROIs, while the second stage performs fine-grained segmentation within

these ROIs to achieve precise spatial delineation. This decoupled design enables task-specific

optimisation and improves memory efficiency and training stability. In the detection stage, we fine-

tuned YOLOv11-seg model[45] (https://github.com/ultralytics/ultralytics) using our own training set,

guided by a composite loss function comprising box loss, segmentation loss, classification loss, and

distribution focal loss. These components respectively optimise object localisation, foreground-

background separation, class prediction, and robustness to complex spatial distributions. Training

was performed using the AdamW optimiser (learning rate = 0.002, momentum = 0.9) with a batch

size of 16 over 300 epochs. The segmentation step 2 employed a U-Net architecture[46] with three

input channels (RGB) and one output channel. Given the small size of centrosomes, segmentation

was formulated as a binary classification task. The model was trained for 100 epochs using the

Binary Cross Entropy with Logits loss function (BCEWithLogitsLoss). Optimisation was performed

using RMSprop (learning rate = 1×10^{-4}, weight decay = 1×10^{-8}, momentum = 0.9), which stabilised

training and improved convergence for small target structures. Training was conducted in Python

(v3.10) using PyTorch (v2.1). We applied data augmentations, such as HSV colour jittering and

geometric transformations, which can be easily implemented using the torchvision.transforms

module in PyTorch. The detection head was trained independently for approximately 24 hours,

followed by an additional 12 hours of training for the segmentation head. All experiments were run

on four NVIDIA A100 GPUs (40 GB VRAM each) using PyTorch's Distributed Data Parallel (DDP)

framework with NVIDIA Collective Communications Library (NCCL) backend for gradient

synchronisation. Input batches were evenly partitioned across GPUs, with local gradient computation

and synchronisation via NCCL's optimised collective communication, achieving near-linear scaling

in throughput. Batch sizes were dynamically adjusted to maximise GPU utilisation while maintaining

training stability.

Comparative segmentation models. To benchmark CenSegNet, we compared its performance

against established segmentation models including U-Net[46], SegNet[48], and DeepLabv3+[49]. Each

model was trained on uniformly sized 256 × 256 image patches cropped from the original dataset.

Official implementations were used without architectural modifications. Comparative experiments

were conducted with and without ImageNet pretraining to assess the impact of transfer learning.

Model outputs were evaluated against ground truth segmentation masks using a composite loss

function combining weighted binary cross-entropy and Dice loss, balancing pixel-wise accuracy with

regional overlap. All models were trained under identical conditions: Adam optimiser (initial learning

rate = 1×10^{-4}), cosine learning rate decay, L2 regularisation (weight decay = 1×10^{-8}), batch size of

16, and early stopping based on validation loss plateauing. A total of 108 immunohistochemistry

images were manually annotated and divided into training (n=80), validation (8), and test (n=20)

datasets. An additional private dataset comprising 25 images was generated and manually curated

for evaluation.

Cell segmentation. For cell nucleus segmentation step 3, we integrated StarDist[47] with the QuPath

platform. StarDist offers state-of-the-art performance in dense and noisy biological environments.

Following accurate nucleus segmentation, cell membrane boundaries were estimated using the

spatial coordinates of nuclei as anchor points. Expansion thresholds of 3, 4, 5, and 6 µm were tested

and validated against KRT8-stained images to determine the optimal value for our dataset.

Evaluation metrics

Centrosome segmentation performance. Centrosome segmentation performance was evaluated

on a test set comprising 25 immunohistochemistry and 17 immunofluorescence images. Metrics

included precision, recall, intersection over union (IoU), and F1 score. IoU was defined as the

common area between the predicted segmentation and the ground truth:

728
$$IoU = \frac{|\text{Prediction} \cap \text{Groundtruth}|}{|\text{Prediction} \cup \text{Groundtruth}|}$$

729   The F1 score, representing the harmonic mean of precision and recall, was calculated as:

730
$$F1\ score = 2 * \frac{\text{Precision} * \text{Recall}}{\text{Precision} + \text{Recall}}$$

731 With:

$$Precision = \frac{TP}{TP + FP}, \qquad Recall = \frac{TP}{TP + FN}$$

Where, TP, FP, and FN denote true positives, false positives, and false negatives, respectively. Both

IoU and F1 scores range from 0 to 1, with higher values indicating superior segmentation

performance.

These metrics provide a comprehensive assessment of spatial overlap, agreement, and pixel-wise

accuracy, with higher values indicating better performance in medical image segmentation.

Epithelial segmentation performance. To assess the performance of epithelial segmentation by

CenSegNet, we quantified the $F1^{IoU50}$ metric[93], a widely adopted benchmark in biomedical image

analysis. IoU50 means the IoU threshold was set to 0.5 (or 50%). A predicted bounding box was

considered a correct detection if: IoU with ground truth ≥0.5.

Statistical analysis

The exact n is stated in the corresponding figure legend. GraphPad Prism 10.3.1 (GraphPad

Software) was used to perform statistical significance analysis. Normality and lognormality were

assessed prior to statistical analysis. For datasets exhibiting a normal distribution, comparisons

between two groups were conducted using an unpaired t-test, while comparisons across multiple

groups were performed using one-way analysis of variance (ANOVA), followed by Tukey's post hoc

test for multiple comparisons or two-way ANOVA, followed by Tukey's post hoc test for multiple

comparisons. For datasets that deviated from normality, non-parametric testing was employed, using

the Kruskal-Wallis test for multiple group comparisons, followed by the Mann-Whitney U test as a

post hoc analysis. For correlation analysis, a two-tailed Pearson test was used followed by simple

linear regression for graphical representation for correlation analysis. To compare the composition

of groups based on categorical variables (e.g., histological tumour grade, T stage, histological

tumour types, hormone status, nodal status, nodal clearance, HER2 status), Fisher's exact test was

employed. This non-parametric test was chosen for its appropriateness with count data and its ability

to provide accurate p-values when expected cell counts are low (<5). All comparisons were two-

sided. All values were presented as mean ± s.e.m. For all statistical tests, $*P \leq 0.05$, $**P \leq 0.01$,

$***P \leq 0.001$ and $****P \leq 0.0001$ were considered significant.

Assistance with manuscript preparation

During the preparation of this manuscript, ChatGPT (OpenAI, version 4) was used to assist with

stylistic and grammatical refinement. All AI-generated content was critically reviewed and edited by

the authors, who take full responsibility for its accuracy and integrity.

Data availability

Examples of immunohistochemistry and immunofluorescence image datasets used for

benchmarking and for testing CenSegNet within the demo version of the pipeline are available on

https://github.com/SKELab/CenSegNet/ and https://zenodo.org/records/17131573. All other

relevant data supporting the key findings of this study are available within the article and its

Supplementary Information files or upon reasonable request. A summary of the human breast

samples used in this study are included in Supplementary Table 1. For participants on the BeGIN

study, further donor anonymised clinicopathological information is available upon reasonable

request, provided all relevant ethics approvals are in place (see "Ethics and human breast tissues"

section for further details). The source data that support the findings in all Figures and

Supplementary Figures are provided as a Source Data file within the paper. All reagents generated

in this study are available upon reasonable request.

Code availability

The source code and software CenSegNet as a ready-to-use executable with a quickstart guide,

example datasets and step-by-step procedures are freely available at

https://github.com/SKELab/CenSegNet/ and https://zenodo.org/records/17131573.

.

Acknowledgements

We would like to thank Dr. Mark Willett at the Imaging and Microscopy Centre for valuable assistance with fluorescence microscopy; Monette Lopez at the Research Histology Facility, University Hospital Southampton for performing immunohistochemistry staining and imaging; Dr Margaret Ashton-Key for classifying breast tumour compartments (normal, edge, core) and providing advice on immunohistochemistry staining conditions; Professor Susana Godinho for providing the MCF10A and MCF10A-PLK4 cell lines. In addition, we thank Dr. Xi Jia, Dr. Colinda Scheele, Dr. Marcin Przewloka, and Dr. Kif Liakath-Ali, for critically reviewing the manuscript. Cartoons in Figs. 1a, 1c and Supplementary Fig. 12 were created using https://www.freepik.com/. This work was supported by an MRC New Investigator Research Grant (MR/R026610/1) awarded to S.E. and an Institute for Life Sciences Pilot Grant awarded to R.C., S.B. and S.E. JC was supported by the Gerald Kerkut Charitable Trust and a China Scholarship Council (CSC) PhD studentship. The BeGIN study was supported by Breast Cancer Now.

Author contributions

J.C. designed and performed experiments, analysed, interpreted the data, and wrote the manuscript.

K.F. trained models and developed CenSegNet. C.S. M.B and X.D. supervised centrosome

annotation and performed transfer annotations from GeoJSON Files to PNG files and data analysis.

M.G. performed immunofluorescence experiments. R.C. provided ethically approved BeGIN breast

cancer tissues. S.B. supervised J.C. and interpreted the data. X.C. supervised K.F. on CenSegNet

development. S.E. conceived and designed the project, analysed and interpreted the data,

supervised J.C., wrote the manuscript, and acquired funding. All the authors provided intellectual

input, edited, and approved the final manuscript.

Competing interests

C.S. reports involvement in a research collaboration with Proteotype and BioNTech. R.C. reports the

following declaration: 1. Grants or Contracts: AstraZeneca, awarded an educational grant to the

University of Southampton in November 2021 for the long-term follow-up of the POSH study. 2.

Leadership or Fiduciary Roles: NICE, acts as a Breast Cancer Topic Advisor, contributing to guideline

development. Association of Breast Surgery, Member of the clinical practice and standards

committee. 3. Receipt of Equipment, Materials, etc.: SECA provided equipment for measuring body

composition to University Hospital Southampton under a model industry collaborative agreement.

References

1.   Conduit, P.T., Wainman, A. & Raff, J.W. Centrosome function and assembly in animal cells.
Nat Rev Mol Cell Biol **16**, 611-624 (2015).

2.   Kim, S. & Dynlacht, B.D. Assembling a primary cilium. *Curr Opin Cell Biol* **25**, 506-511
(2013).

3.   Godinho, S.A. & Basto, R. Centrosomes and cancer: balancing tumor-promoting and
inhibitory roles. *Trends Cell Biol* **35**, 515-526 (2025).

4.   Kiermaier, E., Stotzel, I., Schapfl, M.A. & Villunger, A. Amplified centrosomes-more than just
a threat. *EMBO Rep* **25**, 4153-4167 (2024).

5.   Nigg, E.A., Schnerch, D. & Ganier, O. Impact of Centrosome Aberrations on Chromosome
Segregation and Tissue Architecture in Cancer. *Cold Spring Harb Symp Quant Biol* **82**,
137-144 (2017).

6.   Raff, J.W. & Basto, R. Centrosome Amplification and Cancer: A Question of Sufficiency.
Dev Cell **40**, 217-218 (2017).

7.   Mittal, K. *et al.* Centrosome amplification: a quantifiable cancer cell trait with prognostic
value in solid malignancies. *Cancer Metastasis Rev* **40**, 319-339 (2021).

8.   Boveri, T. Concerning the origin of malignant tumours by Theodor Boveri. Translated and
annotated by Henry Harris. *J Cell Sci* **121 Suppl 1**, 1-84 (2008).

9.   Godinho, S.A. & Pellman, D. Causes and consequences of centrosome abnormalities in
cancer. *Philos Trans R Soc Lond B Biol Sci* **369** (2014).

10. Hsu, L.C., Kapali, M., DeLoia, J.A. & Gallion, H.H. Centrosome abnormalities in ovarian
cancer. *Int J Cancer* **113**, 746-751 (2005).

11. Lingle, W.L., Lutz, W.H., Ingle, J.N., Maihle, N.J. & Salisbury, J.L. Centrosome hypertrophy
in human breast tumors: implications for genomic stability and cell polarity. *Proc Natl Acad
Sci U S A* **95**, 2950-2955 (1998).

12. Pihan, G.A. *et al.* Centrosome defects and genetic instability in malignant tumors. *Cancer
Res* **58**, 3974-3985 (1998).

13. Sato, N. *et al.* Centrosome abnormalities in pancreatic ductal carcinoma. *Clin Cancer Res*
5, 963-970 (1999).

14. Giehl, M. *et al.* Centrosome aberrations in chronic myeloid leukemia correlate with stage of
disease and chromosomal instability. *Leukemia* **19**, 1192-1197 (2005).

15. Kramer, A., Neben, K. & Ho, A.D. Centrosome aberrations in hematological malignancies.
Cell Biol Int **29**, 375-383 (2005).

16. Basto, R. *et al.* Centrosome amplification can initiate tumorigenesis in flies. *Cell* **133**, 1032-
1042 (2008).

17. Lingle, W.L. *et al.* Centrosome amplification drives chromosomal instability in breast tumor
development. *Proc Natl Acad Sci U S A* **99**, 1978-1983 (2002).

18. Levine, M.S. *et al.* Centrosome Amplification Is Sufficient to Promote Spontaneous
Tumorigenesis in Mammals. *Dev Cell* **40**, 313-322 e315 (2017).

19. Chan, J.Y. A clinical overview of centrosome amplification in human cancers. *Int J Biol Sci*
7, 1122-1144 (2011).

20. Shekhar, M.P., Lyakhovich, A., Visscher, D.W., Heng, H. & Kondrat, N. Rad6
overexpression induces multinucleation, centrosome amplification, abnormal mitosis,
aneuploidy, and transformation. *Cancer Res* **62**, 2115-2124 (2002).

21. Meraldi, P., Honda, R. & Nigg, E.A. Aurora-A overexpression reveals tetraploidization as a
major route to centrosome amplification in p53-/- cells. *EMBO J* **21**, 483-492 (2002).

22. Fujiwara, T. *et al.* Cytokinesis failure generating tetraploids promotes tumorigenesis in p53-
null cells. *Nature* **437**, 1043-1047 (2005).

23. Duensing, A., Chin, A., Wang, L., Kuan, S.F. & Duensing, S. Analysis of centrosome
overduplication in correlation to cell division errors in high-risk human papillomavirus
(HPV)-associated anal neoplasms. *Virology* **372**, 157-164 (2008).

24. Duelli, D.M., Hearn, S., Myers, M.P. & Lazebnik, Y. A primate virus generates transformed
human cells by fusion. *J Cell Biol* **171**, 493-503 (2005).
25. Khodjakov, A. *et al.* De novo formation of centrosomes in vertebrate cells arrested during S
phase. *J Cell Biol* **158**, 1171-1181 (2002).
26. La Terra, S. *et al.* The de novo centriole assembly pathway in HeLa cells: cell cycle
progression and centriole assembly/maturation. *J Cell Biol* **168**, 713-722 (2005).
27. Davoli, T. & de Lange, T. Telomere-driven tetraploidization occurs in human cells
undergoing crisis and promotes transformation of mouse cells. *Cancer Cell* **21**, 765-776
(2012).
28. Borel, F., Lohez, O.D., Lacroix, F.B. & Margolis, R.L. Multiple centrosomes arise from
tetraploidy checkpoint failure and mitotic centrosome clusters in p53 and RB pocket
protein-compromised cells. *Proc Natl Acad Sci U S A* **99**, 9819-9824 (2002).
29. Karakaya, K. *et al.* Overexpression of EVI1 interferes with cytokinesis and leads to
accumulation of cells with supernumerary centrosomes in G0/1 phase. *Cell Cycle* **11**, 3492-
3503 (2012).
30. Dwivedi, D., Harry, D. & Meraldi, P. Mild replication stress causes premature centriole
disengagement via a sub-critical Plk1 activity under the control of ATR-Chk1. *Nat Commun*
14, 6088 (2023).
31. Loncarek, J., Hergert, P. & Khodjakov, A. Centriole reduplication during prolonged
interphase requires procentriole maturation governed by Plk1. *Curr Biol* **20**, 1277-1282
(2010).
32. Difilippantonio, M.J. *et al.* Nucleation capacity and presence of centrioles define a distinct
category of centrosome abnormalities that induces multipolar mitoses in cancer cells.
Environ Mol Mutagen **50**, 672-696 (2009).
33. Hut, H.M. *et al.* Centrosomes split in the presence of impaired DNA integrity during mitosis.
Mol Biol Cell **14**, 1993-2004 (2003).
34. Loffler, H., Fechter, A., Liu, F.Y., Poppelreuther, S. & Kramer, A. DNA damage-induced
centrosome amplification occurs via excessive formation of centriolar satellites. *Oncogene*
32, 2963-2972 (2013).
35. Loncarek, J., Hergert, P., Magidson, V. & Khodjakov, A. Control of daughter centriole
formation by the pericentriolar material. *Nat Cell Biol* **10**, 322-328 (2008).
36. Starita, L.M. *et al.* BRCA1-dependent ubiquitination of gamma-tubulin regulates
centrosome number. *Mol Cell Biol* **24**, 8457-8466 (2004).
37. Marteil, G. *et al.* Over-elongation of centrioles in cancer promotes centriole amplification
and chromosome missegregation. *Nat Commun* **9**, 1258 (2018).
38. Tang, C.J., Fu, R.H., Wu, K.S., Hsu, W.B. & Tang, T.K. CPAP is a cell-cycle regulated
protein that controls centriole length. *Nat Cell Biol* **11**, 825-831 (2009).
39. Schmidt, T.I. *et al.* Control of centriole length by CPAP and CP110. *Curr Biol* **19**, 1005-1011
(2009).
40. Burgy, L. *et al.* CenFind: a deep-learning pipeline for efficient centriole detection in
microscopy datasets. *BMC Bioinformatics* **24**, 120 (2023).
41. Sankaran, D.G., Stemm-Wolf, A.J., McCurdy, B.L., Hariharan, B. & Pearson, C.G. A semi-
automated machine learning-aided approach to quantitative analysis of centrosomes and
microtubule organization. *J Cell Sci* **133** (2020).
42. Patel, N. *et al.* Integrated genomics and functional validation identifies malignant cell
specific dependencies in triple negative breast cancer. *Nat Commun* **9**, 1044 (2018).
43. Mittal, K. *et al.* A Quantitative Centrosomal Amplification Score Predicts Local Recurrence
of Ductal Carcinoma In Situ. *Clin Cancer Res* **26**, 2898-2907 (2020).
44. Sauer, C.M. *et al.* Molecular landscape and functional characterization of centrosome
amplification in ovarian cancer. *Nat Commun* **14**, 6505 (2023).
45. Khanam, R. & Hussain, M. YOLOv11: An Overview of the Key Architectural Enhancements.
arXiv e-prints, arXiv:2410.17725 (2024).

46.   Ronneberger, O., Fischer, P. & Brox, T. in Medical Image Computing and Computer-
920   Assisted Intervention – MICCAI 2015. (eds. N. Navab, J. Hornegger, W.M. Wells & A.F.
921   Frangi) 234-241 (Springer International Publishing, Cham; 2015).
922   47.   Schmidt, U., Weigert, M., Broaddus, C. & Myers, G. in Medical Image Computing and
923   Computer Assisted Intervention – MICCAI 2018. (eds. A.F. Frangi, J.A. Schnabel, C.
924   Davatzikos, C. Alberola-López & G. Fichtinger) 265-273 (Springer International Publishing,
925   Cham; 2018).
926   48.   Badrinarayanan, V., Kendall, A. & Cipolla, R. SegNet: A Deep Convolutional Encoder-
927   Decoder Architecture for Image Segmentation. *IEEE Transactions on Pattern Analysis and*
928   *Machine Intelligence* **39**, 2481-2495 (2017).
929   49.   Chen, L.-C., Zhu, Y., Papandreou, G., Schroff, F. & Adam, H. arXiv:1802.02611 (2018).
930   50.   He, L., Zhou, Y., Liu, L., Zhang, Y. & Ma, J. Application of the YOLOv11-seg algorithm for
931   AI-based landslide detection and recognition. *Sci Rep* **15**, 12421 (2025).
932   51.   Sridharan, D.M., Enerio, S., LaBarge, M.A., Stampfer, M.M. & Pluth, J.M. Lesion complexity
933   drives age related cancer susceptibility in human mammary epithelial cells. *Aging (Albany*
934   *NY)* **9**, 665-686 (2017).
935   52.   Shimomura, A., Miyoshi, Y., Taguchi, T., Tamaki, Y. & Noguchi, S. Association of loss of
936   BRCA1 expression with centrosome aberration in human breast cancer. *Journal of Cancer*
937   *Research and Clinical Oncology* **135**, 421-430 (2009).
938   53.   Agnoletto, A. & Brisken, C. Hormone Signaling in Breast Development and Cancer, in *A*
939   *Guide to Breast Cancer Research: From Cellular Heterogeneity and Molecular*
940   *Mechanisms to Therapy*. (eds. T. Sørlie & R.B. Clarke) 279-307 (Springer Nature
941   Switzerland, Cham; 2025).
942   54.   Loibl, S., Poortmans, P., Morrow, M., Denkert, C. & Curigliano, G. Breast cancer. *The*
943   *Lancet* **397**, 1750-1769 (2021).
944   55.   Menarche, menopause, and breast cancer risk: individual participant meta-analysis,
945   including 118 964 women with breast cancer from 117 epidemiological studies. *The Lancet*
946   *Oncology* **13**, 1141-1151 (2012).
947   56.   Type and timing of menopausal hormone therapy and breast cancer risk: individual
948   participant meta-analysis of the worldwide epidemiological evidence. *The Lancet* **394**,
949   1159-1168 (2019).
950   57.   Heer, E. *et al.* Global burden and trends in premenopausal and postmenopausal breast
951   cancer: a population-based study. *The Lancet Global Health* **8**, e1027-e1037 (2020).
952   58.   Delaval, B. & Doxsey, S.J. Pericentrin in cellular function and disease. *J Cell Biol* **188**, 181-
953   190 (2010).
954   59.   Jiang, T. *et al.* Drivers of centrosome abnormalities: Senescence progression and tumor
955   immune escape. *Semin Cancer Biol* **110**, 56-64 (2025).
956   60.   Nigg, E.A. & Raff, J.W. Centrioles, centrosomes, and cilia in health and disease. *Cell* **139**,
957   663-678 (2009).
958   61.   Denu, R.A. *et al.* Centrosome amplification induces high grade features and is prognostic of
959   worse outcomes in breast cancer. *BMC Cancer* **16**, 47 (2016).
960   62.   Van Herck, Y. *et al.* Is cancer biology different in older patients? *The Lancet Healthy*
961   *Longevity* **2**, e663-e677 (2021).
962   63.   Benz, C.C. Impact of aging on the biology of breast cancer. *Critical Reviews in*
963   *Oncology/Hematology* **66**, 65-74 (2008).
964   64.   Lopes, C.A.M. *et al.* Centrosome amplification arises before neoplasia and increases upon
965   p53 loss in tumorigenesis. *J Cell Biol* **217**, 2353-2363 (2018).
966   65.   Monteiro, P., Yeon, B., Wallis, S.S. & Godinho, S.A. Centrosome amplification fine tunes
967   tubulin acetylation to differentially control intracellular organization. *EMBO J* **42**, e112812
968   (2023).
969   66.   Godinho, S.A. *et al.* Oncogene-like induction of cellular invasion from centrosome
970   amplification. *Nature* **510**, 167-171 (2014).

67.　LoMastro, G.M. & Holland, A.J. The Emerging Link between Centrosome Aberrations and
Metastasis. *Dev Cell* **49**, 325-331 (2019).
68.　Rhys, A.D. *et al.* Loss of E-cadherin provides tolerance to centrosome amplification in
epithelial cancer cells. *J Cell Biol* **217**, 195-209 (2018).
69.　Ganier, O. *et al.* Structural centrosome aberrations promote non-cell-autonomous
invasiveness. *EMBO J* **37** (2018).
70.　Garcia-Carpio, I. *et al.* Extra centrosomes induce PIDD1-mediated inflammation and
immunosurveillance. *EMBO J* **42**, e113510 (2023).
71.　Ganier, O., Schnerch, D. & Nigg, E.A. Structural centrosome aberrations sensitize polarized
epithelia to basal cell extrusion. *Open Biol* **8** (2018).
72.　Kahl, I. *et al.* The cell cycle-related genes RHAMM, AURKA, TPX2, PLK1, and PLK4 are
associated with the poor prognosis of breast cancer patients. *J Cell Biochem* **123**, 581-600
(2022).
73.　Toft, D.J. & Cryns, V.L. Minireview: Basal-like breast cancer: from molecular profiles to
targeted therapies. *Mol Endocrinol* **25**, 199-211 (2011).
74.　Cosenza, M.R. *et al.* Asymmetric Centriole Numbers at Spindle Poles Cause Chromosome
Missegregation in Cancer. *Cell Rep* **20**, 1906-1920 (2017).
75.　Ganem, N.J., Godinho, S.A. & Pellman, D. A mechanism linking extra centrosomes to
chromosomal instability. *Nature* **460**, 278-282 (2009).
76.　Ingebriktsen, L.M. *et al.* Elevated expression of Aurora-A/AURKA in breast cancer
associates with younger age and aggressive features. *Breast Cancer Res* **26**, 126 (2024).
77.　Wong, Y.L. *et al.* Cell biology. Reversible centriole depletion with an inhibitor of Polo-like
kinase 4. *Science* **348**, 1155-1160 (2015).
78.　Melichar, B. *et al.* Safety and activity of alisertib, an investigational aurora kinase A inhibitor,
in patients with breast cancer, small-cell lung cancer, non-small-cell lung cancer, head and
neck squamous-cell carcinoma, and gastro-oesophageal adenocarcinoma: a five-arm
phase 2 study. *Lancet Oncol* **16**, 395-405 (2015).
79.　Borisa, A.C. & Bhatt, H.G. A comprehensive review on Aurora kinase: Small molecule
inhibitors and clinical trial studies. *Eur J Med Chem* **140**, 1-19 (2017).
80.　Veitch, Z.W. *et al.* Safety and tolerability of CFI-400945, a first-in-class, selective PLK4
inhibitor in advanced solid tumours: a phase 1 dose-escalation trial. *Br J Cancer* **121**, 318-
324 (2019).
81.　Lei, Q. *et al.* YLT-11, a novel PLK4 inhibitor, inhibits human breast cancer growth via
inducing maladjusted centriole duplication and mitotic defect. *Cell Death Dis* **9**, 1066
(2018).
82.　Watts, C.A. *et al.* Design, synthesis, and biological evaluation of an allosteric inhibitor of
HSET that targets cancer cells with supernumerary centrosomes. *Chem Biol* **20**, 1399-1410
(2013).
83.　Blyth, R.R.R. *et al.* Development and characterisation of a novel 3D in vitro model of
obesity-associated breast cancer as a tool for drug testing. *NPJ Breast Cancer* **11**, 50
(2025).
84.　Daly, A. *et al.* Assessment of body composition in breast cancer patients: concordance
between transverse computed tomography analysis at the fourth thoracic and third lumbar
vertebrae. *Front Nutr* **11**, 1366768 (2024).
85.　Bosy-Westphal, A. *et al.* What makes a BIA equation unique? Validity of eight-electrode
multifrequency BIA to estimate body composition in a healthy adult population. *Eur J Clin
Nutr* **67 Suppl 1**, S14-21 (2013).
86.　Rovero, S. *et al.* DNA vaccination against rat her-2/Neu p185 more effectively inhibits
carcinogenesis than transplantable carcinomas in transgenic BALB/c mice. *J Immunol* **165**,
5133-5142 (2000).
87.　Rakha, E.A. *et al.* UK recommendations for HER2 assessment in breast cancer: an update.
J Clin Pathol **76**, 217-227 (2023).

88.   Ellis, I.O., Rakha, E.A., Tse, G.M. & Tan, P.H. An international unified approach to reporting
and grading invasive breast cancer. An overview of the International Collaboration on
Cancer Reporting (ICCR) initiative. *Histopathology* **82**, 189-197 (2023).
89.   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods*
9, 676-682 (2012).
90.   Bankhead, P. *et al.* QuPath: Open source software for digital pathology image analysis. *Sci*
Rep **7**, 16878 (2017).
91.   Stringer, C. & Pachitariu, M. Cellpose3: one-click image restoration for improved cellular
segmentation. *Nat Methods* **22**, 592-599 (2025).
92.   Stringer, C., Wang, T., Michaelos, M. & Pachitariu, M. Cellpose: a generalist algorithm for
cellular segmentation. *Nat Methods* **18**, 100-106 (2021).
93.   Ong, H.T. *et al.* Digitalized organoids: integrated pipeline for high-speed 3D analysis of
organoid structures using multilevel segmentation and cellular topology. *Nature Methods*
22, 1343-1354 (2025).

1037

Fig. 1

a   Immunohistochemistry training dataset

b   Immunofluorescence training dataset

c   Model comparison

Fig. 1 Development and benchmarking of CenSegNet for centrosome segmentation. a Left: workflow for generating tissue microarrays (TMAs) comprising 911 breast tissue sample cores from normal breast tissue, breast tumours, and adjacent non-tumour tissue, collected from 127 breast cancer patients in the BeGIN cohort. Figure 1a is schematic for illustrative purposes to demonstrate sampling strategy for the TMA. Cores were taken from FFPE blocks following breast cancer surgery. Top right: immunohistochemical staining of all ROIs with pericentrin (centrosome marker) and haematoxylin (nuclear counterstain). Bottom right: random selection of 108 ROIs used to construct a training dataset containing 2,486 epithelial compartments, 108 stromal compartments and 14,679 annotated centrosomes. **b** Left and middle: representative confocal images of human mammary epithelial cells (MECs) and mouse mammary tumour tissues exhibiting normal or amplified centrosomes. Cells and tissues were stained for pericentrin (orange) and counterstained with DAPI (DNA, teal). Scale bars, 10 μm. Right: training dataset derived from these images, comprising 1,285 annotated centrosomes. **c** Left: 256 × 256 pixel cropped patches from immunohistochemistry and immunofluorescence datasets. Middle left: training of existing segmentation architectures (U-Net, SegNet, DeepLabv3+) and the proposed CenSegNet using these datasets. Middle right: CenSegNet performance across both immunohistochemistry and immunofluorescence test sets. CenSegNet operates in three sequential phases: (1) object detection to generate bounding-box predictions for individual centrosomes; (2) pixel-level segmentation of detected objects; (3) StarDist-based whole-cell segmentation.

Fig. 2

a

Performance of published models *versus* CenSegNet

| Raw images | U-Net | SegNet | DeepLabv3+ | CenSegNet (ours) |

Ground-truth   Prediction   Overlap

Immunohistochemistry — DNA Centrosomes

Immunofluorescence — DNA Centrosomes

b Immunohistochemistry
R²=0.9999; *p*<0.0001

Detection efficiency

c Immunofluorescence
R²=0.9873; *p*<0.0001

d Immunohistochemistry
R²=0.9613; *p*<0.0001

Segmentation efficiency

e Overall F1 score (0.82) of Immunohistochemistry and Immunofluorescence datasets

f Annotated 6,475 centrosomes in testing dataset

g Segmented 333,148 centrosomes in testing dataset

Fig. 2 Validation of CenSegNet performance against alternative models and human annotations. a Left: representative immunohistochemistry and immunofluorescence images used for centrosome segmentation. Right: colour overlays showing predictions from U-Net, SegNet, DeepLabv3+, and CenSegNet (teal: ground-truth annotations; orange: predicted segmentation; white: overlap between predictions and ground-truth). Magenta text indicates the F1 score for corresponding images of each method (0: complete disagreement; 1: complete concordance). Yellow arrow marks the overprediction. **b** Pearson correlation between centrosome counts predicted by CenSegNet and ground truth across 25 immunohistochemistry images. Grey circles represent individual patients, purple line represents regression line, Black dot line represents confidence band, and teal dot line represents 1:1 line. Two-tailed Pearson correlation test followed by simple linear regression for visualisation, ****$P < 0.0001$. **c** Pearson correlation between centrosome counts predicted by CenSegNet and ground truth across 17 immunofluorescence images. Two-tailed Pearson correlation test, followed by simple linear regression for graphical representation, ****$P < 0.0001$. **d** Pearson correlation between centrosome size predicted by CenSegNet and measured by annotators (n = 550 centrosomes from >200 ROIs across 20 images). Two-tailed Pearson correlation test, followed by simple linear regression for graphical representation, ****$P < 0.0001$. **e** Heatmap of overall F1 scores (maximum 0.82) from combined immunohistochemistry and immunofluorescence test datasets. The *x*-axis represents the detection confidence threshold, and the *y*-axis represents the segmentation confidence threshold. Each cell represents the mean F1 score for that threshold pair; darker cyan indicates higher values. Optimal performance (F1 = 0.82) was achieved across a broad range of threshold combinations, indicating model robustness to parameter variation. **f, g** Manual annotations of 6,475 centrosomes from normal, edge, and tumour regions—stratified by size (0.5–1.0 μm^2 to >10.5 μm^2) **(f)** were compared with automated segmentations from the full dataset of 333,148 centrosomes **(g)** (0–0.5 μm^2 to >12.0 μm^2). CenSegNet segmentation achieved performance comparable to expert human annotation. Data are presented as individual data points. Source data are provided as Source Data file.

Fig. 3

Fig. 3 Age-related gradients of centrosome aberrations across breast tissue regions. a Percentage of cells with different centrosome numbers in normal, edge, and tumour regions. **b** Representative cell segmentation masks of normal, edge, and tumour regions across patient age groups (Light grey: cells with one centrosome; purple: cells with two centrosomes; cyan: cells with more than two centrosomes). **c** Percentage of patients with structural CA (Stru CA) and Num CA across age groups in edge (≤50 years: n = 8 patients; 50–60 years: n = 17 patients; 60–70 years: n = 10 patients; >70, n = 14 patients) and Tumour (≤50 years: n = 18 patients; 50–60 years: n = 35 patients; 60–70 years: n = 27 patients; >70 years: n = 20 patients) regions. Two-way ANOVA with Tukey's test, *P = 0.0477, **P = 0.0053. Data are presented as mean ± s.e.m. **d** Percentage of cells with Stru CA in edge and tumour regions across age in Edge and Tumour regions. One-way ANOVA (Stru CA$^+$ cells: Edge, P = 0.7637; Tumour, P = 0.4489 with Tukey's test. absence of asterisks indicates no statistically significant correlation. **e** Percentage of cells with Num CA in edge and tumour regions across age. One-way ANOVA (Num CA$^+$ cells: Edge, P = 0.3427; Tumour, P = 0.9086) with Tukey's test, absence of asterisks indicates no statistical significance. Data are presented as mean ± s.e.m. **f, g** Centrosome segmentation results from normal regions across age groups, stratified by size (0.5–1.0 µm^2 to 6.0–6.5 µm^2). One-way ANOVA (****P < 0.0001) with Tukey's test, ****P < 0.0001. Data are presented as violin plots showing the distribution of values; dashed lines indicate median and interquartile ranges. **h, i** Centrosome segmentation results from edge and tumour regions across age groups, stratified by size (6.5–7.0 µm^2 to 12.0 µm^2). One-way ANOVA (****P < 0.0001) with Tukey's test, Edge: ****P < 0.0001; Tumour: *P = 0.0484, ****P < 0.0001. Data are presented as violin plots showing the distribution of values; dashed lines indicate median and interquartile ranges. **j, k** Centrosome number per cell in edge and tumour regions across age groups, stratified by centrosome number (1–8). One-way ANOVA (Edge, ****P < 0.0001; Tumour, ****P < 0.0001) with Tukey's test, Edge: ***P = 0.0006, ****P < 0.0001; Tumour: **P = 0.0054, ****P < 0.0001, absence of asterisks indicates no statistical significance. Data are presented as individual data points and mean ± s.e.m. Source data are provided as Source Data file.

Fig. 4

Fig. 4 Patient stratification by CA burden and correlation with clinicopathological features. a–d Percentage of patients with Stru CA and Num CA across histological tumour grade, nodal status, histological tumour size, and histological tumour type in Edge [Grade: G1 (n= 5), G2 (n=27), G3 (n=17); Nodal status: N0 (n=38), N2 (n=1), N3 (n=10); T stage: T1 (n=18), T2 (n=16), T3 (n=15); Histological type: Invasive ductal (n=36), Lobular (n=8), Mixed (n=5)], and Tumour [Grade: G1 (n=12), G2 (n=52), G3 (n=35); Nodal status: N0 (n=77), N2 (n=4), N3 (n=18); T stage: T1 (n= 30), T2 (n= 29), T3 (n= 40); Histological type: Invasive ductal (n=82), Lobular (n=8), Mixed (n=8)] regions. Two-way ANOVA (Grade, **P = 0.0020; Nodal status, ** P = 0.0030; T stage, P = 0.4024; Histological

type, ** P = 0.0037) with Tukey's test, Grade: *P = 0.0130, **P = 0.0018; Nodal status: *P = 0.0113, **P = 0.0030; Histological type: (Invasive ductal *versus* Lobular) **P = 0.0063, (Invasive ductal *versus* Mixed) **P = 0.0061. Data are presented as mean ± s.e.m. **e, f** Patients classified by composite CA burden into Stru^{high}Num^{low} (A), Stru^{high}Num^{high} (B), Stru^{low}Num^{low} (C), and Stru^{low}Num^{high} (D) groups in Edge (A: n = 14 patients; B: n = 7 patients; C: n = 16 patients; D: n = 7 patients) and Tumour (A: n = 20 patients; B: n = 16 patients; C: n = 35 patients; D: n = 23 patients) regions. Histograms and pie charts show percentages of patients in each group in edge and tumour regions. **g–j** Comparative analysis of patient characteristics, including age, number of involved nodes, and total and histological tumour size, across composite CA groups A–D in edge and tumour regions. One-way ANOVA Age (Edge, P = 0.7719; Tumour, P = 0.6469) Nodal number (Edge, P = 0.3164; Tumour, P = 0.9944); Total tumour size (Edge, *P = 0.0491; Tumour, *P = 0.0318); Histological tumour size (Edge, *P = 0.0403; Tumour, P = 0.0732) with Tukey's test, Total tumour size (Edge, *P (A vs D) = 0.0484; Tumour, *P (B vs D) = 0.0296). Absence of asterisks indicates no statistical significance. Data are presented as individual data points and as box and whiskers plots showing the distribution of values, median and quartiles. **k–o** Percentage of patients with different tumour characteristics (histological tumour grade, T stage, nodal status, nodal clearance, genetic subtype) across composite CA groups A–D in edge and tumour regions. Fisher's exact test, left: *P = 0.0261; middle: ***P = 0.0005 (middle left), ***P = 0.0006 (middle right) and ****P < 0.0001; right: **P = 0.0091. **o** 1st degree, family history with 1st degree relative; 2nd degree, family history with 2nd degree relative. Genetic, pathogenic variant in breast cancer-related gene. Source data are provided as Source Data file.

Fig. 5

Fig. 5 Interplay between Stru CA and Num CA drives centrosome defect spatial heterogeneity in breast edge and tumour regions. a Representative immunohistochemistry images of human breast tumour tissues (stained for pericentrin, counterstained with haematoxylin) and corresponding StarDist cell masks showing differences in Stru CA between edge and tumour regions. Left: Stru CA higher in Tumour than Edge. Right: Stru CA lower in Tumour than Edge (teal: cells with ≥4 centrosomes; purple: cells with three centrosomes; grey: cells with ≤2 centrosomes). Scale bars, 10 µm. **b** Comparative analysis of centrosome number per cell in edge and tumour regions for Stru CA$^{T>E}$ and Stru CA$^{E>T}$ groups. Comparisons between Edge and Tumour were performed separately for each condition using two-sided unpaired t-test, left: ****P < 0.0001; right: ****P < 0.0001. Data

are presented as mean ± s.e.m. **c** Representative immunohistochemistry images of human breast tumour tissues (stained for pericentrin, counterstained with haematoxylin) and corresponding StarDist cell masks showing differences in Stru CA and Num CA between edge and tumour regions. Left: Num CA higher in Tumour than Edge. Right: Num CA lower in tumour than edge (teal: cells with ≥4 centrosomes; purple: cells with three centrosomes; grey: cells with ≤2 centrosomes). **d** Comparative analysis of centrosome size in Edge and Tumour regions for Num $CA^{T>E}$ and Num $CA^{E>T}$ groups. Comparisons between Edge and Tumour were performed separately for each condition using two-sided unpaired *t*-test. Left: ****$P < 0.0001$; right: ****$P < 0.0001$. Data are presented as violin plots showing the distribution of values; dashed lines indicate median and interquartile ranges. **e** Top: correlation between centrosome number and mean centrosome size at the single-cell level in Stru $CA^{T>E}$, Stru $CA^{E>T}$, Num $CA^{T>E}$, and Num $CA^{E>T}$ groups. One-way ANOVA (****$P < 0.0001$ across Stru $CA^{T>E}$, Stru $CA^{E>T}$, Num $CA^{T>E}$, and Num $CA^{E>T}$ groups) with Tukey's test: Stru $CA^{T>E}$ [Edge: ****$P < 0.0001$; Tumour: (top) *$P = 0.0318$, (bottom) *$P = 0.0416$, **$P = 0.0056$, ****$P < 0.0001$]; Stru $CA^{E>T}$ [Edge: *$P = 0.0139$, ****$P < 0.0001$; Tumour: (top) *P (2 vs 5) = 0.0362, (top) *P (2 vs 6) = 0.0144, ****$P < 0.0001$); Num $CA^{T>E}$ [Edge: *$P = 0.0259$, ****$P < 0.0001$; Tumour: *$P = 0.0421$, (top) **$P = 0.0081$, (bottom) **$P = 0.0025$, ****$P < 0.0001$]; Num $CA^{E>T}$ [Edge: ***$P = 0.0003$, ****$P < 0.0001$; Tumour: (top) *$P = 0.0340$, (bottom) *$P = 0.0115$, ****$P < 0.0001$]. Bottom: Corresponding representative StarDist masks showing cells with Stru CA and Num CA in Edge and Tumour regions. Data are presented as individual data points and mean ± s.e.m. Source data are provided as Source Data file.

Fig. 6

a

Fig. 6 Open accessibility and integration of CenSegNet for broad adoption. a CenSegNet can be accessed *via* a Python application programming interface (API). **b** Top: graphical user interface (GUI) of CenSegNet designed for a streamlined workflow, supporting image upload, model-based prediction, and data export. Bottom: interface functions for both immunohistochemistry and immunofluorescence images, enabling users to upload images, apply the relevant prediction models, and export quantitative data including pixel-level centrosome size, localisation, and count.