Endogenous oxytocin-linked heart rate synchrony for social connection in live sports spectators

Takashi Matsui[a,b,1], Takafumi Yamaguchi[a,1], Daisuke Funabashi[a], Shion Takahashi[a],

Hiroki Matsuoka[a,b], Shohei Dobashi[a], Kiyonobu Kigoshi[a], Shinzo Yamada[c], Hideki Takagi[a,b,c]

[a]Institute of Health and Sport Sciences, University of Tsukuba, 1-1-1 Tennoudai, Tsukuba,

Ibaraki 305-8574, Japan

[b]R&D Center for Sport Innovation, University of Tsukuba, 1-1-1 Tennoudai, Tsukuba, Ibaraki

305-8574, Japan

[c]Bureau of Physical Education and Sport, University of Tsukuba, 1-1-1 Tennoudai, Tsukuba,

Ibaraki 305-8574, Japan

[1]These authors contributed equally

Correspondence

Takashi Matsui, Ph.D.

Associate Professor

Exercise Biochemistry & Sport Neurobiology Division

Institute of Health and Sport Sciences, University of Tsukuba

Sport IT Division Leader, University of Tsukuba R&D Center for Sport Innovation

Vice Director, University of Tsukuba R&D Center for Lifestyle Innovation

E-mail: matsui.takashi.ga@u.tsukuba.ac.jp

Abstract

Oxytocin has been proposed as a pharmacological target for social dysfunction. However, clinical trials of exogenous oxytocin have yielded modest and inconsistent benefits, highlighting the need for naturalistic, non-pharmacological approaches that recruit endogenous oxytocin. This prospective observational field study examined whether live sports spectating in a modest collegiate setting engages endogenous oxytocin and heart rate synchrony, and how these processes relate to social bonding. Sixty casual spectators at two university basketball games provided salivary samples for oxytocin and cortisol and completed psychological questionnaires before the game, at halftime, and after the game, while heart rate was continuously recorded. Oxytocin dynamics were baseline dependent: individuals with low pre-game oxytocin showed clear increases during spectating, whereas those with high pre-game oxytocin maintained elevated concentrations; across both groups, cortisol decreased and the oxytocin-to-cortisol ratio increased, indicating a shift toward an oxytocin-dominant, low-stress hormonal profile. Live spectating also enhanced heart rate synchrony between spectators, and higher oxytocin levels and stronger synchrony were associated with greater enjoyment, stronger unity with players and fellow fans, higher flow, and stronger intention to revisit. Mediation analyses indicated that perceived unity mediated links between physiological responses and experiential outcomes. These findings identify coordinated oxytocin dynamics and physiological synchrony as candidate mechanisms through which live sports spectating can foster social connection and positive psychological states in everyday, low-stakes settings. Live sports events may represent a simple, scalable non-pharmacological approach to engaging oxytocin-related systems and interpersonal synchrony to promote social connection in everyday life.

Key words

oxytocin | physiological synchrony | heart rate | spectatorship | social bonding

1. Introduction

Oxytocin is a hypothalamic neuropeptide with potent central effects on social cognition and stress regulation, and has attracted considerable interest as a pharmacological target for social dysfunction in neuropsychiatric disorders such as autism spectrum disorder and schizophrenia[1–3]. Exogenous oxytocin administered intranasally can acutely modulate social attention, emotion recognition, and activity in limbic–striatal circuits, and has therefore been trialled as an adjunctive treatment for social impairments [4–6]. However, large-scale and longer-term clinical trials have yielded small or inconsistent benefits on core social symptoms, and considerable uncertainty remains regarding optimal dosing regimens, individual response profiles, and long-term safety [7–9]. In addition, peripherally administered oxytocin has an unfavourable pharmacokinetic and delivery profile, including a short plasma half-life, limited central penetration, and off-target peripheral effects, and even though its short-term safety profile appears acceptable, rare adverse reactions have been reported with repeated or high-dose use [10,11]. These limitations have stimulated growing interest in naturalistic, non-pharmacological strategies that can recruit endogenous oxytocin to engage oxytocin-linked circuits for social connection without the constraints of exogenous drug administration [12,13].

Beyond its classical roles in parturition and lactation, endogenous oxytocin is released by everyday affiliative cues, particularly low-intensity, non-noxous social touch such as gentle stroking or massage, which has been linked to stress-buffering effects and increased well-being [14,15]. Experimental studies in humans show that affective tactile stimulation, including massage-like foot stimulation, can acutely increase peripheral oxytocin concentrations and concomitantly engage cortical regions implicated in social cognition and reward, such as the orbitofrontal cortex and superior temporal sulcus [14,16]. Similarly, traditional martial arts training, which combines vigorous physical exercise with structured dyadic contact, increases salivary oxytocin immediately after high-intensity sparring, with larger oxytocin responses during close-contact ground grappling than during more distal "punch–kick" sparring [17]. Martial-arts–based interventions in youths at high psychosocial risk further indicate that oxytocin and cortisol responses during training differ between high-risk and low-risk groups, with high-risk youths showing lower baseline oxytocin levels and altered reactivity compared with low-risk peers [18].

Familiar vocal social contact can also trigger oxytocin release after stress; for example, comfort delivered solely via a mother's voice elevates oxytocin and accelerates

cortisol recovery in children, suggesting that social vocalizations provide a potent non-tactile

route for engaging oxytocin-linked bonding and stress-regulation pathways [19,20].

Endogenous oxytocin levels are likewise elevated in naturally occurring affiliative contexts such

as the early stages of romantic attachment, where higher plasma oxytocin covaries with couples'

interactive reciprocity and dyadic synchrony [21]. A recent systematic review and meta-analysis

concluded that endogenous oxytocin concentrations show a modest but reliable association with

human social interaction overall, while also revealing substantial heterogeneity across

individuals, interaction types, and assay methods [22,23]. Collectively, these findings suggest

that multiple sensory, motor, and relational channels, including social touch, vocal

communication, and structured physical practices, can recruit endogenous oxytocin in everyday

life, but they also underscore the need to identify robust, ecologically valid social behaviours

that reliably engage the human oxytocin system despite this heterogeneity and methodological

uncertainty [22].

Large-scale collective events such as religious rituals, concerts, and sporting matches

can generate intense shared emotional experiences and feelings of togetherness, and recent work

suggests that physiological synchrony in crowds may be one mechanism through which such

events promote social bonding [24,25]. Field and laboratory studies show that being part of a

crowd or dyad is associated with increased heart rate synchrony, and that greater physiological

synchrony predicts stronger feelings of connectedness and cooperative attitudes [26,27]. Such

increases in synchrony are likely driven not only by individual arousal but also by shared

external stimuli, including the tension of the game, the roar of the crowd, and collective

emotional reactions, that act as common drivers of cardiac dynamics across individuals. In

parallel, social psychological research on sport fandom has developed the Team Identification–

Social Psychological Health Model, which posits that identifying with a team and engaging in

spectatorship help fans gain social connections that, in turn, enhance social well-being [28]

Consistent with this framework, observational and longitudinal studies indicate that

live sport spectatorship and stronger team loyalty are associated with greater life satisfaction,

perceived emotional support from other fans, and reduced loneliness, particularly among

middle-aged and older adults [29,30]. From a neuroendocrine perspective, theoretical work has

proposed that oxytocin may support coordination, trust, and cohesion in team sport

environments, yet empirical research in sports has focused predominantly on cortisol and

testosterone responses to competition in athletes and, to a lesser extent, fans [31,32]. Despite

growing interest in the psychobiology of fandom, oxytocin in spectators remains largely

unexplored. To date, however, no study has simultaneously examined endogenous oxytocin

dynamics, cardiac synchrony among co-present spectators, and changes in perceived social

bonding within a real-world live sports event, particularly among casual fans who do not belong

to established supporter groups.

We therefore hypothesized that a single session of live sports spectating by casual

spectators induces baseline-dependent increases in endogenous oxytocin in association with

enhancements in interpersonal heart rate synchrony, thereby promoting post-game unity with

fellow fans and loyalty to the team. To test this hypothesis, we conducted a field study of

previously unacquainted casual spectators at collegiate basketball games, integrating repeated

measures of salivary oxytocin and cortisol with continuous heart rate recording and self-

reported indices of social connection and experiential quality.

2. Materials and Methods

2.1. Participants.

We conducted a prospective observational field study with spectators attending home basketball

games at the University of Tsukuba. A total of 62 adults were recruited from attendees of

"TSUKUBA LIVE!", the university's home event series (Fig. 1A). Unlike professional sports

events, these collegiate games attracted modest crowds primarily composed of students and

local residents, reflecting a predominantly casual rather than highly devoted fan base. Each

participant attended one game, which resulted in either a win or a loss for the home team.

Two participants were excluded from all analyses due to insufficient pre-game saliva

samples, leaving 60 participants in the final analytic sample. Based on the median pre-game

salivary oxytocin concentration (27.51 pg/mL; Supplementary Table 1), participants were

classified into a high baseline oxytocin group (High-OT; n = 30; 22.2 ± 8.6 years; 10 females)

and a low baseline oxytocin group (Low-OT; n = 30; 22.9 ± 9.0 years; 11 females). Baseline

oxytocin levels were higher in the High-OT group than in the Low-OT group by design,

whereas there were no significant group differences in sex, age, body composition, resting heart

rate, or habitual physical activity, indicating that the grouping primarily reflected pre-game

oxytocin concentration rather than broader demographic or physiological differences (Table S1).

An a priori power analysis was conducted using G*Power 3.1, based on a partial η^2 of

0.036 derived from a previous study examining the association between baseline oxytocin levels

and responses to video viewing [33]. This analysis indicated that a total sample size of 60

participants would provide 90% power to detect a significant difference between two

independent means in a two-sided test with an α level of 0.05.

Inclusion criteria were as follows: age 18 years or older; ability to understand the

study procedures and provide informed consent; no current medication for neuropsychiatric,

cardiovascular, or metabolic diseases; and ability to attend the stadium for the duration of the

measurements. The study was approved by the Research Ethics Committee of the Institute of

Health and Sport Sciences, University of Tsukuba, and was conducted in accordance with the

Declaration of Helsinki. All participants provided written informed consent prior to

participation. The study is reported in accordance with the STROBE guidelines for

observational studies.

2.2. Study design and procedure

This prospective observational field study was conducted during two home basketball games at

the University of Tsukuba (Fig. 1C). The study was designed to capture coupled endocrine and

cardiovascular responses to live sports spectating by combining repeated saliva sampling with

continuous heart rate recording and self-report measures of mood and social connection.

When inviting participants, an outline of the study and a URL link to an online pre-

survey (see Section 2.3) were sent via e-mail to potential spectators of the home event series

"TSUKUBA LIVE!". Participants were instructed to refrain from consuming alcohol from the

day before the game until the end of the experiment, to avoid caffeine intake and strenuous

physical exercise on the day of the game, and to abstain from eating for at least 2 hours before

arriving at the stadium.

On the day of measurement, participants arrived at the indoor arena approximately 60

minutes before tipoff. Upon arrival, they received a detailed explanation of the procedures again

and provided written informed consent. Participants were then fitted with an optical heart rate

sensor (Polar Verity Sense; Polar, Finland) on the upper arm and completed baseline saliva

sampling (2 mL) for hormone analysis as well as baseline psychological questionnaires in the

concourse area (Fig. 1B). After these baseline assessments, participants moved to their assigned

seats and watched the game under naturalistic conditions. Additional saliva samples and

questionnaires were collected at halftime and immediately after the game, yielding three

repeated measurement points: pre-game (T1), halftime (T2), and post-game (T3).

2.3. Demographic and physical activity assessment

Before the game, participants completed an online pre-survey delivered via Google Forms to

collect demographic characteristics (for example, age and sex) and habitual physical activity.

Habitual physical activity was assessed using the Japanese version of the International Physical

Activity Questionnaire (IPAQ), which estimates average daily time spent in vigorous-intensity,

moderate-intensity, and walking activities, as well as sedentary behaviour.

Some participants did not complete all demographic or physical activity items,

resulting in missing data for certain questionnaire variables. These missing values were not

imputed. Analyses involving those specific variables were conducted using available-case data,

whereas participants' physiological (heart rate, salivary hormones) and psychological

(questionnaire) measures were retained for all other analyses when data quality met the

inclusion criteria.

2.4. Physiological measurements and saliva collection

To evaluate the physiological impact of live sports spectating, we assessed cardiovascular

activity and salivary hormone concentrations (oxytocin and cortisol) across the three

measurement points (pre-game, halftime, post-game). Heart rate (HR) was monitored

continuously using a standalone optical HR sensor worn on the upper arm (Polar Verity Sense;

Polar, Finland), as described in Section 2.2. Participants wore the sensor from approximately 30

minutes before tipoff until 15 minutes after the end of the game. HR data were sampled at 1 Hz

throughout this period. For subsequent analyses, the spectating period was segmented into "in-

game" intervals (when play was in progress and players were on the court) and "out-of-game"

intervals (inter-quarter breaks, halftime, and the 15-minute post-game period).

Salivary oxytocin and cortisol were measured as neuroendocrine markers indexing

social bonding and stress-related responses during spectating. At each time point (pre-game,

halftime, post-game), participants provided approximately 2 mL of whole saliva by passively

drooling through a straw directly into polypropylene collection tubes. Samples were

immediately placed on ice and then stored at $-80\ °C$ until processing. To precipitate mucins and

other debris, frozen samples were thawed and centrifuged at $1500 \times g$ for 20 minutes; the clear

supernatant was then aliquoted and re-frozen at $-80\ °C$ until assay.

For oxytocin quantification, 1000 μL of each saliva sample was concentrated by

centrifugal evaporation and reconstituted according to the manufacturer's instructions before

assay with a commercial competitive enzyme-linked immunosorbent assay (ELISA) kit

(Oxytocin ELISA Kit, Enzo Life Sciences, USA). Salivary cortisol concentration was measured

using a high-sensitivity ELISA kit (Salivary Cortisol ELISA Kit, Salimetrics, USA). All samples

were assayed in duplicate on 96-well plates, and absorbance was read using a multimode

microplate reader (Varioskan LUX, Thermo Fisher Scientific, USA). Hormone concentrations

were calculated from standard curves generated for each plate. Across all assays, the coefficient

of variation for duplicate determinations was below 5%, indicating acceptable analytic

reliability.

2.5. HR data analysis

Heart rate synchrony between spectators was quantified using cross-correlation function (CCF)

analysis of the HR time series [34]. For each participant pair, CCFs were computed within a

temporal lag window of ±30 seconds to identify the maximum correlation between their HR

time series. The temporal lag at which the maximum CCF occurred was extracted, and the

absolute value of this lag was taken as an index of the degree of temporal alignment irrespective

of direction, with smaller values indicating tighter alignment.

Because participants were seated in two separate blocks on opposite sides of the same

courtside, their visual perspectives of the game could differ and potentially influence synchrony.

To account for this, synchrony scores were calculated within each seating block rather than

across all participants. For each participant, an individual synchrony score was obtained by

averaging the dyadic CCF values across all pairings with other participants in the same seating

block.

Baseline HR synchrony was defined as the 5-minute interval during the pre-game

period that showed the lowest within-block CCF values, and synchrony indices during

spectating were expressed relative to this baseline where appropriate. In-game and out-of-game

synchrony measures were derived from the segmented HR data described in Section 2.4,

enabling us to examine whether physiological synchrony differed between active play and non-

play intervals and how these dynamics related to neuroendocrine and psychological measures.

2.6. Psychological questionnaires

Subjective experiences during spectating were assessed using four self-report measures: enjoyment, flow state, intention to revisit, and perceived social unity.

Enjoyment. Momentary enjoyment while watching the game was assessed using a 100-mm visual analogue scale (VAS). The anchors "not enjoyable at all" and "very enjoyable" were printed at the left and right ends, respectively. Participants were instructed to place an "X" on the line at the point that best represented how much they were enjoying the game at that moment. The distance (mm) from the left anchor to the mark was measured and expressed as a percentage of the total line length, with higher scores indicating greater enjoyment.

Flow state. Flow experience during spectating was measured using a brief four-item flow scale developed for this study, which asked participants the extent to which they (1) forgot the time, (2) were completely focused, (3) lost themselves, and (4) enjoyed themselves while watching the game. Each item was rated on a seven-point Likert scale ranging from 1 ("not at all") to 7 ("very much"), and the sum of the four items was used as an index of the subjective flow state during spectating.

Intention to revisit. To capture behavioural intention to re-engage with live sports spectating, we administered a single-item loyalty scale that assessed intention to revisit: "I would like to watch this event again in the future." Responses were given on a seven-point Likert scale from 1 ("not at all") to 7 ("very much"), with higher scores indicating stronger intention to revisit.

Sense of unity (Inclusion of Other in the Self). Perceived social unity with players and fellow spectators was assessed using the Inclusion of Other in the Self (IOS) scale, a widely used pictorial measure of interpersonal closeness [35]. The IOS presents seven pairs of overlapping circles; greater overlap indicates a closer perceived relationship. In this study, participants rated their perceived closeness to four targets: home team players, home team spectators, away team players, and away team spectators. For each target, participants selected the pair of circles that best represented their current sense of unity with that group.

Enjoyment (VAS) and unity (IOS) were assessed at all three time points (pre-game, halftime, post-game) to capture temporal changes across spectating. Flow state and intention to revisit were measured only immediately after the game, reflecting participants' overall experiential and behavioural responses to the event.

2.7 Statistical analysis.

Heart rate synchrony analyses and mediation models were conducted using R statistical

software (version 4.4.0; R Foundation for Statistical Computing, Vienna, Austria). All other

statistical analyses were performed using GraphPad Prism version 10 (GraphPad Software, San

Diego, CA, USA).

Depending on the data structure, group differences and temporal changes were

examined using unpaired t-tests or two-way analyses of variance (ANOVA) with factors for

group (High-OT vs. Low-OT) and time (pre-game, halftime, post-game). When Mauchly's test

indicated violation of sphericity, Greenhouse–Geisser corrections were applied to the degrees of

freedom. When a significant main effect or interaction was detected, Bonferroni's multiple

comparison test was used for post hoc comparisons.

In cases where missing values arose due to incomplete questionnaire responses or

exclusion of physiologically implausible outliers, linear mixed-effects models were employed

instead of repeated-measures ANOVA to maximise the use of available data. For mixed-effects

models, F-statistics and degrees of freedom were used to derive partial η^2 as an approximate

index of effect size. Correlation analyses, for example associations between oxytocin, heart rate

synchrony indices, and psychological measures, were conducted using Pearson's product–

moment correlation coefficient unless otherwise specified.

For mediation analyses, indirect effects were estimated using a nonparametric

bootstrapping procedure with 2000 resamples and bias-corrected 95 per cent confidence

intervals, implemented in the R package lavaan. An indirect effect was considered statistically

significant when the 95 per cent confidence interval did not include zero. All data are presented

as mean ± standard deviation (SD) unless stated otherwise, and the threshold for statistical

significance was set at $p < 0.05$ (two-tailed).

3. Results

3.1. Endogenous oxytocin dynamics shift in a baseline-dependent manner during live sports

spectating

We first tested whether live sports spectating elicits distinct hormonal responses depending on

baseline oxytocin levels. The High-OT group maintained elevated oxytocin concentrations

throughout the game, showing no evidence of further time-dependent change, whereas the Low-

OT group exhibited a significant increase over time that brought post-game oxytocin levels into

the range observed in the High-OT group (High-OT main group effect: $F_{1,58} = 29.06$, $p < 0.0001$,

$\eta p^2 = 0.33$; Low-OT time effect: $F_{2,116} = 11.67$, $p < 0.0001$, $\eta p^2 = 0.16$; Fig. 2A). Cortisol

concentrations decreased across both groups during spectating ($F_{1.8,102.7} = 3.32$, $p = 0.045$, $\eta p^2 =$

0.05; Fig. 2B), resulting in a progressive increase in the oxytocin-to-cortisol ratio ($F_{1.7,95.7} = 14.50$,

$p < 0.0001$, $\eta p^2 = 0.20$; Fig. 2C). Taken together, these findings indicate that sports spectating

modulates hormonal dynamics in a baseline-dependent manner, sustaining high oxytocin levels

in one subgroup and augmenting them in another, while shifting both groups toward an oxytocin-

dominant, low-stress hormonal balance without evidence of stress-related endocrine activation.

3.2. High oxytocin promotes unity through enhanced enjoyment

We next examined how oxytocin dynamics during spectating relate to subjective experiences,

focusing on social bonding and enjoyment. Unity with the home team players and home team

fans increased over time irrespective of baseline oxytocin group (players: $F_{1.48,84.41} = 24.39$, $p <$

0.0001, $\eta p^2 = 0.30$; fans: $F_{1.82,104.10} = 12.36$, $p < 0.0001$, $\eta p^2 = 0.17$; Fig. 3A, B). Enjoyment also

increased in both groups ($F_{1.79,101.70} = 22.18$, $p < 0.0001$, $\eta p^2 = 0.28$), with a more pronounced

effect in the High-OT group than in the Low-OT group (group effect: $F_{1,57} = 6.59$, $p = 0.012$, ηp^2

$= 0.10$; Fig. 3C).

Across all time points, higher oxytocin concentrations were positively associated with

greater enjoyment ($r = 0.21$, $p = 0.006$), stronger unity with players ($r = 0.24$, $p = 0.001$), and

stronger unity with fans ($r = 0.16$, $p = 0.037$; Fig. 3D). Mediation analyses further showed that

oxytocin enhanced perceived unity with players via increased enjoyment (indirect effect: $z = 2.74$,

$p = 0.006$, 95% CI [0.005, 0.023]; Fig. 3E), and similarly enhanced unity with fans through

enjoyment (indirect effect: $z = 2.61$, $p = 0.009$, 95% CI [0.004, 0.022]; Fig. 3E). These results

indicate that baseline-dependent oxytocin dynamics do not merely track social bonding but

contribute to it by amplifying enjoyment during spectating, which in turn promotes social bonding

among spectators.

3.3. Endogenous oxytocin dynamics co-occur with heart rate synchrony

We then investigated whether live sports spectating increases heart rate (HR) synchrony among

spectators and whether this synchrony is associated with oxytocin. Mean HR and peak HR

followed similar trajectories in the High-OT and Low-OT groups, with both indices showing

robust elevations relative to pre-game levels (mean HR: $F_{2.57,146.20} = 84.37$, $p < 0.0001$, $\eta p^2 = 0.59$;

peak HR: $F_{3.14,182.30} = 170.90$, $p < 0.0001$, $\eta p^2 = 0.74$; Fig. 4B, C). Cross-correlation analyses

revealed that HR synchrony, indexed by CCF values, increased significantly during the first and

second halves of the game ($F_{2.97,169.50} = 28.19$, $p < 0.0001$, $\eta p^2 = 0.33$; Fig. 4D). In parallel, the

temporal lag between HR time series shortened during and after the game ($F_{3.03,172.60} = 31.39$, $p <$

0.0001, $\eta p^2 = 0.36$; Fig. 4E), indicating tighter moment-to-moment alignment of cardiac dynamics.

Oxytocin concentrations were positively related to HR synchrony ($r = 0.15$, $p = 0.041$;

Fig. 4F). This association was especially pronounced in the Low-OT group, in which oxytocin

levels were significantly associated with both CCF values ($r = 0.28$, $p = 0.009$) and time-lag

indices ($r = -0.26$, $p = 0.014$; Supplementary Fig. 1B, D). These findings suggest that live sports

spectating promotes HR synchrony among casual spectators regardless of baseline oxytocin level,

and that endogenous oxytocin dynamics, particularly in individuals with lower baseline

concentrations, contribute to the processes supporting this synchrony. Importantly, these effects

were observed in modestly sized collegiate games, indicating that bonding mechanisms based on

HR synchrony generalise beyond the highly fused fan groups typical of professional settings.

3.4. Heart rate synchrony enhances the spectating experience through a sense of unity and

loyalty

Finally, we examined how physiological synchrony relates to social bonding and the overall

spectating experience, in line with the hypothesis that HR synchrony would support psychological

engagement via unity. To characterise the functional role of synchrony, HR data were reclassified

into in-game periods (active play) and out-of-game periods (inter-quarter intervals, halftime, and

post-game). During in-game periods (Fig. 5A), shorter time-lag values between HR time series

were significantly associated with stronger unity toward home team players ($r = -0.45$, $p < 0.001$)

and higher flow state ($r = -0.41$, $p = 0.001$), indicating that tighter moment-to-moment alignment

of cardiac dynamics was linked to more affiliative and immersive responses. During out-of-game

periods (Fig. 5B), increases in CCF relative to pre-game (ΔCCF) were positively correlated with

unity toward home spectators ($r = 0.39$, $p = 0.002$) and with loyalty to the team, indexed by

intention to revisit ($r = 0.27$, $p = 0.044$), suggesting that synchrony outside active play reflects

broader social bonding among spectators.

To test whether social bonding mediated these relationships, we conducted mediation

analyses. In-game, HR synchrony, indexed by shorter time lag, influenced flow state via unity

with players as a mediator (z = −3.32, p < 0.001, 95% CI [−0.69, −0.19]; Fig. 5C). Out-of-game,

∆CCF predicted team loyalty via unity with fellow fans (z = 2.01, p = 0.018, 95% CI [0.52, 4.83];

Fig. 5D). These results suggest that physiological synchrony not only reflects but also facilitates

the formation of social bonds, which in turn shape meaningful psychological experiences such as

immersive flow and loyalty to the team during live spectating, consistent with our hypothesised

pathway from oxytocin-linked HR synchrony to social connection.

378 4. Discussion

Live sports spectating in a modest collegiate setting produced two main findings regarding

endogenous oxytocin, physiological synchrony, and social bonding among casual spectators. First,

oxytocin dynamics during spectating were clearly baseline dependent. Individuals with higher

pre-game oxytocin maintained elevated concentrations throughout the game, whereas those with

lower pre-game oxytocin showed a marked increase that brought their post-game levels into a

similar range (Fig. 2A; Table S1). Across both groups, cortisol decreased and the oxytocin-to-

cortisol ratio increased, indicating a shift toward an oxytocin-dominant, low-stress hormonal

profile rather than a stress-dominated response to the event (Fig. 2B, C).

Second, endogenous oxytocin and heart rate synchrony jointly supported social

connection and experiential outcomes. Higher oxytocin levels were associated with greater

enjoyment and stronger unity with players and fellow fans (Fig. 3C–E). Stronger cardiac

synchrony was likewise associated with greater unity, higher flow, and stronger loyalty to the

team, with distinct in-game and out-of-game relationships (Figs. 4D–F, 5A, B). Mediation

analyses further showed that oxytocin and synchrony facilitated these psychological benefits via

enhanced feelings of unity, such that enjoyment and unity mediated the links between oxytocin

or synchrony and flow or intention to revisit (Figs. 3E, 5C, D). Together, these findings suggest

that live sports spectating can act as a naturalistic, non-pharmacological context in which

endogenous oxytocin and physiological synchrony are recruited to promote social bonding and

positive psychological states in everyday, low-stakes settings.

399 4.2. Baseline-dependent oxytocin dynamics and hormonal profile

The baseline-dependent pattern of oxytocin change observed here complements and extends

previous work on endogenous oxytocin responses to affiliative contexts and physical activity.

Sports spectating promoted oxytocin dynamics in a manner that depended on baseline levels:

individuals with higher pre-game oxytocin maintained elevated concentrations, whereas those

with lower pre-game oxytocin showed marked increases that brought their post-game levels into

a similar range (Fig. 2A; Table S1). Across both groups, cortisol decreased and the oxytocin-to-

cortisol ratio increased (Fig. 2B, C), indicating a shift toward an oxytocin-dominant, low-stress

hormonal profile rather than a stress-dominated response to the event.

Previous research has demonstrated that oxytocin release is facilitated by social

experiences such as eye contact, shared narratives, and storytelling [36,37]. Our findings

suggest that similar mechanisms may be engaged during sports spectating. Elements such as

indirect eye contact mediated through players, the embodied performance conveyed by athletes,

and the unscripted dramatic nature of the game share key features with storytelling. The

collective enjoyment of these elements, as a fundamental human social practice, may serve as a

form of "embodied storytelling" that enhances oxytocin dynamics and provides a basis for

strengthening social bonds among spectators. In this sense, live sports spectating extends a

broader class of everyday affiliative activities, including shared narratives, mutual gaze, and

coordinated movement, that are known to upregulate endogenous oxytocin without

pharmacological manipulation, offering a behaviourally embedded means of engaging the

oxytocinergic system [38–41,37,42].

Mediation analyses further demonstrated that oxytocin fostered unity with players and

fellow fans through enhanced enjoyment (Fig. 3E). Taken together, these results align with

evidence indicating modest but reliable associations between endogenous oxytocin and social

interaction, together with substantial inter-individual heterogeneity and baseline-dependent

responsivity [43,44]. The concurrent decrease in cortisol and increase in the oxytocin-to-cortisol

ratio also contrast with the cortisol elevations typically observed in competitive athletes and

highly invested fans during high-stakes events [45–47], suggesting that modest collegiate

spectating can support social engagement within a hormonal milieu that avoids pronounced

stress-related endocrine costs.

4.3. Neural mechanisms and translational implications of oxytocin dynamics

The observed mediation, whereby oxytocin enhanced unity via enjoyment (Fig. 3E), may be

accounted for by the neural mechanism through which oxytocin potentiates activity in the

brain's reward system. Oxytocin receptors are abundantly expressed in regions central to reward

processing, including the ventral tegmental area (VTA) and nucleus accumbens (NAc) [48].

Animal studies have shown that oxytocinergic neurons in the hypothalamic paraventricular

nucleus project directly to the VTA, where they enhance dopaminergic signalling via oxytocin

receptors [49]. Converging evidence in humans indicates that intranasal oxytocin increases

activation of the VTA and NAc when participants view a partner's face or other socially salient

cues [50]. These findings support the notion that oxytocin enhances the valuation of social

stimuli through reward-related pathways. While sports spectating has long been known to

engage the reward system [51], the neural basis of social bonding during collective spectating

has remained elusive. Our results suggest that elevated oxytocin during sports spectating may

amplify reward system activity as a form of social reward, thereby providing a neural

mechanism for fostering unity with players and fellow fans.

Beyond its actions on mesolimbic structures underlying the reward system, oxytocin

may also influence higher-order regions of the prefrontal cortex. A prominent candidate is the

ventromedial prefrontal cortex (vmPFC). The vmPFC receives projections from the VTA and

NAc and is well established as a core component of the reward system [52], yet it is also

engaged by others' successes and socially salient events, implicating it in vicarious reward

processing [53]. Recent evidence further indicates that oxytocin enhances vmPFC activity in

response to socially salient cues [54,55]. Thus, the facilitation of oxytocin dynamics during

sports spectating may strengthen unity with players and fellow fans through modulation of

mesolimbic and vmPFC function, linking shared emotional engagement with neuroendocrine

and neural mechanisms of social reward.

From a translational perspective, these baseline-dependent and low-stress oxytocin

dynamics support the idea that live sports spectating may serve as a naturalistic, non-

pharmacological means of recruiting endogenous oxytocin in everyday life. Such an approach

could complement pharmacological oxytocin administration by targeting the same neuropeptide

system through behavioural rather than exogenous routes, potentially helping to address some of

the challenges related to efficacy, dosing, and long-term safety that have emerged in clinical

trials of intranasal oxytocin [56,7,5,8,57,11].

4.4. Heart rate synchrony as a component of collective experience

This study also demonstrated that live sports spectating promotes heart rate (HR) synchrony

among spectators irrespective of baseline oxytocin levels (Fig. 4D, E). Such synchrony is likely

driven by the intense collective arousal elicited during spectating. Shared external stimuli, including the tension of the game, the roar of the crowd, and collective reactions, act as powerful forces that align cardiac dynamics across individuals, transcending individual hormonal baselines. Physiological synchrony has similarly been observed in other collective events such as music concerts and religious rituals, where it is regarded as a physiological substrate of shared group experiences [58,59]. Our findings suggest that sports spectating shares this property, functioning like other collective gatherings in fostering synchrony among casual fans.

In line with this literature, HR synchrony increased markedly during the game in our sample, with shorter time lags and higher cross-correlation values indicating tighter temporal alignment of cardiac dynamics among spectators (Fig. 4D, E). Importantly, the present study extends prior work by demonstrating functionally distinct links between synchrony and psychological outcomes in different phases of the event: during in-game periods, higher synchrony, indexed by shorter time lags, was associated specifically with stronger unity toward players and greater flow (Fig. 5A); during out-of-game periods, increases in synchrony relative to pre-game were related to unity with fellow spectators and a stronger intention to revisit (Fig. 5B). These phase-specific associations suggest that synchrony during active play is especially tied to immersive, player-focused engagement, whereas synchrony during breaks and post-game is more closely related to bonding within the spectator community and subsequent loyalty-related motivation.

4.5. Coupling between oxytocin, synchrony, and social experience

Another key finding of this study is the observed association between endogenous oxytocin levels and HR synchrony (Fig. 4F). Although both oxytocin release and physiological synchrony are facilitated in collective events and social interactions such as eye contact and touch [58,36,37], their direct coupling has rarely been tested. Experimental work shows that exogenous oxytocin can increase physiological linkage in small groups during cooperative tasks, and research in parent–infant dyads links endogenous oxytocin to interactive synchrony [60,61]. To our knowledge, however, no study has directly linked moment-to-moment fluctuations in endogenous oxytocin to physiological synchrony in a large, naturalistic spectator setting. Evidence that intranasal oxytocin enhances synchrony in laboratory cooperation tasks is therefore mechanistically consistent with our field data [61,62], yet the present association in

live sports extends these insights to ecologically valid group contexts.

Mediation analyses further showed that unity statistically mediated the relationships

between synchrony and experiential outcomes (Fig. 5C, D). Synchrony enhanced flow via unity

with players during active play, and enhanced team loyalty via unity with fellow fans during

breaks and post-game. Together with the observation that higher oxytocin levels co-occurred

with stronger synchrony and social bonding (Figs. 3D, 4F), these findings support accounts in

which endogenous oxytocin facilitates shared attention, emotional convergence, and alignment

of autonomic states. In this view, oxytocin-linked synchrony can promote both on-court

immersion and off-court social affiliation in spectating contexts. Taken together, our findings

suggest a mutually reinforcing relationship between oxytocin and HR synchrony, situating

sports spectating alongside other collective gatherings as a naturalistic context that both elicits

physiological synchrony and tracks with endogenous oxytocin dynamics.

4.6. Broader social and clinical implications

These observations carry translational implications. Oxytocin has long been considered a

promising pharmacological candidate for improving social functioning in autism spectrum

disorder (ASD) and related conditions, yet large multisite randomised trials have yielded null or

modest effects on core social symptoms when intranasal oxytocin is administered over months

[8]. In parallel, safety and methods reviews emphasise uncertainties regarding nose-to-brain

delivery, individual response profiles, and potential peripheral off-target effects, cautioning

against viewing exogenous oxytocin as a simple "social drug" [57]. By contrast, our data show

that a brief, enjoyable, and widely accessible group activity, live sports spectating, can engage

endogenous oxytocin dynamics and oxytocin-linked HR synchrony in a naturalistic setting,

without pharmacological exposure or stress-related hormonal costs (Figs. 2–5). At the

population level, sports spectating has been associated with higher well-being and lower

loneliness [46,63,64], and ASD research indicates difficulties in spontaneous interpersonal

synchrony [65,66], suggesting that synchrony-supportive environments could serve as low-

burden adjuncts to social habilitation. While causal clinical benefits remain to be established,

the present findings raise the possibility that inclusive spectator contexts might help harness

endogenous oxytocin-linked synchrony to scaffold social engagement in populations with

social-communication difficulties.

Beyond their immediate relevance to oxytocin biology and physiological synchrony,

these findings also inform broader discussions about how everyday collective activities support

social well-being. Social psychological models of sport fandom propose that team identification

and regular spectatorship enhance well-being by providing opportunities to gain and maintain

social connections with other fans [29,30,28]. From this perspective, live sports spectating can

be viewed as a form of embodied storytelling, in which the embodied performances of players,

the indirect eye contact created by shared attention to the court, and the unscripted drama of the

game are collectively experienced and emotionally co-constructed [67,36,37]. Historically,

large-scale collective practices ranging from religious rituals and martial displays to seasonal

festivals and stadium sports have provided embodied arenas for synchronising bodies and

emotions, with modern spectator sports representing a contemporary expression of this long-

standing human tendency toward coordinated collective experience [58,68,67,59].

Against the backdrop of Japanese collegiate sports, these findings carry particular

significance. In the United States, collegiate sports exert far-reaching social influence, supported

by strong athletic performance, substantial financial resources, and extensive entertainment

infrastructures. By contrast, Japanese collegiate sports have not yet achieved comparable

popularity or institutional support [69]. Nevertheless, our study revealed that even in the

relatively local and small-scale context of Japanese university basketball, spectators

synchronised, forged social bonds, and developed stronger intentions to revisit (Fig. 5C, D).

These results indicate that the social benefits of spectating are not confined to professional

sports on grand stages. Rather, they can also emerge from sharing the efforts of familiar

individuals, potentially providing a socio-physiological foundation for community building and

the promotion of well-being in local societies through sports.

Emerging formats of collective viewing, such as e-sports arenas, large-scale online

streaming events, and bio-sharing interfaces that transmit physiological signals between remote

spectators, may reproduce or transform the embodied storytelling processes identified here. For

example, real-time sharing of bodily signals such as facial expressions and HR in e-sports has

been shown to enhance the perceived social presence of others [70]. This suggests that

embodiment, a universally available human resource, serves as a foundation for creating social

connections. Future work should explore how the sharing of embodiments across time and

space can be harnessed to promote collective well-being via endogenous oxytocin dynamics and

oxytocin-linked synchrony in increasingly hybrid physical–digital environments.

4.7. Limitations and future directions

While our field-based approach ensured ecological validity by capturing real-time physiological dynamics during live sports events, several limitations should be acknowledged. First, the determinants of individual differences in baseline oxytocin levels remain unclear. In this study, participants were categorised based on the median pre-game oxytocin concentration, and we verified that this classification was not attributable to physical characteristics such as body composition, resting HR, or habitual physical activity (Table S1). Nonetheless, it remains uncertain whether the observed variability reflects other stable individual traits or transient influences such as recent social interactions.

Second, oxytocin was measured in saliva, leaving the dynamics of oxytocin within the central nervous system unresolved. Salivary oxytocin tracks peripheral oxytocin to some extent: in mothers, concurrently sampled saliva and plasma concentrations covary across tasks, and a preregistered meta-analysis reports modest central–peripheral coupling that emerges after experimental stimulation rather than at rest [71,72]. In humans, plasma oxytocin has been shown to predict cerebrospinal-fluid oxytocin in a lumbar-puncture study [73], although other reports find no baseline association, underscoring ongoing measurement debate [74]. Accordingly, while salivary assays offer a practical window on the oxytocinergic system in the field, inference to brain dynamics remains indirect. Future studies should combine validated peripheral sampling with neuroimaging and, where feasible, pharmacological challenges to delineate the neural mechanisms by which sports spectating fosters social bonding [12].

Third, our analyses focused on spectator dyads sharing similar attributes, such as seating location and game exposure. Whether HR synchrony also occurs between spectators of different demographics, nationalities, or group affiliations remains unknown. Future research should extend these investigations to diverse spectator pairings and, importantly, to synchrony between spectators and players. Finally, the observational design without a non-spectating control group precludes strong causal inferences about the effects of live sports spectating per se, and unmeasured situational factors, including team performance expectations, social relationships among co-attendees, or pre-existing fan loyalty, may have contributed to both hormonal and psychological responses. Combining experimental or quasi-experimental manipulations with more diverse samples, repeated events, and longitudinal follow-up will be essential for testing whether oxytocin-linked synchrony during live spectating prospectively predicts sustained changes in social connection, mental health, and engagement in community

activities.

5. Conclusion

This study showed that a single session of live sports spectating in a real-world collegiate

setting can recruit endogenous oxytocin and interpersonal heart rate synchrony in ways that

support social bonding among casual fans. Spectators with low baseline oxytocin exhibited clear

increases in oxytocin and a shift toward an oxytocin-dominant, low-stress hormonal profile,

while higher oxytocin levels and stronger heart rate synchrony across participants were

associated with greater enjoyment, stronger unity with players and fellow fans, and higher flow

and intention to revisit. These findings suggest that live sports spectating may serve as a simple,

scalable non-pharmacological approach to engaging oxytocin-related systems and interpersonal

synchrony to promote social connection in everyday life. Such contexts may in turn encourage

sustained community engagement and contribute to more active, healthier, and more socially

connected lifestyles.

Acknowledgements

We would like to thank all staff at the Bureau of Physical Education and Sports, University of

Tsukuba, for their dedicated efforts in organizing the home series event "Tsukuba Live!" This

research was supported by a part of 'Sports in Life' project by the Japan Sports Agency to K.K.

and T.M.; a Special Research Grant from the TOYOTA Foundation to T.M. (D20-ST-0034); a

Fusion Oriented Research for disruptive Science and Technology (FOREST) by Japan Science

and Technology Agency (JST) to T.M. (JPMJFR205M).

Data availability

The data supporting the findings of this study are available from the corresponding author upon

reasonable request.

Author contributions

T.M., T.Y., and K.K. conceived and designed the study. T.M., T.Y., S.T., H.M. S.D., K.K., S.Y.,

and H.T. recruited participants. T.M., T.Y., D.F., S.T., H.M., S.D., and K.K. collected the data.

T.M., T.Y., D.F., S.T., H.M., and S.D. conducted the data analysis and interpreted the data. T.M.

and T.Y. drafted the manuscript. T.M., T.Y., D.F., S.T., S.D., K.K., H.T. edited and revised the

manuscript. All authors have read and approved the final version of the manuscript.

Competing interests

The authors declare no competing interests.

Declaration of generative AI in scientific writing.

During the preparation of this work the authors used ChatGPT for only English editing. After

using this service, the authors reviewed and edited the content as needed and took full

responsibility for the content of the published article.

References

1. Goh KK, Chen C-H, Lane H-Y. Oxytocin in Schizophrenia: Pathophysiology and
Implications for Future Treatment. Int J Mol Sci. 2021;22:2146.

2. Feifel D, Shilling PD, MacDonald K. A Review of Oxytocin's Effects on the Positive,
Negative, and Cognitive Domains of Schizophrenia. Biol Psychiatry. 2016;79:222–233.

3. Meyer-Lindenberg A, Domes G, Kirsch P, Heinrichs M. Oxytocin and vasopressin in the
human brain: social neuropeptides for translational medicine. Nat Rev Neurosci.
2011;12:524–538.

4. Yao S, Kendrick KM. Effects of Intranasal Administration of Oxytocin and Vasopressin on
Social Cognition and Potential Routes and Mechanisms of Action. Pharmaceutics.
2022;14:323.

5. Quintana DS, Lischke A, Grace S, Scheele D, Ma Y, Becker B. Advances in the field of
intranasal oxytocin research: lessons learned and future directions for clinical research. Mol
Psychiatry. 2021;26:80–91.

6. Evans SL, Dal Monte O, Noble P, Averbeck BB. Intranasal oxytocin effects on social
cognition: a critique. Brain Res. 2014;1580:69–77.

7. Griffiths JL, Mishaal RA, Nabetani M, Goldman RD. Oxytocin for the treatment of autism
spectrum disorder in children. Can Fam Physician Med Fam Can. 2022;68:103–105.

8. Sikich L, Kolevzon A, King BH, McDougle CJ, Sanders KB, Kim S-J, et al. Intranasal
Oxytocin in Children and Adolescents with Autism Spectrum Disorder. N Engl J Med.
2021;385:1462–1473.

9. Bürkner P-C, Williams DR, Simmons TC, Woolley JD. Intranasal Oxytocin May Improve
High-Level Social Cognition in Schizophrenia, But Not Social Cognition or Neurocognition
in General: A Multilevel Bayesian Meta-analysis. Schizophr Bull. 2017;43:1291–1303.

10. Veening JG, Olivier B. Intranasal administration of oxytocin: behavioral and clinical effects,
a review. Neurosci Biobehav Rev. 2013;37:1445–1465.

11. MacDonald E, Dadds MR, Brennan JL, Williams K, Levy F, Cauchi AJ. A review of safety,
side-effects and subjective reactions to intranasal oxytocin in human research.
Psychoneuroendocrinology. 2011;36:1114–1126.

12. Tabak BA, Leng G, Szeto A, Parker KJ, Verbalis JG, Ziegler TE, et al. Advances in human
oxytocin measurement: challenges and proposed solutions. Mol Psychiatry. 2023;28:127–
140.

13. Bartz JA. Oxytocin and the Pharmacological Dissection of Affiliation. Curr Dir Psychol Sci.
2016;25:104–110.

14. Handlin L, Novembre G, Lindholm H, Kämpe R, Paul E, Morrison I. Human endogenous
oxytocin and its neural correlates show adaptive responses to social touch based on recent
social context. eLife. 2023;12:e81197.

15. Uvnäs-Moberg K, Handlin L, Petersson M. Self-soothing behaviors with particular reference
to oxytocin release induced by non-noxious sensory stimulation. Front Psychol. 2015;5.

16. Li Q, Becker B, Wernicke J, Chen Y, Zhang Y, Li R, et al. Foot massage evokes oxytocin
release and activation of orbitofrontal cortex and superior temporal sulcus.
Psychoneuroendocrinology. 2019;101:193–203.

17. Rassovsky Y, Harwood A, Zagoory-Sharon O, Feldman R. Martial arts increase oxytocin
production. Sci Rep. 2019;9:12980.

18. Harwood-Gross A, Feldman R, Zagoory-Sharon O, Rassovsky Y. Hormonal reactivity during
martial arts practice among high-risk youths. Psychoneuroendocrinology. 2020;121:104806.

19. Olff M, Frijling JL, Kubzansky LD, Bradley B, Ellenbogen MA, Cardoso C, et al. The role
of oxytocin in social bonding, stress regulation and mental health: An update on the
moderating effects of context and interindividual differences. Psychoneuroendocrinology.
2013;38:1883–1894.

20. Seltzer LJ, Ziegler TE, Pollak SD. Social vocalizations can release oxytocin in humans. Proc
Biol Sci. 2010;277:2661–2666.

21. Schneiderman I, Zagoory-Sharon O, Leckman JF, Feldman R. Oxytocin during the initial

stages of romantic attachment: Relations to couples' interactive reciprocity.

Psychoneuroendocrinology. 2012;37:1277–1285.

22. Burenkova OV, Dolgorukova TA, An I, Kustova TA, Podturkin AA, Shurdova EM, et al.

Endogenous oxytocin and human social interactions: A systematic review and meta-analysis.

Psychol Bull. 2023;149:549–579.

23. Crockford C, Deschner T, Ziegler TE, Wittig RM. Endogenous peripheral oxytocin measures

can give insight into the dynamics of social relationships: a review. Front Behav Neurosci.

2014;8:68.

24. Chung V, Mennella R, Pacherie E, Grezes J. Social bonding through shared experiences: the

role of emotional intensity. R Soc Open Sci. 2024;11:240048.

25. Baranowski-Pinto G, Profeta VLS, Newson M, Whitehouse H, Xygalatas D. Being in a crowd

bonds people via physiological synchrony. Sci Rep. 2022;12:613.

26. Flory S, Guglielmini S, Scholkmann F, Marcar VL, Wolf M. How our hearts beat together: a

study on physiological synchronization based on a self-paced joint motor task. Sci Rep.

2023;13:11987.

27. Behrens F, Snijdewint JA, Moulder RG, Prochazkova E, Sjak-Shie EE, Boker SM, et al.

Physiological synchrony is associated with cooperative success in real-life interactions. Sci

Rep. 2020;10:19609.

28. Wann DL, Waddill PJ, Polk J, Weaver S. The team identification–social psychological health

model: Sport fans gaining connections to others via sport team identification. Group Dyn

Theory Res Pract. 2011;15:75–89.

29. Inoue Y, Lock D, Sato M, Funk DC. Psychological processes connecting team identification

and social well-being for middle-aged and older adults: moderated mediation of subjective

and objective on-field performance. Sport Manag Rev. 2022;25:207–233.

30. Inoue Y, Sato M, Filo K, Du J, Funk DC. Sport Spectatorship and Life Satisfaction: A

Multicountry Investigation. J Sport Manag. 2017;31:419–432.

31. Pepping G-J, Timmermans EJ. Oxytocin and the biopsychology of performance in team

sports. ScientificWorldJournal. 2012;2012:567363.

32. Gonzalez-Bono E, Salvador A, Serrano MA, Ricarte J. Testosterone, Cortisol, and Mood in

a Sports Team Competition. Horm Behav. 1999;35:55–62.

33. Comes-Fayos J, Blanco-Gandía MC, Moreno IR, Rodríguez-Arias M, Lila M, Sarrate-Costa

C, et al. Reduced salivary oxytocin after an empathic induction task in Intimate Partner

722  Violence perpetrators: Importance of socio-affective functions and its impact on prosocial

723  behavior. Psychoneuroendocrinology. 2022;137:105644.

724 34. daSilva EB, Wood A. How and Why People Synchronize: An Integrated Perspective.

725  Personal Soc Psychol Rev. 2025;29:159–187.

726 35. Aron A, Aron EN, Smollan D. Inclusion of Other in the Self Scale and the structure of

727  interpersonal closeness. J Pers Soc Psychol. 1992;63:596–612.

728 36. Brockington G, Gomes Moreira AP, Buso MS, Gomes da Silva S, Altszyler E, Fischer R, et

729  al. Storytelling increases oxytocin and positive emotions and decreases cortisol and pain in

730  hospitalized children. Proc Natl Acad Sci. 2021;118:e2018409118.

731 37. Nagasawa M, Mitsui S, En S, Ohtani N, Ohta M, Sakuma Y, et al. Oxytocin-gaze positive

732  loop and the coevolution of human-dog bonds. Science. 2015;348:333–336.

733 38. Eather N, Wade L, Pankowiak A, Eime R. The impact of sports participation on mental health

734  and social outcomes in adults: a systematic review and the 'Mental Health through Sport'

735  conceptual model. Syst Rev. 2023;12:102.

736 39. Hernández-Mendo A, Reigal RE, López-Walle JM, Serpa S, Samdal O, Morales-Sánchez V,

737  et al. Physical Activity, Sports Practice, and Cognitive Functioning: The Current Research

738  Status. Front Psychol. 2019;10.

739 40. Malm C, Jakobsson J, Isaksson A. Physical Activity and Sports—Real Health Benefits: A

740  Review with Insight into the Public Health of Sweden. Sports. 2019;7:127.

741 41. Ostrowsky MK. Sports Fans, Alcohol Use, and Violent Behavior: A Sociological Review.

742  Trauma Violence Abuse. 2018;19:406–419.

743 42. Ostrowsky MK. The social psychology of alcohol use and violent behavior among sports

744  spectators. Aggress Violent Behav. 2014;19:303–310.

745 43. Kawakami R, Kitano N, Fujii Y, Jindo T, Kai Y, Arao T. Association of watching sports games

746  with subsequent health and well-being among adults in Japan: An outcome-wide

747  longitudinal approach. Prev Med. 2024;189:108154.

748 44. Tsuji T, Kanamori S, Watanabe R, Yokoyama M, Miyaguni Y, Saito M, et al. Watching sports

749  and depressive symptoms among older adults: a cross-sectional study from the JAGES 2019

750  survey. Sci Rep. 2021;11:10612.

751 45. Kinoshita K, Nakagawa K, Sato S. Watching sport enhances well-being: evidence from a

752  multi-method approach. Sport Manag Rev. 2024;27:595–619.

753 46. Newson M, Shiramizu V, Buhrmester M, Hattori W, Jong J, Yamamoto E, et al. Devoted fans

release more cortisol when watching live soccer matches. Stress Health J Int Soc Investig

Stress. 2020;36:220–227.

47. Cikara M, Botvinick MM, Fiske ST. Us versus them: social identity shapes neural responses

to intergroup competition and harm. Psychol Sci. 2011;22:306–313.

48. Dölen G, Darvishzadeh A, Huang KW, Malenka RC. Social reward requires coordinated

activity of nucleus accumbens oxytocin and serotonin. Nature. 2013;501:179–184.

49. Hung LW, Neuner S, Polepalli JS, Beier KT, Wright M, Walsh JJ, et al. Gating of social

reward by oxytocin in the ventral tegmental area. Science. 2017;357:1406–1411.

50. Scheele D, Wille A, Kendrick KM, Stoffel-Wagner B, Becker B, Güntürkün O, et al. Oxytocin

enhances brain reward system responses in men viewing the face of their female partner.

Proc Natl Acad Sci. 2013;110:20308–20313.

51. Duarte IC, Afonso S, Jorge H, Cayolla R, Ferreira C, Castelo-Branco M. Tribal love: the

neural correlates of passionate engagement in football fans. Soc Cogn Affect Neurosci.

2017;12:718–728.

52. Haber SN, Kunishio K, Mizobuchi M, Lynd-Balta E. The orbital and medial prefrontal circuit

through the primate basal ganglia. J Neurosci Off J Soc Neurosci. 1995;15:4851–4867.

53. Lockwood PL. The anatomy of empathy: Vicarious experience and disorders of social

cognition. Behav Brain Res. 2016;311:255–266.

54. Gangopadhyay P, Chawla M, Dal Monte O, Chang SWC. Prefrontal-amygdala circuits in

social decision-making. Nat Neurosci. 2021;24:5–18.

55. Aoki Y, Watanabe T, Abe O, Kuwabara H, Yahata N, Takano Y, et al. Oxytocin's

neurochemical effects in the medial prefrontal cortex underlie recovery of task-specific

brain activity in autism: a randomized controlled trial. Mol Psychiatry. 2015;20:447–453.

56. Audunsdottir K, Sartorius AM, Kang H, Glaser BD, Boen R, Nærland T, et al. The effects of

oxytocin administration on social and routinized behaviors in autism: A preregistered

systematic review and meta-analysis. Psychoneuroendocrinology. 2024;167:107067.

57. Leng G, Ludwig M. Intranasal Oxytocin: Myths and Delusions. Biol Psychiatry.

2016;79:243–250.

58. Tschacher W, Greenwood S, Ramakrishnan S, Tröndle M, Wald-Fuhrmann M, Seibert C, et

al. Audience synchronies in live concerts illustrate the embodiment of music experience.

Sci Rep. 2023;13:14843.

59. Konvalinka I, Xygalatas D, Bulbulia J, Schjødt U, Jegindø E-M, Wallot S, et al. Synchronized

arousal between performers and related spectators in a fire-walking ritual. Proc Natl Acad
Sci. 2011;108:8514–8519.

60. Minagawa Y, Hata M, Yamamoto E, Tsuzuki D, Morimoto S. Inter-brain synchrony during
mother-infant interactive parenting in 3-4-month-old infants with and without an elevated
likelihood of autism spectrum disorder. Cereb Cortex N Y N 1991. 2023;33:11609–11622.

61. Thorson KR, McKernan SM, West TV, Woolley JD, Mendes WB, Stauffer CS. Oxytocin
increases physiological linkage during group therapy for methamphetamine use disorder: a
randomized clinical trial. Sci Rep. 2021;11:21004.

62. Mu Y, Guo C, Han S. Oxytocin enhances inter-brain synchrony during social coordination in
male adults. Soc Cogn Affect Neurosci. 2016;11:1882–1893.

63. Keyes H, Gradidge S, Gibson N, Harvey A, Roeloffs S, Zawisza M, et al. Attending live
sporting events predicts subjective wellbeing and reduces loneliness. Front Public Health.
2023;10.

64. van der Meij L, Almela M, Hidalgo V, Villada C, Ijzerman H, van Lange PAM, et al.
Testosterone and cortisol release among Spanish soccer fans watching the 2010 World Cup
final. PloS One. 2012;7:e34814.

65. Carnevali L, Valori I, Mason G, Altoè G, Farroni T. Interpersonal motor synchrony in autism:
a systematic review and meta-analysis. Front Psychiatry. 2024;15.

66. Glass D, Yuill N. Social motor synchrony in autism spectrum conditions: A systematic review.
Autism Int J Res Pract. 2024;28:1638–1653.

67. Sosis R, Kiper J. Sport as a Meaning-Making System: Insights from the Study of Religion.
Religions. 2022;13:915.

68. Pizarro JJ, Zumeta LN, Bouchat P, Włodarczyk A, Rimé B, Basabe N, et al. Emotional
processes, collective behavior, and social movements: A meta-analytic review of collective
effervescence outcomes during collective gatherings and demonstrations. Front Psychol.
2022;13.

69. McCollough B. Effort to Americanize and monetize college sports in Japan faces obstacles.
Los Angel Times. 2020. https://www.latimes.com/sports/story/2020-01-29/effort-
americanize-athletics-japanese-college-campuses-faces-obstacles. Accessed 5 January
2026.

70. Hassan M, Kennard M, Yoshitake S, Ishac K, Takahashi S, Kim S, et al. Augmenting the
Sense of Social Presence in Online Video Games Through the Sharing of Biosignals. IEEE

Access. 2024;12:98977–98989.

71. Valstad M, Alvares GA, Egknud M, Matziorinis AM, Andreassen OA, Westlye LT, et al. The
correlation between central and peripheral oxytocin concentrations: A systematic review
and meta-analysis. Neurosci Biobehav Rev. 2017;78:117–124.

72. Grewen KM, Davenport RE, Light KC. An investigation of plasma and salivary oxytocin
responses in breast- and formula-feeding mothers of infants. Psychophysiology.
2010;47:625–632.

73. Carson DS, Berquist SW, Trujillo TH, Garner JP, Hannah SL, Hyde SA, et al. Cerebrospinal
fluid and plasma oxytocin concentrations are positively correlated and negatively predict
anxiety in children. Mol Psychiatry. 2015;20:1085–1090.

74. Kagerbauer SM, Martin J, Schuster T, Blobner M, Kochs EF, Landgraf R. Plasma oxytocin
and vasopressin do not predict neuropeptide concentrations in human cerebrospinal fluid. J
Neuroendocrinol. 2013;25:668–673.

Figure 1 The experimental protocol.

A. A typical photo of the spectating state. **B.** A typical photo of saliva sampling, questionnaire

completion, and the HR measurement device. **C.** Overview of the experimental protocol. The participants

watch basketball games for approximately 100 minutes. Saliva samples and mood questionnaires were

administered before. Each measurement took approximately 5 minutes to complete. The HR was

measured during spectating.

Figure 2 Endogenous oxytocin dynamics shift in a baseline-dependent manner during live sports spectating.

A. saliva oxytocin levels. **B.** saliva cortisol levels. **C.** oxytocin/cortisol ratio. Standard divisions are shown as red (High-OT) bands and blue (Low-OT) on the graph. Results of two-way ANOVA are shown in the lower left of each graph. Time: main effect of watching time. Group: main effect of the group. Interaction: interaction between time and group. *: $p < 0.05$ vs. pre game for both groups; #: $p < 0.05$ vs. Low-OT.

Figure 3 High oxytocin promotes unity through enhanced enjoyment.

A, B. A sense of unity with home players (A) and home team fans (B). **C**. Enjoyment. Standard divisions are shown as red (High-OT) bands and blue (Low-OT) on the graph. Results of two-way ANOVA are shown in the lower left of each graph. Time: main effect of watching time. Group: main effect of the group. Interaction: interaction between time and group. *: $p < 0.05$ vs. pre game for both groups; #: $p < 0.05$ vs. Low-OT. **D**. Correlation matrix summarizing associations between neuroendocrine responses, mood and a sense of unity. **E**. Mediation model showing the associations between oxytocin, enjoyment, and a sense of unity. *: $p < 0.05$.

Figure 4 Endogenous oxytocin dynamics co-occur with HR synchrony.

A. Typical HR data. **B.** Average HR. **C**. Peak HR. **D.** Similarity of HR dynamics (cross-correlation

function; CCF). **E.** Time lag. Standard divisions are shown as red (High-OT) bands and blue (Low-OT)

on the graph. Results of two-way ANOVA are shown in the lower left of each graph. Time: main effect of

watching time. Group: main effect of the group. Interaction: interaction between time and group. *: p <

0.05 vs. pre game for both groups. **F**. Correlation between oxytocin and HR synchrony. The Pearson's

product-rate correlation coefficient and the results of the single regression analysis are shown in the

upper-left corner of each graph. The computed regression line and 95% confidence intervals are shown as

bands on the graph. *: p < 0.05.

Figure 5 HR synchrony enhances social connection through a sense of unity.

A. Correlation matrix between measurements related to in-game. **B.** Correlation matrix between

measurements related to out of game. **C.** Mediation model of in-game. **D.** Mediation model of out of

game. The significance of the correlation analysis is indicated at the top-right of correlation coefficient. *

$p < 0.05$.

Table S1 Participants' characteristics

	High-OT (n = 30)		Low-OT (n = 30)		p	d
Median oxytocin (pg/ml)		27.5				
Baseline oxytocin levels (pg/ml)	31.9 ±	9.2[#]	21.6 ±	3.7	<0.01	2.46
Age (years)	22.2 ±	8.6	22.9 ±	9.0	0.77	0.07
Gender (n, Male/Female)	20 /	10	19 /	11		
Height (cm)	172.3 ±	9.8	168.6 ±	7.4	0.11	0.42
Body weight (kg)	68.6 ±	15.8	65.3 ±	8.8	0.34	0.25
BMI (kg/m^2)	23.3 ±	1.2	22.9 ±	3.3	0.71	0.11
Resting HR (bpm)	63.2 ±	8.5	67.1 ±	9.4	0.10	0.43
Vigorous-intensity activity (min/day)	87.3 ±	82.6	80.7 ±	80.8	0.76	0.08
Moderate-intensity activity (min/day)	16.8 ±	18.9	20.4 ±	39.7	0.66	0.11
Walking time (min/day)	24.1 ±	29.6	28.1 ±	40.3	0.67	0.11
Exercise time (min/day)	70.0 ±	67.7	70.5 ±	83.3	0.97	<0.01
Sitting time (min/day)	348.6 ±	226.6	284.5 ±	160.9	0.23	0.32
Cheering for home team (score)	6.6 ±	0.5	6.2 ±	0.9	0.10	0.43

Values are presented as mean ± SD. #: p < 0.05 vs. Low-OT.

Figure S1 Stronger relationship between HR synchrony and oxytocin in the Low-OT group.

A, B. Correlation between oxytocin and HR synchrony (CCF) in High-OT group (A), and Low-OT group

(B). **C, D**. Correlation between oxytocin and HR synchrony (time lag) in High-OT group (C), and Low-

OT group (D). The Pearson's product-rate correlation coefficient and the results of the single regression

analysis are shown in the upper-left corner of each graph. The computed regression line and 95%

confidence intervals are shown as bands on the graph. *: $p < 0.05$.

Figure S2 Graphical model of how sports spectating may foster enjoyment-based social connection

via socio-physiological responses.

Sports spectating promotes baseline-dependent oxytocin dynamics: individuals with high baseline

oxytocin maintain elevated concentrations, whereas those with low baseline levels exhibit increases

during spectating. These oxytocin changes, in turn, facilitate heart rate synchrony regardless of baseline

levels, and the two processes may reciprocally reinforce each other. Such socio-physiological bases

ultimately foster enjoyment-based social connection among spectators.