Effects of TMS on the decoding and electrophysiology of priority in working memory

Jacqueline M. Fulvio[1] & Bradley R. Postle[1,2]

[1]Department of Psychology, University of Wisconsin–Madison
[2]Department of Psychiatry, University of Wisconsin–Madison

Corresponding author:
Jacqueline M. Fulvio, PhD
1202 West Johnson St.
Madison, WI 53706-1611
(608) 262-1040
jacqueline.fulvio@wisc.edu

Abstract

The flexible control of working memory (WM) requires prioritizing immediately task-relevant information while maintaining information with potential future relevance in a deprioritized state. Using double-serial retrocuing (DSR) with simultaneous EEG recording, we investigated how single pulses of transcranial magnetic stimulation (spTMS) to right intraparietal sulcus impacts neural representations of unprioritized memory items (UMI), relative to irrelevant memory items (IMI) that are no longer needed for the trial. Twelve human participants (8 female) performed DSR plus a single-retrocue task, while spTMS was delivered during delay periods. Multivariate pattern analysis revealed that spTMS restored decodability of the UMI concurrent with stimulation, and that of the IMI several timesteps later, after the evoked effects of spTMS were no longer present in the EEG signal. This effect was carried by the alpha (8-13 Hz) and low-beta (13-20 Hz) frequency bands. Analyses of the raw EEG signal showed two effects selective to the epoch containing the UMI: the retrocue and spTMS

each produced phase shifts in the low-beta band. These findings demonstrate that deprioritization involves active neural mechanisms distinct from the processing of the IMI, and that these are supported by low-beta oscillatory dynamics in parietal cortex. We hypothesize that the mechanism underlying spTMS-triggered involuntary retrieval of the UMI is the disruption of the encoding of priority status, which may depend on oscillatory dynamics in the low-beta band.

Significance Statement

This study provides key insights into how the brain maintains information at different levels of priority in working memory. Importantly, it shows how deprioritizing information in working memory is different from simply "dropping" it. By combining transcranial magnetic stimulation with high temporal resolution measurement of neural signals (with EEG), we replicate previous findings that spTMS triggers the involuntary retrieval of previously unprioritized information and provide new insight into how it does this. Specifically, at the level of EEG dynamics, spTMS is shown to uniquely alter oscillatory dynamics in the low-beta band.

Introduction

The flexible control of behavior requires controlling the contents of our working memory (WM) to prioritize the information that is critical for in-the-moment behavior, to maintain information with potential future relevance in a way that does not interfere with current

behavior, and to remove information that is no longer useful. These processes have been operationalized with the double-serial retrocuing (DSR) procedure in which the one of two memory items not cued by a first prioritization cue temporarily acquires the status of 'unprioritized memory item' (UMI). Critically, in DSR, the UMI remains relevant because it might be cued later in the trial (Lewis-Peacock et al., 2012).

Initial neuroimaging studies of the DSR task reported that multivariate pattern analysis (MVPA) evidence for an active representation of the UMI dropped to baseline during the delay following the first retrocue (with fMRI, Lewis-Peacock et al., 2012; Rose et al., 2016; LaRocque et al., 2017; with EEG, LaRocque et al., 2014; Rose et al., 2016). Furthermore, when a single pulse of transcranial magnetic stimulation (spTMS) was applied during this time, decodability of the UMI was recovered (Rose et al., 2016), a pattern consistent with the idea that prioritization may be accomplished, in part, via a transformation of the UMI into an "activity silent" state (c.f., Stokes, 2015). More recently, however, several studies have identified evidence for an active representation of the UMI. Some of these have trained and tested the decoder on the UMI itself (e.g., Christophel et al., 2018; Paluch et al., 2025), and some have trained on the PMI and tested on the UMI (i.e., cross-label decoding). Interestingly, cross-label decoding has yielded evidence that the active representation of the UMI is "opposite" that of the PMI, manifested as below-baseline decoding of the UMI (van Loon et al., 2018), as larger multivariate representational dissimilarity between PMI and UMI within category than between category (van Loon et al., 2018), and as 'flipped' inverted encoding model (IEM) reconstructions of the UMI relative to the PMI (Wan et al., 2020; Yu et al., 2020) consistent with a common representational format for UMI and PMI. (But see

Iamshchinina et al. (2021) for cross-label decoding results that do not suggest "opposite" coding of the UMI.)

Although demonstrations of successful decoding of the UMI from neural signals raise questions about the activity-silent hypothesis (c.f., Barbosa et al., 2021, Wolff et al., 2021; Kandemir et al., 2024), understanding the mechanisms underlying the effects of spTMS on working memory remains important because spTMS influences behavior, not just decoding. For example, during blocks of 1-item recall-of-location trials, spTMS delivered during the intertrial interval (ITI) influences the magnitude of the attractive serial bias exerted by the previous trial's location on the subsequent trial's recall (Barbosa et al., 2020). In the DSR task, spTMS affects behavior in priority-dependent ways consistent with findings using brief 20Hz TMS trains to posterior sensory cortex (Zokaei et al., 2014a; Zokaei et al., 2014b). This is demonstrated with a version of the DSR procedure in which the UMI is presented as the recognition probe for a subset of the probes that require a "nonmatch" response. spTMS increases the false-alarm rate (FAR) to these lures when delivered prior to the first of the two recognition probes, but not when delivered later in the trial, when the item not cued by the second retrocue has the status of "irrelevant memory item" (IMI; because it will not be tested; Rose et al., 2016, Fulvio & Postle 2020). This would be expected if spTMS had the effect of reactivating only items currently held in WM.

What might be the neural bases of the effect of spTMS on the UMI? Initial evidence points at an important role for oscillatory dynamics in the beta band. In one study, filtering the EEG signal into classical frequency bands suggested that the spTMS restoration of the decodability of the UMI is restricted to the beta band (Rose et al.,

2016). In another, the decomposition of the EEG signal into a set of discrete coupled-oscillator components revealed that only components with peaks in the beta band, and that were prominent at posterior scalp locations, were sensitive to spTMS in a manner that related to its behavioral effects (Fulvio et al., 2024). These latter results suggested that spTMS may exert its effects on DSR performance by disrupting a beta band-encoded deprioritization mechanism, thereby reducing priority-based discriminability of the cued item ("prioritized memory item" (PMI)) relative to the UMI due to an increased level of interitem competition. In the present study, we recorded the EEG during performance of DSR and a closely-matched single-retrocue (SR) task, to assess the role of low-frequency dynamics, particularly in the low-beta band (13-20 Hz), in priority-related effects of spTMS on WM performance.

Methods

Code accessibility

The code/software described in the paper is freely available online at [URL will be provided prior to publication].

Participants

Twelve neurologically healthy members of the University of Wisconsin–Madison community (8 females, 18-28 years, M = 28.3 years, all right-handed), with no reported contraindications for magnetic resonance imaging (MRI) or transcranial magnetic stimulus (TMS) took part in all portions of the study. The sample size was chosen to match the sample size of Fulvio & Postle (2020), which itself was designed as a replication and extension of Rose et al. (2016), the extension being important for assessing the specificity of the "reactivating" effect of spTMS on the UMI. The sample

size for Fulvio & Postle (2020) was determined using an *a priori* power analysis based on the behavioral results of Experiment 4 of Rose et al. (2016), the results of which indicated that 12 participants would be needed to achieve 80% power in the critical behavioral comparison of interest. (Note that although the Fulvio & Postle (2020) was successful in confirming that the behavioral effect of spTMS was specific to the UMI, and the EEG data that it produced were the basis of (Fulvio et al., 2024), the inclusion in Fulvio & Postle (2020) of lure probes, in which a subset of nonmatching probes were the uncued item (i.e., the UMI), complicated the interpretation of MVPA decoding of the EEG study. The present study did not include UMI lures.)

All participants had normal or corrected-to-normal vision with contact lenses (eyeglasses were not compatible with TMS targeting apparatus), and all reported having normal color vision. The research was approved by the Health Sciences Institutional Review Board at the University of Wisconsin—Madison. All participants gave written informed consent at the start of each session and received monetary compensation in exchange for participation.

Experimental procedure

The study was carried out in four sessions, each on a separate day. The first session was an MRI scan that yielded the anatomical image used to guide spTMS during the ensuing three behavioral EEG sessions. During each behavioral session, participants performed alternating blocks of DSR and SR, with each session comprising eight 30-trial blocks. Block order was counterbalanced across participants and switched across sessions within participants.

Behavioral tasks

The single retrocue (SR) task began with the simultaneous presentation of two sample items drawn from two of three categories (face, word, direction of dot motion), one above and one below central fixation (2 sec), followed by 5 sec of fixation ("*Delay 1.1*"), followed by a dotted line whose location indicated which of the two samples would be tested ("Cue"), followed by 4.5 sec of fixation ("*Delay 1.2*"), followed by a recognition probe (1 sec). Using a standard keyboard, participants made 'Match' ('1' key) /'non-match' ('2' key) responses within a 2-sec window beginning with probe onset, with feedback displayed for remainder of response window (the fixation cross turned green for correct responses, red for incorrect responses). The ITI varied between 2-4 sec (**Fig. 1A**). 50% of probes were matches of the cued item (e.g., the same face that had been shown during the sample period), and 50% of nonmatches were drawn from the same category as the cued item (e.g., a face that was different from the sample face). Throughout each block, the TMS coil was positioned to target area IPS2 in the right hemisphere, and randomization of memory set, cued category, and probe type were constrained so that spTMS was delivered on 50% of trials of each type. spTMS pulses were delivered 2-3 seconds after the offset of the cue, the unpredictable lag determined by selection (with replacement) from a flat distribution of lags that varied in 50 ms steps.

The double serial retrocue (DSR) task followed the procedure of the SR task except that the trial did not end with the first recognition probe. Instead, the 2-sec recognition window for "Probe 1" was followed by a second cue ("Cue 2"), followed by "*Delay 2*" (4.5 sec) and "Probe 2" (1 sec, plus additional 1 sec response capture

window). Cue 2 appeared in the same location as Cue 1 on 50% of trials ("stay" trials), and spTMS was counterbalanced across all levels of Cue 1, Probe 1, Cue 2 ("stay" or "switch") and Probe 2.

Figure 1. Behavioral task schematics. A. Schematic of the single-retrocue (SR) task. Participants memorized the identity of two sample items shown at separate locations and drawn from three possible categories (faces, direction-of-motion, words (not shown)). Partway through the delay (i.e., at the end of *Delay 1.1*) a spatial retrocue indicated which of the two items would be tested, and during the ensuing *Delay 1.2* a single pulse of TMS (spTMS) was delivered on 50% of trials. At test, participants indicated whether the probe stimulus was a match or non-match of the cued memory item and received feedback. **B.** Schematic of the double-serial retrocue (DSR) task, the first half of which was identical to the SR task, followed by a second spatial retrocue (*Cue 2*) indicating which of the two samples would be tested by *Probe 2*, followed by a final delay during which spTMS was delivered with on 50% of the trials.

Stimuli

The experimental stimuli were presented in MATLAB using the Psychophysics Toolbox (Brainard, 1997; Pelli, 1997; Kleiner et al., 2007) on an LCD with a resolution of 1920 x 1080 and background color set to black. The stimuli were viewed from a 70 cm viewing distance. The face stimuli were non-famous faces with neutral expressions selected from the set used by Rose et al. (2016; see also Rose et al., 2012). Matching face probes were exact matches. Nonmatching face probes were 50-70% morphs of the sample face and another face of the same gender. For word stimuli the critical feature was phonology: all probes were a different word from the sample but matches rhymed with the sample and nonmatches did not rhyme. Motion stimuli contained ~125 white dots moving with 100% coherence at a speed of 3 deg/s for the duration of the stimulus presentation (2 sec), each trial in a direction chosen at random from the full 360-degree range. The direction of nonmatching motion probes differed from the sample within a range of 5-45 degrees.

MRI data acquisition and preprocessing

Whole brain images were acquired with a 3T MRI scanner (Discovery MR750; GE Healthcare) at the Lane Neuroimaging Laboratory at the University of Wisconsin–Madison. High resolution T1-weighted images were acquired for all participants with an FSPGR sequence (8.2 ms repetition time (TR), 3.2 ms echo time (TE), 12° flip angle,

172 axial slices, 256×256 in-plane, 1.0 mm isotropic). The T1-weighted images were

processed using the AFNI software program to align each participant's brain with the

MNI152_T1_1mm template. In AFNI, a mark was inserted in right intraparietal sulcus

(rIPS2; coordinate: -22 70 58) and used as the target for spTMS (details below).

spTMS targeting and delivery

rIPS2 was selected as a target for spTMS because of its role in implementing priority

maps (e.g., Jerde et al., 2012), and in binding content to context in visual working

memory (Gosseries, Yu, et al., 2018; Cai et al., 2019; Cai et al., 2020; Fulvio et al.,

2023; Teng & Postle, 2024). It is important to note that because this study was not

designed to test a hypothesis about a particular role for rIPS2 in the control of priority in

working memory, our design did not require spTMS of a control site. Rather, rIPS2 was

chosen as a target for spTMS for the theoretical reasons stated above, and because, in

our experience, its stimulation was likely to produce a robust perturbation of working

memory representations. The critical experimental controls in this study were the two

epochs—*DSR Delay 2* and *SR Delay 1.2*—during which the spTMS-related effect on

the uncued item was predicted to differ from its effects on the UMI (i.e., during *DSR

Delay 1.2*).

spTMS targeting was achieved with a navigated brain stimulation (NBS) system

that uses infrared-based frameless stereotaxy to coregister the location and position the

participant's head and that of the TMS coil according to the individual's high-resolution

MRI (NexStim eXimia, Helsinki, Finland). spTMS was delivered with an eXimia TMS

Focal BiPulse transcranial magnetic stimulator fit with a figure-of-eight stimulating coil.

NBS allows estimation of the electrical field induced by TMS at the targeted cortex using

a model of the participant's head and information about the coil position the distance

from the coil to the cortical target. spTMS was delivered to the target to achieve an

estimated intensity at the stimulation target of 90–110 V/m (60–75% of stimulator

output, depending on the thickness of the participant's scalp, cortex and depth of the

target). The coil was oriented along the sagittal plane to induce an anterior-posterior

direction of current, with individual adjustments to minimize EEG artifact. Stimulator

intensity, coil position, and coil orientation were held constant for each participant for the

duration of each session. To mask the sound of TMS coil discharge, participants were

fitted with earbuds through which white noise was played during task blocks, with

volume titrated such that the participants could not detect the click produced by coil

discharge. Stimulation parameters were in accordance with published safety guidelines.

EEG recording and preprocessing

EEG was recorded with a 60-channel cap and TMS-compatible amplifier, equipped with

a sample-and-hold circuit that held amplifier output constant from 100 µs before

stimulation to 2 ms after stimulation (NexStim eXimia, Helsinki, Finland). Electrode

impedance was kept below 5 kΩ. The reference electrode was placed superior to the

supraorbital ridge. Eye movements were recorded with two additional electrodes, one

placed near the outer canthus of the right eye, and one underneath the right eye. The

EEG was recorded between 0.1 and 350 Hz at a sampling rate of 1450 Hz with 16-bit

resolution.

Data were processed offline using EEGLAB (Delorme & Makeig 2004) with the

TMS-EEG signal analyzer (TESA) open-source EEGLAB extension (Rogasch et al.,

2017; Mutanen et al., 2020) and Fieldtrip (Oostenveld, et al., 2011) toolboxes in

MATLAB. We followed the TESA TMS-EEG analysis pipeline

(http://nigelrogasch.github.io/TESA/). For delay periods for which no spTMS was

delivered, a dummy spTMS event tag was added at a latency that matched the most

recent spTMS-present trial delay period. Then, electrodes exhibiting excessive noise

were removed and the data were epoched to -14 s to 4.5 s around the *Delay 1.2* spTMS

event tag (for both the DSR and SR tasks) and -4.5 s to 4.5 s around the *Delay 2*

spTMS event tag (for the DSR task only). The data were downsampled to 500 Hz. To

minimize the TMS artifact in the EEG signal, the data were interpolated using a cubic

function from -2 to 30 ms around the TMS pulse. This interpolation was also carried out

on delay periods on which TMS was not delivered. Next, the data were bandpass

filtered between 1 and 100 Hz with a notch filter centered at 60 Hz. Independent

components analysis (ICA) was used to identify and remove components reflecting

residual muscle activity, eye movements, blink-related activity, residual electrode

artifacts, and residual TMS-related artifacts. Electrodes with excessive noise were

interpolated using the spherical spline method. Finally, the data were re-referenced to

the average of all electrodes that were included in the ICA.

Decoding

A multivariate decoding analysis was used to address the primary question. The goal of

the decoding analysis was to classify trials at the within-subject level according to

whether or not an item was present in the memory array (at the category level, i.e., face,

word, motion). Classification was carried out as a function of priority status during DSR

Delay 1.2, during DSR *Delay 2*, and during SR *Delay 1.2*. Note that for timepoints prior

to the cues, the items were coded according to the status that they would acquire. Thus,

for each participant, 12 classifiers were trained to decode the presence or absence of

each of 3 categories (face, word, motion), at each of 2 levels of priority (cued, uncued)

and at each of 2 levels of spTMS (delivered ("present"), not delivered ("absent")).

Classifiers were trained on spectrally transformed data. Spectral transforms of

the preprocessed EEG data were calculated for each participant with the FieldTrip

toolbox. A multi-taper-method convolution was performed on the epoched data using a

fixed, Hanning-tapered window of 500 ms was used for integer frequencies from 2 to 50

Hz. This resulted in time-frequency representations consisting of 500 ms windows

centered on timepoints spaced at 500 ms intervals, and epoched to the *Delay 1.2*

spTMS pulse/dummy-code in both tasks, and to the *Delay 2* spTMS pulse/dummy-code

in the DSR task, with the 12 data 'present' trial combinations transformed separately for

each, and their 12 corresponding 'absent' trial combinations also transformed

separately.

Before submitting the spectrally transformed data to classifier training, the power

values were log10 transformed. The data were then temporally smoothed by averaging

the power of each channel and frequency combination at each time point with the power

at the preceding and following time point such that each time point aggregated 1.5

seconds of data. Because our interest was in the theta-, alpha-, and beta-frequency

bands, we carried out a broadband classification including all three (4-30 Hz), and then

we carried out a separate classification analysis in which the classifier was trained on

individual frequency bands. Broadband classification was carried out at each timepoint.

The feature set consisted of 60 channels x 3 frequencies (average theta power (4-8 Hz),

average alpha power (8-13 Hz), and average beta power (13-30 Hz)). For the individual

frequency band classification, we further divided the beta frequency band into low (13-20 Hz) and high (20-30 Hz), and each of the four frequency-band classifiers trained with a feature set of 60 channels. After frequency band averaging, the data were shuffled along the sample (trial) dimension and submitted to classification with approximately 120 samples provided per model. Classification used the MVPA-Light extension for the Fieldtrip toolbox (https://github.com/treder/MVPA-Light; Treder, 2020), using a logistic regression classifier with L2-regularization and lambda = 1. Five-fold classification was carried out with nested z-scoring per fold (i.e., the training data were z-scored and the resulting mean and standard deviation were used to z-score the test data to avoid leakage between the training and test data) and samples were averaged in random groups of two. In cases of uneven numbers of trials within classes, any remaining trials after averaging were discarded. In cases of imbalance across classes, the logistic model included a bias correction. Finally, the five-fold classification was repeated 10 times and the average area under the curve (AUC) was computed across all folds and repetitions for the particular model (combination of stimulus category, status, and TMS delivery). Here, AUC reflects how well the model can distinguish when a memory item was present versus absent from the EEG signal with higher AUC values signaling that the model is better at detecting when a participant was actively remembering something. These analyses were carried out using the HTCondor system deployed at the University of Wisconsin—Madison Center for High Throughput Computing, https://doi.org/10.21231/gnt1-hw21).

Inferential statistical analyses

Behavior. spTMS-related effects on behavior have previously been shown to occur for trials in which the uncued memory item was used as a lure but not for trials in which the probe was drawn from the cued category (Rose et al., 2016; Fulvio & Postle, 2020). Because the current study did not include lure probes, behavioral performance was not expected to be informative. Nevertheless, we analyzed recognition accuracy using two linear mixed effects models. The first analyzed behavioral performance for the first and second probe responses in the DSR task only with main effects of (1) probe response (two levels: first, second) (2) spTMS (delivered, not delivered); and (3) probe type (matching, non-matching) included in the analysis as fixed effects. Additionally, the interaction between probe response, spTMS, and probe type was included. All three fixed effects variables were coded as categorical variables. As random effects, the model included intercepts for the 12 participants. When we included participant-level random slopes for within-subject factors (e.g., spTMS), the model fits did not improve (e.g., with spTMS slope: ΔAIC = 3.9, p = 0.96), therefore, we did not use these more complex models. The second model included the same fixed effects but was fit to the data for all three probe responses (i.e., DSR task probes 1 & 2 and the single probe of the SR task). For this model, the probe response fixed effect had three levels (first (DSR), second (DSR), SR), and because the interaction was not significant, it was not included in the model. The p-values were obtained by F-tests for each term in the linear mixed-effects model.

Decoding. To analyze multivariate decoding effects of spTMS, we used cluster-based permutation tests on logistic regression performance as expressed as area under the ROC curve (AUC). The median AUC was computed for each participant, collapsed

across the three stimulus categories, for each combination of priority status (PMI; UMI; IMI) and TMS delivery (spTMS-present; spTMS-absent). The data were then averaged at the group level across the 12 participants, and cluster-based permutation testing was used to identify timepoints of significant group-level classification performance against an AUC level = .5 (two-tailed) and, separately, to identify timepoints of significant differences in classification performance as a function of priority status. To address the primary question of this study, we report the results of these analyses within three task epochs during which spTMS could be delivered: DSR *Delay 1.2*, DSR *Delay 2*, and SR *Delay 1.2*. Additional follow-up between-task epoch comparisons were carried out on three timepoints: the spTMS delivery timepoint, and the first and second timepoints immediately following spTMS delivery. We report the results of these comparisons using one-tailed Wilcoxon signed-rank tests testing for greater classification AUC, at these time points, in DSR *Delay 1.2* than the other two task epochs. We also report the rank-biserial correlation coefficient (*rbc*) effect size.

EEG dynamics

To gain insight about possible mechanisms underlying effects of spTMS on MVPA decoding, we also carried out analyses of phase and power dynamics in the EEG signal in relation to prioritization cues and to spTMS.

EEG analyses of phase. All phase analyses relied on the results of a Hilbert transform. Prior to the transform, participants' EEG signals were baseline-corrected with the 1000 ms preceding the trial (i.e., during the ITI) and downsampled to 100 Hz. Individual trial data were then averaged across channels. A Butterworth bandpass filter (order = 4) was applied to 4-second epochs centered on the prioritization cues and on

spTMS pulses to isolate each of the four frequency bands that were used for decoding analyses: theta (4-8 Hz), alpha (8-13 Hz), low beta (13-20 Hz), and high beta (20-30 Hz). Filtering was implemented using Matlab's filtfilt() function, which uses zero-phase forward and backward filtering. The filtered data were then analyzed using the Matlab function hilbert() and the instantaneous phase at each time point was computed from the analytic signal using the angle() function.

The resulting phase data were used to derive three measures of interest: (i). within-participant event-induced changes in inter-trial phase-locking; (ii). between-participant event-induced mean phase angle shifts; and (iii). event-related changes in between-participant phase-locking. For prioritization cue-related impacts, we computed these measures relative to *Cue 1* of the DSR task and the single cue of the SR task. (We omitted data surrounding *Cue 2* of the DSR task from this analysis because a response and feedback period immediately preceded the cue, which would complicate the interpretation of the results.) For spTMS-related impacts on phase, we computed these measures for DSR *Delay 1.2*, DSR *Delay 2*, and SR *Delay 1.2*. We additionally computed these measures relative to the dummy-coded spTMS pulse in spTMS-absent trials. Because we could not know *a priori* when each frequency band might show a phase reset effect, we used a data-driven approach to identify an optimal analysis window for each combination of frequency band and event with a sliding time window selection approach. The window selection procedure utilized the combined set of all data around the two cues for the cue-related analysis and separately the combined set of all data around the three spTMS pulses for the spTMS-related analysis. The selected

window was then used to compute each of the three measures for each condition separately. (Note that the window identified in spTMS-present trials was used in analyses of spTMS-absent trial data).

To identify the optimal post-event window via the sliding window approach, for cue-locked analyses, multiple candidate window lengths were defined, corresponding to integer numbers of oscillatory cycles (2–4 cycles for the theta and alpha bands, 3–6 cycles for the low and high beta bands). This ensured that lower frequencies were analyzed with longer windows and higher frequencies with shorter windows to preserve phase estimation reliability. For each candidate window, the "post-cue" period (0-500 ms relative to cue onset, cue duration = 500 ms) was sampled in 10 ms steps (corresponding to the 100 Hz sampling rate of the EEG data), and a pre-event window of equal length was defined immediately preceding the cue. The inter-trial phase-locking value (*PLV*) was computed for each participant in each post-cue window by averaging phase observations across all trials at each timepoint and then averaging all timepoints within the window. Separately, the corresponding baseline PLV was computed in the pre-cue window using the same procedure. Finally, pre-to-post differences were computed (*post-pre*). To select the optimal post-cue window for subsequent analyses, group-level *t*-statistics were computed for participants' post - pre measures for each candidate window and cycle length. The window exhibiting the maximum absolute *t*-statistic was chosen as the representative post-cue window for that frequency band and used in the subsequent condition-specific analyses as described below. The window selection procedure for spTMS-related effects was identical, with the exception that the

post-spTMS period began at 30 ms after the pulse to avoid the portion of data that was interpolated around the pulse during pre-processing (see above).

Having identified the analysis window for each frequency band and event-type (i.e., cue or spTMS) using the sliding-window, frequency-adaptive procedure, we computed the three measures within those windows for each participant and condition separately. For spTMS effects, event-related changes in the three measures were computed relative to a baseline corresponding to the same epoch from the spTMS-absent trial, such that observed effects reflected the impact of spTMS beyond spontaneous or task-related activity. Because our hypotheses related to between-condition differences, especially effects in low-beta, we primarily focused our inferential statistics on between-condition comparisons. For the within-participant event-induced PLV changes and group level mean phase shifts measures, we evaluated cross-condition differences (e.g., DSR *Cue 1* vs. SR *Cue*) within each frequency band using a paired permutation test with 5,000 iterations. In each iteration, the condition labels for each participant were randomly flipped to generate a null distribution of mean differences across participants. The two-tailed *p*-value was computed as twice the smaller proportion of permutations exceeding the observed difference. To assess inter-subject PLV, reflecting the alignment of phase changes across participants, we computed the vector length of the mean phase shift across participants for each condition. This measure ranges from 0 to 1, where 0 indicates no consistency and 1 indicates perfect alignment of phase shifts across participants. Standard errors were estimated via 1,000 bootstrap resamples, and paired permutation tests (5,000

iterations) were used to test for cross-condition differences by randomly swapping condition labels across participants as was done for the other two measures. Finally, to account for multiple comparisons across frequency bands in these, *p*-values were FDR-corrected for the cue-related analyses and for the spTMS analyses.

Finally, we carried out time-resolved phase consistency analyses in 10 time windows from 50 ms to 500 ms post-cue/spTMS in 50 ms increments. At each time window, we computed each participant's mean phase angle (averaged across trials) and tested whether these participant-level angles clustered significantly using the Rayleigh test, with the resultant vector length (*r*-value) quantifying clustering strength. Significant within-condition *r*-values indicating clustering of participant phases at the timepoints post-event were identified using the Rayleigh test with FDR correction across the 10 time windows. To test for between-condition differences for spTMS-present versus spTMS-absent trials, we used a permutation test with 1000 iterations of randomly swapping each participant's condition labels (spTMS-present vs. spTMS-absent) and recomputing the *r*-value for each time point and condition and computed the difference between the *r*-values to form the null difference permutation distribution. The *p*-value was computed as twice the minimum of the proportions where the permuted difference was greater than or equal to, or less than or equal to, the observed difference (two-tailed test) and permutation *p*-values were FDR-corrected over the 10 time windows.

EEG analysis of power. For each participant, for each channel and trial, we isolated the EEG signal for the 1 sec preceding the event of interest and, separately, for

the 1 sec following the event. Individual trial data were then averaged across channels

prior to analysis. Each channel-averaged trial was windowed using a Hanning taper to

minimize spectral leakage. The power spectrum was computed using a fast Fourier

transform (FFT) for each trial. From each trial's resulting power spectrum, we extracted

the relative power within the four frequency bands of interest as the proportion of the

total spectral power (from 1-45 Hz) that fell within the target frequency band. For each

participant, the mean power was computed across trials within each 1-second time

window and each frequency band. Within-subject permutation tests (n = 5000) were

used to assess whether observed pre vs. post changes were significant. These were

carried out by randomly flipping 'pre' and 'post' labels across trials within each

participant's data set, maintaining equal proportions. Between-condition comparisons, in

the case of comparing trials with spTMS delivery and those without, were also

conducted using permutation tests (n = 5000) whereby all data were pooled and the

label of 'with' and 'without' spTMS was randomly flipped, maintaining the original

proportions.

Results

Recognition performance

The linear mixed effects model limited to DSR failed to find evidence that the delivery of

spTMS had a significant effect on memory recognition accuracy (**Fig. 2**). In this

analysis, only a significant probe x probe-type interaction was observed ($F(1,88) =$

4.046, *p* = .047), reflecting the fact that performance on probe 1 was similar for

matching and non-matching probes ($M_{matching\_response1}$ = .884 (*SD* = .049); $M_{non\text{-}}$

$_{matching\_response1}$ = .893 (*SD* = .042), but performance on probe 2 was better for matching probes than for non-matching probes ($M_{matching\_response2}$ = .871 (*SD* = .045); $M_{non-matching\_response2}$ = .858 (*SD* = .054)). The second linear mixed effects model, which included SR task performance, also failed to find evidence of an effect of spTMS. In this analysis, only a main effect of probe was observed (*F*(2,139) = 5.717, *p* = .004), reflecting the fact that performance on the SR task and on Probe 1 of the DSR task (M_{SR} = .883 (*SD* = .046); $M_{DSR\_response1}$ = .889 (*SD* = .038)) were both better than performance on Probe 2 of the DSR task ($M_{DSR\_response2}$ = .864 (*SD* = .045)).

Figure 2. Recognition performance in the experimental tasks. Accuracy in the DSR and SR tasks as a function of probe type (match / non-match) and spTMS delivery. Error bars correspond to +/– 1 standard error of the mean for the n=12 participants.

Decoding stimulus information from the broadband EEG signal.

Beginning with *Delay 1.2* of the DSR task, prior to spTMS delivery neither item could be decoded above chance. For the UMI, classifier performance became significant at the time of spTMS delivery, and this effect persisted into the *Probe 1* epoch. For the PMI, classifier performance became significant one timepoint later than it had for the UMI, and this effect also persisted into the *Probe 1* epoch (**Fig. 3A**). For *Delay 2* of the DSR task, classification prior to spTMS was not different from chance for PMI or IMI, although decoder performance was significantly greater for PMI than for IMI. For the IMI, classifier performance rose above chance beginning with the second timepoint after spTMS (i.e., 1000 ms) and persisted into Probe 2; and for the PMI, classifier performance rose above chance at the time of spTMS and persisted into *Probe 2* (**Fig. 3B**). For the SR task, classifier performance did not differ from chance for either item during *Delay 1.2*. For the IMI, classifier performance rose above chance beginning with the second timepoint after spTMS and persisted into the Probe epoch; and for the PMI, classifier performance rose above chance one timepoint after spTMS and persisted into the Probe epoch (**Fig. 3C**). For trials in which spTMS was not delivered, UMI classifier performance never rose above chance, from the cue to response period, during either task and PMI classifier performance only rose above chance immediately before (*Delay 1.2*) or concurrent with (*Delay 2*) probe onset in the DSR task, and only briefly rose above chance in response to the retrocue in the SR task (**Fig. S1**)*.* Thus, successful UMI/IMI decoding was observed only with spTMS delivery, and concurrent with spTMS only during *Delay 1.2* of the DSR task. One-tailed Wilcoxon signed-rank tests comparing UMI/IMI decoding at the spTMS timepoint confirmed significantly superior

UMI classification performance for *Delay 1.2* relative to *Delay 2* (*W* = 63, *p* = .032, *rbc* =

.615) and relative to SR *Delay 1.2* (*W* = 64, *p* = .026, *rbc* = .641; **Fig. 3D**). None of the

comparable comparisons for subsequent timepoints (i.e., at a longer lag after spTMS)

reached significance.

Figure 3. Broadband power classifier performance. A. Classifier AUC during *Delay*

1.2 of the DSR task for the PMIUMI. Blue highlighting along the x-axis corresponds to

the retrocue, green highlighting corresponds to the probe+feedback periods, and the

spTMS pulse time-locked to 0 seconds. **B.** Same as **A.** for *Delay 2* of the DSR task. **C.**

Same as **A.** & **B.** for *Delay 1.2* of the SR task. Error bands correspond to +/– 1 standard

error of the mean for the n=12 participants. Red squares correspond to significant AUC

clusters for PMI decoding; blue squares correspond to significant AUC clusters for UMI/IMI decoding; black squares correspond to significant AUC difference clusters between PMI and UMI/IMI. Filled markers: *p* < .05; empty markers: *p* < .1. **D.** Between-epoch comparison of UMI/IMI decoding for the three task epochs at the spTMS timepoint (0 s), the timepoint after spTMS delivery (.5 s), and two timepoints after spTMS delivery (1 s). Blue * markers: *p* < .05 based on cluster-based permutation test results in **A.-C.**; black * markers: *p* < .05 based on one-tailed Wilcoxon signed-rank tests. See **Figure S1** for classifier performance for trials without spTMS delivery.

Decoding stimulus information from individual frequency bands. Next, we decomposed the EEG data into classically defined frequency bands--theta ([4,8 Hz)), alpha ([8,13 Hz)), low beta ([13,20 Hz]), and high beta ([(20,30] Hz)—and repeated the analyses summarized in Figure 3. Inspection of the classification results across task epochs revealed no evidence for spTMS reactivation of the UMI/IMI in the theta band (**Fig. 4A**). The pattern of decoding from the alpha band matched what was seen in analyses of the broadband data: successful decoding of the UMI concurrent with the delivery of spTMS that was specific to *Delay 1.2* of the DSR task, with significant IMI decoding during *Delay 2* of DSR and *Delay 1.2* of SR beginning two time points later (**Fig. 4B**). One-tailed Wilcoxon signed-rank tests comparing UMI/IMI classification found significantly better classification performance for DSR *Delay 1.2* compared to DSR *Delay 2* (*W* = 73, *p* = .002, *rbc* = .872) at the spTMS timepoint, but no difference in classification performance when compared to SR *Delay 1.2* (*W* = 53, *p* = .151, *rbc* =

.359; **Fig. 5A**). The overall pattern of decoding of the uncued item from the low beta

band was similar the results from the alpha band, but with the emergence of significant

decoding shifted one timepoint later for all trial types (**Fig. 4C**). However, one-tailed

Wilcoxon signed-rank tests comparing UMI/IMI classification performance for DSR

Delay 1.2 with that of DSR *Delay 2* and SR *Delay 1.2* at each timepoint identified no

significant differences (**Fig. 5B**). Finally, decoding of the UMI/IMI from the high beta

band was unsuccessful throughout DSR *Delay 1.2*, emerged late during DSR *Delay 2*,

and emerged one timepoint after spTMS delivery in SR *Delay 1.2* (**Fig. 4D).** Despite the

lack of statistically significant group-level decoding at the spTMS timepoint in the high

beta band, one-tailed Wilcoxon signed-rank tests comparing UMI/IMI classification

performance found significantly greater UMI classification performance for DSR *Delay*

1.2 compared to IMI classification performance SR *Delay 1.2* at the spTMS timepoint

(W = 65, p = .021, rbc = .667; **Fig. 5A**). Comparisons of the DSR *Delay 1.2*

classification performance with the other epochs at the two timepoints post-spTMS did

not identify any differences. However, a post-hoc comparison found significantly greater

IMI classification performance for DSR *Delay 2* compared to SR *Delay 1.2* (W = 61, p =

.046, rbc = .564). As with the broadband analysis, no successful frequency band-

specific decoding of the UMI or IMI was obtained on trials without spTMS delivery

(**Figure S2**).

Figure 4. Frequency-band specific classifier performance. A. Theta-band Classifier AUC during the DSR *Delay 1.2* epoch (left), DSR *Delay 2* (middle), and SR *Delay 1.2* (right) for the prioritized memory category (PMI) in red and the unprioritized/irrelevant memory category (UMI/IMI) in blue. Gray highlighting along the x-axis corresponds to the retrocue period, green highlighting corresponds to the probe+feedback period, with the delay period occurring between and the spTMS pulse time-locked to 0 seconds. **B.** Same as **A.** for alpha-band classifier AUC. **C.** Same as **A.** & **B.** for low beta-band classifier AUC. **D.** Same as **A.**, **B.**, & **C.** for high beta-band classifier AUC. Error bands correspond to +/– 1 standard error of the mean for the n=12 participants. Red significance squares correspond to significant AUC clusters for PMI decoding; blue squares correspond to significant AUC clusters for UMI/IMI decoding; black squares correspond to significant AUC PMI-UMI/IMI difference clusters. Filled markers: $p < .05$; empty markers: $p < .1$. See **Figure S2** for classifier performance for trials without spTMS delivery.

Figure 5. UMI/IMI classification performance comparisons. A. Between-epoch comparison of alpha-band UMI/IMI decoding for the three task epochs at the spTMS timepoint (0 s), the timepoint after spTMS delivery (.5 s), and two timepoints after spTMS delivery (1 s). **B.** Same as **A.** for low beta-band UMI/IMI decoding. **C.** Same as **A.** and **B.** for high beta-band UMI/IMI decoding. Blue * markers: *p* < .05 based on cluster-based permutation test results in **A.-C.**; black * markers: *p* < .05; black ** markers: *p* < .01 based on one-tailed Wilcoxon signed-rank tests.

Interim Discussion

Whereas there was no evidence for above-chance decoding of either the PMI or the UMI during the early portion of any delay period, patterns of decoding in the second half of delay periods varied with priority status (i.e., UMI vs. IMI vs. PMI). For the uncued item, successful decoding of the UMI returned concurrent with the delivery of spTMS

(i.e., during DSR *Delay 1.2*), whereas for the IMI successful decoding lagged by at least

two timepoints during (i.e., during DSR *Delay 2* and SR *Delay 1.2*). When the data were

decomposed into classical frequency bands, this priority-specific effect was seen in the

alpha band and, with a one-time-window lag, in the low beta band. For the cued item

(the PMI) the return of above-chance decoding during *Delay 1.2* of the DSR task lagged

the UMI by one time window in the analyses of broadband and alpha band-filtered data.

Although it is possible that these results reflect specific "reactivation" effects of spTMS

on the UMI, this interpretation is complicated by two factors. First, pre-spTMS decoding

of the PMI was at chance, and generally not different from the UMI, thus failing to

demonstrate that the two items were, indeed, held at different levels of priority. Second,

the fact that decodability of both items tended to ramp up late in the delay period in

many conditions might be consistent with a nonspecific strengthening of stimulus

information in anticipation of the recognition probe. For these reasons, it was important

to assess whether the retrocues and spTMS had UMI-specific effects on aspects of the

EEG data that may explain the patterns of MVPA decoding—specifically phase and

power dynamics of the EEG signal. Evidence that these measures also responded

differentially for the UMI would support the interpretability of the differential decoding

effects and might also provide insights into the neurophysiological bases of the "spTMS

reactivation" effect.

EEG dynamics

Phase dynamics

Prioritization cues. The cue presented during DSR *Delay 1* (which designated the UMI) was associated with a significant increase in within-participant phase alignment in the low-beta band when compared with the effects of the cue from the SR task (which designated the IMI; p = .044, FDR-corrected; **Fig. 6A**). Accompanying this effect, group-level mean phase shifts differed significantly between *DSR Delay 1* and *SR Delay 1* in the theta, alpha, and low-beta bands (all p-values < .001, FDR-corrected; **Fig. 6B**): in theta, the difference was the result of a larger shift in response to the cue during SR *Delay 1*; in alpha and low-beta, the differences were the result of phase shifts in opposing directions in DSR *Delay 1* vs. SR *Delay 1*. No other significant differences in cue-related effects across tasks and frequency bands were observed for within-participant phase alignment, mean phase shift, or inter-subject PLV (all p-values ≥ .1853, FDR-corrected). Furthermore, time-resolved analyses showed no evidence of between-participant phase clustering at any post-cue timepoint (**Fig. S3**).

spTMS. In general, spTMS-related effects on within-participant phase locking were markedly larger in magnitude in comparison to cue-related effects. The only difference in these effects across conditions was that in the low-beta band. spTMS triggered larger within-participant phase alignment during DSR *Delay 1.2* than during DSR *Delay 2* (p = .026, FDR-corrected; for all other contrasts p-values ≥ .059, FDR-corrected; **Fig. 6C**).

For group-level spTMS-evoked shifts in mean phase, significant differences between conditions were found in the low-beta and high-beta bands (**Fig. 6D**). In low-beta, the effect was significantly larger in the SR Delay 1.2 than in DSR *Delay 1.2* and

DSR *Delay 2,* whereas in high-beta, it was significantly larger during DSR *Delay 1.2*

than during DSR *Delay 2* and SR *Delay 1.2.*

Inter-subject PLV changes were not significantly different across any conditions

within any frequency bands (all *p*-values ≥ .4572, FDR-corrected; results not depicted).

However, time-resolved analyses revealed post-spTMS-specific between-participant

phase clustering in all conditions that was absent in spTMS-absent trials. Clustering

emerged in all four frequency bands following spTMS, but with distinct temporal profiles:

the clustering was persistent throughout the analysis time range (i.e., 500 ms post

spTMS) in theta and alpha, but transient in low-beta and high-beta. (**Fig. S4**).

To summarize, at the level of individual phase dynamics, the effect of the cue

and of spTMS were specific to the low-beta band. At the group level, the cue-related

phase shift for DSR *Cue 1* vs. SR *Cue* was opposite in sign in the low-beta beta band in

comparison to the theta and alpha bands, and a complex pattern of spTMS-related

effects was observed in the low- and high-beta bands.

Figure 6. Analysis of phase dynamics. A. Cue-evoked changes in within-participant phase alignment (post – pre PLV) for each of four frequency bands in response to *Cue 1* during DSR *Delay 1* (indigo) and the cue during SR *Delay 1* (green). **B.** Cue-evoked changes in group-level phase, for each of four frequency bands, in response to *Cue 1* during DSR *Delay 1* (indigo) and the cue during SR *Delay 1* (green). **C.** Same as **A.,** but for spTMS-evoked changes in within-participant phase alignment, normalized by the effects from spTMS-absent trials, for DSR *Delay 1.2* (indigo), DSR *Delay 2* (transparent indigo), and SR *Delay 1.2* (green). **D.** Same as **B.,** but for spTMS-evoked changes

(normalized) in DSR *Delay 1.2* (indigo), DSR *Delay 2* (transparent indigo), and SR

Delay 1.2 (green). * markers: $p < .05$, FDR-corrected; error bars correspond to +/- 1

SEM.

Power dynamics

It was uniformly the case that power declined over time, whether assessed in response

to the cues (**Fig. 7A**), or in the post-cue delay periods on spTMS-present and spTMS-

absent trials (**Fig. 7B**). The magnitude of these declines generally decreased with

increasing frequency. These dynamics are characteristic of active maintenance during

working memory delays and indicate that the phase-related effects reported above were

not confounded by systematic changes in oscillatory power.

Figure 7. Within-subject power. A. Cue-evoked changes in relative power (post – pre) for each of the four frequency bands in response to *Cue 1* (DSR; left) and the single cue in the SR task (right). **B.** Same as **A.** for spTMS-evoked changes of relative power, with comparison to baseline in trials when spTMS was not delivered. *** markers: p < .001; ** markers: p < .01; * markers: p < .05 based on permutation tests; error bars correspond to +/- 1 *SEM.*

Discussion

The results presented here demonstrate patterns of spTMS-related decoding and of EEG dynamics that provide insight into the representation of priority in working memory.

For the uncued item, only during DSR *Delay 1.2*, when it had the status of UMI, was the recovery of decoding concurrent with spTMS, and thus plausibly linked to the instantaneous effects of the perturbation. In the other two delay periods, recovery of decoding of the IMI occurred several timesteps later, whereas for the cued item (the PMI) the latency of spTMS-related decoding was comparable across conditions. Frequency-band specific analyses revealed the alpha and low-beta bands to support decoding of all items in all instances where broadband decoding had been observed. Analyses of phase dynamics in the EEG indicated that only during DSR *Delay 1.2* did the prioritization cue and spTMS produce phase shifts in the low-beta band at the within-participant level.

These results join a large body of evidence indicating that the deprioritization of information is accomplished via a representational transformation that distinguishes it from prioritized information, as well as from information that is no longer needed in working memory (e.g., Lewis-Peacock et al., 2012, van Loon, 2018, Yu, Teng, & Postle, 2020; Wan, 2020; Fulvio & Postle, 2020; Wan et al. 2024). They extend this literature by bringing into focus the idea that beta-band dynamics may play an important role in the encoding of priority. The first demonstration of spTMS-triggered decoding of the UMI isolated this effect to the beta band (Rose et al., 2016). More recently, it has been shown that when the EEG signal on the scalp is decomposed into a set of discrete coupled oscillators, it is components in the beta band whose sensitivity to spTMS correlates with spTMS-related declines in behavior. Here, we provide evidence for a specific role for phase dynamics in the low-beta band at the within-participant level, in

that only the cue designating the UMI is associated with a phase shift in low-beta, and the effect of spTMS that is unique to DSR *Delay 1.2* is a phase shift in low-beta.

The delivery of single pulses of transcranial magnetic stimulation (spTMS) to posterior parietal cortex while the EEG is simultaneously recorded has revealed the "natural frequency" of this region to be in the low-beta range (Rosanova et al., 2009, Kundu et al., 2014), in comparison to alpha when spTMS targets occipital cortex, and high-beta/low-gamma when spTMS targets premotor cortex (Rosanova et al., 2009) and PFC (Ferrarelli et al., 2012). Furthermore, whereas spTMS-evoked phosphene perception is predicted by pre-TMS power and phase in spontaneous alpha when spTMS targets occipital cortex, it depends on pre-TMS power in spontaneous low-beta when PPC is targeted (Samaha et al., 2017). In the present study, and in Fulvio et al. (2024), spTMS was delivered to IPS2. Building from this, our tentative proposal is that priority in working memory may be encoded in a frontoparietal priority map that harnesses endogenous oscillations in the low-beta band to maintain the uncued item in an unprioritized state (c.f., Kundu et al., 2015). From this perspective, the fact that spTMS produces involuntary retrieval of the UMI at the behavioral level (Rose et al., 2016; Fulvio & Postle, 2020) suggests that the spTMS-triggered changes in low-beta band dynamics that are specific to the UMI may reflect a disruption of its low beta-encoded priority status. Importantly, our current construal of this mechanism is that it would be different from the proposed role for beta bursts in the implementation of top-down inhibitory control (e.g., Wessel & Anderson, 2024; Lundqvist et al., 2024), for two reasons. First, decoding of the UMI from the beta band is sustained for periods of time much longer than what would be expected from a phasic signal carried by a burst.

Second, the delivery of spTMS in these studies is unpredictable, and so unlikely to be tightly synchronized with an endogenously formulated, and brief, control signal.

References

Barbosa, J., Stein, H., Martinez, R.L., Galan-Gadea, A., Li, S., Dalmau, J., Adam, K.C.S, Valls-Solé, J., Constantinidis, C., & Compte, A. (2020). Interplay between persistent activity and activity-silent dynamics in the prefrontal cortex underlies serial biases in working memory. *Nature neuroscience, 23*(8), 1016-1024. https://doi.org/10.1038/s41593-020-0644-4

Barbosa, J., Lozano-Soldevilla, D., & Compte, A. (2021). Pinging the brain with visual impulses reveals electrically active, not activity-silent, working memories. *PLoS biology*, *19*(10), e3001436. https://doi.org/10.1371/journal.pbio.3001436

Berens, P. (2009). CircStat: A Matlab Toolbox for Circular Statistics, *Journal of Statistical Software*, *31*(10). http://www.jstatsoft.org/v31/i10

Brainard, D.H. (1997). The psychophysics toolbox. *Spatial vision*, *10*(4), 433-436.

Cai, Y., Sheldon, A.D., Yu, Q., & Postle, B.R. (2019). Overlapping and distinct contributions of stimulus location and of spatial context to nonspatial visual short-term

memory. *Journal of neurophysiology, 121*(4), 1222-1231.

https://doi.org/10.1152/jn.00062.2019

Cai, Y., Fulvio, J.M., Yu, Q., Sheldon, A.D., & Postle, B.R. (2020). The role of location-context binding in nonspatial visual working memory. *Eneuro*, *7*(6) ENEURO.0430-20.2020. https://doi.org/10.1523/ENEURO.0430-20.2020

Christophel, T.B., Klink, P.C., Spitzer, B., Roelfsema, P.R., & Haynes, J.D. (2017). The distributed nature of working memory. *Trends in cognitive sciences, 21*(2), 111-124. https://doi.org/10.1016/j.tics.2016.12.007

Christophel, T.B., Iamshchinina, P., Yan, C., Allefeld, C., & Haynes, J.D. (2018). Cortical specialization for attended versus unattended working memory. *Nature neuroscience*, *21*(4), 494-496. https://doi.org/10.1038/s41593-018-0094-4

Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of neuroscience methods*, *134*(1), 9-21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Ferrarelli, F., Sarasso, S., Guller, Y., Riedner, B., Peterson, M., Bellesi, M., Massimini, M., Postle, B.R., & Tononi, G. (2012). Reduced oscillatory frequency of thalamo-cortical circuits in schizophrenia. *Archives of General Psychiatry, 69*(8):766-74. https://doi.org/10.1001/archgenpsychiatry.2012.147

Fulvio, J.M., & Postle, B.R. (2020). Cognitive control, not time, determines the status of items in working memory. *Journal of Cognition*, *3*(1), 8. https://doi.org/10.5334/joc.98

Fulvio, J.M., Yu, Q., & Postle, B.R. (2023). Strategic control of location and ordinal context in visual working memory. *Cerebral Cortex, 33*(13), 8821-8834. https://doi.org/10.1093/cercor/bhad164

Fulvio, J.M., Haegens, S., & Postle, B.R. (2024). Single-pulse transcranial magnetic stimulation affects working-memory performance via posterior Beta-band oscillations. *Journal of Cognitive Neuroscience, 36*(9), 1827-1846. https://doi.org/10.1162/jocn_a_02194

Gosseries, O., Yu, Q., LaRocque, J.J., Starrett, M.J., Rose, N.S., Cowan, N., & Postle, B.R. (2018). Parietal-occipital interactions underlying control-and representation-related processes in working memory for nonspatial visual features. *Journal of Neuroscience*, *38*(18), 4357-4366. https://doi.org/10.1523/JNEUROSCI.2747-17.2018

Iamshchinina, P., Christophel, T.B., Gayet, S., & Rademaker, R L. (2021). Essential considerations for exploring visual working memory storage in the human brain. *Visual Cognition, 29*(7), 425-436. https://doi.org/10.1080/13506285.2021.1915902

Kandemir, G., Wilhelm, S.A., Axmacher, N., & Akyürek, E.G. (2024). Maintenance of color memoranda in activity-quiescent working memory states: Evidence from impulse perturbation. *Iscience*, *27*(4). https://doi.org/10.1016/j.isci.2024.109565

Kleiner, M., Brainard, D., & Pelli, D. (2007). What's new in Psychtoolbox-3?.

Kundu, B., Johnson, J.S., & Postle, B.R. (2014). Pre-stimulation phase predicts the TMS-evoked response. *Journal of Neurophysiology, 112*(8), 1885-1893. https://doi.org/10.1152/jn.00390.2013

Kundu, B., Chang, J.Y., Postle, B.R., & Van Veen, B.D. (2015). Context-specific differences in fronto-parieto-occipital effective connectivity during short-term memory maintenance. *Neuroimage, 114*(1), 320-327. https://doi.org/10.1016/j.neuroimage.2015.04.001

LaRocque, J.J., Lewis-Peacock, J.A., & Postle, B.R. (2014). Multiple neural states of representation in short-term memory? It's a matter of attention. *Frontiers in human neuroscience, 8,* 5. https://doi.org/10.3389/fnhum.2014.00005

LaRocque, J.J., Riggall, A.C., Emrich, S.M., & Postle, B.R. (2017). Within-category decoding of information in different attentional states in short-term memory. *Cerebral Cortex*, *27*(10), 4881-4890. https://doi.org/10.1093/cercor/bhw283

Lewis-Peacock, J.A., Drysdale, A.T., Oberauer, K., & Postle, B.R. (2012). Neural evidence for a distinction between short-term memory and the focus of attention. *Journal of cognitive neuroscience, 24*(1), 61-79. https://doi.org/10.1162/jocn_a_00140

Lundqvist, M., Miller, E.K., Nordmark, J., Liljefors, J., & Herman, P. (2024). Beta: bursts of cognition. *Trends in Cognitive Sciences, 28*(7), 662-676. https://doi.org/10.1016/j.tics.2024.03.010

Mutanen, T.P., Biabani, M., Sarvas, J., Ilmoniemi, R.J., & Rogasch, N.C. (2020). Source-based artifact-rejection techniques available in TESA, an open-source TMS–EEG toolbox. *Brain Stimulation: Basic, Translational, and Clinical Research in Neuromodulation, 13*(5), 1349-1351. https://doi.org/10.1016/j.brs.2020.06.079

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.M. (2011). FieldTrip: open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational intelligence and neuroscience, 2011*(1), 156869. https://doi.org/10.1155/2011/156869

Paluch, K., Magnuski, M., Średniawa, W., Ivanovski, D., Rysz, A., Służewska-Niedźwiedź, M., Paterski, T., Fortuna, W., Smarzewska, K., Reinacher, P.C., Kaczor, Sz., Tabakow, P., Babu, H., & Kamiński, J. (2025). Unattended working memory items are coded by persistent activity in human medial temporal lobe neurons. *Nature Human Behaviour,* 1-15. https://doi.org/10.1038/s41562-025-02235-0

Panichello, M.F., & Buschman, T.J. (2021). Shared mechanisms underlie the control of working memory and attention. *Nature, 592*(7855), 601-605.

https://doi.org/10.1038/s41586-021-03390-w

Pelli, D.G. (1997). The VideoToolbox software for visual psychophysics: transforming numbers into movies. *Spatial vision, 10*(4), 437-442.

https://doi.org/10.1163/156856897x00366

Rogasch, N.C., Sullivan, C., Thomson, R.H., Rose, N.S., Bailey, N.W., Fitzgerald, P.B., Farzan, F., & Hernandez-Pavon, J.C. (2017). Analysing concurrent transcranial magnetic stimulation and electroencephalographic data: A review and introduction to the open-source TESA software. *Neuroimage, 147*, 934-951.

https://doi.org/10.1016/j.neuroimage.2016.10.031

Rosanova, M., Casali, A., Bellina, V., Resta, F., Mariotti, M., & Massimini, M. (2009). Natural frequencies of human corticothalamic circuits. *Journal of Neuroscience, 29*(24), 7679-7685. https://doi.org/10.1523/JNEUROSCI.0445-09.2009

Rose, N.S., Olsen, R.K., Craik, F.I., & Rosenbaum, R.S. (2012). Working memory and amnesia: the role of stimulus novelty. *Neuropsychologia, 50(*1), 11-18.

https://doi.org/10.1016/j.neuropsychologia.2011.10.016

Rose, N.S., LaRocque, J.J., Riggall, A.C., Gosseries, O., Starrett, M.J., Meyering, E.E., & Postle, B.R. (2016). Reactivation of latent working memories with transcranial magnetic stimulation. *Science, 354*(6316), 1136-1139.

https://doi.org/10.1126/science.aah7011

Samaha, J., Gosseries, O., & Postle, B.R. (2017). Distinct oscillatory frequencies underlie excitability of human occipital and parietal cortex. *Journal of Neuroscience*, *37*(11), 2824-2833. https://doi.org/10.1523/JNEUROSCI.3413-16.2017

Stokes, M.G. (2015). 'Activity-silent'working memory in prefrontal cortex: a dynamic coding framework. *Trends in cognitive sciences, 19*(7), 394-405.

https://doi.org/10.1016/j.tics.2015.05.004

Teng, C., & Postle, B.R. (2024). Investigating the roles of the visual and parietal cortex in representing content versus context in visual working memory. *Eneuro*, *11*(2) ENEURO.0270-20.2024. https://doi.org/10.1523/ENEURO.0270-20.2024

Treder, M.S. (2020). MVPA-Light: A Classification and Regression Toolbox for Multi-Dimensional Data. *Frontiers in Neuroscience, 14,* 289.

https://doi.org/10.3389/FNINS.2020.00289

van Loon, A.M., Olmos-Solis, K., Fahrenfort, J.J., & Olivers, C.N. (2018). Current and future goals are represented in opposite patterns in object-selective cortex. *elife, 7*, e38677. https://doi.org/10.7554/eLife.38677

Wan, Q., Cai, Y., Samaha, J., & Postle, B.R. (2020). Tracking stimulus representation across a 2-back visual working memory task. *Royal Society open science, 7*(8), 190228. https://doi.org/10.1098/rsos.190228

Wan, Q., Menendez, J.A., & Postle, B.R. (2022). Priority-based transformations of stimulus representation in visual working memory. *PLoS Computational Biology, 18(*6), e1009062. https://doi.org/10.1371/journal.pcbi.1009062

Wan, Q., Ardalan, A., Fulvio, J.M., & Postle, B.R. (2024). Representing context and priority in working memory. *Journal of cognitive neuroscience, 36*(7), 1374-1394. https://doi.org/10.1162/jocn_a_02166

Wessel, J.R., & Anderson, M.C. (2024). Neural mechanisms for domain-general inhibitory control. *Trends in Cognitive Sciences 28*(2), 124-143. https://doi.org/10.1016/j.tics.2023.09.008

Wolff, M.J., Akyurek, E.G., & Stokes, M.G. (2021). What is the functional role of delay-related alpha oscillations during working memory?. *PsyArXiv* https://doi.org/10.31234/osf.io/z6y5b

Yu, Q., Teng, C., & Postle, B.R. (2020). Different states of priority recruit different neural

representations in visual working memory. *PLoS biology, 18*(6), e3000769.

https://doi.org/10.1371/journal.pbio.3000769

Zokaei, N., Manohar, S., Husain, M., & Feredoes, E. (2014). Causal evidence for a

privileged working memory state in early visual cortex. *Journal of Neuroscience, 34*(1),

158-162. https://doi.org/10.1523/JNEUROSCI.2899-13.2014

Zokaei, N., Ning, S., Manohar, S., Feredoes, E., & Husain, M. (2014). Flexibility of

representational states in working memory. *Frontiers in Human Neuroscience, 8,*

853. https://doi.org/10.3389/fnhum.2014.00853