SpaceSequest: A unified pipeline for spatial transcriptomics data analysis

Yu H. Sun[1,#,*], Sarbottam Piya[1,#], Zhengyu Ouyang[2,#], Yirui Chen[1], Jake Gagnon[3], Shaolong

Cao[1], Haotian Zhang[1], Baobao Song[1], Jing Zhu[1], Khyati Chandratre[1], Houlin Yu[1], Wenxing Hu[1],

Matthew Ryals[4], Fergal Casey[1], Dann Huh[1], Baohong Zhang[1]

[1] Research, Biogen Inc., Cambridge, MA, 02142, USA

[2] Data Science, BioInfoRx Inc., Madison, WI, 53719, USA

[3] Biostatistics, Biogen Inc., Cambridge, MA, 02142, USA

[4] PharmaLex, Conshohocken, PA, 19428, USA

Contribute equally

* Correspondence: Yu H. Sun, yuhenry.sun@biogen.com

ABSTRACT (limit of 250 words)

Background

Spatial transcriptomics has emerged as one of the most powerful tools for gaining biological insights, enabling researchers to uncover intricate relationships between gene expression patterns and tissue architecture. Recent advances in the field have resulted in a variety of new platforms, including Visium, Visium HD, and Xenium from 10x Genomics, as well as GeoMx and CosMx from NanoString Technologies, which has now been acquired by the Bruker Corporation. However, the existence of diverse spatial transcriptomics platforms and various data formats poses challenges in standardizing data analysis. Thus, there remains a critical gap in the availability of a comprehensive pipeline capable of conducting end-to-end analysis that is necessary to extract biological insights from multiple spatial transcriptomics platforms.

Results

Here, we present SpaceSequest, a tailored pipeline that utilizes cutting-edge computational methodologies to conduct a thorough analysis, enabling the extraction of crucial biological insights from five major spatial transcriptomics technologies. SpaceSequest performs (1) standardized quality control and general data processing, (2) key analyses customized for each spatial platform, (3) automated cell type annotation and deconvolution, and (4) high-quality figure and analysis result generation. In addition, SpaceSequest allows for smooth integration with cellxgene VIP and Quickomics for user-friendly data access and interactive visualization.

Conclusions:

SpaceSequest is a unified and comprehensive pipeline designed for the analysis, visualization, and publication of spatial transcriptomics data from various platforms. The source code is available at https://github.com/interactivereport/SpaceSequest. To facilitate seamless installation and usage, we have also created a detailed Bookdown tutorial that can be accessed through https://interactivereport.github.io/SpaceSequest/tutorial/docs/index.html.

Keywords:

Spatial transcriptomics, Visium, Visium HD, Xenium, GeoMx, CosMx, single-cell RNA-seq, deconvolution

BACKGROUND

The advent of single-cell RNA sequencing (scRNA-seq) has revolutionized our understanding of cellular heterogeneity and gene expression dynamics. It represents a significant advancement from bulk RNA-seq, which only provides average expression profiles from mixed cell populations, thereby obscuring the underlying cellular diversity [1–3]. Although scRNA-seq generates invaluable insights into cellular heterogeneity, it lacks spatial information, as it dissociates cells from their original tissue context [4, 5]. Understanding the intricate spatial organization of gene expression within complex tissues and organs is crucial for unraveling the mysteries of biology [6–8]. Spatial transcriptomics (ST) has emerged as a transformative technology that quantifies RNA within intact tissue sections, enabling researchers to explore gene expression patterns within their native spatial context [6, 9–12].

ST can provide precise insights into disease progression by discerning heterogeneity within diseased tissue, which is powerful for studying complex diseases such as tumor and neurological disorders[13–17]. For example, using ST data, Berglund et al. not only distinguished healthy and diseased areas but were able to identify re-stratified tumor heterogeneity by analyzing gene expression gradients within the tumor microenvironment, specifically in the stroma adjacent to the tumor area [18]. These advancements have unlocked new avenues for addressing critical biological inquiries. First, ST plays a pivotal role in characterizing the cell-type composition of tissues. While definitions are often derived from large-scale single-cell RNA-seq or epigenetics datasets and projected onto ST data, the technology also empowers the direct generation of cell-type definitions from the spatial transcriptomics data itself[19, 20]. This application has been widely utilized, resulting in the creation of compositional atlases for a diverse range of tissues, including the nervous system [21–24], human kidney [25–27], heart [28, 29], testes [30], and lung [31, 32]. Second, ST facilitates the exploration of cellular interactions, revealing the rules and patterns governing spatial covariation among distinct cell types. For instance, studies examining the mouse visual cortex identified a preferential spatial proximity of inhibitory neuron

subtypes, shedding light on their organizational principles[33]. Similar proximity analyses have

been instrumental in elucidating gene expression patterns linked to Alzheimer's disease and in

characterizing histopathological responses to traumatic brain injury[34–37]. Finally, ST's

transcriptome-wide data on in situ gene expression empowers the elucidation of molecular

interactions between tissue components[7]. Through the identification of ligand-receptor pairs

among cell types at varying spatial proximities, we can discern the intricate communication

networks that underlie phenomena such as tumor interactions with the microenvironment,

immune infiltrates in tissues, and the establishment of developmental gradients[38, 39].

There are two distinct ST methods: (i) sequencing-based ST, and (ii) imaging-based ST. In the

sequencing-based ST method, a DNA-barcoded surface is used to capture RNA molecules from

the intact tissue section. These captured RNA molecules are then subjected to next-generation

sequencing to determine their respective gene identities[6]. By contrast, in imaging-based

methods, multiple mRNA transcripts are simultaneously detected within intact tissue using

fluorescence in situ hybridization (FISH) or direct in-situ sequencing[6, 12]. The imaging-based

methods, such as 10x Genomics Xenium and NanoString CosMx, are suitable for hypothesis

testing as these methods allow transcriptomic profiling at single cell resolution albeit for a

limited number of target genes. Conversely, sequencing-based methods which produce an

unbiased, whole-transcriptome profile, are suitable for hypothesis generation. Typical methods

include 10x Genomics Visium, Visium HD, and NanoString GeoMx with sequencing readout.

However, transcriptomics profiles obtained from these methods are not directly resolved at

single cell resolution.

One critical drawback of current grid-based ST profiling methods (Visium, Slide-seq [38, 40],

and HDST [41] is their inability to achieve single-cell resolution. These techniques either

combine multiple cells or fractions of multiple cells at each tissue location, limiting the ability to

discern individual cell contributions. Nevertheless, this limitation can be addressed through

computational integration, leveraging the reference transcriptome signatures of cell types

obtained from coupled single-cell RNA sequencing (scRNA-seq) profiles. By integrating ST data

with scRNA-seq data, researchers can deconvolute cell type composition and retrieve further

information from the spatially resolved gene expression patterns[42, 43].

The landscape of spatial transcriptomics data analysis has seen the development of several data analysis pipelines, including ST Pipeline [44] and Spacemake [45]. ST Pipeline primarily focuses on the essential tools and scripts required to process raw files derived from Visium data in FASTQ format, providing datasets primed for downstream analysis [44]. In contrast, Spacemake represents a more comprehensive solution, offering a wider range of functionalities compared to ST Pipeline. This tool is adept at handling spatial transcriptomic data from diverse platforms, such as Visium, Slide-seq, Seq-scope, etc [45]. Another distinctive feature of Spacemake is its utilization of NovoSpaRc [46] that allows mapping snRNASeq data to spatial coordinates [45]. Recently developed pipelines are focused on specific tasks, for example, PIPEFISH[47] was designed for FISH spatial transcriptomics, ENACT was specifically designed for Visium HD data analysis[48], while Panpipes[49] was created for spatial multi-omics analysis. Although these existing pipelines are useful, their capabilities fall short in extracting comprehensive and meaningful biological insights from diverse spatial transcriptomics datasets.

Here we introduce SpaceSequest, our comprehensive spatial transcriptomics data analysis pipeline tailored for the analysis of several major platforms, spanning 10X Genomics Visium, Visium HD, Xenium, as well as NanoString GeoMx and CosMx platforms. By utilizing cutting-edge computational methodologies, SpaceSequest streamlines a series of essential analyses to extract meaningful biological insights. Compatible with diverse input formats, the pipeline seamlessly navigates through critical steps including clustering, deconvolution, cell type annotation, and differential expression analysis. This integrated approach equips researchers with a powerful toolkit to dissect the spatially defined molecular landscape, facilitating a deeper understanding of complex biological systems and their microenvironmental interactions.

IMPLEMENTATION

General data processing

SpaceSequest supports various input formats and processes the data through a series of automated steps, leveraged by multiple well-designed packages including Seurat 5 [50] and anndata[51]. For sequencing-based strategies (10x Genomics Visium, Visium HD, and NanoString GeoMx with sequencing readouts), FASTQ processing is required before running

SpaceSequest. The input files for running Visium and Visium HD workflows, including

fundamental quality metrics and the unique molecular identifiers (UMI) count matrix, can be

generated through 10x Genomics Space Ranger software. Since our pipeline does not rely on

gene annotation or species-specific details, users are required to accurately select the appropriate

species during Space Ranger execution. For GeoMx, the GeoMx NGS Pipeline developed by

NanoString is required to convert FASTQ to digital count conversion (DCC) files, which can be

further processed by SpaceSequest. In the case of imaging-based technologies such as Xenium

and CosMx, files generated by the platform can be readily used for SpaceSequest data

processing.

Clustering

Clustering is a powerful tool for understanding the structure of the data, as well as the spatial

tissue domains. For general clustering of single cell expression in CosMx and Xenium, or bins in

Visium HD, we utilized the UMAP dimension reduction followed by the FindNeighbors and

FindClusters functions. For Visium technology, we have incorporated two more clustering tools,

SpaGCN [52] and BayesSpace [53]. SpaGCN leverages graph convolutional networks to

integrate gene expression, spatial location, and histology data, allowing for the identification of

spatial domains with distinct expression patterns [52]. In contrast, BayesSpace is a Bayesian

statistical tool that leverages prior spatial knowledge to cluster transcriptomically similar spots in

proximity [53]. Additionally, BayesSpace offers the capability to enhance resolution by

reallocating gene counts from larger spots in array-based data, such as Visium, to more refined

sub-spots based on spatial information from nearby spots [53].

Cell type annotation

Automated cell type annotation can be achieved through transferring labels from a reference

single-cell RNA-seq dataset. For single-cell-level assays, including CosMx and Xenium, we

incorporated Azimuth to predict the most likely cell type for each cell[54]. A UMAP plot will be

generated after running the pipeline for users to examine the cell type annotation results.

Deconvolution

For deconvolution purposes, we have incorporated two state-of-the-art deconvolution tools,

Tangram [55] and Cell2location [42]. We selected these two packages for their outstanding

performance in a recent benchmarking study[43]. Tangram, a deep-learning framework, excelled

in creating comprehensive single-cell-resolution spatial gene-expression maps and integrating

histological and anatomical data from the same specimens. It adeptly handled low-quality genes

in high-resolution methods, offered single-cell resolution for lower-resolution techniques, and

ensured broad genome-wide coverage for targeted approaches[55]. Meanwhile, Cell2location, a

Bayesian model, showcased its ability to discern fine-grained cell types within spatial

transcriptomic data and constructs detailed cellular maps of diverse tissues. In the first step,

Cell2location estimated reference cell type signatures from single-cell and single-nucleus RNA-

seq profiles, aligned with gene expression profiles for user-specified cell types and

subpopulations. In the subsequent stage, Cell2location utilized these reference signatures in

conjunction with one or more spatial transcriptomic datasets to deconstruct mRNA counts at

individual spatial locations into their respective reference cell types[42].

Ligand receptor interactions

Here, we employ SpaTalk to investigate spatially resolved cell-cell interactions that will enable

us to get insights into tissue homeostasis, development, and disease[56]. This tool leverages a

graph network along with a knowledge graph to model and evaluate the ligand-receptor-target

signaling network operating between neighboring cells[56]. SpaTalk serves as an optional step in

the Visium data processing workflow, providing researchers with a powerful means to unravel

the complex dynamics of inter-cellular interactions within tissues[56].

Standardized output

SpaceSequest generates a series of output files that are ready for result query and further

analysis. The R objects are saved as RData files, and the python anndata objects are saved as

h5ad files. These files contain expression matrices, dimension reduction results, and intermediate

data generated during pipeline processing.

RESULTS

Highlights of SpaceSequest

The pipeline incorporates several key workflows to perform various analyses corresponding to
each platform. Key strengths of this pipeline include:

•   1) Self-explainable configuration files that ensure easy set up.
•   2) Standardized and reproducible data processing powered by R and Python.
•   3) Customized tools are integrated in the pipeline for each spatial platform.
•   4) Comprehensive result generation, including R/Python objects and key figures.
•   5) Streamlined data visualization enabled by cellxgene VIP and Quickomics.

Figure 1. Components of SpaceSequest. The pipeline integrates five major workflows tailored to
perform a series of analysis for each platform, including Visium, Visium HD, Xenium, CosMx,
and GeoMx.

<u>Overall pipeline design</u>

SpaceSequest is a unified toolkit which contains a set of workflows. Each workflow can be
activated by calling the workflow script, and necessary configuration files will be created
accordingly. The main configuration files are in YAML format, ensuring standardized formatting
and flexible modification. To execute the pipeline, in most of the times, users are typically
required to fill in a sample metadata file, which documents the details of each data, such as
sample name, file path, and associated sample-level information. The overall pipeline
components are illustrated in Figure 1.

Visium data analysis incorporates multiple tools

The 10x Genomics Visium platform was released in 2019 and continues to serve as a powerful
commercial spatial transcriptomics technology since then. The Visium slides contain an array of
barcoded 55 µm spots, with two major tissue capture sizes: 6.5x6.5 mm or 11x11 mm.
Sequencing data in FASTQ format are processed through Space Ranger, and SpaceSequest
continues the analysis by utilizing the binned output files. Firstly, SpaceSequest performs
standard quality control on the data by removing spots with low gene detection rates, low UMI
counts, etc. The filtered data are then integrated into a new h5ad file, which is piped to
downstream tools for further analysis.

We tested our pipeline on a public dataset generated from the brain tissue of an Alzheimer's
disease mouse model, which has mutations in APP and PSEN1 genes[57]. We also downloaded
the Azimuth mouse motor cortex single-nucleus RNA-seq data as a reference for cell type
annotation[58]. The data analysis is achieved by calling the *visium* command in SpaceSequest,
which reads all input data and performs a series of quality control steps (Figure 2A). The cleaned
data are then integrated for downstream processing, which includes spatial clustering and cell
type deconvolution results. To leverage the power of parallel computing, SpaceSequest is able to
submit sbatch jobs to execute the following four tools simultaneously: BayesSpace[53],
Cell2location[42], SpaGCN[52], and Tangram[55]. The jobs will be monitored until results have
been generated and saved in separate directories. To provide a few examples, BayesSpace
clustering step will cluster the spots across samples and identify comparable spatial clusters
(Figure 2B). While Cell2location results can be visualized by cell abundance heatmaps overlayed
with the tissue image (Figure 2C). Additionally, the pipeline utilizes SpaTalk for cell-cell
communication analysis, which requires Cell2location output so it is only executed once
Cell2location results are available (Figure S1)[56]. Overall, our Visium data analysis workflow
ensures automated data processing and analysis with a streamlined, one-stop-shop design (Figure
2A).

Figure 2. 10x Visium data analysis. (A) Workflow for Visium data processing. (B) Spatial

clustering results generated by BayesSpace. This run contains four samples, and BayesSpace

identified shared clusters across them. (C) Predicted cell type abundance by Cell2location in one

sample (PSAPP-TAM1), highlighting astrocytes, OPCs, microglia, and endothelial cells.

<u>Streamlined Xenium and CosMx data analysis</u>

The 10x Genomics Xenium and NanoString (now acquired by the Bruker Corporation) CosMx

platforms are both imaging-based, high-plex in-situ technologies capable of capturing up to

thousands of RNAs simultaneously at single-cell resolution. They require a set of probes, usually

from pre-designed panels, with additional space for customization as well. Xenium data consists

of a set of well-defined files in various formats, such as the cell_feature_matrix files (a folder

with three associated files, or a single h5 file), transcript files (transcripts.csv.gz,

transcripts.parquet, transcripts.zarr.zip), and cell summary files (cells.csv.gz, cells.parquet), etc.

The LoadXenium function from the Seurat package reads the necessary files from the directory

and converts them into a Seurat object for downstream analysis.

CosMx data are stored as a collection of plain text files in the comma-separated values (csv)

format. These files include a metadata file (metadata_file), a file storing field of view (fov)

positions (fov_positions_file), an expression matrix file (exprMat_file), a polygon file

(polygons), and a transcript data file (tx_file). Data loading is achieved by the LoadNanostring

function, which converts these csv files to a Seurat object.

SpaceSequest contains two workflows—*xenium* and *cosmx* —to streamline the process from

data loading to downstream analysis. Once Xenium or CosMx data are successfully loaded, these

workflows incorporate standard procedures that perform quality control, clustering, cell type

annotation, and differential expression (DE) analysis, etc (Figure 3A). Moreover, the workflows

generate UMAP and tissue visualizations, enabling users to examine the dimension reduction

and cell type annotation results. Example plots generated using public datasets are shown in

Figure 3B-D. The Xenium data were generated from a healthy human brain FFPE sample

profiled with the Xenium Human Brain Gene Expression Panel containing 266 targets. On the

other hand, the CosMx data were derived from human frontal cortex FFPE tissue using >6,000

target genes. After running the workflow, various analysis results and the processed Seurat

object will be saved for future queries and additional analyses, including further exploration by

Cellxgene VIP[59].

Figure 3. Xenium and CosMx data analysis. (A) Streamlined workflows for analyzing Xenium

and CosMx data. (B) UMAP visualization of Xenium data, labelled by predicted cell types. (C)

Tissue and cell visualization of CosMx data. (D) UMAP visualization of CosMx data with

predict cell types labelled.

<u>Enhanced GeoMx data processing and visualization</u>

The NanoString GeoMx platform is a useful technique to compare gene expression differences in

defined regions on the tissue slides. It uses probes to detect the whole transcriptome in the

selected region of interests (ROIs). However, GeoMx is not a single-cell resolution technique;

instead, it requires a sufficient number of cells within the region to generate robust signals. Thus,

GeoMx is considered as a mini-bulk RNA detection platform, capable of profiling >18,000

major genes for human and mouse samples. In our pipeline, we developed *geomx*, a workflow

that takes DCC files, sample metadata, and a set of comparison groups as input. It performs a

series of analysis, and outputs differential expression analysis results (Figure 4A). This workflow

generates four csv files that are fully compatible with other tools including some Shiny

applications[60, 61], for example, Quickomics, an open-source R Shiny application for flexible

data exploration and visualization[60]. After uploading the data, users can examine PCA plots

(Figure 4B), volcano plots (Figure 4C), heatmaps (Figure 4D), and a number of on-the-fly

visualizations through Quickomics. This strategy not only automates GeoMx data analysis,

especially for studies with many samples, but bridges the gap between raw data and insight

generation through Quickomics, which creates a link to facilitate data storage and future

analysis.

Figure 4. GeoMx data analysis. (A) Overview of GeoMx data analysis workflow. (B) PCA plot

showing samples from two kidney regions, glomerulus and tubule. (C) Volcano plot highlighting

differentially expressed genes between glomerulus and tubule. (D) Heatmap displaying the top

differentially expressed genes between glomerulus and tubule, with samples and genes clustered

by dendrograms.

<u>Standardized data analysis of the cutting-edge Visium HD platform</u>

In 2024, 10x Genomics released a new platform called Visium HD, featuring high-resolution,

gap-free detection of tissue RNAs. This platform is now compatible with FFPE, fixed frozen,

and fresh frozen samples. To keep pace with the rapid evolution of spatial transcriptomics

technologies, we incorporated *visiumhd* into SpaceSequest for Visium HD data processing. The

workflow takes Space Ranger outputs, performs bin-level quality control, dimension reduction,

and cell type annotation when an Azimuth reference dataset is available (Figure 5A). By

executing the pipeline, users receive plots to examine the gene counts (Figure 5B) and cell type

annotation results (Figure 5C). Furthermore, we developed a new function,

getVisiumHDspotCol, to extract the mean RGB intensities when a paired high-resolution

immunofluorescence image is available (Figure 5D). This function facilitates accurate bin

selection using nuclei staining, cell type marker staining, or other biomarker staining, for

example, pathological protein markers.

Figure 5. Visium HD data analysis. (A) Overview of the general processing workflow for Visium

HD data. (B) Heatmap showing the number of UMI counts (nCount) detected across the tissue

section at 8 μm bin resolution. (C) Visualization of annotated cell types on the tissue section. (D)

Illustration of a new function that calculates the average RGB intensities across bins.

DISCUSSION

In this study, we developed SpaceSequest, a unified pipeline for spatial transcriptomics data

processing. Currently, it supports five major platforms and incorporates customized software and

packages to maximize the information that can be extracted from the data. To make the pipeline

accessible, we provided a detailed tutorial aimed at lowering the barrier for researchers to apply

this SpaceSequest to their own studies. Guided by the FAIR principles (Findable, Accessible,

Interoperable, and Reusable), SpaceSequest enhances data access, interpretation, and facilitates

data sharing by generating standardized results. In summary, SpaceSequest is a user-friendly,

end-to-end solution for spatial transcriptomics data analysis and interpretation.

Abbreviations

scRNA-seq: Single-cell RNA-seq

NGS: Next-generation sequencing

csv: comma-separated values

fov: field of view

ST: spatial transcriptomics

UMI: unique molecular identifiers

FISH: fluorescence in situ hybridization

DE: Differential expression

DCC: digital count conversion

PCA: Principal component analysis

OPC: Oligodendrocyte progenitor cells

UMAP: Uniform Manifold Approximation and Projection

Cellxgene VIP: Cellxgene visualization in plug-in

LRI: ligand-receptor interaction

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

Y.H.S., S.P., Y.C., J.G., S.C., B.S., J.Z., W.H., F.C., D.H., and B.Z. are employees of Biogen and hold stocks from the company. H.Z., K.C. and H.Y. were co-ops at Biogen and do not hold Biogen's stocks. Z.O. and M.R. are employees of BioInfoRx Inc. and PharmaLex, respectively.

Funding

This study is funded by Biogen Inc.

Author Contributions Statement

Y.H.S., S.P., and Z.O. developed the pipeline and wrote the manuscript, with the help from Y.C., J.G., S.C., H.Z., B.S., J.Z., K.C., H.Y., W.H., M.R., F.C., D.H., and B.Z. All authors reviewed the manuscript.

Acknowledgement

We acknowledge the scientists at Biogen for their valuable feedback on the pipeline.

Availability and Requirements

No new data were generated from this study. Several public datasets were used to demonstrate SpaceSequest. The 10x Visium data were from a public study using mouse brain tissues with NCBI accession number GSE203424 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE203424). Public data were downloaded from 10x Genomics dataset webpage (https://www.10xgenomics.com/datasets) to test Xenium (https://www.10xgenomics.com/datasets/xenium-human-brain-preview-data-1-standard) and Visium HD (FFPE: https://www.10xgenomics.com/datasets/visium-hd-cytassist-gene-expression-libraries-of-mouse-brain-he, Fixed frozen: https://www.10xgenomics.com/datasets/visium-hd-cytassist-gene-expression-mouse-brain-fixed-frozen) workflows, respectively. Moreover, we used a public human kidney GeoMx dataset (http://nanostring-public-share.s3-website-us-west-2.amazonaws.com/GeoScriptHub/Kidney_Dataset_for_GeomxTools.zip) and a human brain CosMx dataset (https://nanostring.com/products/cosmx-spatial-molecular-imager/ffpe-dataset/human-frontal-cortex-ffpe-dataset/) from NanoString websites to test GeoMx and CosMx pipelines respectively. More details related to data downloading and processing can be found at

the tutorial (https://interactivereport.github.io/SpaceSequest/tutorial/docs/index.html) associated

with this pipeline.

Project name: SpaceSequest

Project home page: https://github.com/interactivereport/spacesequest

Operating system(s): Linux, Mac OS

Programming language: Bash (Unix Shell)

Other requirements: Conda

License: MIT license

Any restrictions to use by non-academics: No

Supplementary figure legends

Figure S1: Examples of SpaTalk results. (A) Highlighted pairs between astrocytes (Astro) and
GABAergic neurons. (B) Significantly enriched ligand-receptor interaction (LRI) pairs between
astrocytes and GABAergic neurons. (C) Highlighted pairs between microglia and macrophage
cells (Micro_PVM) and Glutamatergic neurons. (D) Significantly enriched ligand-receptor
interaction (LRI) pairs between microglia/macrophages and Glutamatergic neurons.

REFERENCES

1. Jovic D, Liang X, Zeng H, Lin L, Xu F, Luo Y. Single-cell RNA sequencing technologies and applications: A brief overview. Clinical & Translational Med. 2022;12:e694. https://doi.org/10.1002/ctm2.694.

2. Wang S, Sun S-T, Zhang X-Y, Ding H-R, Yuan Y, He J-J, et al. The Evolution of Single-Cell RNA Sequencing Technology and Application: Progress and Perspectives. IJMS. 2023;24:2943. https://doi.org/10.3390/ijms24032943.

3. Potter SS. Single-cell RNA sequencing for the study of development, physiology and disease. Nat Rev Nephrol. 2018;14:479–92. https://doi.org/10.1038/s41581-018-0021-7.

4. Du J, Yang Y-C, An Z-J, Zhang M-H, Fu X-H, Huang Z-F, et al. Advances in spatial transcriptomics and related data analysis strategies. J Transl Med. 2023;21:330. https://doi.org/10.1186/s12967-023-04150-2.

5. Adema K, Schon MA, Nodine MD, Kohlen W. Lost in space: what single-cell RNA sequencing cannot tell you. Trends in Plant Science. 2024;29:1018–28. https://doi.org/10.1016/j.tplants.2024.03.010.

6. Williams CG, Lee HJ, Asatsuma T, Vento-Tormo R, Haque A. An introduction to spatial transcriptomics for biomedical research. Genome Med. 2022;14:68. https://doi.org/10.1186/s13073-022-01075-1.

7. Longo SK, Guo MG, Ji AL, Khavari PA. Integrating single-cell and spatial transcriptomics to elucidate intercellular tissue dynamics. Nat Rev Genet. 2021;22:627–44. https://doi.org/10.1038/s41576-021-00370-8.

8. Saviano A, Henderson NC, Baumert TF. Single-cell genomics and spatial transcriptomics: Discovery of novel cell states and cellular interactions in liver physiology and disease biology. Journal of Hepatology. 2020;73:1219–30. https://doi.org/10.1016/j.jhep.2020.06.004.

9. Li X, Wang C-Y. From bulk, single-cell to spatial RNA sequencing. Int J Oral Sci. 2021;13:36. https://doi.org/10.1038/s41368-021-00146-0.

10. Moses L, Pachter L. Museum of spatial transcriptomics. Nat Methods. 2022;19:534–46. https://doi.org/10.1038/s41592-022-01409-2.

11. Tian L, Chen F, Macosko EZ. The expanding vistas of spatial transcriptomics. Nat Biotechnol. 2023;41:773–82. https://doi.org/10.1038/s41587-022-01448-2.

12. Wang Y, Liu B, Zhao G, Lee Y, Buzdin A, Mu X, et al. Spatial transcriptomics: Technologies, applications and experimental considerations. Genomics. 2023;115:110671. https://doi.org/10.1016/j.ygeno.2023.110671.

13. Jain S, Eadon MT. Spatial transcriptomics in health and disease. Nat Rev Nephrol. 2024;20:659–71. https://doi.org/10.1038/s41581-024-00841-1.

14. Piwecka M, Rajewsky N, Rybak-Wolf A. Single-cell and spatial transcriptomics: deciphering brain complexity in health and disease. Nat Rev Neurol. 2023;19:346–62. https://doi.org/10.1038/s41582-023-00809-y.

15. Zhang L, Xiong Z, Xiao M. A Review of the Application of Spatial Transcriptomics in Neuroscience.
Interdiscip Sci Comput Life Sci. 2024;16:243–60. https://doi.org/10.1007/s12539-024-00603-4.

16. Jung N, Kim T-K. Spatial transcriptomics in neuroscience. Exp Mol Med. 2023;55:2105–15.
https://doi.org/10.1038/s12276-023-01093-y.

17. Tsai H-H, Piya S, Wang J, Zhu J, Hu W, Gehrke AR, et al. Spatial transcriptomics reveals
heterogeneous cell–cell interactions among brain regions in a cuprizone model consistent with multiple
sclerosis lesions. bioRxiv. 2025;:2025.02.07.637125. https://doi.org/10.1101/2025.02.07.637125.

18. Berglund E, Maaskola J, Schultz N, Friedrich S, Marklund M, Bergenstråhle J, et al. Spatial maps of
prostate cancer transcriptomes reveal an unexplored landscape of heterogeneity. Nat Commun.
2018;9:2419. https://doi.org/10.1038/s41467-018-04724-5.

19. Vandereyken K, Sifrim A, Thienpont B, Voet T. Methods and applications for single-cell and spatial
multi-omics. Nat Rev Genet. 2023;24:494–515. https://doi.org/10.1038/s41576-023-00580-2.

20. Ma Y, Zhou X. Spatially informed cell-type deconvolution for spatial transcriptomics. Nat Biotechnol.
2022;40:1349–59. https://doi.org/10.1038/s41587-022-01273-7.

21. Shi H, He Y, Zhou Y, Huang J, Maher K, Wang B, et al. Spatial atlas of the mouse central nervous
system at molecular resolution. Nature. 2023;622:552–61. https://doi.org/10.1038/s41586-023-06569-
5.

22. Huuki-Myers LA, Spangler A, Eagles NJ, Montgomery KD, Kwon SH, Guo B, et al. A data-driven single-
cell and spatial transcriptomic map of the human prefrontal cortex. Science. 2024;384:eadh1938.
https://doi.org/10.1126/science.adh1938.

23. Qian X, Coleman K, Jiang S, Kriz AJ, Marciano JH, Luo C, et al. Spatial transcriptomics reveals human
cortical layer and area specification. Nature. 2025. https://doi.org/10.1038/s41586-025-09010-1.

24. Han L, Liu Z, Jing Z, Liu Y, Peng Y, Chang H, et al. Single-cell spatial transcriptomic atlas of the whole
mouse brain. Neuron. 2025;113:2141-2160.e9. https://doi.org/10.1016/j.neuron.2025.02.015.

25. Lake BB, Menon R, Winfree S, Hu Q, Melo Ferreira R, Kalhor K, et al. An atlas of healthy and injured
cell states and niches in the human kidney. Nature. 2023;619:585–94. https://doi.org/10.1038/s41586-
023-05769-3.

26. Wu H, Liu F, Shangguan Y, Yang Y, Shi W, Hu W, et al. Integrating spatial transcriptomics with single-
cell transcriptomics reveals a spatiotemporal gene landscape of the human developing kidney. Cell
Biosci. 2022;12:80. https://doi.org/10.1186/s13578-022-00801-x.

27. Chen D, Shao M, Song Y, Ren G, Guo F, Fan X, et al. Single-cell RNA -seq with spatial transcriptomics
to create an atlas of human diabetic kidney disease. The FASEB Journal. 2023;37:e22938.
https://doi.org/10.1096/fj.202202013RR.

28. Mohenska M, Tan NM, Tokolyi A, Furtado MB, Costa MW, Perry AJ, et al. 3D-cardiomics: A spatial
transcriptional atlas of the mammalian heart. Journal of Molecular and Cellular Cardiology.
2022;163:20–32. https://doi.org/10.1016/j.yjmcc.2021.09.011.

29. Lee SE, Joo JH, Hwang HS, Chen S-F, Evans D, Lee KY, et al. Spatial transcriptional landscape of human
heart failure. European Heart Journal. 2025;:ehaf272. https://doi.org/10.1093/eurheartj/ehaf272.

30. Chen H, Murray E, Sinha A, Laumas A, Li J, Lesman D, et al. Dissecting mammalian spermatogenesis
using spatial transcriptomics. Cell Reports. 2021;37:109915.
https://doi.org/10.1016/j.celrep.2021.109915.

31. Madissoon E, Oliver AJ, Kleshchevnikov V, Wilbrey-Clark A, Polanski K, Richoz N, et al. A spatially
resolved atlas of the human lung characterizes a gland-associated immune niche. Nat Genet.
2023;55:66–77. https://doi.org/10.1038/s41588-022-01243-4.

32. Vannan A, Lyu R, Williams AL, Negretti NM, Mee ED, Hirsh J, et al. Spatial transcriptomics identifies
molecular niche dysregulation associated with distal lung remodeling in pulmonary fibrosis. Nat Genet.
2025;57:647–58. https://doi.org/10.1038/s41588-025-02080-x.

33. Zhang Y, Miller JA, Park J, Lelieveldt BP, Long B, Abdelaal T, et al. Reference-based cell type matching
of in situ image-based spatial transcriptomics data on primary visual cortex of mouse brain. Sci Rep.
2023;13:9567. https://doi.org/10.1038/s41598-023-36638-8.

34. Chen W-T, Lu A, Craessaerts K, Pavie B, Sala Frigerio C, Corthout N, et al. Spatial Transcriptomics and
In Situ Sequencing to Study Alzheimer's Disease. Cell. 2020;182:976-991.e19.
https://doi.org/10.1016/j.cell.2020.06.038.

35. Yu M, Risacher SL, Nho KT, Wen Q, Oblak AL, Unverzagt FW, et al. Spatial transcriptomic patterns
underlying amyloid-β and tau pathology are associated with cognitive dysfunction in Alzheimer's
disease. Cell Reports. 2024;43:113691. https://doi.org/10.1016/j.celrep.2024.113691.

36. Navarro JF, Croteau DL, Jurek A, Andrusivova Z, Yang B, Wang Y, et al. Spatial Transcriptomics
Reveals Genes Associated with Dysregulated Mitochondrial Functions and Stress Signaling in Alzheimer
Disease. iScience. 2020;23:101556. https://doi.org/10.1016/j.isci.2020.101556.

37. Zheng P, Zhang N, Ren D, Yu C, Zhao B, Zhang Y. Integrated spatial transcriptome and metabolism
study reveals metabolic heterogeneity in human injured brain. Cell Reports Medicine. 2023;4:101057.
https://doi.org/10.1016/j.xcrm.2023.101057.

38. Ferri-Borgogno S, Zhu Y, Sheng J, Burks JK, Gomez JA, Wong KK, et al. Spatial Transcriptomics Depict
Ligand–Receptor Cross-talk Heterogeneity at the Tumor-Stroma Interface in Long-Term Ovarian Cancer
Survivors. Cancer Research. 2023;83:1503–16. https://doi.org/10.1158/0008-5472.CAN-22-1821.

39. Moreno J, Gluud LL, Galsgaard ED, Hvid H, Mazzoni G, Das V. Identification of ligand and receptor
interactions in CKD and MASH through the integration of single cell and spatial transcriptomics. PLoS
ONE. 2024;19:e0302853. https://doi.org/10.1371/journal.pone.0302853.

40. Rodriques SG, Stickels RR, Goeva A, Martin CA, Murray E, Vanderburg CR, et al. Slide-seq: A scalable
technology for measuring genome-wide expression at high spatial resolution. Science. 2019;363:1463–7.
https://doi.org/10.1126/science.aaw1219.

41. Vickovic S, Eraslan G, Salmén F, Klughammer J, Stenbeck L, Schapiro D, et al. High-definition spatial
transcriptomics for in situ tissue profiling. Nat Methods. 2019;16:987–90.
https://doi.org/10.1038/s41592-019-0548-y.

42. Kleshchevnikov V, Shmatko A, Dann E, Aivazidis A, King HW, Li T, et al. Cell2location maps fine-
grained cell types in spatial transcriptomics. Nat Biotechnol. 2022;40:661–71.
https://doi.org/10.1038/s41587-021-01139-4.

43. Li H, Zhou J, Li Z, Chen S, Liao X, Zhang B, et al. A comprehensive benchmarking with practical
guidelines for cellular deconvolution of spatial transcriptomics. Nat Commun. 2023;14:1548.
https://doi.org/10.1038/s41467-023-37168-7.

44. Navarro JF, Sjöstrand J, Salmén F, Lundeberg J, Ståhl PL. ST Pipeline: an automated pipeline for
spatial mapping of unique transcripts. Bioinformatics. 2017;33:2591–3.
https://doi.org/10.1093/bioinformatics/btx211.

45. Sztanka-Toth TR, Jens M, Karaiskos N, Rajewsky N. Spacemake: processing and analysis of large-scale
spatial transcriptomics data. GigaScience. 2022;11:giac064.
https://doi.org/10.1093/gigascience/giac064.

46. Moriel N, Senel E, Friedman N, Rajewsky N, Karaiskos N, Nitzan M. NovoSpaRc: flexible spatial
reconstruction of single-cell gene expression with optimal transport. Nat Protoc. 2021;16:4177–200.
https://doi.org/10.1038/s41596-021-00573-7.

47. Cisar C, Keener N, Ruffalo M, Paten B. A unified pipeline for FISH spatial transcriptomics. Cell
Genomics. 2023;3:100384. https://doi.org/10.1016/j.xgen.2023.100384.

48. Kamel M, Song Y, Solbas A, Villordo S, Sarangi A, Senin P, et al. ENACT: End-to-End Analysis of Visium
High Definition (HD) Data. Bioinformatics. 2025;41:btaf094.
https://doi.org/10.1093/bioinformatics/btaf094.

49. Curion F, Rich-Griffin C, Agarwal D, Ouologuem S, Rue-Albrecht K, May L, et al. Panpipes: a pipeline
for multiomic single-cell and spatial transcriptomic data analysis. Genome Biol. 2024;25:181.
https://doi.org/10.1186/s13059-024-03322-7.

50. Hao Y, Stuart T, Kowalski MH, Choudhary S, Hoffman P, Hartman A, et al. Dictionary learning for
integrative, multimodal and scalable single-cell analysis. Nat Biotechnol. 2024;42:293–304.
https://doi.org/10.1038/s41587-023-01767-y.

51. Virshup I, Rybakov S, Theis FJ, Angerer P, Wolf FA. anndata: Annotated data. bioRxiv.
2021;:2021.12.16.473007. https://doi.org/10.1101/2021.12.16.473007.

52. Hu J, Li X, Coleman K, Schroeder A, Ma N, Irwin DJ, et al. SpaGCN: Integrating gene expression,
spatial location and histology to identify spatial domains and spatially variable genes by graph
convolutional network. Nat Methods. 2021;18:1342–51. https://doi.org/10.1038/s41592-021-01255-8.

53. Zhao E, Stone MR, Ren X, Guenthoer J, Smythe KS, Pulliam T, et al. Spatial transcriptomics at subspot
resolution with BayesSpace. Nat Biotechnol. 2021;39:1375–84. https://doi.org/10.1038/s41587-021-
00935-2.

54. Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A, et al. Integrated analysis of
multimodal single-cell data. Cell. 2021;184:3573-3587.e29. https://doi.org/10.1016/j.cell.2021.04.048.

55. Biancalani T, Scalia G, Buffoni L, Avasthi R, Lu Z, Sanger A, et al. Deep learning and alignment of
spatially resolved single-cell transcriptomes with Tangram. Nat Methods. 2021;18:1352–62.
https://doi.org/10.1038/s41592-021-01264-7.

56. Shao X, Li C, Yang H, Lu X, Liao J, Qian J, et al. Knowledge-graph-based cell-cell communication
inference for spatially resolved transcriptomic data with SpaTalk. Nat Commun. 2022;13:4429.
https://doi.org/10.1038/s41467-022-32111-8.

57. Castranio EL, Hasel P, Haure-Mirande J, Ramirez Jimenez AV, Hamilton BW, Kim RD, et al. Microglial
INPP5D limits plaque formation and glial reactivity in the PSAPP mouse model of Alzheimer's disease.
Alzheimer's & Dementia. 2023;19:2239–52. https://doi.org/10.1002/alz.12821.

58. Yao Z, Van Velthoven CTJ, Nguyen TN, Goldy J, Sedeno-Cortes AE, Baftizadeh F, et al. A taxonomy of
transcriptomic cell types across the isocortex and hippocampal formation. Cell. 2021;184:3222-
3241.e26. https://doi.org/10.1016/j.cell.2021.04.021.

59. Li K, Ouyang Z, Chen Y, Gagnon J, Lin D, Mingueneau M, et al. Cellxgene VIP unleashes full power of
interactive visualization and integrative analysis of scRNA-seq, spatial transcriptomics, and multiome
data. bioRxiv. 2022;:2020.08.28.270652. https://doi.org/10.1101/2020.08.28.270652.

60. Gao B, Zhu J, Negi S, Zhang X, Gyoneva S, Casey F, et al. Quickomics: exploring omics data in an
intuitive, interactive and informative manner. Bioinformatics. 2021;37:3670–2.
https://doi.org/10.1093/bioinformatics/btab255.

61. Gao B, Sun YH, Zhang X, Lin T, Li W, Admanit R, et al. xOmicsShiny: an R Shiny application for cross-
omics data analysis and pathway mapping. Bioinformatics Advances. 2024;5:vbaf097.
https://doi.org/10.1093/bioadv/vbaf097.

566