Title:

A methodological framework for assessing neural control of shoulder muscles during multi-
directional isometric tasks using high-density surface electromyography.

Authors:

J Greig Inglis[1], Silvia Rio[1], Hélio V. Cabral[1,2,3], Caterina Cosentino[1], Roberto Pagani[4], Clark
R Dickerson[5], Francesco Negro[1]

Affiliations:

[1]Department of Clinical and Experimental Sciences, Università degli Studi di Brescia, Brescia,
Italy
[2]School of Physical Education and Sports, Universidade Federal do Rio de Janeiro, Rio de
Janeiro, Brazil
[3]Biomedical Engineering Program (COPPE), Universidade Federal do Rio de Janeiro, Rio de
Janeiro, Brazil
[4]Department of Mechanical and Industrial Engineering, Università degli Studi di Brescia,
Brescia, Italy
[5]Department of Kinesiology and Health Sciences, University of Waterloo, Waterloo, Canada

Conflict of Interest:

The authors report no conflict of interest.

Acknowledgements:

The authors would like to thank Dr. Tea Lulic-Kuryllo for her input in the protocol and array
placement which were integral to the design of the data collection.

Funding:

This study was funded by the European Research Council Consolidator Grant 'INcEPTION',
contract no. 101045605. J Greig Inglis was supported by the Marie Skłodowska-Curie Actions
Grant 'MUDecomp', agreement no. 101151712.

Corresponding author:
Prof. Francesco Negro
Department of Clinical and Experimental Sciences,
Università degli Studi di Brescia
Viale Europa 11, Brescia, 25121, Italy
Email: francesco.negro@unibs.it

ABSTRACT

The complex movements of the shoulder rely on the coordinated activation of large muscles including the deltoid, pectoralis major, trapezius and latissimus dorsi. However, detailed knowledge of their neuromuscular control remains limited. This study aimed to develop a methodological framework to investigate the neural control of shoulder muscles by combining a 6-degree-of-freedom load cell attached to an industrial robotic arm with high-density surface electromyograms (HDsEMG). Six healthy participants performed isometric contractions (abduction, adduction, flexion and extension) at 30% of maximal voluntary contraction with the shoulder positioned at 30° and 65° of lateral abduction. HDsEMGs were recorded from the four muscles and analysed at both the global activation and motor unit levels. Global activation was quantified using root-mean-square (RMS) amplitude and topographic maps. Moreover, HDsEMGs were decomposed into individual motor unit spike trains and motor unit behaviour was characterized by mean discharge rate, coefficient of variation of inter-spike interval, and spatial distribution of motor unit action potentials (MUAPs). RMS maps revealed movement-specific activation within and between muscles, with the deltoid and trapezius active across all tasks, while the pectoralis and latissimus dorsi were predominantly activated during flexion and extension. Motor unit discharge rates also showed task-dependent recruitment. MUAP spatial distributions further showed distinct motor unit territories within arrays, suggesting region-specific recruitment strategies across movements. In conclusion, this framework demonstrates that individual motor unit activity can be reliably measured non-invasively in the main superficial shoulder muscles. The approach provides a methodological basis for incorporating neural control into biomechanical models of shoulder function.

Keywords: motor unit discharge rate, HDsEMG spatial distribution, neural control, shoulder biomechanics.

1.0 INTRODUCTION

Movement of the humerus requires the complex activation of the muscles of the glenohumeral joint to contract and, ultimately, cause the desired action. During these movements, the activation of large muscles such as the *trapezius, pectoralis, latissimus dorsi,* and *deltoid* must be precisely coordinated with several smaller muscles to generate torque across multiple joints and adapt to changes in both task demands and musculoskeletal alterations. According to Hughes and colleagues (1999), the muscles of the glenohumeral joint (from here on to be referred to as shoulder throughout) and their strength are an important part of functional capacity. Thus, understanding the neuromuscular behaviours of these muscles is essential for the complete understanding of this muscle group which can be used for the prescription of appropriate training and rehabilitation strategies. Biomechanical modelling of the shoulder relies on simple mechanical analysis of the body segments and the internal and external forces acting on the joint to predict the control of the muscles of the shoulder (Hughes & An, 1997; Chang *et al.*, 2000; Dickerson *et al.*, 2008). However, to date, there are restrictions to modelling advances in the shoulder, particularly due to limitations in neural control investigations of the active muscles. Specifically, these limitations stem from the methodological approach of bipolar surface electromyogram recordings to infer the intricate behaviour of the individual motor units involved in muscle activation. Additionally, to date, the addition of motoneuron behaviour from the muscles of the shoulder remains elusive (Hughes & An, 1997). Therefore, shoulder models must rely on the indirect measurement of neural control through the measurement of metrics which has been proven to be ineffective in measuring motor unit behaviour (Farina *et al.*, 2010*a*).

Classically, the approach to assessing the neural control has been through global bipolar electromyography (EMG), using amplitude measures to infer neuromuscular adaptations to specific movements with applications in biomechanics (Lawrence & De Luca, 1983; Woods

& Bigland-Ritchie, 1983; Hermans *et al.*, 1999; Day & Hulliger, 2001; Karlsson & Gerdle,

2001; Keenan *et al.*, 2005; Farina *et al.*, 2010*b*). However, the specificity of the information

gained from the application of the amplitude of EMG activity, in terms of inferring neural drive

to the muscle, is subject to debate (Farina *et al.*, 2010*b*). Waveform cancellation is particularly

a key factor affecting the relationship between EMG amplitude and neural drive (Yao *et al.*,

2000; Day & Hulliger, 2001; Dimitrova & Dimitrov, 2003; Keenan *et al.*, 2005), because as

additional motor units may be recruited during the gradation of force, the superposition of

positive and negative phases of action potentials could result in little to no increase in EMG

amplitude (Day & Hulliger, 2001; Keenan *et al.*, 2005). Additionally, bipolar EMG provides

only muscle activity in the area closest to the small electrode (i.e., local information). In the

case of the shoulder where large muscles produce complex movements in multiple degrees-of-

freedom, it is possible that not all muscle areas are active throughout all movements (i.e.,

within-muscle localized activation). For instance, previous evidence suggested that localized

regions of the deltoid, pectoralis major and latissimus dorsi can be activated based on the

specific shoulder movement direction (Wickham & Brown, 1998, 2012; Brown *et al.*, 2007).

The early 1980's advancements in myoelectric recordings have led to high-density

surface electromyograms (HDsEMG) (Masuda *et al.*, 1983, 1985; Reucher *et al.*, 1987;

Merletti *et al.*, 1999). The benefit to recording with the evolving HDsEMG technology is its

non-invasive nature and the ability to provide the spatial distribution of activation within and

across muscles during complex movements. Additionally, when combined with motor unit

decomposition algorithms (Holobar & Zazula, 2007; Ning *et al.*, 2015; Chen & Zhou, 2016;

Negro *et al.*, 2016), this approach has the potential to record a large number of concurrently

discharging motoneurons during voluntary contractions (Inglis *et al.*, 2025). As the alpha

motoneuron is the final common pathway of the neuromuscular system integrating the synaptic

inputs from spinal and supraspinal pathways (Liddell & Sherrington, 1925; Sherrington, 1925),

the analysis of the discharge behaviour of motor units provides an effective way to assemble information regarding the generation and control of movement. A recent systematic review and meta-analysis revealed that most motor unit behaviour investigations focused on the lower-compared to upper-body which has led to a paucity of data on the upper limb (Inglis *et al.*, 2025). Additionally, it was revealed that of the upper limb muscles, specifically major movers of the shoulder, there are few motor unit behaviour studies investigating the *upper trapezius* (Westgaard & De Luca, 2001; Falla & Farina, 2008*a*, 2008*b*; Gerdle *et al.*, 2008; Holobar *et al.*, 2009; Dideriksen *et al.*, 2016; Kirk *et al.*, 2019) and *clavicular pectoralis* (Lulic-Kuryllo *et al.*, 2021*b*) using invasive or non-invasive techniques*,* none that examine the *latissimus dorsi* (Padovan *et al.*, 2024) and only three examining the *deltoid* (de Luca *et al.*, 1982; Doherty & Stashuk, 2003; Baslo *et al.*, 2024) all which are done with intramuscular means.

Therefore, the aim of this paper was to develop a methodological framework to investigate four isometric shoulder actions (abduction, adduction, flexion and extension) at two shoulder angles to fully assess the neural control of the main superficial shoulder muscles (deltoid, pectoralis major, trapezius and latissimus dorsi) both at the global activation level (signal amplitude) and at the motor unit level (discharge patterns). For this, we developed a framework combining 6-degree-of-freedom load cell attached to an industrial robotic arm while recording EMGs through the evolving HDsEMG technology. Additionally, this paper provides several examples of the type of neural information of shoulder control that can be extracted within the proposed framework. The novel framework proposed here will lead to the establishment of guidelines for future investigations on the shoulder complex and serve to propel shoulder modelling forward in these muscles.

2.0 METHODS

2.1 Participants

Six healthy individuals free of any musculoskeletal or neuromuscular abnormalities (1

female; mean ± SD: age 28 ± 4 years; height 180 ± 7 cm; mass 79 ± 9 kg) volunteered to

participate in this study. All participants provided written informed consent before starting the

experiments that conformed to the standards set by the latest version of the Declaration of

Helsinki which was approved by the ethics committee of the Università degli Studi di Brescia

(code NP5890).

2.2 Experimental protocol

2.2.1 Experimental setup

Each participant visited the lab on a single day where, in a randomized order, they

performed multiple degree-of-freedom isometric contractions of the shoulder in a 2-hour

session. Participants were seated in a rigid testing chair with their hip and knee joints at 90° of

flexion (0° indicates full extension) and a broad Velcro® strap placed around the midsection

to minimize lateral torso movements. The participant's arm was placed slightly (<20°) in the

scapular plane to avoid discomfort when performing the selected movements. For the

movements, the participants' upper arm was placed into a custom-build testing jig

manufactured from rectangular metal tubing to ensure a rigid transfer of forces produced to the

load cell (**Figure 1b**). The testing jig was attached to a 6-degrees of freedom load cell (© 2025

Advanced Mechanical Technology Inc., Watertown MA, USA, model #MC3A) to measure

force output (**Figure 1c**). The load cell has a capacity of F_x, F_y of 225 N and F_z of 445 N and a

sensitivity of F_x, F_y of 5.4 µV/V-N and F_z of 1.35 µV/V-N. The load cell was attached to a

robotic arm (Kawasaki Robotics Inc., Wixom MI, USA; **Figure 1a**) with a custom machined

aluminium coupling (**Figure 1d**). The importance of implementing the use of the robotic arm

was to position the testing jig and loadcell through fine adjustments to accommodate individual

anatomical landmarks to ensure participants performed each movement in the identical relative

position to record with precision the forces produced during each movement.

Figure 1. **Mechanical components designed and manufactured for the robotic setup** *(a)*
Complete robot assembly showing the robot with all components mounted. (b) 3D model and
2D technical drawings of the jig. (c) 3D model and 2D technical drawings of the six degree-
of-freedom load cell. (d) 3D model and 2D technical drawings of the coupling between the
robotic arm and the load cell. Measurements are reported in millimeters (mm).

2.2.2 Experimental tasks

Each participant performed four isometric shoulder contractions at 30º and 65º of lateral

shoulder abduction, the movements were: abduction, adduction, flexion and extension. **Figure**

2 illustrates the position of the participant during the four movements across two shoulder

angels. In the figure the direction of the force vector is depicted with arrows per movement.

Each movement was performed at two shoulder angles to alter the deltoid muscle length (30°

long; 65° short). All experimental actions were repeated twice for each movement and shoulder

angle resulting in a total of 16 trials. Prior to each condition (angle/movement) participants

performed three voluntary maximum isometric contractions (MVCs) for 3-5 s, with a 3-min

rest between each bout. The greatest value across the three MVCs was considered as the

maximal isometric shoulder force output for the individual condition and used as a reference

to set the target of the submaximal contractions. Subsequently, participants underwent a short

familiarization period, during which they were instructed to exert isometric shoulder forces at

10% MVC for approximately 10 s. This familiarization task was repeated up to a maximum of

five times to ensure that participants could maintain the force output as steadily as possible at

a target level, which was visually assessed by a researcher. Following the familiarization

period, participants were instructed to perform two isometric contractions following a

trapezoidal profile. The trapezoidal pathway involved a linear increase at a rate of 3% MVC/s,

a plateau phase for 30 s, and a linear decrease at a rate of 3% MVC/s. The target force output

was set at 30% MVC (i.e., moderate force output) to mimic common outputs during activities

of daily living (Lulic-Kuryllo *et al.*, 2021*a*).

189
Figure 2. Experimental Movements. *Shoulder abduction/adduction (left, anterior view) and*
flexion/extension (right, lateral view) performed at 30° (top) and 65° (bottom) of shoulder
abduction. The arm was secured to a passive robotic interface, which secured the participant
in position for each given movement. A broad black Velcro® strap placed around the
midsection and secured to the testing chair to minimize movements of the trunk.

2.2.3 Force feedback

A novel graphical user interface (GUI) was developed in MATLAB (version 2024a) to

provide visual feedback of the force output and rate at which the force was to be developed

during the trapezoidal profile (**Figure 3e**). Force feedback was provided based on the axis

being investigated relative to the 3-axis load cell connected to the PC by an A-D board

(National Instruments CORP., NI USB-6343). The selected axis was represented by either a

horizontal or vertical line where the Y direction was utilized for abduction and adduction and

the Z direction during flexion and extension (frontal) relative to the load cell. The GUI provided

a green target box with the dimensions set to $\pm$ 2.5 %MVC and a blue circle to represent the

changes in force output. The participant was instructed to maintain the blue circle inside the

square. In cases where the error in force production exceeded 2.5% MVC, the square would

turn red, and the participant would be encouraged to meet the target force output. Visual

feedback was displayed on a monitor placed one meter at eye level in front of the participant.

2.3 Data collection

HDsEMGs were acquired from the four major shoulder girdle movers (deltoid,

pectoralis major, upper trapezius and latissimus dorsi) during the submaximal isometric

movements using 6 two-dimensional adhesive arrays (HD08MM1305 and HD08MM1606; 1

mm diameter; 8 mm inter-electrode distance; OT Bioelettronica, Turin, Italy). The arrays

placed on the deltoid and pectoralis major (see below) consisted of 64 electrodes arranged in

13 rows by 5 columns, with one missing channel in the top left corner. As for the upper

trapezius and the latissimus dorsi, the arrays consisted of 96 electrodes arranged in 16 rows by

6 columns. Prior to array and ground placement, the area was shaved, mildly abraded (EVERI,

Spes Medica, Genova, Italy) and cleaned with distilled water. The electrode-skin contact was

ensured by filling the foam cavities with conductive paste (AC cream, Spes Medica, Genova,

Italy). Two reference electrodes were used; one positioned on the right elbow at the level of

the epicondyles and other on the acromion process.

We piloted several arrays and configurations to achieve the greatest motor unit yield in

the deltoid (see appendix A). For the anterior/middle deltoid, the electrode was placed

approximately 2 cm distal to the proximal deltoid tendon, starting from the anterior border

horizontally towards the posterior portion perpendicular to the insertion of the middle fibres to

the humerus (*array 1*, **Figure 3a**). To cover more of this anterior and middle aspect of the

deltoid a second array was place distally and slightly offset (2 array columns) and parallel to

the proximal array (*array 2*, **Figure 3a**). For the posterior deltoid, the electrode was positioned

in line with the posterior aspect of the deltoid adjacent to the distal deltoid array (*array 3*,

Figure 3a). For the clavicular region of the pectoralis major, the electrode was placed

approximately 2 cm inferior to the clavicle (superior array in (Lulic-Kuryllo *et al.*, 2021*a*);

array 4, **Figure 3b**). For the upper trapezius, the electrode array was positioned to bisect the

7^{th} and 8^{th} row (*array 5*, **Figure 3c**) along the line drawn between C7 and the lateral edge of

the acromion due to the use of the 96-electrode array which is larger (3 extra rows and 1 extra

column) than the standard 64 electrode array used in the deltoid and pectorals above (modified

from (Falla & Farina, 2008*b*). For the latissimus dorsi, a line was drawn approximately 3 cm

inferior and horizontal to the posterior axillary crease and parallel to the superior aspect of the

iliac crest. From these lines a diagonal line was drawn to follow the latissimus dorsi muscle

fibres, and the array was placed in the middle of this area ensuring that majority of the array

covered the bulk of the muscle (Padovan *et al.*, 2024) (*array 6*, **Figure 3d**).

Figure 3. Electrode placement, visual feedback and HDsEMG analysis. *(a) Electrode placement for the deltoid. Array 1 was positioned horizontally approximately 2 cm distal from the insertion of the deltoid tendon. Array 2 was placed distally and slightly offset to cover the middle and anterior deltoid. Array 3 was positioned vertically adjacent to the distal deltoid array. (b) Electrode placement for the clavicular region of the pectoralis major. Array 4 was positioned horizontally and approximately 2 cm inferior to the clavicle, avoiding the sternum. (c) Electrode placement of the trapezius. Array 5 was positioned to bisect the 7^{th} and 8^{th} row of the array along the line between C7 and the acromion process. (d) Electrode placement of the latissimus dorsi. Array 6 was positioned diagonally in the middle of an area delimited by two horizontal lines, drawn approximately 3 cm inferior to the posterior axillary crease and parallel to the superior aspect of the iliac crest, the array was placed in line with the muscle fibers. (e) Visual feedback of the force output. Visual feedback was provided from a line, a square box and a circle representing the pathway, the target force and the force output of the participant, respectively. (f) Schematic representation of the HDsEMG decomposition. HDsEMGs were visually inspected and, subsequently, decomposed into motor unit innervation pulse train (IPTs). The 30-s plateau phase was considered for this analysis (light gray rectangle).*

HDsEMG and force outputs were digitized synchronously at a sampling frequency of 2000 Hz using a 16-bit A/D converter (10-500 Hz bandwidth; Novecento+, OT Bioelettronica, Turin, Italy). HDsEMG signals were recorded in monopolar derivation and amplified (8 V/V) to maximize signal resolution while avoiding saturation.

2.4 Data analysis

HDsEMG recordings were analysed offline in MATLAB (version 2024a) using

custom-written scripts. As potential applications of the developed experimental framework,

analyses were performed both at the global HDsEMG level (signal amplitude) and at the motor

unit level (discharge patterns) to assess the neural control of shoulder muscles. For all analyses,

the 30-s plateau phase of each contraction was selected. HDsEMGs were bandpass filtered

between 20 and 500 Hz using a third-order Butterworth filter and, after that, visually inspected

to identify and remove bad channels due to electrode-skin contact issues or artifacts (mean ±

SD: 2 ± 1 channels removed for all movements and shoulder angles).

2.4.1 HDsEMG root-mean-square amplitude

To assess the global activation of the investigated muscles, the root-mean-square

(RMS) amplitude was used. Initially, single-differential (bipolar) EMGs were calculated as

the algebraic difference between monopolar signals detected by consecutive rows of electrodes.

Then, the RMS amplitude was calculated for each of the 59 (64-electrodes array) and 80 (96-

electrodes array) single-differential channels using non-overlapping windows of 250 ms (Mark

Burden *et al.*, 2014). The average value across windows was considered for further analysis,

providing topographic maps of RMS amplitude for each of the investigated muscles. For

simplicity, only *array 2* of the deltoid was analysed. The average RMS amplitude across all

channels was then considered as the global RMS amplitude of the muscle for each movement

and shoulder angle.

2.4.2 HDsEMG decomposition

HDsEMG signals were completely decomposed into their constituent motor unit spike

trains using a convolutive blind-source separation algorithm (Negro *et al.*, 2016). Briefly, after

extending and whitening the HDsEMG signals, a fixed-point algorithm that seeks sources that

maximize a measure of sparsity was applied to identify the motor unit spike trains (**Figure 3f**).

The spikes were separated from the noise using the K-means algorithm and, while iteratively

updating the motor unit separation vectors, the discharge times estimation was further refined

by minimizing the coefficient of variation of the inter-spike intervals. This decomposition

method has been previously validated (Negro *et al.*, 2016) and extensively applied to assess

the activity of single motor units (Negro *et al.*, 2016; Cogliati *et al.*, 2020; Cabral *et al.*, 2024).

After the automatic identification of motor units, any missed or misidentified motor unit

discharges which produced non-physiological discharge rates (< 4 pulses per second or > 50

pulses per second; (Negro *et al.*, 2009) were manually identified and iteratively edited by an

experienced operator, and the subsequent motor unit spike trains were re-estimated (Martinez-

Valdes *et al.*, 2017; Hassan *et al.*, 2020). This approach has been shown to be highly reliable

across operators (Hug *et al.*, 2021). Only motor units with a silhouette (SIL) value, which is a

quality control measure of the decomposition accuracy, greater than 0.87 (Negro *et al.*, 2016)

(Negro *et al.*, 2016) were included in the further analysis.

2.4.3 Motor unit discharge behaviour

The instantaneous discharge rate of each identified motor unit was determined by

computing the inverse of the inter-spike interval and, subsequently, the mean discharge rate

was calculated and considered for further analysis. Moreover, the coefficient of variation of

the inter-spike interval (CoV-ISI) was calculated by dividing the standard deviation of the

inter-spike interval by its mean.

2.4.4 Motor unit action potential amplitude

To assess the spatial distribution of motor unit action potentials (MUAP), templates

were obtained using the spike-triggered averaging technique (Stein *et al.*, 1972). Specifically,

single-differential EMGs were segmented into 100-ms windows centred on the discharge times

of individual motor units and averaged. The peak-to-peak amplitude of each template was then

calculated across channels, generating topographical MUAP amplitude maps. These maps

were used both to validate the decomposed motor units and to exemplify the type of neural

information that can be extracted with the proposed framework. As for the HDsEMG RMS

amplitude analysis, only *array 2* of the deltoid was analysed. As an additional example

analysis, the centroid of the deltoid MUAPs was calculated from active channels, defined as

those with a peak-to-peak amplitude greater than 70% of the maximum observed across the

grid (Vieira *et al.*, 2010), and compared across movements.

2.5 Statistical analysis

All statistical analyses were performed in R (version 4.2.2), using the RStudio

environment. Linear mixed-effect models (LMM) were applied for all statistical analyses, as

they account for the non-independence of data points within each participant, which is

particularly relevant for motor unit data. To compare global RMS amplitude, mean discharge

rate and CoV-ISI, random intercept models were applied with movement (abduction,

adduction, flexion and extension) and muscle (deltoid, pectoralis, trapezius and latissimus

dorsi) as the fixed effects and participant as the random effect (e.g., *mean discharge rate ~ 1*

+ movement*muscle + (1 | participant)). LMMs were implemented using the package *lmerTest*

(Kuznetsova *et al.*, 2017) with the Kenward-Roger method to approximate the degrees of

freedom and estimate the *p*-values. The *emmeans* package was used, when necessary, for

multiple comparisons and to determine estimated marginal means with 95% confidence

intervals (Lenth *et al.*, 2019).

3.0 RESULTS

3.1 HDsEMG root-mean-square amplitude

Figure 4a depicts the topographical maps of RMS amplitude at 30° shoulder angle

across muscle and movements for a selected participant. During abduction, the deltoid is the

most active muscle, particularly in the central-anterior portion of the *array 2* (see **Figure 3a**),

followed by the caudal area of the trapezius. As for the pectoralis and latissimus dorsi, during

abduction there is no visual evidence of significant activation. During adduction, most of the

activity is in the latissimus dorsi covering much of the array, followed by a small amount of

activity in the medial-caudal part of the array in the pectoralis. There is no evident activity in

the deltoid and trapezius during adduction. During flexion, most of the activity is in the deltoid

anterior region, followed by the pectoralis. During extension, the deltoid was again the most

active, but towards the posterior aspect of the array. There was also significant activity in the

caudal region of the array in the latissimus dorsi and the trapezius, with no evident activity in

the pectoralis. In **Figure 4b,** the topographical RMS amplitude maps for the same selected

participants are depicted but at 65° shoulder angle. The activation patterns during abduction,

adduction, and flexion closely resembled those reported above for the 30° condition During

extension, the deltoid was very active across *array 2*. Additionally, as seen for the 30°

condition, a significant amount of activity was observed in the caudal region of the trapezius,

but with little activity in the latissimus dorsi.

Figure 4. Muscle activation patterns across shoulder movements at 30° and 65°. *RMS amplitude topographical maps for the deltoid, pectoralis major, trapezius, and latissimus dorsi during abduction, adduction, flexion and extension. These movements were performed with the shoulder positioned at 30° of lateral abduction (a) and 65° (b). The x- and y-axes represent the rows and columns of the array, with color intensity (from dark blue to dark red) indicating the level of muscle excitation. (c and d) Group results of the RMS amplitude for each movement and muscle. Horizontal traces, boxes, and whiskers represent the median value, interquartile interval, and distribution range, respectively.*

Figure 4c shows the group results for the global RMS amplitude at 30° shoulder angle. There was a statistically significant interaction between muscle and movement (LMM, F=32.911, p<0.001). For the abduction, the deltoid muscle presented significantly higher global RMS amplitude compared to all the other muscles (p<0.001 for all comparisons). Additionally, trapezius RMS amplitude was significantly greater than pectoralis and latissimus RMS amplitude (p<0.001 for both comparisons). For the adduction, no significant differences were observed across all muscles (p>0.634 for all comparisons). For the flexion, the RMS amplitude of the deltoid and pectoralis major muscles was significantly greater than the trapezius and latissimus dorsi (p<0.031 for all comparisons). Finally, for the extension, the

370 deltoid muscle presented a higher RMS amplitude compared to the pectoralis major and

371 latissimus dorsi ($p<0.041$ for both comparisons), with no other significant differences across

372 muscles ($p>0.069$).

373  Similar results were observed at 65° shoulder angle (**Figure 4d**), where there was a

374 statistically significant muscle*movement interaction (LMM, $F=33.903$, $p<0.001$). For the

375 abduction, the deltoid muscle presented greater RMS amplitude compared to all the other

376 muscles ($p<0.001$ for all comparisons). Furthermore, trapezius RMS amplitude was

377 significantly higher than pectoralis and latissimus ($p<0.001$ for both comparisons). For the

378 adduction, no significant differences were observed across muscles ($p>0.808$ for all

379 comparisons). For the flexion, the global RMS amplitude of the deltoid muscle was

380 significantly greater compared to the trapezius and latissimus dorsi ($p<0.003$ for both

381 comparisons). In addition, the pectoralis major muscle showed a higher RMS amplitude than

382 the latissimus dorsi ($p=0.004$). For the extension, the RMS amplitude of the deltoid muscle was

383 significantly greater compared to all the other muscles ($p<0.001$ for all comparisons).

384 Additionally, the trapezius RMS amplitude was greater than the pectoralis major and latissimus

385 dorsi ($p<0.004$ for both comparisons).

386 *3.2 Motor unit discharge behaviour*

387  **Figure 5** and **Figure 6** illustrate, for different shoulder movements at a 30° shoulder

388 angle, the neural features at the motor unit level that can be extracted using the proposed

389 framework. For each muscle, representative MUAP amplitude maps and the corresponding

390 motor unit instantaneous discharge rates are presented. During abduction (**Figure 5**), motor

391 units could be identified only in the deltoid and trapezius muscles for this representative

392 participant, with 30 and 19 units decomposed, respectively. In the deltoid, motor units were

393 distributed across different regions of the array, including anterior (MUAP 1 and MUAP 30),

central (MUAP 3 and MUAP 5), and posterior regions (MUAP 2 and MUAP 4). In contrast,

most trapezius motor units were located cranially within the array. Additionally, most motor

units from both muscles were tonically active throughout the plateau phase of the task.

Figure 5: Motor unit action potential amplitude maps and motor unit discharge rates during shoulder abduction at 30° Motor unit action potentials were obtained using the spike-triggered averaging technique. For this representative subject, motor units were identified in the deltoid (30 units) and trapezius (19 units). No motor units were decomposed from the pectoralis or latissimus dorsi (left panel). Abduction force profile (gray line), and motor unit instantaneous discharge rates are shown in the right panel. Separately per muscle, each motor unit is represented with a color, used both for the MUAP distribution maps and the instantaneous discharge rates.

Figure 6 shows the same representative participant during the flexion movement. In

this condition, identifiable motor units were found in the deltoid (20 units), pectoralis (21

units), and trapezius (16 units), whereas no motor units were identified in the latissimus dorsi.

Deltoid motor units were predominantly located in the anterior region of the array (e.g., MUAP

1, MUAP 2, and MUAP 5). For both the trapezius and pectoralis, motor units were distributed

across different regions of their respective arrays. While most deltoid and pectoralis motor

units were tonically active throughout the plateau phase, some trapezius units exhibited more

intermittent activity.

Table 1 summarizes the motor unit yield across all muscles, movements, and shoulder

angles. Overall, a higher number of motor units were decomposed at the 30° shoulder angle

compared with 65°. Consistent with the representative case, no motor units were identified in

the pectoralis major and latissimus dorsi during abduction, or in the latissimus dorsi during

flexion. In addition, no units were decomposed from the pectoralis during extension.

420
Figure 6: Motor unit action potential amplitude maps and motor unit discharge rates during
shoulder flexion at 30°. Motor unit action potentials were obtained using the spike-triggered
averaging technique. For this representative subject (same as in Figure 5), motor units were
decomposed from the deltoid (20 units), pectoralis (21 units), and trapezius (16 units). No
motor units were decomposed from the latissimus dorsi (left panel). Flexion force profile (gray
line), and motor unit instantaneous discharge rates are shown in the right panel. Separately
per muscle, each motor unit is represented with a color, used both for the MUAP distribution
maps and the instantaneous discharge rates.

429
430 ***Table 1:*** *Average ± standard deviation motor unit yield across all muscles (deltoid, pectoralis, trapezius and*
431 *latissimus), movements (abduction, adduction, flexion and extension) and shoulder angles (30° or 65°).*

	SHOULDER AT 30 DEGREES				SHOULDER AT 65 DEGREES			
	Abduction	**Adduction**	**Flexion**	**Extension**	**Abduction**	**Adduction**	**Flexion**	**Extension**
Deltoid	12 ± 19	9 ± 15	10 ± 11	12 ± 7	7 ± 7	2 ± 3	10 ± 9	10 ± 11

Pectoralis	0 ± 0	5 ± 3	5 ± 7	0 ± 0	0 ± 0	6 ± 6	5 ± 6	0 ± 0
Trapezius	6 ± 6	5 ± 9	5 ± 5	5 ± 4	5 ± 3	3 ± 1	5 ± 2	4 ± 2
Latissimus	0 ± 0	5 ± 5	0 ± 0	3 ± 3	0 ± 0	3 ± 4	0 ± 0	2 ± 2

Figure 7 shows the group results for motor unit mean discharge rate (top) and CoV-ISI

(bottom). For the 30° shoulder angle (**Figure 7a**), there was a statistically significant

interaction between muscle and movement (LMM, F=3.190, p=0.008). For the abduction, there

was no significant differences between the mean discharge rate of deltoid and trapezius motor

units (p=0.282). Similarly, for the adduction, no significant differences in motor unit mean

discharge rate were observed across muscles (p>0.058 for all comparisons). For the flexion,

the pectoralis major motor units presented significantly greater mean discharge rates compared

to both trapezius and deltoid motor units (p<0.044 for both comparisons). For the extension,

both the deltoid and trapezius motor units showed significantly higher discharge rates than the

latissimus dorsi motor units (p<0.005 for both comparison), with no significant differences

between them (p=0.969). Similarly, for the 65° shoulder angle (**Figure 7b**), there was a

significant muscle*movement interaction (LMM, F=6.821, p<0.001). For the abduction, no

significant differences were found in the mean discharge rate of the deltoid and trapezius motor

units (p=0.308). For the adduction, significant higher discharge rates were found for trapezius

compared to deltoid motor units (p<0.013). For the flexion, both the deltoid and pectoralis

major motor units showed higher discharge rates than the trapezius (p<0.001 for both

comparisons), while no significant differences were observed between deltoid and pectoralis

major motor units (p=0.537). For the extension, deltoid and trapezius motor units presented

significantly greater mean discharge rates compared to the latissimus dorsi (p<0.017), with no

significant differences between them (p=0.311).

When examining the variability of the motor unit discharge rate (CoV-ISI), for the 30°

shoulder angle (**Figure 7c**), there was a statistically significant interaction between muscle and

movement (LMM, F=3.037, p=0.010). For the abduction, the trapezius presented a higher

CoV-ISI compared to the deltoid muscle (p=0.009). Similarly, for the adduction, the trapezius

CoV-ISI was significantly higher than the pectoralis major and latissimus dorsi motor units

(p<0.001 for both comparisons), with no other significant differences across muscles (p>0.262

for all comparisons). For the flexion, the trapezius motor units showed significantly greater

CoV-ISI compared to both the deltoid and pectoralis major (p<0.002). For the extension, no

significant differences in CoV-ISI were observed across muscles (p>0.060). For the 65°

shoulder angle (**Figure 7d**), there was also a significantly muscle*movement interaction

(LMM, F=4.437, p<0.001). For the abduction, no significant differences in the CoV-ISI were

observed between the deltoid and trapezius motor units (p=0.215). For the adduction, the CoV-

ISI of deltoid motor units was significantly greater than the pectoralis major and latissimus

dorsi motor units (p<0.021), with no other significant differences across muscles (p>0.110).

For the flexion, the trapezius motor units presented significantly greater CoV-ISI values

compared to the pectoralis major (p<0.017). For the extension, the latissimus dorsi motor units

showed significantly greater CoV-ISI values than both the deltoid and trapezius (p<0.010).

_Figure 7: Motor unit discharge behavior across movements and elevation angle__. Group_
results for the motor unit mean discharge rate (a and b) and the CoV-ISI (c and d) during
abduction, adduction, flexion, and extension at both 30° (left) and 65° (right) shoulder angles.

Horizontal traces, boxes, and whiskers represent the median value, interquartile interval, and
distribution range, respectively; individual circles correspond to single motor units.

4.0 DISCUSSION

The goal of this methodological paper was to investigate four isometric shoulder

movements at two angles to fully assess the neural control of the main superficial shoulder

muscles (deltoid, pectoralis major, trapezius and latissimus dorsi) at both the global EMG level

(RMS amplitude) and at the motor unit level (discharge patterns). To our knowledge, this is

the first to record motor unit behaviour with non-invasive HDsEMG in the deltoid and

latissimus dorsi. The proposed framework demonstrates the feasibility of integrating global

and motor unit analyses to advance the understanding of shoulder muscle control and provides

a foundation for improving biomechanical modelling of the shoulder.

The RMS topographical maps showed distinct activation patterns across muscles and

movements. As expected, the deltoid was the primary contributor during abduction, while the

pectoralis major and latissimus dorsi were mainly active during flexion and

extension/adduction, respectively. The trapezius displayed consistent activation across

conditions, reflecting its role in stabilizing the shoulder during multi-directional force

generation. Importantly, only minor changes in spatial activation patterns were observed

between 30° and 65° of abduction, although deltoid activity was moderately greater at the

shorted muscle length (see **Figure 4**). These findings align with previous evidence that

localized muscle regions can be selectively recruited depending on shoulder movement

direction (Wickham & Brown, 1998, 2012; Brown *et al.*, 2007). They also highlight the

limitation of relying solely on RMS amplitude from bipolar EMG, which, due to waveform

cancellation, may not fully capture underlying neural activity (Day & Hulliger, 2001; Keenan

et al., 2005).

The analysis of decomposed motor unit discharge behaviour (**Figure 7**) generally

mirrored the RMS amplitude trends but provided a more detailed view of neural control. For

instance, during abduction, motor units were identified only in the deltoid and trapezius, with

similar discharge rates, supporting their dominant role in this task. Conversely, during flexion,

deltoid and pectoralis major motor units exhibited comparable mean discharge rates, whereas

trapezius motor units discharged at significantly lower rates, suggesting a more stabilizing than

prime-mover contribution. These examples illustrate how motor unit analyses may be more

sensitive than global EMG in detecting subtle differences in task-specific muscle contributions.

The spatial distribution of MUAPs further demonstrated movement-specific

recruitment. Although motor units were not tracked across tasks, differences in the sets of

active units between abduction and flexion strongly suggest task-dependent recruitment and

de-recruitment, consistent with previous findings in other upper-limb muscles (ter Haar

Romeny *et al.*, 1984). This capability of the framework to map motor unit territories and

discharge patterns (**Figure 5** and **Figure 6**) provides a novel tool for probing the neural

strategies underlying complex shoulder control.

A notable observation was the large variability in motor unit discharge rate variability

(CoV-ISI), particularly at the 65° shoulder angle. Higher discharge variability may reflect

increased task complexity and the greater need for synergistic adjustments between muscles to

maintain stable force output (Calvin & Stevens, 1968). However, further research is warranted

to clarify how discharge variability changes with contraction intensity and in multiple degrees-

of-freedom tasks.

5.0 LIMITATIONS

Despite the rigor taken to control the specific shoulder movements and position, several

limitations should be acknowledged. First, although participants were stabilized with straps

and a rigid jig, the inherent flexibility of the shoulder joint may have introduced small

variations in movement execution. While this may reduce experimental control, it also

increases ecological validity by better replicating strategies used in daily activities. Second,

only two joint angles within the mid-range were tested, limiting extrapolation to positions

closer to 0° or 180°. Third, contractions were restricted to 30% MVC, chosen to reflect

moderate force demands of daily living, but additional work is needed across lower and higher

force levels. Fourth, the sample included only one female participant, precluding sex

comparisons. Finally, only four major superficial shoulder muscles were recorded. The

inability to assess particularly the scapular stabilizers is a key omission given their likely

contribution during extension and adduction tasks.

6.0 CONCLUSIONS

The present study demonstrates that it is feasible to record and analyse motor unit

behaviour of major shoulder muscles using HDsEMG during controlled isometric tasks. The

framework combines global EMG and motor unit analyses with precise force measurements to

provide complementary insights into both muscle-level activation and neural control strategies.

These methodological advances are essential for the development of comprehensive

biomechanical models of the shoulder that incorporate not only anatomical and mechanical

properties but also neural control mechanisms.

REFERENCES

Baslo SA, Şirin NG, Orhan EK, Baslo MB & Öge AE (2024). Selective Muscle Involvement in Amyotrophic Lateral Sclerosis: Evidence Inferred from the Point of Motor Unit Firing Rates. *Noro Psikiyatr Ars* **61,** 296–305.

Brown JMM, Wickham JB, McAndrew DJ & Huang X-F (2007). Muscles within muscles: Coordination of 19 muscle segments within three shoulder muscles during isometric motor tasks. *J Electromyogr Kinesiol* **17,** 57–73.

Cabral HV, Inglis JG, Cudicio A, Cogliati M, Orizio C, Yavuz US & Negro F (2024). Muscle contractile properties directly influence shared synaptic inputs to spinal motor neurons. *The Journal of Physiology*; DOI: 10.1113/JP286078.

Calvin WH & Stevens CF (1968). Synaptic noise and other sources of randomness in motoneuron interspike intervals. *Journal of Neurophysiology* **31,** 574–587.

Chang YW, Hughes RE, Su FC, Itoi E & An KN (2000). Prediction of muscle force involved in shoulder internal rotation. *J Shoulder Elbow Surg* **9,** 188–195.

Chen M & Zhou P (2016). A Novel Framework Based on FastICA for High Density Surface EMG Decomposition. *IEEE Transactions on Neural Systems and Rehabilitation Engineering* **24,** 117–127.

Cogliati M, Cudicio A, Martinez-Valdes E, Tarperi C, Schena F, Orizio C & Negro F (2020). Half marathon induces changes in central control and peripheral properties of individual motor units in master athletes. *J Electromyogr Kinesiol* **55,** 102472.

Day SJ & Hulliger M (2001). Experimental simulation of cat electromyogram: evidence for algebraic summation of motor-unit action-potential trains. *Journal of Neurophysiology* **86,** 2144–2158.

Dickerson CR, Hughes RE & Chaffin DB (2008). Experimental evaluation of a computational shoulder musculoskeletal model. *Clin Biomech (Bristol)* **23,** 886–894.

Dideriksen JL, Holobar A & Falla D (2016). Preferential distribution of nociceptive input to motoneurons with muscle units in the cranial portion of the upper trapezius muscle. *Journal of Neurophysiology* **116,** 611–618.

Dimitrova NA & Dimitrov GV (2003). Interpretation of EMG changes with fatigue: facts, pitfalls, and fallacies. *J Electromyogr Kinesiol* **13,** 13–36.

Doherty TJ & Stashuk DW (2003). Decomposition-based quantitative electromyography: Methods and initial normative data in five muscles. *Muscle Nerve* **28,** 204–211.

Falla D & Farina D (2008*a*). Motor units in cranial and caudal regions of the upper trapezius muscle have different discharge rates during brief static contractions. *Acta Physiologica* **192,** 551–558.

Falla D & Farina D (2008*b*). Non-uniform adaptation of motor unit discharge rates during sustained static contraction of the upper trapezius muscle. *Exp Brain Res* **191,** 363–370.

Farina D, Holobar A, Merletti R & Enoka RM (2010*a*). Decoding the neural drive to muscles
from the surface electromyogram. *Clinical Neurophysiology* **121,** 1616–1623.

Farina D, Holobar A, Merletti R & Enoka RM (2010*b*). Decoding the neural drive to muscles
from the surface electromyogram. *Clinical Neurophysiology* **121,** 1616–1623.

Gerdle B, Östlund N, Grönlund C, Roeleveld K & Karlsson JS (2008). Firing rate and
conduction velocity of single motor units in the trapezius muscle in fibromyalgia
patients and healthy controls. *Journal of Electromyography and Kinesiology* **18,** 707–
716.

ter Haar Romeny BM, van der Gon JJ & Gielen CC (1984). Relation between location of a
motor unit in the human biceps brachii and its critical firing levels for different tasks.
Exp Neurol **85,** 631–650.

Hassan A, Thompson CK, Negro F, Cummings M, Powers RK, Heckman CJ, Dewald JPA &
McPherson LM (2020). Impact of parameter selection on estimates of motoneuron
excitability using paired motor unit analysis. *J Neural Eng* **17,** 016063.

Hermans V, Spaepen AJ & Wouters M (1999). Relation between differences in
electromyographic adaptations during static contractions and the muscle function.
Journal of Electromyography and Kinesiology **9,** 253–261.

Holobar A, Farina D, Gazzoni M, Merletti R & Zazula D (2009). Estimating motor unit
discharge patterns from high-density surface electromyogram. *Clin Neurophysiol* **120,**
551–562.

Holobar A & Zazula D (2007). Multichannel Blind Source Separation Using Convolution
Kernel Compensation. *IEEE Transactions on Signal Processing* **55,** 4487–4496.

Hug F, Avrillon S, Del Vecchio A, Casolo A, Ibanez J, Nuccio S, Rossato J, Holobar A &
Farina D (2021). Analysis of motor unit spike trains estimated from high-density
surface electromyography is highly reliable across operators. *J Electromyogr
Kinesiol* 102548.

Hughes RE & An K-N (1997). Monte Carlo simulation of a planar shoulder model. *Med Biol
Eng Comput* **35,** 544–548.

Hughes RE, Johnson ME, O'Driscoll SW & An K-N (1999). Age-Related Changes in
Normal Isometric Shoulder Strength. *Am J Sports Med* **27,** 651–657.

Inglis JG, Cabral HV, Cosentino C, Bonardi A & Negro F (2025). Motor unit discharge
behavior in human muscles throughout force gradation: a systematic review and meta-
analysis with meta-regression. *Journal of Applied Physiology* **138,** 1050–1065.

Karlsson S & Gerdle B (2001). Mean frequency and signal amplitude of the surface EMG of
the quadriceps muscles increase with increasing torque — a study using the
continuous wavelet transform. *Journal of Electromyography and Kinesiology* **11,**
131–140.

Keenan KG, Farina D, Maluf KS, Merletti R & Enoka RM (2005). Influence of amplitude
cancellation on the simulated surface electromyogram. *Journal of Applied Physiology*
98, 120–131.

Kirk EA, Gilmore KJ, Stashuk DW, Doherty TJ & Rice CL (2019). Human motor unit
characteristics of the superior trapezius muscle with age-related comparisons. *Journal*
of Neurophysiology **122,** 823–832.

Kuznetsova A, Brockhoff PB & Christensen RHB (2017). lmerTest Package: Tests in Linear
Mixed Effects Models. *J Stat Soft* **82,** 1–26.

Lawrence JH & De Luca CJ (1983). Myoelectric signal versus force relationship in different
human muscles. *Journal of Applied Physiology* **54,** 1653–1659.

Lenth RV, Bolker B, Buerkner P, Giné-Vázquez I, Herve M, Jung M, Love J, Miguez F,
Piaskowski J, Riebl H & Singmann H (2019). emmeans. Available at: https://cran.r-
project.org/web/packages/emmeans/index.html [Accessed September 19, 2024].

Liddell EGT & Sherrington CS (1925). Recruitment and some other features of reflex
inhibition. *Proceedings of the Royal Society of London Series B, Containing Papers*
of a Biological Character **97,** 488–518.

de Luca CJ, LeFever RS, McCue MP & Xenakis AP (1982). Behaviour of human motor units
in different muscles during linearly varying contractions. *The Journal of Physiology*
329, 113–128.

Lulic-Kuryllo T, Negro F, Jiang N & Dickerson CR (2021*a*). Standard bipolar surface EMG
estimations mischaracterize pectoralis major activity in commonly performed tasks.
Journal of Electromyography and Kinesiology **56,** 102509.

Lulic-Kuryllo T, Thompson CK, Jiang N, Negro F & Dickerson CR (2021*b*). Neural control
of the healthy pectoralis major from low-to-moderate isometric contractions. *J*
Neurophysiol **126,** 213–226.

Mark Burden A, Lewis SE & Willcox E (2014). The effect of manipulating root mean square
window length and overlap on reliability, inter-individual variability, statistical
significance and clinical relevance of electromyograms. *Manual Therapy* **19,** 595–
601.

Martinez-Valdes E, Negro F, Laine CM, Falla D, Mayer F & Farina D (2017). Tracking
motor units longitudinally across experimental sessions with high-density surface
electromyography. *J Physiol* **595,** 1479–1496.

Masuda T, Miyano H & Sadoyama T (1983). The propagation of motor unit action potential
and the location of neuromuscular junction investigated by surface electrode arrays.
Electroencephalography and Clinical Neurophysiology **55,** 594–600.

Masuda T, Miyano H & Sadoyama T (1985). A surface electrode array for detecting action
potential trains of single motor units. *Electroencephalogr Clin Neurophysiol* **60,** 435–
443.

Merletti R, Farina D & Granata A (1999). Non-invasive assessment of motor unit properties
with linear electrode arrays. *Electroencephalogr Clin Neurophysiol Suppl* **50,** 293–
300.

Negro F, Holobar A & Farina D (2009). Fluctuations in isometric muscle force can be
described by one linear projection of low-frequency components of motor unit
discharge rates. *The Journal of Physiology* **587,** 5925–5938.

Negro F, Muceli S, Castronovo AM, Holobar A & Farina D (2016). Multi-channel
intramuscular and surface EMG decomposition by convolutive blind source
separation. *J Neural Eng* **13,** 026027.

Ning Y, Zhu X, Zhu S & Zhang Y (2015). Surface EMG Decomposition Based on K-means
Clustering and Convolution Kernel Compensation. *IEEE Journal of Biomedical and*
Health Informatics **19,** 471–477.

Padovan R, Toninelli N, Longo S, Tornatore G, Esposito F, Cè E & Coratella G (2024).
High-Density Electromyography Excitation in Front vs. Back Lat Pull-Down Prime
Movers. *J Hum Kinet* **91,** 47–60.

Reucher H, Rau G & Silny J (1987). Spatial filtering of noninvasive multielectrode EMG:
Part I--Introduction to measuring technique and applications. *IEEE Trans Biomed Eng*
34, 98–105.

Sherrington CS (1925). Remarks on some aspects of reflex inhibition. *Proceedings of the*
Royal Society of London Series B, Containing Papers of a Biological Character **97,**
519–545.

Stein RB, French AS, Mannard A & Yemm R (1972). New methods for analysing motor
function in man and animals. *Brain Research* **40,** 187–192.

Vieira TM, Merletti R & Mesin L (2010). Automatic segmentation of surface EMG images:
Improving the estimation of neuromuscular activity. *J Biomech* **43,** 2149–2158.

Westgaard RH & De Luca CJ (2001). Motor control of low-threshold motor units in the
human trapezius muscle. *J Neuropathol* **85,** 1777–1781.

Wickham JB & Brown JMM (1998). Muscles within muscles: the neuromotor control of
intra-muscular segments. *Eur J Appl Physiol* **78,** 219–225.

Wickham JB & Brown JMM (2012). The function of neuromuscular compartments in human
shoulder muscles. *J Neurophysiol* **107,** 336–345.

Woods JJ & Bigland-Ritchie B (1983). Linear and non-linear surface EMG/force
relationships in human muscles. An anatomical/functional argument for the existence
of both. *Am J Phys Med* **62,** 287–299.

Yao W, Fuglevand RJ & Enoka RM (2000). Motor-Unit Synchronization Increases EMG
Amplitude and Decreases Force Steadiness of Simulated Contractions. *Journal of*
Neurophysiology **83,** 441–452.

AUTHOR CONTRIBUTIONS

- Conceptualization: JGI, SR, HVC, CRD, FN
- Data curation: JGI, SR, HVC
- Formal analysis: JGI, SR, HVC
- Funding acquisition: JGI, FN
- Investigation: JGI, SR, CC
- Methodology: JGI, SR, HVC, RP, CRD, FN
- Project administration: FN
- Writing – original draft: JGI, SR, HVC, FN
- Writing – review and editing: JGI, SR, HVC, CC, RP, CRD, FN