Heritable differences in metabolic stability underpin thermal tolerance of threespine

stickleback (*Gasterostues aculeatus)* ecotypes

Emily V. Kerns*†[1], Kristofer T. Sasser*[1], Jesse N. Weber[1]

*Co-first author

[1]University of Wisconsin-Madison, Department of Integrative Biology

†Corresponding author

Keywords: threespine stickleback, thermal tolerance, oxygen and capacity-limited thermal

tolerance hypothesis, *Schistocephalus solidus*, metabolic plasticity, immunity

Abstract

Assessing intraspecific variation in thermal tolerance can provide insights on how rising global

temperatures will differentially impact individual organisms and populations. The oxygen and

capacity limited thermal tolerance hypothesis (OCLTTH) posits that aerobic scope (AS), the

difference between standard and maximal metabolic rate, underpins other ecologically relevant

metrics of performance. Here we assess short- and long-term thermal tolerance of benthic and

limnetic ecotypes of threespine stickleback (*Gasterosteus aculeatus*) fish through common

garden measurements of heritable variation in AS and parasite immunity. We predicted that

benthic fish would display higher thermal tolerance resulting from adaptation to the relatively

large thermal fluctuations in the shallow lakes that they inhabit. Based on the OCLTTH, we also

expected that AS would decline with temperature and be a precursor for reduced condition.

Although data connecting parasite immunity and temperature are limited, we predicted oxygen

limitation would impair parasite resistance at higher temperature, particularly in the high-

immunity limnetic fish. Despite exposing fish to temperatures below their thermal maxima,

survival decreased with temperature and was most pronounced in limnetic fish at the highest

temperature. The evolved thermal resilience of benthic stickleback was associated with their AS

stability, suggesting that at least some populations of this keystone species are capable of

withstanding hotter climates. However, contrary to the OCLTTH, temperature associated

changes in AS were not associated with declines in fish condition or immunity.

Introduction

Global warming is altering species distributions (Chen et al., 2011) and driving

population declines that can ultimately lead to extinctions (Hof et al., 2011). Phenotypic

plasticity is often proposed as a mechanism for resilience to global warming (Clement et al.,

2023; Seebacher et al., 2015). However, theory regarding how plasticity may influence resilience

to thermal challenge is conflicting. It is often argued that populations that evolve in variable

environments are more likely to display adaptive plasticity, and therefore may be more robust to

rapid environmental fluctuations (Botero et al., 2015). Alternatively, canalized phenotypes can

sometimes be favored under fluctuating selection (Kawecki, 2000). If there is a physiological

tradeoff between plasticity and thermal tolerance, then plastic populations may be more

susceptible to heat stress (Bogan et al., 2024).

The oxygen- and capacity-limited thermal tolerance hypothesis (OCLTTH) posits that

thermal tolerance results from the ability to increase transport of oxygen to tissues under elevated

temperatures (Pörtner, 2001; Pörtner, 2002). The OCLTTH predicts that aerobic scope (AS), the

difference between the minimum and maximum metabolic rate of an organism (Farrell, 2016),

will increase until an optimal temperature (T_{opt}) is reached, after which AS will decline (Fig.

1A). Accordingly, the lower AS observed at extreme temperatures should result from less

oxygen being supplied than is required to maintain homeostasis (i.e. higher standard metabolic

rate (SMR)), as opposed to a change in maximum metabolic rate (MMR). Therefore, decreasing

AS should be among the first indicators of decreased performance. Because of its emphasis on

somatic maintenance, proponents of the OCLTTH highlight its ecological relevance (Pörtner et

al., 2017). However, the OCLTTH remains contentious among ecophysiologists (Jutfelt et al.,

2018). A meta-analysis of 73 studies found that almost half failed to find a thermal optima for

AS, undercutting a major pillar of the hypothesis (Lefevre, 2016). Additionally, physiological

reactions to thermal shifts are often duration dependent. Organisms display variation in their

primary (immediate), secondary (long term maintenance of homeostasis), and tertiary

(dysregulation of biological systems) responses depending on how long they experience an

altered environment (Franke et al., 2024). However, few studies have investigated the effects of

chronic thermal stress on physiological plasticity (Clark et al., 2013). Given that organismal

fitness depends on the duration of environments experienced throughout a lifetime, and that

populations evolve in different environments, to understand how plasticity influences resilience

to climate change requires testing physiological responses to stress in both temporal and

population-specific contexts.

65   While hypotheses connecting respiration and temperature have received substantial

experimental attention, less is known about the effect warming will have on fish immunity and

parasite infection dynamics (Franke et al., 2024). It is hypothesized that global warming will

exacerbate the impacts of parasitism by disrupting host immunity and facilitating parasite

transmission, infectivity, and growth (Lõhmus & Björklund, 2015). The combination of heat and

parasite infection may aggravate the detrimental effects of each stressor alone, amounting to a

synergistically negative physiological response by the host (Earhart et al., 2022; Franke et al.,

2024). Alternatively, heat may detrimentally impact fitness of parasites (Kirk et al., 2022),

hindering their ability to complete their life cycle (Claar & Wood, 2020; Barber et al., 2016).

74   Threespine stickleback (*Gasterosteus aculeatus*) provide an opportunity to test the

OCLTTH in the context of evolved differences in thermal tolerance and immune competency

across populations. Multiple studies have found heritable differences in thermal tolerance of

threespine stickleback ecotypes (Barrett et al., 2011; Dittmar et al., 2014). Marine fish generally

show less tolerance for low temperatures than freshwater populations (Barrett et al., 2011), but

there is little evidence of evolved differences in thermal max (Barrett et al., 2011; Metzger et al.,

2016; Mottola et al., 2022). Benthic ecotypes are so named because they evolve in shallow lakes

that can warm and cool rapidly. In contrast, limnetic ecotypes are adapted to large, thermally

stratified lakes (Willacker et al., 2010). Given that many benthic and limnetic stickleback

populations have lived in divergent temperature environments for thousands of years, they offer

an opportunity to test for the evolution of divergence in thermal tolerance. To our knowledge,

only one study has tested whether thermal tolerance differs among stickleback sampled from

lakes of different size and mean summer temperature (Dammark et al., 2018). This work found

only slight differences in short term thermal maximum and did not explicitly categorize fish by

ecotype.

In addition to inhabiting thermally divergent environments, benthic and limnetic

stickleback differ in diet and, consequently, parasite exposure. Limnetic ecotypes tend to

consume more plankton than benthic ecotypes, which drives higher exposure to the copepod-

transmitted cestode *Schistocephalus solidus* (Bolnick et al., 2024; Stutz et al., 2014). To

minimize the impacts of *S.solidus* infection, soon after parasite exposure many stickleback

populations evolve an intense innate immune response that results in extensive peritoneal fibrosis

(Hund et al. 2022; Weber et al., 2017b). However, severe fibrosis can be costly in terms of

energy required to mount the response (Sasser et al., in review), likely contributing to the

evolved loss of fibrosis in some populations (Weber et al., 2022). Conversely, the protective

effects of fibrosis may be particularly beneficial under warming conditions. Higher temperatures

increase both the growth rate (Franke et al., 2017; Macnab & Barber, 2012) and fecundity

(Scharsack et al., 2021) of *S. solidus*. One caveat is that some stickleback populations exhibit

immune dysregulation at hotter temperatures (Scharsack et al., 2021), even in the absence of

pathogens (Dittmar et al., 2014). Though complicated, these immune-temperature connections

should be most apparent in limnetic stickleback ecotypes, as they are expected to display both

high cestode immunity (i.e., strong fibrosis) and limited thermal resilience.

We conducted an experiment to test three primary questions: 1) Do benthic and limnetic

stickleback differ in their primary and secondary/tertiary responses to heat, 2) Are changes in AS

across temperatures a reliable predictor of fish health (i.e. OCLTTH), and 3) Do increased

temperatures impair fibrotic immune responses, thereby affecting parasite infections, aerobic

capacity, and/or fish condition? We hypothesized that benthic fish would fare better at higher

temperatures due to adaptation to fluctuating thermal environments in shallow lakes. Assuming

that stickleback conform to predictions of the OCLTTH, we expected AS to decline as

temperatures increased due to a higher SMR, leading to concomitant declines in condition

metrics (Fig. 1A). Additionally, benthic thermal tolerance may be reflected by a shallower

change in SMR with temperature. While we initially defined ecotypes based on morphological

characters, in reality populations display a continuum of traits connected to environmental

differences (Haines et al., 2022). Indeed, recent work suggests that fibrosis is best predicted by

physical lake characteristics (Bolnick et al., 2024). Therefore, we also tested whether lake

properties (i.e., depth and surface area) predicted variation in thermal tolerance and survival.

With respect to immunity, we expected that higher temperatures would lead to weakened fibrosis

responses and higher incidence of cestode infections, and that these patterns would be most

pronounced in cestode-immune limnetic fish. We also predicted that prolonged exposure to high

temperatures would enhance parasite growth. Although we were unable to evaluate the energetic

costs of fibrosis when it was induced early in the experiment, its negative effects may lead to

reduced AS and condition.

Figure 1: (A) According to the OCLTTH, SMR oxygen demand is lowest at the thermal
optimum, causing AS to peak before declining at more extreme temperatures. We predict that
limnetic fish (blue line SMR) will exhibit a more narrow thermal curve and have a lower critical
temperature than benthic fish (black line SMR). **(B)** Summary of experimental setup. Graphics
were created on BioRender.

Methods

Animal husbandry and thermal manipulations

We assayed lab-reared fish descended from 4 lakes in southcentral Alaska: Finger (lat:

61.6085, long: -149.2675), Spirit (60.5951, -150.9995), Watson (60.5365, -150.4628), and Wik

(60.7160, -151.2479). Morphological analyses assigned Watson and Finger as benthic ecotypes,

Spirit as moderately limnetic, and Wik as highly limnetic (Haines et al., 2022; Hendry et al.,

2024). Lake depth corresponds with the classical benthic-limnetic paradigm, with Spirit and Wik

being considerably deeper (21m and 24.4m, respectively) than Finger and Watson (13.4m and

4.3m, respectively), but variation in lake surface area is not perfectly associated with depth and

could also affect thermal dynamics. The surface areas (ha) of Spirit, Wik, Finger and Watson

Lakes are, respectively, 123.2, 74.1, 135.7, and 22.8.

During the experiment, fish (n = 32 Wik, n = 72 Spirit, n = 34 Finger, n = 73 Watson)

were placed into mixed-family, population-specific tanks. Each population was represented by 3-

4 families that were spread evenly across temperature treatments. We acclimated the fish to their

experimental treatments (18-26▯) by increasing water temperature 1▯ per day (Fig. 1B) to

prevent mortality associated with a rapid change in temperature (Earhart et al., 2022). The

temperature range was selected because it is below the reported thermal maxima for this species

(Barrett et al., 2011; Dammarck et al, 2018; Dittmar et al., 2014) and spans the projected mean

and maximum air temperatures in the Kenai, AK area during the hottest month of the year in

2060 under a moderate emissions scenario (RCP 6.0; https://snap.uaf.edu/tools/community-

charts). Following thermal acclimations, we maintained fish at experimental temperatures for 31-

32 days to measure primary and secondary physiological responses, as well as cestode resistance

(Fig. 1B; details below). Fish were monitored daily, and any animals showing signs of distress

(i.e., erratic swimming, immobility, loss of equilibrium, or gasping) were immediately

euthanized using an overdose of pH buffered MS-222. If a fish died or was euthanized prior to

the end of the experiment, it was dissected to assess *S. solidus* presence. We similarly euthanized

all fish at the end of the experiment and performed dissections to measure infection outcome. All

procedures were approved by the University of Wisconsin-Madison IACUC (protocol

#L006460).

Respirometry

We assessed primary physiological response to heat 1-2 days after thermal acclimation

and secondary response 31-32 day post acclimation by measuring mass specific SMR and MMR

via intermittent flow respirometry (Supplementary methods). R:FishResp (Svendsen et al., 2019)

was used to calculate AS as the difference between SMR and MMR (Clark et al. 2013).

Respirometry traits were measured in one benthic and one limnetic population at the secondary

timepoint (Supplemental Table 1).

Experimental cestode exposures

Following Weber et al. (2022), batches of cyclopoid copepods (*Acanthocyclops robustus*)

were exposed to lab-reared *S. solidus* coracidea descended from individuals caught in Walby

Lake, Alaska (61.6178, -149.2138). We measured infection rates by screening a subset of each

batch two weeks post exposure. After initial respirometry assays, we transferred individual fish

into temperature-controlled containers and exposed them to infected copepods containing a total

of ~7 cestodes. At the time of exposure, the fish had been fasted for 48 hours. To guarantee the

opportunity to observe uninfected fish, 86% of the fish were exposed to parasites. We could not

distinguish exposed from unexposed individuals in the final analysis.

Health indices, infection outcome, and fibrosis

Spleenosomatic (SSI - a metric of immune condition), hepatosomatic (HSI - a proxy for

energy storage), and gonadosomatic (GSI - for males only, informative for reproductive

investment) index were calculated by dividing the log transformed organ mass by the log

transformed wet mass of the fish (Zapata et al., 2006). Fulton's body condition was calculated

using $K = (Mass/SL^3) \times 100$, where K = body condition. During dissections we screened for

cestode infection and scored fibrosis (Hund et al., 2022). Ordinal scores were 0 for no visible

fibrosis, 1 if organs were attached to the swim bladder, 2 if organs were fused together, 3 if

organs were fused to the side of the body cavity, and 4 if the body cavity was entirely filled with

fibrotic tissue.

Statistical Analysis

We used generalized linear models to test whether primary and secondary (i.e., start and

end of experiment) respirometry traits (MMR, SMR, and AS) responded to temperature, ecotype,

and temperature*ecotype interactions. Although models for secondary traits were limited to data

from two fish populations (Supplemental Table 1), we tested for additive effects of fibrosis

presence and score on respirometry. Stickleback can only be definitively sexed by observing the

gonads during dissection, so sex effects were only included for secondary metabolic and fibrosis

data.

We fit linear mixed effect models (R package nlme v. 31-164; Bates et al., 2015) to test

whether condition metrics (body condition, SSI, GSI, HSI) were predicted by AS, ecotype, and

their interaction. Sex was included as a fixed effect and tank density was a random effect. In the

absence of evidence to support the OCLTTH (i.e., no effect of AS), temperature could still

influence condition metrics through other biological pathways. Therefore, we again used linear

mixed effect models to directly test the effects of temperature, ecotype, temperature*ecotype,

and sex on each condition phenotype, with tank density as a random effect. Cox regression was

used to assess whether survival varied between ecotypes, across temperature treatments, or

between ecotypes within a temperature treatment. Additional details regarding animal husbandry,

respirometry, and analysis can be found in the supplementary information.

Results

Primary Response: Ecotype and lake ecology interact with temperature to explain variation in

SMR and AS

Immediately following acclimation, MMR was best explained by a linear increase with

temperature (Fig. 3A, Supplemental Table 2 & 3). Although there was no effect of ecotype,

MMR did vary depending on lake surface area (Supplemental Table 2 & 4, Supplemental Fig.

1A). Specifically, MMR increased more with temperature in populations that originated from

lakes with a smaller surface area (temperature * surface area: t = -2.393, p = 0.0177).

In contrast to MMR, SMR was best explained as an effect of quadratic temperature and

increased significantly with temperature (Fig. 3B, Supplemental Table 2 & 3). Limnetic fish

tended to have a lower SMR than benthic fish between 18-22°C (ecotype: t = -2.213, p =

0.0281), but experienced a dramatic increase at 24°C (temperature * ecotype: t = 2.266, p =

0.0247). Mean lake depth, not surface area, best predicted temperature-related shifts in SMR

(Supplemental Table 2 & 4, Supplemental Fig. 1B). Fish from shallower lakes tended to have a

higher SMR, but their SMR did not increase with temperature as much as fish from lakes with a

higher depth (temperature*mean depth: t = -2.803, p = 0.00561).

Like SMR, the best model for AS included temperature as a quadratic term (Fig. 3C,

Supplemental Table 2 & 3). AS followed a concave association with temperature, peaking at

22°C. Limnetic individuals displayed ~8% higher AS than benthic fish on average, but limnetic

AS decreased precipitously at 26°C (temperature * ecotype: t = -2.426, p = 0.01626). Surface

area was the best lake attribute to predict AS (Supplemental Table 2 & 4, Supplemental Fig. 1C).

Populations from lakes with a larger surface area tended to have a higher AS, but a more stable

AS across temperature (temperature * surface area: t = -3.482, p = 0.000623).

Secondary & Tertiary Response: Limnetic fish display substantial mortality at high temperature

related to decreases in AS

Although this experiment was not designed to test survival under heat stress, mortality

rates significantly increased with chronic exposure to higher temperature (z = 2.091, p = 0.0365),

with the greatest effect on limnetic fish (ecotype: z = 1.949, p = 0.0512, Fig. 2). Much of this

ecotype difference was driven by the 89% mortality of limnetic fish at 26□, compared to just

11% mortality among benthic fish.

Although we only examined secondary (end of experiment) metabolic rates in two

populations at three temperatures, we found significant linear and quadratic effects of

temperature on MMR, SMR, and AS, with all three metrics reaching a minimum at 22□ (Fig.

3D-F, Supplemental Table 2 & 5). While there was not a significant effect of ecotype or sex

(Supplemental Table 5, Supplemental Fig. 2), when averaged across temperatures limnetic

individuals displayed ~11% higher MMR and ~21% higher AS than benthic fish.

Figure 2: Probability of survival as function of temperature and time for **(A)** benthic and **(B)**
limnetic stickleback. Survival probability significantly decreased at higher temperatures (z =

2.091, p = 0.0365) and there was a marginally significant effect of ecotype (z = 1.949, p =
0.0512).

252
253   **Figure 3**: Metabolic data from the start **(A-C)** and end **(D-F)** of the experiment. **(A)** MMR and
254   **(B)** SMR increased with temperature. **(C)** Limnetic fish displayed an overall higher AS, which
255   increased to a peak at 22☐ followed by a sharp decline. The initial AS of benthic fish increased
256   with temperature. After 31-32 days, **(D)** MMR, **(E)** SMR, and **(F)** AS decreased from 18☐ to
257   22☐ in both ecotypes, but then increased at 26☐ for benthic fish.

258
259   *Limited cestode infection across temperatures and ecotypes*

260   Despite exposing 178 fish to *S. solidus*, we only recovered a single infected fish that died

before the end of the experiment. We did however find that 72 fish exhibited fibrosis, which

includes individuals from all four populations. We also observed a few *S. solidus* encysted in

fibrotic tissue. Although some stickleback populations, including Finger Lake, display fibrosis

without parasite exposure (Bolnick et al., 2024), observing fibrosis in 34% of cases suggests that

exposures were successful but the parasites failed to establish infection after burrowing into the

body cavity of the fish. Contrary to our prediction, temperature had no effect on fibrosis presence

or severity (Supplemental Fig. 3E-F). However, fibrosis presence was associated with lower AS

(t = -2.392, df = 63, p = 0.0197), regardless of fibrosis score or fish ecotype (Fig. 4E-F).

Temperature and AS are poor predictors of condition

Based on the OCLTTH, we predicted that decreased AS would be a precursor to decline

in a number of metrics of somatic condition. Body condition did not vary based on AS.

However, there was a significant difference between ecotypes in the relationship between AS

and condition (AS * ecotype: t = -2.254, df = 61, p = 0.0278, Fig. 4A); condition tended to

decrease with AS in limnetic fish, but this relationship was not significant (t = -1.828, df = 61, p

= 0.0724). Males overall had a higher body condition than females regardless of temperature (t =

5.972, df = 61, p < 0.0001, Supplemental Fig. 4). We found the same ecotype difference in the

slope of AS associations across HSI (AS * ecotype: t = -2.094, df = 60, p = 0.0405, Fig. 4B), GSI

(AS * ecotype: t = -2.016, df = 28, p = 0.0535, Fig. 4C), and SSI (AS * ecotype: t = -1.947473,

df = 60, p = 0.0562, Fig. 4D). There was no relationship between temperature and condition

metrics (Supplemental Figure 3A-D).

Figure 4: Performance metrics as a function of aerobic scope for each ecotype. **(A)** Body condition, **(B)** HSI, **(C)** GSI, and **(D)** SSI decreased slightly with AS in limnetic populations. **(E)** Higher AS was associated with lower incidence of fibrosis, **(F)** but not with fibrosis severity. Significance levels: $*P < 0.05$; $P > 0.05$ (not significant)

Discussion

Thermal tolerance is used to project changes in geographic range (Sunday et al., 2012; Smith et al., 2022), population declines (Roeder et al., 2021), and identify species at risk of either local or global extinction (Turko et al., 2021). However, studies often assess tolerance in relation

to critical thermal maxima (CT_{max}), where temperature is steadily increased until the fish is

unable to maintain equilibrium. While this approach identifies an upper lethal limit, it can

obscure more ecologically relevant differences caused by prolonged exposure to less extreme

temperatures. Additionally, thermal tolerance may be affected by additional stressors that wild

individuals are likely to experience simultaneously, such as pathogen exposure. We therefore

assayed thermal tolerance across four benthic and limnetic populations of threespine stickleback

after exposing them to the common cestode *S. solidus*. Consistent with our prediction, we found

that benthic populations are more resilient to hotter temperatures. However, we found limited

support for the OCLTTH. Aerobic scope (AS) decreased at the highest temperature treatment,

but only for limnetic fish and was a poor predictor of condition. Although we observed no

successful infections, nor any impact of temperature on fibrosis, AS significantly decreased in

fibrotic fish. When combined with temperature-associated declines in AS, this suggests that

variation in immune responses warrant additional attention when estimating thermal tolerance.

Stable AS may enable benthic populations to survive at high temperatures

As noted previously, we designed our temperature challenge to fall well below the

reported CT_{max} for stickleback. The only other comparison of freshwater stickleback from

variable thermal environments suggested that there would be no differences in severe stress

experienced at 26°C (Dammark et al., 2018). Yet, we observed significantly lower survival at

higher temperatures, with the impact being most severe in the 26°C limnetic treatment (Fig. 2). It

is unlikely that wild Alaskan stickleback will be continuously exposed to this high of

temperature, and all populations showed high survival immediately after acute stress (i.e. within

a few days of acclimation). Regardless, our results highlight that stickleback populations respond

differently to prolonged temperature increases, with limnetic fish suffering potentially large and

negative consequences from even relatively small (i.e., compared to CT_{max}) but prolonged

temperature shifts. These longer term consequences were also related to ecotype-specific

metabolism differences observed immediately after acclimation. At the start of the experiment,

AS peaked in limnetic fish at 22°C and declined by 36% at the highest temperature. Another

study on long-term thermal responses used a different limnetic population from Alaska and,

though not spanning the same temperatures, also found substantial plasticity in AS (Ressel et al.,

2021). Meanwhile, initial AS remained relatively stable across all experimental temperatures for

benthic fish. It is important to note that the trajectory of change in AS with temperature was

more telling than absolute AS values. Both ecotypes had equivalent measures of AS at 24°C, but

the downward trajectory of limnetic AS likely reflects the early stages of physiological

dysregulation (i.e. tertiary response).

It is also instructive to consider whether our metabolic results might explain why benthic

populations exhibited low mortality rates even after 30+ days at high temperatures. Although our

secondary data are limited to two populations and three temperatures, they still contain several

interesting trends. As with the primary physiological responses, at the end of the experiment

benthic populations maintained a more stable AS than limnetic populations across temperatures.

The secondary benthic AS-temperature curve does shift to a convex shape, but likely the more

important relationship is the limited change between primary and secondary AS values. When

AS is averaged across temperature in the Watson Lake benthic fish, there is a 19% decrease by

the end of the experiment. Although this suggests that benthic fish suffered consequences from

chronic thermal stress, they generally maintained homeostasis while limnetic fish experienced

physiological dysregulation.

But how did the two ecotypes come to differ in their chronic responses to heat? To

differentiate between canalization in benthic populations or gain of plasticity in limnetic

populations, future research should examine thermal responses of marine stickleback as a proxy

for the ancestral state. However, other empirical examples of thermal canalization, including

adaptation to urban heat islands in anoles (Campbell-Staton et al., 2021), lead us to favor the

hypothesis that benthic populations lost maladaptive plasticity enabling them to better regulate

AS across temperatures. Thermal tolerance is also likely coregulated with other metabolically

costly phenotypes. Previous work identified morphological and physiological changes connected

to stickleback migration, with nonmigratory fish evolving lower MMR than migratory ecotypes

(Dalziel et al, 2012a; Dalziel et al., 2012b). Migration-associated ecotype differences in SMR are

also connected to heritable changes in thyroid hormone signaling (Kitano et al., 2010). Although

all lake fish in the present study are non-migratory, extending our results to a more

geographically diverse set of populations may clarify if thermal tolerance and migration are

corregulated.

Temperature did not affect infection rates or fibrosis

Temperature impacts immune responses across a wide range of teleosts (Scharsack &

Franke, 2022), including stickleback (Dittmar et al., 2014; Franke et al., 2019; Scharsack et al.,

2021). Thus, we were surprised to find no connection between temperature and infection or

fibrosis. However, fibrosis was associated with lower AS at the end of the experiment. While

fibrosis is initiated within days of cestode exposure (Hund et al., 2022), it can have long-term

effects on mobility (Matthews et al., 2023) and gene expression (Weber et al., 2022). Therefore,

lower AS may result from increased energy demands in immobile fish or metabolic costs of

fibrosis maintenance.

Although we did not want to bias our infection results by using cestodes from one of the

four host populations, using a foreign cestode (i.e., from Walby Lake) may explain the total lack

of infection in our study. Stickleback tend to be better at resisting infection from foreign

populations of *S. solidus* (Bolnick et al., 2024; Weber et al., 2017a). Future work that

incorporates *S.solidus* population genetic and host specificity data across benthic and limnetic

lakes will help address the lack of infection in this study.

Limited support for OCLTTH

In line with the OCLTTH, AS declined in limnetic populations as temperature increased

toward their maximum. However, AS peaked at 22°C in limnetic fish at the start of the

experiment, which would indicate that 22°C is the T_{opt} for limnetic populations under the

OCLTTH. This trend agrees with another study showing that AS increases up to at least 20°C

(Ressel et al., 2021). However, both the historical lake temperature and developmental

temperature of these populations is well below 22°C, and previous work in other stickleback

ecotypes suggests their T_{opt} is between 13-17°C (Barrett et al., 2011; Dittmar et al., 2014; Franke

et al., 2017, 2019). Contrary to the OCLTTH, metrics of fish health declined with AS, but this

relationship was only present in limnetic fish and was marginally significant. The lack of change

in benthic fish is clearly connected to their high AS stability, indicating the OCLTTH may have

limited relevance for stickleback. Another explanation involves differences in energetic demand

and uptake between lab and wild environments. Costs may be masked when resources are

plentiful but apparent when they are scarce, as is likely the case in natural settings (Boots, 2011).

Our laboratory fish were fed to satiation, perhaps leading to higher MMR than would be seen in

resource limited animals. Further tests of the OCLTTH will benefit from testing the impact of

resource limitation on performance across temperatures.

Limitations

Several features of our experimental design limited our ability to test connections

between temperature and metabolism. First, we were unable to individually mark the fish used in

this experiment, and therefore could not directly connect AS at the start of the experiment to

survival, growth rate, or individual-specific AS at the end of the experiment. Second, our choice

to acclimate fish to constant temperatures was practical but not ecologically realistic. The

chronic impact of high temperatures in this experiment likely exceeds what stickleback

experience in the wild. Conversely, acclimating fish to high temperatures also appears to buffer

the detrimental impacts of thermal stress compared to rapid environmental change (Earhart et al.,

2022). Rapid temperature fluctuations are expected to become more frequent, eliminating the

opportunity to acclimate (Diffenbaugh & Field, 2013). Although technically challenging, future

studies that mimic daily oscillations in temperature without using an acclimation period may

provide more accurate predictions regarding impacts of natural heat waves. Lastly, when

assaying MMR we were limited to exercising the fish for one minute in compliance with animal

care and ethics guidelines. Ideally, fish will be unresponsive to agitation of the caudal fin before

MMR is measured (Clark et al., 2013). Some, but not all, fish in this experiment reached this

state of exhaustion. Although MMR is expected to remain stable across temperatures, our

observations of increases in primary MMR could have resulted from fish not being reliably

exercised to exhaustion. Despite this limitation, all fish were exercised for an equal amount of

time. Although we did not specifically record the presence of exhaustion, there was no obvious

difference between fish challenged at the start and end of the experiment. It is also noteworthy

that our 20°C MMR values were higher than those reported from wild stickleback assayed at the

same temperature and collected from the same region of south central Alaska (Ressel et al.,

2021). Furthermore, MMR is notoriously difficult to measure using standardized methods

(Ressel et al., 2021). Therefore, while our MMR estimates may not be perfectly comparable with

other studies, they were still comparable between treatment groups.

Conclusions

In contrast to previous work on short term thermal tolerance, this study demonstrates that

threespine stickleback have evolved large differences in long term thermal tolerance. These

differences are closely connected to ecotype variation in thermal habitats, suggesting that they

may result from local adaptation. This work also provides a valuable test of the OCLTTH and

suggests that AS stability, potentially a result of evolved SMR canalization, underpins resilience

to chronic heat in benthic stickleback. Future studies should aim to identify the mechanistic basis

of AS stability and whether benthic and limnetic ecotypes commonly evolve differences in

chronic thermal tolerance.

Data Accessibility

All data and analysis scripts can be found at https://github.com/EmilyKerns/ThermTol and

https://doi.org/10.5061/dryad.f1vhhmh8m.

Permits/Acknowledgments/Funding

All experiments followed the National Research Council's Guide for the Care and Use of

Laboratory Animals and were approved by the Institutional Animal Care and Use Committees of

the University of Wisconsin – Madison (Protocol #L006460). B. Solin, M. Chotlos, J. Abels, M.

Reynolds, C. Wolf, and J. Benavente assisted with animal care and dissections. KTS, EVK, and

JNW were supported by the NIGMS of the NIH (#1R35GM142891-01 to JNW) and lab startup

funding from the University of Wisconsin-Madison (to JNW). EVK was supported by the UW-

Department of Integrative Biology Graduate Summer Research Award.

Author Contributions:

440   All authors contributed to the design of the study and writing of the manuscript. KTS and EVK

441   led data collection and analysis. JNW assisted in analysis and supervised and funded the work.

Conflict of Interest:

444   The authors do not have any conflicts of interest.

References:

Barber I, Berkhout BW, Ismail Z (2016) Thermal Change and the Dynamics of Multi-Host
Parasite Life Cycles in Aquatic Ecosystems. *Integr Comp Biol* 56:561–572.
https://doi.org/10.1093/icb/icw025
Barrett, R. D. H., Paccard, A., Healy, T. M., Bergek, S., Schulte, P. M., Schluter, D., & Rogers,
S. M. (2011). Rapid evolution of cold tolerance in stickleback. *Proc R Soc B*, *278*(1703),
233–238. https://doi.org/10.1098/rspb.2010.0923
Bates D, Maechler M, Bolker B, Walker S. (2015). Fitting Linear Mixed-Effects Models Using
lme4. *Journal of Statistical Software*, 67(1), 1-48. doi:10.18637/jss.v067.i01.
Bogan, S. N., Porat, O. I., Meneses, M. J., & Hofmann, G. E. (2024). Thermal plasticity has
higher fitness costs among thermally tolerant genotypes of *Tigriopus californicus*. *Funct
Ecol*, *38*(7), 1562–1577. https://doi.org/10.1111/1365-2435.14568
Bolnick, D.I., Arruda, S., Polania, C., Simonse, L., Padhiar, A., Rodgers, M.L., Roth-Monzón,
A.J. (2024). The Dominance of Coinfecting Parasites' Indirect Genetic Effects on Host
Traits. *Am Nat*, *204*(5), 482-500. https://doi.org/10.1086/732256

Bolnick, D. I., Barrett, R. D. H., Choi, E., Eckert, L., Hendry, A. P., Kerns, E. V., Lind, Å. J., Milligan-McClellan, K., Peichel, C. L., Sasser, K., Thornton, A. R., Wolf, C., Steinel, N. C., & Weber, J. N. (2024). Destabilized host-parasite dynamics in newly founded populations. *BioRxiv*. https://doi.org/10.1101/2024.06.24.600494

Boots M. (2011). The Evolution of Resistance to a Parasite Is Determined by Resources. *Am Nat, 178*. 214–220. https://doi.org/10.1086/660833

Botero, C. A., Weissing, F. J., Wright, J., & Rubenstein, D. R. (2015). Evolutionary tipping points in the capacity to adapt to environmental change. *Proceedings of the National Academy of Sciences*, *112*(1), 184–189. https://doi.org/10.1073/pnas.1408589111

Chen I-C, Hill JK, Ohlemüller R, et al (2011) Rapid Range Shifts of Species Associated with High Levels of Climate Warming. *Science* 333:1024–1026. https://doi.org/10.1126/science.1206432

Campbell-Staton S.C., Velotta J.P., Winchell K.M. (2021). Selection on adaptive and maladaptive gene expression plasticity during thermal adaptation to urban heat islands. *Nat Commun, 12*(1). DOI: 10.1038/s41467-021-26334-4

Claar DC, Wood CL (2020) Pulse Heat Stress and Parasitism in a Warming World. *Trends Ecol Evol* 35:704–715. https://doi.org/10.1016/j.tree.2020.04.002

Clark, T. D., Sandblom, E., & Jutfelt, F. (2013). Aerobic scope measurements of fishes in an era of climate change: Respirometry, relevance and recommendations. *J Exp Biol*, *216*(15), 2771–2782. https://doi.org/10.1242/jeb.084251

Clement, D. T., Neylan, I. P., Roberts, N. J., Schreiber, S. J., Trimmer, P. C., & Sih, A. (2023). Evolutionary History Mediates Population Response to Rapid Environmental Change through Within-Generational and Transgenerational Plasticity. *Am Nat*, *201*(5), E90–E109. https://doi.org/10.1086/723624

Dalziel, A. C., Ou, M., & Schulte, P. M. (2012a). Mechanisms underlying parallel reductions in aerobic capacity in non-migratory threespine stickleback (*Gasterosteus aculeatus*) populations. *JEB*, 215(Pt 5), 746–759.

Dalziel, A. C., Vines, T. H., & Schulte, P. M. (2012b). Reductions in prolonged swimming capacity following freshwater colonization in multiple threespine stickleback populations: Evolution of swimming performance. *Evolution; International Journal of Organic Evolution,* 66(4), 1226–1239.

Diffenbaugh N.S., Field C.B. (2013). Changes in Ecologically Critical Terrestrial Climate Conditions. *Science, 341*(6145), 486-492. DOI: 10.1126/science.1237123

Dittmar, J., Janssen, H., Kuske, A., Kurtz, J., & Scharsack, J. P. (2014). Heat and immunity: An experimental heat wave alters immune functions in three-spined sticklebacks (*Gasterosteus aculeatus*). *J Anim Ecol*, *83*(4), 744–757. https://doi.org/10.1111/1365-2656.12175

Dammark, K. B., Ferchaud, A.-L., Hansen, M. M., & Sørensen, J. G. (2018). Heat tolerance and gene expression responses to heat stress in threespine sticklebacks from ecologically divergent environments. *Journal of Thermal Biology*, 75, 88–96.

500   Earhart, M. L., Blanchard, T. S., Harman, A. A., & Schulte, P. M. (2022). Hypoxia and High
501   Temperature as Interacting Stressors: Will Plasticity Promote Resilience of Fishes in a
502   Changing World? *Biol Bull*, 000–000. https://doi.org/10.1086/722115
503   Farrell, A.P. (2016). Pragmatic perspective on aerobic scope: peaking, plummeting, pejus and
504   apportioning. *J Fish Biol, 88*(1), 322-343.
505   https://onlinelibrary.wiley.com/doi/10.1111/jfb.12789
506   Franke, A., Beemelmanns, A., & Miest, J. J. (2024). Are fish immunocompetent enough to face
507   climate change? *Biol Lett*, *20*(2), 20230346. https://doi.org/10.1098/rsbl.2023.0346
508   Franke, F., Armitage, S. A. O., Kutzer, M. A. M., Kurtz, J., & Scharsack, J. P. (2017).
509   Environmental temperature variation influences fitness trade-offs and tolerance in a fish-
510   tapeworm association. *Parasit Vectors*, *10*(1). https://doi.org/10.1186/s13071-017-2192-7
511   Franke, F., Raifarth, N., Kurtz, J., & Scharsack, J. P. (2019). Consequences of divergent
512   temperature optima in a host–parasite system. *Oikos*, *128*(6), 869–880.
513   https://doi.org/10.1111/oik.05864
514   Haines, G. E., Moisan, L., Derry, A. M., & Hendry, A. P. (2022). Dimensionality and Modularity
515   of Adaptive Variation: Divergence in Threespine Stickleback from Diverse
516   Environments. *Am Nat*, 000–000. https://doi.org/10.1086/722483
517   Hendry, A.P., Barrett, R.D.H., Bell, A.M., Bell, M.A., Bolnick, D.I., Gotanda, K.M., Haines,
518   G.E., Lind, Å.J., Packer, M., Peichel, C.L., Peterson, C.R., Poore, H.A., Massengill, R.L.,
519   Milligan-McClellan, K., Steinel, N.C., Sanderson, S., Walsh, M.R., Weber, J.N., Derry,
520   A.M. (2024). Designing eco‐evolutionary experiments for restoration projects:
521   Opportunities and constraints revealed during stickleback introductions. *Ecol Evol, 14*(6),
522   e11503. DOI: 10.1002/ece3.11503
523   Hof C, Araújo MB, Jetz W, Rahbek C (2011) Additive threats from pathogens, climate and
524   land-use change for global amphibian diversity. *Nature* 480:516–519.
525   https://doi.org/10.1038/nature10650
526   Hund AK, Fuess LE, Kenney ML, et al (2022) Population‐level variation in parasite resistance
527   due to differences in immune initiation and rate of response. *Evol Lett* 6:162–177.
528   https://doi.org/10.1002/evl3.274
529   Jutfelt F, Norin T, Ern R, et al (2018) Oxygen- and capacity-limited thermal tolerance: blurring
530   ecology and physiology. *J Exp Biol* 221:jeb169615. https://doi.org/10.1242/jeb.169615
531   Kawecki, T. J. (2000). THE EVOLUTION OF GENETIC CANALIZATION UNDER
532   FLUCTUATING SELECTION. *Evolution*, *54*(1), 1–12. https://doi.org/10.1111/j.0014-
533   3820.2000.tb00001.x
534   Kirk D, O'Connor MI, Mordecai EA (2022) Scaling effects of temperature on parasitism from
535   individuals to populations. *J Anim Ecol* 91:2087–2102.
536   https://doi.org/10.1111/1365-2656.13786
537   Kitano, J., Lema, S. C., Luckenbach, J. A., Mori, S., Kawagishi, Y., Kusakabe, M., Swanson, P.,
538   & Peichel, C. L. (2010). Adaptive divergence in the thyroid hormone signaling pathway
539   in the stickleback radiation. *Current Biology: CB*, 20(23), 2124–2130.

Lefevre, S. (2016). Are global warming and ocean acidification conspiring against marine
ectotherms? A meta-analysis of the respiratory effects of elevated temperature, high CO_2
and their interaction. *Conserv Physiol, 4*(1), 1-31. DOI: 10.1093/conphys/cow009

Lõhmus, M., Björklund, M. (2015). Climate change: what will it do to fish-parasite interactions?
Biol. J. Linn. Soc., 116(2), 397-411. DOI: 10.1111/bij.12584

Macnab V, Barber I (2012) Some (worms) like it hot: fish parasites grow faster in warmer water,
and alter host thermal preferences. *Glob Change Biol* 18:1540–1548.
https://doi.org/10.1111/j.1365-2486.2011.02595.x

Matthews D.G., Maciejewski M.F., Wong G.A. Lauder G.V., Bolnick D.I. (2023). Locomotor
effects of a fibrosis-based immune response in stickleback fish. *JEB, 226*(23), jeb246684.
DOI: 10.1242/jeb.246684

Metzger DCH, Schulte PM (2017) Persistent and plastic effects of temperature on DNA
methylation across the genome of threespine stickleback (*Gasterosteus aculeatus*). *Proc*
R Soc B 284:20171667. https://doi.org/10.1098/rspb.2017.1667

Mottola G, López ME, Vasemägi A, et al (2022) Are you ready for the heat? Phenotypic
plasticity versus adaptation of heat tolerance in three☐spined stickleback. *Ecosphere* 13:.
https://doi.org/10.1002/ecs2.4015

Pörtner, H. O. (2001). Climate change and temperature-dependent biogeography: oxygen
limitation of thermal tolerance in animals. *The Science of Nature*, *88*(4), 137–146.

Pörtner, H. O. (2002). Climate variations and the physiological basis of temperature dependent
biogeography: systemic to molecular hierarchy of thermal tolerance in animals. *Comp.*
Biochem. Physiol., Part A Mol. Integr. Physiol., 132(4), 739–761 .

Pörtner, H.-O., Bock, C., & Mark, F. C. (2017). Oxygen- and capacity-limited thermal tolerance:
Bridging ecology and physiology. *J Exp Biol*, *220*(15), 2685–2696.
https://doi.org/10.1242/jeb.134585

R Core Team (2023). R: A Language and Environment for Statistical Computing. *R Foundation*
for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Ressel K.N., Cominassi L., Sarrimanolis J., O'Brien, K.M. (2022). Aerobic scope is not
maintained at low temperature and is associated with cardiac aerobic capacity in the
three☐spined stickleback (*Gasterosteus aculeatus*). *Journal of Fish Biology, 100*(2).
444-453. DOI: 10.1111/jfb.14955

Roeder KA, Roeder DV, Bujan J (2021) Ant Thermal Tolerance: A Review of Methods,
Hypotheses, and Sources of Variation. *Ann Entomol Soc Am* 114:459–469.
https://doi.org/10.1093/aesa/saab018

Sasser KT, De Lisle SP, Thompson KA, Stecyk JAW, Bolnick DI, Weber JN. Surprising
variation in growth and longevity is unconnected to immune fibrosis costs in threespine
stickleback. *In review.*

Scharsack, J. P., & Franke, F. (2022). Temperature effects on teleost immunity in the light of
climate change. *J Fish Biol*, *101*(4), 780–796. https://doi.org/10.1111/jfb.15163

Scharsack, J. P., Wieczorek, B., Schmidt-Drewello, A., Büscher, J., Franke, F., Moore, A.,
Branca, A., Witten, A., Stoll, M., Bornberg-Bauer, E., Wicke, S., & Kurtz, J. (2021).

Climate change facilitates a parasite's host exploitation via temperature-mediated
immunometabolic processes. *Glob Chang Biol*, *27*(1), 94–107.
https://doi.org/10.1111/gcb.15402

Seebacher, F., White, C. R., & Franklin, C. E. (2015). Physiological plasticity increases
resilience of ectothermic animals to climate change. *Nat Clim Chang*, *5*(1), 61–66.
https://doi.org/10.1038/nclimate2457

Smith SJ, Mogensen S, Barry TN, et al (2022) Evolution of thermal physiology alters the
projected range of threespine stickleback under climate change. *Molec Ecol*
31:2312–2326. https://doi.org/10.1111/mec.16396

Stutz, W.E., Lau, O.L., Bolnick, D.I. (2014). Contrasting Patterns of Phenotype-Dependent
Parasitism within and among Populations of Threespine Stickleback. *Am Nat, 183*(6),
810-825. DOI: 10.1086/676005

Sunday JM, Bates AE, Dulvy NK (2012) Thermal tolerance and the global redistribution of
animals. *Nature Clim Change* 2:686–690. https://doi.org/10.1038/nclimate1539

Svendsen MBS, Bushnell P, J. F. Steffensen (2019) AquaResp3

Turko AJ, Leclair ATA, Mandrak NE, et al (2021) Choosing source populations for conservation
reintroductions: lessons from variation in thermal tolerance among populations of the
imperilled redside dace 1. *Can J Fish Aquat Sci* 78:1347–1355.
https://doi.org/10.1139/cjfas-2020-0377

Weber, J.N., Kalbe, M., Shim, K.C., Erin, N.I., Steinel, N. C., Ma, L., Bolnick, D.I. (2017a).
Resist Globally, Infect Locally: A Transcontinental Test of Adaptation by Stickleback
and Their Tapeworm Parasite. *Am Nat, 189*(1), 43-57. https://doi.org/10.1086/689597

Weber, J. N., Steinel, N. C., Peng, F., Shim, K. C., Lohman, B. K., Fuess, L. E., Subramanian,
S., Lisle, S. P. D., & Bolnick, D. I. (2022). Evolutionary gain and loss of a pathological
immune response to parasitism. *Science*, *377*(6611), 1206–1211.
https://doi.org/10.1126/science.abo3411

Weber, J. N., Steinel, N. C., Shim, K. C., & Bolnick, D. I. (2017b). Recent evolution of extreme
cestode growth suppression by a vertebrate host. *Proc Natl Acad Sci*, *114*(25), 6575–
6580. https://doi.org/10.1073/pnas.1620095114

Willacker JJ, Von Hippel FA, Wilton PR, Walton KM (2010) Classification of threespine
stickleback along the benthic-limnetic axis: CLASSIFICATION OF THREESPINE
STICKLEBACK. *Biol J Linn Soc* 101:595–608.
https://doi.org/10.1111/j.1095-8312.2010.01531.x

Zapata, A., Diez, B., Cejalvo, T., Gutiérrez-de Frías, C., & Cortés, A. (2006). Ontogeny of the
immune system of fish. *Fish Shellfish Immunol*, *20*(2), 126–136.
https://doi.org/10.1016/j.fsi.2004.09.005