1

Non-hydrolyzable acetyllysine analogs to study protein acetylation in vitro and in cells

Simon Maria Kienle[1,3,#,*], Matthias Sigg[1,#], Tobias Schneider[2,3], Katrin Stuber[1,2,3], Jan Lehmann[1,3], Jasmin Jansen[1,3], Florian Stengel[1,3], Andreas Marx[2,3], Michael Kovermann[2,3], Martin Scheffner[1,3,*]

[1]Department of Biology, University of Konstanz, Konstanz, Germany
[2]Department of Chemistry, University of Konstanz, Konstanz, Germany
[3]Konstanz Research School Chemical Biology, University of Konstanz, Konstanz, Germany

[#]equal contribution

*Corresponding authors: simon.kienle@uni-konstanz.de , martin.scheffner@uni-konstanz.de

ABSTRACT

Lysine acetylation plays a prominent regulatory role in eukaryotic cells. Yet, determining the functional consequences of acetylation for a given protein represents a considerable challenge. For instance, lysine residues are subject to various posttranslational modifications, rendering interpretation of mutational studies difficult. The genetic code expansion technology enables site-specific incorporation of acetyllysine (AcK) into proteins, but the applicability of AcK is limited, as within cells, the acetyl group is removed by deacetylases. Here, we show that site-specific incorporation of the non-hydrolyzable AcK analog ketolysine (KeK) into ubiquitin closely resembles the structural and functional effects of AcK incorporation. Furthermore, AcK and KeK can be efficiently incorporated into the tumor suppressor p53 in cells. However, whereas AcK becomes deacetylated, KeK remains stable. Accordingly, incorporation of KeK, but not AcK, affects p53-mediated transcription. Thus, we propose that KeK is the AcK surrogate of choice for studying acetylation of a given protein in cells.

INTRODUCTION

The functions and properties of eukaryotic proteins are controlled by various posttranslational modifications (PTMs) that in most cases are reversible[1]. This allows a cell to rapidly adapt its proteome during time-regulated processes such as the cell cycle or differentiation and to respond to environmental changes. The continuous advancement and refinement of mass spectrometry (MS)-based approaches has been instrumental in identifying and cataloging different PTMs, e.g. phosphorylation, acetylation, and modification of proteins by ubiquitin (Ub) and Ub-like proteins, for individual proteins as well as on a proteome-wide scale[2,3]. Yet, while such studies provide invaluable insights into the plasticity of proteomes, how a given PTM affects the properties of a protein of interest (POI) remains unclear in many cases[4].

A general limitation in studying the effect of PTMs on the biochemical and structural properties of a POI is the generation of sufficient amounts of respectively modified proteins[5]. While in principle this can be achieved by incubation of the POI with the respective modifying

enzymes, the efficiency of such reactions is often limited and/or not position-specific (i.e.

residues in addition to, or other than the one of interest are modified), resulting in

heterogeneous populations of the POI. Moreover, evaluation or validation of the results

obtained in vitro may be even more challenging in cells. Common approaches to study the

effect of a PTM in cells include the use of POI variants, in which the respective residue has

been mutated (e.g. in case of lysine acetylation, replacement of lysine by arginine or

glutamine), and modulation of the activity of the respective modifying enzyme[6]. Such

approaches have obvious limitations. Targeting the modifying enzyme does in most, if not all

cases not only affect the modification status of the POI, but also that of other proteins.

Concerning replacement strategies, a main consequence of lysine acetylation, for instance,

is neutralization of the positive charge of the side chain of lysine[7]. To provide evidence for

the importance of a positive charge at the respective position of a POI, lysine is commonly

replaced by arginine. However, lysine is not only subject to acetylation, but also to other

PTMs including modification by methyl groups, Ub, and Ub-like proteins[8]. Thus, it is difficult,

or even not possible to distinguish if a respective effect is due to interference with acetylation

or, for example, with ubiquitination. As surrogate for acetylated lysine (acetyllysine or briefly,

AcK), glutamine is generally employed, which mimics elimination of the positive charge of the

lysine side chain. However, at least in some cases, the functional consequences of lysine

acetylation are not solely explained by charge neutralization[9-11]. Attachment of an acetyl

group results in extension of the length of the side chain, which depending on the position in

a POI, may induce structural changes, an effect that cannot be mimicked by the side chain of

glutamine (see Figure 1a), as we have for instance shown using Ub as model system.[11]

The genetic code expansion technology represents an attractive possibility to

incorporate AcK at the desired position of a POI, e.g. by the so-called amber codon

suppression (ACS) method[5,12-14]. Yet, in applications in eukaryotic cells the use of K-to-Q

variants may seem to be superior to ACS-mediated incorporation of AcK, as the latter is

technically more complex and more importantly, the acetyl group can be readily removed by

the action of deacetylases. To overcome this potential limitation, non-hydrolyzable AcK

analogs, such as trifluoroacetyllysine (TFAcK)[15] and 2-amino-8-oxononanoic acid, or briefly

ketolysine (KeK)[16], should prove helpful (Figure 1a). To address this proposition, we used

Ub[11] and the tumor suppressor p53[17-19] as model proteins in vitro and in cells, respectively.

Our results obtained with Ub in vitro clearly indicate that unlike Q, TFAcK and KeK represent

highly suitable AcK surrogates. Furthermore, we provide evidence that upon incorporation

into p53, AcK, and in part TFAcK, are efficiently deacetylated in cells, while KeK is refractory

to deacetylation. Notably, this difference in deacetylation susceptibility correlates with the

transcriptional activity of p53 variants harboring either AcK, TFAcK or KeK. In conclusion, our

data indicate that KeK is the AcK surrogate of choice for studying protein acetylation in cells.

RESULTS

Incorporation of AcK, TFAcK, or KeK has similar effects on Ub structure and function

Besides its well-established role as a covalent modifier of itself and other proteins, Ub is also

subject to PTMs including acetylation[20-22]. By employing the ACS technology, we have

recently shown that acetylation of K11 affects the structure, and in consequence functional

aspects of Ub[11]. Notably, these effects of acetylation are not caused solely by charge

neutralization, but also involve steric effects presumably imposed by the length of the side

chain (Figure 1a). Indeed, replacement of K11 by the presumed AcK surrogate Q shows

different effects on the structure of Ub, and the respective Ub variant is likewise functionally

different, i.e. it is more efficiently used by the E3 ligase HDM2 for autoubiquitination than

K11-acetylated Ub[11] (see also Figure 1b). Together with results from other groups[9,10], our

studies with Ub therefore indicate that data obtained with protein variants, in which lysine

residues were replaced by glutamine, have to be interpreted with caution, especially in the

absence of structural data.

AcK is likely subjected to deacetylation in cells. Therefore, we resorted to the AcK

analogs TFAcK[23] and KeK[16], as their side chains are similar in length to AcK (Figure 1a), but

are either less (TFAcK)[15] or not (KeK) susceptible to deacetylation (see below). TFAcK or

KeK were incorporated into Ub at position 11 (Ub 11X, where X represents the respective

amino acid) by ACS in bacteria (Supplementary Figure 1), and the respective Ub variants

were first tested for their ability to be used by HDM2 in autoubiquitination experiments

(Figure 1b). In agreement with our previous results[11], Ub 11AcK was less efficiently used by

HDM2 than unmodified Ub and Ub 11R (as judged by the levels of free Ub at 60 min), while

Ub 11Q was more efficiently used than Ub 11AcK (Figure 1b). Importantly, the efficiency of

HDM2 autoubiquitination in the presence of Ub 11KeK or Ub 11TFAcK was similar to that in

the presence of Ub 11AcK (Figure 1b).

The results obtained in HDM2 autoubiquitination assays support the notion that the

effects of K11 acetylation are mediated by a combination of charge neutralization and

structural changes induced by extension of the lysine side chain[11]. To corroborate this

assumption and to show that incorporation of AcK, TFAcK or KeK induces similar structural

changes, we performed high-resolution NMR spectroscopy using isotopically labeled variants

of Ub 11AcK, 11TFAcK, and 11KeK. For all three Ub variants, two-dimensional ^{1}H-^{15}N

heteronuclear single quantum coherence (HSQC) NMR spectra were acquired

(Supplementary Figure 2), and weighted chemical shift perturbations (CSPs) were calculated

for backbone amide resonances with reference to unmodified Ub (Figure 2). By this,

alterations in the chemical environment of corresponding amide proton and nitrogen spins as

consequence of the conformational changes induced by the respective modification at

position 11 of Ub can be monitored on a residue-by-residue basis. As expected by the length

and composition of their side chains, Ub 11AcK, 11TFAcK, and 11KeK show highly similar

CSP patterns, including characteristic perturbations at the C-terminal end of the central α-

helix of Ub[11]. In fact, only slight variations are apparent between the Ub variants in terms of

amplitude and the exact position of the affected amino acids in the primary sequence.

Taken together, the above results indicate that both TFAcK and KeK properly mimic

the effects of acetylation of K11 on Ub structure and that the actual side chain length at

position 11 is the determining factor for the acetylation-induced structural changes resulting

in the inefficient usage of the respective Ub variants by HDM2.

Ub 11AcK, Ub 11TFAcK, and Ub 11KeK share similar interactomes

To further prove that TFAcK and KeK are functionally equivalent to AcK, we performed

affinity enrichment coupled to high-resolution mass spectrometry (AE-MS) experiments with

whole cell extracts to compare the protein-protein interaction properties ("interactomes") of

Ub 11AcK/TFAcK/KeK. As controls, we used Ub 11A, Ub 11R, Ub 11Q, and unmodified Ub

(Ub 11K). In brief, we equipped all Ub variants with an N-terminal Strep tag II for

immobilization on Strep-Tactin® beads. The immobilized Ub variants or Strep beads alone

were incubated with whole cell extracts derived from HEK293T cells, and affinity-enriched

proteins were identified by label-free quantitative MS. Significantly enriched proteins were

determined by ANOVA statistics (S0 = 2, FDR = 0.005) (Figure 3a).

Hierarchical clustering resulted in the identification of 338 significant interactors that

bind to one or more Ub variants (Figure 3b). Besides proteins that show no preference for

any of the Ub variants (Cluster 4), we found 23 proteins that preferentially interact with Ub

11AcK, Ub 11TFAcK, and Ub 11KeK (Cluster 1) and 13 proteins that showed a strong

decrease in binding to these Ub variants (Cluster 5). Similarly, proteins were identified that

preferentially interact with Ub 11R and Ub 11K (Cluster 2, 3), suggesting that in this case the

positive charge at position 11 is an important determinant for binding. To validate the AE-MS

results, we analyzed some of the interactions by AE followed by western blot analysis

(Supplementary Figures 3a, 3b) and by using bacterially expressed proteins (NDP52[11],

UCHL3[11]; Supplementary Figure 3c). Reassuringly, we found a high correlation between the

interaction patterns obtained by the different methods. For instance, NDP52 showed

preferred binding to Ub 11TFAcK, Ub 11KeK, and Ub 11AcK, independent of the method

used. Similarly, UCHL3 showed the strongest interaction with Ub 11K and 11R and the

weakest interaction with Ub 11Q and Ub 11A.

The above results clearly indicate that the identity of the side chain at position 11

affects the protein interaction properties of Ub and that Ub variants with similar side chains

have similar interaction properties (Figure 3b). To further support this notion, we performed a

Pearson correlation analysis, which indicates the degree of similarity or dissimilarity between

two different data sets (Figure 3c). Considering all 338 significant interactors, this revealed a

high correlation between Ub 11R and Ub 11K and between Ub 11A and Ub 11Q, which

reflects the significance of a positive charge at this position. Yet, there is no significant

correlation of Ub 11A/11Q with Ub 11AcK, Ub 11TFAcK, and Ub 11KeK, while there is a high

correlation between the latter three Ub variants in general and between Ub 11AcK und Ub

11KeK in particular (Figure 3c; see also Supplementary Figure 3d). This again indicates that

the structural change induced by elongation of the side chain of K11 by acetylation has a

major impact on the protein interaction properties of Ub and that this structural and, in

consequence, functional change is most reliably mimicked by TFAcK and KeK and not or

only partially by Q.

Taken together, the results obtained show that Ub variants harboring TFAcK, KeK or

AcK are highly similar in their propensity to interact with other proteins.

Incorporation of AcK and its non-hydrolyzable analogs into p53 in human cells

From the results obtained with Ub, we reasoned that KeK and TFAcK are suitable AcK

surrogates for studies in cells. Since Ub is a highly abundant protein in cells rendering

analysis of ectopically expressed Ub rather difficult, we switched to the tumor suppressor p53

to test this assumption. p53 is subjected to various PTMs affecting its stability and/or its

transcriptional activities resulting in different biological outcomes (e.g. cell cycle arrest,

apoptosis, senescence) (summarized in[24-27]). Most importantly for our purpose, the

acetylation status of distinct K residues of p53, including K residues in its DNA binding

domain (DBD), contributes to determining the selectivity or preference of p53 for certain

target genes and thereby the eventual transcriptional program induced by p53[17-19].

To determine whether KeK and TFAcK are suitable AcK surrogates in cells, we

generated expression constructs for human p53, in which the lysine codons at position 120

and/or 164 within the DBD were replaced by the amber stop codon TAG (p53-120 and p53-

164, respectively; note that K120 directly contacts DNA, while K164 does not). We chose

these positions, since besides their physiological relevance[17-19] (see Discussion),

incorporation of unnatural or non-canonical amino acids by ACS is rarely quantitative,

resulting in a mixture of full-length and truncated versions of the POI. The more N-terminal

the truncation occurs, the easier it is to distinguish between the full-length and the truncated

protein, e.g. by western blot analysis. Furthermore, since in our case, the truncations occur in

the DBD, the truncated p53 versions are not of functional relevance - they can neither bind to

DNA nor oligomerize with full-length p53 - and, thus, do not need to be considered in the

analysis.

The p53 expression constructs were transfected together with an expression construct

encoding the CUA-tRNA as well as an aminoacyl-tRNA synthetase for AcK (AcK-RS)[28,29] and

its analogs TFAcK and KeK (for further details, see Methods) in the absence or presence of

increasing concentrations of AcK, TFAcK or KeK. As cell line, we used H1299 cells, since

they do not express endogenous p53[30]. As shown in Figure 4, the three non-canonical amino

acids were incorporated, resulting in the expression of full-length p53, with AcK and KeK

being more efficiently used or incorporated into p53 than TFAcK (Figures 4a-c). Furthermore,

the system is not limited to the incorporation of AcK or its analogs at a single position, but

can also be used to generate p53 variants harboring two acetylated lysine residues

(Supplementary Figure 4), expanding the analytic potential of the system to multiple or at

least double acetylated p53 variants.

Incorporation of AcK, TFAcK or KeK has different effects on the transcriptional

transactivation properties of p53

To test the functionality of the p53-120 variants, we determined their ability to transactivate

the transcription of a reporter construct containing the binding site or response element (RE)

for p53 of the *p21* gene[31,32]. As control, we used expression constructs encoding wild-type

(wt) p53 and the tumor-derived mutant p53-273H (substitution of R273 by H), which has lost

the sequence-specific DNA binding properties (ref[33], and references therein). While the

120AcK and 120TFAcK variants were capable of inducing expression of the reporter

construct, though with somewhat reduced efficiency as compared to wt p53, the 120KeK

variant was not, or only poorly active (Figure 5a). Moreover, the differences in the

transactivational potential of the different p53 variants were also reflected in the levels of

endogenous p21, indicating that the differences are not due to an artifact of the p21 reporter

construct (Figure 5b). Similar results were obtained with the p53-164 variants

(Supplementary Figures 5a, 5c).

At first glance, the finding that the KeK-containing p53 variants are significantly less

active than the AcK and TFAcK variants seems to be surprising. However, it may be

explained by the notion that in contrast to KeK, AcK and TFAcK are deacetylated in cells

(ref[15], and references therein) and, thus, the respective variants regain wt-like activity. To

prove this possibility, we purified the different p53 variants from transfected cells and

subjected them to Parallel Reaction Monitoring (PRM)-MS analysis[34]. The data obtained

indeed suggest that the AcK and TFAcK variants are deacetylated, though to different

extents, while the KeK variant is refractory to deacetylation as expected (Figure 5c;

Supplementary Figure 5d). The "partial resistance" of TFAcK is most likely due to the notion

that Zn-dependent deacetylases can remove the fluorinated acetyl group[15,23], while NAD-

dependent deacetylases cannot do so (see Discussion).

To exclude that the data obtained with KeK are the result of unwanted side effects, we

performed several control experiments. Firstly, we looked into the possibility that the mere

presence of KeK in the cell culture medium interferes with the transactivation activity of wt

p53. Yet, a negative effect was not observed (Supplementary Figure 5e). Secondly, p53-

101KeK was similarly active as wt p53, p53-101AcK, and p53-101TFAcK, showing that

incorporation of KeK per se does not interfere with p53 function (Supplementary Figures 5b,

5c). Thirdly, we previously reported that in vitro, p53-120KeK binds to the p21 response

element (p21-RE) with an efficiency similar to unmodified p53[35], indicating that incorporation

of KeK does not disrupt the conformational integrity of the DBD of p53.

Finally, we asked whether or not Q can serve as substitute for AcK, when studying p53

function(s) in cells. This showed that unlike p53-120KeK, p53-120Q behaved like wt p53 in

the p21 reporter assay (Figure 5d), suggesting that in case of K120 acetylation, Q is not a

suitable surrogate for AcK. In conclusion, our results obtained with Ub and p53 indicate that

KeK is the most suitable AcK analog to study the effect of lysine acetylation of a POI in vitro

and in cells.

DISCUSSION

Lysine acetylation is a prevalent PTM affecting many intracellular proteins[7,36-39]. Yet,

determining the functional consequences as well as the physiological importance of

acetylation for a POI remains a serious challenge. An obvious consequence of acetylation is

neutralization of the positive charge of lysine's side chain, the functional significance of which

can be determined by mutational analysis by replacing the respective lysine residue by

arginine or an uncharged residue such as glutamine or alanine. An alternative, but not

mutually exclusive possibility is that the eventual effect of acetylation is mediated by the

extension of the length of lysine's side chain and/or the actual presence of the acetyl group.

While this cannot be adequately addressed by conventional mutational analysis, the genetic

code expansion technology in general and ACS in particular provide excellent tools to do so.

Our comparative analysis of AcK and its analogs TFAcK and KeK and their effects on distinct

properties of Ub and p53 strongly supports this proposition.

In support of the notion that the effects of acetylation are not necessarily explained by

simple charge neutralization, we recently reported that acetylation of Ub at K11 results in

structural and, in consequence, functional changes that cannot be mimicked by substitution

of K11 by Q[11]. Here, we extended the functional comparison of Ub 11AcK and Ub 11Q by

determining their respective interactomes in vitro (Figure 3). By comparison with unmodified

Ub, Ub 11R, and Ub 11A, this clearly showed that even though the charge status at position

11 affects the Ub interactome (Figure 3c), acetylation of K11 has additional or different

effects on the interaction properties of Ub. Moreover, we show that incorporation of TFAcK

and KeK at position 11 affects the structure of Ub in a manner similar, but not identical to

AcK incorporation (Figure 2). These slight differences, together with intrinsic variations in

proteome-wide affinity enrichments, are reflected in highly similar, but not identical

interactomes of Ub 11AcK, Ub 11TFAcK, and Ub 11KeK (Figure 3, Supplemental Figure 3).

To determine the applicability of ACS-mediated incorporation of AcK and its analogs in

eukaryotic cells, we switched to p53, since acetylation of distinct lysine residues in its DBD

affects the transcriptional properties of p53[17-19] and since p53 activity can be readily

monitored in cells by transcriptional reporter assays. Our results show that AcK, TFAcK, and

KeK can be efficiently incorporated into p53 at different positions, with incorporation of

TFAcK being somewhat less efficient than those of the others for unknown reasons (note

that a similar observation was made for the efficiency of AcK and TFAcK incorporation by

others[15]). However, only p53-120KeK was resistant to the action of deacetylases in cells,

while p53-120AcK was quantitatively deacetylated and p53-120TFAcK was partially

deacetylated (Figure 5c). The "partial resistance" of TFAcK is likely due to the notion that Zn-

dependent deacetylases can remove the fluorinated acetyl group, while NAD-dependent

deacetylases cannot[15,23,40]. From a chemical perspective, TFAcK is more susceptible to

nucleophilic attacks than AcK, as the trifluoroacetyl group is a better leaving group than the

acetyl group[41]. However, while Zn-dependent deacetylases such as KDACs release the

acetyl group via nucleophilic attack of the C atom of the carbonyl group by an activated water

molecule, NAD-dependent deacetylases including sirtuins use NAD as electrophile for

nucleophilic attack of the carbonyl oxygen atom[40]. Thus, due to the electron withdrawing

effect of the three fluorine atoms, the nucleophilicity of the carbonyl oxygen is significantly

decreased[41]. Along this line, it has recently been shown that HDAC2 is able to deacetylate

FOXO4 variants harboring TFAcK or AcK at position 189, while Sirt2 can only deacetylate

the AcK variant[15]. In contrast to TFAcK, KeK is resistant against the attack of Zn-dependent

deacetylases as well[16,40]. An intermediate step in HDAC-catalyzed deacetylation is the

nucleophilic attack of the carbonyl C atom by a water molecule. To do so, a base,

presumably a histidine residue, accepts a proton from the water molecule and transfers it to

the nitrogen atom of the amide bond, which facilitates the final cleavage of the amide bond[40].

In contrast to the nitrogen, the carbon atom of KeK (Figure 1a) can most probably not

function as an acceptor for the proton. Thus, while in the case of KeK the nucleophilic attack

may still occur, though with reduced efficiency, cleavage of the C-C bond is highly unlikely. In

any case, this difference in the deacetylation susceptibility of AcK, TFAcK, and KeK may

represent an attractive possibility to dissect the contribution of the different deacetylase

families to the deacetylation of a POI.

The difference in acetylation status of the different p53-120 variants is also reflected in

different abilities to transactivate a p21-reporter construct as well as to induce expression of

endogenous p21 (Figure 5). p53-120AcK is similarly active as wt p53, p53-120TFAcK is

somewhat less active, and p53-120KeK does neither detectably transactivate the reporter

construct nor induce endogenous p21 expression. In addition and in line with published

data[42], p53-120R (substitution of K by R) efficiently transactivated the reporter construct.

Moreover, the 120Q variant behaved like wt p53 in the reporter assay, suggesting that in this

case, charge neutralization does not suffice to mimic the effect of acetylation. Our data with

p53-120KeK are in line with ChIP data indicating that within cells, p53 acetylated at K120

does not accumulate at the *p21* gene[43] and with the notion that acetylation of K120 does not

result in the induction of expression of the *p21* gene (ref[18,19], and references therein). Yet,

similar to our data with p53-120KeK[35], it has previously been shown that in vitro, p53-120AcK

generated by the ACS method binds to the p21-RE with an affinity similar to unmodified

p53[32]. How can these apparent contradictory data be reconciled? In a simplified view, cell

cycle arrest target genes such as *p21* contain "high affinity" REs for p53, while pro-apoptotic

target genes contain "low affinity" REs (ref[42], and references therein). Furthermore, there is

data to indicate that p53 contacts high affinity REs and low affinity REs in a distinct manner

and that binding of p53-120AcK does not affect the conformation of high affinity REs, but

changes the conformation of low affinity REs[32,42,44]. Moreover, there is evidence that the

conformation of p53-120AcK bound to high affinity REs is different to the one bound to low

affinity REs[32]. Such conformational changes that depend on both p53 acetylation and binding

of p53 to DNA in a sequence-specific manner may affect the protein-protein interaction

properties of p53 and thereby its ability to differentially transactivate p53 target genes.

Whereas K120 acetylation has been associated with the "apoptotic program" of p53 by

inducing expression of pro-apoptotic genes such as *BAX* and *PUMA*, K164 acetylation has

been involved in the "cell cycle program" by inducing the expression of cell cycle arrest

genes including *p21*[17-19,43,45]. Yet, p53-120KeK or p53-164KeK were inactive in our reporter

assays, irrespective of whether a p21 reporter construct or a PUMA reporter construct was

used (Figure 5, Supplementary Figures 5 and 6). As discussed above, all of our data indicate

that KeK structurally and functionally resembles AcK and, thus, its incorporation should affect

p53 activity in a manner similar to lysine acetylation. Yet, we cannot exclude that proteins

exist that bind to KeK with lower affinity than to AcK. For instance, bromodomains are known

binders of AcK-containing proteins[46]. Thus, there may be bromodomain-containing proteins

or other proteins that specifically bind to p53-120AcK, and such interactions may be critically

involved in p53-mediated transactivation of the p21 and/or PUMA reporter constructs.

However, this does not appear to be the case, since p53-120R efficiently induced expression

of both the p21 reporter construct and the PUMA reporter construct. Irrespectively, a likely

conclusion of the data obtained with p53-120KeK (p53-TFAcK), p53-120Q, and p53-120R is

that a positive charge at position 120 is not critical for p53-mediated transcription of the p21

reporter construct and, thus, the negative effect of K120 acetylation cannot be explained by

charge neutralization. For the PUMA reporter construct, charge neutralization at position 120

contributes to the negative effect of KeK incorporation, but additional properties of KeK (and

thus, AcK) appear to be involved. Taken together, our results obtained with the different p53

variants indicate that a comparative analysis of POI variants, in which a distinct K residue is

replaced by R, Q, or AcK surrogates, provides critical information about if and how

acetylation affects known properties/functions of a POI and that KeK represents the AcK

mimic of choice for such studies.

A possibility to explain the apparent discrepancy between our data and published data

is that our reporter constructs miss additional sequence elements of the respective

endogenous genes. At least in case of p21, this seems unlikely, since endogenous p21

expression was also not induced by p53-164KeK (Supplementary Figure 5c). Furthermore,

many of the published data have been obtained by mutational analysis, for instance by

replacing respective lysine residues by arginine (summarized in refs[16-18]). Thus, some of the

effects ascribed to lysine acetylation may in part be due to other lysine modifications such as

mono-ubiquitination[47,48] or lactylation[49]. Other possibilities would be that mono-acetylation of

p53 or acetylation in general is not sufficient for the induction of certain target genes. In other

words, it seems likely that a combination of acetylation with additional PTMs such as

phosphorylation is required to observe the respective transcriptional effects, a scenario that

may not be achieved in our experimental set-up. For instance, it could be envisioned that

such phosphorylation events occur only under certain stress conditions. An alternative, but

not mutual exclusive possibility is that to be functional, such PTMs have to occur in a timely

ordered, i.e. consecutive manner. Along this line, prior lysine acetylation, in particular

incorporation of an AcK mimic that cannot be hydrolyzed as in our case, may prevent

phosphorylation of p53 at residues required for transcriptional transactivation. To address

such possibilities, the continuous development and refinement of the genetic code expansion

technology to allow the combinatorial incorporation of different non-canonical amino acids

resembling non-reversible, yet faithful analogs of respective PTMs should prove highly

valuable.

METHODS

Synthesis of 2-amino-8-oxononanoic acid (Ketolysine; KeK)

KeK was synthesized as described previously[16], except for the final purification step. Instead

of using a Dowex column, KeK was purified twice via MPLC and then lyophilized. For MPLC

purification, a PrepChrom C-700 (Büchi) with SuperVarioFlash® D26 RP18 30 g or D40

RP18 90 g (Götec-Labortechnik GmbH) were used. After sample application, the column was

washed with H_2O for 20 min and then a linear gradient (0%-100% acetonitrile in 20 min) of

acetonitrile in H_2O was applied with a flow rate of 20 ml/min. Runs were monitored by UV

detection at 254 nm. Fractions containing KeK were pooled and lyophilized.

To record NMR spectra of KeK, a Bruker Avance III 400 (400 MHz) was used. Spectra

were measured in the presence of D_2O (the deuterium resonance signal is used as internal

reference frequency for other nuclei) at room temperature. The program MestReNova 14.2

was used for spectrum evaluation. Shifts are indicated in ppm (σ-scale) and coupling

constants in Hertz (Hz).

^{1}H NMR (400 MHz, D_2O): δ [ppm] = 3.88 (t, J = 6.1 Hz, 1H, H-1), 2.62 (t, J = 7.3 Hz,

2H, H-6) 2.25 (s, 3H, CH3), 2.06-1.82 (m, 2H, H-2), 1.61 (p, J = 7.4 Hz, 2H, H-5), 1.53-1.31

(m, 4H, H-3 & H-4).

^{13}C NMR (101 MHz, D_2O): δ [ppm] = 217.56 (CO), 174.68 (CO2H), 54.65 (C-1), 43.13

(C-6), 30.20 (C-2), 29.31 (CH3), 27.88 (C-4), 24.08 (C-3), 22.98 (C-5).

Plasmids

Codon-optimized cDNA encoding Ub with a C-terminal His$_6$-tag was inserted into the pGEX-

2TK backbone replacing the GST cDNA to yield pKS Ub-His[50]. cDNAs of Ub variants, in

which the amber stop codon (TAG), a glutamine, an alanine or arginine codon replaced the

respective lysine codon, were generated via site-directed mutagenesis.

The cDNA encoding the aminoacyl-tRNA synthetase for AcK (AcK-RS) derived from

Methanomethylophilus alvus was kindly provided by Dr. K. Lang (ETH Zürich) and was

inserted into the pRSFDuet vector (Merck Millipore) for bacterial expression. Plasmids

encoding tRNA and AcK-RS derived from *Methanosarcina mazei* for incorporation of

AcK/TFAcK/KeK in H1299 cells were kindly provided by Dr. Simon Elsässer (Karolinska

Institute, Stockholm)[28,29].

cDNAs encoding wild-type p53 or the tumor-derived mutant 273H with and without C-

terminal StrepII tag was inserted into pcDNA3. cDNAs for p53 variants, in which the TAG

stop codon, a glutamine or an arginine codon replaced the respective lysine codon, were

generated via site-directed mutagenesis.

Reporter constructs encoding luciferase were generated by introducing p53-responsive

elements (p21, PUMA)[32] into the pGL4 luciferase reporter system.

Bacterial expression and protein purification

Human His$_6$-tagged UBA1[50,51], N-terminal His$_6$-tagged UCHL3[50], and C-terminal His$_6$-tagged UbcH5b[50,52] were expressed and purified as described.

Expression of the various GST fusion proteins was performed in *E. coli* BL21 (DE3) Rosetta. GST-SUMO fusion proteins of HA-tagged HDM2 and NDP52 were expressed and purified as described[11].

Bacterial generation of modified Ub and unmodified Ub

For generation of Ub site-specifically modified at position 11 (with and without an N-terminal Strep II tag), the respective pKS vector was co-transformed with pRSFDuet harboring the AcK-RS/t-RNA pair for amber codon suppression into *E. coli* B834 (DE3) cells. Cells were grown at 37 °C in 100 ml LB medium containing 100 µg/l carbenicillin/ 34 µg/l kanamycin until OD$_{600}$ = 1 was reached. Then, cultures were diluted to an OD$_{600}$ = 0.1 with minimal media devoid of phenylalanine and further incubated. At an OD$_{600}$ = 0.4, either 10 mM AcK (abcr) or 20 mM TFAcK (abcr) or 10 mM KeK (racemate) and 20 mM nicotinamide (Sigma-Aldrich) were added. Gene expression was induced at OD$_{600}$ = 0.7 by the addition of 1 mM IPTG (final conc.). After incubation at 25 °C for 16 h, cells were harvested by centrifugation, resuspended in lysis buffer (50 mM Tris-HCl pH 7.4, 150 mM NaCl, 30 mM imidazole, 1 % Triton X-100, 1 mg/ml aprotinin/leupeptin, 1 mg/ml Pefabloc) and lysed by sonication. The lysate was cleared, and modified Ub-His was purified using a 5 ml HisTrap HP (Äkta pure25, GE Healthcare) with a step from 30 mM to 242 mM imidazole in 50 mM Tris-HCl pH 7.4, 150 mM NaCl for elution. Protein-containing fractions were pooled, UCHL3 was added to a final concentration of 75 µg/ml to remove the C-terminal His-tag, and the samples were incubated for 4 h. After cleavage, modified Ub variants were dialyzed against 25 mM acetate pH 4.5 overnight at 4 °C. Ub samples were further purified via a 1 ml HiTrap SP HP (GE Healthcare) with a linear gradient from 25 mM acetate pH 4.5 to 25 mM acetate 4.5, 1 M NaCl. Modified Ub containing fractions were pooled and dialyzed against 25 mM Tris-HCl pH 7.5, 50 mM NaCl. The concentration was determined with the BCA Protein Assay Kit (Thermo) and

commercial Ub (Sigma-Aldrich) as standard. The correct masses of the modified Ub variants were confirmed by ESI-MS (Supplementary Figure 1).

Expression and purification of unmodified Ub and the Ub variants containing Q, A or R instead of K at position 11 were performed similar to the modified Ub variants, except that AcK/TFAcK/KEK and nicotinamide were not added to the respective expression cultures.

In vitro autoubiquitination assay

For _in vitro_ autoubiquitination, 100 nM UBA1, 0.5 µM UbcH5b, and 0.55 µM HDM2 were incubated in 25 mM Tris-HCl pH 7.5, 50 mM NaCl, 1 mM DTT, 2 mM ATP, and 2 mM $MgCl_2$. The reaction was started by addition of 7.8 µM (final conc.) of the respective Ub variant and incubated at 37 °C for the times indicated. Reactions were stopped by the addition of 5x reducing SDS loading buffer. Entire reaction mixtures were electrophoresed in 12.5% SDS-polyacrylamide (SDS-PA) gels followed by Coomassie blue staining.

Generation of isotopically labeled Ub for NMR analysis

For preparation of ^{15}N-labeled proteins, _E. coli_ harboring the expression construct for Ub with a TAG stop codon at position 11 was grown in LB medium at 37 °C until OD_{600} = 1. This culture was used to inoculate ^{15}N M9 medium (33.7 mM Na_2HPO_4, 22 mM KH_2PO_4, 8.55 mM NaCl, 9.35 mM $^{15}NH_4Cl$, 1 mM $MgSO_4$, 0.3 mM $CaCl_2$, 134 µM EDTA, 31 µM $FeCl_3$, 6.2 mM $ZnCl_2$, 0.76 µM $CuCl_2$, 0.42 µM $CoCl_2$, 1.62 µM H_3BO_3, 0.081 µM $MnCl_2$, 1 mg/l biotin, 1 mg/l thiamine, 0.4% glucose (w/v)). Upon growth overnight at 37 °C, 100 ml of the culture were diluted with 900 ml of pre-warmed ^{15}N M9 medium. Cells were grown at 37 °C until OD_{600} = 0.4 was reached, and then either 10 mM AcK (abcr), 20 mM TFAcK (abcr) or 10 mM KeK (racemate) and 20 mM nicotinamide (Sigma-Aldrich) (final conc.) were added. Gene expression was induced at OD_{600} = 0.7 by the addition of 1 mM IPTG (final conc.). After incubation at 25 °C for 16 h, cells were harvested by centrifugation. Purification of isotopically labeled Ub was performed as described above for unlabeled Ub.

Acquisition of NMR spectroscopic data

NMR experiments were performed at T = 298 K on a Bruker Avance Neo 800 MHz

spectrometer equipped with a quadruple (QCI) cryo probe with z-gradient. Datasets were

processed using NMRPipe[53] and analyzed by NMRView[54]. Samples were prepared in 20 mM

NaH_2PO_4 (pH 6.8) supplemented with 5% (v/v) D_2O, which serves as frequency lock. All

measurements were performed with singly ^{15}N-labeled variants of Ub 11AcK, 11TFAcK and

11KeK using protein concentrations of 33 µM, 500 µM and 60 µM, respectively. For each

protein, a two-dimensional heteronuclear ^1H-^{15}N HSQC spectrum was collected with 1024

and 256 increments in the direct ^1H and indirect ^{15}N dimension, respectively, using 8-32

transients depending on the protein concentration. WATERGATE pulse sequence and

presaturation were applied for efficient solvent suppression.

Backbone resonance assignment and calculation of chemical shift perturbations

The backbone resonance assignment of Ub 11AcK was transferred from ref[11] and served as

basis for the assignment of backbone amide resonances of Ub 11TFAcK and 11KeK. Due to

the slight differences in peak position, this could be successfully accomplished by visual

estimate when the two-dimensional ^1H-^{15}N HSQC NMR spectra of Ub 11TFAcK and 11KeK

were directly compared to Ub 11AcK.

　　　Weighted chemical shift perturbations between the different Ub variants and

unmodified Ub were calculated according to the following equation[55]:

$$\Delta\omega = \sqrt{\frac{(\Delta^1H)^2 + \frac{1}{25}(\Delta^{15}N)^2}{2}}, \tag{1}$$

where Δ^1H is the peak shift in proton and $\Delta^{15}N$ is the peak shift in nitrogen dimension,

respectively, between two corresponding peaks. Note that the assignment of backbone

resonances arising in the two-dimensional heteronuclear ^1H-^{15}N HSQC NMR spectrum

related to unmodified Ub was also taken from ref[11].

Cell culture and extract preparation

HEK293T and H1299 cells were cultured in Dulbecco's Modified Eagle's Medium (DMEM, Thermo Fisher) supplemented with 10% fetal bovine serum (FBS, Sigma) at 37 °C and 5% CO_2 atmosphere.

For affinity enrichment experiments, HEK293T and H1299 cell pellets were resuspended in cell lysis buffer (1x PBS, 2 mM $MgCl_2$, 1 mM DTT, supplemented with 1 mg/ml aprotinin/leupeptin, 1 mg/ml Pefabloc) and subsequently lysed by sonication. After centrifugation (21,000x g, 30 min, 4 °C), the total protein concentration of the supernatant was determined with the BCA Protein Assay Kit (Thermo Fisher) and freshly dissolved BSA as standard.

For transactivation studies, H1299 cell pellets were resuspended in lysis buffer (100 mM Tris-HCl pH 8.0, 100 mM NaCl, 1 mM DTT, 1% (v/v) Nonidet P-40, supplemented with 1 mg/ml aprotinin/leupeptin, 1 mg/ml Pefabloc) and incubated for 30 min at 4°C prior to centrifugation (16,100x g, 30 min, 4°C). Cleared whole cell extracts were analyzed by western blot and luciferase reporter assay as described below.

Affinity enrichment

For affinity enrichment of Ub interacting proteins, 25 μg of each Strep-Ub variant was incubated with 2.5 mg whole cell lysate for 10 min on ice. After washing 50 μl Strep-Tactin® beads (iba) with three times 200 μl cell lysis buffer, the mixture was added to the beads. Incubation at room temperature for 3 h in an overhead shaker was followed by 5 washing steps with 200 μl cell lysis buffer each. Bound proteins were eluted by incubating the beads for 5 min on ice with two times 150 μl and one time 100 μl elution buffer (100 mM Tris-HCl pH 8.0, 150 mM NaCl, 1 mM EDTA, 2.5 mM desthiobiotin). The eluates were combined and freeze-dried. Each experiment was performed in quadruplicates.

The freeze-dried eluate was resuspended in 100 μl 8 M urea, reduced with 5 mM TCEP for 30 min at 37 °C and alkylated with 10 mM iodoacetamide for 30 min at room temperature. After dilution to 1 M urea with 50 mM NH_4HCO_3, 5 μg trypsin were added, and

proteins were digested for 20 h at 37 °C. The digested samples were freeze-dried and kept

at -20 °C until they were prepared for MS measurement.

Prior to MS measurement, the tryptic peptides were dissolved in 50 µl 0.1% TFA in

H_2O, acidified with 10% TFA and desalted using Pierce™ C18 spin tips (Thermo Fisher).

For affinity enrichment of ectopically expressed Strep-tagged p53, transiently

transfected cells were harvested and lysed as described above. Whole cell extracts were

added to 20 µL Strep-Tactin® XT beads (iba) pre-equilibrated in lysis buffer. After incubation

at 4°C for 3 h in an overhead shaker, beads were washed 5 times with 200 µl lysis buffer

each. Bound proteins were eluted as stated above for Ub with elution buffer (100 mM Tris-

HCl, pH 8.0, 150 mM NaCl, 1 mM EDTA, 50 mM biotin) and lyophilized. Freeze dried eluates

were resuspended in SDS loading buffer and subjected to SDS-PA gel electrophoresis for in-

gel digestion. SDS-PA gels were stained by colloidal Coomassie blue, and gel pieces

containing p53 were excised and digested according to ref[56]. In brief, gel pieces were

washed in 100 mM NH_4HCO_3 for 10 min, destained in 50 mM NH_4HCO_3, 50% (v/v)

acetonitrile and subsequently dehydrated in pure acetonitrile. Disulfide bonds were reduced

by addition of 10 mM TCEP in 100 mM NH_4HCO_3 for 60 min at 56 °C followed by

dehydration in pure acetonitrile and subsequent alkylation in 50 mM iodoacetamide in 100

mM NH_4HCO_3 for 60 min at room temperature in the dark. Following dehydration in

acetonitrile, 5 µg trypsin were added, and proteins digested for 20 h at 37 °C. Peptides were

extracted in 30% acetonitrile, 5% formic acid and 60% acetonitrile, 5% formic acid.

Combined extracts were freeze dried and kept at -20 °C until preparation for MS

measurements. Prior to MS measurement, the tryptic peptides were dissolved and desalted

as stated above.

Mass spectrometric analysis

Peptide samples were analyzed on a Q-Exactive HF mass spectrometer (Thermo Fisher

Scientific, Bremen, Germany) interfaced with an Easy-nLC 1200 nanoflow liquid

chromatography system (Thermo Scientific, Odense, Denmark). Samples were reconstituted

in 0.1% formic acid and loaded onto a C18 analytical column (75 μm × 15 cm). Peptides

were resolved at a flow rate of 300 nl/min using a linear gradient of 6−40% solvent B (0.1%

formic acid in 80% acetonitrile) over 165 min. Data-dependent acquisition with full scans over

a 350−1500 m/z range was carried out using the Orbitrap mass analyzer at a mass

resolution of 60,000 at 200 m/z, automatic gain control target value of $3e^6$, and max injection

time of 60 ms. The 10 most intense precursor ions were selected for further fragmentation.

Only peptides with charge states 2−6 were used, and dynamic exclusion was set to 30 sec.

Precursor ions were fragmented using higher-energy collision dissociation (HCD) with a

normalized collision energy (NCE) set to 28%. Fragment ion spectra were recorded at a

resolution of 15,000, automatic gain control target value of $1e^5$, and max. injection time 60

ms. Each of the biological replicates was measured in technical duplicates.

For label-free quantification, the raw files from LC-MS/MS measurements were

analyzed using MaxQuant[57] (version 1.6.8) with default settings and match between runs and

label-free quantification (LFQ) enabled. For protein identification, the human reference

proteome downloaded from the UniProt database (download date: 2018-02-22) was used.

For further data processing, Perseus software[58] (version 1.6.10.50) was used. Identified

proteins were filtered for reverse hits and common contaminants. LFQ intensities were log2

transformed. Only proteins that were detected in at least 4 out of 6 replicates (three biological

replicates, each measured as technical duplicate) were retained for further analysis. Missing

values were imputed from normal distribution (width = 0.3 and shift = 1.2) based on the

assumption that these proteins were just below the detection limit. Significantly enriched

proteins were identified by an ANOVA test (S0 = 2, FDR = 0.005) and Z-score normalized.

After averaging of groups by using the median of all replicates, enriched proteins were

clustered by correlation and plotted as heatmaps. Additional annotations (GO annotations,

KEGG, Pfam) and identification of enriched terms was done with Perseus, as described

previously[59]. The top 100 significantly enriched proteins were additionally used to compare

interactors between Ub 11AcK and Ub 11KeK and were plotted as Venn diagram.

For intact protein mass measurements, a micrOTOF II (Bruker) interfaced with an

Agilent 1260 Infinity II liquid chromatography system was used. The proteins were loaded

onto an analytical column (Nucleodur 300-5 C4 ec, EC 150/4, Macherey-Nagel) and resolved

at a flow rate of 300 µl/min using a linear gradient from 5% to 100% solvent B (0.1% formic

acid in acetonitrile) over 15 min. The electrospray ion source was operated in positive

ionization mode, scan range 650−2100 m/z at 1 Hz acquisition rate and 2x rolling average

was used. Compass Data Analysis software (Bruker) was used for re-calibration and spectra

deconvolution.

To quantify the p53 modification status, parallel reaction monitoring (PRM) using an

inclusion list and targeted MS2 mode was employed. Peptides were separated on a 15 cm

Acclaim PepMap C18 column (P/N 164943, Thermo Scientific) across a 50 min gradient from

2% to 45% solvent B (0.1% formic acid in 80% acetonitrile). A full mass spectrum was

acquired on an Orbitrap Tribrid Fusion (Thermo Fisher Scientific, Bremen, Germany) at 120

K resolution over a mass range of 330-1250 m/z (adapted to precursors in inclusion list). The

maximum injection time was set to 50 ms and an AGC target of 4e5. Precursors were

isolated sequentially according to the inclusion list with an isolation window width of 1.6 m/z

and fragmented by HCD at 30% NCE. Fragment mass spectra were acquired in the Orbitrap

at 30 K resolution and an AGC target of 5e4. PRM data were analyzed via Skyline software

23.1.0.268. MS1-filtering isotope precursor ion peak was set to 3 and peptides were

quantified on the MS2-level. Obtained values were plotted with GraphPad Prism 6.01.

***In vitro* binding assay with recombinant proteins**

Verifications of the protein interactions with the respective Strep-Ub variants were performed

as described previously[11]. In brief, 5 µM of the respective Strep-Ub variant were incubated 10

min on ice in binding buffer (1x PBS, 10 mM MgCl$_2$, 5 mM DTT) with 5 µM of recombinant

UCHL3 or NDP52. Then, 5 µl Strep-Tactin® beads (iba) pre-equilibrated in binding buffer

were added and the mixture incubated for 3 h at room temperature. After washing, bound

proteins were eluted twice by incubating the beads with 30 µl elution buffer (200 mM Tris-HCl

pH 8.0, 300 mM NaCl, 4 mM EDTA, 10 mM desthiobiotin). Eluates were combined,

subjected to SDS-PA gel electrophoresis, and analyzed by Coomassie blue staining and/or

western blot analysis.

Transient transfection and incorporation of AcK, TFAcK, and KeK into p53

$2\text{-}3 \times 10^5$ cells were seeded in 6-well plates (Sarstedt) 24 h before transfection. Cells were

transfected at an approximate confluency of 80% with Lipofectamine 2000 (Invitrogen-

Thermo Fisher Scientific) according to manufacturer's instructions with plasmids encoding

the AcK-RS/tRNA pair[28,29] required for incorporation of AcK and its analogs TFAcK and KeK

and an expression construct for p53 harboring a TAG stop codon at the position indicated. In

addition, an expression construct encoding β-galactosidase was cotransfected to determine

relative transfection efficiencies. Unless stated otherwise, 1 mM AcK, 8 mM TFAcK or 1mM

KeK (final conc.) were added at the time of transfection, and cells were harvested after 24

hours. After cell lysis, cleared whole cell extracts were subjected to β-galactosidase assay

and β-galactosidase normalized aliquots were electrophoresed on 12.5% SDS-PA gels

followed by western blot analysis with an anti-p53 antibody (DO-I) and an HRP-coupled anti-

mouse secondary antibody.

Luciferase reporter assays

Cells were transfected as stated above with expression constructs for the p53 variants

indicated, the AcK-RS/tRNA pair (if required), and β-galactosidase, and a p53-responsive

luciferase reporter construct (p21, PUMA). 24 h upon transfection, whole cell extracts were

prepared and 10 µl aliquots of each extract were transferred in duplicates to a black 96-well

plate and luciferase activity automatically determined by a micro-plate reader (Wallac 1420

VICTOR³ Multilabel plate reader, PerkinElmer). In brief, 100 µL of luciferase reaction mix

(470 µM D-luciferin, 530 µM ATP, 270 µM CoA, 200 µM EDTA, 20 mM tricine, 2.67 mM

$MgSO_4$, 1.07 mM $MgCO_3$, 33.3 mM DTT, pH 7.5) were dispensed into each well and the

reaction mixture incubated for 5 min. Luciferase activity was assessed by measuring light

emission for 10 seconds.

DATA AVAILABILITY

The data that support this study are available from the corresponding authors upon

reasonable request. The NMR solution structure and the crystal structure of ubiquitin used in

this study are available in the ProteinDataBank under the accession codes1D3Z and 1UBQ,

respectively. The mass spectrometry proteomics data have been deposited to the

ProteomeXchangeConsortium via the PRIDE repository[60] identifiers PXD056021 (Username:

reviewer_pxd056021@ebi.ac.uk; Password: AWE0DQbBpn9D) for the PRM data and

PXD056063 (Username: reviewer_pxd056063@ebi.ac.uk; Password: tCMcGvYVb79f) for

the AE-MS data.

REFERENCES

1. Ramazi, S. & Zahiri, J. Post-translational modifications in proteins: Resources, tools and prediction methods. *Database-the Journal of Biological Databases and Curation* (2021).
2. Aebersold, R. et al. How many human proteoforms are there? *Nat Chem Biol* **14**, 206-214 (2018).
3. Aebersold, R. & Mann, M. Mass-spectrometric exploration of proteome structure and function. *Nature* **537**, 347-355 (2016).
4. Conibear, A.C. Deciphering protein post-translational modifications using chemical biology tools. *Nat Rev Chem* **4**, 674-695 (2020).
5. Peng, T., Das, T., Ding, K. & Hang, H.C. Functional analysis of protein post-translational modifications using genetic codon expansion. *Protein Sci* **32**, e4618 (2023).
6. Barber, K.W. & Rinehart, J. The abcs of ptms. *Nat Chem Biol* **14**, 188-192 (2018).
7. Narita, T., Weinert, B.T. & Choudhary, C. Functions and mechanisms of non-histone protein acetylation. *Nat Rev Mol Cell Biol* **20**, 156-174 (2018).
8. Wang, Z.A. & Cole, P.A. The chemical biology of reversible lysine post-translational modifications. *Cell Chem Biol* **27**, 953-969 (2020).
9. Knyphausen, P., Lang, F., Baldus, L., Extra, A. & Lammers, M. Insights into k-ras 4b regulation by post-translational lysine acetylation. *Biol Chem* **397**, 1071-1085 (2016).
10. Okada, A.K. et al. Lysine acetylation regulates the interaction between proteins and membranes. *Nat Commun* **12**, 6466 (2021).
11. Kienle, S.M. et al. Electrostatic and steric effects underlie acetylation-induced changes in ubiquitin structure and function. *Nat Commun* **13**, 5435 (2022).
12. Liu, C.C. & Schultz, P.G. Adding new chemistries to the genetic code. *Annu Rev Biochem* **79**, 413-444 (2010).
13. Chin, J.W. Expanding and reprogramming the genetic code of cells and animals. *Annu Rev Biochem* **83**, 379-408 (2014).
14. Neumann, H., Peak-Chew, S.Y. & Chin, J.W. Genetically encoding n(epsilon)-acetyllysine in recombinant proteins. *Nat Chem Biol* **4**, 232-234 (2008).
15. Miller, G.M., Flynn, E.M., Tom, J., Song, A. & Cochran, A.G. Trifluoroacetyl lysine as a bromodomain binding mimic of lysine acetylation. *ACS Chem Biol* **17**, 1022-1029 (2022).
16. Huang, Y. et al. Genetic incorporation of an aliphatic keto-containing amino acid into proteins for their site-specific modifications. *Bioorg Med Chem Lett* **20**, 878-880 (2010).

672	17.	Reed, S.M. & Quelle, D.E. P53 acetylation: Regulation and consequences. *Cancers (Basel)* **7**, 30-69 (2014).
674	18.	Wen, J. & Wang, D. Deciphering the ptm codes of the tumor suppressor p53. *J Mol Cell Biol* **13**, 774-785 (2022).
676	19.	Xia, Z., Kon, N., Gu, A.P., Tavana, O. & Gu, W. Deciphering the acetylation code of p53 in transcription regulation and tumor suppression. *Oncogene* **41**, 3039-3050 (2022).
678	20.	Swatek, K.N. & Komander, D. Ubiquitin modifications. *Cell Res* **26**, 399-422 (2016).
679	21.	Yau, R. & Rape, M. The increasing complexity of the ubiquitin code. *Nat Cell Biol* **18**, 579-586 (2016).
681	22.	Dikic, I. & Schulman, B.A. An expanded lexicon for the ubiquitin code. *Nat Rev Mol Cell Biol* **24**, 273-287 (2023).
683	23.	Zhang, F. et al. A genetically encoded (19)f nmr probe for lysine acetylation. *Chem Commun (Camb)* **54**, 3879-3882 (2018).
685	24.	Kruiswijk, F., Labuschagne, C.F. & Vousden, K.H. P53 in survival, death and metabolic health: A lifeguard with a licence to kill. *Nat Rev Mol Cell Biol* **16**, 393-405 (2015).
687	25.	Kastenhuber, E.R. & Lowe, S.W. Putting p53 in context. *Cell* **170**, 1062-1078 (2017).
688	26.	Hafner, A., Bulyk, M.L., Jambhekar, A. & Lahav, G. The multiple mechanisms that regulate p53 activity and cell fate. *Nat Rev Mol Cell Biol* **20**, 199-210 (2019).
690	27.	Liu, Y. & Gu, W. The complexity of p53-mediated metabolic regulation in tumor suppression. *Semin Cancer Biol* **85**, 4-32 (2022).
692	28.	Elsässer, S.J. Generation of stable amber suppression cell lines. *Methods Mol Biol* **1728**, 237-245 (2018).
694	29.	Schmied, W.H., Elsässer, S.J., Uttamapinant, C. & Chin, J.W. Efficient multisite unnatural amino acid incorporation in mammalian cells via optimized pyrrolysyl trna synthetase/trna expression and engineered erf1. *J Am Chem Soc* **136**, 15577-15583 (2014).
697	30.	Mitsudomi, T. et al. P53 gene mutations in non-small-cell lung cancer cell lines and their correlation with the presence of ras mutations and clinical features. *Oncogene* **7**, 171-180 (1992).
700	31.	el-Deiry, W.S. et al. Waf1, a potential mediator of p53 tumor suppression. *Cell* **75**, 817-825 (1993).
702	32.	Arbely, E. et al. Acetylation of lysine 120 of p53 endows DNA-binding specificity at effective physiological salt concentration. *Proc Natl Acad Sci U S A* **108**, 8251-8256 (2011).
704	33.	Bullock, A.N., Henckel, J. & Fersht, A.R. Quantitative analysis of residual folding and DNA binding in mutant p53 core domain: Definition of mutant states for rescue in cancer therapy. *Oncogene* **19**, 1245-1256 (2000).
707	34.	Peterson, A.C., Russell, J.D., Bailey, D.J., Westphall, M.S. & Coon, J.J. Parallel reaction monitoring for high resolution and high mass accuracy quantitative, targeted proteomics. *Mol Cell Proteomics* **11**, 1475-1488 (2012).
710	35.	Julier, A., Radtke, V., Marx, A. & Scheffner, M. Generation and characterization of site-specifically mono-ubiquitylated p53. *Chembiochem* **23**, e202100659 (2022).
712	36.	Glozak, M.A., Sengupta, N., Zhang, X. & Seto, E. Acetylation and deacetylation of non-histone proteins. *Gene* **363**, 15-23 (2005).
714	37.	Choudhary, C., Weinert, B.T., Nishida, Y., Verdin, E. & Mann, M. The growing landscape of lysine acetylation links metabolism and cell signalling. *Nat Rev Mol Cell Biol* **15**, 536-550 (2014).
716	38.	Dancy, B.M. & Cole, P.A. Protein lysine acetylation by p300/cbp. *Chem Rev* **115**, 2419-2452 (2015).
718	39.	Drazic, A., Myklebust, L.M., Ree, R. & Arnesen, T. The world of protein acetylation. *Biochim Biophys Acta* **1864**, 1372-1401 (2016).
720	40.	Seto, E. & Yoshida, M. Erasers of histone acetylation: The histone deacetylase enzymes. *Cold Spring Harb Perspect Biol* **6**, a018713 (2014).
722	41.	Smith, B.C. & Denu, J.M. Mechanism-based inhibition of sir2 deacetylases by thioacetyl-lysine peptide. *Biochemistry* **46**, 14478-14486 (2007).

42. Farkas, M. et al. Distinct mechanisms control genome recognition by p53 at its target genes
linked to different cell fates. *Nat Commun* **12**, 484 (2021).
43. Sykes, S.M. et al. Acetylation of the p53 DNA-binding domain regulates apoptosis induction.
Mol Cell **24**, 841-851 (2006).
44. Vainer, R., Cohen, S., Shahar, A., Zarivach, R. & Arbely, E. Structural basis for p53 lys120-
acetylation-dependent DNA-binding mode. *J Mol Biol* **428**, 3013-3025 (2016).
45. Tang, Y., Luo, J., Zhang, W. & Gu, W. Tip60-dependent acetylation of p53 modulates the
decision between cell-cycle arrest and apoptosis. *Mol Cell* **24**, 827-839 (2006).
46. Mujtaba, S., Zeng, L. & Zhou, M.M. Structure and acetyl-lysine recognition of the
bromodomain. *Oncogene* **26**, 5521-5527 (2007).
47. Marchenko, N.D., Wolff, S., Erster, S., Becker, K. & Moll, U.M. Monoubiquitylation promotes
mitochondrial p53 translocation. *EMBO J* **26**, 923-934 (2007).
48. Nie, L., Sasaki, M. & Maki, C.G. Regulation of p53 nuclear export through sequential changes
in conformation and ubiquitination. *J Biol Chem* **282**, 14616-14625 (2007).
49. Zong, Z. et al. Alanyl-trna synthetase, aars1, is a lactate sensor and lactyltransferase that
lactylates p53 and contributes to tumorigenesis. *Cell* **187**, 2375-2392 e2333 (2024).
50. Stuber, K. et al. Structural and functional consequences of nedd8 phosphorylation. *Nat
Commun* **12**, 5939 (2021).
51. Hammler, D. et al. Fluorescently labelled atp analogues for direct monitoring of ubiquitin
activation. *Chemistry* **26**, 6279-6284 (2020).
52. Götz, K.H. et al. Formation of the alarmones diadenosine triphosphate and tetraphosphate by
ubiquitin- and ubiquitin-like-activating enzymes. *Cell Chem Biol* **26**, 1535-1543.e1535 (2019).
53. Delaglio, F. et al. Nmrpipe: A multidimensional spectral processing system based on unix pipes.
J Biomol NMR **6**, 277-293 (1995).
54. Johnson, B.A. & Blevins, R.A. Nmr view: A computer program for the visualization and analysis
of nmr data. *J Biomol NMR* **4**, 603-614 (1994).
55. Grzesiek, S., Stahl, S.J., Wingfield, P.T. & Bax, A. The cd4 determinant for downregulation by
hiv-1 nef directly binds to nef. Mapping of the nef binding surface by nmr. *Biochemistry* **35**,
10256-10261 (1996).
56. Shevchenko, A., Tomas, H., Havlis, J., Olsen, J.V. & Mann, M. In-gel digestion for mass
spectrometric characterization of proteins and proteomes. *Nat Protoc* **1**, 2856-2860 (2006).
57. Tyanova, S., Temu, T. & Cox, J. The maxquant computational platform for mass spectrometry-
based shotgun proteomics. *Nat Protoc* **11**, 2301-2319 (2016).
58. Tyanova, S. et al. The perseus computational platform for comprehensive analysis of
(prote)omics data. *Nat Methods* **13**, 731-740 (2016).
59. Krüger, L. et al. Chemical proteomic profiling reveals protein interactors of the alarmones
diadenosine triphosphate and tetraphosphate. *Nat Commun* **12**, 5808 (2021).
60. Perez-Riverol, Y. et al. The pride database resources in 2022: A hub for mass spectrometry-
based proteomics evidences. *Nucleic Acids Res* **50**, D543-D552 (2022).
61. Cornilescu, G., Marquardt, J.L., Ottiger, M. & Bax, A. Validation of protein structure from
anisotropic carbonyl chemical shifts in a dilute liquid crystalline phase. *J Am Chem Soc* **120**,
6836-6837 (1998).

ACKNOWLEDGEMENTS

We thank Andreas Marquardt and Anna Sladewska-Marquardt of the Proteomics Center of

the University of Konstanz for assistance with mass spectrometric experiments, and Silke

Büstorf and Nicole Richter-Müller for technical support. We are grateful to Dr. Kathrin Lang

and Dr. Simon Elsässer for providing cDNAs and plasmids encoding the acetyllysyl-tRNA

synthetase and the corresponding tRNA. This work was supported by the DFG (SFB969,

Projects B3 and B9). S.M.K. thanks the doctoral fund of the University of Konstanz and the

Konstanz Research School Chemical Biology for financial support. K.S. acknowledges the

Zukunftskolleg of the University of Konstanz for a doctoral fellowship. F.S. is grateful for

funding by the DFG (project grants 516836828 and 496470458).

AUTHOR CONTRIBUTIONS

S.M.K., M.S., M.Sch. conceived the study and experimental approach, S.M.K. generated all

Ub variants, K.S. provided E1, E2, UCHL3, and established the AE-MS workflow for Ub

variants, M.S. generated all p53 expression constructs, J.L. synthesized KeK. S.M.K.

performed HDM2 autoubiquitination assays, AE-MS with the Ub variants, and interaction

studies for AE-MS validations. T.S. conducted NMR spectroscopic analyses. M.S. performed

all transient transfection experiments with p53. J.J. performed the PRM analysis. S.M.K.,

M.S., T.S., J.L., J.J., F.S., A.M., M.K. and M.Sch. analyzed the data. S.M.K., M.S. and

M.Sch. wrote the paper. All authors provided critical feedback on the paper.

COMPETING INTERESTS

The authors declare no competing interests.

Fig. 1. The length of the side chain at position 11 of Ub affects HDM2 autou-biquitination. a, Comparison of the length (Å) of the side chain of amino acids incorporated at position 11 of Ub. **b,** HDM2 autoubiquitination assay with Ub variants, in which K11 was replaced by AcK, TFAcK, KeK, Q or R, indicate that the length of the side chain at position 11 is the main determinant for the performance of Ub variants in HDM2 autoubiquitination. Autoubiquitination reactions were started by addition of the respective Ub variant. Reactions were stopped at the times indicated. Reactions in the absence of Ub (-Ub) were stopped at 60 min. All reactions were analyzed by SDS-PAGE followed by Coomassie blue staining. Running positions of molecular mass markers (kDa), HDM2, UBA1 (E1), UbcH5b (E2), free Ub (Ub), and autoubiquitinated forms of HDM2 (*) are indicated. Quantitative incorporation of the respective amino acids and the homogeneity of the purified Ub variants were confirmed by ESI-MS analysis (Supplementary Fig. 1).

Fig. 2. Incorporation of AcK, TFAcK or KeK at position 11 of Ub results in similar structural changes. a-c, Weighted chemical shift perturbations (CSPs, Δω) of backbone amide resonances were calculated comparing the differently modified Ub variants (11AcK, 11TFAcK, 11KeK) with unmodified Ub (11K). The corresponding two-dimensional heteronuclear ^{1}H-^{15}N HSQC NMR spectra are shown in Supplementary Fig. 2. In the upper panel, the corresponding CSP plots are presented for Ub 11AcK (a), Ub 11TFAcK (b), Ub 11KeK (c), while in the lower panel the respective CSP values are highlighted on the NMR solution structure of unmodified Ub (PDB ID 1D3Z)[61]. The Ub structure is illustrated in cartoon mode and the respective CSP values are indicated by using a continuous color palette ranging from white (Δω = 0.00 ppm) to red (Δω ≥ 0.15 ppm). Additionally, the position of K11 and its side chain are depicted by a blue circle and by anthracite sticks, respectively. In the CSP plots, the horizontal solid and dashed lines are indicating the mean and the mean plus one standard deviation, respectively, calculated over all CSP values from (a) to (c) excluding values related to position 11. On top of the plots, position 11 is indicated by an arrow and the secondary structural elements are schematically depicted according to the NMR solution structure of unmodified Ub.

Fig. 3. Ub variants containing AcK or its analogs TFAcK and KeK resemble one another with respect to their protein interaction properties. a, Schematic overview of the AE-MS workflow for the identification of interactors of the Ub variants indicated. Unmodified Ub (11K) and each of the six Ub variants were used as bait molecules for affinity enrichment and incubated with HEK293T cell extract. Empty beads (beads) were used as control. **b,** Hierarchical clustering of statistically significant interactors (rows) of the different bait molecules (columns). Red indicates enrichment, whereas blue indicates lack of enrichment. Thresholds for ANOVA statistics were set to S0=2 and FDR=0.005. Cluster numbers, indicating binding to one or multiple Ub variants, are indicated on the left. **c,** Correlation heatmap of significantly enriched interactors of the differently modified Ub variants. Strong correlation is indicated in red, no correlation in white, and inverse correlation in gray.

Fig. 4. Incorporation of AcK and its analogs TFAcK and KeK into p53 by genetic code expansion in H1299 cells. a, Incorporation of AcK, TFAcK, and KeK at position 120 of p53. **b,** Quantification of relative p53 levels in the presence of AcK, TFAcK, and KeK shown in **a**. Levels of wild-type p53 (wt) were set to 100 percent. **c,** Incorporation of AcK, TFAcK, and KeK into position 164 of p53. H1299 cells were transfected with expression constructs encoding wild-type p53 (wt) or p53 with a TAG stop codon at position 120 and 164, respectively, and the AcK-RS/tRNA pair as described in Methods. Growth medium was supplemented with increasing concentrations of AcK, TFAcK, or KeK as indicated. 24 h upon transfection**,** p53 levels were determined by western blot analysis with the p53-specific DO-1 antibody (**a, c**).

Fig. 5. Transactivation activity of p53 variants. a, H1299 cells were transfected with a p53-
responsive expression construct encoding luciferase in the absence (ctrl) or presence of constructs
encoding wild-type p53 (wt), the tumor-derived p53 mutant R273H, p53 with a TAG stop codon at
position 120, and the AcK-RS/tRNA pair. In addition, cells transfected with the p53-120TAG construct
were grown in the absence (120TAG) or presence of AcK (120AcK), TFAcK (120TFAcK) or KeK
(120KeK). 24 h upon transfection, luciferase activity was determined as a measure for p53 activity and
the relative values obtained were adjusted for transfection efficiency (n = 4 ± SD independent
experiments). 20 percent of the extracts were subjected to western blot analysis using the p53-specific
DO-1 antibody. **b,** H1299 cells were transfected as in **a** in the absence of the p53-responsive reporter
construct. 24 h upon transfection, levels of endogenous p21 protein and of ectopically expressed p53
were determined by western blot analysis using a p21-specific antibody and the p53-specific DO-1
antibody, respectively. **c,** H1299 cells were transfected with expression constructs encoding p53-
120TAG and the AcK-RS/tRNA pair. Cells were grown in the presence of AcK, KeK, and TFAcK as
indicated and the presence of AcK, KeK, TFAcK or lysine (unmodified) at position 120 of p53 was
determined by PRM analysis (n = 3 ± SD independent experiments). **d,** Transactivation activity of the
p53 variants 120Q and 120R was determined as described in **a** (n = 4 ± SD independent
experiments).